A conserved HSF:miR169:NF-YA loop involved in tomato and Arabidopsis heat

stress tolerance

Sombir Rao[1], Chandni Bansal[1], Celine Sorin[2,3], Martin Crespi[2,3] and Saloni Mathur[1]*

Author Affiliations:

[1] National Institute of Plant Genome Research, Aruna Asaf Ali Marg, P.O. Box No.

10531, New Delhi - 110 067

[2] Université Paris-Saclay, CNRS, INRAE, Univ Evry, Institute of Plant Sciences Paris-

Saclay (IPS2), 91405, Orsay, France.

[3] Université de Paris, CNRS, INRAE, Institute of Plant Sciences Paris-Saclay (IPS2),

91405, Orsay, France

Email: Sombir Rao (sombirrao97@gmail.com); Chandni Bansal

(bansal16.chandni@gmail.com); Celine Sorin (celine.sorin@universite-paris-saclay.fr);

Martin Crespi (martin.crespi@ips2.universite-paris-saclay.fr) and Saloni Mathur

(saloni@nipgr.ac.in)

*Corresponding author:

Dr. Saloni Mathur

Staff Scientist IV

National Institute of Plant Genome Research

Email: saloni@nipgr.ac.in

Phone: 91-11-26735175

Running title: An HSF: miR169:NF-YA module for thermotolerance

Abstract

Heat stress transcription factors (HSFs) and miRNAs regulate different stress and
developmental networks in plants. Regulatory feedbacks are at the basis of these
networks. Here, we report that plants improve their heat stress tolerance through HSF-
mediated transcriptional regulation of *MIR169* and post-transcriptional regulation of *NF-*
YA transcription factors. We show that HSFs recognize tomato and Arabidopsis *MIR169*
promoters using yeast-one-hybrid/ChIP-qPCR. Silencing tomato HSFs using virus
induced gene silencing (VIGS) reduced *Sly-MIR169* levels and enhanced *Sly-NF-*
YA9/A10 target expression. Further, *Sly-NF-YA9/A10-VIGS* knock-down tomato plants
and Arabidopsis plants overexpressing *At-MIR169d* or *At-nf-ya2* mutants showed a link
with increased heat tolerance. In contrast, Arabidopsis plants overexpressing *At-NF-*
YA2, or those expressing a non-cleavable *At-NF-YA2* form (miR169d-resistant *At-NF-*
YA2) as well as plants inhibited for At-miRNA169d regulation (miR169d mimic plants)
were more sensitive to heat stress, highlighting *NF-YA* as negative regulator of heat
tolerance. Furthermore, post-transcriptional cleavage of *NF-YA* by elevated miR169
levels resulted in alleviating the repression of heat stress effectors HSFA7a/b in tomato
and Arabidopsis revealing a retroactive control of HSFs by the miR169:NF-YA node.
Hence, a regulatory feedback loop involving HSFs, miR169s and NF-YAs plays a critical
role in the regulation of heat stress response in tomato and Arabidopsis plants.

Introduction

Heat stress (HS) adversely affects the distribution and productivity of agriculturally
important plants worldwide by affecting all aspects of plant processes like germination,
growth, development, reproduction and yield (Mittler et al. 2010; Lobell et al. 2011;
Hasanuzzaman et al. 2013). In nature, such HS conditions may be chronic or recurring,
or both (Bäurle 2016); therefore, plants have evolved a variety of responses to elevated
temperatures that minimize damage and ensure protection of cellular homeostasis.

HSFs are important regulators that play critical roles in signal transduction processes to
mediate gene expression in response to multiple abiotic stresses, including cold,

drought, salt, and heat (Collins et al. 1995; Nover et al. 2001; Guo et al. 2016; Jacob et al. 2017). HSFs recognize heat stress elements (HSE: 5′-GAAnnTTC-3′) that are conserved in promoters of heat stress-responsive (HSR) genes including HSP genes (Busch et al. 2005). During the initial stages of HS, HSFs are the primary molecules responsible for relaying signals of cellular stress to the transcriptional apparatus and reprogram gene expression to repair protein damage through elevated synthesis of molecular chaperones and proteases (Morimoto 2002; Scharf et al. 2012; Guo et al. 2016). In plants, there are multiple genes coding for HSFs for example, Arabidopsis has 21 HSF family members (von Koskull-Dˆring et al. 2007) and tomato has 26 HSFs (Yang et al. 2016). Another class of regulatory molecules, the microRNAs (miRNAs) have emerged as a major class of small non-coding RNAs with roles in plant growth and development as well as under stress (Sunkar and Zhu, 2004; Lin et al. 2018; Ravichandran et al. 2019). HSF-mediated induction of MIR398 has been shown to trigger a regulatory loop that is critical for thermotolerance in Arabidopsis (Guan et al. 2013). Yan et al. (2012) have shown that stress-induced alternative splicing of MIR400 provides a mechanism for the regulation of miRNA processing in Arabidopsis during HS. Lin et al. (2018) have shown that miR160 modulates plant development and heat shock protein gene expression to mediate heat tolerance in Arabidopsis. Heat-responsive miRNAs in *Apium graveolens* (Li et al. 2014), switch grass (Hivrale et al. 2015), banana (Vidya et al. 2018) and apple (Niu et al. 2019) have also been reported.

The miR169 family members have significant roles in plant abiotic and biotic stress responses and are involved in various aspects of plant development including root architecture, nodule formation etc. by targeting *Nuclear factor-Y A* (NF-YA) gene family members (Zhao et al. 2011; Ni et al. 2013; Sorin et al. 2014; Zhang et al. 2015; Li et al. 2017). Tomato plants overexpressing miR169 show improved resistance to drought stress (Zhang et al. 2011) while in Arabidopsis enhanced miR169 levels promote leaf dehydration (Li et al. 2008; Ni et al. 2013). Over-accumulation of miR169 is correlated with a reduction of *NF-YA* target transcripts in Arabidopsis seedlings under cold stress (Zhou et al. 2008; Lee et al. 2010). Role of miR169 is also established in long-distance signalling and nitrogen/phosphorus starvation (Pant et al. 2009; Buhtz et al. 2010; Zhao et al. 2011). In a previous publication from our laboratory, we have reported 18 *MIR169*

family members in tomato (Rao et al. 2020). Assessment of *Sly-MIR169* precursors in different heat stress (HS) regimes highlighted that fifteen of the sixteen precursors (two of the loci are bicistronic) are differentially regulated in response to HS. Heat-mediated up-regulation of *MIR169*s has also been reported in Arabidopsis (Li et al. 2010; Guan et al. 2013) however, a detailed mechanism was lacking. Here, we have dissected the transcriptional regulation of *MIR169* family by HSFs in tomato and Arabidopsis HSR. Our laboratory has previously established miR169-mediated transcript cleavage of five classical targets belonging to the *NF-YA* class of transcription factors (*Sly-NF-YA1, Sly-NF-YA3, Sly-NF-YA8, Sly-NF-YA9 and Sly-NF-YA10*) (Rao et al. 2020). Here, we show that miR169 mediated post-transcriptional down-regulation of *At-NF-YA2* in *Arabidopsis* and *Sly-NF-YA9* and *Sly-NF-YA10* in tomato enhances HS tolerance and impacts HSF expression. NF-YA transcriptionally regulates a suite of HSR genes including HSFs which exert a feedback regulation on *NF-YAs* through the control of miR169 expression. This is a new feedback loop involved in the regulation of plant tolerance to HS.

Results

MIR169 members are transcriptionally regulated by HSFs in tomato

The transcriptional activation of heat stress genes is regulated by HSFs that recognize HSE cis-elements in promoters of HSR genes (Busch et al. 2005). All the 15 heat up-regulated *Sly-MIR169* promoters (Rao et al. 2020), contain one or more HSEs indicating possible involvement of HSFs in their heat inducibility (Supplementary figure S1, Supplementary table S1). This prompted us to investigate the transcriptional regulation of *MIR169* promoters by HSFs using yeast-one-hybrid (Y1H) assays. Out of 322 interactions studied between tomato HSFs and 14 heat-responsive *Sly-MIR169* promoters (one promoter could not be assessed as a minimal inhibitory Aureobasidin A antibiotic concentration could not be achieved), we found specific binding of seven HSFs on five *Sly-MIR169* promoters (Figure 1A, Supplementary table S2, Supplementary figure S2). Promoters of *Sly-MIR169a-1* and *Sly-MIR169d-1* are regulated by a single HSF viz., Sly-HSFA1e and Sly-HSFA1a, respectively (Figure 1A, Supplementary table S2). *Sly-MIR169b* promoter interacts with Sly-HSFA1a and Sly-

HSFA7a while *Sly-MIR169c* promoter shows Sly-HSFA4a and Sly-HSFA6b binding.

Transcription of *Sly-MIR169d* is regulated by Sly-HSFA8a and Sly-HSFB1a (Figure 1A).

Further, to confirm the binding of these Sly-HSFs on *Sly-MIR169* promoters, we

performed chromatin immunoprecipitation (ChIP) assays by co-expressing specific *Sly-*

MIR169 promoters and *P35S:HSF:GFP* constructs in *Nicotiana benthamiana* leaves.

ChIP-qPCR was then performed on the HSE-specific regions in the *Sly-MIR169*

promoters. Fold enrichment was calculated by comparing the Ct values of anti-GFP

immunoprecipitated samples with IgG immunoprecipitation control (Figure 1B-I). ChIP-

qPCR analysis of all HSFs with *Sly-MIR169* promoters matched the Y1H results except

for HSFA4a and HSFA8a on *Sly-MIR169c* and *Sly-MIR169d* promoters (Figure 1E, G).

The HSF-mediated regulation of *MIR169* promoters was also assessed in *Nicotiana*

benthamiana leaves using agro-infiltration (Supplementary Figure S3). For this, *Sly-*

MIR169 promoters were fused with *GUS* reporter gene and co-transformed with effector

plasmids containing HSFs under CaMV35S promoter. All the five HSFs showing

promoter enrichment in the ChIP assays were transformed individually or in

combinations (as the case may be) to check their ability to drive *GUS* gene expression

under respective *Sly-MIR169* promoters. qRT-PCR based estimation of *GUS* transcripts

in presence of HSFs confirmed HSF-mediated activation of *Sly-MIR169* promoters for

all the HSFs (Supplementary Figure S3) except Sly-HSFA1a (Supplementary Figure

S3B). However, co-transformation of Sly-HSFA1a with Sly-HSFA7a (Supplementary

Figure S3B) leads to enhanced levels of *GUS* transcripts as compared to individual

HSFs suggesting a cooperative action in regulating *Sly-MIR169b* promoter.

HSFs regulate the orchestration of *miR169:NF-YA* module during heat stress in

tomato

To assess whether the HSF:*miR169:NF-YA* module is functional in tomato HS, we

quantified the transcript abundance of all components of this module in wild type tomato

plants upon HS. Expression profiles of HSFs reveal the heat-mediated up-regulation of

Sly-HSFA6b, Sly-HSFA7a and *Sly-HSFB1a* while *Sly-HSFA1a* and *Sly-HSFA1e* are not

HS responsive (Figure 2A, Supplementary figure S4A). HSFA1s, the master regulators

of HSR response, are present in bound forms with HSPs in cytoplasm. Upon HS, HSFs

dissociate from HSPs and move to nucleus to transcriptionally regulate HSR genes

(Kotak et al. 2004; Chan-Schaminet et al. 2009) hence, HSFAs do not exhibit

transcriptional up-regulation upon HS. However, we note that in non-stressed conditions

the Sly-*HSFA1s* are more abundant in comparison to other HSFs (by comparing Ct

values of these HSFs), suggesting their sufficient levels to orchestrate gene

transcription upon HS (Supplementary table S3A). In agreement with our previous

results (Rao et al. 2020), four miR169 mature forms (Figure 2B, Supplementary figure

S4B) exhibit heat-mediated induction in comparison to non-stressed WT tomato plants.

Comparison of their Ct values in control conditions revealed the abundant presence of

miR169c in non-stressed plants, while imposition of heat stress led to strong induction

of miR169a (Supplementary table S3B). Furthermore, expression analysis of miR169s

targets (NF-YAs) shows reduction of only *Sly-NF-YA9* and *Sly-NF-YA10* transcripts in

HS in WT plants (Figure 2C, Supplementary figure S4C).

To investigate the direct role of HSF-mediated regulation of the *MIR169:NF-YA* module

in tomato, we utilized TRV-based virus induced gene silencing (VIGS) of HSFs. The

VIGS silencing assays were performed 3-weeks-post agro-infiltration. Setting up of

efficient silencing for TRV-HSFs was assessed by the reduction in the endogenous

HSFs transcripts in silenced plants (Figure 2D-G). To ensure the specific

downregulation of selected HSFs, we have examined the expression profiles of putative

off targets HSFs predicted by SGN-VIGS tool (Supplementary table S4) by qRT-PCR.

The unaltered expression of putative off targets in different HSFs-VIGS silenced plants

confirms the specificity of the silencing (Supplementary figure S5A-C). Subsequently,

downstream components in the pathway like *MIR169* precursor levels and their heat

responsive *NF-YA* targets (*Sly-NF-YA9* and *Sly-NF-YA10*) were quantified by qRT-PCR

after these plants were given a HS (Figure 2D-G, Supplementary figure S6A-D).

Silencing of *Sly-HSFA1a* and *Sly-HSFA7a* reduces the transcription of *Sly-MIR169b* by

3 and 5 fold respectively, and leads to more than 2 fold enhanced expression of *Sly-NF-*

YA9 and *Sly-NF-YA10* transcripts upon HS (Figure 2D, E). Transcription of *Sly-*

MIR169c is reduced by 2.6 folds in *Sly-HSFA6b* silenced plants, this in turn leads to up-

regulation of *Sly-NF-YA10* upon HS (Figure 2F). Silencing of *Sly-HSFB1a* (Figure 2G)

reduces the expression of *Sly-MIR169d* by 2.9 folds, resulting in enhanced

accumulation of *Sly-NF-YA9* and *Sly-NF-YA10* transcripts upon HS. *Sly-HSFA6b*
silencing also reduces the levels of *Sly-MIR169d-1* by 4.9 folds that in-turn enhances
the transcript levels of *Sly-NF-YA10* upon HS (Figure 2F). Depressed expression of *Sly-*
MIR169s with concomitant enhanced expression of *Sly-NF-YA9/10* in *Sly-HSF*-silenced
backgrounds upon HS suggests the HSF-mediated regulation of *Sly-MIR169:Sly-NF-*
YA9/10 module in HS.

The miR169:NF-YA module is critical for high temperature tolerance in tomato

To establish the functions of *Sly-NF-YA9* and *Sly-NF-YA10* in heat tolerance, we used
the VIGS-silencing approach to knock-down their expression in tomato plants and
assessed their thermotolerance post HS. Three-weeks-after agro-infiltration, a silencing
efficiency of ~80% was achieved for both *Sly-NF-YA9* and *Sly-NF-YA10* transcripts as
compared to the TRV-vector negative control plants (Supplementary Figure S7A).
Expression analysis of Sly-*NF-YA9* or Sly-*NF-YA10* in WT control (non-stressed) and
TRV-EV infiltrated plants revealed no significant fluctuation in their expression levels
(Supplementary figure S7B), suggesting the specificity of the VIGS silencing constructs.
The *TRV-Sly-NF-YA9* and *TRV-Sly-NF-YA10* plants were given a HS and scored for
heat tolerance six-days-after HS. qRT-PCR based expression analysis revealed that a
strong downregulation of *Sly-NF-YA9* and *Sly-NF-YA10* is maintained in VIGS silenced
plants post HS (Supplementary figure S7C-D), ruling out any time point and treatment
biased alteration in their expression levels. Phenotypic evaluation post-HS showed that
the leaves in vector control plants are highly damaged in these conditions, in contrast to
the *Sly-NF-YA9* and *Sly-NF-YA10* silenced plants which are robust with green, healthy,
erect leaves and also show emergence of new leaves (Figure 3A). The *Sly-NF-YA9* and
Sly-NF-YA10 knock-down plants show ~55% and 80% survival rate (judged by
phenotype of 7^{th} to 9^{th} leaves and emergence of new leaves) in comparison to ~33% for
vector-control plants (Figure 3B). Further, we gauged the physiological performance of
TRV-Sly-NF-YA9 and *TRV-Sly-NF-YA10* silenced plants, six days after HS using a Li-
Cor 6400 photosynthesis measuring system. Both the *NF-YA* silenced plants show
significant enhancement in water use efficiency and net photosynthetic rates as
compared to TRV-control plants under HS (Figure 3C). To rule out that the post HS

phenotyping and survival scoring is not because of an effect of silencing of off targets,
we predicted any possible off targets of *Sly-NF-YA9* and *Sly-NF-YA10* genes. Unaltered
expression patterns of these putative off target genes in *Sly-NF-YA9* and *Sly-NF-YA10*
VIGS silenced plants confirmed that the phenotype observed is specific to the knock-
down of *Sly-NF-YA9* and *Sly-NF-YA10* genes (Supplementary table S4, Supplementary
figure S8A-B).

To further delineate the molecular events driving the improved thermotolerance of *TRV-*
Sly-NF-YA9 and *TRV-Sly-NF-YA10* knock-down plants, we checked whether the
expression of conserved HSR genes like *HSFs (Sly-HSFA3a* and *Sly-HSFA7a), Sly-*
sHSP and *Sly-APX (ascorbate peroxidase)* is significantly modulated upon heat stress*.*
First, to ascertain that these genes are reliable HSR genes in the VIGS conditions, we
show that they all are upregulated in TRV-EV plants under heat stress in comparison to
non-stressed control TRV-EV plants (Figure 3D-H, Supplementary figure S9A-E).
Furthermore, these HSR genes are inducible in *TRV-Sly-NF-YA9* and *TRV-Sly-NF-*
YA10 knock-down plants during heat stress as compared to non-stressed conditions
(Figure 3D-H, Supplementary figure S9A-E). The strong upregulation of HSR genes in
heat stressed *TRV-Sly-NF-YA9* and *TRV-Sly-NF-YA10* knock-down plants in
comparison to TRV-EV HS plants confirms the direct involvement of these HSR genes
in regulating the thermotolerance of tomato plants (Supplementary figure S9F). Further,
since all these HSR genes (except *Sly-APX*) exhibit upregulation in *TRV-Sly-NF-YA9*
and *TRV-Sly-NF-YA10* knock-down plants in comparison to TRV-EV plants in non-
stressed conditions (Supplementary figure S9G), this indicates that these HSR genes
(except *Sly-APX*) are downstream targets of *Sly-NF-YA9* and *Sly-NF-YA10.*
Furthermore, promoter analysis (1kb) of these HSR genes revealed the presence of
CCAAT boxes in *Sly-HSFA3a* and *Sly-HSFA7a* (Supplementary table S5)*,* suggesting a
direct Sly-NF-YA9/10 mediated regulation on these genes and an indirect regulation on
other HSR genes. Since the HSR genes are upregulated in *TRV-Sly-NF-YA9* and *TRV-*
Sly-NF-YA10 knock-down plants, these results suggest that *Sly-NF-YA9* and *Sly-NF-*
YA10 act as negative regulators of thermotolerance.

***MIR169s* are regulated by HSFs during heat stress in Arabidopsis**

To assess whether the HSF:miR169:NF-YA module under HS is active in other plants
too, we decided to evaluate this module in the model plant Arabidopsis. Guan et al.
(2013) have reported heat-mediated induction of miR169 and miR398 by Northern
hybridization in Arabidopsis. However, the regulation of miR169 in the heat responses
of Arabidopsis remains unknown. Using taqman based qRT-PCR analysis of WT control
and heat stress challenged Arabidopsis plants, we show that all four mature miR169
forms are strongly upregulated (≥5 fold) under HS (Figure 4A, Supplementary figure
S10). Further, to support a probable mechanism of their heat inducibility, promoters of
all 14 *At-MIR169* genes (Li et al. 2010) were screened for the presence of HSEs and we
found two to multiple HSEs in all *MIR169* promoters (Supplementary table S6). To
delineate this speculated interplay among Arabidopsis HSFs and *At-MIR169* promoters,
Y1H assays were performed (Supplementary table S7) with 21 At-HSFs and 11 *At-
MIR169* promoters (as six precursors are transcribed as three bi-cistronic units viz., Pro-
At-MIR169i/j, Pro-*At-MIR169k/l*, Pro-*At-MIR169m/n*; Li et al. 2010). This analysis
identified 5 HSFs interacting with 3 *At-MIR169* promoters (Figure 4B). Three HSFs (At-
HSFB3a, At-HSFA7b, At-HSFB2b) regulate *At-MIR169b*; At-HSFA2a controls *At-
MIR169d* transcription and transcription of *At-MIR169h* is regulated by At-HSFA1d. At-
HSFA2 is considered as the key regulator of Arabidopsis HS-response and is required
for the maintenance of acquired thermotolerance (Charng et al. 2007). Therefore, we
decided to dissect the At-HSFA2 mediated *At-MIR169d* module for characterization of
HS-response and thermotolerance in Arabidopsis. In wild type Arabidopsis plants, we
find strong induction of *At-MIR169d* precursor (≥15 fold) upon HS (Figure 4C,
Supplementary figure S11). Transgenic promoter-*At-MIR169d:GUS* plants support this
heat-dependent transcriptional acceleration (Figure 4D, E). Moreover, *GUS* aided
reporter analysis of promoter-*At-MIR169d:GUS* in WT and in the *At-hsfa2* knockout
mutant (Supplementary figure S12A, B, C) background in HS highlights minimal *GUS*
expression in the *At-hsfa2* mutant as compared to WT plants (Figure 4F, G). Further,
there is strong reduction of *At-MIR169d* precursor levels in *At-hsfa2* mutant as
compared to WT plants (Figure 4G) confirming At-HSFA2-mediated regulation of *At-
MIR169d* in Arabidopsis HS and suggesting a general regulatory loop in plants.

The miR169d:NF-YA2 regulatory node is heat responsive and spatially co-expressed in Arabidopsis

To explore the functional significance of heat-mediated regulation of *MIR169s*, we assessed the expression of the eight canonical *NF-YA* targets (Sorin et al. 2014) in HS in Arabidopsis. qRT-PCR based profiling shows significant reduction of *At-NF-YA2* (-3.5 fold), *At-NF-YA3* (-2.2 fold), *At-NF-YA6* (-2.1 fold) and *At-NF-YA10* (-2.1 fold) transcripts in WT plants when exposed to HS (Supplementary figure S13A, B). Further, to assign the functional relevance of the selected *At-MIR169d* isoform in particular, we analyzed target mimicry plants generated by overexpressing MIM169defg constructs under CaMV35S, using constructs from Todesco et al. (2010). The expression of 4 mature forms of At-miR169s in these MIM169defg lines revealed that miR169defg is the most reduced form (Supplementary figure S14A). The target mimic lines sequester the specific mature miRNAs by acting as non-cleavable targets; thereby they protect the target genes from miRNA cleavage (Todesco et al. 2010). The expression of *At-NF-YA2, At-NF-YA3, At-NF-YA6, At-NF-YA8 and At-NF-YA10* is significantly increased in *MIM169defg* lines as compared to WT plants, with *At-NF-*YA2 transcript levels being highest in the *MIM169defg* lines (Supplementary figure S14B). Moreover, overexpressing *At-MIR169d* (At-MIR169d-OE) precursor under constitutive CaMV35S promoter in Arabidopsis strongly represses (-4 to -11-fold in three different *At-MIR169d-OE* lines) *At-NF-YA2* transcripts, establishing it as the major miR169d target (Supplementary figure S14C). Moreover, promoter-*At-NF-YA2* lines do not show HS-mediated up-regulation specifying a pure post-transcriptional regulation (Supplementary figure S15A, B).

To delineate the existence and localization of miR169d:*At-NF-YA2* functional module in plant tissues and development stages, promoter reporter (*GUS)* lines were generated by transforming WT Arabidopsis plants with constructs that contained the 2 kb genomic sequences upstream of the *At-MIR169d* precursor. Parallelly, 2 kb region ahead of the translation start site of *At-NF-YA2* was similarly assessed. The *At-MIR169d* and *At-NF-YA2* promoters show prominent GUS localization in the vasculature of roots and aerial parts of seedling, mature rosette/cauline leaves and sepals in flowers of Arabidopsis,

296 suggesting functional co-existence of the miR169d:*NF-YA2* module (Supplementary

297 figure S16).

298 Arabidopsis plants with enhanced miR169d or reduced *NF-YA2* expression are

299 thermotolerant

To further investigate the miR169d:*NF-YA2* module in HS-response in Arabidopsis, we

assayed 2-weeks-old plants of *At-MIR169d*-OE, *At-NF-YA2*-OE (*NF-YA2*

overexpressed under 35SCaMV promoter, Supplementary figure S17), *At-nf-ya2*

knockout mutant (Supplementary figure S18), an miR169d resistant version of *At-NF-*

YA2 under native promoter (*pNF-YA2:NF-YA2*-r) and MIM169defg lines for their ability

to survive HS. The survival rate of plants was gauged 6-days-after exposure to 2 h of

HS at 42 ☐C. Plants overexpressing *At-MIR169d* (*At-MIR169d*-OE) that have reduced

target *At-NF-YA2* levels and plants that lack *At-NF-YA2* transcripts (*At-nf-ya2* mutant)

exhibit twice the survival rate than WT plants, in contrast, plants with high abundance of

At-NF-YA2 transcripts viz. *At-NF-YA2*-OE, *MIM169defg* and the *pNF-YA2:NF-YA2*-r (a

variant *At-NF-YA2* transcript that allows miRNA169 binding but escapes miRNA-

mediated cleavage) are nearly 3-folds more HS sensitive (Figure 5A, B). While only

~10% plants survive in transgenic plants overexpressing *At-NF-YA2* mRNAs, as much

as 80-90% plants survive the HS in *At-MIR169d-OE* and *At-nf-ya2* mutant plants. This

establishes that HS-tolerance in Arabidopsis by At-miR169d is mediated by down-

regulation of *At-NF-YA2* functions wherein, *At-NF-YA2* acts as negative regulator of

heat stress-response.

To further explore the mechanism underlying the better survival of *At-MIR169d*-OE and

At-nf-ya2 lines as well as the sensitivity of *At-NF-YA2*-OE, MIM169defg and *pNF-*

YA2:*NF-YA2*-r lines, we analyzed the expression of several heat responsive genes

reported in literature (Larkindale and Vierling 2008) in all these lines. Expression

profiles of 11 HSR genes in response to HS (Supplementary table S8) revealed that the

improved thermotolerance in *At-MIR169d*-OE and *At-nf-ya2* mutant plants is correlated

with enhanced expression of *At-Ascorbate peroxidase2* (*At-APX2*), *At-HSFA3a*, *At-*

HSFA7a, *At-HSFA7b* and *At-HSP17.ACI* genes (Figure 5C, Supplementary figure S19).

In contrast, transgenic plants *At-NF-YA2*-OE, *pNF-YA2:NF-YA2*-r, MIM169defg that are

defective in heat-responsive gene regulation, exhibit reduced *At-APX2, At-HSFA3a, At-HSFA7a, At-HSFA7b, and At-HSP17.ACI* transcripts (Figure 5C, Supplementary figure S19). These results ascertain At-NF-YA2 as negative regulator of many HSR genes and thermotolerance in Arabidopsis.

It is well established that imposition of HS can give rise to excess levels of reactive oxygen species (ROS), resulting in cellular oxidative damage. *APX* plays crucial role as H_2O_2-scavenging enzyme in plant cells. Therefore, a higher level of *At-APX2* transcripts in *At-MIR169d*-OE and *At-nf-ya2* mutant plants should effectively maintain the antioxidants that protect plants. Indeed, these plants have low H_2O_2 levels as judged by DAB staining (Figure 5D). In contrast, the overexpression of *At-NF-YA2* gene (*At-NF-YA2-OE*) and chelation of At-miR169d (MIM169defg lines) and *pNF-YA2:NF-YA2*-r show enhanced DAB staining due to higher H_2O_2 levels (Figure 5D). Further, cellular death estimation as a consequence of this in *At-MIR169d*-OE and *At-nf-ya2* lines is negligible while it is much pronounced in the *At-NF-YA2*-OE, *MIM169defg* and *pNF-YA2:NF-YA2*-r lines, as assessed by trypan blue staining (Figure 5E).

Arabidopsis At-NF-YA2 negatively regulates *HSR* genes leading to HS sensitivity

Repressed expression of *HSR* genes in *At-NF-YA2* abundant lines and enhanced expression of *HSR* genes in *At-NF-YA2* down-regulated lines confirms that *At-NF-YA2* is a negative regulator of these HSR genes (Figure 5C, Supplementary figure S19). *At-HSP17.ACI*, *At-HSFA3a*, *At-HSFA7a, At-HSFA7b* and *At-APX2* promoters have NF-YA binding sites (CCAAT-box), making them probable downstream targets of At-NF-YA2 transcription factor (Figure 6A). Indeed, ChIP-qPCR and Y1H assay shows that the At-NF-YA2 protein interacts with *At-HSFA3a* and *At-HSFA7b* promoters, but not with *At-HSP17.ACI*, *At-HSFA7a* and *At-APX2* promoter (Figure 6B, Supplementary figure S20). To further confirm the repression of *At-HSFA3a* and *At-HSFA7b* transcription by At-NF-YA2 *in planta,* GUS promoter reporter constructs Pro-*At-HSFA3a:GUS* and Pro-*At-HSFA7b:GUS* were introduced in WT, *At-NF-YA2*-OE and *At-nfy-a2* mutant backgrounds. Expression analysis and quantification of *GUS* transcription using these lines confirm At-NF-YA2-mediated repression of *At-HSFA3a* and *At-HSFA7b* as *GUS* accumulation is reduced in *At-NF-YA2*-OE line and strongly expressed in *At-nf-ya2*

mutant background as compared to WT (Figure 6C, D). Thus, At-NF-YA2 regulates the

transcription of *At-HSFA3a* and *At-HSFA7b* directly. Thermotolerance assay using *At-*

hsfa3a and *At-hsfa7a* mutants show high sensitivity to HS (Figure 6E, Supplementary

figure S21A, B), only 5% *At-hsfa3a* and 1.7% *At-hsfa7a* plants are able to recover as

compared to 45% WT plants six-days-after HS imposition (Figure 6F, Supplementary

figure S21B). This confirms the positive role of *At-HSFA3a* and *At-HSFA7a* genes in

maintaining thermotolerance. Furthermore, role of *At-HSFA7b* gene in salt and heat

stress tolerance via an E-box-like motif was recently reported by Zang et al. (2019).

Thus, in Arabidopsis, HS induces At-HSFA2 that increases the expression of At-

miR169d which in turn post-transcriptionally down-regulates *At-NF-YA2* transcriptional

repressor. At-NF-YA2 in turn enhances transcription of HSR genes like *At-HSFA3a* and

At-HSFA7b. This regulatory feedback loop plays a critical role in thermotolerance.

Discussion

Previously we have shown heat-responsiveness of *MIR169* members in tomato (Rao et

al. 2020). Moreover, heat-mediated induction of miR169 in Arabidopsis (Guan et al.

2013; Li et al. 2010) and in rice (Zhao et al. 2007; Zhao et al. 2009) is also reported.

This study focuses on defining the underlying upstream regulatory mechanism viz., the

HSF-mediated transcriptional control of *MIR169* genes as well as the post-

transcriptional regulation of downstream targets of miR169s i.e., the *NF-YA* transcription

factors, which in turn transcriptionally regulate HSR genes in both tomato and

Arabidopsis.

Cooperative HSFs binding regulate *MIR169* transcription in HS

At-HSFA1b and At-HSFA7b mediated induction of *At-MIR398b* during heat stress has

been reported earlier in Arabidopsis (Guan et al. 2013). Presence of multiple HSEs in

promoters foster cooperative interactions between multiple HSF trimers (Topol et al.

1985; Xiao et al. 1991; Bonner et al. 1994). The promoters of HS-responsive *MIR169s*

exhibit multiple HSFs binding making these promoters suitable targets for cooperative

binding of HSFs as evidenced by Y1H/ChIP-qPCR assays in both tomato and

Arabidopsis (Figures 1A-I, 4B and 6B). The binding ability of Sly-HSFs on *Sly-MIR169*

promoters in tomato was further confirmed by transient assays in *N. benthamiana*,

where effector plasmids containing different Sly-HSFs were co-transformed with specific *Sly-MIR169* promoters hooked with *GUS* reporter gene. Sly-HSF-mediated activation of *Sly-MIR169a-1*, *Sly-MIR169b*, *Sly-MIR169c*, *Sly-MIR169d* and *Sly-MIR169d-1* promoters was confirmed for all Sly-HSFs except Sly-HSFA1a. Individual co-transfection of Sly-HSFA1a with Pro-*Sly-MIR169b* was unable to induce *GUS* transcription. Co-infiltration of Sly-HSFA1a with Sly-HSFA7a (the other HSF binding on Pro-*Sly-MIR169b*) leads to strong induction of *GUS* transcripts, suggesting the cooperative interactions between Sly-HSFA1a and Sly-HSFA7a. The *Sly-HSFA1a*-VIGS study suggests that threshold level Sly-HSFA1a is sufficient to regulate transcription of its target genes, as portrayed by strong reduction of *Sly-MIR169b* transcription (Figure 2D). Some studies have also reported the stress-mediated hetero-oligomeric complex formation of HSFA1a with other A-class HSFs to synergistically regulate the expression of a number of small HSPs, DnaJ/Hsp40, and HSP70 (Chan-Schaminet et al. 2009; Liu et al. 2011; Hahn et al. 2011; Li et al. 2014).

The HSF 'B-class' is known primarily as a repressor of transcription (Czarnecka-verner et al. 2000; Ikeda et al. 2009). We find Sly-HSFB1a-mediated transactivation of Pro-*Sly-MIR169d* (Supplementary figure S3D). Sly-HSFB1a has been shown to activate transcription of Myc-HSP17.6, alone and synergistically with Sly-HSFA1a as well as with acidic activators like HAC1/CBP (Bharti et al. 2004). The class-A1 HSFs are considered as master regulators of HS response (Liu et al. 2011; Yoshida et al. 2011). In our study we find 'Class-A' HSFs regulate expression of all *MIR169* either singularly or in combination with other HSFs belonging to class-A/-B/-C in both tomato and Arabidopsis. The 'Class-A1' HSFs regulate 2 of 5 and 1 of 3 *MIR169* promoters in tomato and Arabidopsis, respectively reiterating the significance of this class in HS tolerance.

Alleviation of *Sly-NF-YA9*/*Sly-NF-YA10* and *At-NF-YA2* enhances thermotolerance

Further, analysis of thermotolerance of plants with reduced *NF-YA* levels (TRV-*Sly-NF-YA9* and TRV-*Sly-NF-YA10* in tomato and *At-MIR169d*-OE and *At-nf-ya2* knockout mutant in Arabidopsis) and those with higher *NF-YA* expression (At-*NF-YA2*-OX, p*NF-YA2*:*NF-YA2*-r and *MIM169defg*) revealed that reduced *NF-YA* levels increase

thermotolerance (higher survival rate) and there is elevated expression of HSR genes

including *HSF* genes, APX and several HSP genes (Figure 3D-H, Supplementary figure

S9 and Figure 6C, Supplementary figure S19). Thus, establishing *Sly-NF-YA9 and Sly-*

NF-Y10 as negative regulators of HSR gene expression and thermotolerance in tomato

and *At-NF-YA2* in Arabidopsis. Other reports by Ceribelli et al. (2008) and

Leyva-González et al. (2012) have also shown NF-YAs as repressors. Further, we have

demonstrated in both tomato and Arabidopsis that the diminished expression of *Sly-NF-*

YA10 (Rao et al. 2020) and *At-NF-YA2* (Figure 2C, Supplementary figure S13A, B and

supplementary figure S15A, B) upon HS is a post-transcriptional regulation which is a

result of miR169-mediated cleavage and not a transcriptional regulation.

At-NF-YA2 binds to HSR gene promoters

Our study establishes direct At-NF-YA2 mediated transcriptional regulation of *At-*

HSFA3a and *At-HSFA7b* by Y1H/ChIP-qPCR assays. Further, loss of Pro-

HSFA3a:*GUS*, Pro-*HSFA7a*:*GUS* and Pro-*HSFA7b:GUS* activity in *At-NF-YA2*-OE lines

and high *GUS* expression in *At-nf-ya2* mutant background confirms regulation of these

promoters by At-NF-YA2 as a transcriptional repressor (Figure 6C and D). Studies in

literature support the involvement of *HSFA3a* in governing thermotolerance in

Arabidopsis and tomato (Li et al. 2013; Wu et al. 2018). Larkindale and Vierling (2008)

have reported *At-HSFA3a* as a universal candidate gene commonly induced in heat

acclimated plants in different regimes of heat stress treatment. We also observe high

thermo-sensitivity in *At-hsfa3a* mutant plants (Figure 6E). It is known that the NF-YA

subunit forms functional trimer with NF-YB and NF-YC subunits to regulate transcription

of a large number of genes (Warpeha et al. 2007; Kumimoto et al. 2010; Sato et al.

2014). Sato et al. (2014) have shown synergistic interaction of At-DREB2A with trimer of

NF-YA2, NF-YB3 and NF-YC10 to form a transcriptional complex that activates *At-*

HSFA3 expression in Arabidopsis protoplasts. *DREB2A* is also a HS inducible gene and

its overexpression significantly increases HS tolerance (Sakuma et al. 2006). Moreover,

Sato et al. (2014) have found heat-inducible expression of *NF-YB3* and *NF-YC10* but

relatively stable expression of *At-NF-YA2* transcripts upon HS. Here, we conclusively

show that there is heat-mediated down-regulation of *At-NF-YA2* (Supplementary figure

S13A, B) that is a consequence of *miR169*-mediated cleavage and also that At-NF-YA2

acts as a repressor of *At-HSFA3a* transcription (Figures 5C, 6C and 6D). Thus, we

speculate that the reduced availability of NF-YA2 subunits during HS (due to

degradation by miR169d) may favor functional DREB2A:NF-YC-10:NF-YB3 complex

formation, which in turn up-regulates *At-HSFA3a* expression during HS (Figure 7).

An HSF-mediated feedback loop regulates *MIR169* in heat stress

Furthermore, under HS, *Sly-HSFA7a is* up-regulated in *Sly-NF-YA9/A10* silenced

tomato plants and there is enhanced expression of *At-HSFA7b* in *At-nf-ya2* mutant

plants (Figures 3D-H, 5C, and supplementary figure S19 and table S8). We have shown

that these HSFs interact with *MIR169* promoters. This suggests the existence of an

HSF-mediated feedback module regulating HS-mediated transcriptional regulation of

MIR169s directly or indirectly that may be crucial for thermotolerance in both tomato

and Arabidopsis (Figures 1A, 3D-H, 4B, 5C and 7). While our study demonstrates a

direct role of NF-YA class to regulate HSFs, one cannot rule out the possibility that NF-

YAs regulate HSFs indirectly. It is known that *NF-YA* genes regulate abiotic/biotic

stress response mainly via ABA pathway (Zhao et al. 2009; Luan et al. 2015; Ding et

al. 2016). Similarly, HSFs are also induced in response to ABA in Arabidopsis (Huang

et al. 2016). Thus, alteration in HSFs levels in *Sly-NF-YA9/10* down-regulated tomato

plants or *At-nf-ya2* knockdown Arabidopsis plants could also be a result of the

fluctuations in the ABA levels of the silenced plants. Moreover, the HSF perturbation

during HS could also be a result of ROS-mediated HSF induction (Volkov et al. 2006;

Guan et al. 2013), thus feeding to the HSF:*MIR169*:*NF-YA* loop. We find that heat

stress induces At-HSFA2a that in turn transcriptionally activates *At-MIR169d*. At-

HSFA2a has also been shown to be positively regulated by NF-YC10 together with

HSFA1s (Yoshida et al. 2011; Sato et al. 2014). Overexpression of miR156 and the

resultant down-regulation of target SPL genes also induce HSR genes including *At-*

HSFA2a (Stief et al. 2014) (Figure 7). Moreover, our data shows other heat responsive

miR169s may also act parallely to regulate NF-YAs and contribute towards heat

tolerance of Arabidopsis (Figure 7). Thus, multiple pathways work in tandem, interacting

among themselves to regulate cellular homeostasis during HS.

A conserved thermotolerance mechanism in tomato and Arabidopsis

It is also noteworthy, that tomato Sly-NF-YA9/A10 are the best orthologues of At-NF-YA2 (Supplementary figure S22). Also, we show that Sly-HSFA7a and At-HSFA7b regulate the transcription of *MIR169* in both tomato and Arabidopsis, respectively. Interestingly, the heat-responsive *At-MIR398* is also shown to be regulated by an HSFA7 class member (Guan et al. 2013). This suggests that the Sly-NF-YA9/10 and At-NF-YA2 orthologues and Sly-HSFA7-class members have conserved roles in heat tolerance in both plant species.

Our study also highlights the existence of HSF-independent pathways in regulating *MIR169s* transcription during HSR as HSF-mediated regulation could be established for only 5 out of 15 and 3 out of 11 *MIR169* promoters of tomato and Arabidopsis, respectively (Figures 1A, 4B). Though these *MIR169* loci may be indirectly controlled by other cascades regulated by HSFs like the DREB2-mediated *HSFA3* regulation or NF-YA-mediated regulation as we have shown. The present study establishes a conserved strategy that involves HSF-mediated induction of *miR169* in both tomato and Arabidopsis that leads to reduction of target *NF-YA* genes that in-turn regulate HSF transcription which feeds back to the miR169:*NF-YA* pathway. Down-regulation of *At-NF-YA2* homologues may be a viable strategy for improving the thermotolerance and yield stability of crop plants.

Materials and Methods

Plant materials and growth conditions

Tomato cultivar CLN seeds (Balyan et al. 2020) were germinated on filter paper soaked with deionized water at 26 ºC. Three-days-old seedlings of uniform growth were transplanted in plastic pots, filled with soilrite and placed in a plant growth chamber (CMP6050, Conviron, Canada) maintained at 26 ºC/21 ºC (day/night: 16/8 h) relative humidity 60%, light intensity 300 µM per m^2 per sec.

Arabidopsis thaliana (ecotype Columbia) was used as the wild-type in this study. Seeds of the *MIR169d*-OE, *NF-YA2*-OE, Pro-*NF-YA2:NF-YA2*-r, Pro-*At-MIR169d: GUS*, Pro-*At-NF-YA2: GUS* were from Sorin et al. (2014) study. Seeds of *At-nf-ya2*

(SALK_021228) were obtained from the Arabidopsis Biological Resource Center (Columbus, OH). Homozygous plants were identified by diagnostic PCR analysis using the primers listed in Table S6. Arabidopsis plants were grown in soilrite containing plastic pots and kept under cool white light of 100 µmol m−2 sec−1 intensity, with a 16 h light/8 h dark photoperiod at 22 °C/20 °C (day/night) in a plant growth chamber (CMP6050, Conviron, Canada).

One-month old tomato plants were initially exposed for 4 h at a gradual increasing range of temperature from 26 ºC to 45 ºC then exposed additionally for 4.5 h at 45 ºC for HSFs and miR169s expression profiling study. VIGS silenced tomato plants were exposed to heat stress for 4.5 h at 45 ºC while 2-weeks old *Arabidopsis* plants were exposed to 42 ☐C for 2 h heat stress. For phenotype scoring and survival assays, HS treated VIGS silenced plants and *Arabidopsis* plants were allowed to recover for six days at 26 ºC/21 ºC (day/night: 16/8 h) and at 22 °C/20 °C (day/night) in a plant growth chamber (CMP6050, Conviron, Canada). Leaves from these experimental plants were harvested immediately after HS for expression profiling of HSR genes.

Real-time quantitative RT–PCR analysis

Four (expression studies of *HSFs* and miR169s) and five weeks (VIGS silenced) old tomato plants and two-weeks-old seedlings of Arabidopsis were used for total RNA extraction with Trizol reagent (Invitrogen, USA). Total RNA was treated with DNase I (Ambion, USA) to remove genomic DNA contamination. For real-time RT–PCR (qRT–PCR) analysis, 2 µg of total RNA was used for cDNA synthesis by using high capacity cDNA reverse transcription kit (Applied Biosystem, USA) following the manufacturer's protocol. The cDNA reaction mixture was diluted 2- fold, a 0.5 µl aliquot was used as a template in a 10 µl PCR reaction. PCR reactions included a pre-incubation at 95°C for 2 min, followed by 40 cycles of denaturation at 95°C for 15 sec, annealing at 60°C for 40 sec, and extension at 72°C for 45 sec. All the reactions were performed in a CFX-Connect real-time PCR detection system (Bio–Rad, www.bio-rad.com/) using GO Taq qPCR master mix (Promega, USA). Each experiment was repeated at least three times. The relative fold change was calculated by following the $2^{-\Delta\Delta CT}$ method using *Actin* and *Tubulin* gene as endogenous control (Paul et al. 2016). The stability and similarity in the

Ct range of *Actin* and *Tubulin* were assessed across the experimental conditions

(Supplementary table S9). For calculation of fold change in the gene expression the

control value was set as one for all qRT-PCR expression analysis. The primers used in

this study are listed in Supplementary table S10 and all the qRT-PCR primers having a

good amplification efficiency ranging between 90-110% were used (Supplementary

table S11).

For miRNA expression analysis 2 µg of the small RNA was polyadenylated using the

Poly(A) Tailing kit (epicenter- An Illumina company) following the manufacturer's

protocol. The above poly-A tailed small RNA reaction mixture was then reverse

transcribed with a special 100 bp primer i.e. miR_oligodT_RTQ (Mutum et al. 2013)

which is composed of two variable bases at its end followed by the poly(T)25 and then

with a random adapter sequence using the Superscript III Reverse Transcriptase

(Invitrogen, USA). The taqman based q-RT-PCR was performed using the TaqMan Fast

Universal PCR Master Mix (2X, Applied Biosystems) and 'Fam (Fluorescein) dye' and

BHQ labeled taqman universal probe specific to the adapter region of the

miR_oligodT_RTQ (Supplementary table S8) which enable the use of a common

reverse primer (Mutum et al. 2013) and miRNA specific forward primer in the expression

analysis. For each well a total reaction volume of 7 µl was prepared and for each

miRNA two technical with 3 biological replicates were analyzed, *5S rRNA* and

U6snRNA (Supplementary table S8) were used as the endogenous control for data

normalization. For calculation of fold change in the gene expression the control value

was set as one for all qRT-PCR expression analysis.

Promoter sequences analysis

The 2 kb sequences, upstream of the precursor start site of *MIR169* precursors and

translational start codon (ATG) of Arabidopsis HSF genes were downloaded from Sol

Genomic Network database (http://sgn.cornell.edu) and TAIR, respectively and

analyzed for the presence of putative *cis*-regulatory elements by PlantPAN 3.0 (Chow et

al. 2019). Positions of heat stress elements (HSE) were also searched manually in

miRNA169 promoters. We searched for both perfect and imperfect HSE types manually,

as these imperfect variants also have been validated for HSFs binding by different in
vitro and in vivo assays (Santoro et al. 1998; Mittal et al. 2011; Guo et al. 2008).

Yeast one-hybrid assays

The coding sequences for 23 tomato HSFs and 21 Arabidopsis HSFs were cloned into
the pGADT7 vector (Clontech) as a prey (Supplementary table S8). DNA fragments
corresponding to the promoters (250 to 1500 bp) of heat responsive *MIR169* genes
from tomato and Arabidopsis (Supplemental tables S7, S3) were cloned into the pABAI
plasmid (Clontech) as baits. For positive control, sHSP promoter previously reported to
have HSFA3a binding (Li et al. 2013) was cloned. Promoter-specific bait yeast strains
were generated by transforming Y1HGold yeast strain with 1 μg of linearized bait
plasmids and plated on SD-ura medium. Positive bait strains having insertion at the
URA3 locus were selected by performing colony PCR using primers specific to the bait
sequences (Supplementary table S10). To negate this bait specific background *AbAr*
expression, minimal inhibitory concentrations for all 27 baits (miRNA promoters) were
determined by spotting yeast cultures at OD_{600} ranging from 0.1 to 0.0001on SD/-Ura
medium supplemented with different concentration of aureobasidin A (100- 1000 ng/ml;
Supplementary tables S6, S12). For one-on-one bait/ prey interaction, bait strains were
transformed with pGADT7 vector containing CDS of tomato HSFs. Positive bait/prey
containing yeast colonies were selected by plating the transformed yeast cells on SD/-
Leu medium. To study the interaction between different tomato HSFs and miRNA
promoters, specific bait/prey containing yeast strains were spotted on SD/-Leu/AbA
medium having bait specific aureobasidin A concentration.

Chromatin immunoprecipitation (ChIP)-qPCR assays

Promoters for tomato *Sly-MIR169a-1, Sly-MIR169b, Siy-MIR169c, Sly-MIR169d* and
SlyMIR169d-1 were amplified from the genomic DNA and cloned into pbi101 vector
upstream of the *GUS* reporter gene. All transcription factors (*Sly-HSFA1a*, *Sly-HSFA7a*,
Sly-HSFA8a, *Sly-HSFB1a*, *Sly-HSFA4a*, *Sly-HSFA6b* and *Sly-HSFA1e* CDS) were
cloned in pCAMBIA1302 in fusion with the *GFP* reporter. The binary constructs were co-
infiltrated as per the combinations obtained from Y1H assay into 4-weeks-old *N.*
benthamiana leaves via *Agrobacterium tumefaciens* (strain EHA105). These leaves

were harvested 2 days after the infiltration and ChIP assays was performed using four
infiltrated leaves from five independent plants for each co-infiltration. To identify the
binding of At-NF-YA2 on HSR gene promoters, *NF-YA2:GFP* overexpressing transgenic
lines were used. Leaves from *N. benthamiana* plants (for tomato ChIP) and 3-week-old
NF-YA2:GFP overexpressing Arabidopsis plants were cross-linked with 1%
formaldehyde. Chromatin was sheared to an average length of 500bp by sonication,
and immunoprecipitated with GFP tag-specific monoclonal antibody (abcam; catalogue
no. ab290). The fold enrichment was calculated for different promoters by performing
real-time quantitative PCR analysis on immunoprecipitated DNA using a Bio–Rad
CFX96 real-time PCR detection system. For each promoter pair used for ChIP-qPCR
the primer amplification was first tested on input DNA and the fold enrichment was
checked by keeping no-antibody, IgG and input DNA controls. Data for enrichment of
Sly-HSFA6b on Sly-MIR169d-1 promoter is presented as an example in Supplementary
figure S23. All the primers used for ChIP-qPCR are listed in Supplementary Table S10.

Transient promoter reporter assay to confirm HSFs-mediated regulation of *Sly-*
MIR169* promoters in *N. benthamiana

Agro-infiltration was performed in 4-weeks-old *N. benthamiana* leaves. For promoter
reporter assays Pro-*Sly-MIR169s*:GUS and p35S:*Sly-HSFs* were transformed in
Agrobacterium tumefaciens. For co-infiltrations, overnight cultures of individual
constructs were harvested and suspended at OD 1 in infiltration buffer (10 mM MgCl2,
10 mM MES, pH 5.8, 0.5% glucose and 150 µM acetosyringone). After 3 h of incubation
at room temperature, various combinations were mixed at a 1:1:1 ratio; these mixes
were used to infiltrate leaves. These leaves were harvested 2 days after the infiltration
and used for RNA extraction and qRT-PCR as described above. The expression of GUS
transcript is measured by qRT-PCR and presented as fold-change values calculated
using the 2–ΔΔCt method. The values were normalized with *NPTII* that is co-expressed
in the promoter: GUS construct. The GUS expression from the reporter (*pro-Sly-*
MIR169:GUS) infiltrated samples was set to one to normalize the background signal.

Virus induced gene silencing

Transient silencing assays following VIGS was performed in tomato plants as described by Senthil and Mysore (2014). For silencing of selected genes, 300 to 400 bp cDNA fragments designed using VIGS tool (Sol Genomics Network) were amplified and cloned into the *Eco* RI–*Bam* HI site of the TRV2 vector. After confirmation by Sanger sequencing, the TRV2-gene vectors were transformed into *A. tumefaciens* strain GV3101. About 15-days-old tomato plants were inoculated with a mixture of *A. tumefaciens* strains containing the TRV2 vector constructs and TRV1 (a helper plasmid, contains RNA-dependent RNA polymerase (RdRp), movement protein (MP), and a 16 kDa cysteine-rich protein) into the first and second true leaf. Empty TRV2 vector was used as a control. Infected plants were kept in dark and moist conditions for 3 days and then transferred to light; HS was imposed after 3 weeks (the time required for appearance of chlorophyll bleaching phenotype in the leaves of TRV-*Phytoene desaturase*-infiltrated plants used as silencing control) (Supplementary figure S24A, B) at 45 °C for 4.5 h. Leaves were frozen for qRT-PCR analysis. Off targets of the selected VIGS constructs were also predicted at VIGS tool (Supplementary table S4) and validated by qRT-PCR in the silenced plants. To confirm the maintenance of silencing throughout the phenotype scoring and biochemical assays time period, the expression of *Sly-NF-YA9* and *Sly-NF-YA10* was assessed further during the post heat stress period. Survival and cell death was assayed after 6 days' recovery. The experiment was performed in four biological replicates with 8 plants per replicate for each gene.

Gas-exchange parameters measurements

Gas-exchange parameters, like water use efficiency (WUEi), net photosynthetic rate (A), and transpiration rate (E) were measured simultaneously by using a portable Licor 6400 photosynthesis system (LI-6400, Li-Cor Inc., Lincoln NE, USA). WUEi was calculated using the formula A/E. For all measurements, sixth and seventh fully expanded tomato leaves of 6 plants for each VIGS construct were used and the experiment was repeated thrice keeping similar parameters. The measurement conditions were as follows: leaf temperature 27°C, leaf-air vapour pressure deficit 1.5±0.5 kPa, photosynthetic photon flux 300 µmol m^{-2} s^{-1}, relative air humidity 60% and ambient CO2 concentration 400±5 µmolmol^{-1}.

Histochemical detection of H_2O_2 accumulation and cell death in HS

Hydrogen peroxide (H_2O_2) levels and cell death were estimated by using 3,3'-
diaminobenzidine (DAB) and trypan blue stain, respectively on the 6th and 7th leaves of
6-week-old TRV-Vector and TRV-gene plants immediately after heat stress (45 °C for 6
h). DAB staining was performed as previously described (Daudi et al. 2012). The
viability of leaf cells was measured using the trypan blue exclusion method (Koch and
Slusarenko 1990). Experiment was repeated twice with similar parameters having a
sample size of 8 plants per replicate.

GUS assays

To analyze the histochemical localization of GUS reporter protein, tissue samples were
incubated for 12 h at 37 °C with the substrate solution (1 mM 5-bromo-4-chloro-3-
indolyl-β-D-glucuronide, pH 7.0, 100 mM sodium phosphate buffer, 10 mM Na2EDTA,
0.5 mM potassium ferricyanide, 0.5 mM potassium ferrocyanide, and 0.1% Triton X-
100). Subsequently, seedlings and leaves were washed with de-staining solution
containing ethanol:acetone:glycerol (3:1:1) to eliminate chlorophyll. They were
photographed with a Nikon SMZ1000 Stereomicroscope (Tokyo, Japan) and OLYMPUS
SZX16 Stereomicroscope (India). *GUS* transcripts were quantified using qRT-PCR. The
assays were performed with three biological replicates.

Constructs generation

For yeast one hybrid assay, all the promoters and CDS of transcription factors (HSFs
and NF-YA) were cloned in pABAI and PGADT7 vector, respectively. For promoter
reporter assays, all promoters were cloned in pbi101 vector and all transcription factors
(*Sly-HSFA1a*, *Sly-HSFA7a*, *Sly-HSFA8a*, *Sly-HSFB1a*, *Sly-HSFA4a*, *Sly-HSFA6b* and
Sly-HSFA1e CDS) were cloned in pCAMBIA1302. Promoters were amplified as
approximately 2000 bp region upstream of the precursor start site for *MIR169s* (Pro-*Sly-*
MIR169a-1:*GUS*, Pro-*Sly-MIR169b:GUS*, Pro-*Sly-MIR169c:GUS*, Pro-*Sly-*
MIR169d:GUS, Pro-*Sly-MIR169d-1:GUS*, Pro-*At-MIR169d:GUS)* and translation start
site for HSFs (Pro-*At-HSFA3a:GUS* and Pro-*At-HSFA7b:GUS) and* Pro-*NF-YA2:GUS*
from genomic DNA. For overexpressing miR169d (*p35S:At-MIR169d-OE)* a 300 bp
fragment surrounding the miRNA sequence was amplified from genomic DNA, cloned in
pbi121 and transformed into Arabidopsis WT plants by floral dip method. For VIGS,

300-400 bp fragment of *Sly-HSFA1a*, *Sly-HSFA7a*, *Sly-HSFB1a*, *Sly-HSFA6b* and *Sly-*
NF-YA9, *Sly-NF-YA10* were cloned in pTRV-2 vector.

Acknowledgments

This work is supported by grants from DBT-NIPGR, India and grants from DST-SERB,
India. The authors acknowledge phytotron facility, CIF and field area provided by
NIPGR. The authors are thankful to DBT-eLibrary Consortium (DeLCON) for providing
access to e-resources. CLN1621L and CA4 seeds were procured from AVRDC,
Taiwan. SR acknowledges Department of Biotechnology (DBT) Govt. of India and CB
acknowledge University Grants Commission (UGC) for the award of research
fellowships.

Author contributions

695   SM designed and supervised the study; SR designed and performed all the experiments
696   and analyzed the data; CB did mutant genotyping for hsfa2 and nf-ya2; CS generated
697   MIR169, NF-YA2 and NF-YA2-r transgenic lines of Arabidopsis. SR and SM wrote the
698   article; CS and MC complemented the writing and critically reviewed the manuscript.
699   SM agrees to serve as the author responsible for contact and ensures communication.

References:

Balyan S, Rao S, Jha S, Bansal C, Das JR and Mathur S (2020). Characterization of
novel regulators for heat stress tolerance in tomato from Indian sub-continent. Plant
Biotechnology Journal, doi:10.1111/pbi.13371.

Bäurle I (2016) Plant heat adaptation: priming in response to heat stress.
F1000Research, 5.

Bharti K, Von Koskull-Döring P, Bharti S, Kumar P, Tintschl-Körbitzer A, Treuter E, &
Nover L (2004) Tomato heat stress transcription factor HsfB1 represents a novel type of
general transcription coactivator with a histone-like motif interacting with the plant CREB
binding protein ortholog HAC1. The Plant cell, 16(6):1521-1535.

Bonner J J, Ballou C, Fackenthal D L (1994) Interactions between DNA-bound trimers
of the yeast heat shock factor. Molecular Cell Bioogyl,14:501-508.

Buhtz A, Pieritz J, Springer F, Kehr J (2010) Phloem small RNAs, nutrient stress
responses, and systemic mobility. BMC Plant Biology, 10(1), 64.

Busch W, Wunderlich M, Schöffl F (2005) Identification of novel heat shock
factor-dependent genes and biochemical pathways in Arabidopsis thaliana. The Plant
Journal, 41:1-14.

Ceribelli M, Dolfini D, Merico D, Gatta R, Viganò AM, Pavesi G, Mantovani R (2008)
The histone-like NF-Y is a bifunctional transcription factor. Molecular and cellular
biology, 28:2047-2058.

Chan-Schaminet KY, Baniwal SK, Bublak D, Nover L and Scharf KD (2009) Specific
interaction between tomato HsfA1 and HsfA2 creates hetero-oligomeric superactivator
complexes for synergistic activation of heat stress gene expression. Journal of
Biological Chemistry, 284:20848-20857.

Charng YY, Liu HC, Liu NY, Chi WT, Wang CN, Chang SH, Wang TT (2007) A heat-
inducible transcription factor, HsfA2, is required for extension of acquired
thermotolerance in Arabidopsis. Plant physiology, 143(1):251-262.

Chow CN, Lee TY, Hung YC, Li GZ, Tseng KC, Liu YH, Kuo PL, Zheng HQ, Chang WC
(2019) PlantPAN3. 0: a new and updated resource for reconstructing transcriptional
regulatory networks from ChIP-seq experiments in plants. Nucleic acids research,
47(D1): D1155-63.

Collins GG, Nie X, Saltveit ME (1995) Heat shock proteins and chilling sensitivity of
mung bean hypocotyls. Journal of Experimental Botany, 46:795-802.

Czarnecka-Verner E, Yuan CX, Scharf KD, Englich G, Gurley WB (2000) Plants contain
a novel multi-member class of heat shock factors without transcriptional activator
potential. Plant Molecular Biology. 43:459-471.

Daudi A, O'Brien JA (2012) Detection of Hydrogen Peroxide by DAB Staining in
Arabidopsis Leaves. Bio Protocol, 2(18),1-4.

Ding Q, Zeng J, He XQ (2016) MiR169 and its target PagHAP2-6 regulated by ABA are
involved in poplar cambium dormancy. Journal of plant physiology, 198, 1-9.

Guan Q, Lu X, Zeng H, Zhang Y, Zhu J (2013) Heat stress induction of miR398 triggers
a regulatory loop that is critical for thermotolerance in Arabidopsis. The Plant Journal,
74:840-851.

Guo L, Chen S, Liu K, Liu Y, Ni L, Zhang K and Zhang L (2008) Isolation of heat shock
factor HsfA1a-binding sites in vivo revealed variations of heat shock elements in
Arabidopsis thaliana. Plant and Cell Physiology, 49(9), 1306-1315.

Guo M, Liu JH, Ma X, Luo DX, Gong ZH, Lu MH (2016) The plant heat stress
transcription sactors (HSFs): structure, regulation, and function in response to abiotic
stresses. Frontiers in Plant Science, 7:114.

Hahn A, Bublak D, Schleiff E, Scharf KD (2011) Crosstalk between Hsp90 and Hsp70
Chaperones and Heat Stress Transcription Factors in Tomato. Plant Cell, 23:741-55.

Hasanuzzaman M, Nahar K, Alam MM, Roychowdhury R, Fujita M (2013) Physiological,
biochemical, and molecular mechanisms of heat stress tolerance in plants. International
Journal of Molecular Sciences, 14:9643-9684.

Hivrale V, Zheng Y, Puli C, Jagadeeswaran G, Gowdu K, Kakani V, Barakat A, Sunkar
R (2015). Characterization of drought- and heat-responsive microRNAs in switchgrass.
Plant Science, 242, 10:1016.

Huang YC, Niu CY, Yang CR, Jinn TL (2016) The heat stress factor HSFA6b connects
ABA signaling and ABA-mediated heat responses. Plant Physiology, 172, 1182-1199.

Ikeda M, Ohme-Takagi M (2009) A novel group of transcriptional repressors in
Arabidopsis. Plant and Cell Physiology 50:970-975.

Jacob P, Hirt H, Bendahmane A (2017) The heat-shock protein/chaperone network and
multiple stress resistance. Plant Biotechnology Journal, 15:405-414.

Koch E, Slusarenko A (1990) Arabidopsis is susceptible to infection by a downy mildew
fungus. The Plant Cell, 1;2(5):437-45.

Kotak S, Port M, Ganguli A, Bicker F and Von Koskull-Döring P (2004) Characterization of C-terminal domains of Arabidopsis heat stress transcription factors (Hsfs) and identification of a new signature combination of plant class A Hsfs with AHA and NES motifs essential for activator function and intracellular localization. The Plant Journal, 39(1), 98-112.

Kumimoto RW, Zhang Y, Siefers N, Holt III BF (2010) NF−YC3, NF−YC4 and NF−YC9 are required for CONSTANS-mediated, photoperiod-dependent flowering in Arabidopsis thaliana. The Plant Journal, 63(3), 379-391.

Larkindale J, Vierling E (2008) Core genome responses involved in acclimation to high temperature. Plant Physiol, 146:748-761.

Lee H, Yoo SJ, Lee JH, Kim W, Yoo SK, Fitzgerald H, Carrington JC, Ahn JH (2010) Genetic framework for flowering-time regulation by ambient temperature-responsive miRNAs in Arabidopsis. Nucleic acids research, 38: 3081-3093.

Leyva-Gonzalez MA, Ibarra-Laclette E, Cruz-Ramoırez A, Herrera-Estrella L (2012) Functional and transcriptome analysis reveals an acclimatization strategy for abiotic stress tolerance mediated by Arabidopsis NF-YA family members. PLoS ONE, 7: e48138.

Li S, Liu J, Liu, Z, Li X, Wu F, and He Y (2014) HEAT-INDUCED TAS1 TARGET1 Mediates Thermotolerance via HEAT STRESS TRANSCRIPTION FACTOR A1a-Directed Pathways in Arabidopsis. The Plant Cell, 26:1764-1780.

Li WX, Oono Y, Zhu J, He XJ, Wu JM, Iida K, Lu XY, Cui X, Jin H, Zhu JK (2008) The Arabidopsis NFYA5 transcription factor is regulated transcriptionally and post-transcriptionally to promote drought resistance. Plant Cell, 20(8):2238-2251.

Li Y, Fu Y, Ji L, Wu C-A, Zheng C (2010) Characterization and expression analysis of the Arabidopsis mir169 family. Plant Science, 178:271-280.

Li Y, Zhao SL, Li JL, Hu XH, Wang H, Cao XL, Xu YJ, Zhao ZX, Xiao ZY, Yang N, Fan J, Huang F, Wang WM (2017) Osa-miR169 negatively regulates rice immunity against the blast fungus Magnaportheoryzae. Frontiers in Plant Science, 8,2.

Li Z, Zhang L, Wang A, Xu X, Li J (2013) Ectopic overexpression of SlHsfA3, a heat
stress transcription factor from tomato, confers increased thermotolerance and salt
hypersensitivity in germination in transgenic Arabidopsis. PloS one, 8(1), e54880.

Lin JS, Kuo CC, Yang IC, Tsai WA, Shen YH, Lin CC, Liang YC, Li YC, Kuo YW, King
YC, Lai HM, Jeng ST (2018) MicroRNA160 Modulates Plant Development and Heat
Shock Protein Gene Expression to Mediate Heat Tolerance in Arabidopsis. Frontiers in
Plant Science,9:68.

Liu HC, Liao HT, Charng YY (2011) The role of class A1 heat shock factors (HSFA1s) in
response to heat and other stresses in Arabidopsis. Plant Cell and Environment,
34:738-51.

Lobell DB, Schlenker W, Costa-Roberts J (2011) Climate trends and global crop
production since 1980. Science, 333:616-620.

Luan M, Xu M, Lu Y, Zhang L, Fan Y, and Wang L (2015) Expression of zma-miR169
miRNAs and their target ZmNF-YA genes in response to abiotic stress in maize leaves.
Gene, 555:178-185.

Mittal D, Enoki Y, Lavania D, Singh A, Sakurai H and Grover A (2011) Binding affinities
and interactions among different heat shock element types and heat shock factors in
rice (Oryza sativa L.). The FEBS journal, 278(17), 3076-3085.

Mittler R, and Blumwald E (2010) Genetic engineering for modern agriculture:
challenges and perspectives. Annual review of plant biology, 61:443-462.

Morimoto RI (2002) Dynamic remodeling of transcription complexes by molecular
chaperones. Cell 110:281-284.

Mutum RD, Balyan SC, Kansal S, Agarwal P, Kumar S, Kumar M and Raghuvanshi S
(2013). Evolution of variety-specific regulatory schema for expression of osa-miR408 in
indica rice varieties under drought stress. The FEBS journal, 280(7),1717-1730.

Ni Z, Hu Z, Jiang Q, and Zhang H (2013) GmNFYA3, a target gene of miR169, is a
positive regulator of plant tolerance to drought stress. Plant Molecular Biology, 82:113-
129.

Niu C, Li H, Jiang L, Yan M, Li C, Geng D, Xie Y, Yan Y,Shen X, Chen P, Dong J, Ma F,
Guan Q (2019) Genome-wide identification of drought-responsive microRNAs in two
sets of Malus from interspecific hybrid progenies. Horticulture Research, 75:6.

Nover L, Bharti K, Döring P, Mishra SK, Ganguli A, Scharf KD (2001) Arabidopsis and
the heat stress transcription factor world: how many heat stress transcription factors do
we need? Cell Stress Chaperons, 6:177-189.

Pant BD, Musialak-Lange M, Nuc P, May P, Buhtz A, Kehr J, Walther D, Scheible WR
(2009) Identification of nutrient-responsive Arabidopsis and rapeseed microRNAs by
comprehensive real-time polymerase chain reaction profiling and small RNA
sequencing. Plant physiology, 150:1541-1555.

Paul A, Rao S and Mathur S (2016). The α-crystallin domain containing genes:
identification, phylogeny and expression profiling in abiotic stress, phytohormone
response and development in tomato (Solanum lycopersicum). Frontiers in plant
science, 7, 426.

Rao S, Balyan S, Jha S and Mathur S (2020) Novel insights into expansion and
functional diversification of MIR169 family in tomato. Planta, 251:55.

Ravichandran S, Ragupathy R, Edwards T, Michael D, Sylvie C (2019) MicroRNA-
guided regulation of heat stress response in wheat. BMC Genomics, 20:488.

Sakuma Y, Maruyama K, Qin F, Osakabe Y, Shinozaki K, Yamaguchi-Shinozaki K
(2006) Dual function of an Arabidopsis transcription factor DREB2A in
water-stress-responsive and heat-stress-responsive gene expression. Proceedings of
the National Academy of Sciences, 103(49):18822-7.

Santoro N, Johansson N and Thiele DJ (1998) Heat shock element architecture is an
important determinant in the temperature and transactivation domain requirements for
heat shock transcription factor. Molecular and cellular biology, 18(11), 6340-6352.

Sato H, Mizoi J, Tanaka H, Maruyama K, Qin F, Osakabe Y, Morimoto K, Ohori T,
Kusakabe K, Nagata M, Shinozaki K (2014) Arabidopsis DPB3-1 a DREB2A interactor,

specifically enhances heat stress-induced gene expression by forming a heat stress-
specific transcriptional complex with NF-Y subunits. The Plant Cell, 26 4954-4973.

Scharf KD, Berberich T, Ebersberger I, Nover L (2012) The plant heat stress
transcription factor (Hsf) family: structure, function and evolution. *Biochimica et*
Biophysica Acta (BBA)-Gene Regulatory Mechanisms,1819:104-119.

Senthil-Kumar M, Mysore K (2014) Tobacco rattle virus–based virus-induced gene
silencing in *Nicotiana benthamiana*. Nature Protocols, 9:1549-1562.

Sorin C, Declerck M, Christ A, Blein T, Ma L, Lelandais-Brière C, Njo MF, Beeckman T,
Crespi M and Hartmann C (2014) A miR169 isoform regulates specific NF-YA targets
and root architecture in Arabidopsis. New Phytologist, 202:1197-1211.

Stief A, Altmann S, Hoffmann K, Pant BD, Scheible W, Bäurle I (2014) Arabidopsis
miR156 regulates tolerance to recurring environmental stress through SPL transcription
factors. The Plant Cell, 26:1792-1807.

Sunkar R, Zhu JK (2004) Novel and stress-regulated microRNAs and other small RNAs
from Arabidopsis. The Plant Cell, 16:2001- 2019.

Todesco M, Rubio-Somoza I, Paz-Ares J, Weigel D (2010) A collection of target mimics
for comprehensive analysis of microRNA function in Arabidopsis thaliana. PLoS
genetics, 6: e1001031.

Topol J, Ruden D M, Parker C S (1985) Sequences required for *in vitro* transcriptional
activation of a *Drosophila hsp 70* gene. Cell, 42:527-537.

Vidya S, Ravishankar K, Laxman R (2018) Genome wide analysis of heat responsive
microRNAs in banana during acquired thermo tolerance. Journal of Horticultural
Sciences, 13:61-71.

Volkov RA, Panchuk II, Mullineaux PM, Schöffl F (2006) Heat stress-induced H2O2 is
required for effective expression of heat shock genes in Arabidopsis. Plant Molecular
Biology, 61, 733-746.

Von Koskull-Döring P, Scharf KD, and Nover L (2007) The diversity of plant heat stress
transcription factors. Trends in Plant Science, 12:452-457.

Warpeha KM, Upadhyay S, Yeh J, Adamiak J, Hawkins SI, Lapik YR, Anderson MB,
Kaufman LS (2007) The GCR1, GPA1, PRN1, NF-Y signal chain mediates both blue
light and abscisic acid responses in Arabidopsis. Plant physiology, 143(4), 1590-1600.

Wu Z, Liang J, Wang C, Zhao X, Zhong X, Cao X, Li G, He J, Yi M (2018)
Overexpression of lily HsfA3 s in Arabidopsis confers increased thermotolerance and
salt sensitivity via alterations in proline catabolism. Journal of experimental botany,
69(8), 2005-2021.

Xiao H, Perisic O, Lis J T (1991) Cooperative binding of *Drosophila* heat shock factor to
arrays of a conserved 5 bp unit. Cell, 64:585-593.

Yan K, Liu P, Wu CA, Yang G-D, Xu R, Guo Q-H, Huang J-G, Zheng C-C (2012)
Stress-induced alternative splicing provides a mechanism for the regulation of
microRNA processing in Arabidopsis thaliana. Molecular Cell, 48:521-531.

Yang X, Zhu W, Zhang H, Liu N, Tian S (2016) Heat shock factors in tomatoes:
genome-wide identification, phylogenetic analysis and expression profiling under
development and heat stress. PeerJournal,4: e1961.

Yoshida T, Ohama N, Nakajima J, Kidokoro S, Mizoi J, Nakashima K, Maruyama K, Kim
JM, Seki M, Todaka D, Osakabe Y (2011) Arabidopsis HsfA1 transcription factors
function as the main positive regulators in heat shock-responsive gene expression.
Molecular Genetics and Genomics, 286, 321-332

Zang D, Wang J, Zhang X, Liu Z and Wang Y (2019) Arabidopsis heat shock
transcription factor HSFA7b positively mediates salt stress tolerance by binding to an E-
box-like motif to regulate gene expression. Journal of experimental botany, 70(19),
5355-5374.

Zhang B (2015) MicroRNA: a new target for improving plant tolerance to abiotic stress.
Journal of experimental botany, 66 1749-1761.

Zhang X, Zou Z, Gong P, Zhang J, Ziaf K, Li H, Xiao F, Ye Z (2011) Over-expression of
microRNA169 confers enhanced drought tolerance to tomato. Biotechnology Letters,
33(2):403-409

Zhao B, Ge L, Liang R, Li W, Ruan K, Lin H and Jin Y (2009) Members of miR-169
family are induced by high salinity and transiently inhibit the NF-YA transcription factor.
BMC Molecular Biology, 10(1): e29.

Zhao BT, Liang RQ, Ge LF, Li W, Xiao HS, Lin HX, Ruan KC and Jin YX (2007)
Identification of drought-induced microRNAs in rice. Biochemical and biophysical
research communications, 354:585-590.

Zhao M, Ding H, Zhu JK, Zhang F, Li WX (2011) Involvement of miR169 in the nitrogen-
starvation responses in Arabidopsis. New Phytologist, 190:906-1.

Zhou X, Wang G, Sutoh K, Zhu JK, Zhang W (2008) Identification of cold-inducible
microRNAs in plants by transcriptome analysis. Biochimica et Biophysica Acta (BBA)-
Gene Regulatory Mechanisms,11:780-8.

920 **Figure Legends**

921 **Main Figures:**

922 **Figure 1: Yeast one hybrid and Chromatin immunoprecipitation (ChIP-qPCR)**
923 **assay of Sly-MIR169 promoters and HSFs.** (A) Yeast-one-hybrid assay to assess
924 transcriptional regulation of heat-responsive *MIR169* promoters by HSFs. Only the

positive Y1H interactions between different HSFs and *Sly-MIR169* promoters have been shown. The *Sly-MIR169* promoters were cloned in the reporter vector pABAi and the full-length cDNA of HSFs were cloned in the pGADT7 vector. After the co-transformation into Y1H gold yeast strain, serial dilutions (O.D. 0.1 to 0.0001) of yeast cultures were spotted onto plates lacking URA and LEU with specific Aureobasidin A concentration and incubated for 3 days. Promoter of *sHSP* has been used as a positive control to show binding of HSFA3a on *sHSP* promoter (Li et al. 2013) and empty vectors as negative controls. The Aureobasidin A concentrations for all *Sly-MIR169* promoters is presented in supplementary table S7. (B-I) Chromatin immunoprecipitation (ChIP-qPCR) assay of *Sly-MIR169* promoters and HSFs in *Nicotiana benthamiana*. Fold enrichment of *Sly-MIR169a-1* promoter in HSFA1e immunoprecipitated sample (B); Fold enrichment of *Sly-MIR169b* promoter in HSFA1a immunoprecipitated sample (C); in HSFA7a immunoprecipitated sample (D); Fold enrichment of *Sly-MIR169C* promoter in HSFA4a immunoprecipitated sample (E); Fold enrichment of *Sly-MIR169C* promoter in HSFA6b immunoprecipitated sample (F); Fold enrichment of *Sly-MIR169d* promoter in HSFA8a immunoprecipitated sample (G); in HSFB1a immunoprecipitated samples (H); Fold enrichment of *Sly-MIR169d-1* promoter in HSFA6b immunoprecipitated samples (I). These experiments were repeated 3-4 times; the error bars represent standard deviation between the replicates. The values for different promoters in IgG immunoprecipitated sample were set as one for normalization.

Figure 2: Validation of HSF:miR169:*NF-YA* node components in tomato upon heat stress. (A) Expression profiles of *HSF* genes that regulate *Sly-MIR169* transcription in WT tomato leaves during control and heat stress condition by qRT-PCR using the $2-\Delta\Delta Ct$ method. (B) Expression of mature miR169s in response to heat stress as determined by taqman based qRT-PCR in control and heat challenged WT tomato leaves. (C) qRT-PCR based expression profiles of target *NF-YA* transcripts in WT control and heat stressed tomato leaves using the $2-\Delta\Delta Ct$ method. (D-G) The expression analysis of *HSF* genes, *Sly-MIR169* precursors and *NF-YA* transcripts in VIGS plants silenced for *Sly-HSFA1a* (D), *Sly-HSFA7a* (E), *Sly-HSFA6b* (F) and *Sly-HSFB1a* (G). In (D-G) 15-days-old tomato plants were agro-infiltrated with empty vector (TRV-1, TRV-2 VC) or TRV-Sly-HSF VIGS constructs. The VIGS established plants

were subjected to heat stress after 3-weeks-of agro-infiltration, and used for expression
studies of different genes. Graphical data represents mean values of expression of
three to four biological sets. Error bars show standard deviation. To represent the
negative fold-change on y axis in (C-G), the relative expression values of down-
regulated genes were transformed using the formula [-(1 / RQ ≤0.5)]. *Actin* was used as
endogenous control in (A), (C) and (D-G), while *5SrRNA* was used as reference control
in (B). The fold change normalization was done by setting the control value as one for
all qRT-PCR expression analysis.

Figure 3. Functional validation of the role of *Sly-NF-A9* and *Sly-NF-A10* in HS
response in tomato. (A) Phenotype of TRV-VC (Vector control), *TRV-Sly-NF-YA9* and
TRV-Sly-NF-YA10 silenced plants after heat stress. Fifteen-days-old tomato plants were
agro-infiltrated with empty vector (TRV-VC), *TRV-Sly-NF-YA9* or *TRV-Sly-NF-YA10*
silencing constructs. Plants were given HS 3-weeks-after infiltration. Phenotype of HS-
treated silenced plants was scored six-days-post HS. Experiments were repeated 3-4
times with 8 plants per replicate. (B) Percent survival of TRV-EV, *TRV-Sly-NF-YA9* and
TRV-Sly-NFYA10 silenced plants afters HS. (C) Estimation of water use efficiency
(mmol mol^{-1}), net photosynthesis rate (µmol m^{-2}s^{-1}), and transpiration rate of TRV-EV,
TRV-Sly-NF-YA9 and *TRV-Sly-NFYA10* silenced plants afters HS. (D-E) Expression
profiles of HSR genes in control and heat treated *TRV-EV, TRV-Sly-NF-YA9* and *TRV-*
Sly-NF-YA10 silenced plants. The expression levels of genes were calculated using the
2–ΔΔCt method and presented using fold-change values. The fold change
normalization was done by setting the control value as one in (D-H). *Actin* was used as
endogenous control. Error bars represent the standard deviation of two independent
biological replicates. *p<0.05, **p<0.01 and ***p<0.001 vs. wild type, by two-tailed
Student's t-test.

Figure 4. Transcriptional regulation of MIR169s promoters by HSFs in
Arabidopsis. (A) miR169s expression in response to heat stress as determined by
Taqman-based qRT-PCR of control and heat challenged plants. 5S rRNA was used as
the normalization control. Experiment was repeated three times and average values are
plotted as bars, error bars depict standard deviation between the replicates. (B) Y1H

assay showing the positive interaction of HSFs with the promoters of different *At-MIR169s*, Y1H assay shows binding of At-HSFA7b, At-HSFB2b, At-HSFB3a, At-HSFA2 and At-HSFA1b on *At-MIR169b*, d and h promoters. This experiment was repeated at least three times with similar results, and data from one representative experiment is shown. Serial dilutions (O.D. 0.1 to 0.0001) of yeast cultures were spotted onto plates lacking URA and LEU with specific Aureobasidin A concentration and incubated for 3 days. The Aureobasidin A concentrations for all *At-MIR169* promoters is presented in supplementary table S3. (C) Expression levels of *At-MIR169d* in wild-type plants subjected to 0 (control) and 2 h heat stress at 42 °C. Data are shown as means ± SD of three biological replicates and normalized to actin reference gene using (2–ΔΔCT) method. (D) Expression patterns of *Pro-At-MIR169d:GUS* lines of transgenic Arabidopsis plants subjected to 0 (control) or 2 h of heat stress at 42°C. (E) Levels of GUS transcripts in *Pro-At-MIR169d:GUS* lines, measured by qRT-PCR after 0 or 2 h of heat stress at 42°C. (F) Transcriptional activity of *Pro-At-MIR169d:GUS* reporter construct in WT and *At-hsfa2* mutant background during 2 h of heat stress at 42°C. (G) Expression of *GUS* transcript driven by *Pro-At-MIR169d* and *At-MIR169d* precursor in soil-grown wild-type plants of Arabidopsis and *At-hsfa2* mutant. To represent the negative fold-change on y axis in figure (G), the relative expression values of down-regulated genes were transformed using the formula [-(1 divided by RQ ≤0.5)]. The fold change normalization was done by setting the control value as one in Figure (A, C, E and G). Error bars represent the standard deviation. These experiments were repeated at least three times with similar results, and data from one representative experiment are shown in D and F.

Figure 5: Assaying the miR169d:At-NF-YA2 node in heat stress in Arabidopsis. (A) Thermotolerance was gauged in *At-MIR169d-OE*, *At-NF-YA2-OE*, *At-nf-ya2, pNF-YA2:NF-YA2*-r and MIM169defg transgenic plants in comparison to wild-type (WT) plants. Two-weeks-old soil-grown plants were subjected to 0 (control) or 2h heat stress at 42°C, and damage was recorded 6 days later. (B) Estimation of survival (percentage) of HS treated *At-MIR169d-OE*, *At-NF-YA2-OE*, *At-nf-ya2, pNF-YA2:NF-YA2*-r and MIM169defg plants. Plants were assayed for heat stress tolerance after 1 week of HS. These experiments were repeated at least five times with similar results, and average

data of all experiments are shown. Error bars, ± SD of five repeats. (C) Expression

patterns of heat stress-responsive genes in WT, *At-MIR169d-OE*, *At-NF-YA2-OE*, *At-nf-*

ya2, pNF-YA2:NF-YA2-r and MIM169defg plants. Data are shown as means ± SD of

biological replicates and normalized to *Actin* using ($2^{-\Delta\Delta CT}$) method. Log2 transformation

was applied to the RQ data to obtained negative fold change. (D-E) DAB and Trypan

blue staining of HS treated *At-MIR169d-OE*, *At-NF-YA2-OE*, *At-nf-ya2*, *pNF-YA2:NF-*

YA2-r and MIM169defg plants to estimate the production of ROS and cell death,

respectively. *$p<0.05$, **$p<0.01$ and ***$p<0.001$ vs. wild type, by two-tailed Student's t-

test. Three independent lines of *MIR169d-OE* were used and similar results were

obtained for all studied parameters, a single representative picture was shown in the

figure A, B, D and E. The fold change normalization was done by setting the control

value as one for figure (C).

Figure 6: Transcriptional regulation of *At-HSFA3a* and *At-HSFA7b* by At-NF-YA2

during heat stress in Arabidopsis. (A) Graphical representation of CCAAT cis-

elements in At-HSR gene promoters HSFA3a, HSFA7a, HSFA7b, APX and

sHSP17.6A. The 2 kb upstream sequences of start codon were analysed by plant care

and Plant PAN promoter analysis tools. The positions of CCAAT boxes are depicted by

violet coloured tabs. (B) Chromatin immunoprecipitation (ChIP)-qPCR assay of At-NF-

YA2 on At-HSR gene promoters. (C-D) *In-planta* estimation of transcriptional regulation

of *At-HSFA3a* and *At-HSFA7b* transcription by At-NF-YA2 during heat stress. (C) *Pro-*

At-HSFA3a:GUS reporter construct was agro-infiltrated in WT, *At-NF-YA2-OE* and *At-*

nf-ya2 backgrounds. *GUS* was visualized 2-days post-infiltration by histochemical

staining. (D). qRT-PCR based quantification of *GUS* transcripts in promoter reporter

constructs infiltrated plants, 2-days-post infiltration by using *GUS* specific primers. The

NPTII gene present in the same vector backbone was used as reference control. To

represent the negative fold-change on y axis in, the relative expression values of down-

regulated genes were transformed using the formula [-(1 divided by RQ ≤0.5)]. The fold

change normalization was done by setting the control value as one. (E)

Thermotolerance assay of WT and *At-hsfa3a* mutant plants. Two-weeks-old soil-grown

plants were subjected to 0 (control) or 2h heat stress at 42°C, and damage was

recorded 6 days post-stress. (F) Estimation of survival (percentage) of WT and *At-*

hsfa3a plants 6 days post heat stress. The experiments were repeated at least three
times with similar results, and average data of all experiments are shown. Error bars, ±
SD of three repeats. *p<0.05, **p<0.01 and ***p<0.001 vs. wild type, by two-tailed
Student's t-test.

Figure 7: Model for HSF-induced miR169:NF-YA module in mediating
thermotolerance in tomato and Arabidopsis. Heat stress induces the expression of
HSFs by altering the cellular ROS levels as well as other cues independent of ROS
signaling. HSFA1a, HSFA2 and HSFA7a are the key player of plant HSR response.
These HSFs bind to the promoters of *MIR169* leading to transcriptional enhancement of
miR169s in both tomato and Arabidopsis. Enhanced accumulation of miR169s reduces
the levels of *Sly-NF-YA9/A10* in tomato and *At-NF-YA2* in Arabidopsis that leads to
enhancement of HSR genes expression like *HSFA2, HSFA3* and *HSFA7s*, possibly via
altering ABA levels. In Arabidopsis, At-miR169-mediated down-regulation of *At-NF-YA2*
during HS may favor functional DREB2A:NF-YC-10:NF-YB3 complex formation to up-
regulate *At-HSFA3a* expression during HS. Enhanced levels of HSFAs in turn feed the
transcription of *MIR169* genes. The heat-mediated orchestration of parallel miRNA
pathways (miR169:NF-YA, miR398:CSD and miR156:SPL) alters the levels of HSR
genes and regulate plant thermotolerance. Solid lines denote a confirmed interaction
and dotted lines are predictive. Red and black lines denote findings of this study and
existing literature, respectively. Black colored numerals refer to the studies that
established the particular connection between different components of the model.
Reference key- 1: Guan et al. 2013; 2: Stief et al. 2014; 3: Sato et al. 2014; 4: Zhao et
al. 2009, Luan et al. 2015, Ding et al. 2016; 5: Sakuma et al. 2006. CSD: Copper
superoxide dismutase; SPL: SQUAMOSA-promoter binding like; ABA: Abscisic acid;
ROS: Reactive oxygen species; DREB: Dehydration responsive element binding
protein.

Supplementary Figure legends:

Supplementary figure S1: Promoter analysis of *MIR169* genes in tomato.
Distribution of perfect and imperfect heat stress elements (HSE) in *MIR169* promoters

of tomato. HSEs were identified manually by curating a list from published literature and marked as pink (perfect HSE) and turquoise blue (imperfect HSE) colored shapes on black colored lines representing *MIR169* promoters. The red right-handed arrow represents precursor start site. The HSE sequence variants for perfect and imperfect HSEs is provided in supplementary table S1.

Supplementary figure S2: Yeast-one-hybrid assay of 24 tomato HSFs on *Sly-MIR169d-1* promoter. Interaction of only Sly-HSFA6b on *Sly-MIR169d-1* promoter highlights the specificity of the Y1H assay. The co-transformed Y1H gold yeast strain cultures were spotted (O.D. 0.1 to 0.0001) onto plates lacking URA and LEU with and without specific Aureobasidin A concentration and incubated for 3 days. Promoter of *sHSP* has been used as a positive control to show binding of HSFA3a on *sHSP* promoter (Li et al. 2013) and empty vectors as negative controls. The positive binding of HSFA6b on *Sly-MIR169d-1* promoter was assessed at 200 ng/ml Aureobasidin A concentration.

Supplementary figure S3: HSF-mediated transcriptional activation of tomato *MIR169* promoters. *In-planta* transient assays showing HSF-mediated transcriptional activation of *Sly-MIR169:GUS* promoters in *Nicotiana benthamiana* as assessed by qRT-PCR. The fold-change of *GUS* transcripts were calculated using the $2-\Delta\Delta Ct$ method in (A) to (E). (A) Sly-HSFA1e with *pro-Sly-MIR169a-1:GUS*. (B) Sly-HSFA7a and Sly-HSFA1a with *pro-Sly-MIR169b:GUS*. (C) Sly-HSFA6b with *pro-Sly-MIR169c:GUS*. (D) Sly-HSFB1a with *pro-Sly-MIR169d:GUS*. (E) Sly-HSFA6b with *pro-Sly-MIR169d-1:GUS*. The *NPTII* gene co-expressed in the *pro-Sly-MIR169:GUS* construct was taken as the reference gene for normalization. The *GUS* expression from the reporter (*pro-Sly-MIR169:GUS*) infiltrated samples was set to one for normalization (marked as green bars). These experiments were repeated at least six times; the error bars represent standard deviation between the biological replicates.

Supplementary figure S4: Expression profiling of HSFs:miRNAs:NF-YAs during heat stress in tomato leaves. (A) Expression profiles of *HSF* genes that regulate *Sly-MIR169* transcription in WT tomato leaves during control and heat stress condition by

qRT-PCR using the 2−ΔΔCt method. (B) Expression of mature miR169s in response to heat stress as determined by taqman based qRT-PCR in control and heat challenged WT tomato leaves. (C) qRT-PCR based expression profiles of target *NF-YA* transcripts in WT control and heat stressed tomato leaves using the 2−ΔΔCt method. Data represents mean values and standard deviation of biological replicates. *Tubulin* was used as endogenous control in (A) and (C), while *Sly-U6snRNA* was used as normalization control in (B). Error bars in (A), (B) and (C) represent the standard deviation of biological replicates. The fold change normalization was done by setting the control value as one for all qRT-PCR expression analysis.

Supplementary figure S5: Determining the specificity of HSFs silencing in VIGS silenced tomato plants. (A-C) qRT-PCR based expression profiles of putative off target genes using the 2−ΔΔCt method. (A) Relative expression of Sly-*HSFA1b* and Sly-*HSFA1c* off targets in TRV-*Sly-HSFA1a* silenced plants. (B) Relative expression of Sly-*HSFA6a* off target in *TRV-Sly-HSFA6b* silenced plants. (C) Relative expression of Sly-*HSFA4c* and Sly-*HSFB4a* off targets in *TRV-Sly-HSFA7a* silenced plants. Error bars in (A-C) represent the standard deviation of biological replicates. *ACTIN* was used as endogenous control in A-C. Data normalization and fold change values with *TUBULIN* were presented in Supplementary table S3. The fold change normalization was done by setting the TRV-EV expression as one for all qRT-PCR expression analysis.

Supplementary figure S6: In planta validation of heat governed HSFs mediated transcriptional regulation of MIR169s and NF-YAs in tomato leaves.

The expression analysis of *HSF* genes, *Sly-MIR169* precursors and *NF-YA* transcripts in VIGS plants silenced for *Sly-HSFA1a* (A), *Sly-HSFA7a* (B), *Sly-HSFA6b* (C) and *Sly-HSFB1a* (D). 15-days-old tomato plants were agro-infiltrated with empty vector (TRV-EV) or TRV-Sly-HSF VIGS constructs. The VIGS established plants were subjected to heat stress after 3-weeks-of agro-infiltration, and used for expression studies of different genes. Graphical data represents mean values of expression of three to four biological sets. Error bars show standard deviation. To represent the negative fold-change on y axis, the relative expression values of down-regulated genes were transformed using the formula [-(1 / RQ ≤0.5)]. *Tubulin* was used as endogenous reference control. The

fold change normalization was done by setting the control value as one for all qRT-PCR expression analysis.

Supplementary figure S7: Virus induced silencing of Sly-*NF-YA9* and Sly-*NF-YA10*. (A) qRT-PCR analysis of Sly-*NF-YA9* and Sly-*NF-YA10* gene in TRV-EV, *TRV-Sly-NF-YA9* and *TRV-Sly-NF-YA10* plants in control (non-stressed) conditions confirming their silencing. The fold change normalization was done by setting the TRV-EV expression as one for all qRT-PCR expression analysis. (B) Expression of Sly-*NF-YA9* and Sly-*NF-YA10* in WT tomato and TRV-EV infiltrated plants in control (non-stressed) conditions. The fold change normalization was done by setting the WT control value as one for all qRT-PCR expression analysis (C-D) Expression profiles of *Sly-NF-YA9* (C) and *Sly-NFY-A10* (D) in *TRV-Sly-NF-YA9/A10* VIGS silenced plants in heat stress (HS) in comparison to control non-stressed conditions. The fold change normalization was done by setting the control value as one for all qRT-PCR expression analysis. The expression levels of genes were calculated using the $2{-}\Delta\Delta Ct$ method and presented using fold-change values. *Actin* was used for normalising expression in (A-D). Fold change values for (A-D) with *TUBULIN* normalisation were presented in Supplementary table S3. EV: vector control. Error bars represent the standard deviation of biological replicates.

Supplementary figure S8: Determining the specificity of NF-YA9/A10 silencing in VIGS silenced tomato plants. (A-B) qRT-PCR based expression profiles of putative off target genes using the $2{-}\Delta\Delta Ct$ method. (A) Relative expression of Sly-*NF-YA1, Sly-NF-YA3* and Sly-*NF-YA10* off targets in TRV-*Sly-NF-YA9* silenced plants. (B) Relative expression of *Sly-NF-YA9 (*no off targets were predicted for TRV-Sly-NF-YA10, we used *Sly-NF-YA9* to show specificity of silencing) as in *TRV-Sly-NF-YA10-VIGS* silenced plants. Error bars in (A-B) represent the standard deviation of biological replicates. *ACTIN* was used as endogenous control in A-B. Fold change values for (A-B) with *Tubulin* normalisation were presented in Supplementary table S3. The fold change normalization was done by setting the TRV-EV expression as one for all qRT-PCR expression analysis.

Supplementary figure S9: Expression analysis of heat stress responsive genes in control and heat treated TRV-Sly-*NF-YA9* and TRV-Sly-*NF-YA10* silenced tomato plants.

(A-E) Expression profiles of HSR genes in control and heat treated TRV-EV, *TRV-Sly-NF-YA9* and *TRV-Sly-NF-YA10* silenced plants. qRT-PCR based fold change in expression in control and heat stressed plants of *Sly-HSFA2* (A); *Sly-HSFA7a* (B); *Sly-HSFA3a* (C); *Sly-APX* (D) and *Sly-HSP17.6 CII*. (E). Relative expression of HSR genes in *TRV-Sly-NF-YA9* and *TRV-Sly-NF-YA10* silenced plants in control conditions as compared to *TRV-EV* plants. (G) Expression profiles of HSR genes in control and heat treated *TRV-Sly-NF-YA9* and *TRV-Sly-NF-YA10* silenced plants. The expression levels of genes were calculated using the 2–ΔΔCt method and presented using fold-change values. *Tubulin* was used as reference gene for A-E and *Actin* was used as endogenous control for F-G. Fold change values for (F-G) with *TUBULIN* normalisation were presented in Supplementary table S3 The fold change normalization was done by setting the expression in control non-stressed as one for all qRT-PCR expression analysis in (A-E). The fold change normalization was done by setting the TRV-EV heat stressed expression as one for all qRT-PCR expression analysis in (F). The fold change normalization was done by setting the expression of TRV-EV in control non-stressed as one for all qRT-PCR expression analysis in (G). Error bars represent the standard deviation of independent biological replicates.

Supplementary figure S10: Expression profiling of At-miR169s in heat stress.
Expression of mature At-miR169s in response to heat stress as determined by Taqman-based qRT-PCR. *U6snRNA* was used as the endogenous normalization control. The average values of multiple biological replicates are plotted as bars, error bars depict standard deviation between three replicates. The fold change normalization was done by setting the control value as one for all qRT-PCR expression analysis.

Supplementary figure S11: Expression analysis of At-MIR169d in heat stress in Arabidopsis. Expression levels of *At-MIR169d* in wild-type plants subjected to 0

(control) and 2 h heat stress at 42 °C. Data are shown as means ± SE of three
biological replicates and normalized to *At-EF-α* reference gene using (2−ΔΔCT)
method. The fold change normalization was done by setting the control value as one for
the qRT-PCR expression analysis.

Supplementary figure S12: Schematic structure of *At*-HSFA2 gene carrying the T-
DNA insertion and lack of *At*-HSFA2 expression in the *At-hsfA2* mutant
(Salk_008978). (A) The two exons of *At*-HSFA2 genomic DNA are shown according to
the gene features annotated at TAIR database. T-DNA insertion site is shown as
inverted triangle in the second exon of *At*-HSFA2. The exon sequences are depicted by
red and the UTRs are represented by green color. (B) Genotyping PCR of *At-hsfa2*
mutants using genomic DNA, presence of border specific amplification in *At-hsfa2*
confirms their homozygosity, while presence of only gene specific amplification in
control Columbia plants confirms their wild type nature. (C) Expression of *At-HSFA2*
was determined by RT-PCR, using full length cDNA from WT and mutant *At-hsfa2*
plants. Absence of *At-HSFA2* specific band in mutant plants confirms the knockout
nature of *At-hsfa2* homozygous mutants. RNA was isolated from detached mature
leaves of the wild-type (wt) or *At-hsfa2* plants. Expression of *actin* gene is shown as a
loading control.

1218

Supplementary figure S13: Expression pattern of miRNA169d:NF-YA in
Arabidopsis. (A) Expression levels of mature miR169d and eight miR169 target genes
in wild-type plants subjected to 0 (control) and 2 h heat stress at 42°C using qRT-PCR.
Data are shown as means ± SD of biological replicates and normalized to *actin* and
5SrRNA reference genes, for mRNA and miRNA respectively, using $(2^{−ΔΔCT})$ method for
A and with *At-EF-α* reference gene and *U6snRNA* for mRNA and miRNA respectively,
using $(2^{−ΔΔCT})$ method for B. The fold change normalization was done by setting the
control value as one for all qRT-PCR expression analysis.

1227

Supplementary figure S14: Establishing miRNA169:target modules operating in
Arabidopsis. (A) Taqmann based qRT- qPCR analysis of the accumulation of mature
miR169abc, miR169defg, miR169hijklmn and miR171bc (used as negative control) in

Arabidopsis MIM169defg lines. Average values of expression for different miR169s in
three independent lines of MIM169defg transgenic plants were plotted as bars and error
bars deoicts the standard deviation between the replicates. (B) qRT-PCR analysis of the
accumulation of the *At-NF-YA* transcripts in MIM169defg Arabidopsis lines. Data are
presented as average fold induction relative to control where induction is the ratio of
transcript levels in two independent MIM169defg transgenic lines to those in wild-type
plants (WT). Error bars, ± SD of three repeats. (C) qRT-PCR analysis of the reduction of
At-NF-YA2 transcripts in three independent miR169d over expressing (OE) lines. Data
are presented as average fold in two independent *MIM169defg* lines relative to WT
control, that was set as one.

1241

Supplementary figure S15: Heat stress mediated transcriptional regulation of *At-
NF-YA2.* (A) Histochemical GUS expression of Pro-*At-NF-YA2:GUS* transgenic
Arabidopsis plants during control and heat stress. (B) qRT-PCR based determination of
GUS transcripts of Pro-*At-NF-YA2:GUS* transgenic Arabidopsis plants during control
and heat stress. These experiments were repeated 3 times with 20 seedling per
replicate. Error bars depict standard deviation between three replicates. The fold
change normalization was done by setting the control value as one for qRT-PCR
expression analysis.

1250

1251 **Supplementary figure S16: Delineating the existence and localization of**
1252 **miR169d:At-NF-YA2 functional module in Arabidopsis.** Histochemical GUS
1253 expression patterns of *Pro-At-MIR169d:GUS and Pro-At-NF-YA2:GUS* promoter
1254 reporter transgenic Arabidopsis plants in: seedling (2 weeks old), complete rosette (4
1255 weeks), mature rosette leaf, cauline leaf and flower

1256

1257 **Supplementary figure S17: Expression levels of *At-NF-YA2* in *At-NF-YA2***
1258 **overexpressing transgenic plants.** Relative expression of *At-NF-YA2* in transgenic
1259 Arabidopsis plants overexpressing *At-NF-YA2* under constitutive CaMV35S promoter.
1260 Bar represents the average data of three biological replicates and error bar represents
1261 standard deviation of the replicates.

1262

Supplementary figure S18: Schematic structure of *At-NF-YA2* gene carrying the T-DNA insertion and lack of *At-NF-YA2* expression in the *At-nf-ya2* mutant (SALK_021228). (A) The four exons of *At-NF-YA2* genomic DNA are shown according to the gene features annotated at TAIR database. T-DNA insertion site is shown by inverted triangle in the fourth exon of *At-NF-A2*. The exon sequences are depicted by red and the UTR are represented by green color. (B) Genotyping PCR of *At-nf-ya2* mutants, presence of border specific amplification in *At-nf-ya2* plants genomic DNA confirms their homozygosity, while presence of only gene specific amplification in control Columbia plants confirms their wild type nature. (C) Expression of *At-NF-YA2* was determined by RT-PCR, using full length cDNA from WT and mutant *At-nf-ya2* plants. Absence of *At-NF-YA2* specific band in mutant plants confirms the knockout nature of *At-nf-ya2* homozygous mutants. RNA was isolated from detached mature leaves of the wild-type (WT) or *At-nf-ya2* plants. Expression of actin is shown as a loading control.

1277

Supplementary figure S19: Expression profiling of HSR genes in different transgenic lines of Arabidopsis. Expression patterns of five heat stress-responsive genes in WT, *At-MIR169d-OE, At-NF-YA2-OE*, *At-nf-ya2, pNF-YA2:NF-YA2*-r and MIM169defg plants. Data are shown as means ± SD of biological replicates and normalized to *At-EF-α* reference gene using $(2^{-\Delta\Delta CT})$ method. The fold change normalization was done by setting the control value as one for qRT-PCR expression analysis.

1285

Supplementary figure S20: Yeast-one-hybrid assay based transcriptional regulation of *At-HSFA3a, At-HSFA7a* and *At-HSFA7b* promoters by At-NF-YA2. Positive interaction of At-NF-YA2 with the promoters of *At-HSFA3a* and *At-HSFA7b* were obtained. Serial dilutions (O.D. 0.1 to 0.0001) of yeast cultures were spotted onto plates lacking LEU with specific Aureobasidin A concentration and incubated for 3 days.

1291

Supplementary figure S21: Thermotolerance assay of WT and *At-hsfa7a* plants. (A) Two-weeks-old soil-grown WT and *At-hsfa7a* mutant plants were subjected to 0 (control) or 2h heat stress at 42°C, and damage was recorded 6 days later. (B) Estimation of survival (percentage) of heat stress treated WT and *At-hsfa7a* plants. Plants were assayed for heat stress tolerance after 6 days of HS. These experiments were repeated at least three times with similar results, and average data of all experiments are shown. Error bars, ± SD of three repeats. ***$p<0.001$ vs. wild type, by two-tailed Student's t-test.

Supplementary figure S22: At-NF-YA2, Arabidopsis orthologue of tomato Sly-NF-YA9 and Sly-NF-YA10. (A-B) Protein sequence alignment of tomato Sly-NF-YA9 (A) and Sly-NF-YA10 with At-NF-YA2. Protein sequences were aligned with Arabidopsis proteome at TAIR database by using Blastp tool.

Supplementary figure S23: Binding of *Sly-HSFA6b* to Sly-MIR169d-1 promoter. Absence of any amplification in no-Ab (no antibody) and slight amplification in IgG control confirms the specificity of immunoprecipitation. Presence of strong amplification on input DNA confirms the good amplification efficiency of the primer used for analysis. Presence of amplification in Anti-GFP lane confirms the enrichment of Sly-MIR169d-1 promoter in *Sly-HSFA6b:GFP* CHIP. L denotes the 100 bp ladder.

Supplementary figure 24: Virus induced silencing of tomato *Phytoene desaturase* (*PDS*) gene. *PDS* silencing was used as a control for assessing establishment of gene silencing by VIGS. (A) Silencing of the PDS control gene causing photobleaching in tomato plants 3 weeks after agroinfection. (B) Enlarged individual leaves of TRV-PDS and control WT leaves.

Supplementary Table legends:

Supplementary table S1: Distribution of perfect and imperfect heat stress elements (HSE) in*Sly-MIR169* promoters.

Supplementary table S2: Interaction summary of *Sly-MIR169* promoters and Sly-HSFs. Positive HSF interaction is marked in red colored boxes.

Supplementary table S3: All Qrt-PCR raw data, calculations and fold change normalizations with different endogenous controls.

Supplementary table S4: Off targets of candidate VIGS constructs used in the study

Supplementary table S5: Promoter analysis of HSR genes in tomato.

Supplementary table S6: Number of HSE elements and minimal inhibitory concentration of Aureobasidin (Aba) for all At-MIR169 promoters

Supplementary table S7: Interaction summary of At-MIR169 promoters and At-HSFs. Positive HSF interaction is marked in red colored boxes.

Supplementary table S8: qRT-PCR based expression profiles of HSR genes in *At-MIR169d OE, At-NF-YA2 OE, At-nf-ya2, At-pNF-YA2:NF-YA2-r* and *MIM169defg* transgenic lines.

Supplementary table S9: Analysis of Ct similarities across different conditions for the reference genes used in the study.

Supplementary table S10: List of Primers used in the study.

Supplementary table S11: Primer efficiency of all the primers used in the study.

Supplementary table S12: Minimal inhibitory concentration of Aureobasidin (Aba) for all *Sly-MIR169* promoters.

1349

1350

1351

1352

1353

1354

1355

1356

1357
1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

Figure 1:Yeast one hybrid and Chromatin immunoprecipitation (ChIP-qPCR) assay of Sly-MIR169 promoters and HSFs. (A) Yeast-one-hybrid assay to assess transcriptional regulation of heat-responsive *MIR169* promoters by HSFs. Only the positive Y1H interactions between different HSFs and *Sly-MIR169* promoters have been shown. The *Sly-MIR169* promoters were cloned in the reporter vector pABAi and the full-length cDNA of HSFs were cloned in the pGADT7 vector. After the co-transformation into Y1H gold yeast strain, serial dilutions (O.D. 0.1 to 0.0001) of yeast cultures were spotted onto plates lacking URA and LEU with specific Aureobasidin A concentration and incubated for 3 days. Promoter of *sHSP* has been used as a positive control to show binding of HSFA3a on *sHSP* promoter (Li et al. 2013) and empty vectors as negative controls. The Aureobasidin A concentrations for all *Sly-MIR169* promoters is presented in supplementary table S7. (B-I) Chromatin immunoprecipitation (ChIP-qPCR) assay of *Sly-MIR169* promoters and HSFs in *Nicotiana benthamiana* . Fold enrichment of *Sly-MIR169a-1* promoter in HSFA1e immunoprecipitated sample (B); Fold enrichment of *Sly-MIR169b* promoter in HSFA1a immunoprecipitated sample (C); in HSFA7a immunoprecipitated sample (D); Fold enrichment of *Sly-MIR169C* promoter in HSFA4a immunoprecipitated sample (E); Fold enrichment of *Sly-MIR169C* promoter in HSFA6b immunoprecipitated sample (F); Fold enrichment of *Sly-MIR169d* promoter in HSFA8a immunoprecipitated sample (G); in HSFB1a immunoprecipitated samples (H); Fold enrichment of *Sly-MIR169d-1* promoter in HSFA6b immunoprecipitated samples (I). These experiments were repeated 3-4 times; the error bars represent standard deviation between the replicates. The values for different promoters in IgG immunoprecipitated sample were set as one for normalization.

Figure 2: Validation of HSF:miR169:*NF-YA* node components in tomato upon heat stress. (A) Expression profiles of *HSF* genes that regulate *Sly-MIR169* transcription in WT tomato leaves during control and heat stress condition by qRT-PCR using the 2−ΔΔCt method. (B) Expression of mature miR169s in response to heat stress as determined by taqman based qRT-PCR in control and heat challenged WT tomato leaves. (C) qRT-PCR based expression profiles of target *NF-YA* transcripts in WT control and heat stressed tomato leaves using the 2−ΔΔCt method. (D-G) The expression analysis of *HSF* genes, *Sly-MIR169* precursors and *NF-YA* transcripts in VIGS plants silenced for *Sly-HSFA1a* (D), *Sly-HSFA7a* (E), *Sly-HSFA6b* (F) and *Sly-HSFB1a* (G). In (D-G) 15-days-old tomato plants were agro-infiltrated with empty vector (TRV-1, TRV-2 VC) or TRV-Sly-HSF VIGS constructs. The VIGS established plants were subjected to heat stress after 3-weeks-of agro-infiltration, and used for expression studies of different genes. Graphical data represents mean values of expression of three to four biological sets. Error bars show standard deviation. To represent the negative fold-change on y axis in (C-G), the relative expression values of down-regulated genes were transformed using the formula [-(1 / RQ ≤0.5)]. *Actin* was used as endogenous control in (A), (C) and (D-G), while *5SrRNA* was used as reference control in (B). The fold change normalization was done by setting the control value as one for all qRT-PCR expression analysis.

Figure 3. Functional validation of the role of *Sly-NF-A9* and *Sly-NF-A10* in HS response in tomato. (A) Phenotype of TRV-VC (Vector control), *TRV-Sly-NF-YA9* and *TRV-Sly-NF-YA10* silenced plants after heat stress. Fifteen-days-old tomato plants were agro-infiltrated with empty vector (TRV-VC), *TRV-Sly-NF-YA9* or *TRV-Sly-NF-YA10* silencing constructs. Plants were given HS 3-weeks-after infiltration. Phenotype of HS-treated silenced plants was scored six-days-post HS. Experiments were repeated 3-4 times with 8 plants per replicate. (B) Percent survival of TRV-EV, *TRV-Sly-NF-YA9* and *TRV-Sly-NFYA10* silenced plants afters HS. (C) Estimation of water use efficiency (mmol mol^{-1}), net photosynthesis rate (μmol m^{-2}s^{-1}), and transpiration rate of TRV-EV, *TRV-Sly-NF-YA9* and *TRV-Sly-NFYA10* silenced plants afters HS. (D-E) Expression profiles of HSR genes in control and heat treated *TRV-EV, TRV-Sly-NF-YA9* and *TRV-Sly-NF-YA10* silenced plants. The expression levels of genes were calculated using the 2−ΔΔCt method and presented using fold-change values. The fold change normalization was done by setting the control value as one in (D-H). *Actin* was used as endogenous control. Error bars represent the standard deviation of two independent biological replicates. *p<0.05, **p<0.01 and ***p<0.001 vs. wild type, by two-tailed Student's t-test.

Figure 4. Transcriptional regulation of MIR169s promoters by HSFs in Arabidopsis. (A) miR169s expression in response to heat stress as determined by Taqman-based qRT-PCR of control and heat challenged plants. 5S rRNA was used as the normalization control. Experiment was repeated three times and average values are plotted as bars, error bars depict standard deviation between the replicates. (B) Y1H assay showing the positive interaction of HSFs with the promoters of different *At-MIR169s*, Y1H assay shows binding of At-HSFA7b, At-HSFB2b, At-HSFB3a, At-HSFA2 and At-HSFA1b on *At-MIR169b*, d and h promoters. This experiment was repeated at least three times with similar results, and data from one representative experiment is shown. Serial dilutions (O.D. 0.1 to 0.0001) of yeast cultures were spotted onto plates lacking URA and LEU with specific Aureobasidin A concentration and incubated for 3 days. The Aureobasidin A concentrations for all *At-MIR169* promoters is presented in supplementary table S3. (C) Expression levels of *At-MIR169d* in wild-type plants subjected to 0 (control) and 2 h heat stress at 42 °C. Data are shown as means ± SD of three biological replicates and normalized to actin reference gene using (2−ΔΔCT) method. (D) Expression patterns of *Pro-At-MIR169d:GUS* lines of transgenic Arabidopsis plants subjected to 0 (control) or 2 h of heat stress at 42°C. (E) Levels of GUS transcripts in *Pro-At-MIR169d:GUS* lines, measured by qRT-PCR after 0 or 2 h of heat stress at 42°C. (F) Transcriptional activity of *Pro-At-MIR169d:GUS* reporter construct in WT and *At-hsfa2* mutant background during 2 h of heat stress at 42°C. (G) Expression of *GUS* transcript driven by *Pro-At-MIR169d* and *At-MIR169d* precursor in soil-grown wild-type plants of Arabidopsis and *At-hsfa2* mutant. To represent the negative fold-change on y axis in figure (G), the relative expression values of down-regulated genes were transformed using the formula [-(1 divided by RQ ≤0.5)]. The fold change normalization was done by setting the control value as one in Figure (A, C, E and G). Error bars represent the standard deviation. These experiments were repeated at least three times with similar results, and data from one representative experiment are shown in D and F.

Figure 5: Assaying the miR169d:At-NF-YA2 node in heat stress in Arabidopsis. (A) Thermotolerance was gauged in *At-MIR169d-OE*, *At-NF-YA2-OE*, *At-nf-ya2*, *pNF-YA2:NF-YA2*-r and MIM169defg transgenic plants in comparison to wild-type (WT) plants. Two-weeks-old soil-grown plants were subjected to 0 (control) or 2h heat stress at 42°C, and damage was recorded 6 days later. (B) Estimation of survival (percentage) of HS treated *At-MIR169d-OE*, *At-NF-YA2-OE*, *At-nf-ya2*, *pNF-YA2:NF-YA2*-r and MIM169defg plants. Plants were assayed for heat stress tolerance after 1 week of HS. These experiments were repeated at least five times with similar results, and average data of all experiments are shown. Error bars, ± SD of five repeats. (C) Expression patterns of heat stress-responsive genes in WT, *At-MIR169d-OE*, *At-NF-YA2-OE*, *At-nf-ya2*, *pNF-YA2:NF-YA2*-r and MIM169defg plants. Data are shown as means ± SD of biological replicates and normalized to *Actin* using ($2^{-\Delta\Delta CT}$) method. Log2 transformation was applied to the RQ data to obtained negative fold change. (D-E) DAB and Trypan blue staining of HS treated *At-MIR169d-OE*, *At-NF-YA2-OE*, *At-nf-ya2*, *pNF-YA2:NF-YA2*-r and MIM169defg plants to estimate the production of ROS and cell death, respectively. *$p<0.05$, **$p<0.01$ and ***$p<0.001$ vs. wild type, by two-tailed Student's t-test. Three independent lines of *MIR169d-OE* were used and similar results were obtained for all studied parameters, a single representative picture was shown in the figure A, B, D and E. The fold change normalization was done by setting the control value as one for figure (C).

Figure 6: Transcriptional regulation of *At-HSFA3a* and *At-HSFA7b* by At-NF-YA2 during heat stress in Arabidopsis. (A) Graphical representation of CCAAT cis-elements in At-HSR gene promoters HSFA3a, HSFA7a, HSFA7b, APX and sHSP17.6A. The 2 kb upstream sequences of start codon were analysed by plant care and Plant PAN promoter analysis tools. The positions of CCAAT boxes are depicted by violet coloured tabs. (B) Chromatin immunoprecipitation (ChIP)-qPCR assay of At-NF-YA2 on At-HSR gene promoters. (C-D) *In-planta* estimation of transcriptional regulation of *At-HSFA3a* and *At-HSFA7b* transcription by At-NF-YA2 during heat stress. (C) *Pro-At-HSFA3a:GUS* reporter construct was agro-infiltrated in WT, *At-NF-YA2-OE* and *At-nf-ya2* backgrounds. *GUS* was visualized 2-days post-infiltration by histochemical staining. (D). qRT-PCR based quantification of *GUS* transcripts in promoter reporter constructs infiltrated plants, 2-days-post infiltration by using *GUS* specific primers. The *NPTII* gene present in the same vector backbone was used as reference control. To represent the negative fold-change on y axis in, the relative expression values of down-regulated genes were transformed using the formula [-(1 divided by RQ ≤0.5)]. The fold change normalization was done by setting the control value as one. (E) Thermotolerance assay of WT and *At-hsfa3a* mutant plants. Two-weeks-old soil-grown plants were subjected to 0 (control) or 2h heat stress at 42°C, and damage was recorded 6 days post-stress. (F) Estimation of survival (percentage) of WT and *At-hsfa3a* plants 6 days post heat stress. The experiments were repeated at least three times with similar results, and average data of all experiments are shown. Error bars, ± SD of three repeats. *p<0.05, **p<0.01 and ***p<0.001 vs. wild type, by two-tailed Student's t-test.

Figure 7: Model for HSF-induced miR169:NF-YA module in mediating thermotolerance in tomato and Arabidopsis. Heat stress induces the expression of HSFs by altering the cellular ROS levels as well as other cues independent of ROS signaling. HSFA1a, HSFA2 and HSFA7a are the key player of plant HSR response. These HSFs bind to the promoters of *MIR169* leading to transcriptional enhancement of miR169s in both tomato and Arabidopsis. Enhanced accumulation of miR169s reduces the levels of *Sly-NF-YA9/A10* in tomato and *At-NF-YA2* in Arabidopsis that leads to enhancement of HSR genes expression like *HSFA2, HSFA3* and *HSFA7s*, possibly via altering ABA levels. In Arabidopsis, At-miR169-mediated down-regulation of *At-NF-YA2* during HS may favor functional DREB2A:NF-YC-10:NF-YB3 complex formation to up-regulate *At-HSFA3a* expression during HS. Enhanced levels of HSFAs in turn feed the transcription of *MIR169* genes. The heat-mediated orchestration of parallel miRNA pathways (miR169:NF-YA, miR398:CSD and miR156:SPL) alters the levels of HSR genes and regulate plant thermotolerance. Solid lines denote a confirmed interaction and dotted lines are predictive. Red and black lines denote findings of this study and existing literature, respectively. Black colored numerals refer to the studies that established the particular connection between different components of the model. Reference key- 1: Guan et al. 2013; 2: Stief et al. 2014; 3: Sato et al. 2014; 4: Zhao et al. 2009, Luan et al. 2015, Ding et al. 2016; 5: Sakuma et al. 2006. CSD: Copper superoxide dismutase; SPL: SQUAMOSA-promoter binding like; ABA: Abscisic acid; ROS: Reactive oxygen species; DREB: Dehydration responsive element binding protein.