Functional insights of two MATE transporters from *Vibrio fluvialis*

Priyabrata Mohanty[1], Aneri Shah[2] and Ashima Kushwaha Bhardwaj[1]*

1. Department of Human Health and Diseases, Indian Institute of Advanced Research, Koba
Institutional Area, Gandhinagar 382 007, Gujarat, India
2. Clinic of Nephrology and Hypertension, Diabetes and Endocrinology, Otto-von-Guericke
University Magdeburg, 39120 Magdeburg, Germany

Running title: MATE-type efflux pumps from Vibrio fluvialis

To whom correspondence should be addressed: Dr. Ashima Kushwaha Bhardwaj**,** Department
of Human Health and Diseases, Indian Institute of Advanced Research, Koba Institutional Area,
Gandhinagar- 382 007, Gujarat, India;
Tel.: + 91 -079-30514235; Fax: + 91 -079-30514110 ; E mail : ashima.bhardwaj@gmail.com

Keywords: Efflux pumps; Recombinant expression; multi-ion dependence; efflux pump
inhibition

Summary

Functional characterization of H- and D-MATE (Multidrug and Toxin Extrusion) transporters

from clinical isolates of *Vibrio fluvialis* revealed H-type conferred resistance to fluoroquinolones,

ethidium bromide and safranin whereas D-type exhibited marginal resistance towards ethidium

bromide only. Both H-/D-type transporters were inhibited by reserpine resulting in increased

intracellular norfloxacin concentration. The efflux was facilitated by both Na^+/K^+ ions, suggesting

that these efflux pumps were ion-dependent. In presence of various classes of EPIs, there was

decrease in MIC exhibited by H-/D-type efflux pumps towards norfloxacin which didn't translate

into transport inhibition. But reserpine presented a conclusive pattern with decrease in MIC

towards norfloxacin and increased norfloxacin accumulation inferring maximum inhibition.

Substrate binding and electrostatic charge distribution of both the transporters was similar to

other known MATE transporters. The H-type exhibited 10 transmembranes and D-type exhibited

11 TMs which was different from other MATE transporters known to have 12 TMs

(Transmembranes). Data derived from molecular docking and ion binding studies revealed that

Aspartic Acid residue in 1st TM acts as ion binding site with transport mechanism similar to

NorM. Electrostatic potential map of both the transporters revealed that there is a cavity

formation within the transporters surrounded by charged electronegative amino acid residues.

Interestingly, surface models of both transporters revealed that 1[st] TM forms covalent bond with

7[th] TM towards extracellular space. Docking studies also revealed that reserpine covalently

binds to central pocket of both transporters and serves as excellent EPI against these

transporters as evidenced by MIC and drug accumulation assays.

Introduction

Resistance mediated by efflux pumps is one of the non-specific mechanisms that pathogenic

microbes employ to lower intracellular drug concentration (Bhardwaj & Mohanty, 2012, Van

Bambeke *et al.*, 2003, Poole, 2000, Poole, 2004, Poole, 2005). Efflux pumps could be primary

transporters utilizing ATP for transport or could be secondary transporters that utilize sodium-

/proton-motive force for transport of substrates. Till date six families of transporters are known

for their involvement in multidrug resistance (MDR) in prokaryotes (Bhardwaj & Mohanty, 2012,

Van Bambeke *et al.*, 2003, Hassan *et al.*, 2015). Multidrug and toxic compound extrusion

(MATE) family of efflux pumps is one of the newest and least characterized family of

transporters that are proven to be involved in resistance to various antibiotics and dyes

including quinolones (Mohanty *et al.*, 2012, Omote *et al.*, 2006). NorM, the prototype member

was discovered from *Vibrio parahaemolyticus* (Morita *et al.*, 1998) and subsequently its

homologs and orthologs were identified in *Escherichia coli*, *Neisseria gonorrhoeae* and *V.*

cholerae (Long *et al.*, 2008, Morita *et al.*, 2000). The MATE family further consists of three

protein clusters : the NorM branch involved in MDR, a branch containing several eukaryotic

proteins and a branch containing DinF involved in SOS response and MDR (Otsuka *et al.*,

2005). Growing academic interest in this family of transporters is evident from the fact that

structures have been solved for a few within a span of last few years from prokaryotic as well

as eukaryotic sources(Tanaka *et al.*, 2013, He *et al.*, 2010, Lu *et al.*, 2013b, Lu *et al.*, 2013a,

Wong *et al.*, 2014, Tanaka *et al.*, 2017, Miyauchi *et al.*, 2017) . Preliminary *in-silico* and drug

transport studies of MATE family of efflux pumps revealed that these efflux pumps are sodium-

ion coupled and recognize wide spectrum of substrates but neither of these characteristics have

been explicitly explained at molecular level (He *et al.*, 2004). MATE pumps originated from

flippases (Meeske *et al.*, 2015). Both MATE and flippases belong to same family of proteins

multidrug/oligosaccharidyl-lipid/polysaccharide (MOP) transporter superfamily. Divergence can

be seen as functional role has been attributed to aspartic acid in 1st transmembrane of MATE-

pumps, which acts as ion binding site whereas serine at 17 position is known to play functional

role in MurJ flippases (Kuk *et al.*, 2017). From our laboratory, two MATE-type members (H- and

D-) were first identified in *Vibrio fluvialis* (causes cholera-like illness) (Mohanty *et al.*, 2012). The

genes were cloned in pBR322 vector and the protein functions were assessed using MIC and

drug transport assays. Both the pumps conferred resistance to fluoroquinolones norfloxacin

and ciprofloxacin.   In the present study, we cloned both the transporters in arabinose promoter-

based pBAD vector which allowed visualization of recombinant protein and better control over

protein expression. Functional characterization was performed with assays involving MICs, drug

uptake and efflux pump inhibitors (EPIs). Molecular docking, ion binding site predictions and

surface electrostatics of these transporters using the known MATE-type transporters as

reference proteins was carried out to further characterize the functionality of these transporters.

Results and Discussion

Expression of recombinant H- and D- MATE-type efflux pumps in E. coli- The sequenced

genes of D- and H-type efflux pumps were designated as *vcd*, *vfd*, *vch* and vfh, *where* c stands

for *V. cholerae* and *f* stands for *V. fluvialis*. Corresponding to each gene, the proteins were

designated as VCH, VCD, VFH and VFD. Further the genes were cloned in arabinose driven

pBAD expression system and induced as mentioned in materials in methods. Coomassie blue

stained gel of arabinose-induced pBAD-*vfd* samples did not show the band corresponding to

recombinant VFD at the expected position of 55 kDa (Fig. 1A) whereas the expression of

positive control Lac-His was detected at 125 kDa as expected.

Fig. 1. Total cell expression in induced (+) and uninduced (-) cell lysates of different recombinant pBAD clones. **A.** Coomassie blue stained 10% SDS-PAGE for total cell lysates from pBAD-*vfd* along with pBAD-Lac-His positive control in *E. coli* LMG194; **B.** Western blot of induced (+) and uninduced (-) cell lysates of pBAD-*vfd* along with pBAD-Lac-His positive control. **C.** Western blot of recombinant VCH, VFH, VCD, VFD and Lac-His in LMG194. **D.** Western blot of recombinant proteins in K-12 Δ*tolC*. The cells were induced and developed as described in materials and methods. Lane-1 Prestained Spectramulticolour (Fermentas) molecular weight markers have been indicated on the left of each pane. Lane 2,3,4 and 5 represents VCH, VFH, VCD and VFDFor Panels **B** and **D, the** blots were developed with ECL substrates (Novex, Invitrogen) and for Panel **C**, they were developed with DAB and H_2O_2 Goat Ant-V_5 antibody was used as primary antibody.

Therefore, Western Blot was carried out to determine the expression of recombinant

VFD   protein   with   goat   anti-V_5 primary   antibody   and   developed   with   enhanced

chemiluminescence substrates. Results revealed the expression of Lac-His positive control at

right molecular weight but interesting observation was made in case of pBAD-*vfd,* which showed

diffused pattern of bands in the whole lane (Fig 4.16 B). Subsequently, all the four recombinant

clones for H- and D-genes were expressed in *E. coli* LMG194 under same conditions. The

development of Western blots with goat anti-His antibody using colorimetry clearly depicted a

single major band with the mobility greater than 260 kDa in the lanes with induced cells for all

the clones (Fig 1 C) . Therefore, the slow development of the Western blots showed single

major bands whereas the fast/more sensitive ECL detection could also depict the degradation

pattern for the recombinant proteins.

Further, all the four recombinant clones for H- and D-genes (*vch*, *vfh*, *vcd* and *vfd*) were

expressed in *E. coli* LMG194 and Western blots using colorimetry clearly depicted a single

major band with the mobility greater than 260 kDa instead of 55 kDa in the lanes with induced

cells for all the clones (Fig. 1C). The sensitivity of ECL led to the visualization of diffused band

pattern probably indicating either various stages of oligomerisation/recombinant protein

degradation whereas the less sensitive colorimetry showed a single band on Western blot. The

diffused band pattern of the recombinant efflux pumps observed in the current study could be

the final outcome of aberrant mobility due to oligomerisation. The observed aberrant mobility of

recombinant transporter proteins due to oligomerisation has also been reported in case of

ammonium transporter AmtB of *E. coli* (Blakey *et al.*, 2002) where the 45 kDa protein showed

series of bands of varying molecular weights suggesting it to be an oligomer. In our study, the

mobility of the H- and D-recombinant effux pump proteins showed similar pattern on Western

blots where ~55 kDa proteins migrated at >260 kDa molecular weight suggesting that the

recombinant protein existed as oligomers that were stable in the presence of SDS and β-ME.

The recombinant efflux pumps formed a series of aggregated species due to the presence of

hydrophobhic patches that appeared as a ladder of high molecular weight bands under

denaturing and reducing conditions (Blakey *et al.*, 2002). The wild type Na^+/H^+ antiporter of

Methanococcus jannaschii presents itself as a dimer even in the presence of SDS and

undergoes proteolytic degradation (Goswami *et al.*, 2011). The recombinant expression of

membrane proteins is tricky where specialized expression systems have been developed to

counteract protein degradation and oligomerisation (Wagner *et al.*, 2008). Taken together, the

facts from earlier studies and observations from the present study suggested that H- and D-

MATE efflux pumps exhibited aberrant mobility possibly be due to oligomerisation similar to

AcrB due to intra-molecular van der Waals bonding that resists dissociation in presence of SDS

and β-ME.

pBAD recombinant efflux pumps are translocated to the membrane- Upon successful

detection of recombinant proteins in the host cells with aberrant mobility, classical traits of

membrane proteins, further studies were carried out to determine if the recombinant proteins

were actually targeted to the membrane of the *E. coli* host. Two approaches *i.e.* confocal

microscopy and membrane fractionation were utilised to investigate the same.

Earlier bioinformatics studies of both the efflux pumps had predicted that these pumps

were transmembrane proteins (Mohanty *et al.*, 2012). Localization of these efflux pumps in cell

membrane was investigated utilizing confocal microscopy where the laser was programmed to

detect membrane marker Film Tracer 1-43 and CY3 fluorochrome conjugated secondary

antibody from the same pin hole. Results revealed that VFD and VFH were localised in the

membrane of *E. coli* LMG194 (Fig. 2 Panel A: A-F). The recombinant proteins generated red

signal due to CY3 (Fig. 2 A, panelsB and E) and membrane marker generated green signal for

cell membrane (Fig. 2A,panels A and D). DAPI, which was used for nuclear staining was also

employed as counterstain but it did not bind with the nucleoid and was not visible by confocal

imaging. Change in the colour of membrane to yellow in the superimposed image of

recombinant efflux pump protein and membrane marker (Fig. 2A, panels C and F) suggested

that the recombinant proteins were localised in the membrane and were detected by the same

pinhole of the microscope.

Fig. 2. Membrane localisation of recombinant efflux pumps in *E. coli*. **Panel A.** Confocal imaging of pBAD-recombinants in heterologous *E. coli* host. **A/D**:Visualisation of *E. coli* cells harbouring pBAD-*vfh/vfd* stained with membrane marker Film-tracer 1-43; **B/E**:Detection of recombinant pBAD-*vfh/vfd* in *E. coli* labelled with Goat anti-Mouse Cy3 secondary antibody; **C/F** (63X ZOOM 4): Superimposed/merged images from **A/B or D/E** of recombinant pBAD-*vfh/vfd* in *E. coli* LMG194; **G/J**: Visualisation of *E. coli* cells harbouring pBAD-*vfh/vfd* using membrane marker Film-tracer 1-43; **H/K**: Detection of recombinant pBAD-*vfh/vfd* in *E. coli* labelled with Goat anti-Mouse Cy3 secondary antibody; **I/L**: Superimposed/merged images from **G/H** or **J/K** of recombinant pBAD-*vfh/vfd* in *E. coli* K-12 Δ*tolC*2 Δ*tolC*. **Panel B.** Membrane fractionation of *E. coli* harbouring pBAD-*vfd*. 63X Zoom 4 settings were used for obtaining these images. M- Spectramulticolour protein marker (Fermentas). Lane 1- VFD –ve- Total cell lysates for uninduced samples; Lane 2-VFD +ve- Total cell lysates for induced samples; Lane 3- VFD Son (Sup) - total soluble fraction after sonication.; Lane 4- VFD SAR (Sup)- Total soluble proteins after sarkosyl treatment; and Lane 5- VFD SAR Pellet- debris and other subcellular organelles. Lane 6- Lac-HisPositive control. Goat anti-V5 antibody was used as primary antibody. HRP conjugated Mouse Anti-Goat IgG was used as secondary antibody (Jackson Immunoresearch). Prestained Spectramulticolour (Fermentas) molecular weight marker in kDa has been indicated of the left of panel. The blots were developed with Novex ECL substrates (Invitrogen).

To further confirm if the recombinant proteins were localized in the inner membrane, a protocol

utilising a combination of membrane fractionation and selective solubilisation of the inner

membrane with detergents was employed (Hobb *et al.*, 2009, Leyh, 1990, Frankel *et al.*,

1991).Western blot revealed that upon induction, the recombinant protein was detected in total

cell lysate (Fig. 2B, lanes 1 and 2) for VFD. The fraction containing total soluble protein did not

show the presence of recombinant protein indicating that the recombinant protein was not

cytosolic but probably associated with insoluble fraction (Fig. 2B, lane 3). Sarkosyl solubilised

inner membrane fraction (lane 4) revealed the band pattern corresponding to recombinant VFD

confirming that the protein was translocated to the inner membrane. This protein was also

detected in the sarkosyl treated outer membrane pellet indicating partial solubilisation of the

membrane upon sarkosyl treatment (Fig. 2B, lane 5). Apart from the above, partially purified

recombinant VFD from inner membrane also showed aberrant migration and diffused band

pattern as observed earlier.

MIC assays for pBAD recombinants in E. coli K-12 ΔtolC indicated different substrate

profile for H and D pumps - The recombinant constructs for H- and D-type proteins when

introduced into *E. coli* LMG194 did not exhibit any change in the MIC against ethidium bromide

and norfloxacin although both the compounds had been shown as substrates for these

membrane proteins in earlier studies from this laboratory (Mohanty *et al.*, 2012). Earlier studies

with pBR322 expression system had utilized *E. coli* host KAM32 with deletion of two major efflux

pumps AcrAB and YdhE for functional characterization of the recombinant proteins expressed in

this host (Mohanty *et al.*, 2012). In the current study *E. coli* K-12 strains with deletions in genes

for major RND efflux systems; ΔacrA (deletion of connecting protein), ΔacrB (deletion of its

inner protein) and ΔtolC (deletion of its outer membrane protein) were utilized for functional

characterization of pBAD recombinants (Kuroda & Tsuchiya, 2009, Matsuo *et al.*, 2008, Matsuo

et al., 2007, Otsuka *et al.*, 2005, Mohanty *et al.*, 2012).

Results for MICs of ethidium bromide, ciprofloxacin and norfloxacin showed that though

all the three K-12 deletion strains, K-12 ΔacrA, K-12 ΔacrB and K-12 ΔtolC showed reduction in

MIC as compared to the control wild type strains K-12 and LMG194, K-12 ΔtolC exhibited

maximum hypersusceptibilty *i.e.* at least 16-fold decrease in MIC towards ethidium bromide and

norfloxacin and more than 8-fold decrease in MIC towards ciprofloxacin   (Supplementary Table

1). Subsequently, accumulation assays also revealed that *E. coli* K-12Δ*tolC* had the highest

drug accumulation in comparison to the other hosts utilised in the study (Supplementary Fig.1).

Western blot and confocal microscopy (Fig. 1D and Fig. 2A, panels G to L respectively)

revealed that the recombinant proteins were expressed in K-12Δ*tolC* and localized in its inner

membrane. Taken together, the results of Western blots, confocal microscopy, MIC and

accumulation assays (Supplementary Fig.1) indicated that K-12 Δ*tolC* was a suitable

hypersusceptible strain for expression and functional characterization of the recombinant MATE

pumps.

Table 1. MIC for recombinant clones carrying MATE-type efflux pump genes from *V.*
***fluvialis* and *V. cholerae* in pBAD expression vector and *E. coli* Δ*tolC* host**

	MIC in μg/mL for the recombinant clones				
Drugs/ Dyes	Δ***tolC***	Δ***tolC* /VCH**	Δ***tolC* /VFH**	Δ***tolC* /VCD**	Δ***tolC*** **/VFD**
Ethidium Bromide	<3.9	**15.6**	**15.6**	**3.9**	**7.8**
Norfloxacin	<0.0156	**0.03125-0.0625**	**0.03125-0.0625**	<0.0156	<0.0156
Ciprofloxacin	<0.00156	**0.003125-0.00625**	**0.003125-0.00625**	<0.00156	<0.00156
Nalidixic Acid	0.3125-0.625	0.3125-0.625	0.3125-0.625	0.3125-0.625	0.3125-0.625
Ofloxacin	0.0125-0.025	0.0125-0.025	0.0125-0.025	0.0125-0.025	0.0125-0.025
Neomycin	>200	200	200	200	200
Gentamicin	3.125	1.56-3.125	3.125	1.56	1.56
Acridine Orange	37.5	37.5	37.5	37.5	37.5
Acriflavin	6.25	**12.5**	**12.5**	6.25	6.25
Safranin	3.125	**12.5**	**12.5**	3.125	3.125

215
As kanamycin cassette is inserted in place of the gene of interest (Δ*tolC*) to be disrupted, so MIC against
kanamycin could not be evaluated against pBAD recombinants in Δ*tolC* . Increase in MIC is shown as
bold.
Subsequently, *E. coli* K-12 Δ*tolC* expressing pBAD recombinants for VFH, VCH, VFD

and VCD presented an interesting profile in terms of recognition/transport by these two pumps

(Table 1). H-type pumps exhibited increase in MIC for ethidium bromide (at least 4-fold),

norfloxacin and ciprofloxacin (atleast 2-4-fold), acriflavin (2-fold) and safranin (4-fold). As

against this, D-type pumps conferred resistance only towards ethidium bromide with at least 2-

fold increase in MIC when compared to the control. Both H- and D-type pumps showed almost

similar/slight decrease (2-fold) in MIC for the aminoglycosides (neomycin and gentamicin). Both

H- and D-type pumps showed unaltered MIC for nalidixic acid, ofloxacin and acridine orange

while D-type pumps additionally showed unaltered MIC for norfloxacin, ciprofloxacin, acriflavin

and safranin.

NorM, the prototype member of MATE-type efflux pumps from *V. parahaemolyticus* is a

broad spectrum efflux pump reported to confer resistance towards basic lipophilic compounds

like ethidium bromide and rhodamine, hydrophilic aminoglycosides like kanamycin and

neomycin and hydrophilic zwitterionic fluoroquinolones such as norfloxacin and ciprofloxacin,

but not to hydrophobic quinolones such as sparfloxacin and nalidixic acid (Morita *et al.*, 1998).

In the present study, H-type recombinant seemed to be an efflux pump with primary transport

capability towards zwitterionic fluoroquinolones and basic lipophilic compounds. Earlier studies

had revealed that H-type pumps (VCH/VFH) exhibited resistance towards norfloxacin and

ciprofloxacin only when expressed in pBR322 expression system (Mohanty *et al.*, 2012) but

when these pumps were expressed in pBAD/Δ*tolC* system, they exhibited norfloxacin and

ciprofloxacin resistance along with extended resistance towards ethidium bromide, acriflavin

and safranin. The expression of efflux pumps is generally very low and results in a modest

increase of two-four folds in MIC (Long *et al.*, 2008) which was also observed in the current

study (Long *et al.*, 2008). KetM from *Klebsiella pneumoniae* also exhibited narrow spectrum

resistance to fluoroquinolones with modest change in MIC (Ogawa *et al.*, 2015). Asp32, Glu251

and Asp367 from *V. parahaemolyticus* NorM were implicated in sodium-dependent transport of

antimicrobials and Asp32 specifically is known for its role in efflux thus conferring resistance

towards norfloxacin and ethidium bromide (Otsuka *et al.*, 2005). Even though the efflux pumps

in the present study carried only Asp at position 40/38 in TM1 out of three conserved residues

they were able to exhibit marginal MDR phenotype. This also gives a notion that apart from

these three amino acids, other residues might play important role in transport.   It was later

proven in MATE transporter from the plant *Camelina sativa* (Tanaka *et al.*, 2017). Glutamic

Acid , glutamine, asparagine, arginine along with aspartic acid amino acid residues were

implicated for their role in forming hydrophobic cavity formation and transport in MATE

transporter from the plant *Camelina sativa* (Tanaka *et al.*, 2017).

Accumulation assays revealed that the recombinant strains were able to efflux

drugs/dyes- The substrate transport studies revealed that there was an increased

accumulation of ethidium bromide by pBAD recombinants expressing either H-/D-type efflux

pumps in comparison with control as also observed in earlier studies with their corresponding

pBR322 recombinants (Mohanty *et al.*, 2012) (Fig. 3A and 3B).

Fig. 3. **A/D.** Intracellular ethidium bromide/norfloxacin accumulation in recombinant K-12 ΔtolC
harbouring *vfd* and *vfh* genes in pBAD vector. Samples were collected at every 5 mins for processing
and fluorescence was measured A/**B.** Intracellular ethidium bromide accumulation in recombinant K-12
ΔtolC harbouring *vfd* and *vfh* genes in pBAD vector after 15 mins. D/**E.** Intracellular norfloxacin
accumulation in recombinant K-12 ΔtolC harbouring *vfd* and *vfh* genes in pBAD vector after 15 mins. **C/F.**
Ion dependence of pBAD recombinant efflux pumps in the presence of 100 mM or 1 M sodium chloride
(NaCl) or 100 mM or 1 M potassium chloride (KCl) in K-12 ΔtolC cells transformed with *pBAD-vfh/vfdVFD*.

Data is expressed as mean percentage fluorescence intensity of the maximum fluorescence intensity
obtained during the experiment from three independent experiments and * represents significant
difference to control ($P < 0.05$).

The intracellular ethidium bromide accumulation profiles of both the efflux pumps was in

contrast to the results of MIC assays where elevation in MIC was observed towards ethidium

bromide (Fig. 3A and 3B; Table 2). The absence of correlation between the results of MIC and

accumulation assays particularly in ethidium bromide could result due to the time period of both

the assays.

However, in case of norfloxacin, there was a correlation between MIC assays and

accumulation assays. Results revealed that recombinants with H-type pumps which exhibited

increased MIC towards norfloxacin, showed lowering of intracellular norfloxacin accumulation

whereas recombinants harbouring D-type efflux pumps that exhibited unaltered susceptibility

towards norfloxacin showed almost similar intracellular norfloxacin concentration as compared

to the control (Fig. 3D and 3E). Subsequently ion dependence assays were performed for both

the transporters to understand their fueling mechanism.

The Na$^+$/K$^+$-ions equally facilitated the efflux of norfloxacin by H- pumps - Earlier studies

from our laboratory as well as others had revealed that MATE pumps belonging to NorM cluster

could transport their substrates in the presence of monovalent ions like Na$^+$ (Morita *et al.*, 1998,

Mohanty *et al.*, 2012). The K-12 Δ*tolC/* pBAD-*vfh* recombinants were loaded with norfloxacin in

the presence of reserpine, a compound known to block efflux activity. Subsequently, different

concentrations of 0.1 M and 1 M NaCl or KCl were added to the assay mixture and intracellular

norfloxacin was monitored to assess the effect of these ions on the efflux activity of these

pumps. Results showed that there was an increase in intracellular norfloxacin accumulation in

this recombinant in the presence of reserpine compared to reserpine free control (Fig. 3C and

3F, columns 1 and 6 respectively). Though the addition of 100 mM KCl/NaCl did not

significantly induce the efflux, increase in salt concentration to 1M KCl/NaCl resulted in a

greater efflux of the drug from these cells compared to the cells without any salts (Fig. 3C and

3F, columns 2-5).

Earlier studies on MATE- efflux pumps had established that these efflux pumps were

fueled either by Na^+/H^+-ions for transport (Bhardwaj & Mohanty, 2012) which was also observed

in the current study. There was an increase in intracellular drug concentration in H-

recombinants when the Efflux Pump Inhibitor (EPI) reserpine was added. Effluxing commenced

on addition of Na^+/K^+ ions. Newer studies also point out that the transport capability of MATE-

type *E. coli* ClbM transporter involved in transport of precolibactin was equally influenced by

Na^+, K^+ and Rb^+ ions (Mousa *et al.*, 2017). K^+ ions are known to be congener of Na^+ ions.

Another study revealed that NorM from *V. parahaemolyticus* utilises both PMF and Na^+ for

transport (Jin *et al.*, 2014). Overall it seems both the transporters are Na^+-ion dependent which

has been proved earlier (Mohanty *et al.*, 2012).

Changes in MIC of norfloxacin in the presence of EPIs- Different types of EPIs were utilised

to deduce actual transport mechanism of these efflux pumps.. For this, the MIC of these EPIs

was deduced against host *E. coli* K-12 $\Delta tolC$ and sub-lethal concentrations of EPIs (~50% of

MIC) were utilized to monitor changes in MIC/drug accumulation patterns for recombinant efflux

pumps.

Results revealed that compounds like FCCP and CCCP were effective at low

concentrations to host and with norfloxacin a decrease in MIC was observed for H-type (at least

8-folds) and for D-type (2 folds) Table 2). Both compounds are known to kill living cells at low

concentration as they disrupt PMF, and have been extensively utilised for inhibition of efflux

pumps with concentrations varying for each study (Begum *et al.*, 2005, Morita *et al.*, 2000,

Morita *et al.*, 1998). Other compounds like DNP which deplete ATP or compounds like NIG, NA

and VAL which are known as H^+, Na^+ and K^+ specific ionophores respectively, exhibited high

MIC for host. But with norfloxacin, DNP and NIG also decreased the MIC by at least 2-8-folds

for norfloxacin. against both the compounds (Table 2). VAL and NA did not inhibit both the

Table 2. Change in MICs of norfloxacin in pBAD recombinant clones carrying MATE-type
efflux pump genes from *V. fluvialis* and *V. cholerae* with addition of EPIs

Substrate/EPI	Strain/ clones				
Nor	ΔtolC	ΔtolC /VCH	ΔtolC /VFH	ΔtolC /VCD	ΔtolC /VFD
Nor	<0.0156	0.03125-0.0625	0.03125-0.0625	<0.0156	<0.0156
+CCCP	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**
+FCCP	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**
+RES	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**
+NEM	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**
+DNP	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**
+VAL	0.0625	0.0625	0.0625	0.03125	0.0625
+NIG	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**	**<0.0078**
+NA	0.03125-0.0625	0.03125-0.0625	0.03125	0.0156-0.03125	0.0625

321 Carbonyl cyanide-p-trifluoromethoxyphenyl hydrazone (FCCP), Carbonyl cyanide m-chlorophenyl
322 hydrazone (CCCP), Nigericin (NIG), Valinomycin (VAL), N-ethyl maleidamide (NEM), Dintrophenol
323 (DNP), Reserpine (RES), Sodium Ionophore (NA), Norfloxacin (Nor). The concentration of EPIs used in
324 µg/mL are FCCP 3.125, CCCP 1.56, NIG 50, VAL 50, NEM 62.5, DNP 25, RES 125 and NA 50.
325

pumps for norfloxacin (Table 2). These ionophores mostly increase transferability of ions across

membrane involved in generating EMF, which may or may not be important for energy recycling

(Castro *et al.*, 2015). RES and NEM, both of which either bind or covalently modify the efflux

pumps, also exhibited very high MIC for *E. coli* K-12 ΔtolC but in conjunction with norfloxacin,

both compounds decreased the MIC (Table 2). Both NEM and RES decreased the MIC

exhibited by both pumps by at least 2-16-folds whereas (Table 2). It was not surprising that

reserpine increased sensitivity towards fluoroquinolones. Resperine had been earlier reported to

inhibit various efflux pumps resulting in increased susceptibility (Baranova & Neyfakh, 1997,

Brenwald *et al.*, 1997, Aeschlimann *et al.*, 1999, Markham, 1999, Markham *et al.*, 1999,

Gibbons & Udo, 2000, Mullin *et al.*, 2004).

Accumulation of norfloxacin by H-/D-type pumps in the presence of EPIs- The MIC assays

had established that some of the EPIs were active in reducing the MIC of norfloxacin for the

recombinants expressing efflux pumps. Therefore, norfloxacin accumulation studies were

performed to assess the real time effect on drug transport by the H-/D- recombinants in the

presence/absence of EPIs.

Results revealed that *E. coli* K-12 $\Delta tolC$ expressing H-type pumps showed a marginal to

moderate degree of increased norfloxacin accumulation in presence of CCCP and FCCP in

comparison to EPI free control (Fig. 4). These pumps exhibited almost unaltered/marginal

increased accumulation of norfloxacin in presence of NIG, NEM, VAL and DNP (Fig. 4).

Interestingly, both efflux pumps exhibited very high accumulation of norfloxacin in presence of

RES (Fig. 4). MIC studies of norfloxacin against both recombinants in the presence of EPIs had

revealed that compounds like CCCP, RES and FCCP decreased the MICs (Table 4). This was

in sync to the norfloxacin accumulation results, where these compounds increased the

accumulation of norfloxacin. NA increased the intracellular norfloxacin for H-type of efflux which

was in contrast to MIC assays where it did not decrease the MIC for norfloxacin (Table 2).

However, for NEM, DNP and NIG this correlation between MIC and accumulation results

was missing probably due to the kinetics of inhibition by these EPIs and the difference in the

time course for the two type of assays. MATE family of efflux pumps utilise Na^+ ions for

transport which was observed in the current study. There was an increased intracellular

norfloxacin accumulation with utilisation of H^+/ Na^+/K^+ ion specific ionophores such as NA,

CCCP and FCCP. This was in corroboration with our earlier observation in ion dependence

experiments where we observed that both the efflux pumps showed greater efflux in presence

of both Na^+ and K^+ ions also evidenced in MIC experiments (Table 2) where these EPIs

decreased the MIC of norfloxacin. Taken together, there was a relative correlation between drug

accumulation, ion dependence and change in MIC patterns in case of some selected EPIs.

Fig. 4. Accumulation of norfloxacin in presence of various EPIs. Accumulation of norfloxacin by *E. coli* K-12 Δ*tolC* expressing H- and D-type pumps in the presence of EPIs. Recombinant *E. coli* K-12 ΔtolC was loaded with norfloxacin and incubated at 37°C for 15 mins in presence/absence of EPIs. These samples were subsequently processed and analyzed for fluorescence intensity as described in the materials and methods. * represents significant difference to control ($P < 0.05$). The concentration of EPIs used in µg/mL are FCCP 3.125, CCCP 1.56, NIG 50, VAL 50, NEM 62.5, DNP 25, RES 125 and NA 50.

Molecular Modeling and docking- Structural prediction of MATE-type efflux pumps had suggested 12 transmembrane helices in their structure which was further validated with the structure of first Na$^+$ dependent MATE-type efflux pump from *Pyrococcus furiosus* having 12 transmembranes (Lomovskaya & Bostian, 2006, Tanaka *et al.*, 2013). In our earlier attempt in predicting 3D structures of VFH and VFD using I-TASSER server, we had reported that VFD had 11 transmembranes (TMs) whereas VFH had 10 TMs (Mohanty *et al.*, 2012). After 2012, more structures were solved for the members of this family of transporters (Tanaka *et al.*, 2013, He *et al.*, 2010, Lu *et al.*, 2013b, Lu *et al.*, 2013a). Subsequent to these studies, further attempts for protein modeling were made using SwissModel (https://swissmodel.expasy.org/).to

380 understand the topology and mechanism of efflux by these two MATE pumps Both PyMol

381 (https://pymol.org/2/) and Chimera (https://www.cgl.ucsf.edu/chimera/) were used

382 forvisualization of their molecular structures. Results revealed that both the efflux pumps had 10

383 predicted TMs arranged in two bundles of 5 TMs each (Fig. 5.) similar to crystal structures of

384 earlier published MATE transporters (Lu *et al.*, 2013b).

Fig. 5. Docked pose of ethidium bromide and Norfloxacin with *V. fluvialis* MATE efflux pumps (VFH and VFD) forming covalent complex at Asp 38/40 of 1st transmembrane.

They were also outward facing like NorM (Lu *et al.*, 2013b) with distinct internal cavity. It was

both in corroboration (for VFH) and contrast (for VFD) to our earlier finding where VFH model

was similar to NorM and VFD had different predicted protein model (Mohanty *et al.*, 2012).

Subsequently, molecular docking was done using SwissDock (http://www.swissdock.ch/) to

validate binding interactions of both the transporters with ethidium bromide and norfloxacin as

both the transporters showed certain degree of transport for these substrates. Classically, a

MATE pump has 12 TMs arranged in two bundles of six TMs each (Tanaka *et al.*, 2013,

Tanaka *et al.*, 2017, Radchenko *et al.*, 2016, Lu *et al.*, 2013b).

Table 3 **Ligand-docking of VFH and VFD efflux pumps with various substrates**

Protein	Substrate	Orientation	Total Binding sites	Total Hydrogen Bonding	Most plausible binding site
VFH	Nor	Outward	35 clusters with 256 binding sites	11 clusters with 65 sites	1 cluster with 6 binding sites (deltag -8.4)
VFH	EtBr	Outward	30 clusters with 256 binding sites	10 clusters with 69 sites	3 clusters with 20 binding sites (deltag -6.9)
VFD	Nor	Outward	35 clusters with 256 binding sites	23 clusters with 139 sites	1 cluster with 8 binding sites (deltag -7.4 to -7.5)
VFD	EtBr	Outward	34 clusters with 256 binding sites	15 clusters with 80 sites	2 clusters with 10 binding sites (deltag -7.19)

The first TM has an ion binding site represented by D32, the Aspartic Acid residue at 32

position (Otsuka *et al.*, 2005, Long *et al.*, 2008, Lu *et al.*, 2013a, Lu *et al.*, 2013b). With this

input, the protein sequences of both the transporters were submitted to SwissDock ≈server

(http://www.swissdock.ch/). The docking was programmed for finding all possible ligand

interactions. All aspartic acid residues were first identified from the generated models and

hydrogen bonds were calculated. The details have been presented in Table 3 and Fig 5.

Interaction between the protein and ligands was optimum in the first TM at D40 for VFH and

D38 for VFD which was similar to earlier published findings (Tanaka *et al.*, 2013, He *et al.*,

2010, Lu *et al.*, 2013b, Lu *et al.*, 2013a). This also indicated similar transport mechanism for

both the transporters. Both the compounds ethidium bromide and norfloxacin fitted well in the

central cavity of both the transporters. The only noticed difference was that while norfloxacin

fitted in a straight fashion forming bonds with D40/38, Ethidium bromide showed bent fitting from

1st to 7th TM (Fig. 5).

Ion binding and Electrostatic potential surface representation- MATE-type efflux pumps are

secondary pumps that utilize Na$^+$/H$^+$ for transport (Jin *et al.*, 2014, Mohanty *et al.*, 2012). Earlier

studies had proved that these were Na$^+$ ion-driven transporters (Mohanty *et al.*, 2012). In the

ion-dependence assays, we observed the transport capability was equally influenced by both

sodium and potassium ions for VFH (reference). For better understanding of ion dependence

and   amino   acid   residues   involved,   we   utilized   IonCom   server

(https://zhanglab.ccmb.med.umich.edu/IonCom/) to predict their ion dependence and amino

acid residues involved.

Results revealed that both the efflux pumps had no binding site detected for the

following ions:(Cu, Fe, Ca, Mg, Mn, Na, K, CO, NO$_2$, SO$_4$, PO$_4$). The predicted ion binding

amino acid residues were D40, E51, Q82, D88, Y138, H163, N179, N309 and D393 for VFH

and D38, H44, W49, Q81, N163 and D204 for VFD. Interestingly, docking studies clearly

indicated that D38/40 of both the transporters were involved in binding. So, even though we

couldn't predict dependence of ion but D38/40 seems most plausible ion binding site for both

the transporters which was similar to NorM from *V. cholerae* (He *et al.*, 2010).

Electrostatic potential of proteins is caused by charged amino acid residues in side

chains, bound or unbound by various ions and play various roles such as protein folding and

stability and function of the proteins. APBS pluggin in PyMol was used to calculate the

electrostatic potential for both the proteins. It may reflects secondary and tertiary structures of

transporters and their affinity to bind to substrates. Results revealed that both VFH and VFD

had different hydrophobicity patches similar to other known MATE transporters (Fig 6a) (Zheng

et al., 2018). Both VFH and VFD formed a central cavity surrounded by charged electronegative

amino acids. VFH also had an electropositive centre at the end of the cavity (Fig 6a).

Subsequently, VFH and VFD with all possible interaction with norfloxacin via docking were

mapped along and electrostatic potential was calculated, an interesting scenario came out. A

sizable norfloxacin molecule was assembled in the cavity of VFH whereas fewer norfloxacin

molecules were assembled in the cavity Of VFD. NorM which was taken as control had

maximum norfloxacin molecules assembled in the cavity (Fig 6c). This was in sync with our

earlier observation where VFH exhibited broader transport capability than VFD in MIC. The

arrangement was mostly concentrated towards n-lobe which was similar like NorM (Lu *et al.*,

2013a)

Fig. 6. Surface charge differences between VFH, VFD and NorM-VC. Surface electrostatics were calculated by the APBS plugin in PyMOL. Cross-section model of VFH, VFD and NorM-VC surface representation is shown from the membrane region (a) or from the extracellular (b) sides (c) with norfloxacin and (d) Cross-section model of VFH/VFD exposing covalent interaction between TM1 and TM7. C represents cavity. Red arrows represent cavity.

It has been widely accepted that MATE transporters evolved from genetic duplication of flippases (Lomovskaya & Bostian, 2006, Tanaka *et al.*, 2013). Comparison of both the transporters with known structures of MATE transporters from *Camelina sativa,* NorM from *Vibrio cholarae* and MurJ flippases from *Thermosipho africanus* reveals interesting similarities and divergence from both VFH and VFD. D38/40 of both the transporters seemed to more conserved in bacterial transporters (Lu *et al.*, 2013b, He *et al.*, 2010) in comparison to their eukaryotic counterparts (Tanaka *et al.*, 2017). When both these transporters were compared to their evolutionary predecessors like MurJ (Kuk *et al.*, 2017), the difference is quite significant.

MurJ flippases from *Thermosipho africanus* exhibited presence of 14 TMs arranged in 6+6

similar to bacterial transporters and functional role of last two TMs is unknown. Ser17, Arg18,

Arg24, Arg52, and Arg255 in 1st Tm of Murj has been proved important for recognizing the

diphosphate and/or sugar moieties of lipid II whereas Asp in 1st Tm of NorM has been proved for

ion dependence and substrate transport (He *et al.*, 2010). It seems both VFH and VFD has

similar topology like NorM with presence of D38/40 in 1st TM creating a hydrophobic cavity for

substrate recognition and transport.

During mapping of electrostatic potential of VFH and VFD, an aberrant architecture was

also observed in both the transporters. MATE transporters usually have 12 TMs arranged in two

bundles with 6 TMS each. Each forms one lobe and are usually free. The 1st TM upon

protonation displaces to 7th TM to facilitate transport. But in case of both VFH/VFD, the loop

connecting 1st TM to 2nd TM forms a covalent structure with loop of 7th TM connecting 8th TM as

elucidated by surface model (Fig 6d). MATE type transporters are thought to be evolved of

internal genetic duplication, resulting in two lobes. VFH/VFD had 10 TMs, still there was greater

scope for transport. The formation of covalent bond reduces mobility of 1st TM to be displaced

to 6th TM and may overall hinder transport efficiency (Fig 7b).

Mechanism of inhibition of both the transporters by reserpine- Our earlier observation with

electrostatic mapping and inhibition of both the transporters with various EPIs had led to an

indication that the transport of norfloxacin was indeed hampered in presence of these EPIs.

Surprisingly, the similar effect couldn't be observed in case of drug transport assay. Compounds

like CCCP/FCCP, which are extremely toxic to cell at low concentration showed marginal

inhibition. Reserpine which was fairly non-toxic, was able to inhibit both the transporters in real

time drug transport assays. In our earlier study, we had observed 20 microgram/mL was

sufficient to inhibit both the transporters (Mohanty *et al.*, 2012). Even in the current study, the

MIC against the host was fairly high i.e >128 mcg/mL. So, range of concentration was quite

enough to inhibit. Even thoughit has been earlier proved in case of Bmr transporters, where

certain residues were implicated for inhibition (Klyachko *et al.*, 1997), no direct structural studies

have ever proved the actual inhibition by reserpine in MATE efflux pump. To investigate the

exact nature, we tried to dock reserpine with both the transporters along with NorM from *V.*

cholerae.

Results revealed that reserpine could conveniently fit into electronegative cavities of

both the transporters. VFH had 3 binding clusters with 19 binding sites whereas VFD had 2

binding clusters with 15 binding sites. Interestingly, NorM had 26 binding clusters with 190

binding sites.

In some of the earlier studies, MATE transporters have been co-crystalized along with

the inhibitors. (Tanaka *et al.*, 2013) . These studies have delineated the role of hydrogen

bonding between transportersand ligand for exact inhibition mechanism. Our own observation

with NorMhad hinted the same as there were almost 15 clusters and 96 binding sites forming

hydrogen bonds in the cavity of NorM. But in case of VFH there was no predicted hydrogen

bond formation and for VFD, 1 cluster and 3 binding site configuration was observed. Previous

studies regarding co-crystallization of MATE-transporters with certain inhibitors had shed light

on formation of electrostatic bonds for inhibition with verapamil and macrocyclic peptides

(Radchenko *et al.*, 2015, Tanaka *et al.*, 2013). Observation from our earlier studies and current

study reflect that reserpine acts pretty well at different concentrations also.

Fig. 7. A. Docked pose of reserpine with *V. fluvialis* MATE efflux pumps (VFH and VFD) . **B.** Proposed mechanism of transport of NorM-VC and VFD/VFH. Contrasting to NorM-VC, VFH/VFD forms covalent interactions between TM1 and TM7, which along low cavity volume leads to low transport capability of VFH/VFD.

The exact mechanism of reserpine inhibition is binding to phenylalanine, tryptophan and

tyrosine residues of Bmr efflux pumps (Steinfels *et al.*, 2004) and inhibition by reserpine does

not involve ion bioenergetics of the cell and seems to be a non-specific inhibition of efflux

pumps as was also seen in the current study. This led us to believe three important aspects of

transport a. mobility of each lobe plays a vital role in transport b. size and electronegativity of

central cavity plays important role for actual substrate recognition c. binding of ligand in

electronegative pocket is not always electrostatic but may be covalent as reserpine was able to

inhibit without actual hydrogen bond formation in both the transporters.

Experimental procedures

Cloning in pBAD-TOPO-TA vector- The primers used for cloning in pBAD vectors were VFH-

BAD-F/VFH-BAD-R and VFD-BAD-F/VFD-BAD-R mentioned in supplementary table 2. The

amplicons were subsequently purified using PCR purification kit (Qiagen) and ligated in pBAD

vector (Invitrogen) according to manufacturer's protocol.

Expression and detection of recombinants harbouring pBAD efflux pump genes- pBAD

clones for expression of VCH, VFH, VCD and *VFD* (where D- and H- are two MATE-type efflux

pumps, VC stands for *V. cholerae* and VF stands for *V. fluvialis*) and Lac-His (positive control)

were transformed in *E. coli* LMG194 and *E. coli* K-12 Δ *tolC* (JW5503-1, *E. coli* Genetic

Resource Centre, Yale University) for expression as well as functional characterization. For

Western blots, *E. coli* LMG194/K12 Δ*tolC* cells transformed with the recombinant vectors were

grown in Luria-Bertani (LB) medium to an optical density of 0.5 at 600 nm (OD_{600}) and induced

with 0.2% L-arabinose for 4 hours and the total cell lysates prepared in SDS-PAGE lysis buffer

were electrophoresed on 8% SDS-PAGE. Western blot was done with 0.45 micron

nitrocellulose membrane (Whatman/Pall Life sciences) and blocked with 5% fat free milk

powder in Tris buffered saline with 0.05% tween 20 (TBST) for 2 hours at room temperature.

The blot was incubated with goat Anti-His/V_5 primary antibody at 1:5000 dilutions in 0.5% milk

powder in TBST for 2 hours at room temperature with constant shaking followed by washing in

TBST and incubation with anti-Goat IgG-HRP (Horseradish Peroxidase)/Anti-Goat IgG-AP

(Alkaline Phosphatase) (Jackson Immunoresearch) secondary antibody. The blots were

developed either by using Novex ECL HRP chemiluminiscent development kit (Invitrogen) or by

colorimetry using DAB/H_2O_2 (for HRP conjugated secondary antibody) or BCIP/NBT

(Calbiochem) (for AP conjugated secondary antibody) according to the manufacturer's

instructions. For chemiluminiscence, the blot was dried and developed on X-ray.

Detection of expression of recombinant proteins by confocal imaging- Briefly, the cells

were grown and induced as discussed in earlier subsection of Experimental Procedures. 200 µL

of induced cells were centrifuged at 5000 rpm for 5 min, the cells were resuspended in 50 µL of

supernatant and were spotted on coverslips coated with 0.2% L-Poly-lysine (Sigma) in 24-well

tissue culture plates (Corning) for 30 min at room temperature. Cells were then fixed using 4%

paraformaldehyde (PFA) for 15 min followed by washing twice with 250 µL PBST (PBS

containing 0.05% Tween 20) for 5 min without shaking. The cells were blocked with 5% Bovine

Serum Albumin (BSA) in PBST for 1 hour at room temperature followed by washing thrice with

500 µL PBST for 10 min each and solutions were aspirated gently using a pipette. 50 µL mouse

monoclonal anti-6X His tagged primary antibody (Calbiochem) was added at a dilution of 1:150

in 0.5 % BSA in PBST in the respective wells and incubated for 2 hours without shaking at room

temperature. Washing was done thrice as described above. Goat anti-mouse secondary

antibody labeled with Cy3 (Invitrogen) was added at the dilution of 1: 50 in 0.5 % BSA in PBST

and incubated for 2 hours at room temperature followed by one washing step. Cells were

counterstained with 200 µL of 4', 6-diamidino-2-phenylindole (DAPI) at 1 µg mL^{-1} concentration

and 50 µL of Film Tracer 1-43 membrane marker (Invitrogen) at 1 µg mL-1 concentration in

DMSO. Confocal imaging was carried out using CTR 6500 confocal microscope (Leica). Diode

lamp was utilised for detection of DAPI, UV HeNe lamp for CY3 and Argon lamp was used to

detect membrane marker.

Preparation of Membrane fraction- The membrane fraction was prepared according to the

published protocol (Leyh, 1990). The *E. coli* LMG194/ K12 Δ*tolC* recombinant cells were grown

and induced as described earlier. 25 mL of induced cultures were harvested at 5000 rpm, 4°C

for 15 min. The supernatant was discarded and the pellet was washed with 5 mL ice cold PBS

(phosphate-buffered saline) followed by centrifugation at 5000 rpm, 4°C for 10 min. The cells

were resuspended in 1 mL lysis buffer (50 mM potassium phosphate, pH 7.8) having 1 mM

phenylmethanesulfonyl fluoride and incubated in -80°C deep freezer for 2 hours. The samples

were then thawed and sonicated 4 times with 30 sec on and off pulse on ice. The samples were

then centrifuged at 8000 rpm, 4°C for 5 min. The supernatants were saved as soluble fraction

and the pellet was treated with 200 µL of 2% sarcosyl solution (Sigma Aldrich) and incubated at

room temperature for 30 min. The sample was then centrifuged at 15,000 rpm for 30 min at

room temperature to obtain inner membrane proteins as supernatant and the rest of the proteins

as pellet. Samples were subsequently subjected to Western blot for detection of recombinant

proteins.

Minimum Inhibitory Concentration (MIC) assays- The pBAD expression clones *of vfh*, *vch*,

vcd and *vfd* were transformed in *E. coli* LMG194/K-12 $\Delta tolC$ cells, grown in LB broth containing

ampicillin (100 µg mL^{-1}) to an OD$_{600}$ of 0.5 and induced with 0.2% L-arabinose for 4 hours.

Kanamycin (30 µg mL^{-1}) was additionally used for *E. coli* K-12 $\Delta tolC$ cells. Drugs/dyes were

serially double diluted in LB broth containing 0.2% L-arabinose and 2 mL dilution/well were

dispensed in 24 well tissue culture plates (Corning). OD$_{600}$ of the induced sample was adjusted

to 0.08-0.13 and 50 µL of this sample was added to each well. Incubation was done at 37°C.

Readings were observed visually after 16 hours. The assays were repeated at least three times.

Drug/Dye accumulation assays- For *E. coli* K-12ΔtolC cells harboring either pBAD-*vfh* or

pBAD-*vfd* or not harbouring any plasmids were grown and induced as described earlier in

Materials and methods. Ethidium bromide (1.9 µg mL^{-1}) and norfloxacin (0.0156 µg mL^{-1}) were

used for accumulation assays. This was followed by single washing with 0.1 M Tris.HCl, pH 7.0

and resuspension in equal volume of the same buffer. The samples were collected every 15

min. The processing of the cells and monitoring of the fluorescent content was carried out

exactly as described earlier (Mohanty *et al.*, 2012).

Effect of Na$^+$/K$^+$ ions on efflux of drugs/dyes- *E. coli* K-12$\Delta tolC$ cells harboring either *pBAD-*

vfh or *pBAD-vfd* or not harbouring any plasmids were grown in LB broth containing kanamycin

(30 µg mL^{-1}) and induced with 0.02% L-arabinose for 4 hours. Ampicillin was only added to the

cells harbouring recombinant plasmids. The recombinants were grown and processed as

described above in this section. The samples were then harvested by centrifugation, washed

thrice with buffer containing 0.1M Tris.HCl, pH 7.0, and suspended in the same buffer to an

OD_{600} of 1.0.   The drug/dyes were used at 50% of the MIC values for transport studies.

Norfloxacin at a final concentration of 0.0156 µg mL^{-1} (50% MIC concentration)along with 125

µg mL^{-1} of reserpine was added to the cell suspension and incubated for 30 min at 37℃. After

30 min, 1 mL samples were collected and stored in prechilled microcentrifuge tubes. Negative

control (without drug/dyes) and EPI free control were also included in the study.   Appropriate

volume of 5 M and 0.5 M NaCl/KCl stock solution were added to assay mixture to a final

concentration of 1M and 100 mM, incubated at 37℃ for 15 minutes and 1 mL sample was

collected. The volume was kept constant irrespective of salt concentration. Samples were then

centrifuged at 10,000 rpm for 5 min at 4℃, washed once with 0.1 M Tris.HCl, pH 7.0 buffer and

resuspended in 1 mL of 100 mM glycine.HCl, pH 3.0. These samples were subsequently

processed and analyzed for fluorescence intensity as described in the previous sections. The

maximum fluorescence observed in the experiment was considered to be 100%.

Assays to determine efficacy of various EPIs on recombinant efflux pumps- The MIC of

various EPIs was deduced against *E. coli* K-12 Δ*tolC*.Different concentrations of EPIs were

prepared by two-fold dilution method in LB broth containing kanamycin (30 µg mL^{-1}) and 100

µL/well was dispensed in 96-well tissue culture plates (Corning Inc.). The EPIs were procured

from Sigma Aldrich. *E. coli* K-12 Δ*tolC* was grown in LB containing kanamycin (30 µg mL^{-1}) to

an OD_{600} of 0.08-0.13.   Five microlitre of cells were inoculated in 100 µL medium containing

various concentrations of EPIs in a 96 well plate. Observations were made visually after 16

hours. The experiments were repeated at least thrice. 50% concentration of MIC of each EPI

was utilised in combination with various concentrations of norfloxacin. The *E. coli* K-12 Δ*tolC*

carrying recombinant efflux pump plasmids and control *E. coli* K-12 Δ*tolC* were each grown in

LB with kanamycin (30 µg mL^{-1}) and ampicillin (100 µg mL^{-1}). Overnight primary cultures were

done in same way. Then 10 mL of secondary culture was prepared. Upon reaching OD_{600} of 0.5,

the cells were induced with 0.02% L-arabinose for 4 hours with 200 rpm shaking at 37°C. The

optical density of induced cells were adjusted to 0.08-0.13. Five microlitre of cells were

inoculated in 100 µL medium containing 50% concentration of various EPIs along with various

concentrations of norfloxacin in a 96 well plate (Corning Inc). Observations were made visually

after 16 hours. The experiments were repeated at least thrice. The change in MIC was noted to

observe efficacy of each EPI.

Acknowledgments: The authors thank Indian Council of Medical Research, New Delhi for

providing Senior Research Fellowship to PM (F/815/2011/ECD-II), Department of

Biotechnology, New Delhi for Junior Research Fellowship (BT/PR/ 11634 INF/22/104/2008) of

AS, Dr. A.K. Tiwari, Department of Developmental Biology, IIAR for helping in confocal imaging

and the Puri Foundation for Education in India for support.

Author contributions: Conceived and designed the experiments: AKB and PM. Performed the

experiments: PM AS AKB. Analyzed the data: AKB PM. Contributed reagents/materials/analysis

tools: AKB. Wrote the paper: AKB PM.

Conflict of interest: The authors declare that they have no conflicts of interest.

Reference

Aeschlimann, J.R., L.D. Dresser, G.W. Kaatz & M.J. Rybak, (1999) Effects of NorA inhibitors on
636   in vitro antibacterial activities and postantibiotic effects of levofloxacin, ciprofloxacin, and
637   norfloxacin in genetically related strains of Staphylococcus aureus. *Antimicrob Agents*
638   *Chemother* **43**: 335-340.
639   Baranova, N.N. & A.A. Neyfakh, (1997) Apparent involvement of a multidrug transporter in the
640   fluoroquinolone resistance of Streptococcus pneumoniae. *Antimicrob Agents Chemother*
641   **41**: 1396-1398.
642   Begum, A., M.M. Rahman, W. Ogawa, T. Mizushima, T. Kuroda & T. Tsuchiya, (2005) Gene
643   cloning and characterization of four MATE family multidrug efflux pumps from Vibrio
644   cholerae non-O1. *Microbiol Immunol* **49**: 949-957.
645   Bhardwaj, A. & P. Mohanty, (2012) Bacterial efflux pumps involved in multidrug resistance and
646   their inhibitors: rejuvinating the antimicrobial chemotherapy. *Recent patents on anti-*
647   *infective drug discovery* **7**: 73-89.
648   Blakey, D., A. Leech, G.H. Thomas, G. Coutts, K. Findlay & M. Merrick, (2002) Purification of
649   the Escherichia coli ammonium transporter AmtB reveals a trimeric stoichiometry.
650   *Biochem J* **364**: 527-535.

Brenwald, N.P., M.J. Gill & R. Wise, (1997) The effect of reserpine, an inhibitor of multi-drug
efflux pumps, on the in-vitro susceptibilities of fluoroquinolone-resistant strains of
Streptococcus pneumoniae to norfloxacin. *The Journal of antimicrobial chemotherapy*
40: 458-460.
Castro, P.J., A.F. Silva, B.C. Marreiros, A.P. Batista & M.M. Pereira, (2015) Respiratory
complex I: A dual relation with H and Na? *Biochim Biophys Acta*.
Frankel, S., R. Sohn & L. Leinwand, (1991) The use of sarkosyl in generating soluble protein
after bacterial expression. *Proc Natl Acad Sci U S A* **88**: 1192-1196.
Gibbons, S. & E.E. Udo, (2000) The effect of reserpine, a modulator of multidrug efflux pumps,
on the in vitro activity of tetracycline against clinical isolates of methicillin resistant
Staphylococcus aureus (MRSA) possessing the tet(K) determinant. *Phytotherapy*
research : PTR **14**: 139-140.
Goswami, P., C. Paulino, D. Hizlan, J. Vonck, O. Yildiz & W. Kuhlbrandt, (2011) Structure of the
archaeal Na+/H+ antiporter NhaP1 and functional role of transmembrane helix 1. *Embo*
J **30**: 439-449.
Hassan, K.A., Q. Liu, P.J. Henderson & I.T. Paulsen, (2015) Homologs of the Acinetobacter
baumannii AceI transporter represent a new family of bacterial multidrug efflux systems.
MBio **6**.
He, G.X., T. Kuroda, T. Mima, Y. Morita, T. Mizushima & T. Tsuchiya, (2004) An H(+)-coupled
multidrug efflux pump, PmpM, a member of the MATE family of transporters, from
Pseudomonas aeruginosa. *J Bacteriol* **186**: 262-265.
He, X., P. Szewczyk, A. Karyakin, M. Evin, W.X. Hong, Q. Zhang & G. Chang, (2010) Structure
of a cation-bound multidrug and toxic compound extrusion transporter. *Nature* **467**: 991-
994.
Hobb, R.I., J.A. Fields, C.M. Burns & S.A. Thompson, (2009) Evaluation of procedures for outer
membrane isolation from Campylobacter jejuni. *Microbiology* **155**: 979-988.
Jin, Y., A. Nair & H.W. van Veen, (2014) Multidrug transport protein norM from vibrio cholerae
simultaneously couples to sodium- and proton-motive force. *J Biol Chem* **289**: 14624-
14632.
Klyachko, K.A., S. Schuldiner & A.A. Neyfakh, (1997) Mutations affecting substrate specificity of
the Bacillus subtilis multidrug transporter Bmr. *Journal of bacteriology* **179**: 2189-2193.
Kuk, A.C., E.H. Mashalidis & S.Y. Lee, (2017) Crystal structure of the MOP flippase MurJ in an
inward-facing conformation. *Nat Struct Mol Biol* **24**: 171-176.
Kuroda, T. & T. Tsuchiya, (2009) Multidrug efflux transporters in the MATE family. *Biochim*
Biophys Acta **1794**: 763-768.
Leyh, R.D., (1990) Characterization of the outer membrane proteins of Bordetella avium. In:
Veterinary Microbiology and Preventive Medicine. Ames: Iowa State University, pp. 115.
Lomovskaya, O. & K.A. Bostian, (2006) Practical applications and feasibility of efflux pump
inhibitors in the clinic--a vision for applied use. *Biochemical pharmacology* **71**: 910-918.
Long, F., C. Rouquette-Loughlin, W.M. Shafer & E.W. Yu, (2008) Functional cloning and
characterization of the multidrug efflux pumps NorM from Neisseria gonorrhoeae and
YdhE from Escherichia coli. *Antimicrob Agents Chemother* **52**: 3052-3060.
Lu, M., M. Radchenko, J. Symersky, R. Nie & Y. Guo, (2013a) Structural insights into H+-
coupled multidrug extrusion by a MATE transporter. *Nat Struct Mol Biol* **20**: 1310-1317.
Lu, M., J. Symersky, M. Radchenko, A. Koide, Y. Guo, R. Nie & S. Koide, (2013b) Structures of
a Na+-coupled, substrate-bound MATE multidrug transporter. *Proc Natl Acad Sci U S A*
110: 2099-2104.
Markham, P.N., (1999) Inhibition of the emergence of ciprofloxacin resistance in Streptococcus
pneumoniae by the multidrug efflux inhibitor reserpine. *Antimicrob Agents Chemother*
43: 988-989.

Markham, P.N., E. Westhaus, K. Klyachko, M.E. Johnson & A.A. Neyfakh, (1999) Multiple novel inhibitors of the NorA multidrug transporter of Staphylococcus aureus. *Antimicrobial agents and chemotherapy* **43**: 2404-2408.

Matsuo, T., J. Chen, Y. Minato, W. Ogawa, T. Mizushima, T. Kuroda & T. Tsuchiya, (2008) SmdAB, a heterodimeric ABC-Type multidrug efflux pump, in Serratia marcescens. *J Bacteriol* **190**: 648-654.

Matsuo, T., K. Hayashi, Y. Morita, M. Koterasawa, W. Ogawa, T. Mizushima, T. Tsuchiya & T. Kuroda, (2007) VmeAB, an RND-type multidrug efflux transporter in Vibrio parahaemolyticus. *Microbiology* **153**: 4129-4137.

Meeske, A.J., L.T. Sham, H. Kimsey, B.M. Koo, C.A. Gross, T.G. Bernhardt & D.Z. Rudner, (2015) MurJ and a novel lipid II flippase are required for cell wall biogenesis in Bacillus subtilis. *Proc Natl Acad Sci U S A* **112**: 6437-6442.

Miyauchi, H., S. Moriyama, T. Kusakizako, K. Kumazaki, T. Nakane, K. Yamashita, K. Hirata, N. Dohmae, T. Nishizawa, K. Ito, T. Miyaji, Y. Moriyama, R. Ishitani & O. Nureki, (2017) Structural basis for xenobiotic extrusion by eukaryotic MATE transporter. *Nat Commun* **8**: 1633.

Mohanty, P., A. Patel & A. Kushwaha Bhardwaj, (2012) Role of H- and D- MATE-type transporters from multidrug resistant clinical isolates of Vibrio fluvialis in conferring fluoroquinolone resistance. *PloS one* **7**: e35752.

Morita, Y., A. Kataoka, S. Shiota, T. Mizushima & T. Tsuchiya, (2000) NorM of vibrio parahaemolyticus is an Na(+)-driven multidrug efflux pump. *J Bacteriol* **182**: 6694-6697.

Morita, Y., K. Kodama, S. Shiota, T. Mine, A. Kataoka, T. Mizushima & T. Tsuchiya, (1998) NorM, a putative multidrug efflux protein, of Vibrio parahaemolyticus and its homolog in Escherichia coli. *Antimicrob Agents Chemother* **42**: 1778-1782.

Mousa, J.J., R.C. Newsome, Y. Yang, C. Jobin & S.D. Bruner, (2017) ClbM is a versatile, cation-promiscuous MATE transporter found in the colibactin biosynthetic gene cluster. *Biochem Biophys Res Commun* **482**: 1233-1239.

Mullin, S., N. Mani & T.H. Grossman, (2004) Inhibition of antibiotic efflux in bacteria by the novel multidrug resistance inhibitors biricodar (VX-710) and timcodar (VX-853). *Antimicrob Agents Chemother* **48**: 4171-4176.

Ogawa, W., Y. Minato, H. Dodan, M. Onishi, T. Tsuchiya & T. Kuroda, (2015) Characterization of MATE-Type Multidrug Efflux Pumps from Klebsiella pneumoniae MGH78578. *PloS one* **10**: e0121619.

Omote, H., M. Hiasa, T. Matsumoto, M. Otsuka & Y. Moriyama, (2006) The MATE proteins as fundamental transporters of metabolic and xenobiotic organic cations. *Trends Pharmacol Sci* **27**: 587-593.

Otsuka, M., M. Yasuda, Y. Morita, C. Otsuka, T. Tsuchiya, H. Omote & Y. Moriyama, (2005) Identification of essential amino acid residues of the NorM Na+/multidrug antiporter in Vibrio parahaemolyticus. *Journal of bacteriology* **187**: 1552-1558.

Poole, K., (2000) Efflux-mediated resistance to fluoroquinolones in gram-negative bacteria. *Antimicrob Agents Chemother* **44**: 2233-2241.

Poole, K., (2004) Efflux-mediated multiresistance in Gram-negative bacteria. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases* **10**: 12-26.

Poole, K., (2005) Efflux-mediated antimicrobial resistance. *The Journal of antimicrobial chemotherapy* **56**: 20-51.

Radchenko, M., R. Nie & M. Lu, (2016) Disulfide Cross-linking of a Multidrug and Toxic Compound Extrusion Transporter Impacts Multidrug Efflux. *J Biol Chem* **291**: 9818-9826.

Radchenko, M., J. Symersky, R. Nie & M. Lu, (2015) Structural basis for the blockade of MATE multidrug efflux pumps. *Nat Commun* **6**: 7995.

Steinfels, E., C. Orelle, J.R. Fantino, O. Dalmas, J.L. Rigaud, F. Denizot, A. Di Pietro & J.M.
Jault, (2004) Characterization of YvcC (BmrA), a multidrug ABC transporter
constitutively expressed in Bacillus subtilis. *Biochemistry* **43**: 7491-7502.
Tanaka, Y., C.J. Hipolito, A.D. Maturana, K. Ito, T. Kuroda, T. Higuchi, T. Katoh, H.E. Kato, M.
Hattori, K. Kumazaki, T. Tsukazaki, R. Ishitani, H. Suga & O. Nureki, (2013) Structural
basis for the drug extrusion mechanism by a MATE multidrug transporter. *Nature* **496**:
247-251.
Tanaka, Y., S. Iwaki & T. Tsukazaki, (2017) Crystal Structure of a Plant Multidrug and Toxic
Compound Extrusion Family Protein. *Structure* **25**: 1455-1460 e1452.
Van Bambeke, F., Y. Glupczynski, P. Plesiat, J.C. Pechere & P.M. Tulkens, (2003) Antibiotic
efflux pumps in prokaryotic cells: occurrence, impact on resistance and strategies for the
future of antimicrobial therapy. *J Antimicrob Chemother* **51**: 1055-1065.
Wagner, S., M.M. Klepsch, S. Schlegel, A. Appel, R. Draheim, M. Tarry, M. Hogbom, K.J. van
Wijk, D.J. Slotboom, J.O. Persson & J.W. de Gier, (2008) Tuning Escherichia coli for
membrane protein overexpression. *Proc Natl Acad Sci U S A* **105**: 14371-14376.
Wong, K., J. Ma, A. Rothnie, P.C. Biggin & I.D. Kerr, (2014) Towards understanding promiscuity
in multidrug efflux pumps. *Trends Biochem Sci* **39**: 8-16.
Zheng, S., L.T. Sham, F.A. Rubino, K.P. Brock, W.P. Robins, J.J. Mekalanos, D.S. Marks, T.G.
Bernhardt & A.C. Kruse, (2018) Structure and mutagenic analysis of the lipid II flippase
MurJ from Escherichia coli. *Proc Natl Acad Sci U S A* **115**: 6709-6714.