Dynamic persistence of intracellular bacterial communities of uropathogenic Escherichia

coli in a human bladder-chip model of urinary tract infections

Authors: Kunal Sharma, Neeraj Dhar*[&], Vivek V. Thacker*[&], Thomas Simonet, François

Signorino-Gelo, Graham Knott, John D. McKinney*

Affiliations:

School of Life Sciences, Swiss Federal Institute of Technology in Lausanne (EPFL), 1015

Lausanne, Switzerland.

[&] These authors corresponded equally

*Correspondence to: neeraj.dhar@epfl.ch, vivekvthacker@gmail.com , john.mckinney@epfl.ch

Abstract

Uropathogenic *Escherichia coli* (UPEC) is the most common causative agent of urinary tract

infections and is a major cause for antibiotic prescriptions. Previous studies have shown that

infection of terminally differentiated urinary bladder cells leads to the formation of intracellular

bacterial communities (IBCs). However, the precise role of IBCs in recurrence of infection and

antibiotic persistence, is not completely understood in part because the *in situ* dynamic responses

of bacteria within these structures to antibiotic stress is difficult to assess in animal models. Here,

we develop and characterize a human bladder-chip model of UPEC infection wherein superficial

bladder epithelial cells and bladder microvascular endothelial cells are co-cultured under flow in

urine and nutritive media respectively, and the mechanics of bladder filling and voiding cycles

mimicked by application and release of linear strain. Time-lapse microscopy showed that

infection of epithelial cells under shear stress in diluted urine led to the rapid recruitment and

diapedesis of neutrophils across the endothelial-epithelial barrier and the formation of neutrophil

swarms and neutrophil extracellular traps. Subsequently, two cycles of antibiotic administration

interspersed with recovery periods revealed both non-growing and rapidly proliferating IBCs.

Multiple stages of IBC formation captured on-chip with single-cell resolution revealed that that

bacterial killing within IBCs was substantially delayed, outcomes such as shedding of bacteria

and exfoliation are not mutually exclusive and rapidly reseeded the infection, and in rare

instances bacterial growth in IBCs continued for the entire period of antibiotic administration.

These new insights into the early stages of pathogenesis revisit the role of IBCs as harbours of

persistent bacterial populations, with significant consequences for non-compliance with

antibiotic regimens.

Keywords: bladder-chip, uropathogenic *Escherichia coli* (UPEC), intracellular bacterial

communities (IBCs), bacterial persistence, antibiotic treatment, neutrophil swarms, neutrophil

extracellular traps, shear stress, stretching, confocal live-cell imaging

Introduction

Urinary tract infections (UTIs), the second most-common cause for the prescription of antibiotics (Foxman, 2010) are characterized by the high frequency of recurrence, defined as a reappearance of infection within 12 months despite the apparently successful completion of antibiotic therapy. Recurrence occurs in about 25% of all UTIs (Foxman et al., 2000) and strongly impacts the cost of healthcare and reduces the quality of life, particularly considering the fact that more than 60% of women are diagnosed with a UTI at least once in their lifetime (Klein and Hultgren, 2020). Uropathogenic *Escherichia coli* (UPEC), the causative agent for the majority of UTIs, exhibits a complex lifestyle in the bladder, with planktonic sub-populations within the urine co-existing with intracellular bacteria. UPEC invasion of the urinary bladder generates substantial changes to bladder morphology and a robust immune response. Much of our current understanding of early stages of UTI and the intracellular lifestyle is derived from studies in the mouse model (Anderson et al., 2003; Hannan et al., 2012; Hung et al., 2009; Justice et al., 2004) ,. Examinations of mouse bladder explants via microscopy have revealed the formation of intracellular bacterial communities (IBCs) composed of thousands of bacteria within individual superficial bladder cells (Justice et al., 2004). IBCs also play a prominent role in clinical infection and have also been harvested from the urine of cystitis patients (Robino et al., 2013; Rosen et al., 2007). These structures are considered to have a linear progression from an early stage of colonization by single bacteria to an intermediate stage of biofilm-like communities culminating in the release of bacteria at the late stage. However, long-term imaging of infected animals or of explant-tissue is technically challenging and therefore it has been difficult to capture the dynamic changes that underlie the formation of these structures and their consequences to infection.

A key strength of advanced organotypic models such as organ-on-chip (Benam et al., 2016; Huh

et al., 2010a; Jang et al., 2013, 2019; Kim et al., 2016; Novak et al., 2020; Zhou et al., 2016) or

organoids (Cakir et al., 2019; Clevers, 2016; Rossi et al., 2018; Sato et al., 2009) is that they

increasingly recapitulate the complexity of host physiology. The bladder is a particularly

challenging organ to mimic; it has a stratified architecture with well-differentiated cell types; a

lumen filled with urine whose composition and chemical properties can vary depending on the

physiological state of the individual and is subjected to periodic and large changes in organ

volume and surface area (Korkmaz and Rogg, 2007). These features may play important roles in

infection, but they have been hitherto hard to capture outside of a whole animal model, where

they are present in their entirety and cannot be dissected in a modular manner.

Organotypic models are well-suited to address these outstanding questions. Recently, bladder

organoids that mimic the stratified architecture of the urothelium have been used to study

infections (Horsley et al., 2018; Smith et al., 2006; Sharma et al., 2020). However, the organoid

model suffers from certain shortcomings inherent to the 3-D architecture such as the lack of

vasculature, the inability to manipulate the cells mechanically, and the constrained volume of the

lumen. In contrast, organ-on-chip models offer a complementary approach that does not suffer

from these limitations. Organ-on-chip models have been developed for a number of organs either

in isolation or in combination (Novak et al., 2020; Ronaldson-Bouchard and Vunjak-Novakovic,

2018). These systems are being increasingly used to model infectious diseases including viral

and bacterial infections of the respiratory tract (Nawroth et al., 2020; Si et al., 2020; Thacker et

al., 2020b, 2020a), gut (Jalili-Firoozinezhad et al., 2019; Kim et al., 2016; Shah et al., 2016;

Tovaglieri et al., 2019), kidney (Wang et al., 2019), and liver (Kang et al., 2017). However, to

our knowledge, there have been no reports of a bladder-on-chip model. While there have been a

few studies that have developed *in vitro* bladder models that recreate the stratified architecture of

the bladder epithelium (Horsley et al., 2018; Smith et al., 2006; Suzuki et al., 2019), they have

not been used to recapitulate the multiple stages of IBC formation. Similarly, while a few studies

have visualized different stages of IBC development by culturing human bladder cells under

urine flow, these models are restricted to monoculture experiments(Andersen et al., 2012;

Iosifidis and Duggin, 2020). Furthermore, the lack of vasculature in these models restricts the

extent to which immune cell components or drugs can be introduced in a physiologically relevant

manner and none of the systems reported to date offer the possibility of mimicking bladder

mechanics(Andersen et al., 2012; Horsley et al., 2018; Iosifidis and Duggin, 2020; Smith et al.,

2006).

Here we report the development and characterization of a bladder-chip model that mimics the

bladder architecture by co-cultures of a well-characterized human bladder epithelial cell line with

bladder microvascular endothelial cells in a device geometry that allows the two cells to be

exposed to urine and nutritive cell culture media respectively. The flow rates in the apical and

vascular channels can be controlled independently, and multiple rounds of micturition are

recreated via the application of a linear strain to the PDMS membrane used for co-culture. Using

this model, we show that diapedesis of neutrophils to sites of infection on the epithelial side can

lead to the formation of neutrophil swarms and neutrophil extracellular traps (NETs), and that

IBCs offer substantial protection to bacteria from antibiotic clearance. Our observations suggest

that the role of IBCs in reseeding bladder infections upon cessation of antibiotic treatment or

failure to complete a course of antibiotics might be more important than previously assumed and

therefore strategies aimed towards eradication of IBCs are very crucial for treatment efficacy.

Results

Reconstitution of bladder uroepithelium and bladder vasculature

We established a bladder-chip infection model by co-culturing HTB9 bladder epithelial cells

(epithelial cells) and HMVEC-Bd primary human bladder microvascular endothelial cells

(endothelial cells) in a novel bladder-chip approach (Fig. 1A). Immunostaining verified that the

epithelial and endothelial cells formed confluent monolayers (cell densities in Table 1) with high

expression of junction markers such as epithelial cell adhesion molecule (EpCAM) and platelet

endothelial cell adhesion molecule-1 (PECAM-1) and VE-cadherin in the epithelial and

endothelial cell layers respectively (Fig 1B, C, additional images in Fig. S1A-H). Expression of

these and other junction markers such as E-cadherin and zonula occludins-1 (ZO-1) was

consistent across monocultures (Fig. S2A-G) and in co-culture on-chip.

Similarly, a majority of the epithelial cells expressed cytokeratin 7 (CK-7), a general

uroepithelial marker and cytokeratin 8 (CK-8), a differentiated uroepithelial marker both in

monoculture (Fig. S2I, J) and on-chip (Fig. 1B, S1B, S1F). Some endothelial cells were also CK-

7+ (Fig. 1C, S2H). Co-culture of bladder epithelial and bladder endothelial cells in human-

bladder chip therefore did not alter the cellular expression patterns of these markers as compared

to monocultures. We further characterized the HTB9 cell line in monoculture for markers for

urothelial differentiation. HTB9 cells showed high expression of uroplakin-3a (UP3a) which has

been shown to be essential for UPEC infection (Fig. S2K) (Martinez et al., 2000; Mulvey et al.,

1998) but only a small proportion of epithelial cells were positive for the basal cell marker

cytokeratin1 (CK1), in agreement with the umbrella cell nature of the HTB9 cell line (Fig S2L,

M) (Duncan et al., 2004; Smith et al., 2006). Overall, these observations confirmed that the

bladder-chip is populated with cells that mimic the physiology of the bladder vasculature and the

superficial urothelial cell layer. Further, the chip design enables the flow of media with different

compositions through the epithelial and vascular channels. We used this feature to mimic bladder

physiology by perfusion of diluted pooled human urine on the epithelial side and endothelial cell

medium on the vascular side.

Modelling bladder filling and bladder voiding in the bladder-chip

The human bladder experiences vast changes in volume and surface area on a periodic basis. The

first phase of this cycle 'the *filling bladder* state', occurs through a slow and gradual increase in

bladder volume due to addition of urine from kidney via the ureters, resulting in a large increase

in bladder volume. At the level of the uroepithelium, these changes manifest as a 2-dimensional

biaxial stretch with ex vivo studies in rat bladders reporting peak strains of 90 % and 130%

measured in the circumferential and longitudinal directions (Gloeckner et al., 2002; Parekh et al.,

2010). Subsequently, volume growth slows, and the bladder attains a relatively constant volume,

with the bladder epithelium in a corresponding state of maximal stretch '*filled bladder* state'.

Voiding of the bladder through urination rapidly reduces bladder volume ('*voiding bladder*

state') and removes the strain experienced by the bladder cells. The bladder epithelium, as a

viscoelastic material, responds to the removal of strain and relaxes over the subsequent period

(Pascalis et al., 2018) where the bladder volume remains low ('*voided bladder* state') (Fig. 1D).

The architecture of the bladder-chip device allows for a linear strain to be applied to the porous

membrane via negative pressure in the channels adjacent to the main channel of the device ('the

vacuum channels') (Grassart et al., 2019; Huh et al., 2010b) (Fig. 1A). We characterized the

strain experienced by epithelial cells due to application of negative pressure (Fig. S3). A plot of

applied pressure vs. strain was linear, and we could achieve a dynamic range of linear strain (0 –

19 %) by application of negative pressure (0 – 900 mbar). High levels of applied strain were

incompatible with long-term time-lapse microscopy as it generated a significant drift in the axial

position of the membrane. We therefore limited the linear strain applied to a maximum of 10%,

which is nevertheless a significant proportion of the typical strain experienced by the bladder

tissue in vivo. We then modelled the bladder filling and voiding over a 6-hour duty cycle with

different states: *filling bladder* (0 - 2hr, 0 to 10% strain), *filled bladder* (2 – 4hr, 10% strain),

voiding bladder over a period of 2 minutes (4 hr – 4hr + 2min, 10% to 0% strain) and *voided*

bladder (4hr + 2min – 6 hr, 0% strain). This 6-hour duty cycle was then repeated for the

remaining duration of each experiment. The bladder-chip therefore enables co-culture of two

cell-types in nutritionally different microenvironments with a physiologically relevant level of

applied strain.

Determination of optimal UPEC infection conditions in the bladder-chip model

UPEC attachment and invasion of bladder epithelial cells has been shown to be sensitive to

fluidic shear stress (Andersen et al., 2012; Zalewska-Piątek et al., 2020). Although UPEC

infection in the bladder can occur with or without the presence of shear stress, we infected the

epithelial layer on the bladder-chip under flow conditions. The small volumes of the microfluidic

chip motivated this approach, as rapid bacterial growth in the diluted human urine could lead to

acidification of the medium. We therefore maintained a flow rate of 1.2 ml/hour for a period of

1.5-2 hours during infection of the epithelial layer of the bladder-chips with UPEC. Under flow

conditions, an overwhelming majority of bacteria did not attach to the epithelial cells. Time-

lapse microscopy showed that bacterial attachment to the epithelial cells increased steadily over

this period, but that the typical infectious dose at the end of this period was low (<1 bacterium

per epithelial cell, Fig. S4).

Accordingly, we established a 28-hour live-cell imaging experimental protocol as shown in the

schematic in Fig. 1E and described in greater detail in the subsequent sections and in methods.

Briefly, this consisted of an initial period of homeostasis, followed by UPEC infection,

introduction of neutrophils, and two consecutive cycles of antibiotic treatment and recovery to

monitor IBC dynamics at the single-cell level. This experimental protocol therefore mimicked

key aspects of the host-pathogen interactions in early stages of UTIs as well as the response to

antibiotic treatment.

Diapedesis of neutrophils across the epithelial-endothelial barrier in response to UPEC

infection in the bladder-chip

The bladder-chip platform enables continuous imaging while maintaining flow in both epithelial

and vascular channels and mechanically stretching the membrane as part of the bladder voiding

cycle. At the start of each live-cell imaging experiment, bladder epithelial cells in an initial

relaxed state were perfused with sterile diluted pooled human urine on the epithelial side and

endothelial cell medium on the vascular side respectively (Fig. 2A1, B1, S5A1). The epithelial

side was inoculated with a low dose of UPEC in diluted urine and infection was performed under

flow (Fig. 2A2, B2, S5A2) for ca. 1.5 to 2 hours. The first bladder duty cycle also initiated at this

timepoint. We maintained the optical focus of the microscope on the epithelial layer over the

subsequent course of infection. UPEC attachment was evident during this infection phase by

visual inspection (Fig. 2B2, S5A2).

After this period of infection, unattached bacteria in the epithelial channel were removed by

perfusion of diluted urine. Our aim was to study the interaction of UPEC with host cells,

therefore this perfusion was maintained throughout the experiment. Continuous perfusion

reduced the accumulation of bacteria in the urine and enabled imaging of intracellular bacteria

without large background noise in the fluorescence channels from planktonic bacteria. We

subsequently performed a series of interventions that mimic the course of UTI infections and

used time-lapse imaging to monitor the simultaneous changes in host-pathogen interaction

dynamics. As a host response, immune cells were introduced into the vascular channel. Among

innate immune cells, neutrophils are the first line of defense in UTIs (Haraoka et al., 1999). We

isolated neutrophils from human blood through negative depletion and verified that the isolation

procedure yielded neutrophils with a high degree of purity (~96% as reported in (Son et al.,

2017)) through immunostaining for the anti-CD15 a neutrophil specific marker (Zahler et al.,

1997) (Fig. S6). To enable identification and tracking of neutrophils, they were pre-labelled with

CellTracker^{TM} Deep Red dye, which stains the cytoplasm of these cells.

Neutrophils were subsequently introduced in the bladder-chip devices through the vascular

channel, mimicking the natural route of immune cell migration into the bladder (schematic in

Fig. 2A3-A5, snapshots in Fig. 2B3-B5, S5A3-A5). We used a cell concentration of ca. 2 million

cells/mL similar to the neutrophil concentration in human blood (Hsieh et al., 2007). During this

period, the flow rate through the vascular channel was increased to provide a shear stress $\eta = 1$

dyne/cm^2, which aids neutrophil attachment to endothelial cells (Alon et al., 1995). In uninfected

control chips, neutrophil attachment to the endothelial layer was minimal (Fig. S7A, B).

Diapedesis of neutrophils to the epithelial layer was rare (Fig. S8A, 2C). In stark contrast,

infection of the epithelial layer with UPEC elicited a robust attachment of neutrophils to

endothelial cells (Fig. S7C, D) along with a rapid diapedesis across to the epithelial layer (Fig.

2B2-B4, S5A2-A4, S8C). We confirmed that neutrophil diapedesis was also stimulated in the

absence of infection, upon exposure to a gradient of pro-inflammatory cytokines that are

typically upregulated during UTIs (Agace et al., 1993; Hedges et al., 1992; Nagamatsu et al.,

2015; Song et al., 2007) (interleukin-1α, interleukin1-β, interelukin-6 and interleukin-8 each at

100 ng/mL) across the epithelial-endothelial barrier (Fig. S8B). However, neutrophil diapedesis

was more robust under infection as compared to the cytokine gradient, with a higher number of

neutrophils observed across multiple fields of view on the epithelial side (Fig. S8B-D). These

results suggest that infection on-chip generates a strongly pro-inflammatory environment locally

and that neutrophil diapedesis is further stimulated by the presence of UPEC(Agace et al., 1995;

Oliveira et al., 2016).

Using time-lapse imaging, we were also able to quantify the kinetics of neutrophil migration

with a temporal resolution of up to 7.5 minutes. Neutrophil diapedesis to the epithelial sites of

infection was detected as early as ca. 15-17 minutes post-introduction into the vascular channel

(Fig 2B3, S5A3, supplementary movie SMov1, SMov2). Neutrophils that migrated to the

epithelial side aggregated on the epithelial cells (Fig 2B4-2B5, S5A4-A5, supplementary movie

SMov1, SMov2) In some cases, these neutrophils were able to control bacterial growth (Fig.

2B4-2B5, dashed yellow boxes) whereas in others, bacterial growth was uncontrolled (Fig. 2B4-

2B5, dashed white boxes). Diapedesis was observed in every field of view examined across n=4

infected chips, whereas diapedesis occurred infrequently in uninfected controls (Fig. 2C).

Furthermore, the size of these aggregates was significantly larger in infected bladder-chips vs.

uninfected controls, with the aggregates in case of infected-chips exhibiting the characteristic

features of neutrophil swarms (Kienle and Lämmermann, 2016; Kreisel et al., 2010;

Lämmermann et al., 2013). We quantified the size of the swarms formed by either estimating the

maximum number of neutrophils observed in fields of view on the epithelial side (Fig 2D) or by

measuring the total volume occupied by neutrophils normalized to the volume of a 3D field of

view (Fig 2E). Using either metric, infection generated large swarms that did not occur in

uninfected bladder-chips. The dynamic of swarm formation across n=51 and n=40 fields of view

in two bladder-chips is shown in Fig. 2F. In these experiments a fast-imaging frequency of 7.5

minutes was used. In infected bladder-chips, swarms reached a maximum size between 60- and

90-minutes post-introduction of neutrophils (Fig. 2G).

Neutrophil extracellular traps (NETs) formation in response to UPEC infection

Neutrophils have complex responses to infections (Oliveira et al., 2016), including the formation

of Neutrophil extracellular traps (NETs) that occurs by the release of cytosolic azurophilic

granules around a scaffold of decondensed chromatin and potentially serves to trap and kill

extracellular bacteria (Brinkmann et al., 2004). NETs have been shown to consist of

antimicrobial granules such as myeloperoxidase (Metzler et al., 2011) and neutrophil elastase

(Papayannopoulos et al., 2010) and have been observed in urine harvested from UTI patients (Yu

et al., 2017). Interestingly, we observed NET formation by neutrophils in the infected bladder-

chip. We verified the formation of NETs on the epithelial channel through immunostaining of

infected bladder-chips at 3.5 hours post-infection and 2 hours post-introduction of neutrophils in

the vascular channel using antibodies against neutrophil myeloperoxidase (Fig 2H, S9A1-A3)

and elastase (Fig 2I, S9B1-B3). In each case, areas on the epithelial layer with strong

immunostaining were observed (Fig. 2H3, 2I3). Neutrophils in these areas were typically in large

numbers, characteristic of swarms. In addition, by this stage of the experiment, neutrophils that

had high levels of myeloperoxidase or elastase typically did not retain strong expression of the

cytoplasmic CellTracker dye used for identification of neutrophils in live imaging (cf. Fig. 2H2

and 2H3, 2I2 and 2I3). For each example, a zoomed-in image clearly shows long filament-like

structures that extend between cells and are either myeloperoxidase positive (Fig. 2H5) or

neutrophil elastase positive (Fig. 2I5), strongly suggestive of NETs (Brinkmann et al., 2004;

Metzler et al., 2011; Papayannopoulos et al., 2010). NET formation was also observed in some

instances by neutrophils in some locations on the endothelial channel, although large swarms did

not form here (Fig. S9B1-B3).

To further characterize these structures, we imaged the epithelial layer of uninfected and infected

bladder-chips with or without the addition of neutrophils using scanning electron microscopy

(SEM). Representative images of the epithelial layers of uninfected bladder-chip controls (Fig.

S10A) contrast with those from infected bladder-chips after neutrophil diapedesis (Fig. 2J1-J4,

S10C) and clearly highlight the distinctive features of NET formation. In Fig. 2J1, neutrophils on

the surface of the epithelial layer are clearly visible by their distinctive spherical morphology

(amber arrowheads). NETs formed between these neutrophils are shown by magenta arrowheads

and appear to extend between adjacent neutrophils. A majority of the epithelial cells have a

characteristic flattened morphology with cell appendages that can extend between cells (white

arrowheads); a highly infected spherical epithelial cell is indicated with a purple arrowhead.

Zooms in Fig. 2J2-2J4 show the structure of NETs in greater detail. Many individual UPEC

bacteria (green arrowheads) are captured in these NETs. These structures did not form in

infected bladder-chips without the introduction of neutrophils (Fig. S10B), confirming their

identity as NETs identified via immunofluorescence. NET formation was a direct response to

UPEC and not due to hyperactivation of neutrophils due to experimental handling. To verify this,

we collected the neutrophils that passed through an infected bladder-chip without attachment.

Even upon infection, an overwhelming majority of the cells perfused into the chip flow out

without adhering, because of the geometry of the channels in the device. NET formation in these

samples was not observed unless the cells were themselves infected by UPEC shed from the cells

in the bladder-chip (Fig. S11).

Heterogenous dynamics of intracellular bacterial communities within urothelial cells

Urinary tract infections that do not resolve upon intervention by the host immune system often

require treatment with antibiotics. Antibiotics have complex pharmacokinetic and

pharmacodynamic profiles *in vivo*, which we attempted to model through two successive rounds

of high dose administration of antibiotics (40x MIC, Fig. S12A), on both the epithelial and

vascular side, interrupted by periods with no antibiotic (schematics in Fig. 3A). Continuous time-

lapse imaging over this entire period allowed us to capture the responses of bacteria to this

simplified model of antibiotic profiles and to study the persistence of UPEC upon antibiotic

treatment in the different physiological niches (Fig. 3 and Fig. 4).

A consequence of the first cycle of antibiotic administration under flow was the elimination of

extracellular axenic bacterial growth that had occurred in the media in the apical channel, as is

observed within the lumen of the bladder *in vivo*. Many that survived the antibiotic treatment

intact did not regrow after the antibiotic was removed (Fig. S12 B). Regrowth commenced in

only a small fraction of bacteria (Fig. S12C) after a variable lag phase (Fig. S12D) and rapid

regrowth was highly correlated with intracellular location of bacteria within epithelial cells.

These subsequently proceeded to form intracellular bacterial communities (IBCs), with a

variable lag period. An example of an early stage of the IBC formation is shown in the timeseries

in Fig. 3B1-B5, supplementary movie SMov3. This IBC was initially seeded with few bacteria

that are present in Fig. 3B1 and clearly visible in Fig. 3B2. At the end of the timeseries, the

entire host cell was packed with bacteria (Fig. 3B5). The exponential increase in bacterial

numbers frequently led to the saturation of the 8-bit images in the bacterial channel. We

therefore examined similar structures in separate bladder-chips via immunofluorescence (Fig.

3C, S13) staining. Zooms of two of the four IBCs highlighted in Fig. 3C show numerous tightly

packed bacteria within epithelial cells that are both CK7+ (Fig. 3C) and CK8+ (Fig. S13). These

bacterial morphologies are highly similar to early-stage IBCs observed in the bladder(Anderson

et al., 2003; Justice et al., 2004). Higher resolution images of the biofilm-like structures within

IBCs obtained via correlative light (Fig. 3D) and transmission electron microscopy (Fig. 3E)

showed that bacteria in IBCs can be either coccoid or rod shaped, which is also evident in

scanning electron micrographs of bacteria within IBCs (Fig. 3F). IBCs within bladder-chip

model therefore show many of the distinctive morphological features reported from images of

infected bladders in the mouse model (Anderson et al., 2003; Hunstad and Justice, 2010).

Numerous reports from the mouse model of infection have shown IBCs to be dynamic structures;

growth in IBCs eventually culminates in bacterial shedding or the complete exfoliation of the

IBC. The underlying dynamics of these phenotypes is hard to capture in the mouse model, but

the bladder-chip model allowed us to track the dynamics of bacterial growth within individual

IBCs over extended periods of time, providing information on bacterial dynamics within so-

called early, middle and late-stage IBCs. An example of the heterogeneity in growth rates is

evident from the timeseries for three IBCs in Fig. 3G1-G3, supplementary movie SMov4.

IBCs#1 and 2 subsequently began to shed individual bacteria (Fig. 3G4), a phenotype not

observed in IBC#3. At a later timepoint (Fig. 3G5) the fates of all three IBCs were dramatically

different – the cell containing IBC#1 had completely exfoliated and was removed from the field

of the view by flow in the epithelial channel, IBC#2 continued to shed bacteria, whereas growth

remained slow in IBC#3. We monitored the eventual fate of n=100 IBCs during the first and the

second growth cycles which confirmed that shedding and exfoliation were not mutually

exclusive. While some IBCs only shed bacteria (n=28) and others exfoliated without shedding

(n=27), a substantial fraction of IBCs showed shedding prior to exfoliation (n=31) and some

IBCs displayed neither phenotype (n=14). Furthermore, in addition to the coccoid and rod-

shaped morphologies highlighted in Fig. 3C-F, time-lapse imaging also enabled us to capture

events that occurred relatively rarely such as the intracellular growth of filamentous forms of

UPEC (Fig. 3H1-H5, S14, supplementary movie SMov5, SMov6). In this time series, a

filamentous bacteria can be seen and appears to encircle the inner boundary of the cell (Fig. 3H1-

3H4) before filling the volume entirely (Fig. 3H5). Overall, time-lapse imaging confirms that a

significant fraction of bacteria within IBCs can survive the first round of antibiotic treatment and

are the source for reseeding of both the extracellular bacterial populations as well as the

subsequent growth of IBCs in newly infected cells.

Dynamic persistence of intracellular bacterial communities within urothelial cells

A subsequent round of antibiotic treatment provided an opportunity to study the dynamic

persistence of bacteria within IBCs in response to an antibiotic profile that was a closer mimic of

the varying antibiotic concentrations in vivo. Furthermore, at this stage of the experiment, we

were also able to study the responses of bacteria in middle and mature late-stage IBCs to

antibiotic treatment which was not possible during the first round of antibiotic administration

early in the course of infection. In the examples highlighted in Fig. 4A1-A5 and supplementary

movie SMov7, bacterial growth after the first round of antibiotic treatment resulted in the

formation of many large IBCs with tightly packed bacteria (Fig. 4A1-A2). Many of the bacteria

within each of the four IBCs were eliminated by the antibiotic, but a substantial proportion of

bacteria in each IBC nevertheless survived the antibiotic treatment (Fig. 4A3). Each of these

IBCs subsequently regrows when the antibiotic is removed (Fig. 4A4-A5), and two additional

IBCs are seeded during this time (white arrowheads in Fig. 4A5). Overall, the second round of

antibiotic administration led to a sharp decline in instances of subsequent bacterial regrowth (Fig.

S12C). In all instances, regrowth either occurred directly within IBCs, or was caused by the

shedding of bacteria from IBCs to repopulate the extracellular niche and seed new IBCs,

highlighting the dynamic stability of this niche and its importance in establishing persistent

infection. We therefore performed a careful analysis to quantify bacterial growth rate within

IBCs. Growth within IBCs across different time periods was exponential (Fig. 4B). We were

able to track a subset of IBCs across two growth cycles (Fig. 4C). In a majority of instances these

revealed exponential growth in the absence of antibiotic, delayed response to antibiotic, and a lag

phase after the antibiotic was removed. In general, bacterial growth in IBCs in the second growth

cycle was slower than in the first (Fig. 4D). However, this population level statistic may be

influenced by the fact that many of the IBCs monitored in the second growth cycle had been

exposed to two rounds of antibiotic treatment. For this sub-population, in n=16 out of n=18

IBCs, growth was slower after the 2^{nd} round of antibiotic administration (Fig. 4E).

Next, we examined the dynamics of bacterial growth within IBCs during the period of antibiotic

treatment in greater detail. In the IBC shown in the timeseries in (Fig. 4F1-4F5, S15A1-A5,

supplementary movie SMov8, SMov9), growth before antibiotic administration (Fig. 4F1, 4F2)

continued for a considerable period after the antibiotic was administered (Fig. 4F3, Fig. 4F4,

S15A1-S14A3). Eventually, a reduction in bacterial volume towards the end of the antibiotic

administration period was observed (Fig. 4F5, S15A4-S15A5). In contrast, the relatively smaller

IBC in the timeseries in Fig. 4G1-4G5, S15B2-S15B5, supplementary movie SMov10, SMov11

continued to grow throughout the period of antibiotic administration. These different dynamics

are also captured in the plot of bacterial volume for multiple IBCs before, during, and after the

antibiotic treatment. In all cases, bacterial volume continues to increase during a significant

period of antibiotic administration (Fig. 4H), and in 2 out of 18 cases, there was no decrease in

bacterial volume throughout this period. In contrast, extracellular bacteria likely adherent on the

epithelial cells but not internalized within them were rapidly eliminated by combination of

antibiotic treatment and flow (Fig. 4I), in all cases the bacterial volume reduced immediately

upon antibiotic administration. This differential killing resulted in a significantly higher

proportion of extracellular population of bacteria being killed (Fig. 4J). Elimination of bacteria

within IBCs was highly heterogenous. Another consequence of this differential killing was that

after the 2^{nd} round of antibiotic treatment, regrowth was observed in only a very small fraction of

non-IBC UPEC (Fig. S12C). Protection of bacteria within IBCs therefore has a direct outcome of

enabling the reseeding of infection at other locations within the epithelial monolayer.

Lastly, we sought to determine if the bladder duty cycle altered the dynamics of UPEC infection.

A comparison between infected bladder-chips with and without the duty cycle (Fig. 4K, L, S16)

revealed a significant increase in the bacterial burden when the duty cycle was implemented

(Fig. 4M). These results suggest that the bladder filling and voiding cycle influence the uptake

and proliferation of UPEC, possibly through physiological changes in the epithelial cells in

response to applied strains (Apodaca, 2004; Carattino et al., 2013; Truschel et al., 2002; Wang et

al., 2005). In turn, they suggest a deeper connection between the physiology of mechanically

active organs such as the bladder and infection, which can be probed further in the bladder-chip

model.

Discussion

In our human bladder-chip model we focus on recapitulating the key aspects of bladder

physiology relevant to the study of the earliest stages of UPEC infection and IBC formation.

Therefore, superficial bladder epithelial cells, the first cell-type usually infected by UPEC, are

cultured both under flow in diluted pooled urine and in co-culture with bladder microvascular

endothelial cells. The ability to apply a cyclic mechanical stretch to the PDMS membrane,

originally designed to mimic the breathing motion in the lung (Huh et al., 2010b) or the

peristaltic motion in the gut (Kim and Ingber, 2013), has been adapted here to mimic the slow

expansion and rapid contraction of bladder volume. We demonstrate the ability to perform

multiple duty cycles while simultaneously imaging the infected device via long-term time-lapse

imaging, a technical advance that is difficult to achieve with bladder explants (Justice et al.,

2004) or other *in vitro* studies of UPEC infection of bladder epithelial cells (Andersen et al.,

2012; Horsley et al., 2018; Iosifidis and Duggin, 2020; Smith et al., 2006). Using this approach,

we found that the total bacterial burden inside infected bladder-chips was significantly higher at

a late stage of infection if a duty cycle was applied, unlike the non-significant influence of cyclic

stretching on Shigella infection of intestinal-chips under flow conditions (Grassart et al., 2019).

The exact mechanisms underlying this phenomenon remain to be explored, but there is an

increasing understanding of the role of mechanical forces in regulating innate immune function

(Solis et al., 2019). Our results showcase the ability of the bladder-chip model to capture these

interactions between mechanical function, physiology and infection, unlike other infection

models reported thus far.

A primary focus of *in vitro* models developed so far has been to probe specific aspects of UPEC

infection, such as the role of the stratified bladder architecture (Horsley et al., 2018) or the effects

of micturition on IBC formation (Andersen et al., 2012; Iosifidis and Duggin, 2020). However,

the migration of immune cells into the bladder in immune cells is difficult to reproduce in these

models, and in many of these systems, live-cell imaging remains technically challenging

(Horsley et al., 2018; Smith et al., 2006). Furthermore, the models do now allow mechanical

manipulations of the cellular co-culture. In that sense, the bladder-chip model complements these

existing approaches by providing these additional functionalities. Further development,

potentially through the combination of organoid and organ-on-chip approaches, could lead to the

development of a fully stratified urothelium on-chip, although the utility for studying IBC

dynamics is relatively limited.

Neutrophils are the first responders to UPEC infection (Abraham and Miao, 2015; Haraoka et al.,

1999), and neutrophil migration involves a series of steps that commences with attachment to the

endothelium under flow, migration on the endothelial surface, diapedesis across the epithelial-

endothelial cell barrier and movement towards the site of infection (Ley et al., 2007; Nourshargh

et al., 2010). Perfusion of the vascular channel with CD15+ neutrophils isolated from human

blood shows that the bladder-chip model recapitulates all of these phenotypes, with rapid

diapedesis of neutrophils to sites of infection. There, neutrophils aggregate to form swarms

(Kienle and Lämmermann, 2016; Lämmermann et al., 2013) and subsequently, the formation of

NETs (Branzk et al., 2014; Brinkmann et al., 2004; Metzler et al., 2011; Papayannopoulos et al.,

2010) around extracellular bacteria is observed. Characterization of these structures via SEM

shows that they can extend across many tens of microns and stain positive for myeloperoxidase

(Metzler et al., 2011) and neutrophil elastase (Papayannopoulos et al., 2010), indicative of potent

anti-microbial activity. However, neutrophil control of infection on-chip is partial; this is

possibly due to the unrestricted growth of large numbers of extracellular bacteria and in some

instances, exacerbated by the loss of some neutrophils due to flow in the channel. Another

contributing factor could potentially be the architecture of the PDMS membrane, which permits

neutrophil diapedesis only at fixed spatial locations on-chip. Neutrophil migration may also be

impacted by the relative stiffness of the PDMS membrane. Nevertheless, the demonstration of

NET formation is consistent with the occurrence of these structures both in the mouse model

(Ermert et al., 2009) as well as in the urine of infected patients (Yu et al., 2017) and suggests that

the model is able to recapitulate important aspects of disease. It is also an important advance for

the use of organ-on-chip approaches to recapitulate NET formation in infectious disease.

IBC formation begins immediately after infection and is already underway upon neutrophil

administration. However, clear demonstration of IBCs with a high degree of confidence was only

possible after an initial treatment with a high dose of antibiotic that eliminated extracellular

planktonic bacteria and improved the optical imaging of bacteria attached to or within the

epithelial cells. It enabled us to capture the full cycle of IBC growth from 1-2 bacteria to a large

biofilm and subsequent release via shedding and filamentation. Importantly, the compact nature

of the device allowed us to image multiple IBCs concurrently on the same chip with a high

temporal resolution, which is difficult to achieve in bladder explants (Justice et al., 2004). Our

observations reiterate the highly dynamic nature of these structures; growth was asynchronous

and heterogenous and outcomes included bacterial shedding, exfoliation and filamentation

(Hunstad and Justice, 2010; Justice et al., 2004; Scott et al., 2015). Notably, shedding and

exfoliation were not mutually exclusive for the time-period of our observations – we report

examples of IBCs that shed bacteria and contract in volume before exfoliation.

While IBCs are clearly acknowledged as critical players in early infection, the contribution

towards promoting persistence is not completely understood. For example, Blango *et al*. (Blango

and Mulvey, 2010) showed that incubation of bladder explants containing IBCs with a high dose

of antibiotics for a period of 18 hours resulted in a substantial sterilization of these structures and

concluded that other populations, notably quiescent reservoirs (Mysorekar and Hultgren, 2006)

might play a greater role in the establishment of persistent populations. However, the

pharmacokinetic profiles of most antibiotics in the host are not time-invariant. In case of

ampicillin, a standard regimen of ampicillin treatment is typically a bolus of 250-500 mg of

antibiotic administered every 6 hours. Within the serum, ampicillin concentration peaks at a C_{max}

~ 3-40 μg/ml, which is between 1.5 to 20-fold the MIC as measured in human serum (Bryskier,

2020; Putrinš et al., 2015). These values are in good agreement with the concentrations we use,

that are 40-fold higher than the MIC measured in the endothelial medium perfused in the

vascular channel. Ampicillin concentrations in the blood rapidly decay with a half-life of

between 60 and 90 minutes with a characteristic pharmacokinetic/pharmacodynamic profile.

This period is modelled well by our experimental protocol where phases with high concentration

of ampicillin are interspersed with periods with no antibiotic. We are therefore able to model the

delivery of two consecutive doses of antibiotic and find that IBCs offer substantial protection

against sterilization by a short duration of a high-dose antibiotic treatment and in many instances,

bacterial regrowth after two successive rounds of antibiotic administration. Notably, in smaller

'early IBCs', bacterial growth continues throughout the period of ampicillin administration,

suggesting that the intact nature of the cell membrane likely diminishes the effect of the drug.

Our results suggest that, IBCs may continue to play a role in reseeding sites of infection for a

considerable period after the commencement of antibiotic treatment. This has particularly

important implications with regard to the compliance of antibiotic use as proliferating IBCs

could rapidly re-seed sites of infection throughout the bladder if antibiotic doses are missed.

These unique capabilities of the bladder-chip to realistically model antibiotic treatment regimens

for IBCs can in the future be leveraged to screen compounds (Spaulding et al., 2017) and identify

optimal antibiotic regimens, treatments that can eliminate persistent bacterial populations in

IBCs or alter the host-pathogen interaction dynamic in UTIs.

In summary, the bladder-chip model incorporates aspects of bladder physiology highly relevant

to early UPEC infection in a platform amenable to long-term live-cell imaging as well as for the

administration of antibiotics and therapeutics in a physiologically relevant manner. Our results

establish the suitability of this model for immunological and drug-delivery studies and show that

IBCs are highly dynamic structures that offer substantial protection from antibiotic clearance for

an extended period of time.

Figure 1: Human Bladder-chip model of UTI recapitulates the physiology of bladder filling and voiding.

(**A**) Schematic of the human bladder-chip with co-culture of the HTB9 human bladder epithelial cell line (epithelium, top) and primary human bladder microvascular endothelial cells (endothelial, bottom) on either side of the stretchable and porous membrane. Pooled human urine diluted in PBS and endothelial cell medium were perfused in the apical and vascular channels respectively to mimic bladder physiology. A negative pressure in the 'vacuum' channels (magenta) on either side of the main channel was applied to stretch the porous membrane to mimic stretching of the bladder. (**B**, **C**) Immunofluorescence staining of confluent epithelial and endothelial cell monolayers (anti-EpCAM (magenta) and anti-CK-7 (yellow) for the epithelial cells and anti-PECAM-1 (green) for the endothelial cells) in an uninfected control chip. Some endothelial cells also stained positive for CK7. Cell nuclei were labeled with DAPI (azure). (**D**) Schematic of the reconstitution of the bladder filling and voiding cycle via stretching of the membrane with a duty cycle of 6 hours. The cycle consisted of a linear increase in strain through

stretching of the membrane (*filling bladder*, 0 - 2 hours), maintenance of the membrane under

stretch (*filled bladder*, 2 - 4 hours), a quick relaxation of applied strain over 2 minutes (*voiding*

bladder, 4hr+2 min) and maintenance without applied strain (*voided bladder*, 4hr+2min - 6hr).

(**E**) An overview of the timeline of the experimental protocol including infection, addition of

neutrophils via the vascular channel, and two cycles of antibiotic treatment interspersed by two

bacterial growth cycles. The consecutive bladder duty cycles are indicated.

Figure 2: Neutrophil diapedesis and NET formation on-chip

(**A1-A5**) Schematic of the UPEC infection, introduction of neutrophils and diapedesis of

neutrophils across the epithelial-endothelial barrier to sites of infection. Flow in the epithelial

and endothelial channels is indicated by arrows; the flow rate was increased upon introduction of

neutrophils in the vascular channel to enable optimum attachment. (**B1-B5**) Snapshots from

time-lapse imaging highlighting each stage in the infection cycle shown in (**A1**-**A5**). Bladder

epithelial cells (magenta) and neutrophils (amber) were identified with membrane (Cell Mask

Orange) and cytoplasmic (Cell Tracker Deep Red) dyes, respectively. UPEC identified via

constitutive expression of YFP are shown in green. Neutrophil swarms could either control

bacterial growth (yellow dashed square, compare B4 vs B5) or did not manage to restrict

bacterial growth (white dashed square, compare B4 vs B5). (**C**) Bar charts for relative frequency

of neutrophil diapedesis (black) in n=3 uninfected control bladder-chips and n=4 infected

bladder-chips. Data obtained from n=95 and n=116 across 206 x 206 μm^2 fields of view. (**D**)

Quantification of the number of neutrophils detected on the epithelial layer, in control and

infected bladder-chips. The red bar represents the median value. p<1E-15. In many instances in

the uninfected control bladder-chips, no neutrophil diapedesis is detected. (**E**) A plot of the

maximum neutrophil swarm volume on the epithelial layer normalized to the total volume for

n=67 and n=118 fields of view on n=3 uninfected control and n=3 infected bladder-chips. (**F**) A

plot of neutrophil swarm volume on the epithelial layer over time for n=51 and n=40 fields of

view for n=2 technical replicates indicated by squares and circles. For each time profile, the

volume is normalized to the maximum volume attained over the timeseries and t=0 refers to the

timepoint at which neutrophils are introduced into the vascular channel (**G**) Plot of the time to

the reach the maximum swarm volume in n=154 fields of view across n=4 infected bladder-

chips. (**H1-H54, I1-I5**) NET formation by neutrophils on the epithelial layer. The neutrophils are

identified by a cytoplasmic dye (CellTracker Deep Red, false coloured in amber) (**H2, I2**) and

immunostaining with an anti-myeloperoxidase antibody (**H3,** zooms in **H5**) or an anti-neutrophil

elastase antibody (**I3,** zooms in **I5**). Merged images in each case are shown in **H4** and **I4**. UPEC

identified via YFP expression is shown in spring green. Nuclear labelling with DAPI is shown in

Azure. (**H5**) NETs, identified via anti-myeloperoxidase staining or anti-elastase staining are

indicated with white arrows in (**H5**) and (**I5**) respectively. (**J1**) Scanning electron micrographs of

the epithelial layer of an infected bladder chip 2 hours after the introduction of neutrophils in the

endothelial channel. Neutrophils, (amber arrowheads,) are visible above a layer of epithelial

cells. A heavily infected epithelial cell (purple arrowhead), NETs produced by the neutrophils

(magenta arrowheads), individual UPEC bacteria within the NETs (green arrowheads) are

visible. Appendages between epithelial cells are indicated by white arrowheads. (**J2**, **J3**) Zooms

of the regions in **J1** identified by white squares that highlight the structure of two distinct NETs.

(**J4**) Zoom of the region in **J3** identified by a yellow square. Scale bar = 50 μm in **B1**-**B5**, **H1-**

H5, I1-I5 and 20 μm in **J1-J5**.

Figure 3: Bladder-chip reveals dynamics of IBC growth, shedding and exfoliation. (**A**) Schematics of the host-pathogen interactions within IBCs between successive rounds of antibiotic treatment with outcomes including shedding of bacteria and cell exfoliation shown. (**B1**-**B5**) Timeseries for the growth of an IBC from few bacteria (white arrowhead) after the first round of ampicillin treatment. Bacteria grow to completely fill the cell volume (**B5**). (**C**) Immunofluorescence characterization of two IBCs; the intracellular nature of growth is confirmed with staining with an anti-CK7 antibody. Correlative light (**D**) and transmission

electron micrographs (**E**) show two IBCs filled with both rod-shaped and coccoid-shaped

bacteria. (**F**) Coccoid shaped bacteria (yellow arrowheads) and rod-shaped bacteria (magenta

arrowheads) are also visible in a scanning electron micrograph of an infected epithelial cell.

IBCs on infected bladder chips shown in C-F were fixed 13 hours after UPEC infection and 6

hours into the 1st growth cycle. (**G1-G5**) Three examples of IBC growth, labelled 1 – 3 with

differing outcomes. Unrestricted bacterial growth is observed within all IBCs. In IBC#1 at

timepoint (**G4**) the IBC begins to shed bacteria into the surrounding medium. The cell exfoliates

in the time interval between (**G4** and **G5**). In IBC#2, bacterial shedding is visible at (**G4**) and

shedding continues until end of the timeseries with a reduction in bacterial volume. In contrast,

growth within IBC#3 is slower and neither shedding nor exfoliation occurs within the timeseries.

(**H1**-**H5**) Timeseries highlighting an example of filamentous UPEC growth within an IBC. Scale

bars, 10 μm in **B1**-**B5**, **G1**-**G5** and **H1**-**H5**.

Figure 4: IBCs offer a semi-protective niche that delays clearance of bacteria by antibiotics.
(**A1-A5**) Bacteria persist and grow in IBCs despite antibiotic treatment. Snapshots show the growth of four IBCs with variable growth rates (**A1**-**A2**). Ampicillin treatment eliminates some but not all of the bacteria within each IBC (**A3**). Growth resumes at all sites in (**A4, A5**), formation of new IBCs in the 2nd growth cycle is indicated by white arrowheads. (**B**) Plots of

logarithm of bacterial volume within five separate IBCs show a linear increase in volume over

time, suggesting bacterial growth is exponential. IBCs are seeded dynamically; growth can occur

either in the first or second growth cycle, or prior to the administration of antibiotic, or in few

cases continues in the presence of the antibiotic. (**C**) Plots of the logarithm of bacterial volume

vs. time for n=18 IBCs tracked across two growth phases with an intermediate period of

ampicillin treatment. In the growth phases, growth is exponential and bacterial volume continues

to increase for up to ca. 120 minutes after administration of antibiotic before declining due to

loss of bacteria. In each case, growth resumes after the antibiotic is removed. (**D**) Scatter plots of

the doubling time of bacterial volume in IBCs as measured in the first growth cycle (n=102) and

the second growth cycle (n=59). Growth in the second cycle is significantly slower (p=4.7E-7),

red line represents the median value. P-values calculated using a Mann-Whitney test. (**E**)

Doubling time of bacterial volume in IBCs in the first and second growth cycle for some of the

IBCs in (**D**) that survived the antibiotic treatment. In n=16 out of 18 instances, growth is slower

in the second growth cycle. (**F1**-**F5**) High resolution time-series that highlights bacterial growth

within an IBC prior to (**F1**, **F2**) and after (**F3-F5**) administration of ampicillin. Some bacteria

within this IBC are subsequently eliminated (**F5**). (**G1**-**G5**) High resolution time-series that

highlights bacterial growth within an IBC prior to (**G1**, **G2**) and after (**G3-G5**) administration of

ampicillin. The bacterial volume within this IBC is not diminished by antibiotic treatment (**G5**).

(**H**) Plots of logarithm of bacterial volume within n=11 IBCs before, during, and after the second

round of antibiotic treatment. (**I**) Plot of the volume of extracellular bacteria upon antibiotic

administration from n=103 across 206 x 206 μm^2 field of view from n=3 bladder-chips. (**J**)

Scatter plot of the extracellular bacterial volume (n=105) and bacterial volume within IBCs

(n=22) after antibiotic treatment as a fraction of the maximum bacterial volume prior to

antibiotic treatment. Red line represents the median value, p=1E-9 as calculated by Mann-

Whitney test. (**K**, **L**) Representative images from the epithelial face of the infected bladder-chips

with or without duty cycle. (**M**) Scatter plots of the logarithm of the total bacterial area across

n=14 from infected bladder-chips with (n=2) and without (n=2) duty cycle. p= 8.6E-6 as

calculated by the Mann-Whitney test. Infected bladder chips shown in **K**, **L** were fixed 13 hours

after UPEC infection and 6 hours into the 1st growth cycle. Scale bars, 10 μm in panels **A1**-**A5**,

F1-**F5**, and **G1**-**G5**.

Cell type	Cell density/$10^4 \mu m^2$
Epithelial	31.7 ± 5.1
Endothelial	4.0 ± 0.6

639 **Table 1.**

640 Characterization of epithelial and endothelial cell densities from n=2 bladder-chips

Acknowledgements

V.V.T gratefully acknowledges support by a Human Frontier Science Program (HFSP) Long-Term Fellowship (LT000231/2016-L) and a European Molecular Biology Organization (EMBO) Long-Term Fellowship (921-2015). This research was supported by a grant awarded to J.D.M by the Swiss National Science Foundation (SNSF) (Project Funding, 310030B_176397). The authors thank the entire team of EPFL Bioimaging & Optics Core Facility for their assistance in confocal live cell imaging and post analysis in Bitplane Imaris. The authors also acknowledge the tremendous help provided by Dr. Graham Knott and Marie Croisier (at the EPFL Biological Electron Microscopy Facility) for help in optimizing the protocol for TEM of infected bladder-chip. The authors would like to thank and credit BioRender.com for the illustrations and schematics used in this manuscript.

654   **Materials and Methods**

Reagent or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background ()	Uropathogenic Escherichia coli (UPEC) strain CFT073	PMID: 2182540	NCBI: txid199310	originally isolated from a pyelonephritis patient and provided by Prof. H.L.T. Mobley, University of Michigan, USA
strain, strain background ()	CFT073-pZA32-YFP	this paper		constructed by Dr. Neeraj Dhar (this study)
biological sample (*Mus musculus*)	HTB-9™ bladder epithelial cells	ATCC	Cat#:5637	
biological sample (*Mus musculus*)	HMVEC-Bd – Human Bladder Microvascular Endothelial Cells	Lonza	Cat#:7016	
other	RPMI-1640 medium	ATCC	Cat#:30-2001	
other	Gibco™ RPMI 1640 Medium, no phenol red	Thermofisher	Cat#:11835063	
other	Gibco™ Fetal Bovine Serum, Premium Plus	Thermofisher	Cat#: A4766801	
other	EGM™-2 MV Microvascular Endothelial Cell Growth Medium-2 BulletKit™	Lonza	Cat#:CC-3202	
other	EBM-PRF Endothelial Medium Phenol-red free, 500 ML	Lonza	Cat#:CC-3129	
other	EGM™-2 Endothelial SingleQuots™ Kit	Lonza	Cat#:CC-4176	
other	Gibco™ Antibiotic-Antimycotic (100X)	Thermofisher	Cat#:15240062	
other	Gibco™ Trypsin-EDTA (0.05%), phenol red	Thermofisher	Cat#:25300054	
drug	Chloramphenicol			34 mg/ml in ethanol (stored at -20 ^{o}C)
drug	Ampicillin	Sigma-Aldrich	Cat#: A9518-5G	50 mg/ml in ddH$_2$O (stored at -80 ^{o}C)
other	pooled human female urine	Golden West Diagnostics	Cat#: OH2010-pH	
other	Phosphate Buffered Saline	Thermofisher		
other	Gibco™ HEPES	Thermofisher	Cat#: F2006	
other	Invitrogen™ DAPI	Thermofisher	Cat#: D1306	5 mg/ml in DMSO
other	LB Broth (Luria low salt)	Sigma-Aldrich	Cat#: L3397-1KG	
other	Organ-chips (Standard Research Kit - 24 per	Emulate	RE1000001024	Emulate, Inc. 27 Drydock Ave, 5th

				Floor Boston, MA 02210
other	CellMask™ Orange Plasma membrane Stain	Thermofisher	Cat#: C10045	
other	HCS CellMask™ Deep Red Stain	Thermofisher	Cat#: H32721	
other	CellTracker™ Deep Red Dye	Thermofisher	Cat#: C34565	
other	Elveflow® OB1 Pressure Controller OB1 Base MkIII+	Elveflow	Cat#: OB1MKIII+-MIX-	connected to 6bar compressed air channel
other	Elveflow® OB1 Pressure Controller OB1 MkIII Channel-900/1000	Elveflow	Cat#: OB1-Dual-	connected to Elveflow® OB1 Pressure Controller OB1 Base MkIII+
other	Diaphragm moist gas vacuum pump	KNF Neuberger	Cat#:LABOPORT® UN 820.3 FT.40P	connected to Elveflow® OB1 Pressure Controller OB1 MkIII Channel-900/1000
antibody	Anti-EpCAM	abcam	Cat#: ab71916	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Rabbit secondary antibody, Alexa Fluor 647 (2µg/ml) and Alexa Fluor 568 (2µg/ml) in 1%BSA
antibody	Anti-PECAM1 or Anti-CD31	abcam	Cat#: ab24590	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Mouse secondary antibody, Alexa Fluor 647 (2µg/ml) and Goat anti-Mouse Alexa Fluor 488 (2µg/ml) in 1%BSA
antibody	Anti-CK7	abcam	Cat#: ab209599	used at 1:100 dilution in 1% BSA
antibody	Anti-CK8	abcam	Cat#: ab192468	used at 1:100 dilution in 1% BSA
other	Alexa Fluor™ 555 Phalloidin	Thermofisher	Cat#: A34055	used at 1µM concentration in PBS
antibody	Anti-VE-Cadherin	abcam	Cat#: ab33168	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Rabbit secondary antibody, Alexa Fluor 647 (2µg/ml)

				and Alexa Fluor 488 (2µg/mL) in 1%BSA
antibody	Anti-Uroplakin3a	Santa Cruz	Cat#: sc-166808	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Mouse secondary antibody, Alexa Fluor 647 (2µg/ml) in 1%BSA
antibody	Anti-CK1	Thermofisher	Cat#: MA1-06312	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Mouse secondary antibody, Alexa Fluor 647 (2µg/ml) in 1%BSA
antibody	Anti-E-Cadherin	Thermofisher	Cat#: MA5-14408	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Mouse secondary antibody, Alexa Fluor 647 (2µg/ml) in 1%BSA
antibody	Anti-ZO-1	abcam	Cat#: ab216880	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Rabbit secondary antibody, Alexa Fluor 647 (2µg/ml) in 1%BSA
antibody	Anti-myeloperoxidase	abcam	Cat#: ab9535	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Rabbit secondary antibody, Alexa Fluor 568 (2µg/ml) in 1%BSA
antibody	Anti-neutrophil elastase	abcam	Cat#: ab68672	used at 1:100 dilution in 1% BSA, conjugated with Donkey anti-Rabbit secondary antibody, Alexa Fluor 568 (2µg/ml) in 1%BSA
antibody	Anti-CD15	abcam	Cat#: ab665	used at 1:100 dilution in 1% BSA, conjugated with

				Donkey anti-Mouse secondary antibody, Alexa Fluor 568 (2µg/ml) in 1%BSA
antibody	Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 647	Thermofisher	Cat#: A-31571	2 mg/ml
antibody	Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 647	Thermofisher	Cat#: A-31573	2 mg/ml
antibody	Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	Thermofisher	Cat#: A-11029	2 mg/ml
antibody	Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 568	Thermofisher	Cat#: A10042	2 mg/ml
antibody	Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 568	Thermofisher	Cat#: A10037	2 mg/ml
peptide, recombinant protein	Fibronectin from human plasma	Sigma-Aldrich	Cat#: F1056	
peptide, recombinant protein	Native Collagen, Bovine dermis	AteloCell	Cat#: IAC-50	5 mg/ml
peptide, recombinant protein	Human IL-1α, research grade	Miltenyi Biotec	Cat#:130-093-894	1 µg/ml
peptide, recombinant protein	Human IL-1β, research grade	Miltenyi Biotec	Cat#:130-093-895	1 µg/ml
peptide, recombinant protein	Human IL-6, research grade	Miltenyi Biotec	Cat#:130-093-929	1 µg/ml
peptide, recombinant protein	Human IL-8, research grade	Miltenyi Biotec	Cat#:130-122-354	1 µg/ml
commercial assay or kit	MACSxpress® Whole Blood Neutrophil Isolation Kit, human	Miltenyi Biotec	Cat#:130-104-434	
software, algorithm	Imaris 9.5.1	Bitplane		
other	Gas chamber for stages with k-frame insert (160x110mm) - magnetic model with sliding lid.	okolab	Cat#: H201-K-FRAME	
other	Custom holder for 24mm x 60mm coverslip	okolab	Cat#:1x24by60-M	
other	µ-Slide 8 Well	ibidi	Cat#:80826	

other	Masterflex PharMed tubing, 0.89mm ID, 100 ft	Cole-Palmer	Cat#:GZ-95709-26	
other	Masterflex® Transfer Tubing, Tygon® 0.76mm ID	Cole-Palmer	Cat#:GZ-06419-03	
other	1.30 x 0.75 x 10 mm metallic tubes	Unimed	Cat#:9084 / 200.010-A	
other	1.00/0.75 x 20mm metallic tubes	Unimed	Cat#:200.010-A	
other	Aladdin programmable pump	WPI	Cat#: PUMP-NE-1000	

Cell culture of human bladder epithelial and bladder endothelial cells

The HTB9 human bladder epithelial carcinoma cell line (procured from ATCC, 5637) was

cultured in RPMI 1640 medium (ATCC 30-2001) supplemented with 10% Fetal Bovine Serum

(FBS) as recommended by the supplier. Human Bladder Microvascular Endothelial cells

(HMVEC-Bd) (procured from Lonza, CC-7016) were cultured in Lonza EGM-2 MV BulletKit

(CC-4147, CC-3156) medium. Both the epithelial and endothelial cells were generally cultured

in their respective medium supplemented with 1X Antibiotic-Antimycotic (ThermoFisher

15240062). Epithelial cells were passaged by detachment with 0.05% Trypsin (Gibco) at 37°C

for 3-5 minutes followed by neutralization of trypsin with RPMI 1640 medium/10% FBS.

Bladder endothelial cells were passaged and split as recommended by the supplier. The cells

used in all the experiments were at ten passages or fewer. The cell lines were tested routinely for

mycoplasma contamination during passaging.

UPEC culture for infection of bladder epithelial cells in the bladder-chip

Uropathogenic *Escherichia coli* (UPEC) strain CFT073 was originally isolated from a

pyelonephritis patient (Mobley et al., 1990) and provided by Prof. H.L.T. Mobley, University of

Michigan, USA. A derivative strain expressing yellow fluorescent protein (YFP) was generated

by electroporation of CFT073 with the episomal plasmid pZA32-YFP, as described earlier (Dhar

et al., 2015) . To induce expression of type 1 pili, UPEC was grown in LB media containing 25

µg/ml chloramphenicol under non-shaking conditions at 37 ^{0}C for 2 days prior to the experiment,

to achieve a stationary phase culture (OD_{600}=1.5 and corresponding to a concentration of 1.5 x

10^9 bacteria/ml). The bacteria were diluted 10-fold to a final concentration of 1.5 x 10^8 cells/ml)

and resuspended in a solution of pooled human female urine (Golden West Diagnostics

Catalogue OH2010-pH) diluted 10-fold in Phosphate Buffered Saline (PBS, Gibco).

Recapitulation of human bladder physiology in human bladder-on-a-chip device

Bladder-chip devices made of polydimethylsiloxane (PDMS) were purchased from Emulate

(Boston, USA). The dimensions of the microfluidic device were as follows: width of the

channels- 1000 µm, height of the upper channel- 1000 µm, lower channel – 250 µm). For

extracellular matrix (ECM) coating, a coating solution consisting of the ER-1 compound

(Emulate) dissolved in ER-2 solution at 0.5 mg/mL (Emulate) was introduced in both apical and

vascular channels and the chips were subsequently activated by exposing the bladder-chip for 20

minutes under UV light. The channels were then rinsed with fresh coating solution and the

protocol was repeated once. The channels of the bladder-chip were then washed thoroughly with

PBS before incubation with an ECM solution of 150 µg/ml bovine collagen type I (AteloCell,

Japan,) and 30 µg/ml fibronectin from human plasma (Sigma-Aldrich) in PBS buffered with 15

mM HEPES solution (Gibco Catalogue F2006) for 1-2 hours at 37°C as described previously

(Thacker et al., 2020). If not used directly, coated chips were stored at 4°C and pre-activated

before use by incubation for 30 minutes with the same ECM solution at 37°C. Three days before

the day of the experiment, HMVEC-Bd cells were seeded into lower channel of the inverted

bladder-chip device at 0.5 million cells/ml. Two days before the experiment, HTB9 cells were

seeded into upper channel at 5 million cells/ml. The bladder endothelial and bladder epithelial

cells were seeded into the device three and two days before the experiment in their respective

medium supplemented with the 1X Antibiotic-Antimycotic solution. The antibiotics were

removed prior to the experiment.

Characterization and immunostaining of human bladder epithelial and human bladder

epithelial cells in ibidi wells and human bladder-chip

Human bladder epithelial and bladder endothelial cells were cultured inside ibidi 8-wells for one

day. The cells were subsequently fixed with 4% paraformaldehyde (PFA) for 1 hr at room

temperature. Fixed cells were then washed three times with 200 µL of PBS to remove residues of

PFA, permeabilized with 0.15 % Triton X-100 for 15 minutes; washed three times with 200 µL

of PBS to remove residues of detergent and then incubated in a blocking solution of 1% BSA in

PBS for 1 hr. The cells were then incubated with primary antibodies (Anti-EpCAM, Anti-

PECAM1, Anti-CK7, Anti-CK8, Anti-VE-Cadherin, Anti-Uroplakin3a, Anti-CK1, Anti-E-

Cadherin, Anti-ZO-1, Anti-myeloperoxidase, Anti-neutrophil elastase, Anti-CD15) at a dilution

of 1:100 in an antibody incubation buffer comprising 1% BSA and 0.01% Triton-100 in PBS.

The ibidi wells were subsequently washed three times with PBS for 10 minutes each. Incubation

with secondary antibody (Donkey anti-Mouse Alexa Fluor 647, Donkey anti-Mouse Alexa Fluor

568, Goat anti-Mouse Alexa Fluor 488, Donkey anti-Rabbit Alexa Fluor 647, Donkey anti-

Rabbit Alexa Fluor 568, Donkey anti-Rabbit Alexa Fluor 488) at a concentration of 2 µg/mL in

antibody incubation buffer was subsequently performed for 1 hour at room temperature. Excess

antibody was removed by washing three times with PBS for 10 minutes each. Cell nuclei were

stained with DAPI (5 µg/mL, Sigma) for 30 minutes. Ibidi 8-wells were washed three times to

remove unbound DAPI. Cells were covered with appropriate volume of PBS until imaging. Cells

were imaged with 63X oil objective on Leica SP8 confocal microscope. Images were

deconvolved using SVI Huygens (Quality, 0.05; Iterations, 40).

For characterization of the cell types on-chip, human bladder-chips with co-culture of bladder

epithelial and bladder endothelial cells were kept under pooled diluted urine and EBM2 medium

using P200 pipette tips for 2 days prior to PFA fixation. The cells inside human bladder chip

were then immunostained following the protocol described above. Images were acquired with

Leica HC FLUOTAR 25X (NA 0.95) multi-immersion objective on Leica SP8 confocal

microscope.

Characterization of strain-pressure curve in the bladder-chip model

Elveflow OB1 MK3 – Microfluidic flow control system was used to control the negative

pressure applied to the human bladder-chip. The control system was connected to the

compressed air line (6 bar) for the positive pressure and diaphragm vacuum pump (KNF

Neuberger) for the negative pressure. A negative pressure (0 to -900 mbar) with step function of

-100mbar was subsequently applied to the vacuum channels in the bladder-chip using a Pressure

Controller (Elveflow OB1 pressure controller). For these experiments, both bladder epithelial

and endothelial cells were seeded on the respective sides in the devices. This experiment was

performed with the chip maintained on the stage of the microscope and a brightfield image was

acquired at each step increase of -100mbar, on Leica SP8 confocal microscope. The PDMS inter-

740 pore-to-pore distance was measured for 14 pore-to-pore combinations at each input of negative

741 pressure. Linear fitting was performed using GraphPad Prism (version 9).

743 **Mimicking bladder filling and voiding cycle in human bladder-chip**

744 The stretching of the human bladder-chip was done using the Elveflow OB1 MK3 – Microfluidic

745 flow control system connected to the compressed air line (6 bar) for the positive pressure and

746 diaphragm vacuum pump (KNF Neuberger) for the negative pressure. Human bladder-chip in the

747 relaxed (*voided bladder*) state was slowly perturbed with a linear ramp function to reach a

748 stretched (*filled bladder*) state over a period of 2 hr. This period corresponded to the *filling*

749 *bladder* state in Figure 1D. Bladder-chip was kept under stretched (*filled bladder*) state over the

750 subsequent 2 hours. Micturition or urination was recapitulated by rapidly reducing the applied

751 strain on the bladder-chip from the stretched (*filled bladder*) state to relaxed (*voided bladder*)

752 state over a period of 2 minutes. Subsequently, the bladder-chip was maintained with no negative

753 pressure applied in the vacuum channel under relaxed (*voided bladder*) state from (4hr + 2min to

754 6hr). This 6-hour bladder filling and voiding cycle with a duty cycle of 6hr was repeated

755 continuously for the rest of the experiment.

756 **Experimental setup and imaging parameters for time lapse imaging for UPEC infection in**

757 **the human bladder-chip devices**

758 The bladder-chip device was connected to the syringe pumps via gas impermeable PharMed®

759 tubing (inner diameter =0.89 mm, Cole palmer GZ-95809-26) along with longer transparent

760 Tygon tubings (internal diameter of 0.76mm, Masterflex transfer tubing Cole palmer GZ-06419-

03). The Harmed® tubing was connected to the inlet and outlet of the bladder-chip with 1.30 x

0.75 x 10 mm metallic tubes (Unimed) and transparent tubing with 1.00/0.75 x 20mm metallic

tubes (Unimed). The bladder-chip connected to external sources of flow was then mounted onto

a 24 x 60 mm No. 1 glass coverslip for microscopy imaging. When required for stretching

experiments, tubing was also connected to the stretching channels on either side of the main

channel. The connected device was subsequently assembled inside a temperature-controlled

microscope environmental chamber at $37^{o}C$ supplemented with 5% CO_2 (OKO labs H201-K-

Frame). Time-lapse imaging was conducted using a Leica HC FLUOTAR 25X (NA 0.95) multi-

immersion objective within custom made environmental chamber set at $37^{o}C$ for infusion with

syringe pumps. Water was pumped to the ring around the water objective at 9 Hz with pumping

duration of 9 seconds and pumping interval 30 minutes, controlled by SRS software (HRZ=9,

VPP=95). The autofocus mode (best focus, steps =9, range = 30 µm) was used to maintain the

optical focus on the apical side of the PDMS membrane. The experiments were monitored

frequently to ensure that the optical focus was maintained, and the experiment was halted and

restarted if the focus was lost.

To enable rapid 3-D imaging across multiple spatial locations on-chip, we utilized the capability

of the white light laser on the Leica SP8 confocal microscope to image at multiple wavelengths

simultaneously. The excitation wavelengths were grouped into two sequences to minimize the

spectral overlap. In the first sequence, laser emission at 555 nm was used to excite the Calmest

Orange stain in the bladder epithelial cells. In the second laser excitation sequence, laser

emission at 500 nm and 630 nm were used to excite the YFP within the bacteria and the

cytoplasmic CellTracker[TM] Deep Red in the human neutrophils. Images were acquired with a

scan speed of 400-700 Hz and a zoom factor of 2.25 (206.67 µm x 206.67 µm) resulting in an

XY resolution of 450 nm depending on the number of pixels acquired per field of view. Z-stacks

were acquired with 1 µm step sizes. Time lapse images were acquired with interval duration of

ca. 15-17 minutes, in a subset of experiments this was further reduced to 7.5 minutes by imaging

with the second sequence only at a z-step size of 2 µm.

Time lapse imaging for UPEC infections in the human bladder-chip devices

<u>Pre-infection stage</u>

The time-course of the entire experimental protocol is shown in a schematic in Figure 1. The

bladder-chip device was perfused with diluted urine in the apical channel and EBM Endothelium

phenol-red free medium supplemented with EGM-2 endothelial SingleQuots kit in the vascular

channel. Prior to the commencement of infection, the chip was maintained at homeostasis and

the epithelial cells were imaged for a period of two hours.

<u>Infection stage</u>

Stationary phase UPEC in diluted pooled urine at a concentration of 150 million cells/mL were

flowed through the apical channel of the device at 1200 µL/hour for 1.5 hours. During this

period, EBM2 media was flowed through the endothelial channel at 600 µL/hour.

<u>Bacterial washout and neutrophil introduction stage</u>

Next, the syringe connected to the apical channel was replaced with a fresh syringe containing

pooled diluted urine, this solution was then flowed in the apical channel over the epithelial layer

of the chip at a flow rate of 600 µL/hour over the next 3 hours. This allowed for the continuous

removal of extracellular planktonic bacteria in the apical channel. At the same time, human

neutrophils were introduced into the endothelial channel of the bladder-chip via flow. A solution

of human neutrophils at a density of 2 million cells/mL isolated via negative selection was

flowed through the vascular channel for 2 hours in EBM2 medium with at higher shear stress of

$\eta=1.0$ dyne/cm^2 to enhance neutrophil attachment to endothelial cells.

<u>Neutrophil diapedesis stage</u>

Diapedesis of human neutrophils to epithelial side and subsequent interactions of neutrophils

with UPEC was observed for the subsequent 3 hours since introduction of neutrophils into the

vasculature side of the chip. During this period, the flow rate in the apical and vascular channels

were maintained at 600 µL/hr and 3000 µL/hr respectively. During this period, in a subset of

experiments, only the channels that were part of the second laser scanning sequence were

imaged. Images of the CellTracker Orange dye for epithelial cell identification were not acquired

(first laser scanning sequence). This enabled a number of fields of view to be captured with an

enhanced temporal resolution and a frame rate of up to 7.5 minutes. For experiments studying

the formation of NETs, the experiment was halted at this stage and the infected chips were fixed,

permeabilized, blocked, and immunostained with anti-myeloperoxidase (abcam) or anti-

neutrophil elastase (abcam) using the procedure described earlier. For all other experiments, live

824 imaging continued to the subsequent antibiotic treatment phase, Immunofluorescence for anti-

825 myeloperoxidase and anti-neutrophil elastase was done as the procedure described previously.

827 <u>Antibiotic treatment and growth cycles</u>

828 Thereafter, syringes connected to both apical and vascular channels were changed and ampicillin

829 at 250 µg/mL (used at ~40-fold over the minimum inhibitory concentration (MIC) of ampicillin

830 against UPEC grown in EBM2 medium) was introduced in the diluted urine and the EBM2

831 media perfused into the apical and vascular channels, respectively, for 3 hours at a flow rate of

832 600 µl/hour. This was the first ampicillin treatment cycle; whose purpose was to eliminate

833 extracellular bacteria in the apical channel and allow intracellular bacterial colonies (IBCs) to be

834 identified. The medium was subsequently switched with antibiotic-free medium for next 8 hours

835 to allow for IBC growth within epithelial cells. During this period, the flow within the apical

836 channel was maintained, to remove bacteria that grew extracellularly either because they

837 survived the antibiotic treatment or that were released from infected epithelial cells. The

838 ampicillin and growth cycle were then repeated, to allow the assessment of the response of

839 bacteria within IBCs to antibiotic treatment as well as characterize the subsequent regrowth.

841 **Image analysis of confocal live-cell images**

842 Image analysis was performed with Bitplane Imaris 9.5.1. The time-lapse imaging stack included

843 five channels: uroepithelial cells (epithelial channel), transmitted light (bright field channel),

844 UPEC (bacterial channel), neutrophils (neutrophil channel), and transmitted light (bright field

channel). Neutrophils were identified via the spot detection algorithm in Imaris used on the

images from the neutrophil channel with the following parameters (Size, 8 µm; Quality, 4 to 8).

Swarms of neutrophils generate dense aggregates that are ill-suited to quantification with the

spot detection algorithm. To quantify the size of the swarms, we therefore segmented images in

the neutrophil channel to generate surfaces via the automatic segmentation tool in Imaris. with

the following parameters (Threshold, 10; Smooth Surfaces Detail, 0.5 or 1.0 µm). Unfortunately,

the time resolution was insufficient to track individual neutrophils over time particularly since

neutrophils formed small clumps and rapidly changed their cell shapes.

For UPEC volume inside IBCs, necessary 3D volume was cropped in Imaris to ease image

analysis. Total bacterial volume inside IBC was detected by creating the surface (Threshold, 15;

Smooth Surfaces Detail, 0.5 or 1 µm) on UPEC channel.

In cases of measuring extracellular UPEC growth, surface was generated (Threshold, 15; Smooth

Surfaces Detail, 0.5 or 1 µm) on UPEC channel to calculate total extracellular volume in the

field of view.

Infection experiments with relaxed and pre-stretched bladder chip

Human bladder-chip was kept in stretching state (ca. 10 % strain) for 2hr using Elveflow OB1

microfluidic control system. Post attainment of the stretched state, 200 µL of UPEC inoculum

(150 million cells per mL) was introduced into the apical channel with 200 µL tips. During this

period, EBM2 medium was maintained under static culture conditions. UPEC infection and

attachment to the epithelial layer was then performed as described earlier during the stretched

state. Subsequently, the pre-stretched bladder-chip was brought to relaxed state (0% strain)

within 2 minutes. The bladder-chip was then incubated with Gentamicin (100 µg/ml) for 3 hours

introduced both on the apical and vascular channels of the bladder-chip to prevent extracellular

growth. Similarly, a bladder-chip without any stretch applied to the membrane was also infected

and processed as above. The bladder-chip was then washed three times with PBS, fixed,

permeabilized, and then stained with the HCS Cell Mask ^{TM} Deep red and imaged on a Leica

SP8 confocal microscope. Bacterial surfaces were generated in Imaris (Threshold, 15; Smooth

Surfaces Detail, 1.0 or 2.0 µm).

Isolation and Labelling of human neutrophils from fresh peripheral human blood

Primary human neutrophils were isolated via negative depletion method from human peripheral

blood with MACSxpress® Whole Blood Neutrophil Isolation Kit (Miltenyi Biotec), following

the manufacturers instructions. Isolation was performed without the use of a density-based

centrifugation method. Isolated human neutrophils were then incubated with a 1µM solution

CellTracker^{TM} Deep Red in a serum free RPMI phenol red free medium for 30 minutes in a cell

culture incubator maintained at 37 C and 5% CO_2 Labelled human neutrophils were then washed

with 10ml of 20% FBS in RPMI phenol red free medium twice to remove the unbound dye. The

human neutrophils were suspended in Lonza EBM2 medium at a cell density of 1 million

cells/ml. In some instances, labelled human neutrophils were passed through a filter with 40 µm

pores (Thermofisher) to remove neutrophil clusters that may have formed during the isolation

process.

Scanning electron microscopy of human bladder-chip

UPEC were allowed to grow inside UPEC-infected epithelial cells in human bladder-chip was

allowed for 6 hours (14 hr from start of UPEC infection) during the IBC 1st growth cycle. After 6

hours, human bladder-chip was fixed at room temperature for 1 hr with a mix of 1%

glutaraldehyde and 2% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4). The fixed bladder-

chip was kept in the fixative overnight (at 4°C). Post overnight fixation, human bladder-chip was

cut open from the apical channel side. A scalpel was used to cut approximately in the middle of

the apical channel side (height-1mm) to expose the bladder epithelial cells. The fixed chip was

further fixed for 30 minutes in 1% osmium tetroxide in 0.1 M cacodylate buffer followed by

washing with the distilled water. Next, the bladder-chip was dehydrated in a graded alcohol

series and dried by passing them through the supercritical point of carbon dioxide (CPD300,

Leica Microsystems). Finally, the bladder-chip was attached to an adhesive conductive surface

followed by coating with a 3 – 4 nm layer of gold palladium metal (Quorum Q Plus, Quorum

Technologies). Images of the cells were captured using a field emission scanning electron

microscope (Merlin, Zeiss NTS).

In case of uninfected human bladder-chip (control), the human bladder chip was fixed at the

same time point (14 hr from start of the experiment). For the case of NETs formed on epithelial

layer, human bladder chip was fixed at 2 hours post neutrophil introduction into the endothelium

side of the chip.

Preparation for transmission electron microscopy (TEM)

UPEC within infected human bladder-chips were allowed to proliferate for 6 hours (14 hr from

start of UPEC infection) during the IBC 1st growth cycle. After 6 hours, human bladder-chip was

fixed at room temperature for 1 hr with a mix of 1% glutaraldehyde and 2% paraformaldehyde in

0.1 M phosphate buffer (pH 7.4). The fixed bladder-chip was kept in the fixative overnight (at

4°C). Post overnight fixation, human bladder-chip was cut open from the apical channel side. A

scalpel was used to cut approximately in the middle of the apical channel side (height-1mm) to

expose the bladder epithelial cells. The bladder-chip was then washed in cacodylate buffer

(0.1M, pH 7.4), postfixed for 40 minutes in 1.0 % osmium tetroxide with 1.5% potassium

ferrocyanide, and then fixed again with 1.0% osmium tetroxide alone. The bladder-chip was

finally stained for 30 minutes in 1% uranyl acetate in water before being dehydrated through

increasing concentrations of alcohol and then embedded in Durcupan ACM (Fluka, Switzerland)

resin. The bladder-chip was then placed in Petri dishes so that approximately 1 mm of resin

remained above the cells, and the dish than left in an oven at 65°C for 24 hours. Regions on

interest, and corresponding to structures imaged with light microscopy were trimmed from the

rest of the device, once the resin had hardened, and thin, 50 nm-thick sections were cut with a

diamond knife, and collected onto single-slot copper grids with a pioloform support film. These

were contrasted with lead citrate and uranyl acetate, and images taken with a transmission

electron microscope at 80 kV (Tecnai Spirit, FEI Company with Eagle CCD camera).

Data availability statement
The datasets generated during and analysed during the current study are available from the

corresponding authors on reasonable request and will be uploaded to Zenodo prior to publication.

Bibliography

Abraham, S.N., and Miao, Y. (2015). The nature of immune responses to urinary tract
infections. Nat Rev Immunol *15*, 655–663.

Agace, W.W., Hedges, S.R., Ceska, M., and Svanborg, C. (1993). Interleukin-8 and the
neutrophil response to mucosal gram-negative infection. J Clin Invest *92*, 780–785.

Agace, W.W., Patarroyo, M., Svensson, M., Carlemalm, E., and Svanborg, C. (1995).
Escherichia coli induces transuroepithelial neutrophil migration by an intercellular adhesion
molecule-1-dependent mechanism. Infect Immun *63*, 4054–4062.

Alon, R., Hammer, D.A., and Springer, T.A. (1995). Lifetime of the P-selectin-carbohydrate
bond and its response to tensile force in hydrodynamic flow. Nature *374*, 539–542.

Andersen, T.E., Khandige, S., Madelung, M., Brewer, J., Kolmos, H.J., and Møller-Jensen, J.
(2012). Escherichia coli uropathogenesis in vitro: invasion, cellular escape, and secondary
infection analyzed in a human bladder cell infection model. Infect Immun *80*, 1858–1867.

Anderson, G.G., Palermo, J.J., Schilling, J.D., Roth, R., Heuser, J., and Hultgren, S.J. (2003).
Intracellular Bacterial Biofilm-Like Pods in Urinary Tract Infections. Science *301*, 105–107.

Apodaca, G. (2004). The Uroepithelium: Not Just a Passive Barrier. Traffic *5*, 117–128.

Benam, K.H., Villenave, R., Lucchesi, C., Varone, A., Hubeau, C., Lee, H.-H., Alves, S.E.,
Salmon, M., Ferrante, T.C., Weaver, J.C., et al. (2016). Small airway-on-a-chip enables analysis
of human lung inflammation and drug responses in vitro. Nat Methods *13*, 151–157.

Blango, M.G., and Mulvey, M.A. (2010). Persistence of Uropathogenic Escherichia coli in the
Face of Multiple Antibiotics ▯. Antimicrob Agents Ch *54*, 1855–1863.

Branzk, N., Lubojemska, A., Hardison, S.E., Wang, Q., Gutierrez, M.G., Brown, G.D., and
Papayannopoulos, V. (2014). Neutrophils sense microbe size and selectively release neutrophil
extracellular traps in response to large pathogens. Nat Immunol *15*, 1017–1025.

Brinkmann, V., Reichard, U., Goosmann, C., Fauler, B., Uhlemann, Y., Weiss, D.S., Weinrauch,
Y., and Zychlinsky, A. (2004). Neutrophil Extracellular Traps Kill Bacteria. Science *303*, 1532–
1535.

Bryskier, A. (2020). Antimicrobial Agents: Antibacterials and Antifungals. pp. 113–162.

Cakir, B., Xiang, Y., Tanaka, Y., Kural, M.H., Parent, M., Kang, Y.-J., Chapeton, K., Patterson,
B., Yuan, Y., He, C.-S., et al. (2019). Engineering of human brain organoids with a functional
vascular-like system. Nat Methods *16*, 1169–1175.

Carattino, M.D., Prakasam, H.S., Ruiz, W.G., Clayton, D.R., McGuire, M., Gallo, L.I., and
Apodaca, G. (2013). Bladder filling and voiding affect umbrella cell tight junction organization
and function. Am J Physiol-Renal *305*, F1158–F1168.

Clevers, H. (2016). Modeling Development and Disease with Organoids. Cell *165*, 1586–1597.

Duncan, M.J., Li, G., Shin, J.-S., Carson, J.L., and Abraham, S.N. (2004). Bacterial Penetration
of Bladder Epithelium through Lipid Rafts. J Biol Chem *279*, 18944–18951.

Ermert, D., Urban, C.F., Laube, B., Goosmann, C., Zychlinsky, A., and Brinkmann, V. (2009).
Mouse Neutrophil Extracellular Traps in Microbial Infections. J Innate Immun *1*, 181–193.

Foxman, B. (2010). The epidemiology of urinary tract infection. Nat Rev Urol *7*, 653–660.

Gloeckner, D.C., SACKS, M.S., FRASER, M.O., SOMOGYI, G.T., GROAT, W.C. de, and
CHANCELLOR, M.B. (2002). Passive Biaxial Mechanical Properties of the Rat Bladder Wall
After Spinal Cord Injury. J Urology *167*, 2247–2252.

Grassart, A., Malardé, V., Gobba, S., Sartori-Rupp, A., Kerns, J., Karalis, K., Marteyn, B.,
Sansonetti, P., and Sauvonnet, N. (2019). Bioengineered Human Organ-on-Chip Reveals
Intestinal Microenvironment and Mechanical Forces Impacting Shigella Infection. Cell Host
Microbe *26*, 435-444.e4.

Hannan, T.J., Totsika, M., Mansfield, K.J., Moore, K.H., Schembri, M.A., and Hultgren, S.J.
(2012). Host–pathogen checkpoints and population bottlenecks in persistent and intracellular
uropathogenic Escherichia coli bladder infection. Fems Microbiol Rev *36*, 616–648.

Haraoka, M., Hang, L., Frendéus, B., Godaly, G., Burdick, M., Strieter, R., and Svanborg, C.
(1999). Neutrophil Recruitment and Resistance to Urinary Tract Infection. J Infect Dis *180*,
1220–1229.

Hedges, S., Svensson, M., and Svanborg, C. (1992). Interleukin-6 response of epithelial cell lines
to bacterial stimulation in vitro. Infect Immun *60*, 1295–1301.

Horsley, H., Dharmasena, D., Malone-Lee, J., and Rohn, J.L. (2018). A urine-dependent human
urothelial organoid offers a potential alternative to rodent models of infection. Sci Rep-Uk *8*,
1238.

Hsieh, M.M., Everhart, J.E., Byrd-Holt, D.D., Tisdale, J.F., and Rodgers, G.P. (2007).
Prevalence of Neutropenia in the U.S. Population: Age, Sex, Smoking Status, and Ethnic
Differences. Ann Intern Med *146*, 486.

Huh, D., Matthews, B.D., Mammoto, A., Montoya-Zavala, M., Hsin, H.Y., and Ingber, D.E.
(2010a). Reconstituting Organ-Level Lung Functions on a Chip. Science *328*, 1662–1668.

Huh, D., Matthews, B.D., Mammoto, A., Montoya-Zavala, M., Hsin, H.Y., and Ingber, D.E.
(2010b). Reconstituting Organ-Level Lung Functions on a Chip. Science *328*, 1662–1668.

Hung, C.-S., Dodson, K.W., and Hultgren, S.J. (2009). A murine model of urinary tract
infection. Nat Protoc *4*, 1230–1243.

Hunstad, D.A., and Justice, S.S. (2010). Intracellular Lifestyles and Immune Evasion Strategies
of Uropathogenic Escherichia coli. Annu Rev Microbiol *64*, 203–221.

Iosifidis, G., and Duggin, I.G. (2020). Distinct Morphological Fates of Uropathogenic
Escherichia coli Intracellular Bacterial Communities: Dependency on Urine Composition and
pH. Infect Immun *88*.

Jalili-Firoozinezhad, S., Gazzaniga, F.S., Calamari, E.L., Camacho, D.M., Fadel, C.W., Bein, A.,
Swenor, B., Nestor, B., Cronce, M.J., Tovaglieri, A., et al. (2019). A complex human gut
microbiome cultured in an anaerobic intestine-on-a-chip. Nat Biomed Eng *3*, 520–531.

Jang, K.-J., Mehr, A.P., Hamilton, G.A., McPartlin, L.A., Chung, S., Suh, K.-Y., and Ingber,
D.E. (2013). Human kidney proximal tubule-on-a-chip for drug transport and nephrotoxicity
assessment. Integr Biol *5*, 1119–1129.

Jang, K.-J., Otieno, M.A., Ronxhi, J., Lim, H.-K., Ewart, L., Kodella, K.R., Petropolis, D.B.,
Kulkarni, G., Rubins, J.E., Conegliano, D., et al. (2019). Reproducing human and cross-species
drug toxicities using a Liver-Chip. Sci Transl Med *11*, eaax5516.

Justice, S.S., Hung, C., Theriot, J.A., Fletcher, D.A., Anderson, G.G., Footer, M.J., and Hultgren,
S.J. (2004). Differentiation and developmental pathways of uropathogenic Escherichia coli in
urinary tract pathogenesis. Proc National Acad Sci *101*, 1333–1338.

Kang, Y.B. (Abraham), Rawat, S., Duchemin, N., Bouchard, M., and Noh, M. (2017). Human
Liver Sinusoid on a Chip for Hepatitis B Virus Replication Study. Micromachines-Basel *8*, 27.

Kienle, K., and Lämmermann, T. (2016). Neutrophil swarming: an essential process of the
neutrophil tissue response. Immunol Rev *273*, 76–93.

Kim, H.J., and Ingber, D.E. (2013). Gut-on-a-Chip microenvironment induces human intestinal
cells to undergo villus differentiation. Integr Biol *5*, 1130–1140.

Kim, H.J., Li, H., Collins, J.J., and Ingber, D.E. (2016). Contributions of microbiome and
mechanical deformation to intestinal bacterial overgrowth and inflammation in a human gut-on-
a-chip. Proc National Acad Sci *113*, E7–E15.

Klein, R.D., and Hultgren, S.J. (2020). Urinary tract infections: microbial pathogenesis, host-
pathogen interactions and new treatment strategies. Nat Rev Microbiol *18*, 211–226.

Korkmaz, I., and Rogg, B. (2007). A simple fluid-mechanical model for the prediction of the
stress–strain relation of the male urinary bladder. J Biomech *40*, 663–668.

Kreisel, D., Nava, R.G., Li, W., Zinselmeyer, B.H., Wang, B., Lai, J., Pless, R., Gelman, A.E.,
Krupnick, A.S., and Miller, M.J. (2010). In vivo two-photon imaging reveals monocyte-
dependent neutrophil extravasation during pulmonary inflammation. Proc National Acad Sci
107, 18073–18078.

Lämmermann, T., Afonso, P.V., Angermann, B.R., Wang, J.M., Kastenmüller, W., Parent, C.A.,
and Germain, R.N. (2013). Neutrophil swarms require LTB4 and integrins at sites of cell death
in vivo. Nature *498*, 371–375.

Ley, K., Laudanna, C., Cybulsky, M.I., and Nourshargh, S. (2007). Getting to the site of
inflammation: the leukocyte adhesion cascade updated. Nat Rev Immunol *7*, 678–689.

Martinez, J.J., Mulvey, M.A., Schilling, J.D., Pinkner, J.S., and Hultgren, S.J. (2000). Type 1
pilus-mediated bacterial invasion of bladder epithelial cells. Embo J *19*, 2803–2812.

Metzler, K.D., Fuchs, T.A., Nauseef, W.M., Reumaux, D., Roesler, J., Schulze, I., Wahn, V.,
Papayannopoulos, V., and Zychlinsky, A. (2011). Myeloperoxidase is required for neutrophil
extracellular trap formation: implications for innate immunity. Blood *117*, 953–959.

Mulvey, M.A., Lopez-Boado, Y.S., Wilson, C.L., Roth, R., Parks, W.C., Heuser, J., and
Hultgren, S.J. (1998). Induction and Evasion of Host Defenses by Type 1-Piliated Uropathogenic
Escherichia coli. Science *282*, 1494–1497.

Mysorekar, I.U., and Hultgren, S.J. (2006). Mechanisms of uropathogenic Escherichia coli
persistence and eradication from the urinary tract. Proc National Acad Sci *103*, 14170–14175.

Nagamatsu, K., Hannan, T.J., Guest, R.L., Kostakioti, M., Hadjifrangiskou, M., Binkley, J.,
Dodson, K., Raivio, T.L., and Hultgren, S.J. (2015). Dysregulation of Escherichia coli α-
hemolysin expression alters the course of acute and persistent urinary tract infection. Proc
National Acad Sci *112*, E871–E880.

Nawroth, J.C., Lucchesi, C., Cheng, D., Shukla, A., Ngyuen, J., Shroff, T., Varone, A., Karalis,
K., Lee, H.-H., Alves, S., et al. (2020). A Microengineered Airway Lung Chip Models Key
Features of Viral-induced Exacerbation of Asthma. Am J Resp Cell Mol *63*, 591–600.

Nourshargh, S., Hordijk, P.L., and Sixt, M. (2010). Breaching multiple barriers: leukocyte
motility through venular walls and the interstitium. Nat Rev Mol Cell Bio *11*, 366–378.

Novak, R., Ingram, M., Marquez, S., Das, D., Delahanty, A., Herland, A., Maoz, B.M., Jeanty,
S.S.F., Somayaji, M.R., Burt, M., et al. (2020). Robotic fluidic coupling and interrogation of
multiple vascularized organ chips. Nat Biomed Eng *4*, 407–420.

Oliveira, S. de, Rosowski, E.E., and Huttenlocher, A. (2016). Neutrophil migration in infection
and wound repair: going forward in reverse. Nat Rev Immunol *16*, 378–391.

Papayannopoulos, V., Metzler, K.D., Hakkim, A., and Zychlinsky, A. (2010). Neutrophil
elastase and myeloperoxidase regulate the formation of neutrophil extracellular traps. J Cell
Biology *191*, 677–691.

Parekh, A., Cigan, A.D., Wognum, S., Heise, R.L., Chancellor, M.B., and Sacks, M.S. (2010).
Ex vivo deformations of the urinary bladder wall during whole bladder filling: Contributions of
extracellular matrix and smooth muscle. J Biomech *43*, 1708–1716.

Pascalis, R.D., Parnell, W.J., Abrahams, I.D., Shearer, T., Daly, D.M., and Grundy, D. (2018).
The inflation of viscoelastic balloons and hollow viscera. Proc Royal Soc *474*, 20180102.

Putrinš, M., Kogermann, K., Lukk, E., Lippus, M., Varik, V., and Tenson, T. (2015). Phenotypic
Heterogeneity Enables Uropathogenic Escherichia coli To Evade Killing by Antibiotics and
Serum Complement. Infect Immun *83*, 1056–1067.

Robino, L., Scavone, P., Araujo, L., Algorta, G., Zunino, P., and Vignoli, R. (2013). Detection of
intracellular bacterial communities in a child with Escherichia coli recurrent urinary tract
infections. Pathog Dis *68*, 78–81.

Ronaldson-Bouchard, K., and Vunjak-Novakovic, G. (2018). Organs-on-a-Chip: A Fast Track
for Engineered Human Tissues in Drug Development. Cell Stem Cell *22*, 310–324.

Rosen, D.A., Hooton, T.M., Stamm, W.E., Humphrey, P.A., and Hultgren, S.J. (2007). Detection
of Intracellular Bacterial Communities in Human Urinary Tract Infection. Plos Med *4*, e329.

Rossi, G., Manfrin, A., and Lutolf, M.P. (2018). Progress and potential in organoid research. Nat
Rev Genet *19*, 671–687.

Sato, T., Vries, R.G., Snippert, H.J., Wetering, M. van de, Barker, N., Stange, D.E., Es, J.H. van,
Abo, A., Kujala, P., Peters, P.J., et al. (2009). Single Lgr5 stem cells build crypt-villus structures
in vitro without a mesenchymal niche. Nature *459*, 262–265.

Scott, V.C.S., Haake, D.A., Churchill, B.M., Justice, S.S., and Kim, J.-H. (2015). Intracellular
Bacterial Communities: A Potential Etiology for Chronic Lower Urinary Tract Symptoms.
Urology *86*, 425–431.

Shah, P., Fritz, J.V., Glaab, E., Desai, M.S., Greenhalgh, K., Frachet, A., Niegowska, M., Estes,
M., Jäger, C., Seguin-Devaux, C., et al. (2016). A microfluidics-based in vitro model of the
gastrointestinal human–microbe interface. Nat Commun *7*, 11535.

Si, L., Bai, H., Rodas, M., Cao, W., Oh, C.Y., Jiang, A., Nurani, A., Zhu, D.Y., Goyal, G.,
Gilpin, S.E., et al. (2020). Human organs-on-chips as tools for repurposing approved drugs as
potential influenza and COVID19 therapeutics in viral pandemics. Biorxiv 2020.04.13.039917.

Smith, Y.C., Grande, K.K., Rasmussen, S.B., and O'Brien, A.D. (2006). Novel Three-
Dimensional Organoid Model for Evaluation of the Interaction of Uropathogenic Escherichia
coli with Terminally Differentiated Human Urothelial Cells. Infect Immun *74*, 750–757.

Solis, A.G., Bielecki, P., Steach, H.R., Sharma, L., Harman, C.C.D., Yun, S., Zoete, M.R. de,
Warnock, J.N., To, S.D.F., York, A.G., et al. (2019). Mechanosensation of cyclical force by
PIEZO1 is essential for innate immunity. Nature *573*, 69–74.

Son, K., Mukherjee, M., McIntyre, B.A.S., Eguez, J.C., Radford, K., LaVigne, N., Ethier, C.,
Davoine, F., Janssen, L., Lacy, P., et al. (2017). Improved recovery of functionally active
eosinophils and neutrophils using novel immunomagnetic technology. J Immunol Methods *449*,
44–55.

Song, J., Duncan, M.J., Li, G., Chan, C., Grady, R., Stapleton, A., and Abraham, S.N. (2007). A
Novel TLR4-Mediated Signaling Pathway Leading to IL-6 Responses in Human Bladder
Epithelial Cells. Plos Pathog *3*, e60.

Spaulding, C.N., Klein, R.D., Ruer, S., Kau, A.L., Schreiber, H.L., Cusumano, Z.T., Dodson,
K.W., Pinkner, J.S., Fremont, D.H., Janetka, J.W., et al. (2017). Selective depletion of
uropathogenic E. coli from the gut by a FimH antagonist. Nature *546*, 528–532.

Suzuki, K., Koyanagi-Aoi, M., Uehara, K., Hinata, N., Fujisawa, M., and Aoi, T. (2019).
Directed differentiation of human induced pluripotent stem cells into mature stratified bladder
urothelium. Sci Rep-Uk *9*, 10506.

Thacker, V.V., Sharma, K., Dhar, N., Mancini, G.-F., Sordet-Dessimoz, J., and McKinney, J.D.
(2020a). Rapid endothelial infection, endothelialitis and vascular damage characterise SARS-
CoV-2 infection in a human lung-on-chip model. Biorxiv 2020.08.10.243220.

Thacker, V.V., Dhar, N., Sharma, K., Barrile, R., Karalis, K., and McKinney, J.D. (2020b). A
lung-on-chip model of early M. tuberculosis infection reveals an essential role for alveolar
epithelial cells in controlling bacterial growth. Elife 9, e59961.

Tovaglieri, A., Sontheimer-Phelps, A., Geirnaert, A., Prantil-Baun, R., Camacho, D.M., Chou,
D.B., Jalili-Firoozinezhad, S., Wouters, T. de, Kasendra, M., Super, M., et al. (2019). Species-
specific enhancement of enterohemorrhagic E. coli pathogenesis mediated by microbiome
metabolites. Microbiome 7, 43.

Truschel, S.T., Wang, E., Ruiz, W.G., Leung, S.-M., Rojas, R., Lavelle, J., Zeidel, M., Stoffer,
D., and Apodaca, G. (2002). Stretch-regulated Exocytosis/Endocytosis in Bladder Umbrella
Cells. Mol Biol Cell 13, 830–846.

Wang, E.C.Y., Lee, J.-M., Ruiz, W.G., Balestreire, E.M., Bodungen, M. von, Barrick, S.,
Cockayne, D.A., Birder, L.A., and Apodaca, G. (2005). ATP and purinergic receptor–dependent
membrane traffic in bladder umbrella cells. J Clin Invest 115, 2412–2422.

Wang, J., Wang, C., Xu, N., Liu, Z.-F., Pang, D.-W., and Zhang, Z.-L. (2019). A virus-induced
kidney disease model based on organ-on-a-chip: Pathogenesis exploration of virus-related renal
dysfunctions. Biomaterials 219, 119367.

Yu, Y., Kwon, K., Tsitrin, T., Bekele, S., Sikorski, P., Nelson, K.E., and Pieper, R. (2017).
Characterization of Early-Phase Neutrophil Extracellular Traps in Urinary Tract Infections. Plos
Pathog 13, e1006151.

Zahler, S., Kowalski, C., Brosig, A., Kupatt, C., Becker, B.F., and Gerlach, E. (1997). The
function of neutrophils isolated by a magnetic antibody cell separation technique is not altered in
comparison to a density gradient centrifugation method. J Immunol Methods 200, 173–179.

Zalewska-Piątek, B., Olszewski, M., Lipniacki, T., Błoński, S., Wieczór, M., Bruździak, P.,
Skwarska, A., Nowicki, B., Nowicki, S., and Piątek, R. (2020). A shear stress micromodel of
urinary tract infection by the Escherichia coli producing Dr adhesin. Plos Pathog 16, e1008247.

Zhou, M., Zhang, X., Wen, X., Wu, T., Wang, W., Yang, M., Wang, J., Fang, M., Lin, B., and
Lin, H. (2016). Development of a Functional Glomerulus at the Organ Level on a Chip to Mimic
Hypertensive Nephropathy. Sci Rep-Uk 6, 31771.

1149

1150