An increase in atypical petal numbers during a shift to autogamy in a coastal sand verbena and

potential evolutionary mechanisms

Eric F. LoPresti[1,2]*; James G. Mickley[3], Caroline L. Edwards[1,4] & Marjorie G. Weber[1]

1: Dept. Plant Biology and Program in Ecology, Michigan State University

2: Dept. Plant Biology, Ecology, and Evolution, Oklahoma State University

3: Dept. of Botany and Plant Pathology, Oregon State University

4: Dept. of Biology, Indiana University

*Address for correspondence: lopresti.eric@gmail.com

Running title: Atypical flower production in Abronia umbellata

Manuscript received _______; revision accepted _______.

Abstract

Premise:

The evolution of variation in reproductive traits is of longstanding interest in biology. In plants,

meristic traits, such as petal and sepal numbers, are usually considered invariant within taxa.

However, certain species consistently exhibit great variability in these traits, though the factors

contributing to "atypical" counts are not well-known. The sand verbenas, *Abronia*

(Nyctaginaceae), usually have five perianth lobes ('petals') in their fused corollas and are self-

incompatible, thus departures from either of these norms in populations, varieties, or species are

of evolutionary interest.

Methods:

To characterize and understand an increase in atypical petal numbers during a transition from

xenogamy (outcrossing) to autogamy (selfing) in the coastal sand verbena *Abronia umbellata*,

we integrated common garden studies with analysis of over 11,000 photographed flowers from

iNaturalist, a citizen science project. Here we evaluate several adaptive and nonadaptive

explanations for the production of these 'atypical' flowers.

Key results

Our photo analysis and common garden show that the nominate xenogamous variety has 5 petals

with very little variation, however, an autogamous, geographically separated variety, *A. u.* var.

breviflora has a high preponderance of four-petalled morphs. Flower morph did not affect

successful autogamy, and petal numbers were not related to environmental factors, hybridization,

or flower size in the ways hypothesized.

Conclusions

We conclude that this loss of petals is consistent with relaxation of selection on petal number in

selfers, inbreeding leading to a loss of developmental stability, or correlated selection on another

trait. This study strongly demonstrates the power of data available from public citizen databases

for easily scored traits, such as petal number.

Keywords: Abronia; autogamy; citizen science; floral traits, hybridization, iNaturalist, meristic

variation, Nyctaginaceae, petal number,

INTRODUCTION

Understanding variation in reproductive traits has long been a focus in evolutionary biology. Taxonomic and geographic variation in reproductive phenotypes can provide key insights into fundamental evolutionary processes such as selection, speciation, and adaptation. In plants, petal number and other meristic characters are easily quantified reproductive traits. On one hand, these characters are often considered fixed within species, genera, or families and, as such, are commonly used in species descriptions, keys, and other taxonomic and identification resources. Any departure from a 'normal' meristic character within a taxon, whether species, genus, or even family, is thus of evolutionary, developmental, morphological, and taxonomic interest (Ellstrand, 1983). On the other hand, interspecific transitions in petal number or intraspecific variants in petal number are quite common in wild plants of many families, and may take the form of a change in mean or variance within or among species (e.g. Stark, 1918; Lowndes, 1931; Saunders, 1934; Roy, 1962; Huether, 1968; Schemske, 1978; Ellstrand, 1983; Lehmann, 1987; Shepard et al., 2005). Despite interest in the development, breeding, genetics, and evolution of of meristic variation, critical examination of evolutionary drivers of patters of variation, whether adaptive or non-adaptive, are scarce. Here, we integrate lab, field, and citizen-science data to examine support for several *a priori* hypotheses (Table 1) about patterns of petal number variation using a coastal sand verbena (*Abronia umbellata*, Nyctaginaceae) where allopatric varieties show remarkably different patterns of variation in perianth lobe (hereafter petal, as functionally equivalent) number.

Several hypotheses concern petal number evolution in plants, broadly falling into adaptive and non-adaptive categories. The invariance of petal number, notably across many taxa in the ancestrally five-petalled Pentapetalae (a group which includes much of Eudicot diversity:

Cantino et al., 2007; Soltis and Soltis, 2013), suggests a highly conserved developmental

program (Endress 2001), or selection maintaining constancy (Stebbins, 1974; Herrera, 2009). If

the mechanism is selective, pollinator choice or innate preference would be obvious candidates

for the maintenance of petal number. While pollinator selection has been long-hypothesized (e.g.

Leppik, 1953), positive evidence for this hypothesis remains elusive. Certain pollinators may

have the capability to 'count' petals, though many others were not found to differentiate petal

numbers (e.g., Leppik, 1953; Golding et al., 1999; Mickley, 2017). In this vein, Mickley and

Schlichting (2018) devised an alternate way of examining pollinator-mediated stabilizing

selection on petal number. They reasoned that if pollinators were maintaining a fixed petal

number, autogamous species would lack constancy due to relaxed selection on petal number.

Using several closely-related species of Polemoniaceae, they found no evidence that

xenogamous species had greater constancy in petal number than autogamous congeners.

However, using many species in a comparative phylogenetic analysis would be necessary to

robustly test this hypothesis.

An alternative, but related, hypothesis is that loss of petals in autogamous taxa may be adaptive,

but unrelated to pollinator preferences, if petals are costly (e.g., Galen, 1999; Strauss & Whittall,

2007; Lambrecht 2013) or if loss of petals increases autogamous efficiency. The autogamous

efficiency hypothesis was formulated by Monnaiux et al. (2016) as a potential mechanism by

which *Cardame hirsuta*, an autogamous mustard, has reduced petals from the four-petals typical

of Brassicaceae. Specifically, they hypothesize that petal loss slows bud opening and promotes

more effective selfing, though they do not test the proposed hypothesis (Table 1: H1). However,

a later paper found that a genotype with reduced petal numbers had a lower outcrossing rate in

the field, but did not differ in seed set in the lab (Monnaiux et al., 2018). Therefore, it is unlikely

that low petal number increased autogamous success in *C. hirsuta*, as hypothesized, but this

intriguing hypothesis certainly deserves testing in other plant species.

Non-adaptive evolutionary or phenomenological hypotheses have also been proposed to drive

petal number variation. For example, inbreeding itself, independent of pollinator-mediated

selection could reduce meristic stability (Table 1: H2). Another set of non-adaptive hypotheses

focus on hybridization. Hybridization, especially between distantly-related taxa, may cause

breakdown in the canalization of normal development due to genetic incompatibilities, e.g.

outbreeding depression, Dobzhansky-Muller incompatibilities (Bachman et al., 1981; Pelabon et

al., 2003). Hybridization led to an increased number of atypical petals in intrageneric hybrids of

both *Gilia* and *Rubus* (Grant, 1956; Bammi and Olmo, 1966). Hybridization between *Microseris*

species with 5 and 10 pappus parts respectively led to complete breakdown of canalization, with

all combinations between the parents represented (Bachmann et al., 1981). However, Choi et al.

(2001) found that, in Pseudostellaria, crosses between a species with variable petal number and

one with invariable petal number produced hybrids without petal number variation, a similar

result to certain *Solanum* hybrids (Bletsos et al., 1998). In systems where hybridization occurs, it

is possible that incompatibilities due to hybridization could drive atypical petal counts (Table 1:

H3); alternately, different developmental processes might cause hybridization to stabilize petal

number.

Regardless of the specific hypothesized mechanism, if evolution is to change petal number in

any way, it must have a heritable component. A preponderance of published evidence bears this

assumption out. Petal number was found to be heritable or genetic control was demonstrated in

Begonia semiovata, *Cardamine hirsuta*, *Dianthus cariophyllus*, *Gilia* spp*., Impatiens balsamina,*

Linanthus androsaecus, Portulaca grandiflora, Rosa wichurana, Spergularia maritima, P.

drummondii (Grant, 1956; Alpi et al., 1968; Katsuyoshi and Harding, 1969; Stevens et al., 1972;

Agren and Schemske, 1995; Delesalle and Mazer, 1995; Mazer et al., 1999; Tooke and Battey,

2000; Byerley, 2006, Pieper et al., 2015; Roman et al., 2015; Monnaiux et al., 2016, 2018;

Mickley 2017). Together with the invariance of meristic traits within most taxa, these studies

show that some level of heritability in petal number and other meristic traits is expected across

plants (and indeed, our ability to describe and identify plants based on these characters depends

on it!). If heritable, these same genes could be responsible for other traits involved in meristic

development and therefore, selection on other traits could correlate with shifts in petal number

(e.g. Table 1:H4).

Finally, while petal number is demonstrably heritable in most systems, several studies have

additionally found evidence of plastic variation in petal number (Table 1: H5). While the genetic

control of atypical petal numbers in *Cardamine hirsuta* is well-known, there is a seasonal

component to the expression, with atypical petal numbers being more common in summer than

spring (McKim et al., 2017). Huether (1968, 1969) found that herbivory, day length, and

temperature increased petal number variation, though only in some plants, suggesting GxE

interactions similar to that of *C. hirsuta*. Hoffman et al. (2009) found that an elevational gradient

in atypical floral number in *Stylidium armeria* disappeared in a greenhouse common garden,

suggesting an environmentally plastic component. In populations of *Spergularia maritima*,

greenhouse grown plants consistently had fewer petals than plants in the populations from which

seeds were collected, however, petal number had the highest broad-sense heritability of all

reproductive traits they measured (Delesalle and Mazer, 1995). Roy (1962), with the help of a

dedicated field assistant, one JBS Haldane (!), found that while mean petal number of individual

Nyctanthes arbor-tristis plants did not change markedly through the season ($< 5\%$), the variance

increased substantially (20-40%) from beginning to end of season. Developmental changes could

also explain variation in certain cases. Older clones of a buttercup (*Ranunculus repens*) had

higher proportions of atypical petal numbers, a result which Warren (2009) ascribes to the

accumulation of somatic mutations over time, but this result could also be driven by other age-

correlated factors, such as hormonal changes. The production of repeated units in an organism,

i.e. many flowers on a plant, inevitably leads to some variation in these parts (Herrera, 2009). A

model of developmental stochasticity over repeated floral development led to realistic variation

in petal numbers (Kitazawa and Fujimoto, 2014). This suggests that no matter the characteristics

of the plant; one with many flowers will have a higher likelihood of a greater range of atypical

petal numbers than without (sampling effects of this sort also discussed in Roy 1962).

Despite the various hypotheses proposed for petal number variation, no investigations have

addressed multiple hypotheses in a single system or characterized variation across species' entire

range in the field. Petal number can be scored on live plants, herbarium specimens, or

photographs. Rapidly growing georeferenced photographic databases from citizen science

initiatives (e.g. iNaturalist, CalFlora, etc.), hold massive amounts of easily quantified data,

extraordinarily well-suited for studies of petal number variation. Photographic citizen science

data have been used recently in other studies for easily scored traits including plant pathogen

infection rates (Kiddo and Hood, 2020) and lizard ectoparasite presence (Putnam et al., 2020).

Therefore, these databases represent an exciting complement to herbarium and other natural

history collections for scoring of characterization of traits.

However, scoring phenotypes from field photos alone cannot disentangle evolutionary and

plastic drivers of trait variation. As such, integrating large, range-wide field photography datasets

with carefully controlled common garden experiments is a powerful approach for evaluating

wide-scale patterns of natural field variation in light of genetic variation in the lab. Here, we

employ this integrative approach to address several proposed hypotheses for atypical petal

numbers (Table 1) using the naturally variable species *Abronia umbellata*. In particular, we

capitalize on natural variation across two a natural transition, a southern, outcrossing variety

(=subspecies in Nyctaginaceae taxonomic conventions) with typical 5-lobed flowers, and a

northern selfing variety with 4-lobe flowers (any deviation from five is considered 'atypical' for

our purposes). We integrated a greenhouse common garden of *A. umbellata* subspecies and

hybrids, and photographic analysis of over 11,000 citizen science (iNaturalist) photos spanning

the entire *A. umbellata* range to characterize the petal number variation and evaluate several

adaptive and non-adaptive hypotheses (Table 1). Together, this study represents a comprehensive

integrative analysis of petal number variation in a single system, and characterizes an

evolutionarily compelling transition in reproductive morphology in association with the

evolution of selfing.

MATERIALS AND METHODS

***The geographic range of* Abronia umbellata *and its varieties —* A. *umbellata* grows on

beaches and coastal dunes – never more than a few hundred meters inland - from British

Columbia, Canada, south to Baja California, Mexico. It has two geographically separated,

genetically-distinct varieties (=subspecies) across its range on the Pacific Coast of North

America (Baja California, Mexico – British Columbia, Canada) (Greer, 2016; Van Natto, 2020).

The southern variety (*A. u. umbellata*) is self-incompatible, and therefore xenogamous, whereas

the northern variety (*A. u. breviflora*) is self-compatible and probably nearly completely

autogamous in nature (Doubleday et al., 2013; Doubleday and Eckert, 2018; LoPresti, pers.

obs.). Initial observations (E.F.L.) suggested that the southern xenogamous (outcrossing) *A. u.*

umbellata has very little variation in perianth lobe (hereafter, petal) number, almost all flowers

having the typical five petals. In contrast, the northern autogamous *A. u. breviflora* has strikingly

more variable petal number (Fig. 1: A-D). The autogamy, and petal number variation, of this

variety is likely derived. Almost all *Abronia* species are self-incompatible (Tillett, 1967;

Galloway, 1975; Williamson et al., 1994; Darling et al., 2008; LoPresti, unpublished data), and

thus, the shift in production of atypical flowers in *A. u. breviflora* occured with a transition to

autogamy (Doubleday et al., 2012; Greer, 2016; Van Natto, 2020).

The opening of San Francisco Bay (~37.8° N) separates the northern *A. u. breviflora* from *A. u.*

umbellata (Doubleday et al., 2013; Greer 2016; Van Natto 2020). Various authors have posited

that specimens, especially from southern Sonoma county (~38.3° N), do not fit this break point

(e.g. Tillett 1967). However, all plants from north of the bay, even if not fitting the Galloway

(2003) or Murdock (2012) measurements for *A. u. breviflora*, appear to be selfing, both from

common garden, and field collections (a high proportion of filled fruit is indicative of autogamy

in coastal *Abronia*), a result also found by Darling et al. (2008) and Van Natto (2020).

Population genetic data also shows a cluster from the north and a cluster from the south (Greer,

2016), though the break was not entirely possible to ascertain completely as it did not include

any samples close to the boundary.

Darling et al. (2008), in an examination of range limits and dispersal of the species, treated the

northern range limit as southern Oregon, USA (~43° N), as populations north of that point are

small, rare, and protected from collecting. They considered the southern limit in Baja California,

Mexico as ~30° N (these range limits also used in Samis, 2007; Greer, 2016; Van Natto 2020).

We consider the range limit differently in this study, using 49° N to 32.5° S, with the caveat that

our southern limit is conservative. Our northern range limit is accurate, as both historical

herbarium collections and a few extant populations range northward to British Columbia, Canada

(49° N); sometimes these northernmost populations are treated as a separate variety, *A. u.* var.

aculata, though we do not treat this as a separate variety from *A. u.* var. *breviflora* as they are

morphologically indistinguishable in our experience (also supported by the analyses of Van

Natto, 2020; though taxonomic priority belongs to *A. u.* var. *aculata*).

The southern range limit is truly unknown because of a gradation in morphology into *A. gracilis*

and, probably, *A. villosa*, almost certainly indicative of an extensive hybrid zone (see Ann

Johnson's extensive collections and determinations of hybrids in Baja California at the UC-Davis

herbarium). This accords well with previous knowledge of the genus; when growing in

sympatry, many sand verbenas hybridize naturally, and hybrids are easy to make and usually

successful in artificial crosses (Tillett, 1967; Galloway, 1974; Johnson, 1978; Van Natto, 2020;

Edwards & LoPresti, unpublished data). While we have seen herbarium specimens from as far

south as 24° N have been annotated as *A. umbellata*, these are unlikely to be accurate or of non-

hybrid origin. Tillett (1967), in the most comprehensive examination of coastal *Abronia* species,

found *A. gracilis* at San Simon (30.5° N) and apparent hybrid *A. gracilis* x *A. umbellata* at San

Quintin, Baja, MX (30.6° N; both records north of the farthest south populations in Darling et al

2008). However, from herbarium specimen examination, he notes that *A. gracilis* sometimes

occurs as far north as San Diego county (i.e. north of 32.5° N), a result which we doubt. The

farthest north record we find convincing is a record from iNaturalist at 31° north (record

#16085273) which is morphologically indistinguishable from *A. gracilis*, though could certainly

be of hybrid origin. In the only population genetics examination, Greer (2016) found that Baja

populations "clustered apart from [*A. u. umbellata*] in adjacent Southern California, with very

high levels of polymorphism" and Van Natto (2020) found much genetic variation in these

populations not shared by farther north populations. Neither of these works mentioned *A. gracilis*

or the possibility of hybridization but is seems possible introgression or a hybrid zone a likely

source of the unique SNPs found in these populations. Further complicating identifications, the

only noted character differentiating *A. umbellata* and *A. gracilis* is leaf shape (Tillett 1967,

Galloway 2004) which is immensely variable within and among individuals, even within the core

range of *A. umbellata* (and leads some individuals of *A. umbellata* and *A. villosa* to key to *A.*

gracilis in various keys, pers. obs.). For these reasons, we conservatively used the US-MX

border, 32.5° N, as the southern edge of this study. Further study is necessary to determine the

actual southern range limit of *A. umbellata* (Fig. 2) and any hybrid zone in that area.

***Broad survey of petal number across the range of* Abronia umbellata (*~H2)* —**To quantify

patterns of petal number variation in the field, we manually extracted data from iNaturalist

photographic records across the entire range of the species. All records identified as "research

grade" *Abronia umbellata* as of 13-Jan-2020 were assessed (n=1096). Of these, 865 records were

able to be scored, comprising 1,808 inflorescences, and 11,428 individual flowers. For each

inflorescence in the photograph able to be scored, we recorded petal number of each flower, as

well as record number, latitude, and the date of the record. For those records where the location

was obscured by iNaturalist, the latitude listed was taken, as the obscuring occurs within a 10 km

radius, a relatively small distance compared to the range-wide scale of this study and therefore,

this uncertainty should not affect our results. Because interdigitation between prostrate stems of

closely-packed plants is common, we were unable to conclusively assign inflorescences to

individual plants therefore, multiple inflorescences in the same photograph were scored

separately, but with the iNaturalist record number the same; the record number was used as a

random effect in analyses when necessary (as a rough proxy for individual).

Two-hundred and two records were not used for the following reasons: the plants were not in

flower, the flowers were out of focus, flowers were turned away from the camera, the photograph

was too distant to count petals on individual flowers, the inflorescence had been consumed by

herbivores, only one flower was visible, the photograph was duplicated across records, or the

record occurred south of this study's border. Twenty-nine 'research-grade' records were also

discarded because of misidentifications: one *Abronia maritima* was misidentified as *A. umbellata*

and 28 records were obvious hybrids between *A. umbellata* and co-occurring congeners. EFL

added the correct identifications as part of the review process and these records will likely

eventually get a correct determination in the iNaturalist database (pending other users' 'votes').

To determine whether variety predicted atypical petal counts (i.e. not 5-lobed), we used a

binomial mixed model (R 3.6.2: package *lme4*) of with typical, atypical petal number as a

response variable and variety as a fixed effect and record number as a random factor. This model

was compared to a null model without variety using a likelihood ratio test.

Common garden petal counts (~H2)— To quantify variation in petal counts and minimize the

effect of plasticity, we scored plants in the shared environment of an *Abronia* common garden at

Michigan State University. As part of a genus-wide growout, we grew ten var. *breviflora*

individuals and six var. *umbellata* individuals from seed collected by EFL and one USDA

accession of var. *breviflora* (Supplementary Table 1). These plants were maintained in the

greenhouse and laboratory under artificial lights with a 12-hour light:dark cycle (though

greenhouse plants got additional natural light in summer). Plants were maintained in 8-18" pots,

grown in a mixture of sand and potting soil (~1:1), and kept ~70° F with 12-hour artificial

lighting, though individuals in the greenhouse experience longer days naturally during spring and

summer seasons. Plants were watered every 3-5 days and fertilized (Miracle-Gro, 10:10:10) once

a month. Plants were rotated between the greenhouse and laboratory often, as we took all

measurements in the lab (for this and other studies). Petal number of flowers in all flowers in

mature inflorescences were counted between July 2019 and May 2020 when each plant was

flowering as part of the broader grow-out. We compared summed counts of petal numbers of

each variety in the common garden, however statistics were not performed because of low

number of individuals involved (in contrast to the iNaturalist data set).

Hybridization effects (H3) — Hybridization may alter meristic characters (e.g. Vlot et al., 1992)

and therefore, may result in the production of atypical petal morphs in regions where

hybridization and introgression occurs (H3). Hybridization occurs commonly in *Abronia*,

especially in the xenogamous *A. u.* var. *umbellata,* which hybridizes naturally with *A. latifolia,*

A. maritima, A. villosa, A. gracilis, and probably *A. pogonantha* (Tillett, 1967; Johnson, 1977;

Van Natto, 2020; LoPresti, pers. obs.). While Tillett (1967) hypothesized that *A. u.* var.

breviflora may be the result of a hybridization event between *A. umbellata* var. *umbellata* and *A.*

latifolia, using genetic methods Van Natto (2020) found this hypothesis extremely unlikely. We

experimentally crossed *Abronia* species across the phylogeny in genus-wide hybridization study;

for this study we scored petal numbers of F1 hybrids, grown under the same conditions as

parents, which had a parent either of *A. u. umbellata* or *A. u. breviflora* (10 combinations, n = 44

total hybrid individuals; parental collection locations in Supplementary Table 1). These plants

did not necessarily reflect naturally-occurring hybrids, instead these hybrid combinations were

designed to score traits across as broad a range of relatedness as we could in the genus. However,

we see no reason to assume that these particular combinations would have meristic

incompatibilities not representative of other combinations within the genus, and therefore we do

not expect the exact combinations to affect the overall results in a particular direction. We asked

whether each hybrid combinations had a consistent pattern of production of atypical flowers

compared to its' parental *umbellata* variety using a chi-squared test.

Laboratory test of autogamy success with different petal numbers (H1) — In order to the

adaptive hypothesis that decreased petal number increases autogamous success, we compared the

success of autogamy in flowers differing in petal number. Between 7-November-2019 and 09-

March-2020 we marked all individual flowers in 58 inflorescences of the autogamous variety *A.*

u. var. *breviflora* in common garden study and scored the success of autogamy when fruit had

developed (Fig. 1: E, F). In total, these 58 inflorescences comprised 597 flowers (417 5-petalled,

169 4-petalled, 6 3-petalled). As all *Abronia* develop one seed per fruit/flower, the response

variable was binomial (seed/no seed). We used a binomial mixed model with petal number as a

factor and plant ID and inflorescence ID as random variables.

Environmental effects (H5) — To determine whether hypothesized seasonal and latitudinal

factors affected production of atypical petal number morphs (H5), we tested for correlation

between these environmental variables and each of atypical petal production and inflorescence

mean petal number using the iNaturalist data set. Latitidue and season have been suggested to

affect petal number in other plants (Mathiason, 1982; McKim et al., 2017). As the varieties

showed such different patterns of petal number and do not overlap in latitude, variation is

confounded with latitude (and flowering time correlates with latitude), we analyzed the varieties

separately for each analysis. For seasonal effects, we used mixed models with Julian date (from

Jan 1 of each year) as a continuous predictor and iNaturalist record as a random variable to

account for non-independence of inflorescences in the same photo being likely from the same

plant. We tested linear and second-order polynomial (in case of changes at start and end of

flowering period; i.e. as found for Cardamine in McKim et al., 2017) for each variety.

Correlation of floral size and petal numbers (H4) — To test whether flower petal number

correlates with flower size, we combined our iNaturalist data with previously published data. As

measuring floral size is not possible from photos, and we had only one population of var.

umbellata, and three of var. *breviflora* in the common garden, data on corolla diameter and

number flowers in the inflorescence from 12 populations was extracted from Doubleday et al.

(2013) using the data extraction program: (https://apps.automeris.io/wpd/). The petal number data

for these populations was taken from the iNaturalist photo analysis, using +/- 0.0025 degrees

from their coordinates as the range for each population, since the nearest populations in their

analysis were 0.0050 apart. This minimum grain ensures that no records are individually counted

in multiple populations. By this population grain, we had petal number records for 7 of

Doubleday et al.'s populations, all of var. *umbellata*, with a range of 2 to 56 iNaturalist records

falling within each population. We analyzed data using mixed models with petal number as a

response, either floral face diameter or flowers per umbel fit as fixed effects and record as a

random effect. There was not sufficient overlap between the populations in our sample and

Doubleday et al.'s to perform a similar analysis for variety *breviflora*.

RESULTS

Because our hypotheses incorporated evidence from common garden and the iNaturalist data set,

we have organized the results by hypothesis and were specific about from where the data came.

Characterization of petal numbers of autogamous and xenogamous varieties — Petal number

counts from both common garden and field iNaturalist plants showed that the autogamous *A. u.*

breviflora had far fewer 5-petalled (and more 4-petalled) flowers than the xenogamous *A. u.*

umbellata. The varieties differed significantly in their production of atypical petals in the range-

wide iNaturalist data set; the random-effects model including variety as a fixed effect fit

significantly better than a null model without variety (X^2=93.6, df=1, p<0.0001). A mean

percentage per record of 80.2% of the autogamous *A. u. breviflora* flowers were five-petalled

and 93.1% of the xenogamous *A. u. umbellata* were five-petalled (Fig. 2,3,4). In the common

garden, *A. u. breviflora* had a mean percentage per individual of 74% 5 petalled, and 25% four-

petalled flowers (Table 2), whereas *A. u. umbellata* had 93% 5-petalled and 7% four-petalled

flowers. While lab-grown plants scored for petal number were of varying age, the fact that field

results showed the same qualitative pattern as that in lab-grown plants suggest that plant age is

not a large factor in petal number variation between the two varieties.

Hypothesis 1: autogamous success of differing petal numbers — Petal number had no effect on

autogamous success (4: 76.3% & 5: 78.4% set a seed; n=58 inflorescences, 586 flowers: six

flowers with three petals and five which did not open were excluded from analyses). A binomial

mixed model with petal number as a predictor fit no better than a null with just random effects -

inflorescence ID nested in plant ID (Likelihood ratio test, df=1, X^2=0.41).

Hypothesis 3: hybridization effects on petal numbers — Hybrids did not consistently produce

more atypical flowers compared to parents in fact, almost all hybrids that differed significantly

from their parental variety had lower than expected atypical morphs (Table 2). Contrary to the

prediction of hybridization reducing meristic stability (i.e. increasing atypical morphs), it is

noteworthy that most crosses involving variety *umbellata* with already low levels of atypical

petal morphs, had even lower rates of atypical petals than expected (significantly different in 4 of

the 8 crosses: Table 2). However, these results are not drawn from large numbers of samples

(n=44 plants). Crosses of the two umbellata varieties showed atypical petal numbers which did

not differ from variety *breviflora*, with 27% of flowers being four-petalled, but differed

significantly from variety *umbellata*.

Hypothesis 4: correlation of flower size and petal number — Among the populations (n=11) of

A. u. var. *umbellata* with flowers measured by Doubleday et al. (2013) for which we also had

petal number data in the iNaturalist data set (n = 7), there was no correlation between corolla

width and production of atypical petals. Models with either floral face diameter or flowers per

umbel fit no better than a null with only a random effect of record (X^2=1.48, df=1, p=.22;

X^2=1.07, df=1, p=0.30).

Hypothesis 5: environmental drivers of petal number — We found no significant effect of

phenology on atypical petal number of either variety from the iNaturalist data set. Date (days

after 1-Jan each year) had no correlation with atypical petal production either linearly or with a

second-order polynomial (in case of changes at start and end of flowering period; i.e. as found

for *Cardamine* in McKim et al., 2017) for either variety. Mixed models including only the

random effect of the record ID fit better than those containing the day for both varieties (var.

breviflora: X2=1.46, df=1, p=0.23; var. *umbellata*: X2=0.07, df=1, p=0.79).

Latitude predicted atypical petal number better than a null model in each variety (var. *breviflora*:

X^2=23.2, df=1, p < 0.001; var. *umbellata*: X^2, df=1, p < 0.001); however, the amount of variance

explained by latitude in these best-fitting models was small (marginal pseudo-R^2, package

sjstats: *breviflora* = 0.055; *umbellata* = 0.016). In both models, the random effects of record (a

proxy for individual) explained far more variance (conditional pseudo-R^2, *breviflora* = 0.195,

umbellata = 0.260). Using mean petal number instead to characterize the trend, we found mean

petal number decreased with increasing latitude. In *A. u.* var. *breviflora*, the relationship between

mean petal number and latitude was strong; for every increase of 1° latitude, mean petal number

decreased by 0.042 ($\pm$ 0.008); from the southernmost populations (~38 N) to the northernmost

(~49 N), this equates to a drop in mean petal number of almost 0.5! In contrast, while the

correlation between latitude and mean petal number was significant in *A. u.* var. *umbellata* as

well and this model fit better than a null; the effect was extremely small – a predicted drop of

0.01 petals per every increase of 1° latitude (i.e. 0.05 drop across the entire range) and explained

only ~1% of the variance observed (r^2=0.0111). Regardless, this result demonstrates that the

direction, but not magnitude, of correlation is consistent between the varieties.

DISCUSSION

Using a combination of a large citizen-science dataset, common garden grow-outs, and

laboratory studies of autogamous success and of hybridization, we characterized petal number

variation across the entire range of a widespread species and critically examined several

published hypotheses about the evolutionary drivers of petal number variation in plants. Broadly,

we found that the autogamous *Abronia umbellata* var. *breviflora* had far higher proportions of

atypical flowers than the xenogamous *A. u. umbellata*. Overall, we documented a pattern

consisten with relaxation of selection in petal numbers in selfers, correlated selection on another

trait (but probably not flower size), or inbreeding leading to a loss of developmental stability

(H2). We did not find support for the hypotheses of increase in autogamous efficiency of reduced

petal numbers (H1), hybridization reducing meristic stability and increasing atypical morphs

(H3), correlated selection on flower size (H4), and evidence for environmental drivers of

plasticity was mixed, and the effect was too small to account for the overall pattern (H5). While

we were unable to conclusively identify any mechanism as contributing to the pattern; our

integration of citizen science and common garden studies was able to eliminate several

hypotheses and we believe the approach we used can be applied to other systems to better

understand this interesting, yet understudied, phenomenon.

Interpretation of the pattern — We found a robust pattern of increased atypical petal production

in the self-compatible *Abronia umbellata* var. *breviflora* compared to the outcrossing *A. u.* var.

umbellata in both field and lab. This pattern is consistent with several, but not all, of the

hypotheses proposed in the literature to explain variation in petal number (Table 1). In particular,

the increase in atypical flowers in the selfing var. *breviflora* is at least partially consistent with at

least three, non-exclusive, mechanistic hypotheses: relaxation of selection on petal number in

selfers (implying some adaptive function of five-petals in outcrossers, for example via pollinator-

mediated selection), inbreeding leading to a loss of meristic stability (H2), or correlated selection

on an unmeasured trait (H4). We were, however, able to provisionally eliminate several proposed

mechanisms (autogamous efficiency: H1, hybridization: H3, seasonality/phenology: H5) in this

system. Below we discuss the rationale for each of the three remaining hypotheses, including

what additional evidence might be garnered to support each of them.

It is possible that inbreeding may be responsible for the observed high rates of atypical petal

production in selfers. In contrast to the self-incompatible *A. u. umbellata*, *A. u.* var. *breviflora*

reproduces almost solely autogamously (Doubleday et al., 2013; Van Natto, 2020), and

therefore, almost all individuals in our study – in nature and the lab – were probably inbred for

several generations. Greer (2016) and Van Natto (2020) found that there was little genetic

variation in *A. u.* var. *breviflora*. Inbreeding may increase aberrant morphs in meristic traits

(Sherry and Lord, 1996), and potentially carry little cost in this autogamous population.

Therefore, it seems possible that inbreeding may be responsible for the observed high rates of

atypical petal production. While this is a viable hypothesis in our system, the only other

evaluation of this hypothesis that we know of, which used data from Schlichting and Levin

(1986) on xenogamous *Phlox drummondii*, found that inbreeding decreased the proportion of

atypically petalled flowers (Mickley and Schlichting, unpublished data). Allowing lines of var.

breviflora to self for several generations as well as manually outcrossing those lines could be

used to rigorously evaluate this hypothesis in future studies.

A relaxation of pollinator-mediated selection maintaining five-petalled flower (Leppik, 1953,

1956), or petal number constancy (Herrera, 2009) during the evolution of autogamy in var.

breviflora also remains a possibility as an explanatory mechanism of the high rates of atypical

petal numbers in var. *breviflora*. Correlated selection on numbers of other meristic traits by

pollinators also is possible, though the lack of a reduction in stamen number at the same time

makes this less likely (virtually all four-petalled flowers in *Abronia* have five stamens: LoPresti,

unpublished data) Any pollinator-mediated selection on petal number would require pollinators

to be able to count petals, which bees have been found to be capable of (Leppik, 1953, 1956;

Lehrer et al., 1995) but Lepidoptera have not (Leppik, 1954, 1955). *A. umbellata* is likely

primarily moth pollinated; Doubleday and Eckert (2018) found that *A. umbellata* had higher

floral visitation during the day, but all but a very small amount of pollination occurred at night.

Given these lines of evidence, it seems unlikely that the lower production of atypical morphs in

var. *umbellata* is due to pollinator-mediated selection, though it is consistent with our results and

we cannot rule it out. Further investigations into pollination success of atypical flowers in the

field would be necessary to rigorously examine this hypothesis.

The two varieties of *A. umbellata* differ greatly in floral size, and, given that close relatives are

also outcrossing and large flowered, the small flowers and autogamy in var. *breviflora* are

derived states (Doubleday et al., 2013). Large-flowered, close relatives, including *Abronia*

villosa var. *villosa*, *Abronia villosa* var. *aurita,* and *Abronia gracilis,* have extremely low rates of

atypical petal morphs (LoPresti, unpublished data), therefore, the evolution of atypical petals

occurred alongside the evolution of autogamy and small flowers in this lineage. We did not find

evidence that within var. *umbellata*, petal morphs correlated with corolla width; however, the

macro-evolutionary and population-level processes may be decoupled, and rigorous

interpretation of those data, from seven populations in a narrow part of the range of this variety

(34.22 –36.95 ° N) is limited. Because the atypical petal number is confounded with floral size

differences between the morphs, analyzing across the range is inappropriate. Therefore, whether

the lack of correlation within the narrow range of flower sizes within var. *umbellata* is

informative in explaining the marked transition to smaller flowers and fewer petals during the

evolution of var. *breviflora* remains an open question and cannot be tested with a single

transition. Instead, either correlations across multiple transitions to selfing across more species or

an artificial selection experiment could be used more rigorously examine this hypothesis.

No effect of petal number on autogamy — Petal expansion assists in opening of the floral bud,

leading Monnaiux et al. (2016) to hypothesize that a reduction in petal number might delay bud

opening and increase successful autogamous fertilization. In our laboratory study, we found no

support for increased autogamous success of four-petalled morphs compared to five-petalled

morphs in the autogamous *A. u.* var. *breviflora*. There is a possibility that in the field, four-

petalled morphs may get fewer, shorter, or less effective pollinator visits and thus, may have

lower pollen removal and therefore higher autogamous success; our laboratory study could not

examine this hypothesis. Additionally, we attempted and failed to find any distinguishing

characteristics in fruit resulting from flowers of differing petal numbers, meaning a *post hoc*

analysis could not be done from field seed collections, as we had initially hoped. Any test for

autogamy in a field setting would therefore require marking individual flowers, as we did in the

laboratory. Nevertheless, this study was the first to test this broad hypothesis and we found no

support. Our study design is simple and could be easily replicated in other taxa with variable

petal number (species in the Polemoniaceae, Ranunculaceae, etc.) to help draw larger

conclusions than from this test alone.

Environmental or phenological effects — In other systems, changes in petal numbers occur

plastically in response to environmental variables (e.g. Huether, 1968; McKim et al., 2017), the

two which we could test with the large iNaturalist data set were latitude and phenology (both as

proxies for climatic variables or day length). In *A. u.* var. *breviflora*, latitude was correlated with

atypical petal number; as latitude increased mean petal number decreased. Whether this effect

was due to environmental factors associated with latitude or some genetic variation, due to any

number of factors, including drift or founder effects of those small, very isolated northern

populations, is impossible to tell with certainty. Our common garden data is consistent with the

observed gradient in the field, though it is extremely limited, suggesting that this gradient may

not be entirely due to plasticity. Individuals from a population from Oregon (n=3) had a higher

proportion of 4- (36.5%) and 3-petalled (5.3%) flowers compared to a population from Sonoma

County, CA (n=6) with 23.5% and 0.8%, respectively. Further common gardens drawn from

populations across the entire range could shed light on to what extent this latitudinal gradient is

heritable and/or environmental.

Latitude was also a 'significant' negative predictor of atypical and mean petal number in *A. u.*

var. *umbellata*. However, any biological meaning is difficult to interpret, given that it explained

~1% of the variation with minimal effect on petal number across the much larger range of this

variety; this was a pattern that would likely not have been picked up in any smaller data set than

the >11,000 flowers we were able to score from iNaturalist. We therefore hesitate to ascribe any

biologically relevancy to this small an effect, though the consistent direction between the

varieties does suggest a common mechanism.

In contrast to McKim et al. (2017)'s characterization of phenology causing changes in petal

number in *Cardamine hirsuta*, we did not find any effect of phenology on either variety. We

analyzed this with linear and polynomial models; none fit better than a null model. Coupled with

the consistent pattern between field and common garden plants, we believe that this lack of a

seasonal correlation is further evidence that the observed production of petal morphs is largely

heritable and the latitudinal effect should be investigated to parse out population differentiation

across latitudes or a plastic response to environmental differences across latitudes.

No consistent effect of hybridization on petal number variation — Hybrids of *A. umbellata* did

not consistently exhibit increases in atypical petal number (only one of 9 crosses did). While we

cannot completely exclude hybridization as a mechanism behind the variable petal numbers in *A.*

u. var. *breviflora*, especially given the lack of crosses involving other coastal species, it seems

extremely unlikely that hybridization leads to consistent meristic instability. Most tellingly in

our system, Van Natto (2020) found no measureable levels of introgression in var. *breviflora*,

which they deduced was likely due to its autogamous reproduction system. Therefore, since the

non-introgressed variety has the higher atypical petal morphs and hybridization does not increase

atypical petal numbers in most crosses, hybridization is an unlikely mechanism behind the

increased atypical petal numbers in *A. u.* var. *breviflora*.

Conversely, could hybridization reduce petal number variation in *A. u.* var. *umbellata*?

Consistent with this hypothesis, most of the hybrids created in this study had reduced atypical

petal numbers; somewhat contrary to this broader hypothesis is the stability of meristic traits

across all *Abronia* species, a result very unlikely to be driven by hybridization (LoPresti, pers.

obs./unpublished data). Demonstrating that the low levels of introgression into this variety

(Tillett, 1967; Van Natto, 2020) stabilizes petal number would require far more investigation

than this survey study could provide, and would need to be investigated in far more detail.

Whether the identity of the hybrid species pairs also may be informative in predicting stability or

lack thereof, Pelabon et al., (2003) found that in crosses of intermediate distance (within a

species) the resulting hybrids had reduced variation in floral traits. We did not find that result;

our crosses of var. *umbellata* did not show a pattern with relatedness (unpublished data),

however, we caution interpretation of this result both as Pelabon's comparisons were between

populations of the same species, and the small sample size of our crossing data. The ongoing

larger hybridization study from which these data were extracted holds the potential to test the

shape of this relationship across a greater range of relatedness of many species pairs across the

Abronia phylogeny.

More broadly, hybridization deserves increased study in *Abronia umbellata* var. *umbellata*; Van

Natto's (2020) findings that "early-generation hybridization was occurring anywhere [var.

umbellata] was sympatric with another species" suggests the importance of considering

hybridization in evolution and ecology of this species. While Van Natto examined *A. umbellata*

hybrids with only *A. maritima* and *A. latifolia*, Tillett (1967) also suggested hybridization with *A.*

villosa. We consider this additional hybrid extremely likely in several locations where *A. villosa*

var. *aurita* occurs near the coast (i.e. Camp Pendleton); material from these sites is not easily

ascribable morphologically to one or the other species. We also add another intriguing

possibility: specimens collected by William Palmer in 1876 from an inland site in eastern San

Luis Obispo county (the 'McGinnis Ranch') were described as *Abronia minor* by Standley

(1909). Tillett considered *A. minor* a hybrid of *A. umbellata* x *A. latifolia*, though neither species

occurs inland and morphologically, these specimens have no characteristics indicative of *latifolia*

parentage. Most tellingly, the McGinnis ranch falls into the current and historic range of *A.*

pogonantha. Most species of *Abronia* are pollinated commonly by large, migratory hawkmoths,

(Keeler, 1978; Douglas, 2008; Doubleday and Eckert, 2018; LoPresti, pers. obs) and pollen

movement from over this <50km distance from *A. umbellata* to *A. pogonantha*, but not fruit

movement, seems most likely. For these reasons, from our examination of Standley's

illustrations and scans of the series collected, we find it far more likely that this material

represents backcrosses of *A. umbellata* (♂) x *A. pogonantha* (♀) with *A. pogonantha* (F1 hybrids

are unlikely since Palmer collected several individual plants).

The invaluable contribution of iNaturalist citizen science data — Initial observations of

atypical petal production were confined to the few plants of each variety we had in the Michigan

State University greenhouse during a common garden grow-out of many *Abronia* species.

Herbarium sheets often have very few easily scoreable flowers on them, and therefore, the scale

of this study and conclusions reached would have been greatly reduced without the immense

iNaturalist repository of identified photographs. Field sampling across the geographic range

during all twelve months of the year, would have been logistically and financially impossible;

yet, the analysis of these flowers from iNaturalist data took EFL less than six weeks! This study

adds to a burgeoning body of literature using iNaturalist data to extract scoreable traits on varied

organisms (e.g. Kido and Hood, 2020; Putman et al., 2020). Petal number could easily be scored

across various clades with transitions to selfing and would be informative for more evolutionary

correlations than the one transition studied here.

Conclusions — Integrating a common garden study, crossing experiment, and large-scale field

citizen science data, we characterized an intriguing shift in atypical petal number during a

transition from outcrossing to autogamy in a coastal sand verbena. The increase of atypical petal

number of the selfing variety was not consistent with directional selection caused by increased

autogamous success, nor was it driven by seasonal environmental changes, as in other similar

systems. While we did not find total support for any hypothesis, we believe the totality of the

data presented suggests that some combination of inbreeding leading to loss of meristic stability,

correlated selection on another trait, or relaxation of stabilizing selection on petal number in this

selfing variety are potential mechanisms. We also demonstrate the power and ease of using high-

quality photographs from citizen-science efforts to score easily quantifiable traits across range

and seasons to generate a data set which would have been logistically difficult to get with

fieldwork or herbarium specimens.

ACKNOWLEDGEMENTS

The authors thank the many iNaturalist contributors for detailed photographs and identifications;

the 'data' here was gathered by hundreds of people with cameras! We thank past and present

members of the Weber lab, especially Carina Baskett, and Michael Foisy, for comments on the

manuscript and on the project overall. Laura Doubleday gave us invaluable information allowing

us to meaningfully incorporate her data on flower size. Rick Karban, Patrick Grof-Tisza, Dena

Grossenbacher, Kimiora Ward, and Kathy Toll accompanied and assisted EL on fieldwork and

seed collecting of *Abronia umbellata* during 2015-2018 seasons and Carolyn Graham and Bruce

Martin assisted in caring for the common garden growouts. EFL was funded by NSF-PRFB

#1708942, MGW was funded by NSF-DEB #1831164.

DATA AVAILABILITY

Data sheets will be archived as supplements, see online supporting information section.

Hybridization data is in Table 2; more data (including more crosses of *A. umbellata*) will be

available when the whole-genus study from which those data were taken is completed.

ONLINE SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information section at

the end of the article, Four supplementary data sheets and one file of code are included: (1) the

iNaturalist data, (2) autogamy data, and (3) parsed iNaturalist data to the populations in

Doubleday et al (2013) for the flower size comparison, (4) collection locations and herbarium

specimen numbers for plants used in the common garden studies, and (5) an R script uses the

attached data sheets for the analyses in the paper.

LITERATURE CITED

Agren, J., and D.W. Schemske. 1995. Sex Allocation in the Monoecious Herb Begonia
semiovata. Evolution 49: 121–130.

Alpi, A., M. Buiatti, and S. Baroncelli. 1968. Some data on the polygenic control of two
quantitative traits in a vegetatively propagated flower plant, the carnation. Theoretical
and Applied Genetics 38: 298–300.

Bachmann, K., K. Chambers, and H. Price. 1981. Genetic determination of pappus part number
in the annual hybrid Microseris B 87 (Asteraceae - Lactuceae). Plant Systematics and
Evolution 138: 235–246.

Bammi, R.K., and H.P. Olmo. 1966. Cytogenetics of Rubus. V. Natural Hybridization Between
R. procerus P. J. Muell. and R. laciniatus Willd. Evolution 20: 617–633.

Bletsos, F.A., D.G. Roupakias, M.L. Tsaktsira, A.B. Scaltsoyjannes, and C.C.

Thanassoulopoulos. 1998. Interspecific hybrids between three eggplant (Solanum

melongena L .) cultivars and two wild species (Solanum torvum Sw . and Solanum

sisymbriifolium Lam .). Plant Breeding 164: 159–164.

Byerley, M. 2006. Patterns and consequences of floral formula variation in Phlox

(Polemoniaceae). Colorado State University.

Cantino, P.D., J.A. Doyle, S.W. Graham, W.S. Judd, R.G. Olmstead, D.E. Soltis, P.S. Soltis, and

M.J. Donoghue. 2007. Towards a phylogenetic nomenclature of Tracheophyta. Taxon.

Choi, K., J.S. Kim, and J.H. Pak. 2001. Natural hybridization between Pseudostellaria davidii

and Pseudostellaria palibiniana (Caryophyllaceae). Plant Species Biology.

Delesalle, V.A., and S.J. Mazer. 2014. The structure of phenotypic variation in gender and floral

traits within and among populations of Spergularia marina (Caryophyllaceae). American

Journal of Botany 82: 798–810.

Doubleday, L.A.D., and C.G. Eckert. 2018. Experimental evidence for predominant nocturnal

pollination despite more frequent diurnal visitation in Abronia umbellata

(Nyctaginaceae). Journal of Pollination Ecology 22: 67–74.

Doubleday, L.A.D., R.A. Raguso, and C.G. Eckert. 2013. Dramatic vestigialization of floral

fragrance across a transition from outcrossing to selfing in Abronia umbellata

(Nyctaginaceae). American Journal of Botany 100: 2280–2292.

Douglas, N.A. 2008. Tripterocalyx carneus (Nyctaginaceae) Is Self-Compatible. The

Southwestern Naturalist 53: 403–406.

Ellstrand, N.C. 1983. Floral Formula Inconstancy Within and Among Plants and Populations of
Ipomopsis aggregata. Botanical Gazette 144: 119–123.

Endress, P.K., 2001. Origins of flower morphology. Journal of Experimental Zoology 291: 105–
115

Galen, C. 1999. Why Do Flowers Vary? BioScience 49: 631.

Golding, Y.C., M.S. Sullivan, and J.P. Sutherland. 1999. Visits to manipulated flowers by
Episyrphus balteatus (Diptera: Syrphidae): partitioning the signals of petals and anthers.
Journal of Insect Behavior 12: 39–45.

Grant, V. 1956. The genetic structure of races and species in Gilia. In Advances in Genetics, 55–
87. Academic Press.

Greer, S.U. 2006. Genomic consequences of mating system evolution in the Pacific coastal dune
endemic Abronia umbellata (Nyctaginaceae). Queens University, Kingston, Ontario,
Canada.

Herrera, C.M. 2009. Multiplicity in Unity. Chicago, U. University of Chicago Press.

Hoffmann, A.A., P.C. Griffin, and R.D. MacRaild. 2009. Morphological variation and floral
abnormalities in a trigger plant across a narrow altitudinal gradient. Austral Ecology 34:
780–792.

Huether, C.A. 1968. Exposure of natural genetic variability underlying the penamerous corolla
constancy in Linanthus androsaceus ssp androsaceus. Genetics 60: 123–146.

Huether, C.A. 1969. Constancy of the pentamerous corolla phenotype in natural populations of
Linanthus. Evolution 23: 572–588.

Katsuyoshi, K., and J. Harding. 2015. Genetic and environmental variation for corolla traits in

656   Portulaca grandiflora. Journal of Horticultural Science 44: 37–47.

657   Keeler, K.H. 1981. Function of Mentzelia nuda (Loasacae) postfloral nectaries in seed defense.

658   American Journal of Botany 68: 295–299.

659   Keeler, K.H. 1979. Nocturnal pollination of Abronia fragrans (Nyctaginaceae). Southwestern

660   Naturalist 24: 692–693.

661   Kido, A., and M. Hood. 2020. Mining new sources of natural history observations for disease

662   interactions. American Journal of Botany 107: 3–11.

663   Kitazawa, M.S., and K. Fujimoto. 2014. A developmental basis for stochasticity in floral organ

664   numbers. Frontiers in Plant Science 5: 1–14.

665   Lambrecht, S.C. 2013. Floral water costs and size variation in the highly selfing Leptosiphon

666   bicolor (Polemoniaceae). International Journal of Plant Sciences.

667   Lehmann, N.L., 1987. Floral formula variation in Phlox drummondii Hook. University of Texas

668   at Austin.

669   Leppik, E.E., 1953. The Ability of Insects to Distinguish Number. American Naturalist 87: 229–

670   236.

671   Lowndes, A.G., 1931. Note on Individual Variation in Paris quadrifolia L. New Phytologist 30:

672   298–299.

673   Mathiasen, R.L. 1982. Taxonomic studies of dwarf mistletoes (Arceuthobium ssp.) parasitizing

674   Pinus strobiformis. Great Basin Naturalist 42: 120–127.

675   Mazer, S.J., V.A. Delesalle, and P.R. Neal. 1999. Responses of Floral Traits to Selection on

676   Primary Sexual Investment in Spergularia marina: The Battle between the Sexes.

677   Evolution 53: 717–731.

678   McKim, S.M., A.L. Routier-Kierzkowska, M. Monniaux, D. Kierzkowski, B. Pieper, R.S. Smith,

679   M. Tsiantis, and A. Hay. 2017. Seasonal regulation of petal number. Plant Physiology.

680   Mickley, J., 2017. The Adaptive Nature of Stasis for Petal Number: Can Pollinator-Mediated

681   Stabilizing Selection Explain Five-petaled Flowers? Storrs: University of Connecticut.

682   Mickley, J., and C. Schlichting. 2018. Revisiting an old question in California botany : Why do

683   many plant species have five-petaled flowers? Mojave National Preserve Science

684   Newsletter 2018: 13–16.

685   Monniaux, M., B. Pieper, and A. Hay. 2016. Stochastic variation in Cardamine hirsuta petal

686   number. Annals of Botany.

687   Monniaux, M., B. Pieper, S.M. McKim, A.L. Routier-Kierzkowska, D. Kierzkowski, R.S. Smith,

688   and A. Hay. 2018. The role of APETALA1 in petal number robustness. eLife.

689   Pélabon, C., M.L. Carlson, T.F. Hansen, N.G. Yoccoz, and W.S. Armbruster. 2004.

690   Consequences of inter-population crosses on developmental stability and canalization of

691   floral traits in Dalechampia scandens (Euphorbiaceae). Journal of Evolutionary Biology.

692   Pieper, B., M. Monniaux, and A. Hay. 2016. The genetic architecture of petal number in

693   Cardamine hirsuta. New Phytologist 209: 395–406.

694   Roman, H., M. Rapicault, A.S. Miclot, M. Larenaudie, K. Kawamura, T. Thouroude, A.

695   Chastellier, et al. 2015. Genetic analysis of the flowering date and number of petals in

696   rose. Tree Genetics and Genomes 11: . Available at: http://dx.doi.org/10.1007/s11295-

697   015-0906-6.

698 Roy, S. 1963. The variation of organs of individual plants. Journal of Genetics 58: 147–176.

699 Saunders, E.R. 1934. A study of Veronica from the viewpoint of certain floral characters. Journal

700   of the Linnean Society of London, Botany 49: 453–493.

701 Schemske, D.W. 1978. Sexual reproduction in an Illinois population of Sanguinaria canadensis

702   L. The American Midland Naturalist 100: 261–268.

703 Shepherd, K.A., T.D. Macfarlane, and M. Waycott. 2005. Phylogenetic analysis of the Australian

704   Salicornioideae (Chenopodiaceae) based on morphology and nuclear DNA. Australian

705   Systematic Botany 18: 89–115.

706 Soltis, P.S. & Soltis, D.E., 2013. Flower diversity and angiosperm diversification. In J. L.

707   Riechmann & F. Wellmer, eds. Flower Development: Methods and Protocols. New York:

708   Springer New York, pp. 85–102.

709 Stark, P., 1918. Die Blütenvariationen der Einbeere. Zeitschrift für induktive Abstammungsund

710   Vererbungslehre 19: 241–303.

711 Stebbins, G.L., Jr., 1974. Flowering Plants: Evolution Above the Species Level, Cambridge:

712   Harvard University Press.

713 Stevens, P.T., C.A. Huether, and T.K. Wilson. 1972. Apical size in the determination of corolla

714   lobe number in Linanthus androsaecus ssp androsaecus. American Journal of Botany 59:

715   989–992.

716   Strauss, S.Y., and J.B. Whittall. 2006. Non-pollinator agents of selection on floral traits. Ecology

717   and evolution of flowers.

718   Tooke, F., and N.H. Battey. 2000. A leaf-derived signal is a quantitative determinant of floral

719   form in Impatiens. The Plant Cell 12: 1837–1847.

720   Vlot, E.C., W.H.J. van Houten, S. Mauthe, and K. Bachmann. 1992. Genetic and Nongenetic

721   Factors Influencing Deviations from Five Pappus Parts in a Hybrid between Microseris

722   douglasii and M. bigelovii (Asteraceae, Lactuceae). International Journal of Plant

723   Sciences.

724   Warren, J. 2009. Extra petals in the buttercup (Ranunculus repens) provide a quick method to

725   estimate the age of meadows. Annals of Botany 104: 785–788.

Table 1: Evolutionary and ecological hypotheses for atypical petal counts, with predictions for

Abronia umbellata, and references.

Table 2: Summed counts of floral morphs in common garden *Abronia umbellata* (both varieties)

and hybrids of this species. Statistics not performed because of low sample sizes.

Figure 1. A: *Abronia umbellata* var. *breviflora* with many 4-petalled and a 3-petalled (arrow)

flower. Doran Beach, Sonoma County, CA. B. *A. u.* var. *umbellata* with a 6-petalled flower

(arrow). Morro Beach, San Luis Obispo County, CA. (C) *A. u.* var. *breviflora* with an atypically

low number of 4-petalled flowers (none visible). Doran Beach, Sonoma County, CA. D. A. u.

var. *umbellata* with a six-petalled flower (arrow). Laboratory plant. E. Individually marked

flowers in autogamy experiment on *A. u.* var. *breviflora*. F. Developing fruit with tape denoting

petal counts. All photos by EFL.

Figure 2: The range of iNaturalist records used in this study, with proportions of floral morphs as

depicted for visualization as pie charts for ease of visualization of each variety.

Figure 3: Summation of iNaturalist data. Left: histograms of proportion of flowers 5-petalled for

each variety, by infloresence (var. *breviflora*, n=238 inflorescences, var. *umbellata* n=1570);

Right: histogram of petal numbers for all flowers of each variety comprising (var. breviflora,

n=1355 flowers; var. umbellata n=10,070 flowers).

Figure 4: Box plot of mean petal numbers, per inflorescence, of each variety, by inflorescence,

from the iNaturalist data set (derived from the same data as in Figure 3, left).

Figure 1:

Figure 2:

766   Figure 3:

769   Figure 4

Hypotheses from literature tested here:

(H1) Loss of petals increases autogamous success

(H2) Inbreeding increases developmental instability

(H3) Incompatibilities due to hybridization increase atypical petal morphs

(H4) Correlated selection on other traits (here: flower size)

(H5) Environmentally-driven

Predictions for *Abronia umbellata*

In *A. u*. var. *breviflora*, fewer petalled-flowers will have higher reproductive success in greenhouse

A. u. var. *breviflora* (autogamous) will have increased atypical petal morphs

Artificially-created hybrids will have increased atypical petal morphs

A. u. var. *umbellata* (xenogamous), which hybridizes with congeners commonly in the field, will have an increased atypical petal morphs in field compared to autogamous variety

A. u. var. *breviflora* (smaller flowered) will have fewer mean petals

Within each variety, populations with smaller flowers will have fewer mean petals

A correlation between phenology and atypical petal morphs for each variety

A correlation between latitude and atypical petal morphs for each variety

	# individuals, inflorescences	8 Petals	7 Petals	6 Petals	5 Petals
umbellata var. *breviflora*	10, 63	0	0	0	493
umbellata var. *umbellata*	6,16	0	0	0	259
hybrids					
umbellata x *breviflora*	3,3	0	0	0	33
hybrids of umbellata					
bigelovii x *umbellata*	2,3	0	0	0	56
pogonantha x *umbellata* [2,3]	12,51	1	0	3	674
umbellata x *pogonantha* [3]	1,4	0	0	0	56
umbellata x *ameliae*	3,10	0	0	1	231
umbellata x *turbinata*	9,42	0	2	11	890
umbellata x *villosa* var. *villosa* [3]	2,5	0	0	0	56
villosa var. *villosa* x *umbellata* [3]	7,19	2	0	8	241
villosa var. *aurita* x *umbellata* [3]	1,1	0	0	2	5
hybrids of breviflora					
pogonantha x *breviflora*	4,9	0	0	0	106

1: cumulative of all scored flowers, expected from the parental variety, 2: see text for information on this hyl
1967, 4: expected calculated from var. *breviflora* , 5: expected calculated from var. *umbellata* .

4 Petals	3 Petals	Proportion 5-petalled	Proportion 4-petalled	Chi-squared value[1]	p =	Greater or fewer atypical petals than expected?
163	9	0.74	0.25			
20	1	0.93	0.07			
12	0	0.73	0.27	0.70^{4}	0.353	no difference
				25.96^{5}	**<0.001**	greater
10	0	0.85	0.15	6.58	**0.019**	greater
2	0	0.99	0.00	50.37	**<0.001**	fewer
0	0	1.00	0.00	4.54	0.052	no difference
1	0	0.99	0.00	16.65	**<0.001**	fewer
2	0	0.98	0.00	67.37	**<0.001**	fewer
2	0	0.97	0.03	1.42	0.246	no difference
3	0	0.95	0.01	13.67	**<0.001**	fewer
0	0	0.71	0.00	0.86	0.325	no difference
2	0	0.98	0.02	32.49	**<0.001**	fewer

brid, specifically 3: hybrid occurs or is very likely to occur in nature; see above note and Tillett