***In vitro* pectate lyase activity and carbon uptake assays and whole genome sequencing of**

***Bacillus amyloliquefaciens* subsp.** *plantarum* **strains for a pectin defective pathway**

Mohammad K. Hassan[1*],

[1]Department of Plant Sciences and Plant Pathology, Montana State University, 119 Plant

BioScience Building, Bozeman, MT 59717-3150, USA.

Correspondence: * E-mail: mohammad.hassan@montana.edu

Abstract

The pectin lyase activity of 59 *Bacillus amyloliquefaciens* subsp. *plantarum (Bap)*

strains was tested *in vitro* on Pectate Agar (PA) and Tris-Spizizen Salts (TSS) medium. Bap

strains were cultured on TSA medium and washed three times with sterile water before the

inoculation on PA media. Higher and lower pectate lyase activity were observed in six

(AP193, AP203, AP299, AP80, AP102, and AP52) and four (AP 194, AP214, AP215, and

AP305) *Bap* strains compared to other Bap strains. A total of 12 Bap strains (AP67, AP71,

AP77, AP78, AP85, AP102, AP108, AP135, AP143, AP189, AP192, and AP193) grew

vigorously on TSS medium. A total of six *Bap* strains (AP194, AP204, AP214, AP216,

AP219, and HD73) had lower growth compared to other *Bap* strains. Pectin (1%) were used

for *in vitro* PA and TSS medium. Pectate lyase and utilization activity were not found in

Bacillus thuringiensis subsp. *kurstaki* strain HD73 compared to *Bap* strains. A draft genome

sequence for strains AP194 and AP214 that were negative for pectin utilization were

generated using an Illumina MiSeq. RAST analysis revealed that the bpectin-associated gene

altronate hydrolase (*uxaA*) absent in AP 214 strain [Is this true?]. Multiple amino acid

alignments of *exuT* and *uxuB* gene sequence showed dissimilarities among AP194, AP214,

and reference *Bap* strains.

1. Introduction

Pectate lyase enzyme was first discovered in *Erwinia carotovora* and *Bacillus polymyxa* in 1962 (Starr, 1962) and has since been reported in many plant pathogenic and non-pathogenic bacteria such as *E. aroideae* (Kamimiya et al., 1977); *E. chrysanthemi* (Starr, 1972); *Clostridium multifermentas* (Macmillan, 1964); *Fusarium solani* (Crawford, 1987)*, B. amyloliquefaciens* (Fan X., 2008); and *B. subtilis* (Soriano et al., 2006). Mechanism of this enzyme activity in pathogenic and non-pathogenic bacteria could have a subtle difference in the mode of action. Pectate lyase breaks down polygalacturonate into D-galacturonate and D-glucuronate, making it available form for bacteria to use as a carbon source. Previous studies have reported that *B. subtilis* (Mekjian et al., 1999), *E. coli* K-12 (Nemoz et al., 1976), *E. carotovora* (Abbott, 2008), and *E. chrysanthemi* (Abbott, 2008) are capable of utilizing pectin as a sole carbon source and energy.

Whole genome sequencing is a useful method to find out the high resolution, base by base view, gene expression, and regulation of the strains (Anonymous, 2016). Rapid Annotation using Subsystem Technology (RAST) is designed to annotate the gene of complete prokaryotic genomes, and it uses the highest confidence first assignment that guarantees a high degree of genome consistency (Anonymous, undated). This study was designed to screen 59 Bap strains for pectate lyase and utilization of pectin as a sole carbon source. In addition, this study also included whole genome sequences analysis of two Bap strains AP194 and AP214 in figuring out their pectin defective genes. The purpose of this research was i) to screen a large collection of *Bap* strains for pectin degradation and utilization as a sole carbon source, ii) to conduct a comparative genomic analysis that includes *Bap* strains that lack the capacity for pectin utilization.

2. Material and methods

2.1. Bap strains preparation

A total of 59 Bap strains were streaked onto Tryptic Soy Agar (TSA) plates from the cryo stocks stored in the -80^{o}C and incubated at 28°C for 24 hours. One loopful of bacteria was inoculated into 10ml TSB in a glass test tube and placed into a shaking incubator at 28°C overnight using 220 Revolutions Per Minute (RPM) for 24-48 hours. Three replicates were used for each of the Bap strains.

2.2. Pectate lyase activity test

The bacteria grew from cryo stocks in the -80^{o}C on Tryptic Soy Broth (TSB) at 28°C overnight using 220 rpm for 5 ml culture. A one-ml aliquot was pipetted into the 1.5 ml microcentrifuge tube, and centrifugation was done for 5 minutes at 10,000 x g speed. The supernatant was discarded, and the process was repeated three times using sterile water. In the final bacterial pellet, 1 ml of the sterile water was added to a microcentrifuge tube and vortexed thoroughly to uniform the bacterial suspension. Then, it was transferred to 1 ml of the sterile water containing test tube to measure the turbidity of a bacterial suspension of all strains until the optical density at 600 nm was approximately 0.5. Twenty µl of this standardized bacterial suspension was used in triplicate onto pectate-agar (Pa) media (Kobayashi et al., 1999) to determine the pectin lyase activity. The pH 8.0 of 0.1M Tris-HCl buffer was adjusted for the medium and sterilized using 0.45 µm Nalgene syringe filter (Thermo Scientific, USA) separately. The pectate-agar media plates were incubated at 28°C for 24-48 hours and then 1% Cetyltrimethyl ammonium bromide (CTAB) was poured over the surface of the plate at room temperature.

2.3.Pectin carbon uptake test

The capacity of the 59 *Bap* strains to utilize as a sole carbon source were assessed

using a minimal Tris-Spizizen salt (TSS) (Shingaki et al., 2003) as the base media

supplemented with 1% plant pectin (citrus source) (Tokyo Chemical Industry Co., Ltd). The

TSS media was filter-sterilized using a 0.2 µm polyethersulfone (PES) vacuum filter unit

(VWR, USA) and adjusted the medium p^H 7.0 using 10N NaOH. Each of the bacterial

cultures was prepared for the pectin lyase assays with bacterial suspensions washed three

times in sterile water, normalized to an $OD_{600} = 0.5$, and then 100 microliters (µl) of a 1:100

dilution were used to inoculate 1.9 ml TSS + 1% pectin cultures to adjust the $OD_{600} = 0.030$,

in triplicate. Broth cultures were incubated at 28°C with 220 rpm, and OD_{600} readings were

reocorded over a 40 hr period. *Bap* strains AP 193 was used as the positive control and

Bacillus thuringiensis subsp. *kirstaki* HD 73 was used as the negative control. *Bacillus*

thuringiensis subsp. *kirstaki* HD 73 was obtained from the USDA-ARS culture collection

(Ames, Iowa) that was identified as a non-pectin utilizing strain based on its genome

sequence. This strain was not observed to growth using pectin as a sole carbon source.

2.4.Whole genome sequencing of Bap strains

The genomic DNA of AP194 and AP214 was extracted by E.Z.N.A. ® DNA Isolation Kit

and the DNA concentration was measured by Qubit® 2.0 Fluorometer (Thermo Fisher

Scientific, USA). Nextera DNA Library Preparation Kit (Illumina, Inc.) was used for

Illumina MiSeq® sequencing. A total of 50 ul reaction (3.75 µl genomic DNA, 25 µl TD

buffer, 5 µl enzyme, and 16.25 µl nuclease free water) was prepared using Nextera sample

prep kit. The prepared reaction was vortexed and centrifuged at 10,000x rpm before the

incubation for 5 minutes at 55°C. Nextera PCR program cycle was used for PCR

amplification using Eppendorf Thermal Mastercycler (Eppendorf AG, Hamburg). The

holding temperature was 10°C. Zymo Research DNA Clean & Concentrator™-5 was used

for purifying the tagmented DNA. A total of 250 µl DNA binding buffer was used for the 50

ul sample (5:1) and centrifuged for 30 seconds using 10,000 x g. A total of 200 µl DNA wash

buffer was added and centrifuged it for 30 seconds. This process was repeated twice. The

empty column was used for drying the sample and centrifuged for 2 minutes. A total of 25 µl

DNA wash buffer was added and kept room temperature for one minute. Then, 50 µl reaction

(20 µl tagmented DNA, 5 µl index 1, 5 µl index 2, 15 µl PCR master mix, and 5 µl PCR

primer coctail) was prepared using Nextera prep kit. The temperatures for the PCR cycles

were 12°C (3 min), 98°C (30 sec) for one cycle, 98°C (10 sec), 63°C (30 sec), and 72°C (3

min) for five cycles. The size-select purification kit was used for the cleaning of the 50 µl

PCR products. The last step was a measurement of the DNA library concentration. DNA

concentration was followed 16ng/µl, and 3.75µl genomic DNA were used for 60 ng

concentration. Before running the MiSeq system, all instruments were cleaned using 1X

Tween buffer. Then, all the samples were loaded to run the MiSeq® system. After successful

running the system, fastq.gz files were generated and saved as output files. Then, fastq.gz

output files were imported, trimmed, and de novo assembled.

2.5.RAST analysis

Rapid Annotation using Subsystem Technology (RAST) version 2.0 was used for

annotation of the whole genome sequences for strains AP194 and AP214. Fasta files of

AP194 and AP214 were uploaded in RAST server. The annotated genome was viewed in a

seed viewer link for AP194 and AP214.

2.6. CLC genomics analysis of *exuT* and *uxuB* protein gene sequence

Two pectin-associated genes (*exuT* and *uxuB*) sequences of 15 Bap reference strains were collected from the NCBI database. Each gene sequence was translated into an amino acid sequence using CLC Genomics Workbench 4.9 (CLC bio, Cambridge, MA, USA) software. Reference strains gene sequences were aligned with *exuT* and *uxuB* gene sequences of AP193, AP194, and AP214 strains.

3. Results

3.1. Bap strains preparation

A total of 59 Bap strains grew profusely within 24 hours in TSB and TSA plates. Most strains exhibited similar growth rate, but strains AP194, AP214, and AP52 grew slower than the rest. None of the Bap strains were cross contaminated.

3.2. Pectate lyase activity test

Clear zone appeared around bacterial colonies after 30 minutes. The magnitude of the zone of clearing was measured in milliliters (mm) and recorded in an Excel spreadsheet, with average zones of clearing determined for each of the *Bap* strains. The clear zone images of each plate were photographed using the AlphaImager® HP high-performance imaging System. The pectinase clear zone diameter of each strain is shown in Table 1.

158 **Table 1. The clear zones range from 22 mm to 55 mm in Bap strains.**

Bap strain	Pectinase clear zone diameter (mm)
AP52	52
AP67	41
AP71	32
AP75	42
AP76	40
AP77	38
AP78	39
AP79	52
AP80	50
AP81	39
AP85	38
AP86	35
AP87	50
AP108	39
AP112	42
AP135	37
AP136	39
AP143	42
AP150	40
AP183	41
AP184	35
AP188	55
AP189	40
AP190	45
AP191	41
AP192	42
AP193	39
AP194	36
AP195	38
AP196	43
AP197	43
AP198	40
AP199	41
AP200	44
AP201	40
AP203	52
AP205	40
AP207	35
AP208	56
AP210	39

159 **Table 2. The clear zones ranges from 22 mm to 55 mm in Bap strains**

Bap strain	Pectinase clear zone diameter (mm)
AP211	38
AP212	44
AP213	35
AP214	34
AP215	32
AP216	25
AP218	37
AP219	36
AP241	40
AP260	33
AP295	40
AP296	45
AP297	38
AP298	42
AP299	50
AP300	30
AP301	35
AP304	30
AP305	22

161   The highest clear zone activity was observed in AP193, AP203, AP299, AP80,

162 AP102, and AP52. The lowest clear zone was observed in AP 194, AP214, AP215, and

163 AP305. The clear zone was zero in HD73. The average pectate lyase activity was 35.45 mm.

170   **3.3.Pectin carbon uptake test**

171   **Table 3. Pectin utilization of 40 Bap strains.**

Bap strain	600nm OD value
AP52	0.36
AP67	0.51
AP71	0.51
AP75	0.49
AP76	0.39
AP77	0.55
AP78	0.49
AP79	0.4
AP80	0.32
AP81	0.35
AP85	0.57
AP86	0.4
AP87	0.45
AP108	0.66
AP112	0.52
AP135	0.52
AP136	0.44
AP143	0.49
AP150	0.35
AP183	0.54
AP184	0.6
AP188	0.72
AP189	0.37
AP190	0.27
AP191	0.67
AP192	0.66
AP193	0.68
AP194	0.33
AP195	0.36
AP196	0.34
AP197	0.38
AP198	0.35
AP199	0.29
AP200	0.24
AP201	0.33
AP203	0.46
AP205	0.34
AP207	0.24

173   **Table 4. Pectin utilization of 21 Bap strains.**

Bap strain	600nm OD value
AP208	0.38
AP210	0.15
AP211	0.2
AP212	0.22
AP213	0.29
AP214	0.2
AP215	0.09
AP216	0.38
AP218	0.1
AP219	0.21
AP241	0.1
AP260	0.17
AP295	0.18
AP296	0.11
AP297	0.22
AP298	0.22
AP299	0.19
AP300	0.05
AP301	0.09
AP304	0.2
AP305	0.11

175   **Table 5. 12 strains showed the highest level of pectin utilization based on OD.**

Bap strain	600nm OD value
AP67	0.51
AP71	0.51
AP77	0.55
AP78	0.49
AP85	0.57
AP108	0.66
AP135	0.52
AP143	0.49
AP189	0.37
AP192	0.66
AP193	0.68

Bap strain	600nm OD value
AP194	0.1
AP 205	0.24
AP214	0.09
AP216	0.1
AP218	0.21
AP219	0.1
HD73	0.09

Table 6. Six strains showed the lowest level of pectin utilization based on OD.

3.4.Whole genome sequencing of Bap strains

The GC (%) contents, genome size, largest contig, sequence reads, and average

coverage of AP 194 strains were 46.33, ~3.988 Mbp, 688,671bp, 1,179,872, and 28.77x. The

total read count, mean read length, and total read length were 1,132,586, 100.91, and

114,293,488. The total contig length was 3,971,326 and mean contig length was 86,333. The

GC (%) contents was 46.23, genome size was 4.039 Mbp, largest contig was 486,108 bp,

sequence reads was 1,056,594 bp, and average coverage was 28.12x in AP 214 strains. The

total read count was 1,015,508 bp, mean read length was 111.75, and total read length was

113,479,797 bp. The total contig lenth was 4,034,213 and mean contig length was 65,067.

3.5.RAST analysis

The number of contigs with protein coding genes was 59, the number of subsystems

was 462, the number of coding sequences was 4014, and the number of RNAs was 98 in

AP194 bap strain. The number of contigs with protein coding genes was 51, the number of

subsystems was 462, the number of coding sequences was 4060, and the number of RNAs

was 75 in AP214 bap strain. A total of 14 and 13 pectin-associated subsystem feature were

found in AP194 and AP214. The altronate hydrolase (uxaA) gene was found in AP194 Bap

strain. The uxaB gene was not found in AP214 Bap strain.

3.6.CLC genomics analysis of exuT and uxuB protein gene sequence alignment

In translation frame three of *exuT* gene sequence alignment, glycine (G) amino acid

was found in AP194 and AP214 Bap strains. However, cysteine (C) amino acid was found in

the same translation frame of reference strains CC178, FZB42, Trigocor1448, CAU-B946,

IT-45, and AS43.3. cysteine (C) also was available in AP193 Bap strains. Isoleucine (I) was

found only in AP214 Bap strains. phenylalanine (F) was found in AP193, AP194, and other

reference strains.

In translation frame one and two of *exuT* gene sequence alignment, arginine (R)

amino acid was found in AP194 and AP214 Bap strains. On the contrary, glycine (G), and

leucine (L) amino acid were found in the same translation frame of reference strains CC178,

FZB42, Trigocor1448, CAU-B946, IT-45, and AS43.3. Arginine (R) also was found in

AP193 Bap strains.

In translation frame two and three of *uxuB* gene sequence alignment, isoleucine (I),

tryptophan (W), methionine (M), tyrosine (Y), and histidine (H) were found in AP 194, and

AP214 Bap strains. However, threonine (T), cysteine (C), valine (V), leucine (L), and

arginine (R) amino acid were found in the same translation frame of reference strains

UCMB5113, UCMB5033, CC178, FZB42, Trigocor1448, CAU-B946, IT-45, NAU-B3, Y2,

and AS43.3.

4. Discussion

The presented results indicate that Bap strains had slower growth rates on pectate agar

media than on TSA media. From a total of 59 Bap strains, three strains (AP207, AP214, and

AP215) did not grow very well on PA media. The remaining 56 Bap strains growth had

started after 6 hr inoculation time. The pectate lyase test results demonstrate that six Bap

strains had the largest clear zone around the bacterial colony. Two Bap strains (AP194 and

AP214), and one strain (HD73) had the lowest and zero clear zone around the colony. Based

on the results, it can be concluded that the highest clear zone showing strains have the highest

pectate lyase activity. In contrast, the lowest clear zone showing strains had the lowest

pectate lyase activity, and zero clear zone strain has no activity. Previous studies found that

the clear zone formed in *Bacillus sp.* Strain KSM-P15 (Kobayashi et al., 1999) in 10

minutes. However, the clear zone around the Bap strains colony were observed after 30

minutes.

Based on the pectin carbon uptake test, 12 *Bap* strains grew vigorously in 1% pectin.

The remaining *Bap* strains have showed lowest and average growth in TSS medium. Two

Bap strains AP194 and AP214 did not grow in TSS medium. *Btk* strain HD73 also did not

grow in TSS medium. Because, *Btk* strain HD73 had no pectin-associated genes. It can be

inferred from the results that *exuT* and *uxuB* pectin-associated genes from the fastest growing

strains might transport and metabolize faster than other strains.

Rast results of AP193, AP194, and AP214 Bap strains have revealed that they have

differences among them. Pectin pathways associated subsystem features have counted in

three Bap strains are 16, 14, and 13. Altronate hydrolase (*uxaB*) gene has found in AP193 and

AP194 only, not present in AP214 Bap strain. However, pairwise Blastx with different

reference genome sequences has indicated that it is present in three Bap gene sequences. It is

also found that pectin-associated rest of the genes are present in three Bap strains.

Based on the *exuT* protein gene sequence alignment results (figure 17), it can be

concluded that Glycine (G), and Arginine (R) amino acid differences exist in Bap strains

AP194 and AP214 in comparison with reference strain FZB42 (Chen et al., 2007). This type

of differences might affect the hexuronate transport of D-glucuronate and D-galacturonate

chemical compounds into the bacterial cell.

The *uxuB* protein gene sequence alignment results (figure 16) indicate that Isoleucine

(I), Tryptophan (W), Methionine (M), Tyrosine (Y), and Histidine (H) amino acid changes

occurred in AP194 and AP214 in comparison with reference strain FZB42 (Chen et al.,

2007). This change could hamper the overall metabolic activity of strains AP194 and AP214.

Reference

Abbott, D.W.B., Alisdair B., 2008. Structural Biology of Pectin Degradation by

Enterobacteriaceae. Microbiology and Molecular Biology Reviews : MMBR 72**,** 301-

316.

Anonymous, 2016. Advantages of Whole-Genome Sequencing.

http://www.illumina.com/techniques/sequencing/dna-sequencing/whole-genome-

sequencing.html.

Anonymous, undated Using the RAST prokaryotic genome annotation server

www.nmpdr.org.

Chen, X.H., Koumoutsi, A., Scholz, R., Eisenreich, A., Schneider, K., Heinemeyer, I.,

Morgenstern, B., Voss, B., Hess, W.R., Reva, O., Junge, H., Voigt, B., Jungblut, P.R.,

Vater, J., Sussmuth, R., Liesegang, H., Strittmatter, A., Gottschalk, G., Borriss, R.,

2007. Comparative analysis of the complete genome sequence of the plant growth-

promoting bacterium Bacillus amyloliquefaciens FZB42. Nat Biotechnol 25**,** 1007-

1014.

Crawford, M.S.K., P. E., 1987. Pectate lyase from Fusarium solani f. sp. pisi: purification,

characterization, in vitro translation of the mRNA, and involvement in pathogenicity.

Arch Biochem Biophys 258**,** 196-205.

Fan X., Q.S., Hu F, 2008. Bacillus amyloliquefaciens strain TB-2 pel gene for pectate lyase.

EMBL/GenBank/DDBJ databases.

Kamimiya, S., Itoh, Y., Izaki, K., Takahashi, H., 1977. Purification and Properties of a

Pectate Lyase in *Erwinia aroideae*. Agricultural and Biological Chemistry 41**,** 975-

981.

Kobayashi, T., Koike, K., Yoshimatsu, T., Higaki, N., Suzumatsu, A., Ozawa, T., Hatada, Y.,

Ito, S., 1999. Purification and properties of a low-molecular-weight, high-alkaline

pectate lyase from an alkaliphilic strain of Bacillus. Biosci Biotechnol Biochem 63**,**

65-72.

Macmillan, J.D.V., R. H., 1964. PURIFICATION AND PROPERTIES OF A

POLYGALACTURONIC ACID-TRANS-ELIMINASE PRODUCED BY

CLOSTRIDIUM MULTIFERMENTANS. Biochemistry 3**,** 564-572.

Mekjian, K.R., Bryan, E.M., Beall, B.W., Moran, C.P., Jr., 1999. Regulation of hexuronate

utilization in Bacillus subtilis. J Bacteriol 181**,** 426-433.

Nemoz, G., Robert-Baudouy, J., Stoeber, F., 1976. Physiological and genetic regulation of

the aldohexuronate transport system in Escherichia coli. J Bacteriol 127**,** 706-718.

Shingaki, R., Kasahara, Y., Iwano, M., Kuwano, M., Takatsuka, T., Inoue, T., Kokeguchi, S.,

Fukui, K., 2003. Induction of L-form-like cell shape change of Bacillus subtilis under

microculture conditions. Microbiology 149**,** 2501-2511.

Soriano, M., Diaz, P., Pastor, F.I., 2006. Pectate lyase C from Bacillus subtilis: a novel endo-

cleaving enzyme with activity on highly methylated pectin. Microbiology 152**,** 617-

625.

Starr, M.P.C., A. K., 1972. The genus Erwinia: enterobacteria pathogenic to plants and

animals. Annu Rev Microbiol 26**,** 389-426.

Starr, M.P.M., F., 1962. Eliminative split of pectic substances by phytopathogenic soft-rot

bacteria. Science 135**,** 920-921.

309 ***In vitro* pectin utilization tests on Bap strains**

310 Figure 1: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

311 AP75, AP76, AP77, and AP215).

321 Figure 2: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP52,

322 AP67, AP71, and AP193).

332 Figure 3: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

333 AP73, AP78, AP79, and AP80).

Figure 4: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193, AP81, and AP85).

Figure 5: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193, AP86, AP87, AP102, and AP108).

Figure 6: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

AP112, AP135, AP136, and AP150).

Figure 7: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193, AP190, AP191, AP192, AP194, AP214, AP188, and AP189).

Figure 8: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193, AP195, AP196, and AP197).

410   Figure 9: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

411   AP198, AP199, and AP200).

421   Figure 10: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

422   AP201, AP202, and AP203).

436   Figure 11: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

437   AP204, AP205, and AP207).

446   Figure 12: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

447   AP208, AP210, and AP211).

461   Figure 13: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

462   AP212, AP213, and AP214).

471   Figure 14: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

472   AP214, AP215, and AP216).

487   Figure 15: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

488   AP218, AP219, AP241, and AP260).

498   Figure 16: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

499   AP295, AP296, and AP297).

513   Figure 17: *in vitro* Bap bacterial growth using pectin as a sole carbon source (HD73, AP193,

514   AP298, AP299, AP300, AP301, AP304, and AP305).

539 **Pectate lyase activity of Bap strains**

540 Figure 18: Pectate lyase activity of Bap strains (AP52, AP67, AP71, and AP75).

546 Figure 19: Pectate lyase activity of Bap strains (AP76, AP73, HD73, and AP78).

548 Figure 20: Pectate lyase activity of Bap strains (AP79, AP80, AP81, and AP85).

557   Figure 21: Pectate lyase activity of Bap strains (AP86, AP87, AP102, and AP108).

Figure 22: Pectate lyase activity of Bap strains

560   (AP112, AP135, AP136, and AP143).

562   Figure 23: Pectate   562   lyase

activity of Bap strains (AP180, AP183, AP184, and AP188).

573   Figure 24: Pectate lyase activity of Bap strains (AP190, AP191, AP192, and AP194).

575   Figure 25: Pectate lyase activity of Bap strains (AP195, AP197, AP198, and AP199).

577   Figure 26: Pectate lyase activity of Bap strains (AP200, AP193, AP201, and AP202).

584   Figure 27: Pectate lyase activity of Bap strains (AP203, AP204, AP205, and AP208).

586   Figure 28: Pectate lyase activity of Bap strains (AP210, AP211, AP212, and AP213).

588   Figure 29: Pectate lyase activity of Bap strains (AP215, AP216, AP218, and AP241).

598   Figure 30: Pectate lyase activity of Bap strains (AP260, AP295, AP219, and AP296).

600   Figure 31: Pectate lyase activity of Bap strains (AP298, AP297, AP304, and AP297).

602   Figure 32: Pectate lyase activity of Bap strains (AP299, AP214, AP301, and AP305).

616   Figure 33: Multiple amino acid alignment of *uxuB* gene (AP193, AP194, and AP214) with
617   reference strains.

620 Figure 34: Multiple amino acid alignment of *exuT* gene (AP193, AP194, and AP214) with
621 reference strains.