A FIELD-WIDE ASSESSMENT OF DIFFERENTIAL HIGH THROUGHPUT SEQUENCING REVEALS WIDESPREAD BIAS

Taavi Päll
Department of Microbiology
Institute of Biomedicine and Translational Medicine, University of Tartu, Estonia
Ravila 19, 50411, Tartu, Estonia
taavi.pall@ut.ee

Hannes Luidalepp
Quretec (https://www.quretec.com)
Ülikooli 6a, 51003, Tartu, Estonia
luidale@gmail.com

Tanel Tenson
Institute of Technology, University of Tartu, Estonia
Nooruse 1, 50411, Tartu, Estonia
tanel.tenson@ut.ee

Ülo Maiväli *
Institute of Technology, University of Tartu, Estonia
Nooruse 1, 50411, Tartu, Estonia
ymaivali@gmail.com

January 26, 2021

ABSTRACT

Here we assess reproducibility and inferential quality in the field of differential HT-seq, based on analysis of datasets submitted 2008-2019 to the NCBI GEO data repository. Analysis of GEO submission file structures places an overall 59% upper limit to reproducibility. We further show that only 23% of experiments resulted in theoretically expected p value histogram shapes, although both reproducibility and p value distributions show marked improvement over time. Uniform p value histogram shapes, indicative of <100 true effects, were extremely few. Our calculations of π_0, the fraction of true nulls, showed that 36% of experiments have π_0 <0.5, meaning that in over a third of experiments most RNA-s were estimated to change their expression level upon experimental treatment. Both the fraction of different p value histogram types and π_0 values are strongly associated with the software used for calculating these p values by the original authors, indicating widespread bias.

Introduction

Over the past decade a feeling that there is a crisis in experimental science has increasingly permeated the thinking of methodologists, captains of industry, working scientists, and even the lay public (Ioannidis 2005; Baker 2016; Begley and Ellis 2012; Prinz, Schlange, and Asadullah 2011; Harris 2017). This manifests in poor statistical power to find true effects (Button et al. 2013), in poor reproducibility (defined as getting identical results when reanalysing the original data by the original analytic workflow), and in poor replicability (defined as getting similar results after repeating the entire experiment) of the results (Goodman, Fanelli, and Ioannidis 2016). While reproducibility depends on the availability of data and on the quality of description of the data analytic workflow, replicability depends on the quality of experiments, on the quality of data analysis, including data pre-processing, and on the power of the experiment to detect true effects. The proposed reasons behind the crisis include sloppy experimentation, selective publishing, perverse

---

*corresponding author

incentives, difficult-to-run experimental systems, insufficient sample sizes, over-reliance on null hypothesis testing, and much-too-flexible analytic designs combined with hypothesis-free study of massively parallel measurements (Grimes, Bauch, and Ioannidis 2018; Maiväli 2015; Munafò et al. 2017; Szucs and Ioannidis 2017; Botvinik-Nezer et al. 2020; Leng and Leng 2020). Although there have been attempts at assessing experimental quality through replication of experiments, prohibitive cost and theoretical shortcomings in analysing concordance in experimental results have encumbered this approach (Hardwicke et al. 2020). Here we assess *in silico* the reproducibility and replicability of high throughput differential expression studies by next-generation sequencing (HT-seq). We concentrate on the HT-seq field for two reasons. HT-seq has become the gold standard for whole transcriptome gene expression quantification, both in research and in clinical applications. And secondly, due to the massively parallel testing in individual studies of tens of thousands of features per experiment, we have access to study-wide unbiased lists of p values. From the shapes of histograms of p values we can find the experiments where p values were calculated apparently correctly, and from these studies we can determine the study-wise relative frequencies of true nulls (the π_0-s). Also, we believe that the very nature of the HT-seq field, where a single biological experiment entails comparing the expression levels of about 20,000 features (e.g. RNA-s) on average, predicates that the quality of data analysis, and specifically statistical inference based on p values (directly, or indirectly through FDR) must play a decisive part in scientific inference. Simply, one cannot analyse an HT-seq experiment intuitively, without resorting to formal statistical inference. Therefore, quality problems of statistical analysis would very likely directly and substantially impact the quality of science. Thus we use the quality of statistical analysis as a proxy for the quality of science, with the understanding that this proxy may work better for modern data-intensive fields, where scientist's intuition has a comparatively smaller role to play.

Results

Assessing reproducibility by NCBI GEO database supplementary files

We queried the NCBI GEO database for "expression profiling by high throughput sequencing", retrieving 32,017 datasets (GEO series) from 2006, when first HT-seq dataset was submitted to GEO, to Dec-31, 2019. The number of yearly new HT-seq submissions increased from 1 in 2006 to 8770 by 2019, making up 27.4% of all GEO submissions in 2019. Most of the GEO HT-seq submissions are from human and mouse, and they increase in a much faster rate than submissions from other taxa (data not shown). We filtered the GEO series containing supplementary processed data files. Processed data are a required part of GEO submissions, defined as the data on which the conclusions in the related manuscript are based. The format of processed data files submitted to GEO is not standardized, but in case of expression profiling such files include, but are not limited to, quantitative data for features of interest, e.g. mRNA, in tabular format. Sequence read alignment files and coordinates (SAM, BAM, and BED) are not considered as processed data by GEO. As reproducibility by definition entails arriving at the original conclusions by independently repeating original analysis (Peng 2011), we surmise that the fraction of GEO submission with processed data files gives an upper limit of reproducibility (upper limit, because processed data files *per se* do not guarantee reproducibility). According to our analysis 17,920 GEO series, containing 43,340 supplementary processed data files, conform with GEO submission guidelines. After further excluding the submissions with potentially non-tabular supplementary files, based on file extensions (see Methods for details), the number of GEO series was reduced to 15,520, containing 31,862 supplementary files, including tar.gz archives, which we downloaded from the GEO server. From those we programmatically imported 32,764 files as tables, resulting in 32,414 (99%) successfully imported files. For the purpose of reproducibility analysis, we considered all 17,920 GEO series with supplementary processed files out of 32,017 published GEO series in our time window as potentially reproducible. Therefore, the observed overall 56% retention rate of GEO submissions with processed data files should be seen as an upper bound for reproducibility. There is a substantial increase of the retention rate over time, from 16% (3/19) in 2008 to 62% (5,465/8,770) in 2019, indicating increasing reproducibility of the HT-seq field (Figure 1).

According to GEO submission requirements, the processed data files may contain raw counts of sequencing reads, and/or normalized abundance measurements. Therefore, a valid processed data submission may or may not contain lists of p values. We identified p values from 2,109 GEO series, from which we extracted 6,267 unique p value sets. While the mean number of p value lists, each list corresponding to a separate experiment, per 2,109 GEO submissions was 2.97 (max 66), 49% of submissions contained a single p value list and 78% contained 1-3 p value lists. For further analysis we randomly selected one p value list per GEO series.

P value histograms

We algorithmically classified the p value histograms into five classes (Breheny, Stromberg, and Lambert 2018) (See materials and methods for details and Figure 2 for representative examples). The "Uniform" class contains flat p value histograms indicating no true effects (at the sample sizes used to calculate these p values). The "Anti-Conservative" class

Figure 1: The increasing proportion of GEO submissions conforming with submission guidelines in regard of inclusion of processed data is estimated from a bernoulli logistic model. Line denotes linear model best fit using formula *conforms ~ year*, bernoulli likelihood. Shaded area denotes 95% credible region. N = 32,017.

Figure 2: Classes of p value histograms. A. Summary of p value histograms identified from GEO supplementary files. One p value set was randomly sampled from each GEO series where p values were identified. N = 2,109. 95% CI denotes credible intervals calculated from a binomial model. B. Examples of p value histogram classes. Red line denotes QC threshold used for dividing p value histograms into discrete classes.

Figure 3: Association of the p value histogram class with differential expression analysis tool. A. The increase in the proportion of anti-conservative histograms is accompanied by decreases mostly in the class "other", irrespective of the DE analysis tool. Lines denote best fit of linear model *class ∼ year + (year | de_tool)*, categorical likelihood. Shaded areas denote 95% credible regions. N = 2,109. B. Association of p value histogram type with DE analysis tool; data is restricted to 2018-2019 GEO submissions. Points denote linear model fit *class ∼ de_tool*, categorical likelihood. Error bars denote 95% credible intervals. N = 980.

contains otherwise flat histograms that contain a spike near zero. The "Conservative" class contains histograms that have a distinct spike close to one. The "Bimodal" histograms have two peaks, one at either end. The class "Other" contains a panoply of malformed histogram shapes (humps in the middle, gradual increases towards one, spiky histograms, etc.). The "Uniform" and "Anti-Conservative" histograms are the theoretically expected shapes of p value histograms.

We found that overall, 23% of the histograms fall into anti-conservative class, 12% were conservative, 30% bimodal and 36% fell into class "other". Only 3 of the 2,109 histograms were classified as "uniform". Logistic regression reveals a clear trend for increasing proportion of anti-conservative histograms, starting from <10% in 2010 and topping 25% in 2018 (Figure 3–figure supplement 1). Hierarchical modelling indicates that all differential expression (DE) analysis tools and sequencing platforms exhibit similar temporal increases of anti-conservative p value histograms (Figure 3–figure supplement 2-3). Multinomial hierarchical logistic regression further demonstrated that the increase in the fraction of anti-conservative histograms is accomplished by decreases mostly in the class "other", irrespective of the DE analysis tool (Figure 3A).

This positive temporal trend in anti-conservative p value histograms suggests improving quality of the HT-seq field. Rather surprisingly, Figure 3A also indicates that different DE analysis tools are associated with very different proportions of p value histogram classes, suggesting that quality of p value calculation, and therefore, quality of scientific inferences based on these p values depends on DE analysis tool. We further tested this conjecture in a simplified model, restricting our analysis to 2018-2019, the final years in our dataset (Figure 3B). As no single DE analysis tool dominates the field – cuffdiff 23%, deseq 33%, edgeR 13%, limma 2%, unknown 29% (see Figure 3–figure supplement 4 for temporal trends) –, a state of affairs where proportions of different p value histogram classes do not substantially differ between analysis tools would indicate lack of DE analysis tool-specific bias to the results. However, we find by multinomial regression that all p value histogram classes, except "uniform", which is largely unpopulated, depend strongly on the DE analysis tool used to calculate the p values (Figure 3B). This is confirmed by modelling the frequency of the anti-conservative p value histograms in binomial logistic regression (Figure 3–figure supplement 5A). Using the whole dataset of 6,267 p value histograms, or controlling for GEO publication year, taxon (human,

Figure 4: Association of the proportion of true null effects (π_0) with DE analysis tool. A. Histogram of π_0 values estimated from anti-conservative and uniform p value sets. N = 488. B. Robust linear model (*pi0 ∼ de_tool*, student's t likelihood) indicates association of π_0 with DE analysis tool. Points denote best estimate for the mean π_0 and error bars denote 95% credible intervals. N = 488. C. Hierarchical model (*pi0 ∼ year + (year | de_tool)*, student's t likelihood) reveals that all DE analysis tools are associated with temporally increasing proportion of true null effects. Lines denote best estimate for the mean π_0 and error bars denote 95% credible regions. N = 488.

mouse, and pooled other), of the RNA source or sequencing platform does not change this conclusion (Figure 3–figure supplement 5B-E). These results indicate that DE analysis tools bias the analysis of HT-seq experiments.

Proportion of true nulls

To further enquire into DE analysis tool-driven bias we estimated from user-submitted p values the fraction of true null effects (the π_0) for each HT-seq experiment. As non-anti-conservative sets of p values (excepting the "uniform") indicate problems during the respective experiments and/or data analyses, we only calculated the π_0 for datasets with anti-conservative and uniform p value distributions (n = 488). Nevertheless, the π_0-s show an extremely wide distribution, ranging from 0.999 to 0.06. Remarkably, 36% of the π_0 values are smaller than 0.5, meaning that in those experiments over half of the features (e.g. mRNA-s) are estimated to change their expression levels upon experimental treatment (Figure 4A). Conversely, only 21% of π_0-s exceed 0.8, and 8.5% exceed 0.9. Intriguingly, the peak of the π_0 distribution is not near 1, as might be expected from experimental design considerations, but there is a wide peak between 0.5 and 0.8 (median and mean π_0-s are both at 0.59). The median π_0-s range over 20 percentage points, from 0.5 to 0.7, depending on the DE analysis tool (Figure 4B). Using the whole dataset confirms this analysis, also producing narrower credible intervals, due to larger sample (N = 1,567) (Figure 4–figure supplement 1A).

In addition, mean π_0 tend to rise in time, similarly to fraction of anti-conservative p value histograms, and this increase is also common to all DE analysis tools (Figure 4C). Controlling for time, taxon or sequencing platform, did not substantially change the association of DE analysis tools with the π_0-s, except for the multilevel models, which resulted in substantially larger estimation uncertainty (Figure 4–figure supplement 1B-E). Recalculating the π_0-s with a different algorithm (Storey 2002) and reanalysing the data did not change these conclusions (data not shown). As there is a strong association between both proportion of anti-conservative p value histograms and π_0 with DE analysis tool, we further checked for, and failed to see, similar associations with variables from raw sequence metadata, such as the sequencing platform, library preparation strategies, library sequencing strategies, library selection strategies, and library layout (single or paired) (Figure 3–figure supplement 6-9, Figure 4–figure supplement 2-5). These negative results support the conjecture of specificity of the associations with DE analysis tools.

Discussion

In this work we have calculated five indicators of health of the HT-seq field: (i) the proportion of GEO submissions with published analysis endpoints in a tabular format, (ii) the relative fractions of classes of p value distribution shapes, (iii) the association of p value distribution shapes with DE analysis tool, (iv) the estimated proportion of true null effects (the π_0) for each experiment, and (v) the association of π_0 values with DE analysis tool. We believe that (i) indicates reproducibility, (ii) the quality of p value calculations (use of correct distributional models, etc.), (iv) the experimental design and data pre-processing choices, and that (iii) and (v) indicate bias.

Our analysis puts an upper bound of about 60% to differential HT-seq reproducibility in recent years, based on the presence of processed data files in the GEO submissions. To elucidate the meaning of this estimate, we must look into the relationship between the GEO submission data structures and the HT-seq workflows. The GEO repository requires three components for data submission: a metadata spreadsheet providing experimental design details, tabular processed data files, defined as the data on which the conclusions in the related manuscript are based (containing normalized abundance measurements or raw counts of sequencing reads), and raw data files (reads and quality scores as generated by the sequencing instrument). Even in the presence of raw data files, processed data files are practically necessary for assessing and reproducing the evidence behind the conclusions of a study, because of the large number of choices available to a discerning analyst. To put this in another way: scientific inference in the functional genomics field is based on null hypothesis testing and p values, and there isn't a single correct method to calculate them (nor is there one for FDR). Therefore, there is no guarantee that two independent analyses of the same data will obtain the same p values, and thus the same conclusions (Maiväli 2015). This means that access to original analytic choices (analysis code), or at least to p values, is needed for reproducibility.

The major points of divergence in the analytic pathway include aligning the sequences to genomic DNA, counts normalization, and differential expression testing (Nookaew et al. 2012; Sun and Zhu 2012). The analysis tools that are used for differential expression testing allow for a plethora of choices, including different distributional models, data transformations and basic analytic strategies, which can lead to different results through different trade-offs (Everaert et al. 2017; Nookaew et al. 2012). A similar state of affairs exists in the field of fMRI experiments, where reproducibility is questionable, and where 70 teams, testing the same hypotheses on the same data, used 70 different workflows, getting variable results (Carp 2012a, 2012b; Botvinik-Nezer et al. 2020). Taken together, this leads us to conclude that access to correctly annotated read counts is a minimum requirement for assessing reproducibility of read pre-processing and alignment. More desirably, also raw p values are necessary to judge the evidence behind conclusions of a paper. For full reproducibility, starting from raw sequencing data, complete analysis instructions and modelling choices should be provided, which is currently not a part of the GEO submission protocol. Thus our reproducibility estimate does not refer to full reproducibility, but to mere potential of independently recreating conclusions of a paper. Although full reproducibility of the HT-seq field still seems elusive, the robust temporal trend of improvement, documented by us, gives reason for optimism.

While there is also a positive trend for the increasing fraction of anti-conservative p value sets, a strong majority of them yet fall into shapes that are considered problematic for successful analysis, including FDR/q value calculations. In fact, the most common class of p value histograms, "other", encompasses a diverse mixture of unruly shapes least likely to lead to good downstream analysis and interpretation of these p values.

There are very few uniform p value distributions, suggesting relatively few true effects. This surprising result was confirmed by visual re-examination of p value histograms. As a technical comment, it should be noted, that the assigned class of the p value histogram depends on arbitrarily set bin size. Our use of 40 bins leads to histograms, where an experiment with even around 100 true effects could reasonably lead to uniform histogram shape, because of swamping of the lowermost bin with p values emanating from true null effects (see Figure 2–figure supplement 1).

Importantly, the proportions of different classes of p value distributions differ greatly between DE analysis tools, indicating analysis tool-specific bias. We see similar tool-specific bias (and similar temporal increase) in the value of estimated fractions of true null effects (π_0). As π_0 -s were calculated from anti-conservative p value sets only, this bias could mean (i) that the anti-conservativeness of a p value set is not a sufficient predictor of its quality in terms of further analysis, and/or (ii) that the algorithms implemented in differential expression analysis tools differ to a degree that introduces such bias downstream of the p value calculation.

The DE analysis tool-specific medians of π_0 values range from 0.43 (tool "unknown") to 0.68 (tool "cuffdiff"), and the total median π_0 is 0.52, showing that by this criterion in an average experiment about half of the cellular RNA-s are expected to change their expression levels.

In a biased situation the measured effects are a mixture of effects of the intended experimental treatment and of the undesirable effects of a more-or-less accidental choice of DE analysis tool, as no analytic workflow has been shown to

systematically outperform the others (Everaert et al. 2017; Nookaew et al. 2012). Determining the relative weight of DE analysis tool in this mixture requires careful case-by-case study, as this depends both on the performed experiment (its variations, effect sizes, and actual laboratory implementation), and on the particular analytic choices reflected in the shapes of p value distributions and in π_0 values. However, from the evidence at hand, notably from the strong associations with DE analysis tool, in combination with the preponderance of extreme p value distributions and low π_0-s, we can conclude that such bias must be substantial, as it is widespread. Another limitation of our study, also due to its large-scale nature, is our inability to pinpoint the sources of DE analysis tool-specific bias. However, a recent close study of 35 datasets shows data normalization to be a potentially important source of bias, which cannot be corrected by many of the currently widely used data normalization methods (Mandelboum et al. 2019; Quinn et al. 2019). Indeed, the widely used DE analysis tools use different RNA-seq data pre-processing strategies, all vulnerable to the situation where a large fraction of features change expression (McGee et al. 2019). In the light of our findings, the sources of bias clearly merit further case-by-case study of individual GEO submissions.

Could the relatively high frequency of low π_0-s reflect reality in the sense that in many experiments most RNA-s actually change expression levels? Although there is a small number of well-supported examples of this (Lin et al. 2012; Nie et al. 2012; Hu et al. 2014), it has been argued that the vast majority of genome wide differential gene expression studies ever conducted, including by HT-seq, have used experimental designs that would make it impossible to uncover such global effects, at least in a qualitatively accurate way (Lovén et al. 2012; Chen et al. 2016). The issue lies in the use of internal standards in normalizing the counts for different RNA-s (commonly normalizing for total read counts in each sample), which leads to completely wrong interpretations, if most genes change expression in one direction. To overcome this problem, one could use spike-in RNA standards like the External RNA Controls Consortium (ERCC) set (Ambion) or the Spike-in RNA Variants (SIRV) set (Lexogen) (Lun et al. 2017) and compositional normalization (McGee et al. 2019). However, even spike-in normalization requires great care to properly work in such extreme cases (Risso et al. 2014; Quinn et al. 2019), and outside single-cell RNA-seq it is used infrequently (McGee et al. 2019). In the absence of spike-in normalization, it seems likely that many or most of the low π_0 experiments represent technical failures, most likely during data normalization (McGee et al. 2019).

While the aim of this paper was to provide a birds-eye view on the HT-seq field, the dataset created to support this work provides added value by allowing access at individual GEO submission/experiment level. The dataset that accompanies this study allows to get a first estimate for the reproducibility and quality of conclusions of more than 30,000 HT-seq studies deposited in NCBI GEO. The dataset contains for every GEO submission the date of submission, association with publications, organisms studied, association with tabular files; and for every submitted tabular file it contains the name of file, the number of p value columns, the number of features, the p value histogram type and depiction of the histogram, π_0.

Finally, we note that our methodology can be adapted to study any type of experiment that results in at least several hundreds of parallel measurements/ p values, such as quantitative protein mass spectroscopy and metabolomics.

Strengths of the study:

- Our study is based on a large unbiased dataset, which allows for reliable quantification.
- All steps of the analysis are transparent and reproducible, as we provide full workflow and code for data mining, p value histogram classification, π_0 calculations, modelling, and figures.
- To the best of our knowledge this is the first large-scale study to offer quantitative insight into general quality of experimentation/data analysis/reproducibility of a fairly large field of biomedical science.
- The accompanying full dataset includes, for individual GEO submissions, useful quality control measures, like presence of tabular files describing analysis end-points, the shapes of the p value distributions and the estimated proportions of true null effects, providing evidence on the quality of inference of almost any HT-seq work in the literature.

Limitations of the study:

- It is a general study, whose measure of bias (statistical association inferred from many experiments) cannot be directly extended to single experiments. To do so requires (at least) incorporating additional information about the effect sizes encountered in the study of interest.
- From our dataset we cannot determine, which features of which differential expression analysis tools are responsible for the inferred bias. Neither can we say, which (if any) of the tools are better than the others. However, these questions have been addressed in the literature, and we hope that by stressing the seriousness of the problem, our study will inspire further work on the subject.
- Our estimate for reproducibility only provides an upper limit (presence of tabular files, which have the mere potential of containing enough pertinent information). Obtaining the true level of reproducibility requires

actual reproduction of individual analyses, which we feel, cannot be automated. Although presence of tabular supplementary files is not a good measure of reproducibility of an individual study, their absence give a strong indication for irreproducibility of a study.

Methods

NCBI GEO database queries were performed using Bio.Entrez Python package and by sending requests to NCBI Entrez public API. The query was 'expression profiling by high throughput sequencing[DataSet Type] AND ("2000-01-01"[PDAT] : "2019-12-31"[PDAT])'. FTP links from GEO datasets document summaries were used to download supplementary file names. Supplementary file names were filtered for downloading, based on file extensions, to keep file names with "tab", "xlsx", "diff", "tsv", "xls", "csv", "txt", "rtf", and "tar" file extensions. We dropped the file names where we did not expect to find p values using regular expression "filelist.txt|raw.tar$|readme|csfasta|(big)?wig|bed(graph)?|(broad_)?lincs".

Downloaded files were imported using Python pandas package, and searched for unadjusted p value sets. Unadjusted p value sets and summarized expression level of associated genomic features were identified using column names. P value columns from imported tables were identified by regular expression "p[^a-zA-Z]{0,4}val", from these, adjusted p value sets were identified using regular expression "adj|fdr|corr|thresh" and omitted from further analysis. Columns with expression levels of genomic features were identified by using following regular expressions: "basemean", "value", "fpkm", "logcpm", "rpkm", "aveexpr".

Raw p value sets were classified based on their histogram shape. Histogram shape was determined based on the presence and location of peaks. P value histogram peaks (bins) were detected using a quality control threshold described in (Breheny, Stromberg, and Lambert 2018), a Bonferroni-corrected alpha-level quantile of the cumulative function of the binomial distribution with size m and probability p. Histograms, where none of the bins were over QC-threshold, were classified as "uniform". Histograms, where bins over QC-threshold started either from left or right boundary and did not exceeded 1/3 of the 0 to 1 range, were classified as "anti-conservative" or "conservative", respectively. Histograms with peaks or bumps in the middle or with non-continuous left- or right-side peaks were classified as "other". Histograms with peaks on both left- and right-side were classified as "bimodal".

Raw p value sets with anti-conservative shape were used to calculate the π_0 statistic. The π_0 statistic was calculated using local FDR method implemented in limma::PropTrueNullByLocalFDR (Ritchie, Phipson, Wu, et al. 2015b) and, independently, Storey's global FDR smoother method (Storey 2002) as implemented in gdsctools (Cokelaer et al. 2017) Python package. Differential expression analysis tools were inferred from column names pattern for cuffdiff (column name = "fpkm" and "p_value") (Trapnell et al. 2013), DESeq/DESeq2 (column name = "basemean") (Love, Huber, and Anders 2014), EdgeR (column name = "logcpm") (McCarthy, Chen, and Smyth 2012), and limma (column name = "aveexpr") (Ritchie, Phipson, Wu, et al. 2015a), all other unidentified sets were binned as "unknown".

Publication data were downloaded from NCBI PubMed database using PubMedId-s from GEO document summaries. Citation data were downloaded from Elsevier Scopus database using doi-s. Sequence read library statistics were downloaded from NCBI SRA database using GEO accessions.

The code to produce raw dataset is available as a snakemake workflow (Köster and Rahmann 2012) on rstats-tartu/geo-htseq Github repo (`https://github.com/rstats-tartu/geo-htseq`). Raw dataset produced by the workflow is deposited in Zenodo `https://zenodo.org` with doi: 10.5281/zenodo.4046422 (`http://doi.org/10.5281/zenodo.4046422`).

RNA-seq experiment simulation was done with polyester R package (Frazee et al. 2015) and differential expression was assessed using DESeq2 R package (Love, Huber, and Anders 2014) using default settings. Code and workflow used to run and analyse RNA-seq simulations is available on Github: `https://github.com/rstats-tartu/simulate-rnaseq`. Processed data, raw data and workflow with input fasta file is deposited in Zenodo with doi: 10.5281/zenodo.4463803 (`http://doi.org/10.5281/zenodo.4463803`).

The code to produce article's figures and fit models is deposited on rstats-tartu/geo-htseq-paper Github repo (`https://github.com/rstats-tartu/geo-htseq-paper`). Article's input data, code, software required to produce all models and figures in Linux, along with fitted model objects is deposited in Zenodo with doi: 10.5281/zenodo.4469911 (`https://zenodo.org/deposit/4469911`). Statistical analysis was done in R vers. 4.0.3. Bayesian modelling was done using R libraries rstan (vers. 2.21.2; `http://mc-stan.org/`) and brms (vers. 2.13.3) (Bürkner 2018). ggplot2 (vers. 3.3.1) (Wickham 2016) R library was used for graphics. Data wrangling was done using tools from tidyverse package (Wickham et al. 2019).

Acknowledgments

We are grateful for Toomas Mets (University of Tartu) for critically reading the manuscript, and Niilo Kaldalu (University of Tartu) and Margus Pihlak (Tallinn University of Technology) for useful discussions. The work was supported by the European Union from the European Regional Development Fund through the Centre of Excellence in Molecular Cell Engineering (2014-2020.4.01.15-0013) and by the grants from the Estonian Research Council (PRG335, PUT1580).

References

Baker, M. 2016. "1,500 scientists lift the lid on reproducibility." *Nature* 533 (7604): 452–54. `https://doi.org/10.1038/533452a`.

Begley, C. G., and L. M. Ellis. 2012. "Drug development: Raise standards for preclinical cancer research." *Nature* 483 (7391): 531–33. `https://doi.org/10.1038/483531a`.

Botvinik-Nezer, Rotem, Felix Holzmeister, Colin F Camerer, Anna Dreber, Juergen Huber, Magnus Johannesson, Michael Kirchler, et al. 2020. "Variability in the Analysis of a Single Neuroimaging Dataset by Many Teams." *Nature*, 1–7. `https://doi.org/10.1038/s41586-020-2314-9`.

Breheny, Patrick, Arnold Stromberg, and Joshua Lambert. 2018. "P-Value Histograms: Inference and Diagnostics." *High-Throughput* 7 (3): 23. `https://doi.org/10.3390/ht7030023`.

Bürkner, Paul-Christian. 2018. "Advanced Bayesian Multilevel Modeling with the R Package brms." *The R Journal* 10 (1): 395–411. `https://doi.org/10.32614/RJ-2018-017`.

Button, Katherine S, John PA Ioannidis, Claire Mokrysz, Brian A Nosek, Jonathan Flint, Emma SJ Robinson, and Marcus R Munafò. 2013. "Power Failure: Why Small Sample Size Undermines the Reliability of Neuroscience." *Nature Reviews Neuroscience* 14 (5): 365–76. `https://doi.org/10.1038/nrn3475`.

Carp, Joshua. 2012a. "On the Plurality of (Methodological) Worlds: Estimating the Analytic Flexibility of Fmri Experiments." *Frontiers in Neuroscience* 6: 149. `https://doi.org/10.3389/fnins.2012.00149`.

———. 2012b. "The Secret Lives of Experiments: Methods Reporting in the fMRI Literature." *Neuroimage* 63 (1): 289–300. `https://doi.org/10.1016/j.neuroimage.2012.07.004`.

Chen, Kaifu, Zheng Hu, Zheng Xia, Dongyu Zhao, Wei Li, and Jessica K Tyler. 2016. "The Overlooked Fact: Fundamental Need for Spike-in Control for Virtually All Genome-Wide Analyses." *Molecular and Cellular Biology* 36 (5): 662–67. `https://doi.org/10.1128/MCB.00970-14`.

Cokelaer, Thomas, Elisabeth Chen, Francesco Iorio, Michael P Menden, Howard Lightfoot, Julio Saez-Rodriguez, and Mathew J Garnett. 2017. "GDSCTools for mining pharmacogenomic interactions in cancer." *Bioinformatics* 34 (7): 1226–8. `https://doi.org/10.1093/bioinformatics/btx744`.

Everaert, Celine, Manuel Luypaert, Jesper LV Maag, Quek Xiu Cheng, Marcel E Dinger, Jan Hellemans, and Pieter Mestdagh. 2017. "Benchmarking of Rna-Sequencing Analysis Workflows Using Whole-Transcriptome Rt-qPCR Expression Data." *Scientific Reports* 7 (1): 1–11. `https://doi.org/10.1038/s41598-017-01617-3`.

Frazee, Alyssa C., Andrew E. Jaffe, Ben Langmead, and Jeffrey T. Leek. 2015. "Polyester: simulating RNA-seq datasets with differential transcript expression." *Bioinformatics* 31 (17): 2778–84. `https://doi.org/10.1093/bioinformatics/btv272`.

Goodman, S. N., D. Fanelli, and J. P. Ioannidis. 2016. "What does research reproducibility mean?" *Sci Transl Med* 8 (341): 341ps12. `https://doi.org/10.1126/scitranslmed.aaf5027`.

Grimes, D. R., C. T. Bauch, and J. P. A. Ioannidis. 2018. "Modelling science trustworthiness under publish or perish pressure." *R Soc Open Sci* 5 (1): 171511. `https://doi.org/10.1098/rsos.171511`.

Hardwicke, Tom E, Stylianos Serghiou, Perrine Janiaud, Valentin Danchev, Sophia Crüwell, Steven N Goodman, and John PA Ioannidis. 2020. "Calibrating the Scientific Ecosystem Through Meta-Research." *Annual Review of Statistics and Its Application* 7: 11–37. `https://doi.org/10.1146/annurev-statistics-031219-041104`.

Harris, Richard. 2017. *Rigor Mortis: How Sloppy Science Creates Worthless Cures, Crushes Hope, and Wastes Billions*. Basic Books.

Hu, Zheng, Kaifu Chen, Zheng Xia, Myrriah Chavez, Sangita Pal, Ja-Hwan Seol, Chin-Chuan Chen, Wei Li, and Jessica K Tyler. 2014. "Nucleosome Loss Leads to Global Transcriptional up-Regulation and Genomic Instability During Yeast Aging." *Genes & Development* 28 (4): 396–408. https://doi.org/10.1101/gad.233221.113.

Ioannidis, John PA. 2005. "Why Most Published Research Findings Are False." *PLoS Medicine* 2 (8): e124. https://doi.org/10.1371/journal.pmed.0020124.

Köster, Johannes, and Sven Rahmann. 2012. "Snakemake—a Scalable Bioinformatics Workflow Engine." *Bioinformatics* 28 (19): 2520–2. https://doi.org/10.1093/bioinformatics/bts480.

Leng, Gareth, and Rhodri Ivor Leng. 2020. *The Matter of Facts: Skepticism, Persuasion, and Evidence in Science*. Mit Press.

Lin, Charles Y, Jakob Lovén, Peter B Rahl, Ronald M Paranal, Christopher B Burge, James E Bradner, Tong Ihn Lee, and Richard A Young. 2012. "Transcriptional Amplification in Tumor Cells with Elevated c-Myc." *Cell* 151 (1): 56–67. https://doi.org/10.1016/j.cell.2012.08.026.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold Change and Dispersion for Rna-Seq Data with Deseq2." *Genome Biology* 15 (12): 550. https://doi.org/10.1186/s13059-014-0550-8.

Lovén, Jakob, David A Orlando, Alla A Sigova, Charles Y Lin, Peter B Rahl, Christopher B Burge, David L Levens, Tong Ihn Lee, and Richard A Young. 2012. "Revisiting Global Gene Expression Analysis." *Cell* 151 (3): 476–82. https://doi.org/10.1016/j.cell.2012.10.012.

Lun, Aaron TL, Fernando J Calero-Nieto, Liora Haim-Vilmovsky, Berthold Göttgens, and John C Marioni. 2017. "Assessing the Reliability of Spike-in Normalization for Analyses of Single-Cell Rna Sequencing Data." *Genome Research* 27 (11): 1795–1806. https://doi.org/10.1101/gr.222877.117.

Maiväli, Ülo. 2015. *Interpreting Biomedical Science: Experiment, Evidence, and Belief*. Academic Press.

Mandelboum, Shir, Zohar Manber, Orna Elroy-Stein, and Ran Elkon. 2019. "Recurrent Functional Misinterpretation of Rna-Seq Data Caused by Sample-Specific Gene Length Bias." *PLoS Biology* 17 (11): e3000481. https://doi.org/10.1371/journal.pbio.3000481.

McCarthy, Davis J, Yunshun Chen, and Gordon K Smyth. 2012. "Differential Expression Analysis of Multifactor Rna-Seq Experiments with Respect to Biological Variation." *Nucleic Acids Research* 40 (10): 4288–97. https://doi.org/10.1093/nar/gks042.

McGee, Warren A, Harold Pimentel, Lior Pachter, and Jane Y Wu. 2019. "Compositional Data Analysis Is Necessary for Simulating and Analyzing Rna-Seq Data." *bioRxiv*. https://doi.org/10.1101/564955.

Munafò, Marcus R, Brian A Nosek, Dorothy VM Bishop, Katherine S Button, Christopher D Chambers, Nathalie Percie Du Sert, Uri Simonsohn, Eric-Jan Wagenmakers, Jennifer J Ware, and John PA Ioannidis. 2017. "A Manifesto for Reproducible Science." *Nature Human Behaviour* 1 (1): 1–9. https://doi.org/10.1038/s41562-016-0021.

Nie, Zuqin, Gangqing Hu, Gang Wei, Kairong Cui, Arito Yamane, Wolfgang Resch, Ruoning Wang, et al. 2012. "C-Myc Is a Universal Amplifier of Expressed Genes in Lymphocytes and Embryonic Stem Cells." *Cell* 151 (1): 68–79. https://doi.org/10.1016/j.cell.2012.08.033.

Nookaew, Intawat, Marta Papini, Natapol Pornputtapong, Gionata Scalcinati, Linn Fagerberg, Matthias Uhlén, and Jens Nielsen. 2012. "A Comprehensive Comparison of Rna-Seq-Based Transcriptome Analysis from Reads to Differential Gene Expression and Cross-Comparison with Microarrays: A Case Study in Saccharomyces Cerevisiae." *Nucleic Acids Research* 40 (20): 10084–97. https://doi.org/10.1093/nar/gks804.

Peng, Roger D. 2011. "Reproducible Research in Computational Science." *Science* 334 (6060): 1226–7. https://doi.org/10.1126/science.1213847.

Prinz, Florian, Thomas Schlange, and Khusru Asadullah. 2011. "Believe It or Not: How Much Can We Rely on Published Data on Potential Drug Targets?" *Nature Reviews Drug Discovery* 10 (9): 712–12. https://doi.org/10.1038/nrd3439-c1.

Quinn, Thomas P, Ionas Erb, Greg Gloor, Cedric Notredame, Mark F Richardson, and Tamsyn M Crowley. 2019. "A Field Guide for the Compositional Analysis of Any-Omics Data." *GigaScience* 8 (9): giz107. https://doi.org/10.1093/gigascience/giz107.

Risso, Davide, John Ngai, Terence P Speed, and Sandrine Dudoit. 2014. "The Role of Spike-in Standards in the Normalization of Rna-Seq." In *Statistical Analysis of Next Generation Sequencing Data*, 169–90. Springer.

Ritchie, Matthew E, Belinda Phipson, Di Wu, Yifang Hu, Charity W Law, Wei Shi, and Gordon K Smyth. 2015a. "limma Powers Differential Expression Analyses for RNA-Sequencing and Microarray Studies." *Nucleic Acids Research* 43 (7): e47. `https://doi.org/10.1093/nar/gkv007`.

Ritchie, Matthew E, Belinda Phipson, DI Wu, Yifang Hu, Charity W Law, Wei Shi, and Gordon K Smyth. 2015b. "Limma Powers Differential Expression Analyses for Rna-Sequencing and Microarray Studies." *Nucleic Acids Research* 43 (7): e47–e47. `https://doi.org/10.1093/nar/gkv007`.

Storey, John D. 2002. "A Direct Approach to False Discovery Rates." *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 64 (3): 479–98. `https://doi.org/10.1111/1467-9868.00346`.

Sun, Zhaonan, and Yu Zhu. 2012. "Systematic Comparison of Rna-Seq Normalization Methods Using Measurement Error Models." *Bioinformatics* 28 (20): 2584–91. `https://doi.org/10.1093/bioinformatics/bts497`.

Szucs, Denes, and John Ioannidis. 2017. "When Null Hypothesis Significance Testing Is Unsuitable for Research: A Reassessment." *Frontiers in Human Neuroscience* 11: 390. `https://doi.org/10.3389/fnhum.2017.00390`.

Trapnell, C., D. G. Hendrickson, M. Sauvageau, L. Goff, J. L. Rinn, and L. Pachter. 2013. "Differential analysis of gene regulation at transcript resolution with RNA-seq." *Nat Biotechnol* 31 (1): 46–53. `https://doi.org/10.1038/nbt.2450`.

Wickham, Hadley. 2016. *Ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. `https://ggplot2.tidyverse.org`.

Wickham, Hadley, Mara Averick, Jennifer Bryan, Winston Chang, Lucy D'Agostino McGowan, Romain François, Garrett Grolemund, et al. 2019. "Welcome to the tidyverse." *Journal of Open Source Software* 4 (43): 1686. `https://doi.org/10.21105/joss.01686`.

Figure supplements

Figure 2–figure supplement 1. Simulated RNA-seq data shows that histograms from p value sets with around one hundred true effects out of 20,000 features can be classified as "uniform". RNA-seq data was simulated with polyester R package (Frazee et al. 2015) on 20,000 transcripts from human transcriptome using grid of 3, 6, and 10 replicates and 100, 200, 400, and 800 effects for two groups. Fold changes were set to 0.5 and 2. Differential expression was assessed using DESeq2 R package (Love, Huber, and Anders 2014) using default settings and group 1 versus group 2 contrast. Effects denotes in facet labels the number of true effects and N denotes number of replicates. Red line denotes QC threshold used for dividing p histograms into discrete classes. Code and workflow used to run these simulations is available on Github: https://github.com/rstats-tartu/simulate-rnaseq. Raw data of the figure is available on Zenodo https://zenodo.org with doi: 10.5281/zenodo.4463803.

Frazee, Alyssa C., Andrew E. Jaffe, Ben Langmead, and Jeffrey T. Leek. 2015. "Polyester: simulating RNA-seq datasets with differential transcript expression." *Bioinformatics* 31 (17): 2778–84. https://doi.org/10.1093/bioinformatics/btv272.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold Change and Dispersion for Rna-Seq Data with Deseq2." *Genome Biology* 15 (12): 550. https://doi.org/10.1186/s13059-014-0550-8.

Figure 3–figure supplement 1. The increasing proportion of anti-conservative histograms. Binomial logistic model: *anticons ~ year*, N = 2,109. Lines denote best fit of linear model. Shaded area denotes 95% credible region.

Figure 3–figure supplement 2. A 2-level binomial logistic model *anticons ~ year + (year | de_tool)* reveals that all differential expression analysis tools are associated with temporally increasing anti-conservative p value histograms, N = 2,109. Lines denote best fit of linear model. Shaded area denotes 95% credible region.

Figure 3–figure supplement 3. A 2-level binomial logistic model *anticons ~ year + (year | model)* reveals that all sequencing instrument models are associated with temporally increasing anti-conservative p value histograms, N = 1,718. Only GEO submissions utilizing single sequencing platform were used for model fitting. Lines denote best fit of linear model. Shaded area denotes 95% credible region.

Figure 3–figure supplement 4. No single differential expression analysis tool dominates the field. Y-axis shows the proportion of analysis platforms, x-axis shows publication year of GEO submission, N = 1,733.

Figure 3–figure supplement 5. Binomial logistic models for proportion of anti-conservative p value histograms. A, simple model *anticons ~ de_tool*, N = 2,109. B, simple model *anticons ~ de_tool* fitted on complete data, N = 6,267. C, model conditioned on year of GEO submission: *anticons ~ year + de_tool*, N = 2,109. D, model conditioned on studied organism (human/mouse/other): *anticons ~ organism + de_tool*, N = 1,733. E, varying intercept model *anticons ~ de_tool + (1 | model)* where "model" stands for sequencing instrument model, N = 1,718. F, varying intercept and slope model *anticons ~ de_tool + (de_tool | model)*, N = 1,718. Points denote best fit of linear model. Error bars, 95% credible interval.

Figure 4–figure supplement 1. Robust (student's t likelihood) modeling of π_0. A, simple model *pi0 ~ de_tool* fitted on complete data, N = 1,567. B, model conditioned on year of GEO submission: *pi0 ~ year + de_tool*, N = 488. C, model conditioned on studied organism (human/mouse/other): *pi0 ~ organism + de_tool*, N = 400. D, varying intercept model *pi0 ~ de_tool + (1 | model)* where 'model' stands for sequencing instrument model, N = 396. E, varying intercept/slope model *pi0 ~ de_tool + (de_tool | model)*, N = 396. Points denote best fit of linear model. Error bars denote 95% credible interval.

Figure 4–figure supplement 2. Modeling dependency of π_0 on sequencing instrument model: *pi0 ~ (1 | model)*, N = 396. Points denote best fit of linear model. Error bars denote 95% credible interval.

Figure 4–figure supplement 3. Modeling dependency of π_0 on library strategy: *pi0 ~ (1 | library_strategy)*, N = 396. Points denote best fit of linear model. Error bars denote 95% credible interval.

Figure 4–figure supplement 4. Modeling dependency of π_0 on library selection: *pi0 ~ (1 | library_selection)*, N = 396. Points denote best fit of linear model. Error bars denote 95% credible interval.

Figure 4–figure supplement 5. Modeling dependency of π_0 on library layout: *pi0 ~ (1 | library_layout)*, N = 396. Points denote best fit of linear model. Error bars denote 95% credible interval.

Figure 3–figure supplement 6. Modeling dependency of proportion of anti-conservative histograms on sequencing platform: *anticons ~ (1 | model)*, N = 1,718. Points denote best fit of linear model. Error bars denote 95% credible interval.

Figure 3–figure supplement 7. Modeling dependency of proportion of anti-conservative histograms on library strategy: *anticons ~ (1 | library_strategy)*, N = 1,718. Points denote best fit of linear model. Error bars denote 95% credible interval.

Figure 3–figure supplement 8. Modeling dependency of proportion of anti-conservative histograms on library selection: *anticons ~ (1 | library_selection)*, N = 1,718. Points denote best fit of linear model. Error bars denote 95% credible interval.

Figure 3–figure supplement 9. Modeling dependency of proportion of anti-conservative histograms on library layout: *anticons ~ (1 | library_layout)*, N = 1,718. Points denote best fit of linear model. Error bars denote 95% credible interval.