Extreme parallel evolution of flagellar motility is facilitated by synonymous sequence

Authors: James S. Horton[1,*], Robert W. Jackson[2], Nicholas K. Priest[1,a], and Tiffany B. Taylor[1,*,a]

[1]Milner Centre for Evolution, Department of Biology & Biochemistry, University of Bath, Claverton Down, Bath BA2 7AY,

UK

[2]School of Biosciences and Birmingham Institute of Forest Research (BIFoR), University of Birmingham, Edgbaston,

Birmingham, B15 2TT, UK

*Corresponding authors: j.s.horton@bath.ac.uk; t.b.taylor@bath.ac.uk

[a] These authors share senior authorship

Keywords: Parallel evolution, Predicting evolution, Synonymous sequence, Mutation rate

heterogeneity, Experimental evolution

<u>Abstract</u>

In recent years increasing interest has been shown in 'forecasting' evolution. Parallel evolution studies

are crucial towards realising this goal because they are, by their very nature, easy to forecast. However,

the causal role of identified drivers of parallel evolution – competitive selective advantage, mutational

accessibility or heterogeneity in *de novo* mutation – remain unresolved. Here we show that synonymous

sequences facilitates extreme parallel evolution during the evolutionary rescue of flagellar motility. An

immotile variant of the soil microbe, *Pseudomonas fluorescens*, swiftly recovers flagellum-dependent

motility through parallel *de novo* mutation. This typically manifests within 96 h under strong selection

through repeatable mutation within the nitrogen pathway's histidine kinase gene, *ntrB*. We found that

mutation was often parallel to nucleotide resolution, with lineages fixing the same mutation (*ntrB*

A289C) in over 95% of cases in minimal medium (M9). This repeatable *de novo* mutation was robust

to nutrient condition despite evidence for antagonistic pleiotropy across nutrient regimes. Competition

assays against alternative motile alleles revealed some evidence for selection enforcing repeated

fixation of *ntrB* mutants, but there was no evidence for clonal interference driving parallel evolution to

nucleotide resolution. Instead, we discovered evidence for extremely localised heterogeneity in *de novo*

mutation. The introduction of 6 synonymous substitutions surrounding the mutational hotspot reduced

parallel evolution from >95% to 0% at the site. Our results reveal that unique quirks in how DNA is

structured at specific loci can strongly bias evolutionary outcomes and ensure extreme repeatability.

Introduction

Evolution is sometimes remarkably repeatable, but the determining factors underlying extreme parallel evolution events are not well understood. Acquiring a better understanding of how certain populations are constrained in the ways they can evolve could prove especially insightful, as it would improve our ability for 'evolutionary forecasting'. Over the past decade there has been growing interest in the idea of predicting evolution (for review see, Lässig, Mustonen and Walczak, 2017). Parallel evolutionary studies, where independent lines evolve in identical ways when placed under selection for the same trait, provide examples where evolutionary forces are sufficiently powerful to exclude all but one mutational path. Such studies therefore illuminate the key evolutionary mechanisms which must be understood if we are ever to form a generalisable and accurate predictive model for adaptive evolution.

There have been many examples of experimental systems evolving in parallel. Microbes evolving under strong selection often rapidly adopt similar novel phenotypes (Fong et al. 2005; Ostrowski et al. 2008). More interestingly, however, is the observation that these phenotypes are often underpinned by clustered genetic changes within the same region of the genome (Riehle et al. 2001; Fraebel et al. 2017), or within limited pockets of loci (Bull et al. 1997; Wichman et al. 1999; Herron and Doebeli 2013; Kram et al. 2017). Sometimes viable mutations are limited to genes that comprise a single regulatory pathway (Notley-McRobb and Ferenci 1999; Miller et al. 2013) or a single protein complex (Avrani et al. 2017). In extreme cases, evolutionary events can be seen to repeatedly target just a handful of sites within a single locus (Meyer *et al.*, 2012; van Ditmarsch *et al.*, 2013). Parallel genetic evolution typically becomes less common as the degree of parallelism descends from broader genomic regions to the nucleotide (Tenaillon et al. 2012; Bailey et al. 2015). However, despite frequent descriptions of parallel evolutionary events, a detailed understanding of the evolutionary forces driving their occurrence is often lacking.

There are three primary drivers of parallel evolution in experimental systems: (*i*) Fixation bias, which skews evolution toward mutations that enjoy a higher likelihood of dominating the population pool. Not all facilitators of fixation bias are considered adaptively advantageous (Eyre-Walker and Hurst 2001), but in instances where we observe rapid and highly parallel sweeps it will likely take the form of

selection, which drives the fittest competing genotypes in the population to fixation (see Wood, Burke

and Rieseberg, 2005; Woods *et al.*, 2006). (*ii*) Mutational accessibility, as there may be only a small

number of readily accessible mutations a genotype can undergo to improve fitness (Weinreich et al.

2006). And, (*iii*) Mutation rate heterogeneity, where genetic and molecular features scattered

throughout the genome cause sites to radiate at different rates, introducing a mutation bias toward a

particular outcome (Bailey et al. 2017). Previous research shows that mutation rate heterogeneity can

be influenced by the arrangement of nucleotides surrounding a particular site (Long et al. 2014), and

genetic quirks such as the secondary structure of DNA (Duan et al. 2018) including the formation of

single-stranded DNA hairpins (De Boer and Ripley 1984). Nevertheless, the nature of genetic sequence

in driving parallel evolutionary outcomes remains unknown.

To establish which mechanisms are at play, it is important to consider whether parallel outcomes are

robust to experimental conditions such as environment (Turner et al. 2018) and to account for clonal

interference, which can alter the chance of observing parallel evolution (Bailey et al. 2017; Lässig et al.

2017). Clonal interference can occur either due to standing genetic variation in the founder population

which yields multiple adaptive genotypes in a novel environment (i.e. a soft selective sweep; Hermisson

and Pennings, 2005) or when mutation rate is high relative to the selective coefficient (Barrett et al.

2006). However, clonal interference does not often play an important role when founding experimental

lines with clonal samples, performing experimental procedures over short timescales, and ensuring

rapid fixation of adaptive mutants e.g. through spatial separation and/or introducing an artificial

bottleneck.

The ideal system for evaluating evolutionary drivers of parallel evolution would involve a model system

that evolves in an extremely parallel manner i.e. reliably fixing the same single nucleotide

polymorphism across a range of environmental conditions and independent lines, while limiting clonal

interference. The merit of this approach is that it would allow to test whether an extremely parallel

evolutionary event is either a consequence of an extremely determinant evolutionary force, or a

combination of multiple harmonious factors. We have been able to address this challenge by employing

two engineered non-flagellate variants of the soil bacterium *Pseudomonas fluorescens* strain SBW25.

These variants lack the master regulator of flagella-dependent motility, FleQ (SBW25Δ*fleQ*). One

variant additionally has a transposon inserted within *viscB* that prevents production of the biosurfactant

viscosin, denying the bacteria an alternative form of sliding motility and rendering the variant

completely immotile (SBW25Δ*fleQ* IS-Ω-Km/hah: PFLU2552, hereafter AR2; Alsohim *et al.*, 2014).

AR2 has previously been shown to rapidly re-evolve flagella-mediated motility under strong directional

selection (Taylor et al. 2015). This phenotype was achieved in independent lineages via repeatable *de*

novo mutation in the *ntrB* locus of the nitrogen regulatory (ntr) pathway. Mutations in *ntrB* resulted in

the constitutive expression and subsequent over-activation of NtrC, the ntr pathway's response regulator

and FleQ homolog, allowing NtrC to act as a surrogate for the deleted flagella regulator (Taylor et al.

2015). The parallel evolution of *ntrB* mutants was noteworthy, as the locus was constantly targeted

during adaptation despite the observation of other mutational routes during the evolution of motility

within a closely related *P. fluorescens* strain, Pf0-1 (giving rise to slow moving mutants, collectively

dubbed Pf0-2xS). These mutations occurred within the ntr regulatory hierarchy which is shared by AR2

and Pf0-2x, suggesting other viable routes also existed for SBW25 derivatives (Taylor et al. 2015).

Here we show that not only does motility evolve in SBW25-derived strains in an extremely parallel

manner, this adaptation is reliant on a previously uncharacterised genetic quirk. The evolution of

flagella motility was found to target the same nucleotide substitution in over 95% of cases in minimal

media (M9). This outcome was found to be robust across multiple nutrient regimes both in the immotile

SBW25 variant (AR2) and another variant that was able to access biosurfactant-mediated motility prior

to evolution (Δ*fleQ*). The role of selection and the number of viable mutational routes in ensuring the

parallel outcome were found to provide some explanation for parallel evolution to the level of the *ntrB*

locus, but not the nucleotide. This implied that intra-locus mutation rate heterogeneity was playing a

critical role. We then genetically augmented the *ntrB* locus to indirectly incriminate mutation bias and

revealed a key underlying genetic driver of parallel evolution. Synonymous changes introduced within

the local region around the frequently targeted site were found to reduce parallel evolution at the

mutational hotspot from >95% to 0%. This result reveals that synonymous sequence can play a

dominant role in ensuring parallel evolutionary outcomes, and shines a spotlight on the overlooked

mechanistic drivers behind parallel evolutionary events.

Materials and Methods

Model System

Our model system employs strains of the soil microbe *Pseudomonas fluorescens* SBW25 that lack motility through partial gene deletion of *fleQ,* the master regulator of flagellar motility (Alsohim et al., 2014). Two ancestral strains were used in this study: SBW25 Δ*fleQ* (hereafter Δ*fleQ*) and a Δ*fleQ* variant with a functional *viscB* knockout isolated from a transposon library (SBW25Δ*fleQ* IS-ΩKm-hah: PFLU2552, hereafter AR2; Alsohim *et al.*, 2014). Δ*fleQ* can migrate on soft agar prior to mutation via a form of sliding motility, which is owed to the strain's ability to produce viscosin. AR2 cannot produce viscosin and is thus rendered completely immotile prior to mutation. All cells were grown at 27°C and all strains used throughout the study (ancestral, evolved and engineered) were stored at -80°C in 20% glycerol until required. The nutrient conditions used throughout the work were lysogeny broth (LB) and M9 minimal media containing glucose and 7.5 mM NH_4. The minimal media was used in isolation or supplemented with either glutamate (M9+glu) or glutamine (M9+gln) at a final supplement concentration of 8 mM unless stated otherwise.

Motility Selection Experiment

Immotile variants were placed under selection for flagella-mediated motility using LB and M9 soft agar motility plates (0.25%). Details of agar preparation are described in Alsohim et al., 2014. Supplemented concentrations of glutamate (glu)/glutamine (gln) in M9 soft agar were expanded to include final concentrations at 4 mM, 8 mM and 16 mM, as it was observed that biosurfactant-mediated dendritic motility in Δ*fleQ* lines was enhanced at higher supplement concentrations, which masked any emergent blebs (data not shown). Lowering the gln supplement concentration improved the likelihood of observing an emergent flagella bleb in M9+gln motility plates (16 mM: 4/12, 8 mM: 9/20, 4 mM: 7/12 independent lines). However, dendritic motility remained high on all supplements of M9+glu and persistently masked blebbing (16 mM: 2/12, 8 mM: 3/20, 4 mM: 2/11 independent lines). Although gln/glu supplementation had no bearing on motility in AR2 lines, supplement conditions across both gln/glu were expanded for consistency. Single clonal colonies were inoculated into the centre of the

agar using a sterile pipette tip and monitored daily until emergence of motile bleb zones (as visualised

in fig. 1A). Samples were isolated from the leading edge, selecting for the strongest motility phenotype

on the plate, within 24 h of emergence and streaked onto LB agar (1.5%) to obtain a clonal sample. As

Δ*fleQ* lines were motile via dendritic movement prior to re-evolving flagella motility and could visually

mask flagella-mediated motile zones, samples were left for 120 h prior to sampling from the leading

edge of the growth. An exception was made in instances where blebbing motile zones were observed

solely further within the growth area, in which case this area was preferentially sampled.

Sequencing

Motility-facilitating changes were determined through PCR amplification and sequencing of *ntrB, glnK*

and *glnA* genes (supplementary table S1). Polymerase chain reaction (PCR) products and plasmids were

purified using Monarch® PCR & DNA Cleanup Kit (New England Biolabs) and Sanger sequencing

was performed by Eurofins Genomics. A subset of AR2 samples evolved on different nutritional

backgrounds was additionally screened through Illumina Whole-Genome Sequencing by the Milner

Genomics Centre and MicrobesNG (LB: n = 5, M9: n = 6, M9+gln: n = 6, M9+glu: n = 7). This allowed

us to screen for potential secondary mutations and to identify rare changes in motile strains with

wildtype *ntrB* sequences. *P. fluorescens* SBW25 genome was used as an assembly template (NCBI

Assembly: ASM922v1, GenBank sequence: AM181176.4) and single nucleotide polymorphisms were

called using Snippy with default parameters (Seemann 2015) through the Cloud Infrastructure for

Microbial Bioinformatics (CLIMB; Connor *et al.*, 2016). In instances where coverage at the called site

was low (≤10x), called changes were confirmed by Sanger sequencing.

Assessing Pleiotropy via Growth Rate

Cryopreserved samples of AR2 and derived *ntrB* mutants were streaked and grown for 48 h on LB agar

(1.5%). Three colonies were then picked, inoculated in LB broth and grown overnight at an agitation of

180 rpm to create biological triplicates for each sample. Overnight cultures were pelleted via

centrifugation, their supernatant withdrawn and the cell pellets re-suspended in phosphate buffer saline

(PBS) to a final concentration of OD1 cells/ml. The cultures were corrected to OD1 by first determining

ODx cells/ml and adjusting the collected volume of culture so that $x = v$, with v being the volume (ml)

of PBS used to re-suspend the cell pellet. The resuspension was subsequently diluted 100-fold into a

96-well plate (Costar®) containing nutrient broth. The plates were analysed in a Multiskan™ FC

Microplate Photometer (Thermo Fisher Scientific) for 24h, with autonomous optical density readings

every 10 min without agitation. Growth values were determined by calculating area under the curve

using the trapezoidal rule (approached outlined in Huang and Pang, 2012). This allowed us to

incorporate all elements of the pleiotropic consequences to metabolism and the benefit of swimming

motility, including prolonged lag phases and differing eventual yields achieved by mutant populations

relative to the ancestral strain (growth curves not shown). This process was repeated with an

independent batch of biological triplicates to produce a total of 6 biological replicates for each sample.

Soft Agar Motility Assay

OD-corrected biological triplicates were prepared as outlined above, which allowed us to standardise

the starting population count for the assay. 1 µl of each replicate was inoculated into soft-agar by

piercing the top of the agar with the pipette tip and ejecting the culture into the cavity as the tip was

withdrawn. Plates were then incubated for 48 h and photographed. Diameters of concentric circle

growths were calculated laterally and longitudinally, allowing us to calculate an averaged total surface

area using A= πr^2. This process was repeated as several independent lines underwent a second-step

mutation (Taylor et al. 2015) within the 48 h assay. This phenotype was readily observable as a blebbing

that appeared at the leading edge along a segment of the circumference, distorting the expected

concentric circle of a clonal migrating population. As such these plates were discarded from the study.

By completing additional sets of biological triplicates, we ensured that each sample had at least three

biological replicates for analysis.

Invasion Assay

OD-corrected biological quadruplets of both *ntrB* mutant lines were prepared as outlined above. For

each pair of biological replicates, 1 µl of *ntrB* A683C was first inoculated as outlined above and

incubated, followed by *ntrB* A289C's inoculation into the same cavity after the allotted time had elapsed

(3 h, 6 h, 9 h and 12 h). When inoculated at 0 h, *ntrB* A289C was added to the plate immediately after

ntrB A683C. In instances where *ntrB* A289C was added to the plate ≤6 h after *ntrB* A683C, overgrowth

of culture was avoided by incubating *ntrB* A289C cultures at 22°C at 0 h until cell pelleting and re-

suspension approximately 1 h prior to inoculation. When *ntrB* A289C cultures were added to the plate

≥9 h after *ntrB*-A683C culture, overgrowth of culture was avoided by diluting the culture of *ntrB*-

A289C 100-fold into fresh LB broth at 0 h. The same 'angle of attack' was used for both instances of

inoculation (i.e. the side of the plate that the pipette tip travelled over on its way to the centre), as small

volumes of fluid falling from the tip onto the plate could cause local satellite growth. To avoid the risk

of satellite growths affecting results, isolated samples were collected from the leading edge 180° from

the angle of attack after a period of 24 h. The *ntrB* locus of one sample per replicate was determined by

Sanger sequencing to establish the dominant genotype at the growth frontier.

***ntrB* loci analysis**

Theoretical hairpin stem-loop structures were generated using the *mfg* tool and methodology developed

by Wright *et al.*, 2003. The *mfg* tool is used in conjunction with the Quikfold tool on the DINAMelt

Web Server (Markham and Zuker 2005). Default parameters were used for Quikfold with the exception

of temperature, which was amended to 27^0C. The first 400 nucleotides of the open reading frames of

P. fluorescens SBW25 *ntrB* and Pf0-1 *ntrB* were used as input sequences, and AR2-sm's input sequence

was created by manually editing SBW25's *ntrB* sequence. The *mfg* application generates the most stable

stem-loop structure for each base in which the selected base remains unpaired and so is at a higher

likelihood of mutation. The window size of neighbouring nucleotides that are used to form the stem-

loop structure can be adjusted, and a window length of 40 nucleotides was used for the analysis in this

study.

Genetic engineering

A pTS1 plasmid containing *ntrB* A683C was assembled using overlap extension PCR (oePCR) cloning

(for detailed protocol see Bryksin and Matsumura, 2010) using vector pTS1 as a template. The *ntrB*

synonymous mutant and synonymous mutant with A289C pTS1 plasmids were constructed using

oePCR to assemble the insert sequence for allelic exchange, followed by amplification using nested

primers and annealed into a pTS1 vector through restriction-ligation (for full primer list see

supplementary table. S1). pTS1 is a suicide vector, able to replicate in *E. coli* but not *Pseudomonas*,

and contains a tetracycline resistance cassette as well as an open reading frame encoding SacB. Cloned

plasmids were introduced to *P. fluorescens* SBW25 strains via puddle mating conjugation with an

auxotrophic *E. coli* donor strain ST18. Mutations were incorporated into the genome through two-step

allelic exchange, using a method outline by Hmelo *et al.*, 2015, with the following adjustments: (*i*) *P.*

fluorescens cells were grown at 27^0C. (*ii*) An additional passage step was introduced prior to

merodiploid selection, whereby colonies consisting of *P. fluorescens* cells that had incorporated the

plasmid (merodiploids) were allowed to grow overnight in LB broth free from selection, granting extra

generational time for expulsion of the plasmid from the genome. (*iii*) The overnight cultures were

subsequently serially diluted and spot plated onto NSLB agar + 15% (wt/vol) sucrose. Positive mutant

strains were identified through targeted Sanger sequencing of the *ntrB* locus.

Analysis of molecular data

All statistical tests and figures were produced in R (R Core Team 2014). Figures were created using the

ggplot package (Wickham 2016). A simulated dataset was produced for the Bootstrap test by randomly

drawing from a pool of 3 values with equal weights 24 times for 1 million iterations. Note that as the

simulated dataset draws from a pool of 3 values, it encodes that no other mutational routes are possible

aside from the observed 3. As such the derived statistic is an underestimate, with additional routes at

any weight lowering the likelihood of repeat observations of a single value. All other tests were

completed using functions in base-R aside from the Dunn test, which was performed using the *FSA*

package (Ogle et al. 2020).

Results

Remarkable Parallel Evolution

We evolved 24 independent replicates under strong directional selection in a minimal medium environment (M9) to quantify the degree of parallel evolution of flagellar motility within the immotile SBW25 model system (AR2). Motile mutants were readily identified through emergent motile zones that migrated outward in a concentric circle (fig. 1A). Clonal samples were isolated from the zone's leading edge within 24 h of emergence and their genotypes analysed through either whole-genome or targeted Sanger sequencing of the *ntrB* locus. Motile strains evolved rapidly (fig. 1B) and each independent line was found to be a product of a one-step *de novo* mutation. All 24 lines had evolved in parallel at the locus level: each had acquired a single, motility-restoring mutation within *ntrB* (fig. 1C). More surprising however, was the level of parallel evolution within the locus. 23/24 replicates had acquired a single nucleotide polymorphism at site 289, resulting in a transversion mutation from A to C (hereafter referred to as *ntrB* A289C). This resulted in a T97P missense mutation within NtrB's PAS domain. The remaining sample had acquired a 12-base-pair deletion from nucleotide sites 406-417 (Δ406-417), resulting in an in-frame deletion of residues 136-139 (ΔLVRG) within NtrB's phospho-acceptor domain.

Robust parallel evolution across environments

Parallel evolution could be robust or highly context-dependent, especially when it occurs via *de novo* mutations with antagonistic pleiotropic effects (Mcgrath et al. 2011; Mcgee et al. 2016; Sackman et al. 2017). However, we found that the repeatability of the *ntrB* A289C mutation was robust across all tested conditions, despite evidence of antagonistic pleiotropic effects on growth. We tested for environment-specific antagonistic pleiotropy by measuring relative growth of the ancestral line and both evolved *ntrB* mutants on rich lysogeny broth and minimal medium containing either ammonia as the sole nitrogen source or supplemented with either glutamate (M9+glu) or glutamine (M9+gln), both of which are naturally assimilated and metabolised by the ntr system. Though large fitness costs were evident in M9 minimal medium, supplementing M9 with glu or gln reduced levels of antagonistic pleiotropy for

both the *ntrB* A289C and the Δ406-417 mutants (supplementary fig. S1). Indeed, the antagonistic

pleiotropy of impaired metabolism was sufficiently low in M9 supplemented with the amino acid

glutamine (M9+gln) that motile mutants had increased fitness over the ancestral line in static broth,

which was significant in *ntrB* A289C (supplementary fig. S1). These findings show that antagonistic

pleiotropy has the potential to influence the robustness of parallel evolution.

We then tested whether antagonistic pleiotropy interferes with the robustness of parallel evolution in

our system. Our expectation was that supplemented nutrient regimes would lower pleiotropic costs and

thus unlock alternative routes of adaptation. We additionally hypothesised that the ability of the Δ*fleQ*

strain, which is able migrate prior to mutation (see materials and methods), would also ease starvation-

induced selection pressures and could facilitate yet more mutational routes. We observed a 'blebbing'

phenotype (fig. 1A) in Δ*fleQ* lines despite their ability to migrate in a dendritic fashion; however, we

also found blebbing was less frequent under richer nutrient regimes (where populations migrated more

rapidly utilising viscosin, see materials and methods). Overall, there was no evidence that competitive

ability of the *ntrB* A289C mutation changed with nutrient condition (Gene-by-environment interaction:

χ^2= 0.9375, df = 7, P = 0.9958, see fig. 2). Instead, we observed that the *ntrB* A289C mutation was

robust across all tested conditions, featuring in 90-100% of the Δ*fleQ* strains and 80-100% of AR2

strains (fig. 2).

Additionally, three novel mutational routes were observed in a small number of mutants (fig. 2),

revealing that mutational accessibility could not explain the level of observed parallel evolution. Most

notably was a non-synonymous A-C transversion mutation at site 683 (*ntrB* A683C) in a Δ*fleQ* line

evolved on M9+gln, resulting in a missense mutation within the NtrB histidine kinase domain. As a

single A-C transversion within the same locus, we may expect A683C to mutate at a similar rate to

A289C. We also observed a 12 base-pair deletion from sites 410-421 (*ntrB* Δ410-421) in an AR2 line

evolved on M9+gln. Furthermore, we discovered a double mutant in an AR2 line evolved on M9+glu:

one mutation was a single nucleotide deletion at site 84 within *glnK,* and the second was another A to

C transversion at site 688 resulting in a T230P missense mutation within RNA polymerase sigma factor

54. GlnK is NtrB's native regulatory binding partner and repressor in the ntr pathway, meaning the

frameshift mutation alone likely explains the observed motility phenotype. However, as this mutant

underwent two independent mutations we will not consider it for the following analysis. In addition,

ntrB Δ410-421 and *ntrB* Δ406-417, despite targeting different nucleotides, translate into identical

protein products (both compress residues LVRGL at positions 136-140 to a single L at position 136).

Therefore, we will also group them for the following analysis. Under the assumptions that the three

remaining one-step observed mutational routes to novel proteins are (*i*) equally likely to appear in the

population and (*ii*) equally likely to reach fixation, the original observation of *ntrB* A289C appearing

in 23/24 cases becomes exceptional (Bootstrap test: n = 1000000, $P < 1 \times 10^{-6}$). The likelihood of our

observing this by chance, therefore, is highly unlikely. This means that one or both assumptions are

almost certainly incorrect. Either the motility phenotype facilitated by the mutations may be unequal,

leading to fixation bias. Or the mutations may appear in the population at different rates, resulting in

mutation bias. One or both of these elements must be skewing evolution to such a degree that parallel

evolution to nucleotide resolution becomes highly predictable.

Assessing fixation bias

The Darwinian explanation for parallel evolution is that the observed mutational path is outcompeting

all others on their way to fixation. If selection alone was driving parallel evolution, the superior fitness

of the *ntrB* A289C genotype should have allowed it to out-migrate other motile genotypes co-existing

in the population. To test if the *ntrB* A289C mutation granted the fittest motility phenotype, we allowed

the evolved genotypes (A289C, Δ406-417, A683C and *glnK* Δ84) to migrate independently on the four

nutritional backgrounds and measured their migration area after 48 h. To allow direct comparison, we

first engineered the *ntrB* A683C mutation, which originally evolved in the Δ*fleQ* background, into an

AR2 strain. We observed that the non-*ntrB* double mutant, *glnK* Δ84, migrated significantly more

slowly than *ntrB* A289C in all four nutrient backgrounds (fig. 3A). However, *ntrB* A289C did not

significantly outperform either of the alternative *ntrB* mutant lines in any environmental condition (fig.

3A). This suggests that selection may have played a role in driving parallel evolution to the level of the

ntrB locus, but it cannot explain why nucleotide site 289 was so frequently radiated.

To determine if this result remained true when mutant lines were competing in the same population, we

directly competed *ntrB* A289C against *ntrB* A683C on M9 minimal medium. In brief, we co-inoculated

the two mutant lines on the same soft agar surface and allowed them to competitively migrate before

sampling from the leading edge after 24 h of competition. The length of competition was maintained

throughout the assay, but *ntrB* A683C lines were allowed to migrate for between 0 and 12 h before the

addition of *ntrB* A289C to the agar. We observed that *ntrB* A289C was found predominantly on the

leading edge (3/4 replicates) when the mutants were inoculated concurrently, but invading populations

of the common genotype swiftly became unable to establish themselves at the leading edge within a

narrow time window of 3 h (fig. 3B). This result highlights that in minimal medium *ntrB* A289C does

offer a slight dominant phenotype, but to ensure establishment at the leading edge the genotype would

need to appear in the population within a handful of generations of a competitor. Given that the range

in time before a motility phenotype was observed could vary considerably between independent lines

(fig. 1B), our data do not support the hypothesis that global mutation rate could be high enough to allow

multiple phenotype-granting mutations to appear in the population almost simultaneously. More likely

is that each independent line adhered to the "early bird gets the word" maxim, i.e. the *ntrB* mutant which

was the first to appear in the population was the genotype that reached fixation. This therefore suggests

that the reason *ntrB* A289C is so frequently collected when sampling is due to an evolutionary force

other than selection and mutational accessibility.

Identifying mutation bias

Local mutational biases can play a key role in evolution (Bailey et al. 2017; Lind et al. 2019). Such

biases can be introduced by changing DNA curvature (Duan et al. 2018) or through neighbouring tracts

of reverse-complement repeats (palindromes and quasi-palindromes), which have been shown to invoke

local mutation biases by facilitating the formation of single-stranded DNA hairpins (De Boer and Ripley

1984). Therefore we next searched for a local mutation bias at *ntrB* site 289. Previously, we re-evolved

motility in two engineered immotile strains of *Pseudomonas fluorescens*, SBW25 and Pf0-1 (Taylor et

al. 2015). Although evolved lines in SBW25 (namely AR2) frequently targeted *ntrB*, Pf0-1 lines (Pf0-

2xS) fixed mutations across the ntr pathway. Furthermore, although Pf0-2x did acquire *ntrB* mutations

in multiple independent lines, we observed no evidence of *ntrB* site 289 being targeted in Pf0-2xS

(Taylor et al. 2015). The NtrB proteins of SBW25 and Pf0-1 are highly homologous (95.57% identity)

but share less identity at the genetic level (88.88% identity). A considerable portion of this genetic

variation is explained by synonymous genetic variation (8.34%) rather than non-synonymous variance

(2.76%). Synonymous mutations can play a role in altering local mutation rates. This may occur by

altering the nucleotide-triplet to one with a higher mutation rate (Long et al. 2014) or by altering the

secondary structure of longer DNA tracts via the mechanisms outlined above. Nucleotides that remain

unpaired when their neighbouring nucleotides form hairpins with nearby reverse-complement tracts

have been observed to exhibit increased mutation rates (Wright et al. 2003). Both SBW25 and Pf0-1

were found to have short reverse-complement tracts that flanked site 289, however the called hairpins

were not entirely identical in their composition owing to synonymous variance (supplementary fig. S2).

Overall, there are 6 synonymous nucleotide substitutions ± 5 codons of site 289 (C276G, C279T,

C285G, C291G, T294G and G300C), which may have been affecting such hairpin formations and

impact local mutation rate.

To test if synonymous sequence was biasing evolutionary outcomes, we replaced the 6 synonymous

sites in an AR2 strain with those from a Pf0-1 background (hereafter AR2-sm). Not all these sites

formed part of a theoretically predicted stem that overlapped with site 289, but all were targeted due to

their close proximity with the site. AR2-sm lines were placed under selection for motility and we

observed that these lines evolved motility significantly more slowly (fig. 4A), both in M9 minimal

medium and LB (Wilcoxon rank sum tests with continuity correction: M9, $W = 44.5$, $P < 0.001$; LB,

$W = 22$, $P < 0.001$). Evolved AR2-sm lines that re-evolved motility within 8 days were sampled and

their *ntrB* locus analysed by Sanger sequencing (fig. 4B). We observed some similar *ntrB* mutations to

those identified previously: the *ntrB* A683C mutation was observed in one independent line evolved on

LB, and *ntrB* Δ406-417 was also observed in both strain backgrounds. However, the most common

genotype of *ntrB* A289C fell from being observed in over 95% of independent lines to 0. Furthermore,

we observed multiple previously unseen *ntrB* mutations, while a considerable number of lines reported

wildtype *ntrB* sequences, instead either targeting another gene of the ntr pathway (*glnK*) or unidentified

targets that may lay outside of the network (fig. 4B).

To test that the A289C transition remained a viable mutational target in the AR2-sm genetic

background, we subsequently engineered the AR2-sm strain with this additional mutation. We observed

that AR2-sm *ntrB* A289C was both motile and comparable in phenotype to *ntrB* A289C in the ancestral

AR2 genetic background (supplementary fig. S3). We additionally found that AR2-sm *ntrB* A289C

retained comparable motility to the other *ntrB* mutants evolved from AR2-sm (supplementary fig. S3).

Therefore, we can determine that the AR2-sm genetic background would not prevent motility following

mutation at *ntrB* site 289, nor does it render such a mutation uncompetitive. This therefore infers that

the sole variable being altered between strains (the 6 synonymous changes) are precluding radiation at

site 289. Taken together these results strongly suggest that the synonymous sequence immediately

surrounding *ntrB* site 289 facilitates its position as a local mutational hotspot, and that local mutational

bias is imperative for realising extreme parallel evolution in our model system.

Discussion

Understanding why certain mutations reach fixation in adapting populations remains an immense challenge. This is true even in simple systems such as the one employed in this study, where clonal bacterial populations were evolved under strong directional selection for very few phenotypes, namely motility and nitrogen metabolism. Here we took immotile variants of *Pseudomonas fluorescens* SBW25 that had been observed to repeatedly target the same locus (*ntrB*) during the re-evolution of motility (Taylor et al. 2015), and found that evolving populations adapted via *de novo* substitution mutation at the same nucleotide site in over 95% of cases (A289C) in minimal medium. We found that populations were somewhat limited in which genetic avenues they could take to access the phenotype under selection, but fitness differences alone could not explain the high degree of parallel evolution. Instead we observed that introducing synonymous changes around the mutational hotspot pushed evolving populations away from the parallel path, revealing that synonymous sequence is an integral factor toward realising extreme parallel evolution in our system.

Models looking to describe adaptive evolution often place precedence on fitness and the number of accessible adaptive routes (Orr 2005; Krug 2019) yet pay little attention to local mutational biases (however, see Sackman *et al.*, 2017). However, mutation rate heterogeneity becomes of paramount importance when systems adhere to the Strong Selection Weak Mutation model (SSWM), which describes instances when an advantageous mutation undergoes a hard sweep to fixation before another beneficial mutation appears (Gillespie 1984). In such cases relative fitness values between adaptive genotypes are relegated to secondary importance behind the likelihood of a genotype appearing in the population. Indeed, experimental systems that adhere to the SSWM maxim have been observed to evolve in parallel despite the option of multiple mutational routes to improved fitness (Vogwill et al. 2014). This suggests that uneven mutational biases can be a key driver in realising parallel evolution, a conclusion which has been reinforced theoretically (Bailey et al. 2017) although empirical data is still lacking. Understanding the mechanistic causes of mutation rate heterogeneity, therefore, will be essential if we are to determine when and if accurate predictions of evolution are possible (Bailey et al.

2018; Lind et al. 2019). The challenge remains in identifying what these mechanistic quirks may be,

and where they may be found.

Synonymous changes are often treated as adaptively neutral, offering no discernible changes to fitness

post-translation. However, more recent studies have revealed that synonymous changes have an

underestimated effect on fitness through their perturbances before and during translation. Synonymous

sequence variance can impact fitness by changing the stability of mRNA (Kudla et al. 2009; Kristofich

et al. 2018; Lebeuf-Taylor et al. 2019) and altering codons to perturb or better match the codon-

anticodon ratio (Frumkin et al. 2018). To our knowledge, we have shown here for the first time that

synonymous sequence can also be essential for realising extreme parallel evolution. We believe this is

due to its impact on local mutational biases, which mechanistically may be owed to the formation of

single-stranded hairpins that form between short inverted repeats (De Boer and Ripley 1984; Fieldhouse

and Golding 1991). The formation of these secondary DNA structures provides a mechanism for intra-

locus mutation bias that can operate with extremely local impact and is contingent on DNA sequence

variation, as introducing synonymous changes could readily perturb the complementarity of

neighbouring inverse repeats (e.g. supplementary fig. S2). Furthermore, the finding of just six

synonymous mutations having a significant impact on DNA structure would not represent a surprising

result, as secondary structures can be altered by single mutations (Dong et al. 2001).

We have further shown that the frequently observed *ntrB* A289C mutation remains both a viable and

competitive mutational route within the synonymous mutant background (supplementary fig. S3). As

the competitive fitness and broad genetic background remains the same, the six synonymous mutations

are the sole variable between the two AR2 strains (AR2 and AR2-sm), and therefore are solely

responsible for both the delayed evolution and the loss of a mutational hotspot.

One caveat of this finding is that local mutation bias occurring via single stranded hairpins is only

indirectly inferred by our results. We can confidently assert, however, that the altered mutational bias

is owed to an intralocus effect, owing to the six synonymous sites all residing within 14 bases at either

flank of site 289. Interestingly, our data suggest that the mutational hotspot typically mutates so quickly

as to mask mutations appearing elsewhere and outside of the nitrogen regulatory pathway, which only

appear when the hotspot is perturbed (fig. 4B). However, the exact degree of bias owed to synonymous

sequence variation remains an open question.

One unexpected result was that parallel evolution was robust to changes in nutrient regime and

starvation-induced selection. We expected the selective environments to hold some influence over

evolutionary outcomes (Bailey et al. 2015) mostly owing to varying levels of antagonistic pleiotropy,

which has been found to be a key driver in similar motility studies (Fraebel et al. 2017). However, we

have not yet assessed robustness across broad genetic backgrounds. We have previously highlighted

that two immotile strains of *P. fluorescens* (AR2 and Pf0-2x) evolve in parallel only so far as the

targeted regulatory pathway (Taylor et al. 2015). However, what is less clear is how much of the

observed divergence between the two strains is owed to synonymous sequence variation within the *ntrB*

locus, and the resultant mutational hotspot, and how much to genetic variation throughout the rest of

the genome. Enquiries into this area may prove especially insightful, as the evolutionary hotspot-lacking

strain (AR2-sm) more closely aligns to the Pf0-1 variant's mutational trajectory than AR2 (fig. 4B;

Taylor *et al.*, 2015). Parallel evolution can sometimes be impressively robust across genetic

backgrounds (Vogwill et al. 2014) but some innovations can be strongly determined by an organism's

evolutionary history (Blount et al. 2012). In addition, genomic variation typically combines with

environmental differences to drive populations down diverse paths (Spor et al. 2014), so robustness

may be hampered in a novel genetic background. Determining whether genetic background or the

presence of a mutational hotspot is the stronger deterministic evolutionary force in our model system

awaits further study.

The ultimate end goal of predictive evolution is to make an accurate evolutionary forecast prior to

performing the experiment and without any foreknowledge of the expected outcome. This is an

ambitious task because mutation is inherently a random process, but not all sites in the genome possess

equal fixation potential. Most changes will not improve a phenotype under selection, and those that do

will not necessarily mutate at the same rates. Therefore, we can increase our predictive power

dramatically, permitting we have a detailed understanding of the evolutionary variables at play.

Considerable inroads have already been made toward realising this goal. It has been highlighted that

loss-of-function mutations are the most frequently observed mutational type under selection (Kimura,

1968; Lind, Farr and Rainey, 2015) and that a gene's wider position within its regulatory network

determines its propensity in delivering phenotypic change (McDonald et al. 2009). It has also been

shown that parallel evolution at the level of the locus is partially determined by gene length (Bailey et

al. 2018) and that global mutation bias can strongly influence the likelihood of a given nucleotide

substitution (Lind and Andersson 2008; Stoltzfus and McCandlish 2017). Here, we show that

synonymous sequence warrants consideration alongside these other variables by highlighting its impact

on the realisation of extreme parallel evolution.

Acknowledgments and funding information

The authors would like to thank Laurence Hurst for comments on earlier versions of this manuscript.

In addition we thank members of the Taylor lab, Matthew Shepherd and Louise Flanagan, for insightful

comments and discussion. This work was supported by the University of Bath University Research

Studentship Account (URSA) awarded to TBT and NKP; and a Royal Society Dorothy Hodgkin

Research Fellowship awarded to TBT (DH150169). Bioinformatics analysis of the paper was carried

out using MRC CLIMB Infrastructure, and Illumina Whole-Genome Sequencing by the Milner

Genomics Centre, Bath, UK and MicrobesNG, Birmingham, UK.

References

Alsohim AS, Taylor TB, Barrett GA, Gallie J, Zhang X, Altamirano-Junqueira AE, Johnson LJ,

Rainey PB, Jackson RW. 2014. The biosurfactant viscosin produced by Pseudomonas

fluorescens SBW25 aids spreading motility and plant growth promotion. *Environ. Microbiol.*

16:2267–2281.

Avrani S, Bolotin E, Katz S, Hershberg R. 2017. Rapid Genetic Adaptation during the First Four

Months of Survival under Resource Exhaustion. *Mol. Biol. Evol.* 34:1758–1769.

Bailey SF, Blanquart F, Bataillon T, Kassen R. 2017. What drives parallel evolution?: How

population size and mutational variation contribute to repeated evolution. *BioEssays* 39:1–9.

Bailey SF, Guo Q, Bataillon T. 2018. Identifying drivers of parallel evolution: A regression model

approach. *Genome Biol. Evol.* 10:2801–2812.

Bailey SF, Rodrigue N, Kassen R. 2015. The effect of selection environment on the probability of

parallel evolution. *Mol. Biol. Evol.* 32:1436–1448.

Barrett RDH, M'Gonigle LK, Otto SP. 2006. The distribution of beneficial mutant effects under

strong selection. *Genetics* 174:2071–2079.

Blount ZD, Barrick JE, Davidson CJ, Lenski RE. 2012. Genomic analysis of a key innovation in an
experimental Escherichia coli population. *Nature* 489:513–518.

De Boer JG, Ripley LS. 1984. Demonstration of the production of frameshift and base-substitution
mutations by quasipalindromic DNA sequences. *Proc. Nail. Acad. Sci. USA* 81:5528-5531.

Bryksin A V, Matsumura I. 2010. Overlap extension PCR cloning: a simple and reliable way to create
recombinant plasmids. *Biotechniques* 48:463–465.

Bull JJ, Badgett MR, Wichman HA, Huehenbeck JP, Hillis DM, Gulati A, Ho C, Molineux IJ. 1997.
Exceptional Convergent Evolution in a Virus. *Genetics* 147:1497–1507.

Connor TR, Loman NJ, Thompson S, Smith A, Southgate J, Poplawski R, Bull MJ, Richardson E,
Ismail M, Elwood-Thompson S, et al. 2016. CLIMB (the Cloud Infrastructure for Microbial
Bioinformatics): an online resource for the medical microbiology community. *Microb.*
Genomics 2:6.

Van Ditmarsch D, Boyle KE, Sakhtah H, Oyler JE, Carey D, Déziel É, Dietrich LEP, Xavier JB.
2013. Convergent Evolution of Hyperswarming Leads to Impaired Biofilm Formation in
Pathogenic Bacteria. *Cell Rep* 4:697–708.

Dong F, Allawi HT, Anderson T, Neri BP, Lyamichev VI. 2001. Secondary structure prediction and
structure-specific sequence analysis of single-stranded DNA. *Nucleic Acids Res.* 29:3248–3257.

Duan C, Huan Q, Chen X, Wu S, Carey LB, He X, Qian W. 2018. Reduced intrinsic DNA curvature
leads to increased mutation rate. *Genome Biol.* 19:1–12.

Eyre-Walker A, Hurst LD. 2001. The evolution of isochores. *Nat. Rev. Genet.* 2:549–555.

Fieldhouse D, Golding B. 1991. A source of small repeats in genomic DNA. *Genetics* 129:563–572.

Fong SS, Joyce AR, Palsson BØ. 2005. Parallel adaptive evolution cultures of Escherichia coli lead to
convergent growth phenotypes with different gene expression states. *Genome Res.* 858:1365–
1372.

Fraebel DT, Mickalide H, Schnitkey D, Merritt J, Kuhlman TE, Kuehn S. 2017. Environment

determines evolutionary trajectory in a constrained phenotypic space. *eLIfe* 6:e24669.

Frumkin I, Lajoie MJ, Gregg CJ, Hornung G, Church GM, Pilpel Y. 2018. Codon usage of highly

expressed genes affects proteome-wide translation efficiency. *Proc. Natl. Acad. Sci. U. S. A.*

115:E4940–E4949.

Gillespie JH. 1984. Molecular Evolution Over the Mutational Landscape. *Evolution (N. Y).* 38:1116.

Hermisson J, Pennings PS. 2005. Soft sweeps: Molecular population genetics of adaptation from

standing genetic variation. *Genetics* 169:2335–2352.

Herron MD, Doebeli M. 2013. Parallel Evolutionary Dynamics of Adaptive Diversification in

Escherichia coli. *PLoS Biol.* 11:e1001490.

Hmelo LR, Borlee BR, Almblad H, Love ME, Randall TE, Tseng BS, Lin CY, Irie Y, Storek KM,

Yang JJ, et al. 2015. Precision-engineering the Pseudomonas aeruginosa genome with two-step

allelic exchange. *Nat. Protoc.* 10:1820–1841.

Huang S, Pang L. 2012. Comparing statistical methods for quantifying drug sensitivity based on in

vitro dose-response assays. *Assay Drug Dev. Technol.* 10:88–96.

Kram KE, Geiger C, Ismail WM, Lee H, Tang H, Foster PL, Finkel SE. 2017. Adaptation of

Escherichia coli to Long-Term Serial Passage in Complex Medium: Evidence of Parallel

Evolution. *mSystems* 2:1–12.

Kristofich J, Morgenthaler AB, Kinney WR, Ebmeier CC, Snyder DJ, Old WM, Cooper VS, Copley

SD. 2018. Synonymous mutations make dramatic contributions to fitness when growth is limited

by a weak-link enzyme.Matic I, editor. *PLOS Genet.* 14:e1007615.

Krug J. 2019. Accessibility percolation in random fitness landscapes. *bioRxiv*. Available from:

http://arxiv.org/abs/1903.11913

Kudla G, Murray AW, Tollervey D, Plotkin JB. 2009. Coding-sequence determinants of gene

543   expression in Escherichia coli. *Science.* 324:255–258.

544 Lässig M, Mustonen V, Walczak AM. 2017. Predicting evolution. *Nat. Ecol. Evol.*1:1–9.

545 Lebeuf-Taylor E, McCloskey N, Bailey SF, Hinz A, Kassen R. 2019. The distribution of fitness

546   effects among synonymous mutations in a gene under directional selection. *eLIfe* 8:e45952.

547 Lind PA, Andersson DI. 2008. Whole-genome mutational biases in bacteria. *Proc. Natl. Acad. Sci. U.*

548   *S. A.* 105:17878–17883.

549 Lind PA, Farr AD, Rainey PB. 2015. Experimental evolution reveals hidden diversity in evolutionary

550   pathways. *eLIfe* 4:e07074.

551 Lind PA, Libby E, Herzog J, Rainey PB. 2019. Predicting mutational routes to new adaptive

552   phenotypes. e*LIfe* 8:e38822.

553 Long H, Sung W, Miller SF, Ackerman MS, Doak TG, Lynch M. 2014. Mutation rate, spectrum,

554   topology, and context-dependency in the DNA mismatch repair-deficient Pseudomonas

555   fluorescens ATCC948. *Genome Biol. Evol.* 7:262–271.

556 M. Kimura. 1968. Evolutionary Rate at the Molecular Level. *Nature* 217:624–626.

557 Markham NR, Zuker M. 2005. DINAMelt web server for nucleic acid melting prediction. *Nucleic*

558   *Acids Res.* 33:577–581.

559 McDonald MJ, Gehrig SM, Meintjes PL, Zhang XX, Rainey PB. 2009. Adaptive divergence in

560   experimental populations of Pseudomonas fluorescens. IV. Genetic constraints guide

561   evolutionary trajectories in a parallel adaptive radiation. *Genetics* 183:1041–1053.

562 Mcgee LW, Sackman AM, Morrison AJ, Pierce J, Anisman J, Rokyta DR. 2016. Synergistic

563   Pleiotropy Overrides the Costs of Complexity in Viral Adaptation. *Genetics* 202:285–295.

564 Mcgrath PT, Xu Y, Ailion M, Garrison JL, Butcher RA, Bargmann CI. 2011. Parallel evolution of

565   domesticated Caenorhabditis species targets pheromone receptor genes. Nature 477: 321-325.

566 Meyer JR, Dobias DT, Weitz JS, Barrick JE, Quick RT, Lenski RE. 2012. Repeatability and

contingency in the evolution of a key innovation in phage lambda. *Science.* 335:428–432.

Miller C, Kong J, Tran TT, Arias CA, Saxer G, Shamoo Y. 2013. Adaptation of Enterococcus faecalis

to daptomycin reveals an ordered progression to resistance. *Antimicrob. Agents Chemother.*

57:5373–5383.

Notley-McRobb L, Ferenci T. 1999. Adaptive mgl-regulatory mutations and genetic diversity

evolving in glucose-limited Escherichia coli populations. *Environ. Microbiol.* 1:33–43.

Ogle DH, Wheeler P, Dinno A. 2020. FSA: Fisheries Stock Analysis. Available from:

https://github.com/droglenc/FSA

Orr HA. 2005. THE PROBABILITY OF PARALLEL EVOLUTION. *Evolution (N. Y).* 59:216.

Ostrowski EA, Woods RJ, Lenski RE. 2008. The genetic basis of parallel and divergent phenotypic

responses in evolving populations of Escherichia coli. *Proc. R. Soc. B Biol. Sci.* 275:277–284.

R Core Team. 2014. R: A language and environment for statistical computing. *R Found. Stat.*

Comput. Vienna, Austria. Available from: http://www.r-project.org/.

Riehle MM, Bennett AF, Long AD. 2001. Genetic architecture of thermal adaptation in Escherichia

coli. *Proc. Natl. Acad. Sci. U. S. A.* 98:525–530.

Ripley LS. 1982. Model for the participation of quasi-palindromic DNA sequences in frameshift

mutation. *Proc. Natl. Acad. Sci. U. S. A.* 79:4128–4132.

Sackman AM, McGee LW, Morrison AJ, Pierce J, Anisman J, Hamilton H, Sanderbeck S, Newman

C, Rokyta DR. 2017. Mutation-driven parallel evolution during viral adaptation. *Mol. Biol. Evol.*

34:3243–3253.

Seemann T. 2015. Snippy: fast bacterial variant calling from NGS reads. Available from:

https://github.com/tseemann/snippy

Spor A, Kvitek DJ, Nidelet T, Martin J, Legrand J, Dillmann C, Bourgais A, De Vienne D, Sherlock

G, Sicard D. 2014. Phenotypic and genotypic convergences are influenced by historical

contingency and environment in yeast. *Evolution (N. Y)*. 68:772–790.

Stoltzfus A, McCandlish DM. 2017. Mutational biases influence parallel adaptation. *Mol. Biol. Evol.*

34:2163–2172.

Taylor TB, Mulley G, Dills AH, Alsohim AS, McGuffin LJ, Studholme DJ, Silby MW, Brockhurst

MA, Johnson LJ, Jackson RW. 2015. Evolutionary resurrection of flagellar motility via rewiring

of the nitrogen regulation system. *Science*. 347:1014–1017.

Tenaillon O, Rodríguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, Gaut BS. 2012.

The molecular diversity of adaptive convergence. *Science*. 335:457–461.

Turner CB, Marshall CW, Cooper VS. 2018. Parallel genetic adaptation across environments differing

in mode of growth or resource availability. *Evol. Lett.* 2:355–367.

Vogwill T, Kojadinovic M, Furió V, Maclean RC. 2014. Testing the role of genetic background in

parallel evolution using the comparative experimental evolution of antibiotic resistance. *Mol.*

Biol. Evol. 31:3314–3323.

Weinreich DM, Delaney NF, De Pristo MA, Hartl DL. 2006. Darwinian Evolution Can Follow Only

Very Few Mutational Paths to Fitter Proteins. *Science*. 312.

Wichman HA, Badgett MR, Scott LA, Boulianne CM, Bull JJ. 1999. Different trajectories of parallel

evolution during viral adaptation. *Science*. 285:422–424.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Available from:

https://ggplot2.tidyverse.org

Wood TE, Burke JM, Rieseberg LH. 2005. Parallel genotypic adaptation: When evolution repeats

itself. *Genetica* 123:157–170.

Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. 2006. Tests of parallel molecular

evolution in a long-term experiment with Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.*

103:9107–9112.

Wright BE, Reschke DK, Schmidt KH, Reimers JM, Knight W. 2003. Predicting mutation

frequencies in stem-loop structures of derepressed genes: Implications for evolution. *Mol.*

Microbiol. 48:429–441.

 Figures and figure legends:

Fig. 1. Extremely parallel evolution of flagella-mediated motility in immotile variants of *P. fluorescens* SBW25 (AR2). (A) Immotile populations evolved on soft agar (left) re-evolved flagella-mediated motility through one-step *de novo* mutation (right). (B) Phenotype emergence appeared rapidly, typically within 3-5 days following inoculation (box edges represent the 25th and 75th percentiles and the whiskers show the observed range). (C) The underlying genetic changes were highly parallel, with all independent lines targeting one of two sites (left circle, A289C and right circle Δ406-417) within the *ntrB* locus at the expense of other sites within the nitrogen (ntr) pathway. (D) A single transversion mutation, A289C, was the most common mutational route, appearing in over 95% of independent lines (23/24).

Fig. 2. Repeatability of the A289C *ntrB* mutation across genetic background and nutrient environment (total $N = 116$). The proportion of each observed mutation is shown on the y axis. *ntrB* mutation A289C was robust across both strain backgrounds (SBW25Δ*fleQ* shown as Δ*fleQ*, and AR2) and the four tested nutritional environments, remaining the primary target of mutation in all cases (>87%). Lines were evolved using 4mM, 8mM and 16mM of amino acid supplement (see materials and methods). No significant relationship between supplement concentration and evolutionary target was observed (Kruskal-Wallis chi-squared tests: AR2 M9+glu, df = 2, $P > 0.2$; AR2 M9+gln, df = 1, $P > 0.23$; Δ*fleQ* M9+gln, df = 1, $P > 0.3$), as such they are treated as independent treatments for statistical analysis but visually grouped here for convenience. Δ*fleQ* lines evolved on LB were able to migrate rapidly through sliding motility alone, masking any potential emergent flagellate blebs (see Alsohim et al., 2014). Sample sizes (N) for other categorical variables: Δ*fleQ* – M9: 25, M9+gln: 20, M9+glu: 7; LB: 5, M9: 24, M9+gln: 17, M9+glu: 18.

Fig. 3. Selection does not strongly favour *ntrB* A289C motility over alternative *ntrB* mutations. (A)

Surface area of motile zones following 48h of growth across four environmental conditions. Individual

data points from biological replicates are plotted and each migration area has been standardised against

the surface area of a *ntrB* A289C mutant grown in the same environment (*ntrB* A289C growth mean =

0). Significance values: * = $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$ (Kruskal-Wallis post-hoc Dunn

test). (B) *ntrB* A289C lines fail to reach the growth frontier within 6 h of competitor pre-inoculation.

Two *ntrB* mutant lines, A289C and A683C, were co-inoculated in equal amounts on soft agar, either

immediately (0 h) or with A289C being added at 3 h time points up to 12 h (x-axis) into the centre of

an A683C inoculated zone. The strains were competed for 24 h prior to sampling from the motile zone's

leading edge. Genotype establishment at the frontier across the four replicates is shown on the y-axis

with the number of lineages at the leading edge represented as 0-4.

Fig. 4. Loss of repeatable evolution conferred by a synonymous sequence mutant (AR2-sm). (A) Histogram of motility phenotype emergence times across independent replicates of immotile SBW25 (AR2) and AR2-sm in two nutrient conditions. (B) Observed mutational targets across two environments (AR2: LB $N = 5$, M9 $N = 24$; AR2-sm: LB $N = 8$, M9 $N = 8$). Note that characterised genotypes were sampled within 8 days of experiment start date. Unidentified mutations could not be distinguished from wild type sequences of genes belonging to the nitrogen regulatory pathway *(ntrB, glnK* and *glnA)* which were analysed by Sanger sequencing (supplementary table. S1). *ntrB* Δ406-417 was the only mutational target shared by both lines within the same nutritional environment.

Supplementary figures and tables:

Fig. S1. Growth kinetics of mutant AR2 lines in static liquid culture over 24h. Nutrient environments:

M9 = M9 minimal media supplemented with NH_4 at 7.5 mM. M9+glu = additional glutamate added

at 8 mM. M9+gln = additional glutamine added at 8 mM. LB = lysogeny broth. Growth yield was

determined using area under the curve, and each yield has been standardised against the yield of the

AR2 ancestral strain grown in the same environment (AR2 ancestor growth mean = 0). Individual

data points from biological replicates are plotted, and ranges around the mean growth of the ancestral

strain are shown in column one of each frame. Plots are the means of six biological replicates.

Significance values: * = $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$ (one-way ANOVA post-hoc Tukey

HSD test).

Fig. S2. Quasi-palindromic sequences flank *ntrB* site 289 in both *P. fluorescens* strains SBW25 and

Pf0-1. Theoretical hairpin formations were generated using the *mfg* program (Wright et al. 2003).

This software calculates the most stable hairpin formed between neighbouring tracts ($\pm$ 40 nucleotides

from site 289) in which the site of interest (in this case site 289, highlighted in red) remains unpaired.

In these examples the nucleotides are forced into stem-loop structures that have been documented to

comprise hairpins (Ripley, 1982). The stability, structure and included nucleotide tracts that form

the predicted hairpins differ between strains (Maximum-energy (ΔG) of secondary structures: Pf0-1 =

-13.2. SBW25 = -8.0. AR2-sm = -11.6). These differences are partially owed to synonymous

sequence variation, as highlighted by the unique hairpin formation exhibited by AR2-sm after the

introduction of 6 synonymous substitutions. Although the *mfg* program only calls the most stable

hairpin configuration and therefore may miss alternative structures that temporarily form and raise

mutation rate, the tool exemplifies the power of synonymous variance in altering hairpin stability.

Fig. S3. *ntrB* A289C in AR2-sm retains comparative fitness to its ancestral counterpart. The motility

phenotype of AR2 *ntrB* A289C and alternative AR2-sm *ntrB* mutants (Δ406-417-sm and A683-sm)

were measured against an engineered AR2-sm *ntrB* A289C mutant (A289C-sm) in minimal medium.

A289C-sm was not significantly outperformed by any strain, instead showing a significantly superior

motility phenotype to A683-sm in M9. Although the two motile lines displayed comparable motility

in an AR2 background (fig. 3A), the inferior phenotype observed here may be owed to an

uncharacterised secondary mutation. Individual data points from biological replicates are plotted and

each migration area has been standardised against the surface area of a *ntrB* A289C-sm mutant grown

in the same environment (*ntrB* A289C-sm growth mean = 0). Significance values: * = $P < 0.05$, ** =

$P < 0.01$, *** = $P < 0.001$ (Kruskal-Wallis post-hoc Dunn test).

For use in:	Primer description:	Sequence:
Sanger sequencing of ntr pathway / Invasion assay	SBW25 *ntrB* locus (forward)	5'- GAGGTCCCAATGACCATCAG -3'
	SBW25 *ntrB* locus (reverse)	5'- GACGATCCAGACGGTTTCAC -3'
	SBW25 *glnK* locus (forward)	5'-GTGGGCAAAGGACTGATTTC-3'
	SBW25 *glnK* locus (reverse)	5'-GATGATGGCGAAGGTCATCT-3'
	SBW25 *glnA* locus (forward)	5'-CGGAAATCGCTCAAGGTTTA-3'
	SBW25 *glnA* locus (reverse)	5'-CTGATAATCCCCAGGCAAAA-3'
ntrB A683C integration into pTS1 backbone (allelic exchange)	Upstream fragment (forward)	5'- GAAATTAATAGGTTGTATTGATGTTGATGACCATCAGCGATGCACTG -3'
	Upstream fragment (reverse)	5'- GAATGCTCGGGGCGTAGTCGC -3'
	Downstream fragment (forward)	5'- GCGACTACGCCCCGAGCATTC -3'
	Downstream fragment (reverse)	5'- GCCGTTTCTGTAATGAAGGAGAAAACTCATGTCGATGGGGCTCCTTG -3'
ntrB synonymous substitution sequence integration into pTS1 backbone (allelic exchange)	Upstream fragment (forward)	5'- GAAATTAATAGGTTGTATTGATGTTGTGCCAAATGCCGCCTACATC -3'
	Upstream fragment (reverse)	5'- CGTTGCTGAGGATCGGCGTCACCGCGTAATCCACCGTCAG -3'
	Downstream fragment (forward)	5'- CTGACGGTGGATTACGCGGTGACGCCGATCCTCAGCAACG -3'
	Downstream fragment (reverse)	5'- GCCGTTTCTGTAATGAAGGAGAAAACGTTGATCAGCACGGTGATGT -3'
	SBW25 *ntrB* nested primer (forward)	5'- AATTTGGATCCATGACCATCAGCGATGCACTG -3'
	SBW25 *ntrB* nested primer (reverse)	5'- AATTTAAGCTTGATCCAGACGGTTTCACTACG -3'
ntrB synonymous substitution sequence with A289C	Upstream fragment (reverse)	5'- CGTTGCTGAGGATCGGCGGCACCGCGTAATCCACCGTCAG -3'
	Downstream fragment (forward)	5'- CTGACGGTGGATTACGCGGTGCCGCCGATCCTCAGCAACG -3'

704   Table. S1. List of primers used throughout the study.