Aurora kinases promote mitotic progression and asymmetric cell division through

activation of Polo kinase

Emmanuel Gallaud[1], Aude Pascal[1], Mathieu Métivier[1]*, Laurent Richard-Parpaillon[1], Vincent

Archambault[2] and Régis Giet[1]*

[1] Univ Rennes, CNRS, IGDR (Institut de Génétique et Développement de Rennes) - UMR

6290, F-35000 Rennes, France

[2] Institute for Research in Immunology and Cancer, Université de Montréal, Montréal, QC,

Canada.

* Corresponding author: regis.giet@univ-rennes1.fr

Abstract

Organ development and integrity requires the maintenance of a defined and restricted pool of

dividing stem cells. This process requires the coupling of mitosis progression to the

establishment of stem cells polarization and spindle orientation to ensure that they retain the

ability to proliferate while their descendant cells do not proliferate excessively. How this

coupling occurs is unknown. Here we show that downstream of Aurora kinases, Polo activity

levels are crucial for timely mitotic progression independently from the Spindle Assembly

Checkpoint (SAC). In addition, Polo functions downstream of Aurora A to ensure cortical

polarity and spindle orientation in neural stem cells, thereby preventing excessive cellular

proliferation in developing larval brains, and suppressing tumor development. In addition,

induction of aneuploidy in *polo* mutant brain by inactivation of the SAC leads to an increase

in tumor development Altogether, our results reveal that the Aurora-Polo kinase axis is an

essential module coupling mitotic progression to asymmetric division in NSCs.

Introduction

Correct chromosome segregation in many cell types depends on the Spindle Assembly

Checkpoint (SAC) that delays Cyclin B degradation and anaphase onset until all kinetochores

are properly attached to spindle microtubules. As a consequence, ablation of SAC genes in

dividing cancer cells induces premature anaphase onset and chromosome segregation defects

[1–5]. In *aurA* mutant NSCs, mitosis is prolonged and it was initially thought that this effect was

due to activation of the SAC[6,7]. Consistent with this idea, these cells are delayed in

assembling a functional bipolar spindle and Cyclin B-Cdk1 levels persist longer than normal.

However, SAC ablation in *aurA* mutant does not restore normal mitotic progression nor lead

to severe aneuploidy; indicating that the prolonged time in mitosis does not result from SAC

activation in *aurA* mutants[4]. Thus, despite increased knowledge in the control of mitotic

progression, how Aurora A kinase regulates this aspect remains elusive. A possible

downstream target is Polo kinase, which is known to be activated by Aurora A and Aurora

B[8–13]. In addition, other reports have suggested that Polo kinase contributes to mitotic exit by

phosphorylation of several subunits of the APC/C (Anaphase Promoting

Complex/Cyclosome), which triggers Cyclin B degradation and anaphase[14–20]. Moreover

mitosis is also prolonged in Polo-depleted *Drosophila* S2 cells[21].

In the fly NSCs, polarization of the apical cortex is an essential step that ensures asymmetric

localization of PAR proteins at the apical cortex and cell fate determinants at the basal cortex.

After mitotic spindle alignment along the apico-basal axis and cytokinesis, one cell retains the

NSC proliferation fate while the other cell (GMC for Ganglion Mother Cell) inherits the cell

fate determinants anchored at the basal cortex and is subjected to differentiation (reviewed in

[22]). Both AurA and Polo are needed for cell polarization and for mitotic spindle alignment[23–27]. As a consequence, the two daughter cells acquire the NCS proliferative fate in *aurA* and

polo mutants and a strong increase of NSC-like population occurs leading to tumor formation

in *aurA* and *polo* mutant neural tissues. Therefore, in the context of central brain

development, *aurA* and *polo* genes are both tumor suppressors[23,24,27]. Altogether, the

contributions of a potential cascade linking Aurora A and Polo kinases ([28] for review) to the

control of time in mitosis, spindle orientation, cell polarization and their consequences for

tissue development remains unknown. In the context of the stem-cell based *Drosophila*

central brain development, we show here that these key features of NSCs are mainly regulated

by the activation of Polo by Aurora A and Aurora B.

polo mutants NSCs exhibit a SAC-independent mitotic delay.

We first wished to determine if Polo activity promotes mitotic progression in NSCs

independently from its role in ensuring correct spindle assembly and chromosome

attachments in mitosis. To this end, we examined how time in mitosis was regulated in

different *polo* mutants associated with a loss of the SAC (*polo,mad2^P*). Western blotting

analyses of brain tissues confirmed the absence of Mad2 in the double mutants (Figure 1A).

Using time lapse-imaging microscopy, we found that the weak *polo^1/polo^{10}* and *polo^1/polo^9*

hypomorphic mutants showed a prolonged mitosis that was mostly SAC-independent (Figure

1B and C). The time in mitosis was strongly increased in strong hypomorphic *polo^9/polo^{10}*

mutants but, by contrast to weak *polo* hypomorphs, ablation of the SAC decreased the time in

mitosis, indicating a higher SAC contribution to mitotic delay in strong *polo* mutants. These

results were confirmed by analyses of the delay between SAC silencing (as detected by the

loss of GFP::Mad2 at kinetochores) and anaphase onset in weak and strong *polo* mutants.

This time was 0.8 min in controls, 2.1 to 2.4 min in *polo^1/polo^{10}* and *polo^1/polo^9* and 8.6 min

in *polo^9/polo^{10}* NSCs. This result reveals that weak interference with Polo kinase prolonged

mitosis in a mostly SAC-independent manner but stronger decrease in Polo kinase activity

triggers a mitotic delay that is also due to SAC activation. Altogether, these results indicate

that the SAC is satisfied well before anaphase onset and cannot account for the prolonged

mitosis of *polo* mutant NSCs, similarly to *aurA* mutant NSCs[4].

Constitutively active Polo partially rescues mitotic delays in *polo* and *aurA* mutants.

In human cells, the activation of Plk1 is triggered by the phosphorylation of the Thr210

residue in its activation loop by Aurora A[8–10,28]. Therefore, we wondered if this Aurora A-

mediated activation of Polo kinase was conserved in the fly and could also regulate the time

in mitosis in NSCs. We hypothesized that the expression of a constitutively active Polo kinase

(Polo^{T182D} in flies) should decrease the duration of mitosis in *aurA* mutants. In order to test

this idea, we generated transgenic flies allowing the expression of different GFP-tagged

variants of Polo kinase in brain tissues including wild-type (Polo^{WT}), kinase-dead (Polo^{K54M}),

constitutively active (Polo^{T182D}) and constitutively inactive (Polo^{T182A}). We first analyzed the

functionality of Polo::GFP variants in NSCs. Their localization was similar to what was

previously observed in early embryos and S2 cells (Figure S1 and [29]). Moreover, only

Polo^{WT}::GFP was able to restore viability of the strong *polo$^{9/10}$* hypomorph (hereafter referred

to as *polo* mutant) when expressed under the control of a ubiquitous Actin5C promoter

(Figure 2A-B). We then assayed the ability of each of these variants to reduce the time in

mitosis of *polo* NCSs by live microscopy (Figure 2C-D). Expression of Polo^{T182D}, Polo^{T182A}

and Polo^{K54M} in a WT background slightly increased the time in mitosis of NSCs when

compared to the expression of Polo^{WT}, suggesting hypermorphic or dominant-negative

effects. Moreover, only expression of Polo^{WT} was able to restore a normal time in mitosis

when expressed in *polo* mutants (8.0 vs 45.6 min). Interestingly, expression of the Polo^{T182D}

variant decreased by more than half the time in mitosis of *polo* mutant NSCs. Given the

literature, we envisaged that the excess of Polo kinase activity in WT and *polo* mutant cells

may perturb kinetochore attachments to spindle microtubules and activate the SAC[30–34]. We

therefore decided to examine the time in mitosis of NSCs upon Polo^{T182D} expression, in

mad2^P mutant and *polo,mad2^P* double mutant. Interestingly, the increased time in mitosis

resulting from Polo^{T182D} expression in WT cells was not abrogated following deletion of the

SAC, indicating that this delay did not result from aberrant kinetochore attachments.

However, expression of Polo^{T182D} was able to reduce by 34.2% the time in mitosis in

polo,mad2^P double mutant NSCs. These results suggest that Polo^{T182D} is not a fully functional

protein, in agreement with a previous study in human cells[32]. Finally, the ability of Polo^{T182D}

to delay mitotic exit (anaphase entry) in a SAC-independent manner (in WT NSCs) while

accelerating mitotic exit in *polo,mad2^P* double mutants, suggests that Polo activation levels

must be finely controlled to set the normal duration of mitosis.

To investigate if the mitotic delay observed in *aurA* and in *aurA,mad2^P* mutants was Polo-

kinase-dependent, we expressed Polo^{WT} and Polo^{T182D} in these backgrounds (Figure 3A-B and

S2A). We found that expression of Polo^{WT} and Polo^{T182D} decreased the time in mitosis of

aurA and *aurA,mad2^P* mutants NSCs, suggesting that Polo kinase T182 phosphorylation by

Aurora A contributes to mitotic exit.

AurA and Ial are both required for Polo-dependent mitotic progression.

The mitotic phosphorylation of Polo on T182 is also mediated by Aurora B kinase/Ial (Ipl1-

Aurora-like in flies), a component of the Chromosome Passenger Complex (CPC) at

centromeres and during cytokinesis[11,12,29]. We therefore wondered if Aurora B was also

contributing, with Aurora A, to the control of the time in mitosis through Polo kinase

phosphorylation on T182. As described before for *CPC* (*survivin*) mutants, *ial* mutant NSCs

displayed prolonged mitosis (9.2 min for *ial^{2443}/ial^{1689}* vs 6.0 min in a control, Figure 3C-D,

S2C and [35]). This delay was significantly reduced upon Polo^{T182D} expression (8.5 min)

indicating that the mitotic delay in *ial* mutant is partially Polo kinase-dependent. Strikingly,

combination of *ial* and *aurA* mutations triggered a synergistic delay in the time in mitosis, as

most of the double mutant NSCs failed to exit mitosis after more than 150 min (Figure S2B-

C).

Polo and the SAC protect NSCs against aneuploidy and excessive proliferation.

In addition to their role for the control of the time in mitosis, AurA and Polo are both required

for cell polarization and mitotic spindle alignment in NSCs[23,24,27]. Thus, we hypothesized that

activation of Polo by AurA may partly underly their requirements in the asymmetric division

of NSCs. We first analyzed the larval brain phenotypes of the weak and strong *polo*

hypomorphs. We found that NSC numbers (Deadpan-positive cells) increased in the central

brain of both weak and strong *polo* mutants. Interestingly, ablation of the SAC with the

mad2^P mutation abrogated this increase in a strong *polo* hypomorph but not in weak *polo*

hypomorphs (Figure 4A-B). As NSCs proliferation can be perturbed by aneuploidy in flies[4,5],

we also monitored the aneuploidy and polyploidy levels of weak and strong *polo* hypomorphs

associated with loss of the SAC (Figure 4C-D). We found that 36 to 41% of the mitotic cells

in *polo,mad2[P]* double mutants were either aneuploid or polyploid, indicating that *polo* mutant

brain tissues were prone to chromosome segregation errors.

To check if the number of NSCs reflects their ability to proliferate, we performed brain

transplantation experiments. We transplanted weak or strong *polo* mutant brain tissues in

which the SAC was active or not, in the abdomen of host WT flies and monitored the growth

of the transplants[36,37]. Surprisingly, inactivation of the SAC and the subsequent increase in

aneuploidy correlated with severely enhanced tumor growth of weaker *polo* mutant NSCs

(45.1 and 51.5% vs. 22.0 and 24.7% of transplanted flies). SAC inactivation had no effect on

strong *polo* tumor growth as it was already close to maximal (85.2 vs. 86.5% of transplanted

flies) (Figure 4E-F). These results suggest that Polo collaborates with the SAC to protect

NSCs against chromosome segregation errors and excessive proliferation. In this way, Polo is

akin to a tumor suppressor.

AurA-dependent tumor suppression is largely mediated by Polo.

Given that Polo and AurA both act like tumor suppressors in NSCs, we decided to investigate

if the tumor suppression potential of *aurA* mutants was Polo-dependent. We therefore

expressed a Polo[T182D] kinase in an *aurA* mutant background and quantified the number of

NSCs in the central brain region. Consistent with several studies[4,23,27], the NSC number in

aurA mutants was increased 10-25 times compared to control brains (2050 NSCs per lobe,

Figure 5 A-B). Strikingly, while expression of Polo[T182D] had no effect on its own (95 NSCs

per lobe), it was sufficient to strongly diminish the NSC amplification incurred in *aurA*

mutant tissues (390 NSCs). We also compared tumor growth progression after transplantation

of *aurA* mutant neural tissues expressing or not Polo[T182D] (Figure 5C-D). We found that 90%

of transplanted *aurA* mutant brain tissues labeled with mCherry produced large tumors that

killed host flies after 15 days. When Polo[T182D] was expressed in *aurA* mutant, only 10 % and

55 % of the injected hosts developed tumors after 15 and 30 days respectively, indicating that

the tumor suppressor potential of AurA is Polo-kinase dependent (Figure 5D). Importantly,

the ploidy defects caused by *aurA* mutation (11.8% of the cells) were unchanged following

Polo^{T182D} expression (13.1% of the cells), ruling out differences in ploidy as the cause of the

aurA mutant tumor growth restriction by Polo^{T182D} (Figure 5E).

Aurora A is required to regulate larval brain development by controlling mitotic spindle

orientation and cell polarization in NSC asymmetric division [23,25–27]. We therefore analyzed

these two processes in *aurA* mutant NSCs expressing Polo^{T182D}. We first monitored NSC

polarization by analyzing the localization of aPKC and Miranda apical and basal crescents

respectively. We found that both aPKC and Miranda crescents were properly localized in 100

% of control or Polo^{T182D} expressing NSCs. By contrast and in agreement with previous

reports, most *aurA* mutant NSCs (78%) displayed abnormally weak homogenous cortical

aPKC while Miranda was properly localized at the basal cortex (Figure 5F-H). Strikingly,

expression of Polo^{T182D} in *aurA* mutant restored the presence of the aPKC apical crescent in

84% of the NCSs. In parallel, we also measured mitotic spindle alignment along the apico-

basal axis in *aurA* mutant with or without Polo^{T182D} expression. In control and Polo^{T182D}

expressing NSCs, the vast majority of mitotic spindles were properly oriented in metaphase

(angle < 30°: 94% and 100%) while *aurA* mutant NSCs displayed a random mitotic spindle

alignment (angle < 30°: 41.8%) in agreement with previous reports [23,27]. Strikingly, Polo^{T182D}

expression was sufficient to significantly restore the *aurA* mutant-dependent spindle

alignment defects (angle < 30°: 76.3%; Figure 5F, I and J). Altogether, these results indicate

that Polo functions downstream of AurA to ensure the asymmetry of NSC divisions that is

required to prevent their excessive proliferation during larval brain development.

The regulation of the time in mitosis is a complex process that is regulated by the

maintenance of Cyclin-CDK activity, satisfaction of the SAC and activation of the APC/C by

phosphorylation/dephosphorylation events[17–20] ([38] for review). Here we present several

evidence supporting the idea that the Aurora-Polo signaling cascade is pivotal to control the

duration of mitosis. Firstly, the loss of Aurora A or Aurora B leads to a synergistic extension

of the time in mitosis, much like inactivation of Polo. This mitotic delay can be rescued by the

expression of a constitutively active Polo^{T182D}, consistent with the idea that Aurora-dependent

phosphorylation of Polo at T182 is the underlying connection. Interestingly, the use of weak

or strong hypomorphic *polo* mutants and the controlled expression of a dominant Polo^{T182D}

active kinase allowed to establish the importance of Polo kinase activity thresholds. We found

that weak *polo* hypomorphs are characterized by near normal spindle formation and a

prolonged mitosis in which the SAC has little if no contribution. By contrast, strong *polo*

hypomorphs exhibit spindle defects associated with a prolonged mitosis that is partially SAC-

independent. The use of active Polo^{T182D} has enabled us to highlight the subversive function

of Polo kinase in the control of the time in mitosis. Indeed, when expressed in a wild type

background, Polo^{T182D} significantly lengthens mitosis whereas it shortens it in a null genetic

background (Figure 2D-E, Figure 3B and D and [33]). This apparent antagonism could be

explained by the fact that Polo inhibits the APC/C activator Cdc20 when the SAC is

unsatisfied but is also necessary to activate the phosphorylation of APC/C subunits and thus

the exit from mitosis[19,20,39].

This work also is first to demonstrate the essential role of the activation of Polo kinase by

Aurora A for the orientation of the mitotic spindle and the polarization of neural stem cells.

We show here that tumor growth of *aurA* mutant brains is strongly inhibited by the persistent

activation of Polo. This restoration of tissue homeostasis is likely caused by the almost

complete rescue of *aurA* mutant cell polarity defects following constant Polo activation.

Given that a past study showed that Aurora A phosphorylation of the PAR complex is the

main player leading to cell polarization at least *in vitro*[25], we speculate that yet

uncharacterized Polo-dependent phosphorylation of cortical or polarity proteins are also

essential to mediate NSC polarization.

In addition, rescue of tissue homeostasis by Polo[T182D] may also be mediated by a direct

regulation of the spindle orientation machinery by Polo together with Aurora A[26] or by

stimulation of centrosomal microtubule nucleation and interaction with the cell cortex ([13,40]

and [41] for review). In conclusion, this work has revealed that essential characteristics of

NSCs, such as the duration of mitosis and cell polarization, are in fact regulated by a signaling

cascade leading to the spatio-temporal activation of the Polo protein kinase by the Aurora

kinases, with a profound impact on tissue development. Further studies will be required to

identify and investigate the functions of the Polo targets in these essential processes.

Acknowledgements:

This work was supported by Fondation pour la Recherche Médicale (DEQ20170336742). We

thank members of the "Cytoskeleton and Cell Proliferation" lab as well as Roland Le Borgne

and Damien Coudreuse for helpful discussions. We are grateful to Renata Basto for advices

on tumor growth assays, Clemens Cabernard, David Sharp, Juliette Mathieu and Jean-René

Huynh for antibodies, flies and plasmids. We also acknowledge Stéphanie Dutertre and

Xavier Pinson for their help using the microscopes of the Microscopy Rennes imaging center

facility (MRic).

Author Contributions:

R.G. and E.G. conceived the project. E.G. performed experiments and quantifications with the

help of L.R.-P. (Figure 4E-F), A.P. (Figure 2A), R.G. (Figure 1D-E, Figure 4A-D) and M.M.

(initial experiment corresponding to Figure 3A-B). R.G. and E.G. discussed the data and

wrote the manuscript with the help of V.A.

Competing Interests statement:

The authors declare no competing interests.

Data availability:

Numerical data are available in the Table S1 and all other data are available upon request.

References

1.   Dominguez-Brauer, C. *et al.* Targeting Mitosis in Cancer: Emerging Strategies. *Mol.*
Cell **60**, 524–536 (2015).
2.   Janssen, A., Kops, G. J. P. L. & Medema, R. H. Elevating the frequency of
chromosome mis-segregation as a strategy to kill tumor cells. *Proc. Natl. Acad. Sci.* **106**,
19108–19113 (2009).
3.   Silk, A. D. *et al.* Chromosome missegregation rate predicts whether aneuploidy will
promote or suppress tumors. *Proc. Natl. Acad. Sci.* **110**, E4134–E4141 (2013).
4.   Caous, R. *et al.* Spindle assembly checkpoint inactivation fails to suppress neuroblast
tumour formation in aurA mutant Drosophila. *Nat. Commun.* **6**, (2015).
5.   Gogendeau, D. *et al.* Aneuploidy causes premature differentiation of neural and
intestinal stem cells. *Nat. Commun.* **6**, 8894 (2015).
6.   Glover, D. M., Leibowitz, M. H., McLean, D. A. & Parry, H. Mutations in aurora
prevent centrosome separation leading to the formation of monopolar spindles. *Cell* **81**, 95–
105 (1995).
7.   Giet, R. *et al.* Drosophila Aurora A kinase is required to localize D-TACC to
centrosomes and to regulate astral microtubules. *J. Cell Biol.* **156**, 437–451 (2002).
8.   Macůrek, L. *et al.* Polo-like kinase-1 is activated by aurora A to promote checkpoint
recovery. *Nature* **455**, 119–123 (2008).
9.   Seki, A., Coppinger, J. A., Jang, C.-Y., Iii, J. R. Y. & Fang, G. Bora and the Kinase
Aurora A Cooperatively Activate the Kinase Plk1 and Control Mitotic Entry. *Science* **320**, 5
(2008).
10.   Bruinsma, W., Macůrek, L., Freire, R., Lindqvist, A. & Medema, R. H. Bora and
Aurora-A continue to activate Plk1 in mitosis. *J. Cell Sci.* 11 (2014).
11.   Carmena, M. *et al.* The Chromosomal Passenger Complex Activates Polo Kinase at
Centromeres. *PLoS Biol.* **10**, 15 (2012).
12.   Kachaner, D. *et al.* Interdomain allosteric regulation of Polo kinase by Aurora B and
Map205 is required for cytokinesis. *J. Cell Biol.* **207**, 201–211 (2014).
13.   Joukov, V., Walter, J. C. & De Nicolo, A. The Cep192-Organized Aurora A-Plk1
Cascade Is Essential for Centrosome Cycle and Bipolar Spindle Assembly. *Mol. Cell* **55**, 578–
591 (2014).
14.   Shirayama, M., Zachariae, W., Ciosk, R. & Nasmyth, K. The Pololike kinase Cdc5p
and the WDrepeat protein Cdc20p/fizzy are regulators and substrates of the anaphase
promoting complex in Saccharomyces cerevisiae. *EMBO J.* **17**, (1998).
15.   Descombes, P. & Nigg, E. A. The pololike kinase Plx1 is required for M phase exit
and destruction of mitotic regulators in Xenopus egg extracts. *EMBO J.* **17**, (1998).
16.   Charles, J. F., Jaspersen, S. L., Tinker-Kulberg, R. L., Hwang, L. & Morgan, D. O.
The Polo-related kinase Cdc5 activates and is destroyed by the mitotic cyclin destruction
machinery in S. cerevisiae. *Curr. Biol.* **8**, (1998).
17.   Golan, A., Yudkovsky, Y. & Hershko, A. The Cyclin-Ubiquitin Ligase Activity of
Cyclosome/APC Is Jointly Activated by Protein Kinases Cdk1-Cyclin B and Plk. *J. Biol.*
Chem. **277**, (2002).
18.   Kraft, C. *et al.* Mitotic regulation of the human anaphase-promoting complex by
phosphorylation. *EMBO J.* **22**, 6598–6609 (2003).
19.   Qiao, R. *et al.* Mechanism of APC/CCDC20 activation by mitotic phosphorylation.
Proc. Natl. Acad. Sci. **113**, 9 (2016).
20.   Zhang, S. *et al.* Molecular mechanism of APC/C activation by mitotic
phosphorylation. *Nature* **533**, 19 (2016).
21.   Conde, C. *et al.* Drosophila Polo regulates the spindle assembly checkpoint through

Mps1dependent BubR1 phosphorylation. *EMBO J.* **32**, (2013).

22.   Gallaud, E., Pham, T. & Cabernard, C. Drosophila melanogaster Neuroblasts: A
Model for Asymmetric Stem Cell Divisions. in *Asymmetric Cell Division in Development,*
Differentiation and Cancer (eds. Tassan, J.-P. & Kubiak, J. Z.) vol. 61 183–210 (Springer
International Publishing, 2017).

23.   Wang, H. *et al.* Aurora-A acts as a tumor suppressor and regulates self-renewal of
Drosophila neuroblasts. *Genes Dev.* **20**, 3453–3463 (2006).

24.   Wang, H., Ouyang, Y., Somers, W. G., Chia, W. & Lu, B. Polo inhibits progenitor
self-renewal and regulates Numb asymmetry by phosphorylating Pon. *Nature* **449**, 96–100
(2007).

25.   Wirtz-Peitz, F., Nishimura, T. & Knoblich, J. A. Linking Cell Cycle to Asymmetric
Division: Aurora-A Phosphorylates the Par Complex to Regulate Numb Localization. *Cell*
135, 161–173 (2008).

26.   Johnston, C. A., Hirono, K., Prehoda, K. E. & Doe, C. Q. Identification of an Aurora-
A/PinsLINKER/ Dlg Spindle Orientation Pathway using Induced Cell Polarity in S2 Cells.
Cell **138**, 1150–1163 (2009).

27.   Lee, C.-Y. *et al.* Drosophila Aurora-A kinase inhibits neuroblast self-renewal by
regulating aPKC/Numb cortical polarity and spindle orientation. *Genes Dev.* **20**, 3464–3474
(2006).

28.   Joukov, V. & Nicolo, A. D. Aurora-PLK1 cascades as key signaling modules in the
regulation of mitosis. *Sci. Signal.* 26 (2018).

29.   Kachaner, D. *et al.* Coupling of Polo kinase activation to nuclear localization by a
bifunctional NLS is required during mitotic entry. *Nat. Commun.* **8**, (2017).

30.   Foley, E. A., Maldonado, M. & Kapoor, T. M. Formation of stable attachments
between kinetochores and microtubules depends on the B56-PP2A phosphatase. *Nat. Cell*
Biol. **13**, (2011).

31.   Beck, J. *et al.* Ubiquitylation-dependent localization of PLK1 in mitosis. *Nat. Cell*
Biol. **15**, (2013).

32.   Paschal, C. R., Maciejowski, J. & Jallepalli, P. V. A stringent requirement for Plk1
T210 phosphorylation during K-fiber assembly and chromosome congression. *Chromosoma*
121, (2012).

33.   Barbosa, J. *et al.* Polo regulates Spindly to prevent premature stabilization of
kinetochore–microtubule attachments. *EMBO J.* **39**, (2020).

34.   Liu, D., Davydenko, O. & Lampson, M. A. Polo-like kinase-1 regulates kinetochore–
microtubule dynamics and spindle checkpoint silencing. *J. Cell Biol.* **198**, 491–499 (2012).

35.   Mathieu, J. *et al.* Aurora B and Cyclin B Have Opposite Effects on the Timing of
Cytokinesis Abscission in Drosophila Germ Cells and in Vertebrate Somatic Cells. *Dev. Cell*
26, 250–265 (2013).

36.   Caussinus, E. & Gonzalez, C. Induction of tumor growth by altered stem-cell
asymmetric division in Drosophila melanogaster. *Nat. Genet.* **37**, 1125–1129 (2005).

37.   Castellanos, E., Dominguez, P. & Gonzalez, C. Centrosome Dysfunction in
Drosophila Neural Stem Cells Causes Tumors that Are Not Due to Genome Instability. *Curr.*
Biol. **18**, 1209–1214 (2008).

38.   Corbett, K. D. Molecular Mechanisms of Spindle Assembly Checkpoint Activation
and Silencing. in *Centromeres and Kinetochores* (ed. Black, B. E.) vol. 56 429–455 (Springer
International Publishing, 2017).

39.   Jia, L., Li, B. & Yu, H. The Bub1–Plk1 kinase complex promotes spindle checkpoint
signalling through Cdc20 phosphorylation. *Nat. Commun.* **7**, 10818 (2016).

40.   Conduit, P. T. *et al.* The Centrosome-Specific Phosphorylation of Cnn by Polo/Plk1
Drives Cnn Scaffold Assembly and Centrosome Maturation. *Dev. Cell* **28**, 659–669 (2014).

41.   Magnaghi-Jaulin, L., Eot-Houllier, G., Gallaud, E. & Giet, R. Aurora A Protein
Kinase: To the Centrosome and Beyond. *Biomolecules* **9**, 28 (2019).
341
342

344   **Figure 1: *polo* mutant NSCs display a SAC-independent mitotic delay.**

Figure 1

A) Polo, Mad2 and Actin Western blot analyses in *polo* and *polo,mad2^P* mutant brain extracts.

The genotypes are indicated for each lane. *polo^1/polo^{10}* and *polo^1/polo^9* are weak hypomorphs

and *polo^9/polo^{10}* is a strong hypomorph. **B)** Time-lapse imaging of mitosis in *polo* mutants in

the absence or presence of Mad2. Selected image series of dividing NSCs of the indicated

genotypes expressing mCherry::Tubulin (from top to bottom). The white dashed line outlines

the dividing NSCs. Scale bar: 5 μm. Time is min:s (t=00:00 is NEBD). **C)** Quantification of

time in mitosis from NEBD to anaphase onset for NSCs of the indicated genotypes. Means

are shown as black bars. Mann-Whitney unpaired test: ns: *p*>0.05; *: *p*<0.05; ****: *p*<0.0001.

D) GFP::Mad2 kinetochore localization in control, colchicine-treated, and *polo* mutant NCSs

expressing His2Av::mRFP1. Colchicine treatment (20μM) was used as a control for

GFP::Mad2 accumulation on kinetochores. Yellow arrowheads indicate the last kinetochore

labelled with GFP::Mad2 before anaphase onset (AO). Scale bar: 5 μm. Time is min:s

(t=00:00 is the beginning of the experiment). **E)** Quantification of the time between SAC

satisfaction (disappearance of the last GFP::Mad2-labeled kinetochores) and anaphase onset

in NSCs of the indicated genotypes. Means are shown as black bars. Mann-Whitney unpaired

test: ****: *p*<0.0001. Numerical data (mean, SD and *n*) are available in Table S1.

363 **Figure 2: Loss or excess of Polo activity is not compatible with normal mitosis duration**

364 **and *Drosophila* viability.**

365

<p style="text-align:center">**Figure 2**</p>

366 **A)** Polo and Actin Western blot analyses showing the expression of endogenous Polo kinase

367 in control and GFP-tagged Polo variants (WT, T182D, T182A and K54M) expressed

368 ubiquitously in *polo* mutant brains. The genotypes are indicated for each lane. Actin was used

369 as a loading control. **B)** Hatching rates of *polo* mutants expressing the indicated Polo variants.

370 **C)** Time-lapse imaging of mitosis in *polo* mutant NSCs expressing the indicated Polo variants

371 and mCherry::Tubulin. The white dashed lines outline the dividing NSCs. Scale bar: 5 μm.

372 Time is min:s (t=00:00 is NEBD). **D-E)** Quantification of the time in mitosis of NSCs of the

373 indicated genotypes. Means are shown as black bars. Mann-Whitney unpaired test: ns:

374 p>0.05; *: *p*<0.05; ****: *p*<0.0001. Numerical data (mean, SD and *n*) are available in Table

375 S1.

376 **Figure 3: Mitotic delays in *aurA* and *aurB* mutants are Polo-dependent.**

Figure 3

A, C) Time-lapse imaging of mitosis in *aurA* and *aurB/ial* mutant NSCs expressing the

indicated Polo variants and mCherry::Tubulin. The white dashed lines outline the dividing

NSCs. Scale bar: 5 μm. Time is min:s (t=00:00 is NEBD). **B, D)** Quantification of time in

mitosis for the indicated genotypes. Means are shown as black bars. Mann-Whitney unpaired

tests: ns: *p*>0.05; *: *p*<0.05; **: *p*<0.01; ****: *p*<0.0001. Numerical data (mean, SD and *n*)

are available in Table S1.

Figure 4: Impairing the SAC in *polo* mutants induces aneuploidy and promotes tumor

growth.

Figure 4

A) Representative images of larval brain lobes of the indicated genotypes stained for the NSC

marker Deadpan. The white dashed lines outline the central brain area where NSCs were

scored. Note the reduced brain lobe size in the *polo$^{9/10}$* mutant. Scale bar: 100 μm. **B)**

Quantifications of Deadpan-positive NSCs per brain lobe for the indicated genotypes. Mann-

Whitney unpaired test: *: $p<0.05$; ****: $p<0.0001$. Means are shown in black. **C)**

Representative images of the three classes of mitotic figures observed from squashed brains

stained with DAPI: euploid (8 chromosomes), aneuploid (more or less than 8 chromosomes)

and polyploid (multiple of 8 chromosomes) cells. Scale bar: 10 μm. **D)** Percentage of mitotic

brain cells with ploidy defects (aneuploidy and polyploidy) for the indicated genotypes. For

statistical analyses, the proportion of euploidy was compared to the proportion of ploidy

defects. Fisher's exact test: ns: $p>0.05$; *: $p<0.05$; ****: $p<0.0001$. **E)** Examples of wild-type

adult host flies 30 and 15 days after transplantation with control and *polo* mutant brain lobe

expressing His2Av::GFP as a readout for tumor development (bottom). Asterisks indicate

transplantation scars. **F)** Proportion of host flies developing tumors following the

transplantation of brain lobes from the indicated genotypes. Fisher's exact test; ns: $p>0.05$; *:

$p<0.05$; ***: $p<0.001$. Numerical data (mean, SD and *n*) are available in Table S1.

Figure 5: Activation of Polo restores NSCs polarization, spindle orientation and inhibits

tumor growth in an *aurA* mutant.

Figure 5

A) Representative images of fixed control and *aurA* mutant larval brain lobes with or without

Polo^{T182D}::GFP expression and stained for the NSC marker Deadpan. The white dashed lines

outline the central brain area where NSCs were scored. Note that the expression Polo^{T182D} in

aurA mutant restores the delimitation between the central brain and neuroepithelium. Scale

bar: 100 μm. **B)** Quantification of the NSC number in each brain lobe for the indicated

genotypes. **C)** Examples of wild-type adult host flies 15 and 30 days after transplantation with

aurA mutant or *aurA* mutant expressing Polo^{T182D}::GFP brain lobes expressing

mCherry::Tubulin as a readout to follow tumor development. Red asterisks indicate

transplantation scars. **D)** Percentage of host flies developing tumors 15 and 30 days after the

transplantation of brain lobes from the indicated genotypes. **E)** Quantification of ploidy

defects scored in neural tissue for the indicated genotypes. **F)** Representative images of

control, Polo^{T182D}::GFP, *aurA* mutant and Polo^{T182D}::GFP;*aurA* metaphase NSCs stained for

Miranda (magenta in the merge), aPKC and Phospho-Histone H3 (green in the merge), and α-

Tubulin (white in the merge). Yellow arrowheads indicate the edges of aPKC crescent. Scale

bar: 5 μm. **G)** Schematic representation of aPKC localization patterns observed in NSCs: left:

aPKC forms a crescent which defines the apical cortex of the control cell; right: aPKC is

unpolarized and exhibits an homogenous cytoplasmic and cortical localization. **H)** Percentage

of NSCs of the indicated genotypes with either a polarized apical crescent (light grey) or a

homogenous cortical and cytoplasmic aPKC localization (dark grey). **I)** Schematic illustration

of the method used to measure the spindle alignment α angle (orange) between the polarity

axis (black dashed line), determined by aPKC (green) and Miranda (magenta), and the spindle

axis (blue dashed line) determined by spindle poles (See Methods). **J)** Quantification of the

α angle between polarity and spindle axes in metaphase NSCs for the indicated genotypes.

Mann-Whitney unpaired test: ns: $p>0.05$; ****: $p<0.0001$. Means are shown as black bars.

For panels D), E) and H): Fisher's exact test: ns: $p>0.05$; *: $p<0.05$, ***: $p<0.001$, ****:

$p<0.0001$. Note that in panel E, the proportion of euploidy was compared to the proportion of

ploidy defects for statistical analyses. Numerical data (mean, SD and *n*) are available in Table

S1.

Materials and Methods

Fly strains and alleles combinations

The following strains were used for this study:

UAS-Cherry::Tubulin (BDSC #25774), Ubi-RFP::Tubulin (gift from Renata Basto), UASt-

mCherry::Jupiter (gift from Clemens Cabernard), UAS, His2Av::mRFP1 (BDSC #23651),

His2Av::GFP[1] and GFP::Mad2[2] were used to monitor by live-imaging the microtubule

network, chromosomes and unattached kinetochores. Wild-type (UASp-Polo^{WT}::GFP),

phosphomutant (UASp-Polo^{T182A}::GFP), phosphomimic (UASp-Polo^{T182D}::GFP) and kinase-

dead (UASp-Polo^{K54M}::GFP) Polo variants tagged with GFP were expressed using the Gal4-

UAS system. All four constructs were inserted on the second chromosome (attP40, see below)

and on the third chromosome (attP154,[3]).

Inscuteable-Gal4 (Insc-Gal4; BDSC #8751) and Worniu-Gal4 (Wor-Gal4; BDSC #56554)

allowed the expression of UAS driven fluorescent markers in the neuroblasts; and Actin5C-

Gal4 (BDSC #4414) was used for ubiquitous expression in somatic cells. The expression of

Polo variants for protein level and rescue assays in Figure 2A and 2B as well as the

expression of Polo^{T182D}::GFP for ploidy analysis (Figure 5E) were performed using the

Actin5C-Gal4 driver. The expression of mCherry::Jupiter and Polo^{T182D}::GFP in *ial* mutant

background were driven by Wor-Gal4. For other experiments (live-imaging,

immunofluorescence and transplantation assay), Polo variants and the microtubule marker

were expressed under the control of Insc-Gal4.

The following alleles were used in this study and described before: *aurA^{8839}* (referred as *aurA*

mutant, [4,5]), *mad2^p* ([6]), *polo^1* ([7]), *polo^9* and *polo^{10}* ([8]), *ial^{1689}*, *ial$^{35.33}$* and *ial^{2443}* ([9]). The allelic

combination *polo^9/polo^{10}* (referred as *polo* mutant) corresponds to a strong *polo* hypomorph

mutant and the flies do not reach the pupal stage, as described before[10]. The allelic

combinations *polo^1/polo^9* and *polo^1/polo^{10}* mutant flies are viable and corresponds to weak

polo hypomorph mutants[10]. The phenotype of Ial/Aurora B loss-of-function was assessed

during the larval stage using two heteroallelic combinations: $ial^{35.33}/ial^{1689}$ and ial^{2A43}/ial^{1689}.

Flies were maintained and crossed under standard conditions at 25°C.

Generation of transgenes

The constructs containing the coding sequence of Polo variants (WT, T182A, T182D and

K54M) C-terminally tagged with GFP were cloned into the plasmid pUAS-K10-attB[3] and

were injected in embryos for targeted insertion on the attP40 site located on second

chromosome (BestGene).

Immunohistochemistry and antibodies

Third instar larval brains (120 to 144 h after egg lying depending on overall growth delay)

were dissected in Schneider medium supplemented with 10% FCS and fixed in Testis Buffer

(183 mM KCl, 47 mM NaCl, 10 mM Tris, and 1 mM EDTA, pH 6.8) containing 10%

formaldehyde and 0.01% Triton X-100 at 25°C for 20 minutes, as described in a previous

study[11].

Primary antibodies: monoclonal anti-α-tubulin DM1A (mouse, 1:500, Sigma T6199), anti-

PKCζ (C-20) (rabbit, 1:250, Santa Cruz sc-216), anti-Miranda (rat, 1:200, Abcam 197788),

polyclonal anti-phosphorylated Ser10 of Histone H3 (rabbit, 1:500, Millipore 06-570) and

anti-Deadpan (rat, 1:200, Abcam 195172).

Secondary antibodies: anti-rat, anti-mouse and anti-rabbit conjugated to 488, 568, 647 Alexa-

Fluor (1:500 to 1:1000, Life Technologies).

Images were acquired on a Leica SP5 confocal microscope using 40X (1.25 N.A.) and 63X

(1.40 N.A.) objectives, controlled with Leica LAS software and analyzed on Fiji or Imaris.

Western blotting and antibodies

Two to three brains were dissected, resuspended in Laemmli Buffer and loaded on a 4-15%

Mini-PROTEAN TGX Precast Protein Gel (BioRad). After migration, proteins are transferred

on a nitrocellulose membrane using a Trans-Blot Turbo (BioRad). Membranes were washed

with PBS+0.1% Tween20 (PBST) and blocked with PBST+5 to 10% powder milk. The

membranes were then incubated in the following primary antibodies, diluted in TBST+2.5 to

5% powder milk: Rabbit polyclonal anti-Actin (I-19, Santa Cruz sc-1616-R; 1/4000), anti-

Mad2 (Gift from David Sharp; 1/500), mouse monoclonal anti-Polo MA294 (Gift from David

Glover; 1/100). Anti-mouse and anti-Rabbit secondary antibodies were purchased from

Jackson ImmunoResearch. Western blots were revealed with SuperSignal West Dura or

Femto (ThermoFisher).

Live cell imaging

Third instar larval brains (120 to 144h after egg lying depending on overall growth delay)

were dissected in Schneider medium containing 10% FCS and transferred to 50 μL wells

(Ibidi, μ-Slide Angiogenesis) for live imaging. Mutant and control brains were imaged in

parallel at 25°C. Z-series (thickness of 20 μm with 1 μm spacing) were acquired with a

temporal resolution of 30 to 90 s for 1 to 2.5 hours. Alternatively, samples were mounted on a

stainless-steel slide, between coverslip and mineral oil as described in a previous study[11].

Images were acquired with a spinning disk system consisting of a DMi8 microscope (Leica)

equipped with a 63X (1.4 N.A.) oil objective, a CSU-X1 spinning disk unit (Yokogawa) and

an Evolve EMCCD camera (Photometrics). The microscope was controlled by the Inscoper

Imaging Suite and the dedicated software (Inscoper). Alternatively, a CSU-X1 spinning-disk

unit mounted on an inverted microscope (Elipse Ti; Nikon) equipped with a 60X (1.4 N.A.)

oil objective, a sCMOS ORCA Flash 4.0 (Hamamatsu) and controlled by MetaMorph; was

also used for some experiments.

Images were processed with Fiji or Imaris softwares.

Time in mitosis and spindle orientation angle measurements

The time in mitosis was determined by the duration between NEBD (tubulin entry in the

nuclear space) and anaphase onset. Note that only the cells undergoing complete mitosis

within the span of the experiments were scored. For *ial;aurA* double mutant (Supplementary

Figure 2B-C) displaying the most severe phenotype, the cells could not complete mitosis

throughout the experiment (2.5 h) and the time in mitosis could not be measured. Spindle

orientation on fixed tissue was measured on metaphase cells using the "spot" tool of Imaris.

Briefly, two spots were placed in the center of the apical aPKC and basal Miranda crescents

to determine the polarity axis; two spots positioned at the spindle poles determined the mitotic

spindle axis. For *aurA* mutant that did not exhibit an apical aPKC crescent, the polarity axis

was determined by a straight line between the center of the Miranda crescent and a second

spot at the opposite cortex. The 3D coordinates of the spots were used to calculate polarity

and spindle axes vectors and the angle between them as described in [12].

Ploidy analysis in brains

Third instar larval brains were dissected in PBS, incubated for 8 minutes in 0.5% sodium

citrate solution, fixed in acetic acid (45% for 45 seconds and 60% for 2 minutes), squashed

between slide and coverslip and flash-frozen in liquid nitrogen (Caous et al., 2015). After

coverslip flip off, the squashed cells on the slide were washed in PBS and mounted in

ProlonGold supplemented with DAPI. Images were acquired using a Leica DMRXA2

microscope equipped a 63X (1.32 N.A.) objective, a CoolSnap HQ2 camera and controlled by

MetaMorph. Images were processed with Fiji software.

Tumor transplantation assay

The protocol to assess proliferation ability of brain tissues was originally set up in the lab of

Cayetano Gonzalez[13]. Briefly, a central brain fragment (expressing a fluorescent probe) was

transplanted in the abdomen of a live wild-type adult female fly and the growth of the

transplant was monitored over time. Control or mutant third instar larval brains, expressing

either His2Av::GFP or UAS-Cherry::Tubulin under the control of Insc-Gal4 were dissected in

sterile PBS (Sigma). One brain lobe was transplanted in the abdomen of a wild-type adult fly

using a pulled capillary with a beveled tip (around 150 μm diameter) adjusted to a

microinjection system (IM-9B; Narishige). After transplantation, host flies were maintained at

18°C overnight and transferred at 25°C for a month. Tumor growth was monitored every one

or two days upon 30 days for the appearance of a mCherry or GFP signal in the abdomen. For

aurA and strong *polo* mutants, transplant growth was assessed over a 15-days period because

these tumors were particularly aggressive and killed the host flies before 30 days. The

neuroblast-specific Insc-Gal4 was used to assay the effect of Polo^{T182D}::GFP variant for tumor

growth.

Statistical analyses

Sample size and statistical test used for each analysis are found in the corresponding figure

legends. Statistical analyses were performed using the Prism software (Version 7, GraphPad).

Two tests were used according to the data: to compare ranks between two unpaired samples

with variable variances and non-Gaussian distributions, a nonparametric Mann-Whitney test

was used; while the comparisons of proportions were assessed by a two-sided Fisher's exact

test with a confidence index of 95%. Significance are displayed on the figure: ns: $p>0.05$; *:

$p<0.05$; **: $p<0.01$; ***: $p<0.001$; ****: $p<0.0001$.

References

1. Clarkson, M. & Saint, R. A His2AvDGFP Fusion Gene Complements a Lethal His2AvD
Mutant Allele and Provides an in Vivo Marker for Drosophila Chromosome Behavior.
DNA Cell Biol. **18**, 457–462 (1999).
2. Buffin, E., Lefebvre, C., Huang, J., Gagou, M. E. & Karess, R. E. Recruitment of Mad2 to
the Kinetochore Requires the Rod/Zw10 Complex. *Curr. Biol.* **15**, 856–861 (2005).
3. Kachaner, D. *et al.* Coupling of Polo kinase activation to nuclear localization by a
bifunctional NLS is required during mitotic entry. *Nat. Commun.* **8**, (2017).
4. Wang, H. *et al.* Aurora-A acts as a tumor suppressor and regulates self-renewal of
Drosophila neuroblasts. *Genes Dev.* **20**, 3453–3463 (2006).

5. Lee, C.-Y. *et al.* Drosophila Aurora-A kinase inhibits neuroblast self-renewal by
regulating aPKC/Numb cortical polarity and spindle orientation. *Genes Dev.* **20**, 3464–
3474 (2006).
6. Buffin, E., Emre, D. & Karess, R. E. Flies without a spindle checkpoint. *Nat. Cell Biol.* **9**,
565–572 (2007).
7. Sunkel, C. E. & Glover, D. M. polo, a mitotic mutant of Drosophila displaying abnormal
spindle poles. *J. Cell Sci.* **89**, (1988).
8. Donaldson, M. M., Tavares, Á. A. M., Ohkura, H., Deak, P. & Glover, D. M. Metaphase
Arrest with Centromere Separation in polo Mutants of Drosophila. *J. Cell Biol.* **153**, 663–
676 (2001).
9. Mathieu, J. *et al.* Aurora B and Cyclin B Have Opposite Effects on the Timing of
Cytokinesis Abscission in Drosophila Germ Cells and in Vertebrate Somatic Cells. *Dev.*
Cell **26**, 250–265 (2013).
10. Carmena, M., Lombardia, M. O., Ogawa, H. & Earnshaw, W. C. Polo kinase regulates the
localization and activity of the chromosomal passenger complex in meiosis and mitosis in
Drosophila melanogaster. *Open Biol.* **4**, 140162 (2014).
11. Gallaud, E. *et al.* Ensconsin/Map7 promotes microtubule growth and centrosome
separation in Drosophila neural stem cells. *J. Cell Biol.* **204**, 1111–1121 (2014).
12. Loyer, N. The last-born daughter cell contributes to division orientation of Drosophila
larval neuroblasts. *Nat. Commun.* 12 (2018).
13. Caussinus, E. & Gonzalez, C. Induction of tumor growth by altered stem-cell asymmetric
division in Drosophila melanogaster. *Nat. Genet.* **37**, 1125–1129 (2005).
151