The impact of COVID-19 vaccination campaigns accounting for antibody-dependent enhancement

Nessma Adil M. Y.[1,2❍], H. Christian Jr. Tsoungui Obama[1,2❍], Jordan Y. Ngucho[2❍], S. Frank Kwamou[1,2], Loyce Kayanula[1,2], George Kamanga[1,2], Toheeb B. Ibrahim[1,2], Patience Bwanu Iliya[1,2], Sulyman Iyanda[1,2], Looli Alawam[1,2], Miranda I. Teboh-Ewungkem[3], Kristan A. Schneider[1❍*]

1 Department of Applied Computer- and Biosciences, University of Applied Sciences Mittweida, Mittweida, Germany
2 African Institute for Mathematical Sciences Cameroon, Limbe, Cameroon
3 Department of Mathematics, Lehigh University, Bethlehem, PA, USA

❍These authors contributed equally to this work.
*kristan.schneider@hs-mittweida.de

Abstract

Background: COVID-19 vaccines are approved, vaccination campaigns are launched, and worldwide return to normality seems within close reach. Nevertheless, concerns about the safety of COVID-19 vaccines arose, due to their fast emergency approval. In fact, the problem of antibody-dependent enhancement was raised in the context of COVID-19 vaccines.

Methods and findings: We introduce a complex extension of the model underlying the pandemic preparedness tool CovidSim 1.1 (http://covidsim.eu/) to optimize vaccination strategies with regard to the onset of campaigns, vaccination coverage, vaccination schedules, vaccination rates, and efficiency of vaccines. Vaccines are not assumed to immunize perfectly. Some individuals fail to immunize, some reach only partial immunity, and – importantly – some develop antibody-dependent enhancement, which increases the likelihood of developing symptomatic and severe episodes (associated with higher case fatality) upon infection. Only a fraction of the population will be vaccinated, reflecting vaccination hesitancy or contraindications.

We parameterized the model to reflect the situation in Germany and predict increasing incidence (and prevalence) in early 2021 followed by a decline by summer. Assuming contact reductions (curfews, social distancing, etc.) to be lifted in summer, disease incidence will peak again. Fast vaccine deployment contributes to reduce disease incidence in the first quarter of 2021, and delay the epidemic outbreak after the summer season. Higher vaccination coverage results in a delayed and reduced epidemic peak. A coverage of $75\% - 80\%$ is necessary to prevent an epidemic peak without further drastic contact reductions.

Conclusions: With the vaccine becoming available, compliance with contact reductions is likely to fade. To prevent further economic damage from COVID-19, high levels of immunization need to be reached before next year's flu season, and vaccination strategies and disease management need to be flexibly adjusted. The predictive model can serve as a refined decision support tool for COVID-19 management.

Introduction

Interventions such as curfews, lockdowns, cancellations of mass events, in response to the COVID-19 pandemic caused massive losses in revenue for whole economic sectors [1], [2]. They were justified as legitimate measures to delay the epidemic to increase healthcare capacities, develop effective treatments, and vaccines to immunize the population. Despite all efforts, the spread of COVID-19 since fall 2020 drives healthcare systems to their limits in Europe and North America, underlining the urge for an effective and safe vaccine.

Governments across the globe were ambitious in facilitating SARS-CoV-2 vaccine development, by large-scale programs such as *Operation Warp Speed* launched by the US Government [3]. Currently, more than 227 vaccine-development projects against SARS-CoV-2 are ongoing [4]. There are four major vaccination platforms to stimulate antibody production triggered by the SARS-CoV-2 spike protein: (i) viral vectors fused with a gene that encodes for the SARS-CoV-2 spike protein; (ii) inactivated SARS-CoV-2 variants; (iii) protein subunits of SARS-CoV-2 antigens; and (iv) a rather new technique, where lipid nanoparticles encapsulate nucleoside-modified mRNA (modRNA) encoding mutated forms of the SARS-CoV-2 spike protein. The most promising candidates typically follow a 2-3 week vaccination schedule, after whose completion the protective effect is reached within 2-3 weeks [5], [6].

Russia was ambitious to release the world's first SARS-CoV-2 vaccine *Gam-COVID-Vac*. In homage named *Sputnik V*, the vaccine is based on two human adenovirus (common cold) vectors [7], costs less than 20 USD per dose, and is highly controversial due to its fast emergency use authorization (EUA) without a phase III study [8]. Nevertheless, mass vaccination started in Russia on December 5, 2020. The modified chimpanzee adenovirus vector-based candidate from AstraZeneca, *AZD1222*, is currently under phase III study, will cost only 4 USD per dose, and has a capacity of 400 million doses for Europe and 300 million doses for the USA.

China gave EUA to two vaccines that trigger an immune response by inactivated SARS-CoV-2 variants. *BBIBP-CorV* has a capacity of 1 billion doses for China in 2021 at a cost of less than 75 USD per dose and was fully approved, while *CoronaVac* costs 30 USD per dose. *Covaxin (BBV152)* is a cheap vaccine (1.36 USD per dose) currently under EUA developed by the Indian Council of Medical Research, based on inactivated SARS-CoV-2 variants.

NVX-CoV2373 by Novavax, seeking approval in Mexico, is a vaccine that uses SARS-CoV-2 recombinant spike protein nanoparticles with adjuvants to trigger an immune response [9].

Two modRNA-based candidates are currently in phase III studies, which either seek approval or were granted EUA. *Tozinameran (BNT162b2)* by BioNTech (20 USD per dose), was approved in Canada and Europe, and received EUA in the UK and the USA. A second modRNA-based candidate, *mRNA-1273* by Moderna, is currently in phase III trials, and received EUA in Canada and the USA.

Vaccination campaigns aim for herd immunity. There is an ongoing debate on the optimal deployment of the vaccine. Some countries have ambitious deployment strategies, e.g., Morocco plans to immunize up to 80% of the population. Globally the trend is to deploy vaccines voluntarily and free of charge, with a general agreement to prioritize vulnerable risk groups (e.g., senior citizens, people with co-morbidities, etc.) and people of systemic importance (e.g., healthcare workers, police, public services) before making the vaccine available to the general public [10]. Incentives for getting voluntary vaccines have been proposed, e.g., recently Qantas airlines announced to make the vaccine mandatory for their passengers [11, 12].

Nevertheless, skepticism about vaccines and their potential side effects are widespread, resulting in vaccine hesitancy [13]. One of the potentially negative effects of

a vaccine is the occurrence of antibody-dependent enhancement (ADE) or, more general, enhanced respiratory disease (ERD) [14, 15]. ADE is best understood in Dengue fever and was observed also in SARS-CoV and MERS-CoV both in vitro and in vivo [16]. In SARS-CoV-2, ADE occurs most likely via enhanced immune activation [17]. Here, sub-optimal antibodies form immune complexes with the virus that deposit into airway tissues and activate cytokine and complement pathways. This triggers inflammation, airway obstruction, and even acute respiratory distress syndrome [17]. By this mechanism, vaccines could potentially result in more severe symptoms upon infection with SARS-CoV-2.

Here, we employ predictive modelling to explore the outcome of vaccination strategies on the COVID-19 pandemic. We explore the influence of (i) the vaccination coverage reflecting vaccine hesitancy, the commonness of contraindications, and access to the vaccine; (ii) the vaccination rate, reflecting supply and infrastructure; and (iii) the immunizing effect on disease incidence, prevalence, and mortality. We further investigate the impact of the launch of the vaccination campaign relative to the epidemic peak. Intentionally, our model accounts for the occurrence of ADE, ERD, and other deleterious side effects of the vaccine – subsumed here as ADE. By this approach, we seek to address the following questions: What is the benefit of launching the vaccination campaign early (at full scale) to substantially reduce disease incidence? Can side effects of the vaccine result in overall higher mortality than the virus itself? How fast should the vaccine be deployed? Which vaccination coverage should be aimed at?

We use an extension of the SEIR model underlying the pandemic preparedness tool CovidSim 1.1 [18] to predict the outcome of vaccination campaigns. The model incorporates general contact reduction (hard and soft lockdowns, social distancing, etc.) and case isolation in a time-dependent manner. Unvaccinable persons, summarizing anti-vaxxers, individuals with contraindications, and individuals that do not have access to the vaccine, are properly addressed. Furthermore, immunization after receiving the vaccine does not occur instantaneously, reflecting vaccination schedules and immunogenicity. Importantly, our model allows individuals to be vaccinated during the infection (if it is asymptomatic and undetected) and to be infected before the outcome of the vaccination manifests. As an example, we use model parameters that reflect the situation in the Federal Republic of Germany. In the main text, we present only a verbal description of the model and refer to S1 Appendix for a concise formal description, dedicated to readers interested in the technical aspects of the model.

Methods

We model the occurrence of ADE during vaccination campaigns in the ongoing COVID-19 pandemic by an extended SEIR model. More precisely, we generalize the compartmental model underlying the pandemic preparedness tool CovidSim (cf. [18]), which is formulated as a system of ordinary differential equations. Modelling ADE requires a substantial extension of the original model. We describe it verbally in a simplified form here and refer readers interested in a concise description to S1 Appendix.

Course of the disease

We divide a population of size N into susceptible, infected, and recovered individuals, each of which is subdivided into numerous compartments. During the course of the infection (Fig 1), individuals pass through the (i) latent phase (no symptoms, not yet infectious); (ii) prodromal phase (no symptoms, infectious but not yet to the fullest extent); (iii) fully infectious phase (symptoms might start, infectious to the fullest extent); (iv) late infectious phase (infectiousness is reduced). The fully infectious and

late infectious phases can be asymptomatic or symptomatic with mild or severe symptoms. At the beginning of the fully infectious phase a fraction of infections becomes symptomatic (mild or severe symptoms), whereas the remaining fraction remains asymptomatic. At the end of the late infectious phase individuals either recover and obtain full immunity or they die. Only symptomatic infections can result in death, asymptomatic infections are not lethal (except individuals get vaccinated during the infection and this results in complications).

Fig 1. Phases of the infection. Schematic representation of the disease phases and at which stages individuals become invective and potentially symptomatic.

Susceptible individuals

Susceptible individuals are sub-divided (see Fig 2) into those that are: (i) not vaccinable because they do not have access, refuse to be vaccinated, or cannot be vaccinated for medical reasons (e.g., allergies, or the vaccine is not approved for them); (ii) waiting to be vaccinated; (iii) already vaccinated, but the outcome of the vaccination is still pending; (iv) already vaccinated, but only partially immunized; (v) vaccinated, but the vaccination completely failed to immunize; (vi) vaccinated, but the vaccine caused ADE. Vaccinable susceptibles have to wait before being vaccinated. The outcome of the vaccination does not occur immediately but after some waiting time (pending outcome). Vaccination either results in (i) full immunity, (ii) partial immunity, (iii) no immunity (the vaccine had no effect), or (iv) ADE (the vaccine had a deleterious effect). Partial immunity gives some protection from infection and manifests in a lower likelihood to develop symptoms upon infection. ADE also protects partially from infections, but upon infection increases the likelihood of severe disease (and death).

Fig 2. Groups of susceptible and immunized individuals. Schematic representation of different compartments of susceptible individuals. Unvaccinated susceptibles are either unvaccinable (NV) or wait to be vaccinated (U). Vaccinated susceptibles (V) remain susceptible immediately after vaccination. After the vaccination outcome manifested, they remain susceptible at different levels if immunization failed (NI), only partial immunization was achieved (PI), or they developed ADE. Individuals that are successfully immunized are no longer susceptible (R). Arrows show how individuals move between groups. All susceptibles can be infected (not indicated).

Effect of vaccination

Individuals are vaccinated only once (one vaccination cycle, subsuming all necessary doses). The waiting time during which immunization is pending reflects the waiting time for the vaccination cycle to be completed (it can be one or several doses). In the model we need to distinguish between infections of individuals that are (i) partially immunized, (ii) not immunized or unvaccinable, (iii) developed ADE, and (iv) vaccinable, but still waiting to be vaccinated. Vaccinable individuals (those that wait for vaccination) can also be vaccinated when they are already infected (i.e., during the latent, and prodromal phases, and during the fully infectious and late infectious phases if the infections remain asymptomatic and undetected; see Figure 3). After some waiting period (during which the effect of the vaccination is pending) these individuals are either successfully, partially, or not immunized, or they might be affected by ADE. The likelihood of these outcomes depends on the phase of the infection. Particular consideration is given to individuals that are vaccinated during the fully infectious or late infectious phase. Namely, when the disease has already progressed, the effect of the vaccine potentially changes.

Fig 3. Simplified model flow. Susceptible individuals are infected and ultimately recover or die. Individuals are infected either after or before the outcome of the vaccination is determined. In the latter case the outcome of the vaccine is not determined upon infection (they might get vaccinated before or during the infection), or they are not vaccinated before or during the infection (this includes unvaccinable individuals that get infected).

Case isolation

A fraction of symptomatic infections seeks medical help and will be isolated in quarantine wards, until the wards are full, in which case they are sent in home isolation. Quarantine wards guarantee perfect isolation, whereas home isolation does not eliminate all infectious contacts with the isolated individuals.

General contact reduction

During designated time intervals general contact reduction (curfews, social distancing, cancellation of mass events, etc.) is sustained. These phases reduce the number of contacts between individuals in the population. These contact reductions are time dependent. For the simulations here we assume an initial "hard lockdown" followed by a phase of relief, a "soft lockdown", a second "hard lockdown", and a final relief phase before contact reductions are lifted.

Contact rate

Susceptible individuals get infected by random contacts with infected individuals that can transmit the disease and are not isolated. The contact rates are mediated by general contact reduction. The basic reproductive number R_0 is assumed to fluctuate seasonally.

Model implementation

The model as described in detail in S1 Appendix was implemented in Python 3.8. We used a 4th order Runge-Kutta method using the function *solve ivp* as part of the library Scipy. Graphical output was created in R [19].

Results

Here, we study the effectiveness of different hypothetical vaccination programs/strategies to immunize the population in the ongoing COVID-19 pandemic in terms of disease incidence (or prevalence) and overall mortality. Vaccination campaigns differ in (i) their onset and vaccination rate reflecting the available infrastructure (availability of the vaccine and medical infrastructure); (ii) the vaccination coverage, reflecting the willingness of the population to get vaccinated and access to medical care; and (iii) vaccination schedules, immunogenicity, and efficacy/effectiveness summarizing vaccine-specific properties. The ultimate goal of any vaccination campaign is to reach herd immunity, mitigate SARS-CoV-2, and "return to normality" as soon as possible. We explore how fast herd immunity is reached assuming that contact reductions cannot be sustained for too long. The approach is conservative, as we focus on the potential negative effects of the vaccine.

We report disease incidence and prevalence. Disease prevalence is defined as the sum of all infected individuals (in the latent, prodromal, fully infectious, and late infectious phases). Incidence, i.e., the number of new cases per time interval, is notoriously difficult to define in an SEIR-type model since individual infections are not modelled explicitly. We approximated incidence as prevalence divided by the duration of an

average disease episode during which the infection is detectable. More concretely, we defined incidence as prevalence divided by 10 days. Hence, incidence and prevalence differ only by a multiplicative factor here an can be used synonymously. Both are reported for the readers convenience to facilitate comparison with publicly accessible data.

Regarding the results, we use model parameters reflecting the situation in the Federal Republic of Germany that so far intervened successfully in the COVID-19 pandemic. The parameters used are listed in S1 Table - S5 Table. The population size of Germany was set to $N = 83$ million. We assumed the first COVID-19 cases were introduced in late February 2020 (corresponding to $t = 0$). A basic yearly average reproductive number of $\bar{R}_0 = 3.5$ was assumed, which fluctuates seasonally by 43% with a peak in late December ($t_{R_{0_{\max}}} = 300$). The average durations for the latent, prodromal, fully infectious and late infectious phases were set to $D_E = 3.7$, $D_P = 1$, $D_I = 5$, $D_L = 5$ days, respectively. Individuals in the prodromal and late infectious phases were half as infectious as in the fully infectious phase ($c_P = c_L = 0.5$). If individuals developed only partial immunity or ADE, they were assumed to be half as susceptible to SARS-CoV-2 than unimmunized individuals ($p_{PI} = p_{ADE} = 0.5$). A percentage $f_{Sick} = 58\%$ of infections became symptomatic if immunity was not mediated by the vaccine. This percentage decreased for partially immunized individuals and increased for those that developed ADE (see S4 Table). Regarding general contact reduction, we assumed a "hard lockdown" from late March ($t_{Dist_1} = 21$) to early May 2020 ($t_{Dist_2} = 75$) that reduced $p_{Dist_1} = 70\%$ of all contacts, followed by a period of relaxation until the end of October ($t_{Dist_3} = 246$) during which $p_{Dist_2} = 48\%$ of the contacts were avoided. The first "soft lockdown" from late October was sustained until the end of December ($t_{Dist_4} = 300$) with a contact reduction of $p_{Dist_3} = 60\%$. This was followed by a second "hard lockdown" from late December until late January ($t_{Dist_5} = 330$) and a phase of relief resulting in a $p_{Dist_4} = 60\%$ contact reduction sustained until June 2021 ($t_{Dist_6} = 450$), after which all general contact reductions are lifted reflecting the worst-case scenario, in which compliance with social distancing can no longer be sustained in the face of the vaccine becoming available.

The simulation results until late December ($t = 300$) match the disease incidence in Germany. Notably, only the number of confirmed, not of actual cases is known. The true incidence (which is modelled) is obviously unobservable and might be substantially higher than the confirmed incidence (true incidence multiplied with the probability of detection). Consequently, the incidences in Figs 4-9 exceed the number of reported cases in Germany - however, the numbers are plausible and match in the order of magnitude.

Regarding the vaccination campaigns, we use the following reference scenario: 60% of the population will get vaccinated, a vaccination rate of 1/180 (i.e., an average time of 180 days to get vaccinated), a launch of the campaign in early January 2021 ($t = 300$), and a vaccination schedule of 28 days (the vaccine becomes effective after 28 days).

No vaccination

We consider the situation in which no vaccination campaign (VC) is launched as a reference case for comparisons. Figs 4A-D show disease incidence, which matches the reported numbers in Germany. The base reproductive number is at the seasonal maximum at times $t = 300$ and $t = 665$. Although R_0 declines after $t = 300$, the end of the second "hard lockdown" in late January ($t = 330$) leads to an increase in incidence. In fact, it will more than double by the end of March 2021 ($t = 390$), after which it will decline due to seasonal reductions in R_0. After contact reductions are lifted at time $t = 450$, disease incidence would increase drastically and the epidemic peak would be reached around $t = 590$ (late October 2021) with almost 40% of the population being infected at this time point. This is a hypothetical worst-case scenario.

Fig 4. Onset of the vaccination campaign: Plots show (total) incidence, i.e. prevalence (all latent, prodromal, fully infectious, and late infectious individuals) divided by 10 (panels A-B), prevalence per 100 000 individuals (panels C-D), and mortality, i.e., the cumulative deaths (D), per 100 000 individuals (panels C-D) as a functions of time t for different onsets of the vaccination campaign (colors). In panels E and F the effect of increased ADE-induced mortality is shown (dashed lines $f_{\text{Dead}}^{(\text{ADE})} = 20\%$ *vs.* solid lines $f_{\text{Dead}} = f_{\text{Dead}}^{(\text{ADE})} = 7\%$). (Note the values of $f_{\text{Dead}}^{(\text{ADE})}$ only affects mortality, not incidence.) As a baseline comparison, the black lines show incidence and mortality in the absence of the vaccine. Seasonal fluctuations in R_0 are shown by the grey dashed lines corresponding to the y-axis on the right-hand side. Plot parameters are given in S1 Table - S3 Table

Onset of vaccination campaigns

The onset of the vaccination campaign (VC) has a profound effect on disease incidence (see Fig 4). The earlier the onset of the VC, the earlier and the stronger the reduction in incidence after the second "hard lockdown". Even if the VC starts at $t = 300$ incidence will increase after the second "hard lockdown". However, the increase is less than half of that observed without a VC. Launching the campaign in March 2021 ($t = 360$) hardly leads to a reduction in disease incidence until summer 2021 (Figs 4A, C). (The launch of the VC has to be interpreted as the time when it becomes fully effective.)

A much stronger effect of the VC's onset manifests after contact reduction is no longer sustained. High epidemic peaks will emerge at the end of 2021 or in early 2022. Later launches of the VC result in earlier and higher epidemic peaks (Figs 4B, D). The differences in height of the epidemic peaks are substantial in comparison to disease incidence in 2020. In any case, the benefit of the VCs is clear. The epidemic peaks will be substantially lower compared to the situation without a VC. The earlier the vaccination campaign starts, the higher the reduction in mortality. Indeed the differences by the end of June 2021 ($t = 450$) are visible (Fig 4E). The reduction in mortality by the end of March 2022 is substantial (Fig 4F).

Fraction of the population being vaccinated

Also, the proportion of the population being vaccinated has a profound effect on disease incidence and mortality. Even if only 25% of the population gets vaccinated, there is a substantial decrease in incidence by time $t = 450$ (June 2021). The higher the proportion of individuals being vaccinated, the stronger the reduction in incidence (Figs 5A, C). Incidence will not substantially decrease further if the proportion of the population being vaccinated exceeds 80%. This directly translates into mortality (Figs 5E, F).

It takes a vaccination coverage of about 70% to avoid a high epidemic peak after the contact reductions are lifted. The lower the proportion of the population being vaccinated, the earlier and higher will the epidemic peak be (Fig 5B). If 75% of the population gets vaccinated, an epidemic peak emerges in spring 2022 ($t = 750$) - notice this prediction assumes that no contact reducing interventions are in place after summer 2021 ($t = 450$). This peak will exceed the one in early 2021. If 80% of the population is vaccinated, a pandemic peak that is in between the first and second wave of 2020 emerges. This peak will be avoided if 85% of the population gets vaccinated.

Mortality substantially decreases as the proportion of the population getting vaccinated increases (Figs 5C, D). The reduction occurs in a nonlinear fashion and shrinks with higher proportions of the population being vaccinated.

Note that vaccination coverage is not the same as the herd immunity threshold. The latter is the percentage of the population that needs to be immune for the disease to vanish. The vaccination coverage is the the percentage of the population that will be vaccinated throughout the epidemic. A higher coverage reflects faster deployment of the

vaccine.

Fig 5. Vaccination coverage. As Fig 4, but for different vaccination coverage (colors).

Rate of vaccination

Not surprisingly, the faster the population is vaccinated the better. Assuming a realistic average waiting time to be vaccinated ranging between 180 - 300 days and 60% vaccination coverage, the effects are immediately visible in the short term (from $t = 300$ to $t = 450$, see Figs 6A, C). The benefit of vaccinating the population fast becomes substantial in the long term. In fact, the epidemic peak that emerges at around generation $t = 700$ (early 2022) is twice as high if the waiting time to get vaccinated is 300 days rather than 180 days on average (see Figs 6B, D). This translates directly into mortality (see Figs 6E, F).

Fig 6. Vaccination rate. As Fig 4, but for different vaccination rates (colors), i.e., average waiting times (shown in legend) to get vaccinated.

Vaccination schedule and immunogenicity

Another important factor in VCs is the time until immunization is reached, as determined by the vaccination schedule and immunogenicity. There is a visible effect if the time to immunization is increased from 28 days to 42 days. However, the effect is not as strong as that of the rate of infection or the proportion being vaccinated. In relative terms, the short-term effect is more pronounced (cf. Figs 7A with B and C with D). The reason is that the number of infections is rising at the onset of the vaccination campaign. During this period early immunization reduces the spread of COVID-19. Once incidence is low, the time to immunization is not as important until the final epidemic peak emerges. This will emerge earlier and will be higher if the time to immunization is longer (see Figs 7B, D). Again, incidence directly translates into mortality (see Figs 7E, F). The long-term effect on mortality of longer times to immunization is by far less pronounced than those of the vaccination rate and proportion of vaccinated individuals.

Fig 7. Vaccination schedule and time to immune response. As Fig 4, but for different average waiting times (D_A) until the vaccinations manifests its outcome (immunizing effect), corresponding to vaccination schedules and times to immune response.

Effectiveness of the vaccine

The better the immunizing effect of the vaccine, the more effective is the VC. We compared four scenarios summarized in Table 1, one being a worst-case scenario which is unrealistic and just used as a comparison. When effectiveness increases from 78% (low) to 95% (high) full immunization there will be a clear drop in disease incidence and mortality in the long run (see Figs 8B, D, F). The short-term effects are not as pronounced (see Figs 8A, C, E). The hypothetical worst-case scenario, in which the effectiveness is only 50%, leads to substantially higher infections and deaths.

Fig 8. Vaccine effectiveness. As Fig 4, but for different vaccination coverage and effectiveness (quality) of the vaccine as summarized in Table 1. No ADE-induced increased mortality is assumed.

Table 1. Vaccine effectiveness.

Effectiveness	$f_.^{(R)}$	$f_.^{(NI)}$	$f_.^{(PI)}$	$f_.^{(ADE)}$
High	0.94	0.02	0.03	0.01
Medium	0.90	0.04	0.05	0.01
Low	0.78	0.10	0.10	0.02
Poor	0.50	0.24	0.24	0.02

Parameter values corresponding to vaccine effectiveness, where the subscript "." is a placeholder for susceptibles (S), latent (E), prodromal (P) and fully infectious (I), respectively. After, the outcome of the vaccine manifests, $f_.^{(R)}$, $f_.^{(PI)}$, $f_.^{(NI)}$, $f_.^{(ADE)}$ are the fractions of individuals who become completely immune, partially immune, fail to immunize, and develop ADE, respectively.

Severity of ADE

By default, we assumed that ADE manifests in a higher probability of developing symptomatic infections, i.e., $f_{\text{Sick}}^{(ADE)} = 92\%$ *vs.* $f_{\text{Sick}} = 58\%$. Additionally, we investigated the effect of higher mortality of symptomatic infections in individuals that developed ADE. Figs 4-8 contrast the situations with and without ADE-induced increased mortality ($f_{\text{Dead}}^{(ADE)} = 20\%$ *vs.* $f_{\text{Dead}} = f_{\text{Dead}}^{(ADE)} = 7\%$). This increased mortality does not change incidence. For a vaccine with high effectiveness, the impact of ADE-induced increased mortality is marginal. If effectiveness is low, the effect is visible. This is not surprising, because the occurrence of ADE and vaccine effectiveness are not independent, namely, ADE occurs only if the vaccine fails to immunize properly. Fig 9 shows the effect of different amounts of mortality induced by ADE for different vaccination coverage. The effects are not immediate and only visible in the long run. The lower the vaccination coverage, the higher the increase in mortality. This is not surprising: if a higher proportion of the population is immunized, it is less likely that individuals that developed ADE get infected. Thus, the deleterious effects of ADE cannot manifest.

Discussion

With COVID-19 vaccines being approved swift and efficient action is required to immunize populations around the globe to further contain health and economic damages caused by the pandemic. Vaccination strategies need likely to be adjusted flexibly in response to the current status of the pandemic, feasibility of logistics, further vaccines becoming available, potential contraindications, side and long-term effects that were not recognized during ongoing phase III clinical studies, and the willingness in the population to get vaccinated. The purpose of this study was to tailor predictive modelling to optimize vaccination strategies for COVID-19 management and eradication. More precisely, we developed a realistic pandemic-preparedness model to study the influence of the onset of vaccination campaigns, the vaccination rate (i.e., the average time to get vaccinated), the vaccination coverage (the fraction of the population

Fig 9. ADE-induced increased mortality. As Fig 8, but for different levels of ADE-induced mortality instead of vaccination coverage. A vaccination coverage of 60% was assumed. Panels show only mortality. For the corresponding incidence see Fig 8.

that can get vaccinated), vaccination schedules, effectiveness of the vaccine, and adverse ₃₂₇
side effects, particularly ADE. The model is a complex extension of the model ₃₂₈
underlying CovidSim 1.1 http://covidsim.eu/ [18]. We parameterized the model to ₃₂₉
reflect the situation in the Federal Republic of Germany, where vaccination campaigns ₃₃₀
started in late December 2020. Notably, although the results reported reflect the ₃₃₁
situation in Germany, the results apply qualitatively to any other country. Although, ₃₃₂
the model itself can be adapted to other countries to obtain quantitative results, their ₃₃₃
appropriateness needs to be taken with caution. Namely, the model neglects an explicit ₃₃₄
age-structure. Therefore, it is applicable to industrial nations with demographics similar ₃₃₅
to Germany. Adaptations will be necessary for low and middle-income countries with a ₃₃₆
large young population. These adaptations can be done similarly as in CovidSim 2.0 ₃₃₇
http://covidsim.eu/. ₃₃₈

The impact of the vaccination in terms of incidence and mortality depends on the ₃₃₉
contact-reducing interventions in place. Here, it was assumed that a "hard lockdown" ₃₄₀
will be sustained until the end of January 2020 followed by a "relief period" still with ₃₄₁
relatively strong contact reduction, until summer 2021. Afterwards no contact reduction ₃₄₂
was assumed. These assumptions are obviously questionable. The rationale behind ₃₄₃
them was that case numbers in 2021 will first require action to reduce disease incidence ₃₄₄
by sustaining contact reduction. Once people get vaccinated and incidence is decreasing, ₃₄₅
compliance with distancing measures will fade after the summer season. As soon as ₃₄₆
attendance of cultural events (e.g., concerts, museums, sports events, etc.) and ₃₄₇
unrestricted air travel will be possible and mandatory wearing of facial masks will be ₃₄₈
lifted for vaccinated individuals, it will be difficult to control distancing interventions in ₃₄₉
the population. Notably, we assume that case isolation (quarantine and home isolation ₃₅₀
of confirmed cases) is further sustained. ₃₅₁

Our simulations adequately reflect the dynamics of the COVID-19 epidemic in ₃₅₂
Germany in 2020. The future predictions, however, depend on assumptions regarding ₃₅₃
COVID-19 management. The model parameters need to be dynamically updated as ₃₅₄
COVID-19 management interventions are altered. Particularly, parameters concerning ₃₅₅
vaccination campaigns are likely to vary over time and can be modelled in a ₃₅₆
time-dependent fashion. This is a straightforward generalization. Here, we decided to ₃₅₇
simulate a plausible range of fixed parameters to quantify their effects, rather than ₃₅₈
making assumptions on time-dependence that would be purely speculative. According ₃₅₉
to our predictions, vaccination campaigns will have a strong impact on the reduction in ₃₆₀
disease incidence. In the short term, a swift onset of the vaccination campaign ₃₆₁
contributes to a substantial reduction in incidence and mortality in the first quarter of ₃₆₂
2021. The later mass vaccination starts, the smaller the reduction in incidence or ₃₆₃
mortality. The onset of the campaign mainly depends on the logistics to initially ₃₆₄
distribute the vaccine efficiently. ₃₆₅

The vaccination schedule also has a substantial impact on incidence and mortality. ₃₆₆
Our simulations showed a substantial improvement if the time to immunization is ₃₆₇
shortened from 42 to 28 days. This depends on the vaccination schedule, which requires ₃₆₈
typically two doses with a waiting time of two to three weeks between them. After that, ₃₆₉
the immunizing effect is reached within about 14 days. Efficient logistic planning and ₃₇₀
properly scheduling appointments to receive the two required doses can help to ₃₇₁
minimize the time to immunization. ₃₇₂

Not surprisingly, the higher the vaccination rate (i.e., the faster the population is ₃₇₃
vaccinated), the stronger the benefit. The vaccination rate depends crucially on the ₃₇₄
available infrastructure. The modRNA-based vaccine of BioNTech (cf. Introduction) ₃₇₅
requires storage between -60 to $-80°C$ [20]. Therefore, it needs to be distributed ₃₇₆
through a specific infrastructure (vaccination centers/units) limiting the vaccination ₃₇₇
rate. The achievable vaccination rate hence depends on the approval and availability of ₃₇₈

competing vaccines. Vaccine efficiency differs across competing products and has a strong effect on incidence and mortality. In the simulations, by default, we assumed an ambitious vaccination rate, with an average waiting time of 180 days to get vaccinated. With two doses per person vaccinating 60% of the German population within one year requires a capacity of 270 000 injections per day. With around 400 vaccination centers that have been established, this requires a daily average capacity of 680 injections per center. While the capacity is realistic, also the willingness to get vaccinated must be high to efficiently utilize the capacities.

Another important factor is vaccination coverage. Our simulations suggest that a vaccination coverage of $75\% - 80\%$ is necessary to mitigate the epidemic by summer 2021, without further strong lockdowns and contact restrictions. The reason is that a sufficiently high level of immunity needs to be reached by the onset of the 2021 flu season in order to prevent another epidemic outbreak. Such an outbreak might be more difficult to control as interventions will be a balancing act between restrictions tolerable by the vaccinated part and necessary to protect the unvaccinated part of the population. Notably, even if the herd immunity threshold to prevent a COVID-19 outbreak is substantially lower than 75%, this threshold must be reached on time. Importantly, the vaccination coverage in our predictions also reflects the propensity to get vaccinated early. Furthermore, vaccination does not lead to immunization in all cases, a fact that was addressed in our model. In particular, we studied the consequences of varying vaccine efficiencies, which have a substantial effect. According to clinical trials, efficacy varies between $78\% - 95\%$ among the most promising vaccines [21–24]. With lower efficiency vaccination coverage must increase to reach herd immunity, i.e, to reach immunization in 60% if the population 63% needs to be vaccinated if efficiency is 95%, while 77% need to be vaccinated if efficiency is just 78%.

Our predictions are conservative in as far as negative side effects of the vaccine were considered. In particular, we incorporated ADE or more generally ERD. These are well known in corona viruses and it has been explicitly warned about ADE in the context of vaccination campaigns [17]. The effects of ADE are notoriously difficult to predict [25]. Here, we assumed that a fraction of vaccinated individuals develops ADE, which results in a higher likelihood to develop symptomatic infections and higher mortality. Although we assumed mortality to be substantially increased (20% rather than 7% mortality in symptomatic infections), the overall effect was minor. More precisely, the reduction in mortality due to immunization achieved through vaccination always outweighs increased mortality due to ADE. This is an encouraging result that justifies neglecting ADE in future models. In fact, our model can be substantially simplified if ADE is neglected.

While our predictions adequately reflect disease incidence, the predictions for mortality have to be interpreted with caution. Namely, vaccination campaigns will target risk groups suffering from elevated mortality first. In fact, 50% of COVID-19-related deaths occur in long-term care facilities (LTCFs), although less than 1% of the German populations live inside such a facility. The spread of COVID-19 inside LTCFs can be efficiently maintained by regularly testing employees [26]. Concerning the interpretations of our results, mortality has to be understood qualitatively rather than quantitatively. However, adequate quantitative predictions can be easily deduced from our simulations by multiplying mortality with an adjustment factor. The relative effect of model parameters remains unaffected by an increase or decrease in mortality.

In conclusion, vaccination campaigns should be launched as early as possible. Logistics should be well planned to utilize the maximum capacity of the vaccination infrastructure. Failure to immunize a sufficient part of the population by the beginning of the flu season in 2021 will result in a high endemic peak, by far exceeding current levels of incidence. Adverse effects of the vaccine such as ADE are by far outweighed by the benefits of the vaccine. In fact, the higher vaccination coverage, the lower the risks

associated with ADE. We predict that vaccination coverage of 80% would result in 431
sufficiently high levels of herd immunity to allow a return to normality by summer 2021. 432
Nevertheless, it is important to sustain the vaccination campaign until the herd 433
immunity threshold is actually reached. This will require sustained incentives to get 434
vaccinated after disease incidence drops, e.g., through general contact reduction 435
measures being tight to vaccination coverage. 436

Acknowledgements 437

We want to dedicate this work to all voluntary participants in the COVID-19 438
vaccination trials. We also want to dedicate it to the victims of the SARS-CoV-2 virus. 439
Our grief is with the friends and families of the dreadful disease. The authors gratefully 440
acknowledge the discussion with Prof. Martin Eichner, Prof. Gideon Ngwa, and Dr. 441
Kristina B. Helle on the model. K. Schneider could not have put together the research 442
team without the support of the German Academic Exchange (Project-ID 57417782), 443
the SMWK-SAB project "Innovationsvorhaben zur Profilschärfung an Hochschulen für 444
angewandte Wissenschaften" (Project number 100257255), the Federal Ministry of 445
Education and Research (BMBF) and the DLR (Project number 01DQ20002), the ESF 446
Young Investigator Group "Agile Publika" funded by ESF, SMWK, SAB (SAB Project 447
100310497). 448

Supporting information 449

S1 Fig. Simplified model flow. Susceptible individuals are infected and ultimately 450
recover or die. Individuals are infected either after or before the outcome of the 451
vaccination is determined. In the latter case the outcome of the vaccine is not 452
determined upon infection (they might get vaccinated before or during the infection), or 453
they are not vaccinated before or during the infection. 454

S1 Appendix. Mathematical Description 455

S1 Table. (Sub-) population sizes used in simulations. 456

S2 Table. Parameters describing disease progression. 457

S3 Table. Model compartments. 458

S4 Table. Parameters describing disease severity and mortality. 459

S5 Table. Parameters describing contact behavior and force of infection. 460

References

1. Blustein DL, Duffy R, Ferreira JA, Cohen-Scali V, Cinamon RG, Allan BA. Unemployment in the time of COVID-19: A research agenda. Journal of Vocational Behavior. 2020;119:103436. doi:https://doi.org/10.1016/j.jvb.2020.103436.

2. Talevi D, Socci V, Carai M, Carnaghi G, Faleri S, Trebbi E, et al. Mental health outcomes of the CoViD-19 pandemic. Rivista di psichiatria. 2020;55(3):137–144.

3. O'Callaghan KP, Blatz AM, Offit PA. Developing a SARS-CoV-2 Vaccine at Warp Speed. JAMA. 2020;324(5). doi:10.1001/jama.2020.12190.

4. Organization WH. Draft landscape of COVID-19 candidate vaccines; 2020.

5. Anderson EJ, Rouphael NG, Widge AT, Jackson LA, Roberts PC, Makhene M, et al. Safety and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine in Older Adults. New England Journal of Medicine. 2020;doi:10.1056/nejmoa2028436.

6. Walsh EE, Frenck RW, Falsey AR, Kitchin N, Absalon J, Gurtman A, et al. Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. New England Journal of Medicine. 2020;doi:10.1056/nejmoa2027906.

7. Mahase E. Covid-19: Russia approves vaccine without large scale testing or published results. BMJ. 2020;doi:10.1136/bmj.m3205.

8. Burki TK. The Russian vaccine for COVID-19. The Lancet Respiratory medicine. 2020;doi:10.1016/S2213-2600(20)30402-1.

9. Novavax. Evaluation of the Safety and Immunogenicity of a SARS-CoV-2 rS Nanoparticle Vaccine With/Without Matrix-M Adjuvant. US National Library of Medicine. 2020;.

10. Organization WH, Others. Guidance on developing a national deployment and vaccination plan for COVID-19 vaccines: interim guidance, 16 November 2020. World Health Organization; 2020.

11. Independent. Qantas to only accept passengers who have had coronavirus vaccine; 2020. Available from:
https://www.independent.co.uk/travel/news-and-advice/
qantas-passengers-coronavirus-vaccine-australia-b1760415.html.

12. CNN. Qantas boss says passengers will need to be vaccinated for international flights; 2020. Available from: https://edition.cnn.com/travel/article/
qantas-coronavirus-vaccination-intl-hnk-scli/index.html.

13. Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, Rabin K, et al. A global survey of potential acceptance of a COVID-19 vaccine. Nature Medicine. 2020;doi:10.1038/s41591-020-1124-9.

14. Wen J, Cheng Y, Ling R, Dai Y, Huang B, Huang W, et al.. Antibody-dependent enhancement of coronavirus; 2020.

15. Beretta A, Cranage M, Zipeto D. Is Cross-Reactive Immunity Triggering COVID-19 Immunopathogenesis? Frontiers in Immunology. 2020;doi:10.3389/fimmu.2020.567710.

16. Wu F, Yan R, Liu M, Liu Z, Wang Y, Luan D, et al. Antibody-dependent enhancement (ADE) of SARS-CoV-2 infection. medRxiv. 2020;.

17. Lee WS, Wheatley AK, Kent SJ, DeKosky BJ. Antibody-dependent enhancement and SARS-CoV-2 vaccines and therapies. Nature Microbiology. 2020;doi:10.1038/s41564-020-00789-5.

18. Schneider, Kristan and Ngwa, Gideon A and Schwehm, Markus and Eichner, Linda and Eichner M. The COVID-19 Pandemic Preparedness Simulation Tool: CovidSIM. SSRN. 2020;4/16/2020. doi:http://dx.doi.org/10.2139/ssrn.3578789.

19. R Core Team (2019). R: A language and environment for statistical computing. Accessed 1st April 2019. 2019;.

20. Pfizer Biontech. Checklist for Storage, Handling, and Preparation of the Pfizer-BioNTech COVID-19 Vaccine; 2020. Available from: `https://www.cvdvaccine-us.com/product-storage-and-dry-ice`.

21. Mahase E. Covid-19: Pfizer vaccine efficacy was 52% after first dose and 95% after second dose, paper shows. BMJ. 2020;371. doi:10.1136/bmj.m4826.

22. Mahase E. Covid-19: Moderna vaccine is nearly 95% effective, trial involving high risk and elderly people shows. BMJ. 2020;doi:10.1136/bmj.m4471.

23. Ryan Cross. Pfizer and BioNTech say their COVID-19 vaccine is successful in late-stage trial. Chemical & Engineering News. 2020;doi:10.47287/cen-09844-buscon1.

24. Voysey M, Clemens SAC, Madhi SA, Weckx LY, Folegatti PM, Aley PK, et al. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. Lancet (London, England). 2020; p. S0140–6736(20)32661–1. doi:10.1016/S0140-6736(20)32661-1.

25. Arvin AM, Fink K, Schmid MA, Cathcart A, Spreafico R, Havenar-Daughton C, et al. A perspective on potential antibody-dependent enhancement of SARS-CoV-2. Nature. 2020;584(7821):353–363. doi:10.1038/s41586-020-2538-8.

26. Tsoungui Obama NM Henri Christian Jr andain Adil Y, Alawam LN, Ngougoue N PM, Ngwa GA, Teboh-Ewungkem M, Schneider KA. Preventing COVID-19 spread in closed facilities by regular testing of employees-an efficient intervention in long-term care facilities and prisons. medRxiv. 2020;.