The RAG1 Ubiquitin Ligase Domain Enhances Assembly and Selection of T Cell Receptor Genes to Restrain the Autoimmune Hazard of Generating T Cell Receptor Diversity

Thomas N. Burn*,†,[1]; Charline Miot‡,[1]; Scott Gordon*,§; Erica J. Culberson‡; Tamir Diamond *,¶; Portia Kreiger‡; Katharina E. Hayer‖; Anamika Bhattacharyya#,**, Jessica M. Jones#, Craig H. Bassing*,‡,[2]; Edward M. Behrens*,†,[2]

*Institute for Immunology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104
†Division of Rheumatology, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
‡Department of Pathology and Laboratory Medicine, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
§Division of Neonatology, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
¶Division of Gastroenterology, Hepatology, and Nutrition, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
‖Department of Biomedical and Health Bioinformatics, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
#Department of Biochemistry and Molecular and Cellular Sciences, Georgetown University, Washington, DC, 20057
**Present Address: Vyome Therapeutics Lt., 501 Patparganj Industrial Area, Dehli, India
††Department of Pathology and Laboratory Medicine, The Perelman School of Medicine of the University of Pennsylvania, Philadelphia, PA 19104

[1]Co-first Authors

[2]Co-corresponding Authors:
        Craig H. Bassing, Ph.D.
        Children's Hospital of Philadelphia
        4054 Colket Translational Research Building
        3501 Civic Center Blvd.
        Philadelphia, PA 19104
        Phone: 267-426-0311
        Email: bassing@email.chop.edu

        Edward M. Behrens, M.D.
        Children's Hospital of Philadelphia
        1102 Abramson Research Center
        3615 Civic Center Boulevard
        Philadelphia, PA 19104
        Phone: 267-426-0142
        Email: behrens@email.chop.edu

Running Title: RAG1 enhances TCR assembly and thymic selection

Abstract

RAG1/RAG2 (RAG) endonuclease-mediated assembly of diverse lymphocyte antigen receptor genes by V(D)J recombination is critical for the development and immune function of T and B cells. However, this process creates highly self-reactive cells that must be negatively selected to suppress autoimmunity. The RAG1 protein contains a ubiquitin ligase domain that stabilizes RAG1 and stimulates RAG endonuclease activity *in vitro*. We report that mice with a mutation that inactivates the RAG1 ubiquitin ligase *in vitro* exhibit decreased assembly of T cell receptor (TCR) β and α genes in thymocytes and impaired thymocyte developmental transitions that require assembly of these genes and signaling by their proteins. The mice also exhibit a combined phenotype of: 1) decreased expression of TCR signaling proteins in thymocytes, 2) less efficient positive selection of conventional αβ T cells and iNKT cells, 3) impaired negative selection of highly self-reactive thymocytes, and 4) CD4$^+$ αβ T cells with elevated autoimmune potential. Our findings demonstrate that the RAG1 ubiquitin ligase domain functions *in vivo* to stimulate efficiencies of TCRβ and TCRα gene assembly and αβ TCR selection, thereby establishing replete αβ TCR diversity and αβ T cell lineages while restraining the inherent autoimmune hazard of creating diverse antigen specificities.

Key Points

The RAG1 ubiquitin ligase domain stimulates TCR α and β gene assembly *in vivo*.

The RAG1 ubiquitin ligase domain enhances αβ TCR selection of thymocytes.

The RAG1 ubiquitin ligase domain limits autoimmune potential of αβ T cells.

Introduction

The ability of T and B lymphocyte populations to express antigen receptors able to recognize a potentially unlimited number of distinct pathogens is the fundamental basis of adaptive immunity. RAG1/RAG2 (RAG) endonuclease-mediated assembly of T cell receptor (TCR) and immunoglobulin (Ig) genes through V(D)J recombination establishes this vast pool of diverse receptors. Germline TCR and Ig loci consist of variable (V), joining (J), and sometimes diversity (D), gene segments residing upstream of constant (C) region exons. During T and B cell development, the lymphocyte-specific RAG complex cooperates with lineage- and developmental stage-specific activation of TCR and Ig loci to perform V(D)J recombination (1). RAG induces DNA double strand breaks (DSBs) between coding sequences and flanking recombination signal sequences of two participating gene segments, liberating hairpin-sealed coding ends and blunt signal ends (2). In a subsequent reaction phase guided by RAG, DNA repair factors ligate signal ends to generate a signal join and open, process, and ligate coding ends to create a V(D)J coding join, which comprises the second exon of an assembled V(D)J-C antigen receptor gene (2). The high number of V(D)J joining events, broad utilization of gene segments, and inherent imprecision in opening and processing coding ends all cooperate to yield a vast population and diversity of receptors. The imprecise formation of coding joins creates the third hypervariable complementarity-determining region (CDR3)-encoding nucleotides of Ig and TCR variable regions. Due to imprecise means of opening and processing coding ends, the outcomes of V(D)J recombination include out-of-frame genes unable to make protein and receptors that recognize self-antigens and have potential for autoimmunity. Accordingly, developing T and B lymphocytes engage quality control checkpoints to select for potentially beneficial antigen receptor proteins and against hazardous highly self-reactive receptors.

Both immature T and B cells employ developmental programs that link functional protein expression from in-frame V(D)J rearrangements to signal either survival and continued differentiation or apoptosis (1).

The maturation of αβ T cells in the thymus provides a paradigm for how V(D)J recombination and antigen receptor protein quality control checkpoints cooperate to create a large pool of lymphocytes with diverse receptors that provides immunity from pathogens without causing overt autoimmunity. After entering the thymus, early thymocyte progenitor cells differentiate into $CD44^+CD25^+$ stage 2 and then into $CD44^-CD25^+$ stage 3 $CD4^-CD8^-$ double negative (DN) thymocytes (3, 4). Both DN2 and DN3 cells express RAG and conduct V(D)J recombination of *Tcrb*, *Tcrd*, and/or *Tcrg* loci (3, 4). Although DN2-to-DN3 thymocyte maturation can occur independent of RAG and V(D)J recombination, an in-frame *Tcrb* rearrangement is necessary for development beyond the DN3 stage (3, 4). The resultant TCRβ protein associates with invariant pTα protein to produce pre-TCR complexes that signal antigen-independent survival, proliferation, and differentiation of DN3 cells (β-selection) into $CD44^-CD25^-$ stage 4 DN thymocytes and then $CD4^+CD8^+$ double-positive (DP) thymocytes (3, 4). DP cells express RAG and conduct V(D)J recombination of *Tcra* loci, which typically proceeds through multiple successive rounds of V-to-J rearrangement on both alleles (5). After an in-frame VJ rearrangement, the resulting TCRα protein can pair with TCRβ protein and yield surface αβ TCRs, which must signal to prevent death by neglect (6-9). This signal activation requires appropriate physical interactions between αβ TCRs and self-peptide/MHC (pMHC) complexes displayed on thymic epithelial cells (TECs) or dendritic cells (DCs), meaning that a functional αβ TCR is inherently self-reactive (6-9). Interactions below a particular low threshold of affinity (or avidity) cannot activate signalling to block apoptosis (6-9). At the other extreme, contacts above a higher threshold trigger strong signals that causes apoptosis (negative selection) to delete highly self-reactive cells with substantial autoimmune potential (6-9). Contacts between these thresholds activate TCR signals of a strength within a range that promotes survival and differentiation of DP cells (positive selection) into $CD4^+$ or $CD8^+$ single-positive (SP) thymocytes, which lack RAG expression and exit the thymus as mature αβ T cells (6-9). Strong TCR signals in DP cells can also activate an alternative agonist selection process that promotes differentiation of unconventional αβ T cell lineages: regulatory T

(T reg) cells, intestinal intraepithelial lymphocytes (IELs), and invariant natural killer T (iNKT) cells (6, 7). Notably, the αβ TCRs of iNKT cells have limited diversity and recognize lipids displayed by MHC-like CD1d proteins on another DP thymocyte (10). This limited diversity renders developing iNKT cells a useful thymocyte population to probe issues of TCR signal strength without confounding by contribution of TCR specificity. A notable aspect of the αβ TCR quality control checkpoint is that thymocytes are more sensitive than mature αβ T cells as positively-selecting pMHC activates TCR signaling only in the former (9, 11-15). This differential is thought to enable positive selection from low affinity/avidity interactions, prevent the same contacts from activating mature αβ T cells, and provide a safety net for negative selection to more effectively suppress autoimmunity (6-9). Although the T lineage-specific Themis protein contributes to this differential in TCR sensitivity between immature and mature αβ T cells (9, 16, 17), the precise mechanisms by which this protein and possibly additional factors "tune" TCR signaling and control αβ TCR selection remain to be elucidated.

The RAG1 protein N-terminus has a Zinc-binding RING finger ubiquitin ligase domain of undetermined relevance for antigen receptor gene assembly. *In vitro,* RAG1 autoubiquitylates on lysine 233 (18), which enhances RAG endonuclease activity *in vitro* as evidenced by impaired cleavage upon mutation of this residue (19). *In vitro,* RAG1 also ubiquitylates other proteins including histone H3 within chromatin (20, 21). RAG1 RING domain mutations that disrupt ubiquitin ligase activity impair RAG-mediated cleavage and joining of chromatin substrates in cell lines, which could be due to loss of histone H3 ubiquitylation and/or alteration in RAG1 protein structure (21-23). The C328Y mutation of a critical structural zinc-binding cysteine of human RAG1 impairs V(D)J recombination and causes severe T and B cell immunodeficiency associated with uncontrolled proliferation of activated oligoclonal αβ T cells (24). The analogous C325Y mutation of mouse Rag1 protein (C325Y) *in vitro* destabilizes Rag1 tertiary structure, abrogates ubiquitin ligase activity, reduces V(D)J recombination (22). Mice with homozygous Rag1

C325Y mutation have a near complete block of αβ T cell development at the DN3 thymocyte stage due to impaired *Tcrb* recombination (22, 23). RING domains have a conserved proline (proline 326 in mouse Rag1) that is critical for ubiquitin ligase activity by permitting functional interaction with ubiquitin-conjugating enzymes (25, 26). Relative to C325Y mutation, P326G mutation abrogates Rag1 ubiquitin ligase activity equivalently, but has substantially less severe effects on both destabilizing tertiary structure of the Rag1 RING domain and reducing RAG endonuclease activity (23). Accordingly, P326 mutation of endogenous mouse Rag1 protein allows an approach to elucidate potential physiologic roles of the RAG1 ubiquitin ligase domain beyond promoting V(D)J recombination by stabilizing RAG1 protein structure.

To determine potential *in vivo* functions of RAG1 ubiquitin ligase activity, we created and analyzed mice carrying a homozygous Rag1 P326G mutation. We show that these mice exhibit decreased levels of *Tcrb* or *Tcra* rearrangements in DN3 or DP thymocytes, respectively, correlating with lower fractions of these cells continuing to differentiate. When assaying αβ TCRs expressing specific TCRβ or TCRα chains, we observe diminished efficiencies for superantigen-mediated negative selection of conventional αβ T cells and differentiation of iNKT cells. We find that the αβ TCR signaling protein Zap70 is expressed at lower than normal levels in DP cells but not mature αβ T cells, providing one possible reason for less efficient selection of αβ TCRs. Finally, we show that mature CD4$^+$ effector αβ T cells of these mice exhibit normal immune responses to activation, yet confer greater potential for autoimmunity. Our findings demonstrate that the RAG1 ubiquitin ligase domain functions *in vivo* to stimulate efficiencies of TCRβ and TCRα gene assembly and αβ TCR selection, thereby establishing replete αβ TCR diversity and αβ T cell lineages while restraining the inherent autoimmune hazard of creating diverse TCR specificities. We discuss how the RAG1 ubiquitin ligase domain might function *in vivo* to regulate the assembly of TCRβ and TCRα genes and the selection of αβ TCRs.

Materials and Methods

Mice

C57BL/6 background *PG* mice were made through gene-targeting as performed by InGenious Targeting Laboratory (2200 Smithtown Avenue, Ronkonkoma, NY). The 9.5 kb genomic DNA used to construct the targeting vector was sub-cloned from a 129Svev BAC clone. The CC>GG double point mutation was shuttled from pJMJ029[P326G](23) into the targeting vector using unique restriction sites, then confirmed by sequencing. Linearized targeting vector was electroporated into BA1 (C57BL/6 x 129/SvEv) embryonic stem (ES) cells. After selection with G418 antibiotic, clones were expanded for PCR analysis to identify recombinant ES clones. Positive clones were screened for short arm integration and then confirmation of the point mutation was performed by PCR and sequencing. Positive clones were analyzed by Southern blot to confirm long arm integration. Injected blastocysts were grown in foster mothers and chimeras were backcrossed against C57BL/6 mice to generate heterozygotes. These were crossed with CMV-*cre* mice (JAX #006054)(27) to remove the *neo* reporter cassette and generate the *Rag1^{P326G}* allele. After confirmation of *neo* cassette removal, the strain was backcrossed against C57BL/6, and removal of CMV-*cre* was confirmed by PCR. Sequencing of the entire *Rag1* locus in progeny confirmed the intended CC>GG mutation and an additional T>C point mutation that generates a V238A substitution outside of the RING finger domain. The strain was deposited at the Mutant Mouse Regional Resource Center at the University of North Caroline (RAG1-tm1, MMRRC #: 37105). Before initiating studies described herein, the *Rag1^{P326G}* allele was backcrossed for nine generations onto the C57BL/6 background. To generate NOD background *PG* mice, CRISPR/Cas9-mediated gene editing was conducted in zygotes from *in vitro* fertilization of NOD eggs with NOD sperm. A guide RNA (5'-AAGCACGGATATCGGCAAGA-3') that hybridizes near Rag1 sequences that encode proline 326 was generated as described (28). The following mix was electoporated into zygotes: 4μM of purified guide RNA, 4μM of Cas9-NLS nuclease (qb3 MacroLab Berkeley), 10μM of ssDNA 5'-

C*A*G*AAAGGACTTCACTGGGCTCTCCAGGTCAGTAGGGAAGCACGGATATCGGCAGGAT
CCACAATAGCTGCCCATGACTTTGAGACATCTGAGAATGCAGATCCTACAGAATAGATGCT
T-3' (IDT) that contains nucleotides encoding glycine in place of proline at amino acid 326 and a silent point mutation of the next codon which introduce a cleavage site for BamHI restriction enzyme. Mice were screened for introduction of the mutation by PCR with primers 5'-CGTCGCAAGAGAACTCAGGCT-3' and 5'-TCTCGTCAGTGACAGGAGATGCT-3', followed by BamHI digestion of the PCR product. Sanger sequencing of PCR products confirmed incorporation of the desired mutation and no other mutations in the *Rag1* coding sequence. Founders with mutations were mated with NOD mice to establish NOD *Rag1$^{P326G/+}$* mice, which were bred with each other to generate NOD *PG* mice and *WT* littermate controls. NOD mice, C57BL/6 background wild-type and *Rag1$^{-/-}$* mice and BALB/c background *Rag1$^{-/-}$* mice were purchased from Jackson Laboratories. All experimental mice were littermates or age-matched with control wild-type animals. All experiments were conducted in accordance with national guidelines and approved by the Institutional Animal Care and Use Committee (IACUC) of the Children's Hospital of Philadelphia.

Flow Cytometry

Single cell suspensions of all organs were ACK lysed to remove erythrocytes before being stained with LIVE/DEAD™ Fixable Aqua Dead Cell Stain Kit (Invitrogen) and antibodies against respective surface antigens (BD Biosciences, eBioscience, and Biolegend). Cells were stained for respective intracellular proteins using Cytofix/Cytoperm Fixation/Permeabilization solution kit (BD Biosciences). For staining of intracellular cytokines, cells (1 x10^6) were cultured in the absence or presence of 50 ng/ml PMA (Sigma) and 1μg/ml Ionomycin (Cell Signaling Technology), with 2 μg/ml brefeldin A (Sigma) and 2 μM monensin (eBioscience) for 4 hours at 37°C. After staining for LIVE/DEAD and surface antigens, cells were stained for respective cytokines using the Cytofix/Cytoperm kit (BD Bioscience) or transcription factors using the FoxP3/Transcription Factor staining kit (eBiosciences). Data were acquired on a

MACSQuant flow cytometer (Miltenyi Biotec) or LSRII Fortessa (BD Biosciences) and analyzed using Flowjo software version 10.5.3 (Tree Star). Analyses were conducted with the following antibodies: anti-CD4 (APC-eFluor 780, clone RM4-5, 1:100), anti-CD25 (BUV395, clone PC61, 1:100), anti-TCRβ (BV711, clone H57-597, 1:100), anti-CD19 (PE, clone 1D3, 1:100), anti-CD11b (PE, clone M1/70, 1:100), anti-CD11c (PE, clone HL3, 1:100), anti-TER (PE, clone TER-119, 1:100), anti-TCRγ/δ (PE, clone GL3, 1:100), anti-NK-1.1 (PE, clone PK136, 1:100), and anti-B220 (PE, clone RA3-6B2, 1:100) from BD Biosciences; or anti-CD8α (Pacific Blue, clone 53-6.7, 1:100), anti-CD44 (APC, PerCP/Cy5.5, clone IM7, 1:100), anti-CD24 (PE/Cy7, clone M1/69, 1:100), anti-CD28 (FITC, clone E18, 1:50), anti-CD45.1 (APC, clone A20, 1:100), anti-C45.2 (FITC, clone 104, 1:100) from BioLegend.

Cell Sorting

To isolate DN3 cells, thymocytes were stained with PE-labelled CD4, CD8, CD11b, CD11c, NK1.1, Gr1, and Ter119. Non-labelled cells were enriched by MACS depletion using anti-PE microbeads and LS columns (Miltenyi). Enriched cells were then stained with CD4, CD8, CD44, CD25 and anti-lineage/stage markers (TCRβ, B220, CD19, CD11b, CD11c, TCRδ, NK1.1, Ter119) antibodies. DN3 thymocytes were sorted on Lin⁻CD4⁻CD8⁻CD44⁻CD25⁺ phenotype. To isolate DP cells, thymocytes were stained with CD4, CD8, and anti-lineage (B220, CD19, CD11b, CD11c, TCRδ, NK1.1, Ter119) antibodies. DP thymocytes were sorted on Lin⁻CD4⁺CD8⁺ phenotype. The cells of interest were isolated using a FACSAria Fusion sorter (BD Biosciences).

Analyzing *Tcrb* and *Tcra* Rearrangements

Genomic DNA was extracted from DN3 or DP cells using the DNeasy Blood and Tissue kit (Qiagen). To measure *Tcrb* rearrangements in DN3 cells, a Taqman PCR assay was used to quantify Vβ-to-DJβ1.1 and Vβ-to-DJβ2.1 rearrangement levels with a primer specific for each Vβ paired and a probe, FAM or HEX,

specific for Jβ1.1 or Jβ2.1, respectively. Taqman PCR was performed with conditions according to the manufacturer's instructions (IDT DNA) on the ViiA 7 system (Applied Biosystems). PCR of CD19 was used for normalization. Primers, probes, and reaction conditions are as described (29). To assay *Tcrb* repertoire, DNA from DN3 thymocytes were sent to Adaptive Biotechnologies, who used multiplex PCR to amplify and deep sequence Vβ-Dβ-Jβ rearrangements. Gene segment usage was analyzed by ImmunoSEQ Analyzer software (Adaptive Biotechnologies). To assess *Tcra* rearrangements in DP cells, representative Vα-Jα rearrangements were quantified using a QuantiFast SYBR Green PCR kit (Qiagen) on the ViiA 7 system (Applied Biosystems) as described (30, 31). PCR of β2M was used for normalization.

RT-qPCR for mRNA Expression

Respective cell populations were lysed in RLT buffer (Qiagen) containing 2-mercaptoethanol or TRIzol (Life Technologies) immediately after cell sorting. Total RNA was isolated using RNeasy Mini kit (Qiagen), treated with DNase (RNase-Free DNase Set, Qiagen), and reverse transcribed to generate cDNA with High-Capacity RNA-to-cDNA™ Kit (Themo-Fisher Scientific) according to manufacturer's directions. The cDNA was subjected to qPCR using Power SYBR Green kit (Applied Biosystems) and actin, HPRT, ZAP70, and SYK primers (Qiagen). Relative expression was calculated using the ddCt method, using actin or HPRT as a housekeeping gene, and relevant calibrator sample (explained in figure legends for respective samples).

Competitive Mixed Bone Marrow Chimeric Mice

6-week-old CD45.1 B6.SJL (WT) hosts were lethally irradiated (950 Rad on an X-RAD irradiator). Bone marrow from at least 3 pooled donors (CD45.1 WT only, CD45.2 RAG P326G mutant only, or mixed) was sterilely prepped and RBCs lysed. Acceptable mixing (CD45.1/CD45.2 ratio) was confirmed by flow cytometry. Then, $3-5 \times 10^6$ total cells were injected intravenously into hosts 6 hours after irradiation. Mice were monitored over 8 weeks for graft rejection and graft-versus-host disease prior to euthanasia for

analysis. Engraftment was measured using the following antibodies from Biolegend: anti-CD45.1 (APC-Cy7, clone A20, 1:100), anti-CD45.2 (FITC, Clone 104, 1:100), anti-NK1.1 (PE, clone PK136, 1:200), anti-CD44 (PerCP-Cy5.5, clone IM7, 1:100), anti CD24 (PE-Cy7, clone M1/69, 1:100), and anti-TCRβ (Pacific blue, clone H57-597, 1:100). CD1d tetramer from the NIH Tetramer Core (APC, 1:100) was a gift of Dr. Hamid Bassiri. Data were collected on a MACSQuant 10 (Miltenyi Biotec) and were analyzed using FlowJo software, version 10.

In vitro T Cell Differentiation

Splenocytes from C57BL/6 background *Rag1*$^{-/-}$ mice were irradiated with 2,500 Rads (X-rad irradiator) and 3 x 10^5 cells were seeded into each well of a 96 well plate. Naïve T cells (CD90.2$^+$, CD4$^+$, CD62L$^+$, CD44$^+$, CD25$^-$) were sorted from C57BL/6 background *WT* or *Rag1*$^{-/-}$ mice using a FACSaria Fusion cell sorter (BD Biosciences), and 3 x 10^4 sorted cells were added to each well containing irradiated feeder cells. All cells were cultured with 10 μg/ml anti-CD3 (Biolegend) and 3 μg/ml anti-CD28 (Biolegend) in RPMI with 10% FBS, 1% PSG, and 1x NEAA (Invitrogen), 1x Sodium Pyruvate (Invitrogen), and 0.001% 2-mercaptoethanol. rhIL-2 was used at 30 IU/mL, rmIL-12 and rmIL-4 at 10 ng/mL, rmIL-6 at 20 ng/mL, mTGFβ1 at 1 ng/mL, anti-mIL-4, anti-mIFNγ and anti-mIL-12 at 10 μg/mL (from Biolegend except rhIL-2, rmIL-12, and rmIL-6 from Prepotech). The following cytokines/blocking antibodies were used to skew towards respective Th-subsets: Th0 condition: rhIL-2; Th1 condition: rhIL-2, rmIL-12 and anti-mIL-4; Th2 condition: rhIL-2, rmIL-4, anti-mIFNγ and anti-mIL-12; Th17 condition: rhIL-2, mIL-6, mTGFβ1, anti-mIL-4 and anti-mIL-12; Treg condition: rhIL-2, mTGFβ1, anti-mIL-4 and anti-mIL-12.

LCMV Infections

Mice were infected intraperitoneally with 2 x 10^5 plaque-forming units of LCMV-Armstrong and euthanized at indicated timepoints. Virus and gp33-tetramer were kindly provided by E. John Wherry (University of Pennsylvania).

T Cell Transfer Model of Colitis

This model was performed as described (32). Briefly, naïve CD4$^+$ T cells were sorted from respective donor mice (CD90.2$^+$, CD4$^+$, CD62L$^+$, CD44$^-$, CD25$^-$) on a FACSAria Fusion cell sorter. 0.5 x 10^6 cells were transferred in 100 µl of PBS into each of the donor mice via intraperitoneal injection. Non-transferred mice were injected with PBS alone. Mice were weighed weekly for the first 4 weeks, and then twice weekly for 4 weeks. Mice were euthanized 8 weeks post-transfer unless severe morbidity enforced premature euthanasia (in accordance with CHOP IACUC approved guidelines) or spontaneous death occurred.

Diabetes Onset in NOD Background Mice

NOD background *WT* and *PG* mice were aged and assessed for spontaneous development of diabetes. Mice were aged and monitored for disease to a maximum of 30 weeks of age. Mice older than 10 weeks of age were weekly tested for glycosuria (Uristix$^®$, Siemens). For positive mice, glycosuria was assayed by a second test 24 hours later. Capillary blood glucose of mice testing positive twice for glycosuria was tested via facial vein bleeding. Mice with blood glucose $\geq$250 mg/dl (13.9 mmol/l) were considered as diabetic and euthanized.

Statistical analysis

All data was analyzed in Graphpad Prism 8 using statistical tests indicated in the figure legends. Error bars indicate mean +/- SEM unless otherwise stated. n.s.=$p>0.05$, *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Homozygous Rag1 P326G mutation diminishes TCR gene assembly and subsequent thymocyte developmental stage transitions.

To determine potential *in vivo* functions of the RAG1 ubiquitin ligase, we created and analyzed C57BL/6 strain mice homozygous for the Rag1 P326G mutation (*Rag1$^{P326G/P326G}$* mice; see Materials and Methods) that abrogates RAG1 ubiquitin ligase activity but minimally destabilizes RAG1 protein (23). We initially analyzed αβ T cell development and TCR gene rearrangements in *Rag1$^{P326G/P326G}$* (*PG*) mice and wild-type (*WT*) mice. As compared to *WT* mice, *PG* mice have a ~2-fold higher percentage and number of DN3 thymocytes and correspondingly lower percentage and number of DN4 thymocytes (Fig. 1, A and B). *PG* mice also exhibit ~10% reductions in the percentage and number of DP thymocytes and the ratio of CD4$^+$ SP thymocytes to DP thymocytes (Fig. 1, C-E). Despite these minor impairments of the DN3-to-DN4 and DP-to-SP thymocyte developmental transitions, *PG* mice have normal numbers of total thymocytes and splenic αβ T cells (Fig. 1 F). Using Taqman PCR to amplify Vβ rearrangements to Dβ1Jβ1.1, Dβ1Jβ2.1, or Dβ2Jβ2.1 complexes in non-selected DN3 cells, we observed ~50% lower levels of rearrangement for nearly all individual Vβ segments in *PG* mice (Fig. 2, A and B). Notably, the recombination of some Vβ segments involving Jβ2.1 were undetectable in *PG* mice (Fig. 2, A and B), suggesting altered Jβ repertoire from Rag1^{P326G} mutation. *Rag1* or *Rag2* mutations that substantially decrease RAG endonuclease activity, thymocyte development, and thymic cellularity also alter the usage of individual Vβ segments (33, 34). However, high-throughput sequencing of *Tcrb* rearrangements in DN3 thymocytes shows similar usage of each Vβ segment in *PG* and *WT* mice (Fig. 2 C), reflecting the modest reduction of *Tcrb* gene assembly in *PG* mice. This analysis also reveals modest alterations in Dβ and Jβ usage, but no difference in CDR3β lengths, in *Tcrb* rearrangements of *PG* mice (Fig. 2, D and E; data not shown). Using Taqman PCR to amplify a subset of possible Vα-to-Jα rearrangements in non-selected DP thymocytes, we detected ~50% lower levels of rearrangements for many of the Vα/Jα combinations in *PG* mice (Fig. 2, F). Together,

these data indicate that homozygous Rag1^{P326G} mutation produces RAG endonuclease complexes that catalyze decreased levels of *Tcrb* and *Tcra* rearrangements, alter the primary repertoires of *Tcrb* and *Tcra* genes, and support less efficient thymocyte developmental transitions that depend on the functional assembly of a *Tcrb* or *Tcra* gene. The lower efficiency of overall *Tcrb* and *Tcra* gene assembly certainly contributes to the diminished DN3-to-DP and DP-to-SP thymocyte developmental progressions.

Notably, the degrees to which TCR gene rearrangements and αβ T cell development are reduced in *PG* mice are dramatically less severe than reported for homozygous Rag1 C235Y mutation (22). For example, Rag1^{P236G} mice generate 90% of the normal number of DP thymocytes, whereas homozygous Rag1^{C235Y} mice produce only 1% of the normal number of DP thymocytes. The markedly different severities of these phenotypes suggest that abrogation of RAG1 ubiquitin ligase activity without substantial disruption of RAG1 protein structure only slightly impairs V(D)J recombination and αβ T cell development. Notably, the Rag1^{P326G} protein is expressed at modestly higher levels in thymocytes than wild-type Rag1 protein (Beilinson *et al.*, manuscript in revision), indicating that reduced Rag1 protein levels do not contribute to the slightly impaired development of αβ T cells in *PG* mice. We were not able to test the expected abrogation of *in vivo* Rag1 ubiqutin ligase activity in *PG* mice that is seen in *in vitro* data, leaving open the possibility that the mutant Rag1^{P326G} protein exhibits some level of ubiquitin ligase activity *in vivo*.

To confirm that the impaired development of αβ T cells in PG mice is due to cell intrinsic properties, we made and analyzed competitive bone marrow chimeric mice. These data show no signficiant differences in contributions of *WT* cells and *PG* cells at each of DN thymocyte stages (Supplemental Figure 1A). In marked contrast, there were greater contributions of *WT* cells at the DP and both SP thymocyte stages and within the CD4$^+$ and CD8$^+$ splenic αβ T cell populations (Supplemental Figure 1A, B). The data support

the notion that the decreased progression of Rag1^{P326G} thymocytes beyond the DN stage is due to cell intrinsic alterations, almost certainly the decreased assembly of TCRβ genes.

Reduced *Tcrb* gene assembly impairs DN thymocyte development in homozygous Rag1 P326G mice. The greater than normal number of DN3 thymocytes in *PG* mice is not typical of a mutation that impairs V(D)J recombination. It is notable that thymocytes lacking the Syk pre-TCR signaling protein display this phenotype (35, 36), and that RAG DSBs induced in pre-B cells trigger a genetic program that transcriptionally represses Syk protein expression to dampen pre-BCR signaling (37). As *PG* DN cells have reduced numbers of RAG DSBs from decreased *Tcrb* rearrangements, we determined whether these cells express lower than normal levels of Syk protein. We detected a slightly reduced level of Syk protein in *PG* mice (Fig. 3, A and B); however, unexpectedly, this correlated with an increase in the expression of Syk mRNA (Fig. 3 C). Thus, it appears that Rag1 P326G mutation results in a dissociation between the levels of transcript and protein of the Syk signaling molecule through as yet undetermined mechansisms. Regardless, these data open the possibility that pre-TCR signaled developmental progression of DN3 cells is impaired in *PG* mice.

To distinguish potential contributions of reduced *Tcrb* gene assembly versus decreased pre-TCR-signaling on impaired DN thymocyte development in PG mice, we conducted a more refined analysis of DN stage cells based on CD28 protein expression. After successful *Tcrb* gene assembly in CD28⁻CD25^{high} DN3a thymocytes, pre-TCR signaling drives differentiation of DN3b and then DN3c thymocytes characterized by upregulation of CD28 and downregulation of CD25 (38). Mirroring relative levels of *Tcrb* rearrangements in DN3 thymocytes, we observed decreased progression of DN3a cells to the DN3b and DN3c stages, as well as a smaller fraction of DN3a cells expression TCRβ protein, in *PG* mice relative to *WT* mice (Fig. 3, D - I). We compared the percentage of DN3a cells expressing TCRβ protein to the percentage of these cells that develop into DN3b cells. We found equivalent values between *PG* and *WT*

mice (Fig. 3, J). We also compared the percentage of total DN3 cells expressing TCRβ protein to the fraction of these cells that progress to the DN4 stage. Here again we found equivalent values between *PG* and *WT* mice (Supplemental Figure 2). Together, these data provide no evidence for diminished pre-TCR signaled development of β-selected DN3 thymocytes. Therefore, the most plausible explanation for impaired DN thymocyte development in *PG* mice is decreased levels of TCRβ gene rearrangements.

Homozygous Rag1 P326G mutation lowers expression of the Zap70 signaling protein in thymocytes.
Given the evidence of the decreased levels of Syk protein but not mRNA in DN thymocytes of PG mice, we assessed whether the related Zap70 protein that transduces αβ TCR signals might show a similar pattern in DP thymocytes. To this end, we quantified levels of Zap70 protein and mRNA in DP cells at each of four stages of αβ TCR-signaled positive selection as defined by expression of CD69 and CCR7 proteins (Supplemental Figure 3) (39). In *PG* mice, we observed lower than normal Zap70 protein levels, but normal levels of TCRβ protein, at each stage of selection (Fig. 4, A-D). We detected normal levels of Zap70 mRNA in cells at each stage assayed in *PG* mice (Fig. 4 E), implying that lower Zap70 mRNA translation and/or Zap70 protein stability is the basis for lower Zap70 expression. RAG DSBs induced during lymphocyte development signal gene expression changes that persist in mature NK cells lacking RAG1 and RAG2 expression (40). We therefore considered whether the Zap70 alterations caused by mutant RAG1 during development would also persist in mature T cells after RAG1 is no longer expressed. We detected normal Zap70 protein levels in splenic CD4$^+$ and CD8$^+$ αβ T cells of *PG* mice (Fig. 4, F and G), indicating that Rag1^{P326G} mutation decreases Zap70 expression specifically in immature αβ T cells. These data imply that Zap70-mediated αβ TCR signaling might be weaker than normal in thymocytes and contribute to impaired DP-to-SP thymocyte transition in *PG* mice, beyond that resulting from reduced *Tcra* gene assembly and altered primary repertoires of assembled *Tcrb* and *Tcra* genes.

Homozygous Rag1 P326G mutation decreases efficiencies of positive, negative, and agonist selection.

In developing B cells, RAG DSBs generated during *Igk* gene assembly signal transcriptional upregulation of a proteins involved in lymphocyte selection and activation to modulate pre-BCR signaling (41). Considering this, the decreased levels of RAG DSBs in *PG* thymocytes, and the important roles of Zap70 in mediating positive, negative, and agonist selection (42, 43), we investigated whether any of these αβ TCR quality control processes might be impaired in *PG* mice. As a proxy to evaluate positive selection in *PG* mice, we determined the frequency of αβ TCR expressing (TCRβ$^+$) Stage 1 thymocytes that initiate positive selection as indicated by progression to Stage 2 in *PG* and *WT* mice (Fig. 5 A). By analyzing only αβ TCR expressing cells, we circumvent potential effects of decreased *Tcra* gene rearrangements reducing the fraction of DP thymocytes that express αβ TCRs and are subject to selection. This frequency is lower in *PG* mice (Fig. 5 B), consistent with modestly less efficient positive selection of αβ TCR expressing DP thymocytes upon homozygous Rag1^{P326G} mutation. Yet, we note that this is an imperfect means to quantify positive selection because it cannot account for possible differences in progression through subsequent stages or cellular fate decisions from specific TCR/pMHC interactions. Thus, we also quantified numbers of cells at each of the four stages of positive selection, finding fewer cells at each stage in *PG* mice relative to *WT* mice (Fig. 5C). Importantly, there is a greater decrease of cellularity in stage 2 (1.71x) as compared to stage 1 (1.25x), indicating fewer cells are moving from stage 1 to stage 2 in *PG* mice. Finally, we quantified CD69 expression and the ratio of CD69 to TCRβ expression at each stage, finding no difference of CD69 expression but a decrease of CD69 expression relative to TCRβ expression in stage 2 cells of *PG* mice (Fig. 5D). Collectively, these analyses indicate that positive selection is less efficient in *PG* mice. However, they do not distinguish between potential effects from altered primary repertoires of *Tcrb* and *Tcra* genes versus dulled αβ TCR signals. They also do not address the possibility that the <10% reduction in numbers of total thymocytes in *PG* mice alters thymic architecture in a manner that increases the fraction of DP cells that engage pMHC presented by TECs or DCs.

To determine whether the Rag1^{P326G} mutation impairs negative selection , we utilized a naturally occurring system of superantigen-mediated negative selection (44). Laboratory mice express remnant mouse mammary tumor virus (MMTV) retroviral products. Some of these products can bind the MHCII I-E^d protein and certain Vβ peptides, including Vβ5.1-5.2 (encoded by *Trbv12.1-12.2*) and Vβ12 (encoded by *Trbv15*), to trigger exceptionally strong αβ TCR signaling that promotes negative selection (44, 45). Our sequencing analysis of *Tcrb* gene rearrangements in DN3 thymocytes shows equivalent representation of *Trbv12.1*, *Trbv12.2*, and *Trbv15* in the primary Vβ repertoire (Fig. 2 C). Considering that Vβ repertoire remains constant during DN-to-DP thymocyte development (46), assaying superantigen-mediated deletion of Vβ5.1-5.2 or Vβ12 accounts for differences of Jβ repertoire of *Tcrb* genes assembled in *WT* and *PG* mice. Moreover, the reduced level and altered repertoire of *Tcra* genes assembled in DP cells of *PG* mice would not influence superantigen-mediated deletion of thymocytes expressing αβ TCRs with specific Vβ peptides. BALB/c, but not C57BL/6, strain mice express I-E^d protein. As a result, thymocytes expressing Vβ5.1-5.2 or Vβ12 are efficiently deleted in BALB/c, but not B6, genetic backgrounds. We bred BALB/c background *Rag1$^{-/-}$* mice with C57BL/6 background Rag1 *PG* or *WT* mice to generate and analyze *Rag1$^{PG/-}$* and *Rag1$^{WT/-}$* mice expressing I-E^d protein. To ascertain whether these mice exhibit a difference in superantigen-mediated negative selection, we measured the frequencies of pre-selected DP thymocytes and post-selection SP thymocytes expressing I-E^d:MMTV-reactive (Vβ5$^+$ or Vβ12$^+$) or -unreactive (Vβ6$^+$, Vβ8$^+$, or Vβ14$^+$ encoded by *Trbv19*, *Trbv13.1-13.3*, or *Trbv31*, respectively) αβ TCRs (Fig. 6, A-F). We calculated the percentages of DP and SP thymocytes expressing each Vβ peptide (Fig. 6, B-F). Our analysis indicates that greater fractions of I-E^d:MMTV-reactive thymocytes proceed through DP-to-SP development in *Rag1$^{PG/-}$* mice as compared to *Rag1$^{WT/-}$* mice (Fig. 5, B, C, and G), consistent with impaired negative selection of highly self-reactive thymocytes. In contrast, we found that the types of un-reactive thymocytes assayed develop equivalently in each genotype (Fig. 5, D -G). While the latter

finding does not reflect less efficient positive selection, we note that this experiment monitors a small subset of thymocytes expressing specific Vβ peptides and thus is not as sensitive as determining the frequencies of overall DP cells that initiate positive selection. Regardless, the data indicate that *PG* mice exhibit less efficient deletion of thymocytes that express αβ TCRs containing superantigen-reactive Vβ peptides. The plausible reasons for this impaired form of negative selection include that lower numbers of DP thymocytes prevents some from contacting or maintaining interaction with superantigen and/or αβ TCR signaling capacity is dampened.

Finally, to ascertain potential effects of Rag1^{P326G} mutation on agonist selection, we focused on assaying iNKT cell development. This lineage develops when TCRs containing an invariant Vα14-Jα18 TCRα chain and TCRβ chains with Vβ regions encoded by *Trbv1*, *Trbv13.1-13.3*, or *Trbv19* bind lipids presented by the MHC-like CD1d protein on different DP thymocytes (10, 47, 48). These interactions induce strong TCR signals that drive agonist selection and differentiation of iNKT cells through distinct stages that can be identified by binding lipid/tetramers and expression of defined proteins (10, 47, 48). We used a CD1d tetramer loaded with PBS57, an analogue of the α-galactosylceramide (α-GalCer) glycolipid, to identify α-GalCer-reactive iNKT cells in *WT* and *PG* mice. As compared to *WT* mice, *PG* mice had fewer absolute numbers of thymic α-GalCer-reactive iNKT cells (Fig. 7, A and B). While *PG* mice also tended to have fewer splenic iNKT cells (Fig. 7, A and B), this difference did not reach statistical significance (p=0.06) with the numbers of mice analyzed. The lower number of thymic iNKT cells in *PG* mice could arise solely from decreased levels of *Tcra* rearrangements. However, one advantage of using tetramers is that they identify DP thymocytes that have assembled and express the appropriate *Tcrb* and *Tcra* genes for development into iNKT cells. Thus, as a means to quantify agonist selection of iNKT cells independent of Vα14-Jα18 recombination levels, we quantified α-GalCer-reactive cells at each stage of iNKT cell development. We observed no significant differences in numbers of Stage 0 (CD24^{hi}CD44$^-$NK1.1$^-$) and

Stage 1 (CD24^{lo}CD44^{-}NK1.1^{-}) iNKT cells in *WT* and PG mice (Fig. 7, C and D). However, *PG* mice had lower numbers of Stage 2 (CD24^{lo}CD44^{+}NK1.1^{-}) and Stage 3 (CD24^{lo}CD44^{+}NK1.1^{+}) iNKT cells (Fig. 7, C and D) and diminished expansion of iNKT cells during progression from Stage 0 through Stage 4 (Fig. 7, E). Notably, the median fluorescence intensities of TCRβ expression and PBS57/CD1d tetramer binding of thymocytes was similar or slightly higher in *PG* mice relative to *WT* mice (Fig. 7 A, data not shown), indicating that α-GalCer-reactive iNKT cells of *PG* mice have a normal range of intrinsic affinity for this lipid presented by CD1d. Therefore, the plausible reasons for the impaired development of iNKT cells in *PG* mice are diminished engagements of TCRs with lipid/CD1d complexes and/or TCR signaling following lipid/CD1d recognition. The molecular basis for this could be decreased expression of Zap70 and/or other TCR signaling proteins, altered CDR3-encoding sequences that create TCRs less effective at signaling upon recognition of lipids presented by CD1d, or both.

We analyzed iNKT development in the competitive setting of mixed bone marrow chimeras comparing mice that received donor bone marrow that was either 100% *PG* or at 1:1 mix of *PG* and *WT* marrow. The very few remnant *WT* cells in the 100% PG receipient mice very effectively outcompeted the PG cells starting at Stage 1 and increasing in later stages (Supplemental Figure 1, C). The same effect can be seen in the 1:1 mix recipient mice (Supplemental Figure 1, C). Thus, there is a cell-intrinsic defect in *PG* thymocytes for iNKT development, consistent with a signlaing defect rather than an extrinsic, ligand based defect in their development.

Mature αβ T cells of homozygous Rag1 P326G mice exhibit normal responses to activation.

RAG endonuclease activity during lymphocyte ontogeny triggers permanent changes in gene expression that correlate with impaired responses of mature lymphocytes (40). Considering that RAG endonuclease function is diminished in thymocytes of *PG* mice, we sought to elucidate whether the mature αβ T cells

of these animals have normal or altered response to activation. We first conducted *in vitro* analyses. Upon

stimulation with PMA and ionomycin, splenic CD44^{hi} αβ T cells from *PG* mice produce normal levels of

IFNγ, TNFα, IL-2, and IL-17A (Fig. 8 A). These cells also show normal proliferation upon anti-CD3/anti-

CD28 stimulation (Fig. 8 B). Moreover, stimulated CD4$^+$ and CD8$^+$ αβ T cells of *PG* mice normally

express their canonical transcription factors (Fig. 8, C and D) and cytokines (Fig. 8, E and F) when cultured

in respective skewing conditions. We next monitored αβ T cell responses *in vivo* following infection of

mice with an acute strain of lymphocytic choriomeningitis virus (LCMV-Armstrong). We analyzed mice

either seven days later to monitor at the immune response peak or 35 days later to monitor memory cells.

At each timepoint, we detected similar frequencies and numbers of antigen-specific CD8$^+$ αβ T cells that

stain with gp33:H-2Db tetramer in *PG* and *WT* mice (Fig. 8 G), indicating that *PG* mice display normal

numbers and dynamic responses of αβ T cells that recognize LCMV-Armstrong. We also observed

equivalent dynamic expression of Tbet, Eomes, KLRG1, and CD127 proteins on αβ T cells from *PG* and

WT mice (Fig. 8 H), indicating normal contraction of short-lived effector cells (SLECs), and sustained

memory/memory precursor effector cells (MPECs) in *PG* mice. Finally, we detected similar numbers of

antigen specific αβ T cells able to respond to the gp33, NP396, or gp61 viral peptides within a peptide re-

stimulation assay (Fig. 8, I and J). Together, these data demonstrate that the mature αβ T cells of *PG* mice

have normal responses to activation, at least for the stimuli assayed.

.

Mature αβ T cells of homozygous Rag1^{P326G} mice exhibit increased predisposition to autoimmunity.
Within a setting of less efficient negative selection as with homozygous Rag1^{P326G} mutation, an increased

predisposition to αβ T cell mediated autoimmunity would be expected. However, we did not observe overt

autoimmune symptoms within *PG* mice, possibly because mice of the C57BL/6 background are resistant

to spontaneous autoimmune diseases (49, 50). Thus, to investigate whether the Rag1^{P326G} mutation can

predispose to autoimmunity, we utilized an induced model of autoimmune colitis in C57BL/6 mice and a spontaneous model of autoimmune diabetes in NOD mice.

The colitis model involves transferring into $Rag1^{-/-}$ mice naïve CD4$^+$ effector $\alpha\beta$ T cells, which become activated and trigger cytokine-driven colitis and associated systemic pathologies including weight loss due to the absence of Tregs (32, 51, 52). We observed accelerated wasting of $Rag1^{-/-}$ mice receiving PG cells relative to WT cells or no transfer (Fig. 9 A). We analyzed mice at eight weeks after transfer. As compared to WT recipients, we detected greater numbers of total splenic CD4$^+$ T cells and of CD4$^+$ T cells producing IFNγ, TNFα, and/or GM-CSF cytokines (Fig. 9, B and C). Colon length is used as a measure of disease severity in this model, and mice receiving transferred PG cells had shorter colons relative to mice receiving WT cells, consistent with worse disease caused by PG cells (Fig. 9 D). Histologic assessment and scoring in this colitis model has been formalized (52), and can be grossly broken down into scores related to inflammatory cell infiltration and scores related to injury of tissue architecture. There was no statistical difference in inflammation scores between mice receiving WT and PG cells. However, tissue damage scores were higher in mice receiving PG cells, suggesting that, on a per cell basis, PG cells induce greater tissue injury in this model (Fig. 9, E-F). Notably, specifically in PG-transferred animals, we observed autoimmune pathologies in addition to colitis, including dermatitis, chylous ascites, severe small intestine enlargement, and neurological stress (Fig. 9 H). Dermatitis was most common and involved massive inflammatory infiltration and tissue damage (Fig. 9 I). Consistent with more widespread autoimmune conditions in PG-transferred mice, we significant splenomegaly in these mice (Fig. 9 J). Finally, while not statistically significant, we observed an increased rate of premature death in the mice transferred with PG cells compared to WT cells (Fig. 9 K), which was especially surprising because death is an unexpected outcome in this model. Collectively, the data from this colitis model indicate that CD4$^+$ effector T cells from PG mice have greater than normal intrinsic autoimmune potential.

For the spontaneous model of autoimmune diabetes, we utilized CRISPR/Cas9-mediated gene editing to establish NOD mice carrying the $Rag1^{P326G}$ mutation. NOD mice develop spontaneous αβ T cell driven diabetes at an incidence of 60–80% in females and 20–30% in males (53, 54). We bred together NOD:$Rag1^{P326G/WT}$ mice to create cohorts of NOD *PG* and *WT* females, which we aged and screened for diabetes by monitoring glycosuria and blood glucose levels, euthanizing diabetic mice and confirming disease by blood glucose levels. We observed accelerated onset of diabetes in *PG* mice compared to *WT* mice, with median age of disease-free survival 120 and 175 days, respectively (Fig. 9 L). We started this study with the assumption that NOD $Rag1^{P326G}$ mice will experience an early event due to autoimmunity risk as compared to NOD $Rag1^{WT}$ mice, and if events do not occur early then they are unlikely to occur at all. Therefore, we evaluated potential significance of the different median disease-free survival using the Gehan-Breslow-Wilcoxon test that models such un-proportional hazard, finding the difference significant (p=0.049). However, we also applied the long-rank Mantel-Cox test that is more stringent, but should be used when the assumption is of proportional hazards between groups, finding the difference near, but not reaching, significance (p=0.1151). Unfortunately, the disruption of research from the Covid19 pandemic prevented us from continuing to generate and monitor mice so that we could more rigorously evaluate the difference in diabetes-free survival, as well as pathologically and immunologically characterize disease in afflicted animals.

Discussion

Our analysis of homozygous Rag1^{P326G} mice indicates roles for the integrity of the RAG1 RING domain ubiquitin ligase in stimulating TCRα and TCRβ gene asssembly and enhancing both positive selection of conventional αβ T cells and agonist selection of iNKT cells. The assembly and expression of functional Tcrb genes is rate limiting for thymocyte development from the DN3a stage through the DP stage. Our data show that the lower levels of Vβ rearrangements predominantly, if not solely, underlie the impaired progression of thymocytes through these early developmental stages in *PG* mice. On average, multiple rounds of successive Vα-to-Jα rearrangements occur in DP thymocytes before resulting αβ TCRs signal positive selection, negative selection, or agonist selection. For agonist selection of iNKT cells, Vα-to-Jα rearrangements must proceed until assembly of Vα14-Jα18 rearrangements, which can produce αβ TCRs that recognize lipid/CD1d complexes and signal development of iNKT cells from Stage 0 cells. The lower levels of Vα rearrangements in *PG* mice likely contributes to less robust positive selection of conventional αβ T cells by reducing the generating αβ TCR-expressing DP thymocytes. Although the normal numbers of α-GalCer-reactive Stage 0 iNKT cells does not support the notion that decreased levels of overall Vα rearrangements impairs the genesis of iNKT cells, our analysis of iNKT cells of a single specificity leaves open this possibility for the overall iNKT cell population. The amino acids encoded by individual gene segments and CDR3 nucleotides influence αβ TCR antigen recognition and ensuing TCR signaling. While we did not comprehensively study *Tcrb* and *Tcra* gene repertoires, our Taqman and Adaptive Immunoseq analyses of *Tcrb* gene rearrangements indicate minor differences of Dβ and Jβ repertoire between *WT* and *PG* mice. These in turn would generate subtle differences in TCRβ CDR3 repertoire that might contribute to decrease efficiencies of positive selection and agonist selection. Based on *in vitro* experiments (23), the molecular basis for reduced RAG endonuclease activity in *PG* mice could result from P326G mutation altering RAG complex tertiary structure, abrogating Rag1 ubiquitin ligase activity, and/or impairing Rag1

interaction with a protein that regulates RAG endonuclease activity. Moreover, while an intrinsically less efficient RAG endonuclease could account entirely for the reduced assembly of *Tcrb* and *Tcra* genes in *PG* mice, our analyses do not address potential roles of the RAG1 ubiquitin ligase in promoting chromatin accessibility and/or RAG binding of these loci. Distinguishing among all of these possibilities requires identifying all thymocyte proteins that bind Rag1 and/or are substrates of the Rag1 ubiquitin ligase, and then determining the effects of preventing their binding or ubiquityation by Rag1.

Our findings of less efficient superantigen-mediated negative selection, impaired iNKT cell development, and αβ T cells of higher autoimmune potential in *PG* mice imply roles for the RAG1 RING domain that transcend stimulating TCR gene assembly. The basis for superantigen-mediated negative selection is that these antigens interact simultaneously with MHC proteins on TECs and specific Vβ peptides on DP cells to illicit very strong TCR signaling that triggers apoptosis. The reduced levels of Vβ gene rearrangements in *PG* mice do not alter primary Vβ repertoire, and to our knowledge there is no evidence that differences among Dβ-, Jβ-, or CDR3β encoded nucleotides influence superantigen-directed negative selection. The initiation and progression of iNKT cell development is driven by strong TCR signaling upon recognition of lipid/CD1d complexes. In theory, ongoing *Tcra* gene recombination before Stage 0 or Stage 1 iNKT cells silence RAG expression could replace the invariant Vα14-Jα18 rearrangement and divert these cells into conventional αβ T cells. If this scenario occurs, the lower levels of *Tcra* gene rearrangements in *PG* mice would be predicted to diminish the chance of this happening and thereby promote developmental progression of iNKT cells. However, the slightly altered Dβ and Jβ usage in *Tcrb* gene rearrangements of *PG* mice could generate αβ TCRs less effective at signaling upon recognition of lipid/CD1d complexes, leading to impaired iNKT cell development. The higher autoimmune potential of αβ T cells in *PG* mice must result from impaired negative selection of high self-reactive thymocytes. In mice with normal RAG endonuclease activity, highly self-reactive αβ TCRs signal negative selection by apoptosis before further

Tcra gene recombination can replace these hazardous receptors with new ones (56). The reduced levels of *Tcra* gene rearrangements in *PG* mice would be expected to increase the likelihood that negatively-selecting TCRs remain expressed and trigger apopotisis. Considering all of the above, the most plausible reasons for less efficient negative selection, impaired iNKT cell development, and higher autoimmune potential of CD4$^+$ T cells in *PG* mice are: 1) decreased frequencies or durations that αβ TCRs engage self-antigens due to the 50% fewer DP thymocyes and potentially altered thymic microenvironment, and/or 2) less robust TCR signaling upon recognition of self-antigens. Genetic mutations in mice that substantially impair V(D)J recombination also dramatically (>10-fold) reduce thymic cellularity and consequently alter thymic architecture through defective maturation of TECs and/or distribution of DCs, which collectively impairs negative selection and increases development of hazardous autoreactive αβ T cells (60). Although the slightly lower than normal levels of TCR gene rearrangements in *PG* mice cause only a minor decrease in DP thymocyte cellularity, it will be important to determine if the Rag1^{P326G} mutation has any effects on the thymic microenvironment that could underlie less efficient positive selection, negative selection, and agonist selection.

The lower than normal expression of Zap70 protein specifically in thymocytes of *PG* mice offers support and a potential basis for impaired αβ TCR signaling contributing to less efficient thymocyte selection. As reduced Zap70 activity impairs TCR signaling, positive selection, negative selection, and agonist selection of iNKT cells (57-59), attenuated TCR signaling from lower Zap70 expression, even the subtle decrease observed, is a plausible explanation for these phenotypes of *PG* mice. However, definitive conclusions necessitate elucidating how Rag1^{P326G} mutation decreases Zap70 protein levels and determining effects of disrupting these mechanisms without lowering RAG endonuclease activity. Considering that RAG DSBs induced during Igκ gene recombination in pre-B cells activate signaling pathways that alter the expression of antigen receptor signaling proteins (37, 61), RAG DSBs induced during *Tcra* gene rearrangements in DP thymocytes could trigger nucleus-to-cytoplasm signaling cascades that post-transcriptionally increase

Zap70 protein expression. In this scenario, the reduced number of RAG DSBs generated during *Tcra* gene rearrangements in *PG* thymocytes might impair the activation of such signals. A not-mutally exclusive possibility is that Rag1^{P326G} mutation prevents the Rag1-directed ubiquitylation of nuclear proteins that amplify the signals from RAG DSBs that up-regulate Zap70 protein expression. Notably, we did not detect differences in Zap70 activation or other aspects of TCR signaling between *WT* and *PG* thymocytes after anti-CD3 stimulation (data not shown). However, this approach for activating αβ TCR signaling is not as sensitive for detecting differences as using tetramers or DCs to present peptides to transgenic αβ TCRs of fixed specificity (9, 11, 17). Such experiments, which are beyond the scope of our current study, are critical to determine whether Rag1^{P326G} mutation diminishes αβ TCR signaling spefiically in DP thymocytes. Considering that Zap70 might not be the only, or even the relevant, protein affected by this Rag1 mutation, findings of reduced TCR signaling would warrant identifying all changes in thymocyte protein expression triggered by RAG DSBs and elucidating the effect of Rag1^{P326G} mutation on these. This information would suggest approaches to disrupt signaling from RAG DSBs without reducing RAG endonuclease activity to separate impacts of Rag1^{P326G} mutation on altering expression of TCR signaling proteins versus reducing *Tcrb* and *Tcra* gene rearrangements, altering assembled *Tcrb* and *Tcra* gene repertoires, and decreasing thymic cellularity.

The requirement for TCR selection on self-peptides displayed by self-proteins presents major challenges for establishing conventional and non-conventional αβ T cell populations that recognize and respond to a vast array of diverse foreign peptides, but not self-antigens. An inability to positively-select αβ TCRs with low affinity/avidity for self-peptide/protein complexes could eliminate cells capable of binding foreign antigens with no similarities to self-peptides, generating holes in adaptive immunity. At the same time, any positively-selected TCR is able to recognize a self-peptide and thus could cause autoimmunity. To meet these challenges, cell intrinsic mechanisms have evolved to render thymocytes more sensitive to

TCR stimulation than their mature αβ T cell counterparts (11-15). This "TCR signal tuning" elevates the affinity/avidity thresholds for both positive and negative selection because any functional TCR interaction triggers a stronger signal in thymocytes versus splenic αβ T cells (6-9). This lower sensitivity of mature αβ T cells prevents their activation when encountering self-peptide/protein complexes identical to that on which they were positively-selected. While several factors have been implicated in regulating TCR signal tuning, each is expressed in both thymocytes and mature T cells (17, 63-67). Our observations of reduced thymic Zap70 expression and less efficient positive selection, negative selection, and agonist selection in *PG* mice are consistent with the possibility that the RAG1 protein functions as a thymocyte specific TCR signal tuning factor. In this model, the efficient activation of signaling cascades from RAG DSBs and/or functions of the RAG1 ubiquitin ligase domain might be important to modulate expression of Zap70, and likely many other factors, to raise affinity/avidity thresholds for positive selection and negative selection. Critically, our study provides a phenotypic framework from which to elucidate the molecular mechanisms by which the integrity of the RAG1 ubiquitin ligase domain promotes TCR gene assembly and transcends this process to govern the proper selection of developing αβ T cells.

Acknowledgements

We thank Kyutae Lee for designing the gene-editing strategy and Adele Harman of the CHOP Transgenic Core to generate the NOD *PG* mice. The authors have no competing financial interests to declare.

References

1.	Bassing, C. H., W. Swat, and F. W. Alt. 2002. The mechanism and regulation of chromosomal V(D)J recombination. *Cell* 109 Suppl: S45-55.

2.	Schatz, D. G., and P. C. Swanson. 2011. V(D)J recombination: mechanisms of initiation. *Annu Rev Genet* 45: 167-202.

3.	Bhandoola, A., H. von Boehmer, H. T. Petrie, and J. C. Zuniga-Pflucker. 2007. Commitment and developmental potential of extrathymic and intrathymic T cell precursors: plenty to choose from. *Immunity* 26: 678-689.

4.	von Boehmer, H., and F. Melchers. 2010. Checkpoints in lymphocyte development and autoimmune disease. *Nat Immunol* 11: 14-20.

5.	Carico, Z., and M. S. Krangel. 2015. Chromatin Dynamics and the Development of the TCRalpha and TCRdelta Repertoires. *Adv Immunol* 128: 307-361.

6.	Hogquist, K. A., and S. C. Jameson. 2014. The self-obsession of T cells: how TCR signaling thresholds affect fate 'decisions' and effector function. *Nat Immunol* 15: 815-823.

7.	Gascoigne, N. R., V. Rybakin, O. Acuto, and J. Brzostek. 2016. TCR Signal Strength and T Cell Development. *Annu Rev Cell Dev Biol* 32: 327-348.

8.	Klein, L., B. Kyewski, P. M. Allen, and K. A. Hogquist. 2014. Positive and negative selection of the T cell repertoire: what thymocytes see (and don't see). *Nat Rev Immunol* 14: 377-391.

9.	Fu, G., V. Rybakin, J. Brzostek, W. Paster, O. Acuto, and N. R. Gascoigne. 2014. Fine-tuning T cell receptor signaling to control T cell development. *Trends Immunol* 35: 311-318.

10.	Krovi, S. H., and L. Gapin. 2018. Invariant Natural Killer T Cell Subsets-More Than Just Developmental Intermediates. *Front Immunol* 9: 1393.

11.	Davey, G. M., S. L. Schober, B. T. Endrizzi, A. K. Dutcher, S. C. Jameson, and K. A. Hogquist. 1998. Preselection thymocytes are more sensitive to T cell receptor stimulation than mature T cells. *J Exp Med* 188: 1867-1874.

12.	Lucas, B., I. Stefanova, K. Yasutomo, N. Dautigny, and R. N. Germain. 1999. Divergent changes in the sensitivity of maturing T cells to structurally related ligands underlies formation of a useful T cell repertoire. *Immunity* 10: 367-376.

13.	Yachi, P. P., C. Lotz, J. Ampudia, and N. R. Gascoigne. 2007. T cell activation enhancement by endogenous pMHC acts for both weak and strong agonists but varies with differentiation state. *J Exp Med* 204: 2747-2757.

14.	Pircher, H., U. H. Rohrer, D. Moskophidis, R. M. Zinkernagel, and H. Hengartner. 1991. Lower receptor avidity required for thymic clonal deletion than for effector T-cell function. *Nature* 351: 482-485.

15.	Sant'Angelo, D. B., and C. A. Janeway, Jr. 2002. Negative selection of thymocytes expressing the D10 TCR. *Proc Natl Acad Sci U S A* 99: 6931-6936.

16.	Choi, S., R. Cornall, R. Lesourne, and P. E. Love. 2017. THEMIS: Two Models, Different Thresholds. *Trends Immunol* 38: 622-632.

17.	Fu, G., J. Casas, S. Rigaud, V. Rybakin, F. Lambolez, J. Brzostek, J. A. Hoerter, W. Paster, O. Acuto, H. Cheroutre, K. Sauer, and N. R. Gascoigne. 2013. Themis sets the signal threshold for positive and negative selection in T-cell development. *Nature* 504: 441-445.

18.	Jones, J. M., and M. Gellert. 2003. Autoubiquitylation of the V(D)J recombinase protein RAG1. *Proc Natl Acad Sci U S A* 100: 15446-15451.

19.	Singh, S. K., and M. Gellert. 2015. Role of RAG1 autoubiquitination in V(D)J recombination. *Proc Natl Acad Sci U S A* 112: 8579-8583.

20.	Jones, J. M., A. Bhattacharyya, C. Simkus, B. Vallieres, T. D. Veenstra, and M. Zhou. 2011. The RAG1 V(D)J recombinase/ubiquitin ligase promotes ubiquitylation of acetylated, phosphorylated histone 3.3. *Immunol Lett* 136: 156-162.

21. Grazini, U., F. Zanardi, E. Citterio, S. Casola, C. R. Goding, and F. McBlane. 2010. The RING domain of RAG1 ubiquitylates histone H3: a novel activity in chromatin-mediated regulation of V(D)J joining. *Mol Cell* 37: 282-293.

22. Deng, Z., H. Liu, and X. Liu. 2015. RAG1-mediated ubiquitylation of histone H3 is required for chromosomal V(D)J recombination. *Cell Res* 25: 181-192.

23. Simkus, C., P. Anand, A. Bhattacharyya, and J. M. Jones. 2007. Biochemical and folding defects in a RAG1 variant associated with Omenn syndrome. *J Immunol* 179: 8332-8340.

24. Villa, A., C. Sobacchi, L. D. Notarangelo, F. Bozzi, M. Abinun, T. G. Abrahamsen, P. D. Arkwright, M. Baniyash, E. G. Brooks, M. E. Conley, P. Cortes, M. Duse, A. Fasth, A. M. Filipovich, A. J. Infante, A. Jones, E. Mazzolari, S. M. Muller, S. Pasic, G. Rechavi, M. G. Sacco, S. Santagata, M. L. Schroeder, R. Seger, D. Strina, A. Ugazio, J. Valiaho, M. Vihinen, L. B. Vogler, H. Ochs, P. Vezzoni, W. Friedrich, and K. Schwarz. 2001. V(D)J recombination defects in lymphocytes due to RAG mutations: severe immunodeficiency with a spectrum of clinical presentations. *Blood* 97: 81-88.

25. Zheng, N., P. Wang, P. D. Jeffrey, and N. P. Pavletich. 2000. Structure of a c-Cbl-UbcH7 complex: RING domain function in ubiquitin-protein ligases. *Cell* 102: 533-539.

26. Albert, T. K., H. Hanzawa, Y. I. Legtenberg, M. J. de Ruwe, F. A. van den Heuvel, M. A. Collart, R. Boelens, and H. T. Timmers. 2002. Identification of a ubiquitin-protein ligase subunit within the CCR4-NOT transcription repressor complex. *EMBO J* 21: 355-364.

27. Schwenk, F., U. Baron, and K. Rajewsky. 1995. A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. *Nucleic Acids Res* 23: 5080-5081.

28. Lee, K. D., and C. H. Bassing. 2020. Two Successive Inversional Vbeta Rearrangements on a Single Tcrb Allele Can Contribute to the TCRbeta Repertoire. *J Immunol* 204: 78-86.

29. Gopalakrishnan, S., K. Majumder, A. Predeus, Y. Huang, O. I. Koues, J. Verma-Gaur, S. Loguercio, A. I. Su, A. J. Feeney, M. N. Artyomov, and E. M. Oltz. 2013. Unifying model for molecular determinants of the preselection Vbeta repertoire. *Proc Natl Acad Sci U S A* 110: E3206-3215.

30. Shih, H. Y., J. Verma-Gaur, A. Torkamani, A. J. Feeney, N. Galjart, and M. S. Krangel. 2012. Tcra gene recombination is supported by a Tcra enhancer- and CTCF-dependent chromatin hub. *Proc Natl Acad Sci U S A* 109: E3493-3502.

31. Chen, L., Z. Carico, H. Y. Shih, and M. S. Krangel. 2015. A discrete chromatin loop in the mouse Tcra-Tcrd locus shapes the TCRdelta and TCRalpha repertoires. *Nat Immunol* 16: 1085-1093.

32. Ostanin, D. V., J. Bao, I. Koboziev, L. Gray, S. A. Robinson-Jackson, M. Kosloski-Davidson, V. H. Price, and M. B. Grisham. 2009. T cell transfer model of chronic colitis: concepts, considerations, and tricks of the trade. *Am J Physiol Gastrointest Liver Physiol* 296: G135-146.

33. Liang, H. E., L. Y. Hsu, D. Cado, L. G. Cowell, G. Kelsoe, and M. S. Schlissel. 2002. The "dispensable" portion of RAG2 is necessary for efficient V-to-DJ rearrangement during B and T cell development. *Immunity* 17: 639-651.

34. Horowitz, J. E., and C. H. Bassing. 2014. Noncore RAG1 regions promote Vbeta rearrangements and alphabeta T cell development by overcoming inherent inefficiency of Vbeta recombination signal sequences. *J Immunol* 192: 1609-1619.

35. Colucci, F., D. Guy-Grand, A. Wilson, M. Turner, E. Schweighoffer, V. L. Tybulewicz, and J. P. Di Santo. 2000. A new look at Syk in alpha beta and gamma delta T cell development using chimeric mice with a low competitive hematopoietic environment. *J Immunol* 164: 5140-5145.

36. Palacios, E. H., and A. Weiss. 2007. Distinct roles for Syk and ZAP-70 during early thymocyte development. *J Exp Med* 204: 1703-1715.

37. Bednarski, J. J., R. Pandey, E. Schulte, L. S. White, B. R. Chen, G. J. Sandoval, M. Kohyama, M. Haldar, A. Nickless, A. Trott, G. Cheng, K. M. Murphy, C. H. Bassing, J. E. Payton, and B. P.

Sleckman. 2016. RAG-mediated DNA double-strand breaks activate a cell type-specific checkpoint to inhibit pre-B cell receptor signals. *J Exp Med* 213: 209-223.

38.   Williams, J. A., K. S. Hathcock, D. Klug, Y. Harada, B. Choudhury, J. P. Allison, R. Abe, and R. J. Hodes. 2005. Regulated costimulation in the thymus is critical for T cell development: dysregulated CD28 costimulation can bypass the pre-TCR checkpoint. *J Immunol* 175: 4199-4207.

39.   Kimura, M. Y., J. Thomas, X. Tai, T. I. Guinter, M. Shinzawa, R. Etzensperger, Z. Li, P. Love, T. Nakayama, and A. Singer. 2016. Timing and duration of MHC I positive selection signals are adjusted in the thymus to prevent lineage errors. *Nat Immunol* 17: 1415-1423.

40.   Karo, J. M., D. G. Schatz, and J. C. Sun. 2014. The RAG recombinase dictates functional heterogeneity and cellular fitness in natural killer cells. *Cell* 159: 94-107.

41.   Bednarski, J. J., A. Nickless, D. Bhattacharya, R. H. Amin, M. S. Schlissel, and B. P. Sleckman. 2012. RAG-induced DNA double-strand breaks signal through Pim2 to promote pre-B cell survival and limit proliferation. *J Exp Med* 209: 11-17.

42.   Negishi, I., N. Motoyama, K. Nakayama, K. Nakayama, S. Senju, S. Hatakeyama, Q. Zhang, A. C. Chan, and D. Y. Loh. 1995. Essential role for ZAP-70 in both positive and negative selection of thymocytes. *Nature* 376: 435-438.

43.   Zhao, M., M. N. D. Svensson, K. Venken, A. Chawla, S. Liang, I. Engel, P. Mydel, J. Day, D. Elewaut, N. Bottini, and M. Kronenberg. 2018. Altered thymic differentiation and modulation of arthritis by invariant NKT cells expressing mutant ZAP70. *Nat Commun* 9: 2627.

44.   Jordan, M. S., J. E. Smith, J. C. Burns, J. E. Austin, K. E. Nichols, A. C. Aschenbrenner, and G. A. Koretzky. 2008. Complementation in trans of altered thymocyte development in mice expressing mutant forms of the adaptor molecule SLP76. *Immunity* 28: 359-369.

45.   Surh, C. D., and J. Sprent. 1994. T-cell apoptosis detected in situ during positive and negative selection in the thymus. *Nature* 372: 100-103.

46.   Wilson, A., C. Marechal, and H. R. MacDonald. 2001. Biased V beta usage in immature thymocytes is independent of DJ beta proximity and pT alpha pairing. *J Immunol* 166: 51-57.

47.   Rossjohn, J., D. G. Pellicci, O. Patel, L. Gapin, and D. I. Godfrey. 2012. Recognition of CD1d-restricted antigens by natural killer T cells. *Nat Rev Immunol* 12: 845-857.

48.   Dashtsoodol, N., S. Bortoluzzi, and M. Schmidt-Supprian. 2019. T Cell Receptor Expression Timing and Signal Strength in the Functional Differentiation of Invariant Natural Killer T Cells. *Front Immunol* 10: 841.

49.   Jiang, W., M. S. Anderson, R. Bronson, D. Mathis, and C. Benoist. 2005. Modifier loci condition autoimmunity provoked by Aire deficiency. *J Exp Med* 202: 805-815.

50.   Jin, Y., A. Lee, J. H. Oh, H. W. Lee, and S. J. Ha. 2019. The R229Q mutation of Rag2 does not characterize severe immunodeficiency in mice. *Sci Rep* 9: 4415.

51.   Mottet, C., H. H. Uhlig, and F. Powrie. 2003. Cutting edge: cure of colitis by CD4+CD25+ regulatory T cells. *J Immunol* 170: 3939-3943.

52.   Eri, R., M. A. McGuckin, and R. Wadley. 2012. T cell transfer model of colitis: a great tool to assess the contribution of T cells in chronic intestinal inflammation. *Methods Mol Biol* 844: 261-275.

53.   Anderson, M. S., and J. A. Bluestone. 2005. The NOD mouse: a model of immune dysregulation. *Annu Rev Immunol* 23: 447-485.

54.   Delovitch, T. L., and B. Singh. 1997. The nonobese diabetic mouse as a model of autoimmune diabetes: immune dysregulation gets the NOD. *Immunity* 7: 727-738.

55.   Huang, C. Y., B. P. Sleckman, and O. Kanagawa. 2005. Revision of T cell receptor {alpha} chain genes is required for normal T lymphocyte development. *Proc Natl Acad Sci U S A* 102: 14356-14361.

56. Kreslavsky, T., H. J. Kim, S. B. Koralov, D. Ghitza, T. Buch, H. Cantor, K. Rajewsky, and H. von Boehmer. 2013. Negative selection, not receptor editing, is a physiological response of autoreactive thymocytes. *J Exp Med* 210: 1911-1918.

57. Sakaguchi, N., T. Takahashi, H. Hata, T. Nomura, T. Tagami, S. Yamazaki, T. Sakihama, T. Matsutani, I. Negishi, S. Nakatsuru, and S. Sakaguchi. 2003. Altered thymic T-cell selection due to a mutation of the ZAP-70 gene causes autoimmune arthritis in mice. *Nature* 426: 454-460.

58. Mallaun, M., G. Zenke, and E. Palmer. 2010. A discrete affinity-driven elevation of ZAP-70 kinase activity initiates negative selection. *J Recept Signal Transduct Res* 30: 430-443.

59. Hsu, L. Y., Y. X. Tan, Z. Xiao, M. Malissen, and A. Weiss. 2009. A hypomorphic allele of ZAP-70 reveals a distinct thymic threshold for autoimmune disease versus autoimmune reactivity. *J Exp Med* 206: 2527-2541.

60. Marrella, V., P. L. Poliani, L. D. Notarangelo, F. Grassi, and A. Villa. 2014. Rag defects and thymic stroma: lessons from animal models. *Front Immunol* 5: 259.

61. Bredemeyer, A. L., B. A. Helmink, C. L. Innes, B. Calderon, L. M. McGinnis, G. K. Mahowald, E. J. Gapud, L. M. Walker, J. B. Collins, B. K. Weaver, L. Mandik-Nayak, R. D. Schreiber, P. M. Allen, M. J. May, R. S. Paules, C. H. Bassing, and B. P. Sleckman. 2008. DNA double-strand breaks activate a multi-functional genetic program in developing lymphocytes. *Nature* 456: 819-823.

62. Van, H. T., and M. A. Santos. 2018. Histone modifications and the DNA double-strand break response. *Cell Cycle* 17: 2399-2410.

63. Fu, G., S. Vallee, V. Rybakin, M. V. McGuire, J. Ampudia, C. Brockmeyer, M. Salek, P. R. Fallen, J. A. Hoerter, A. Munshi, Y. H. Huang, J. Hu, H. S. Fox, K. Sauer, O. Acuto, and N. R. Gascoigne. 2009. Themis controls thymocyte selection through regulation of T cell antigen receptor-mediated signaling. *Nat Immunol* 10: 848-856.

64. Lesourne, R., S. Uehara, J. Lee, K. D. Song, L. Li, J. Pinkhasov, Y. Zhang, N. P. Weng, K. F. Wildt, L. Wang, R. Bosselut, and P. E. Love. 2009. Themis, a T cell-specific protein important for late thymocyte development. *Nat Immunol* 10: 840-847.

65. Johnson, A. L., L. Aravind, N. Shulzhenko, A. Morgun, S. Y. Choi, T. L. Crockford, T. Lambe, H. Domaschenz, E. M. Kucharska, L. Zheng, C. G. Vinuesa, M. J. Lenardo, C. C. Goodnow, R. J. Cornall, and R. H. Schwartz. 2009. Themis is a member of a new metazoan gene family and is required for the completion of thymocyte positive selection. *Nat Immunol* 10: 831-839.

66. Wang, D., M. Zheng, L. Lei, J. Ji, Y. Yao, Y. Qiu, L. Ma, J. Lou, C. Ouyang, X. Zhang, Y. He, J. Chi, L. Wang, Y. Kuang, J. Wang, X. Cao, and L. Lu. 2012. Tespa1 is involved in late thymocyte development through the regulation of TCR-mediated signaling. *Nat Immunol* 13: 560-568.

67. Li, Q. J., J. Chau, P. J. Ebert, G. Sylvester, H. Min, G. Liu, R. Braich, M. Manoharan, J. Soutschek, P. Skare, L. O. Klein, M. M. Davis, and C. Z. Chen. 2007. miR-181a is an intrinsic modulator of T cell sensitivity and selection. *Cell* 129: 147-161.

Footnotes

This work was supported by the Fondation ARC pour la recherche sur le cancer (C.M.), La Ligue contre le cancer (C.M.), Fondation Pasteur Mutualite (C.M.), Nancy Taylor Foundation for Chronic Disease (E.M.B.) and NIH grants R03 AI097611 (J.M.J.), R01 HL112836 (E.M.B.), R01 AI121250-A1 (E.M.B.), R21 A1 135435 (C.H.B.), and RO1 AI 112621 (C.H.B.).

Figure 1. *PG* mice exhibit impaired thymocyte developmental transitions that require TCR gene assembly. **(A)** Representative flow cytometry plots of DN stages of thymocyte development showing the frequency of cells at each stage. **(B)** Quantification of numbers of cells in each DN stage. **(C)** Representative flow cytometry plots of DN, DP, CD4$^+$ SP, and CD8$^+$ SP thymocytes. **(D)** Quantification of numbers of cells in each stage. **(E)** Quantification of the ratio of CD4$^+$ SP or CD8$^+$ SP thymocytes to DP thymocytes. **(F-G)** Quantification of the numbers of total thymocytes (F) and splenic αβ T cells (G). (A-G) All data were collected from 6-10 weeks old mice and (B, D, E, F, and G) are combined from four independent experiments including a total of 14-16 mice per genotype. Bars indicate mean +/- SEM. Stats: 2-way ANOVA with Sidak multiple comparison test. n.s. $p > 0.05$; * $p < 0.05$, **$p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure 2. Thymocytes of *PG* mice support diminished levels of *Tcrb* and *Tcra* recombination. **(A-B)** Taqman qPCR quantification of rearrangements of each Vβ segment to DJβ1.1 (A) or DJβ2.1 (B) complexes conducted on genomic DNA from DN3 thymocytes. Signals from each assay were normalized to values from an assay for the invariant CD19 gene. Shown are average values +/- SEM from three independent DN3 thymocyte preparations, each from a different mouse. **(C-E)** Adaptive Immunoseq quantification of the percentage usage of each Vβ (C), Dβ (D), or Jβ (E) segment in *Tcrb* rearrangements of DN3 thymocytes. Shown are average values from two independent DN cell preparations. **(F)** qPCR quantification of specific Vα/Jα rearrangements peformed on genomic DNA from DP thymocytes. Signals from each assay were normalized to values from an assay for the invariant β2m gene. Shown are the average values +/- SEM from three independent DP cell preparations. For all graphs in figure: #, not detected, n.s. $p > 0.05$; * $p < 0.05$, **$p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$

Figure 3. *PG* DN3 thymocytes exhibit reduced Syk protein expression but normal efficiency of pre-TCR signaled differentiation. **(A-B)** Representative flow cytometry histogram plots (A) and quantification (B) of Syk protein in DN3 thymocytes. (B) Shown are all data points and average values +/- SEM from one of two independent experiments (n = 3-4 mice per genotype, unpaired t-test). **(C)** qRT-PCR quantification of Syk mRNA in DN3 cells. Relative mRNA levels were calculated using ddCt method with signals from each assay normalized to values from an assay for the *Hprt* gene. **(B-C)** Shown are all data points and average values +/- SEM from two independent experiments (n = 8-9 mice per genotype, 2-way ANOVA with experiment an independent variable, stats indicate difference as a result of genotype). **(D)** Representative flow cytometry plots of DN3a, DN3b, and DN3c stages of thymocyte development showing the frequency of cells at each stage. **(E-F)** Quantification of percentages (E) or numbers (F) of total DN3 cells in each stage. Shown are all data points and average values +/- SEM from three independent experiments each with three mice of each genotype (n = 9 mice per genotype, analyzed by multiple t tests using the Holm-Sidak method; n.s. $p>0.05$, * $p<0.05$, ** $p<0.005$, **** $p<0.00005$). **(G)** Quantification of percentages of DN3a cells that progress into DN3b cells. Shown are all data points and average values +/- SEM from three independent experiments each with three mice of each genotype (n = 9 mice per genotype, analyzed by unpaired t-test with Welch's correction; ** $p<0.005$). **(H)** Representative flow cytometry plots of intracellular TCRβ protein expression in DN3a thymocytes. **(I)** Quantification of percentages of DN3a cells expressing intracellular TCRβ protein. **(J)** Ratios comparing the percentages of DN3a cells expressing intracellular TCRβ protein to the percentages of DN3a cells that transition into DN3b cells. **(H-J)** Shown are all data points and average values +/- SEM from three independent experiments each with three mice of each genotype (n = 9 mice per genotype, analyzed by unpaired t-test with Welch's correction; n.s. $p>0.05$, *** $p<0.0005$).

Figure 4. Thymocytes of PD mice have reduced Zap70 protein levels. **(A-D)** Representative flow cytometry histogram plots (A, C) and quantification (B, D) of Zap70 (A, B) and TCRβ (C, D) protein in

Stages 1-4 of DP thymocyte selection. (C, D) Shown are all data points and average values +/- SEM from one of three independent experiments (n = 4-5 mice per genotype, 2-way ANOVA with Sidak's multiple comparison post-test) **(E)** qRT-PCR quantification of Zap70 mRNA in DN3 cells or Stage 2 or 4 DP cells. Relative mRNA levels were calculated using ddCt method with signals from each assay normalized to values from an assay for the Hprt gene. Shown are all data points and average values +/- SEM from one experiment (n = 4-5 mice per genotype, 2-way ANOVA with sidak multiple comparison post-test. **(F-G)** Representative flow cytometry histogram plots (F) and quantification (G) of Zap70 protein in naïve CD4$^+$ or CD8$^+$ T cells. Shown are all data points and average values +/- SEM from two independent experiments (n = 3 mice per genotype, 2-way ANOVA with Sidak multiple comparison post-test). For all graphs in figure: n.s. $p > 0.05$; * $p < 0.05$, **$p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure 5. *PG* mice exhibit impaired positive selection. **(A)** Gating strategy for measuring positive selection of thymocytes showing Stage 1 and 2 gates. Pre-gated on live singlets and CD4$^+$ and/or CD8$^+$. **(B)** Quantification of the ratio of the number of Stage 2 cells to the total number of Stage 1 and 2 cells. Shown are all data points and global mean from three independent experiments (n = 10-11 mice per genotype, 2-way ANOVA indicating effect of genotype). **(C)** Quantification of numbers of cells in each stage of positive selection. Dots indicate individual mice. Data pooled from 2-independent experiments (n = 7-8 mice per genotype, bars indicate mean +/- SEM analyzed by Man-Whitney test). **(D)** Quantification of CD69 expression alone and in comparison to TCRβ expression on cells at each stage of positive selection. Dots indicate individual mice. Data pooled from 2-independent experiments (n = 8 mice per genotype, bars indicate mean +/- SEM. Analysed by 2-way ANOVA and Sidak multiple comparison post-test) For all graphs in figure: n.s. $p > 0.05$; * $p < 0.05$, **$p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure 6. *PG* mice exhibit impaired superantigen-mediated negative selection. **(A)** Gating strategy form

measuring negative selection of specific Vβ$^+$ thymocytes in F1 progeny of C57BL/6 background *WT* or

PG mice crossed with BALB/c background *Rag1$^{-/-}$* mice. Pre-gated on live singlets. **(B-F)** Representative

analyses and quantification of the indicated Vβ$^+$ cells in pre-selection or post-selection thymocytes. **(G)**

Data from B-F displayed as LOG$_2$ ratio of respective percent Vβ$^+$ cells in SP versus DP gates. Shown are

all data points and average +/- SEM from two independent experiments (n = 9 mice per genotype 2-way

ANOVA with Sidak multiple comparison post-test). For any significantly different combinations, fold

change is indicated. For all graphs in figure: * p<0.05, **** p<0.0001.

Figure 7. *PG* mice have impaired agonist selection of iNKT cells. (**A-B**) Representative flow cytometry

plots (**A**) and quantification (**B**) of PBS57/CD1d tetramer binding iNKT cells in thymuses or spleeens.

Gated on live, singlet lymphocytes. (**B**) Data from three independent experiments with 1-4 mice per group.

Mann-Whitney test: *, p=0.01 (thymus); ns, not significant (spleen). (**C-D**) Representative flow

cytometry plots and gating strategy (**C**) or quantification (**D**) of Stage 0 (CD24^{hi} CD44-NK1.1-), Stage 1

(CD44-NK1.1-), Stage 2 (CD44+NK1.1-), and Stage 3 (CD44+NK1.1+) iNKT cells. (**D**) Data from threee

independent experiments with 1-4 mice per group. Mann-Whitney test: **, p<0.01. ns, not significant.

Ed- The BM chimera data are not shown in results or described here.

Figure 8. Mature αβ T cells of PG mice exhibit normal immune response. **(A)** Representative plot of

cytokine expression on CD4$^+$CD44^{hi} αβ T cells following *ex vivo* stimulation of splenocytes with PMA

and ionomycin. **(B-F)** Representative analyses of naïve splenic CD4$^+$ or CD8$^+$ T cells after *ex vivo*

simulation with anti-CD3/anti-CD28 antibodies and additional antibodies to skew towards indicated

lineages. Shown is CFSE incorporation (B) or expression of relevant transcription factors (C, D) or

cytokines (E, F). **(G, H)** Representative flow data showing (G) frequencies of splenic αβ T cells that stain

with the gp33:H-2Db tetramer or (H) express indicated cytokines after LCMV infection of mice. Data are representative of two independent experiments with at least four mice of each genotype. **(I, J)** Representative plots (J) and (I) quantification of antigen-specific αβ T cells able to respond to indicated viral peptides within a peptide re-stimulation assay. (A-F) Data are representative of two independent experiments. (G-J) Data are combined from four independent experiments (two D7 and two D35 timepoints) with 8-11 mice of each genotype per time-point.

Figure 9. Mature αβ T cells of *PG* mice possess higher than normal autoimmune hazard**. (A)** Graph of body weights of *Rag1⁻/⁻* hosts over time after transfer of *WT* or *PG* CD4⁺ T cells or no cells (NT). Data combined from three independent experiments with a total of 13-17 mice of each genotype. Lines indicate fitted segmented-linear regression lines +/- 95% confidence interval. Day 15 was fixed as the inflection point. Difference between gradients of second slope was determined by linear regression analysis. **(B)** Numbers of CD4⁺ T cells in spleen eight weeks post-transfer. **(C)** Total number of IFNγ, TNFα, and/or GM-CSF producing cells per spleen. Shown are all data points from three independent experiments with 12-14 mice per group. Bar indicates global mean. Pre-gating on live singlets, CD90.2⁺, CD4⁺, and CD44⁺. **(D)** Colon length of mice sacrificed at eight weeks post-infection. **(E)** Representative hemotoxylin and eosin stained colon sections from respective mice at eitght weeks post-transfer. 10x magnification. **(F-G)** Total colitis score as assessed by blinded pathologist. Scored as per previously described guidelines for the five phenotypes (52): (F) Colon inflammation score (contribution of scores for crypt abcesses and inflammatory cell infiltrate). (G) Colon tissue damage score (contribution of scores for crypt architecture, tissue damage, goblet cell loss). **(H)** Non-colitis pathologies described for mice transferred with *WT* versus *PG* cells. **(I)** Hemotoxylin and eosin stained skin section from *PG*-transferred mouse with severe dermatitis at eight weeks post-transfer. 10x magnification. **(J)** Spleen weight at eight weeks post-transfer normalized to total mouse bodyweight prior to transfer. **(K)** Survival curves of mice transferred with *WT* or *PG* cells. Data combined from three independent experiments. Statistical analysis = Log-rank Mantel-

Cox test. Overall: 0/13 NT mice died, 1/15 *WT*-transferred mice died, and 4/17 *PG*-transferred mice died. (B, C, F, G, and J) Shown are all data points from three independent experiments, 12-14 mice per group. Experiment indicated by shape of data point. Bar indicates global mean. Analyzed by 2-way ANOVA with Tukey HSD post-test. n.s. $p > 0.05$; * $p < 0.05$, **$p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ (**L**) Curves showing diabetes free survival of female NOD *WT* (n=16) and *PG* (n=14) mice. Mice were generated by mating together NOD *Rag1$^{WT/PG}$* mice and monitored for their first 30 weeks of life. The percent diabetes survival curves include mice that have passed 20 weeks of age or gotten diabetes before 20 weeks of age. Analyzed by Gehan-Breslow-Wilcoxon test (p=0.049) and log-rank Mantel Cox test (p=0.1151).

Figure 1

Figure 2

Figure 3

Figure 3

Figure 5

Figure 5

Figure 7

Figure 8

Figure 9

Supplemental Materials

Supplemental Figure 1. *PG* thymocytes exbibit cell intrinsic deficiencies in developmental progression. **(A-B)** Shown are percent of donor *PG* (CD45.2$^+$) or *WT* (CD45.1$^+$) cells within indicated (A) thymocyte or (B) splenocyte populations after transfer of a 1:1 mix of *PG* and *WT* bone marrow into lethally irradiated *WT* congenic (CD45.1$^+$) recipients. Bone marrow was pooled from three donors of each genotype and injected into five recipients, all analyzed on the same day. Data represented as mean +/- SEM. Statistics calculated with multiple t tests using the Holm-Sidak method. *p<0.05, ***p<0.0005, ****p<0.00005. **(C)** Shown are percent donor *PG* (CD45.2$^+$) chimerism in indicated compartments after lethal irradiation of *WT* congenic (CD45.1$^+$) recipients. Open circles show percent *PG* chimerism after transfer of only *PG* bone marrow, while filled circles show *PG* chimerism in a 1:1 competitive setting with *WT* cells. *PG* bone marrow reconstitutes the blood NK cell compartment at expected ratios. In mice receiving only *PG* bone marrow, *PG* chimerism in bulk splenic iNKT (p<0.0001), bulk thymic iNKT (p<0.0001), Stage 2 iNKT (p=0.001), and Stage 3 iNKT (p<0.0001) are all significantly reduced relative to blood NK cells by one-way ANOVA with Dunnett multiple comparisons test. In mice receiving 1:1 mixed *PG* and *WT* bone marrow, *PG* chimerism in bulk splenic iNKT (p=0.0002), bulk thymic iNKT (p<0.0001), Stage 0 iNKT (p<0.0001), Stage 1 iNKT (p<0.0001), Stage 2 iNKT (p=0.001), and Stage 3 iNKT (p<0.0001) are all significantly reduced relative to blood NK cells. Bone marrow was pooled from three donors and injected into 4-5 mice per group, all analyzed on the same day.

Supplemental Figure 2. *PG* mice exhibit normal efficiency of pre-TCR signaled DN3-to-DN4 thymocyte differentiation. **(A)** Representative flow cytometry plots of DN stages of thymocyte development showing the frequency of cells at the DN3 and DN4 stages. **(B)** Quantification of percentages of DN3 cells that transition into DN4 cells. Shown are all data points and average values +/- SEM from three independent experiments, each with three mice of each genotype (n = 9 mice per genotype, analyzed by multiple t tests using the Holm-Sidak method; *** $p < 0.0005$). **(C)** Representative flow cytometry plots of intracellular TCRβ protein expression in DN3 thymocytes showing the frequency of TCRβ+ cells in each gate. **(D)** Quantification of percentages of DN3 cells expressing intracellular TCRβ protein. **(E)** Ratios comparing the percentages of DN3 cells expressing intracellular TCRβ protein to the percentages of DN3 cells that transition into DN4 cells. (D-E) Shown are all data points and average values +/- SEM from three independent experiments each with three mice of each genotype (n = 9 mice per genotype, analyzed by unpaired t-test with Welch's correction; n.s. $p > 0.05$, *** $p < 0.0005$).

Supplemental Figure 3. Example gating strategy for Stage 1 - Stage 4 selecting thymocytes as depicted in Figure 4 and 5. Pre-gating: Live, singlets, CD4$^+$ AND/OR CD8$^+$. Stage 1 = CD69$^-$CCR7$^-$, Stage 2 = CD69$^+$CCR7$^-$, Stage 3 = CD69$^+$CCR7^{int}, Stage 4 = CD69$^+$CCR7^{hi}.