1

The RAG1 Ubiquitin Ligase Domain Enhances T Cell Receptor Gene Assembly

and Thymic Selection

Thomas N. Burn*,†,**; Charline Miot‡,**; Portia Kreiger‡; Katharina E. Hayer§; Anamika
Bhattacharyya¶,‖, Jessica M. Jones¶, Craig H. Bassing*,‡,#,††; Edward M. Behrens*,†,††

*Institute for Immunology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104
†Division of Rheumatology, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
‡Department of Pathology and Laboratory Medicine, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
§Department of Biomedical and Health Bioinformatics, The Children's Hospital of Philadelphia, Philadelphia, PA 19104
¶Department of Biochemistry and Molecular and Cellular Sciences, Georgetown University, Washington, DC, 20057
‖Present Address: Vyome Therapeutics Lt., 501 Patparganj Industrial Area, Dehli, India
#Department of Pathology and Laboratory Medicine, The Perelman School of Medicine of the University of Pennsylvania, Philadelphia, PA 19104

**Co-first Authors

††Co-corresponding Authors:
        Craig H. Bassing, Ph.D.
        Children's Hospital of Philadelphia
        4054 Colket Translational Research Building
        3501 Civic Center Blvd.
        Philadelphia, PA 19104
        Phone: 267-426-0311
        Email: bassing@email.chop.edu

        Edward M. Behrens, M.D.
        Children's Hospital of Philadelphia
        1102 Abramson Research Center
        3615 Civic Center Boulevard
        Philadelphia, PA 19104
        Phone: 267-426-0142
        Email: behrens@email.chop.edu

Abstract

RAG1/RAG2 (RAG) endonuclease-mediated assembly of diverse lymphocyte antigen receptor genes by

V(D)J recombination is critical for the development and immune function of T and B cells. However, this

process creates highly self-reactive cells that must be negatively selected to suppress autoimmunity. The

RAG1 protein contains a ubiquitin ligase domain that stabilizes RAG1 and stimulates RAG endonuclease

activity. We report here that mice lacking RAG1 ubiquitin ligase activity exhibit diminished

recombination of T cell receptor (TCR) β and α loci, and impaired thymocyte developmental transitions

that require the assembly of these genes and signaling by their proteins. The mice also have reduced

expression of TCR signaling proteins within thymocytes, less efficient negative selection of highly self-

reactive thymocytes, and mature αβ T cells of elevated autoimmune potential. Thus, we propose that the

RAG1 ubiquitin ligase domain provides αβ T cell developmental stage-specific means to augment TCR

signaling and thereby enhance selection for beneficial TCR genes and against αβ TCRs possessing high

autoimmune potential.

Introduction

The ability of T and B lymphocyte populations to express antigen receptors able to recognize a potentially unlimited number of distinct pathogens is the fundamental basis of adaptive immunity. RAG1/RAG2 (RAG) endonuclease-mediated assembly of T cell receptor (TCR) and immunoglobulin (Ig) genes through V(D)J recombination establishes this vast pool of diverse receptors. Germline TCR and Ig loci consist of variable (V), joining (J), and sometimes diversity (D), gene segments residing upstream of constant (C) region exons. During T and B cell development, the lymphocyte-specific RAG complex cooperates with lineage- and developmental stage-specific activation of TCR and Ig loci to perform V(D)J recombination (1). RAG induces DNA double strand breaks (DSBs) between coding sequences and flanking recombination signal sequences of two participating gene segments, liberating hairpin-sealed coding ends and blunt signal ends (2). In a subsequent reaction phase guided by RAG, DNA repair factors ligate signal ends to generate a signal join and open, process, and ligate coding ends to create a V(D)J coding join, which comprises the second exon of an assembled V(D)J-C antigen receptor gene (2). The high number of V(D)J joining events, broad utilization of gene segments, and inherent imprecision in opening and processing coding ends all cooperate to yield a vast population and diversity of receptors. Due to imprecise means of opening and processing coding ends, the outcomes of V(D)J recombination include out-of-frame genes unable to make protein and receptors that recognize self-antigens and have potential for autoimmunity. Accordingly, developing T and B lymphocytes engage quality control checkpoints to select for potentially beneficial antigen receptor proteins and against hazardous highly self-reactive receptors.

Both immature T and B cells employ developmental programs that link functional protein expression from in-frame V(D)J rearrangements to signal either survival and continued differentiation or apoptosis (1). The maturation of $\alpha\beta$ T cells in the thymus provides a paradigm for how V(D)J recombination and antigen receptor protein quality control checkpoints cooperate to create a large pool of lymphocytes with diverse

receptors that provides immunity from pathogens without causing overt autoimmunity. After entering the

thymus, early thymocyte progenitor cells differentiate into CD44$^+$CD25$^+$ stage 2 and then into CD44$^-$

CD25$^+$ stage 3 CD4$^-$CD8$^-$ double negative (DN) thymocytes (3, 4). Both DN2 and DN3 cells express RAG

and conduct V(D)J recombination of *Tcrb*, *Tcrd*, and/or *Tcrg* loci (3, 4). Although DN2-to-DN3

thymocyte maturation can occur independent of RAG and V(D)J recombination, an in-frame *Tcrb*

rearrangement is necessary for development beyond the DN3 stage (3, 4). The resultant TCRβ protein

associates with invariant pTα protein to produce pre-TCR complexes that signal antigen-independent

survival, proliferation, and differentiation of DN3 cells (β-selection) into CD44$^-$CD25$^-$ stage 4 DN

thymocytes and then CD4$^+$CD8$^+$ double-positive (DP) thymocytes (3, 4). DP cells express RAG and

conduct V(D)J recombination of *Tcra* loci, which typically proceeds through multiple successive rounds

of V-to-J rearrangement on both alleles (5). After an in-frame VJ rearrangement, the resulting TCRα

protein can pair with TCRβ protein and yield surface αβ TCRs, which must signal to prevent death by

neglect (6-9). This signal activation requires appropriate physical interactions between αβ TCRs and self-

peptide/MHC (pMHC) complexes displayed on thymic epithelial cells or dendritic cells, meaning that a

functional αβ TCR is inherently self-reactive (6-9). Interactions below a particular low threshold of

affinity (or avidity) cannot activate signalling to block apoptosis (6-9). At the other extreme, contacts

above a higher threshold trigger strong TCR signalling that causes apoptosis (negative selection) to delete

highly self-reactive cells with substantial autoimmune potential (6-9). However, interactions between

these two thresholds activate TCR signalling of a strength within a range that promotes survival and

differentiation of DP cells (positive selection) into CD4$^+$ or CD8$^+$ single-positive (SP) thymocytes, which

lack RAG expression and exit the thymus as mature αβ T cells (6-9). A notable aspect of this αβ TCR

quality control checkpoint is that thymocytes are more sensitive than mature αβ T cells as positively-

selecting pMHC activates TCR signaling only in the former (9-14). This differential is thought to enable

positive selection from low affinity/avidity interactions, prevent the same contacts from activating mature

αβ T cells, and provide a "safety net" for negative selection to more effectively suppress autoimmunity

(6-9). Although the T lineage-specific Themis protein contributes to this differential in TCR sensitivity

between immature and mature αβ T cells (9, 15, 16), the precise mechanisms by which this protein and

possibly additional factors "tune" TCR signaling and control αβ TCR selection remain to be elucidated.

The RAG1 protein N-terminus has a Zinc-binding RING finger E3 ubiquitin ligase domain of

undetermined relevance for antigen receptor gene assembly. RAG1 autoubiquitylates on lysine 233 (17),

which enhances RAG endonuclease activity as evidenced by impaired cleavage upon mutation of this

residue (18). RAG1 also ubiquitylates other proteins including histone H3 within chromatin (19, 20).

RAG1 RING domain mutations that disrupt ubiquitin ligase activity impair RAG-mediated cleavage and

joining of chromatin substrates, which could be due to loss of histone H3 ubiquitylation and/or alteration

in RAG1 protein structure (20-22). The C328Y mutation of a critical structural zinc-binding cysteine of

human RAG1 impairs V(D)J recombination and causes severe T and B cell immunodeficiency associated

with uncontrolled proliferation of activated oligoclonal αβ T cells (23). The analogous C325Y mutation

in mouse Rag1 destabilizes Rag1 tertiary structure, abrogates ubiquitin ligase activity, reduces V(D)J

recombination *in vitro,* and causes a near complete block of αβ T cell development at the DN3 thymocyte

stage due to impaired *Tcrb* recombination (21, 22). RING domains have a conserved proline (proline 326

in mouse Rag1) that is critical for ubiquitin ligase activity by permitting functional interaction with

ubiquitin-conjugating enzymes (24, 25). Relative to C325Y mutation, P326G mutation abrogates RAG1

ubiquitin ligase activity equivalently, but has substantially less severe effects on destabilizing RAG1

protein and reducing RAG endonuclease activity (22). Accordingly, P326 mutation of endogenous mouse

RAG1 protein allows an approach to elucidate potential physiologic roles of the RAG1 ubiquitin ligase

domain beyond promoting V(D)J recombination by stabilizing RAG1 protein structure.

To determine potential *in vivo* functions of RAG1 ubiquitin ligase activity, we created and analyzed mice carrying a homozygous Rag1 P326G mutation. We show that these mice exhibit reduced recombination of *Tcrb* and *Tcra* loci and lower efficiencies of β-selection, positive selection, and negative selection. We find that inactivation of the RAG1 ubiquitin ligase lowers expression of the pre-TCR signaling protein Syk in DN3 thymocytes and of the αβ TCR signaling protein Zap70 in DP thymocytes. Finally, we demonstrate that mature αβ T cells of these mice exhibit normal Zap70 expression and immune responses to activation, yet nevertheless confer greater potential for autoimmunity. Although reduced V(D)J recombination alone could underlie impaired β-selection and positive selection, this does not provide a plausible explanation for less efficient negative selection and greater autoimmunity hazard of αβ T cells. Thus, we propose that the RAG1 ubiquitin ligase domain might provide developmental stage-specific means to increase TCR signaling sensitivities in thymocytes and thereby enhance selection for beneficial *Tcrb* genes and against highly self-reactive αβ TCRs with autoimmune potential.

Materials and Methods

Mice

C57BL/6 background *PG* mice were made through gene-targeting as performed by InGenious Targeting

Laboratory (2200 Smithtown Avenue, Ronkonkoma, NY). The 9.5 kb genomic DNA used to construct

the targeting vector was sub-cloned from a 129Svev BAC clone. The CC>GG double point mutation was

shuttled from pJMJ029[P326G](22) into the targeting vector using unique restriction sites, then confirmed

by sequencing. Linearized targeting vector was electroporated into BA1 (C57BL/6 x 129/SvEv)

embryonic stem (ES) cells. After selection with G418 antibiotic, clones were expanded for PCR analysis

to identify recombinant ES clones. Positive clones were screened for short arm integration and then

confirmation of the point mutation was performed by PCR and sequencing. Positive clones were analyzed

by Southern blot to confirm long arm integration. Injected blastocysts were grown in foster mothers and

chimeras were backcrossed against C57BL/6 mice to generate heterozygotes. These were crossed with

CMV-*cre* mice (JAX #006054)(26) to remove the *neo* reporter cassette and generate the *Rag1^{P326G}* allele.

After confirmation of *neo* cassette removal, the strain was backcrossed against C57BL/6, and removal of

CMV-*cre* was confirmed by PCR. Sequencing of the entire *Rag1* locus in progeny confirmed the intended

CC>GG mutation and an additional T>C point mutation that generates a V238A substitution outside of

the RING finger domain. The strain was deposited at the Mutant Mouse Regional Resource Center at the

University of North Caroline (RAG1-tm1, MMRRC #: 37105). Before initiating studies described herein,

the *Rag1^{P326G}* allele was backcrossed for nine generations onto the C57BL/6 background. To generate

NOD background *PG* mice, CRISPR/Cas9-mediated gene editing was conducted in zygotes from *in vitro*

fertilization of NOD eggs with NOD sperm. A guide RNA (5'-AAGCACGGATATCGGCAAGA-3') that

hybridizes near Rag1 sequences that encode proline 326 was generated as described (27). The following

mix was electoporated into zygotes: 4μM of purified guide RNA, 4μM of Cas9-NLS nuclease (qb3

MacroLab Berkeley), 10μM of ssDNA 5'-

C*A*G*AAAGGACTTCACTGGGCTCTCCAGGTCAGTAGGGAAGCACGGATATCGGCAGGAT

CCACAATAGCTGCCCATGACTTTGAGACATCTGAGAATGCAGATCCTACAGAATAGATGCT

T-3' (IDT) that contains nucleotides encoding glycine in place of proline at amino acid 326 and a silent

point mutation of the next codon which introduce a cleavage site for BamHI restriction enzyme. Mice

were screened for introduction of the mutation by PCR with primers 5'-

CGTCGCAAGAGAACTCAGGCT-3' and 5'-TCTCGTCAGTGACAGGAGATGCT-3', followed by

BamHI digestion of the PCR product. Sanger sequence of PCR products confirmed incorporation of the

desired mutation and no other mutations in the *Rag1* coding sequence. Founders with mutations were

mated with NOD mice to establish NOD *Rag1$^{P326G/+}$* mice, which were bred with each other to generate

NOD *PG* mice and *WT* littermate controls. NOD mice, C57BL/6 background wild-type and *Rag1$^{-/-}$* mice

and BALB/c background *Rag1$^{-/-}$* mice were purchased from Jackson Laboratories. All experimental mice

were littermates or age-matched with control wild-type animals. All experiments were conducted in

accordance with national guidelines and approved by the Institutional Animal Care and Use Committee

(IACUC) of the Children's Hospital of Philadelphia.

Flow Cytometry

Single cell suspensions of all organs were ACK lysed to remove erythrocytes before being stained with

LIVE/DEAD™ Fixable Aqua Dead Cell Stain Kit (Invitrogen) and antibodies against respective surface

antigens (BD Biosciences, eBioscience, and Biolegend). Cells were stained for respective intracellular

proteins using Cytofix/Cytoperm Fixation/Permeabilization solution kit (BD Biosciences). For staining of

intracellular cytokines, cells (1 x10^6) were cultured in the absence or presence of 50 ng/ml PMA (Sigma)

and 1μg/ml Ionomycin (Cell Signaling Technology), with 2 μg/ml brefeldin A (Sigma) and 2 μM

monensin (eBioscience) for 4 hours at 37^{o}C. After staining for LIVE/DEAD and surface antigens, cells

were stained for respective cytokines using the Cytofix/Cytoperm kit (BD Bioscience) or transcription

factors using the FoxP3/Transcription Factor staining kit (eBiosciences). Data were acquired on a

MACSQuant flow cytometer (Miltenyi Biotec) or LSRII Fortessa (BD Biosciences) and analyzed using

Flowjo software version 10.5.3 (Tree Star).

Cell Sorting

To isolate DN3 cells, thymocytes were stained with PE-labelled CD4, CD8, CD11b, CD11c, NK1.1, Gr1,

and Ter119. Non-labelled cells were enriched by MACS depletion using anti-PE microbeads and LS

columns (Miltenyi). Enriched cells were then stained with CD4, CD8, CD44, CD25 and anti-lineage/stage

markers (TCRβ, B220, CD19, CD11b, CD11c, TCRδ, NK1.1, Ter119) antibodies. DN3 thymocytes were

sorted on Lin$^-$CD4$^-$CD8$^-$CD44$^-$CD25$^+$ phenotype. To isolate DP cells, thymocytes were stained with CD4,

CD8, and anti-lineage (B220, CD19, CD11b, CD11c, TCRδ, NK1.1, Ter119) antibodies. DP thymocytes

were sorted on Lin$^-$CD4$^+$CD8$^+$ phenotype. The cells of interest were isolated using a FACSAria Fusion

sorter (BD Biosciences).

Analyzing *Tcrb* and *Tcra* Rearrangements

Genomic DNA was extracted from DN3 or DP cells using the DNeasy Blood and Tissue kit (Qiagen). To

measure *Tcrb* rearrangements in DN3 cells, a Taqman PCR assay was used to quantify Vβ-to-DJβ1.1 and

Vβ-to-DJβ2.1 rearrangement levels with a primer specific for each Vβ paired and a probe, FAM or HEX,

specific for Jβ1.1 or Jβ2.1, respectively. Taqman PCR was performed with conditions according to the

manufacturer's instructions (IDT DNA) on the ViiA 7 system (Applied Biosystems). PCR of CD19 was

used for normalization. Primers, probes, and reaction conditions are as described (28). To assay *Tcrb*

repertoire, DNA from DN3 thymocytes were sent to Adaptive Biotechnologies, who used multiplex PCR

to amplify and deep sequence Vβ-Dβ-Jβ rearrangements. Gene segment usage was analyzed by

ImmunoSEQ Analyzer software (Adaptive Biotechnologies). To assess *Tcra* rearrangements in DP cells,

representative Vα-Jα rearrangements were quantified using a QuantiFast SYBR Green PCR kit (Qiagen)

on the ViiA 7 system (Applied Biosystems) as described (29, 30). PCR of β2M was used for normalization.

RT-qPCR for mRNA Expression

Respective cell populations were lysed in RLT buffer (Qiagen) containing 2-mercaptoethanol or TRIzol

(Life Technologies) immediately after cell sorting. Total RNA was isolated using RNeasy Mini kit

(Qiagen), treated with DNase (RNase-Free DNase Set, Qiagen), and reverse transcribed to generate cDNA

with High-Capacity RNA-to-cDNA™ Kit (Themo-Fisher Scientific) according to manufacturer's

directions. The cDNA was subjected to qPCR using Power SYBR Green kit (Applied Biosystems) and

actin, HPRT, ZAP70, and SYK primers (Qiagen). Relative expression was calculated using the ddCt

method, using actin or HPRT as a housekeeping gene, and relevant calibrator sample (explained in figure

legends for respective samples).

In vitro **T Cell Differentiation**

Splenocytes from C57BL/6 background *Rag1$^{-/-}$* mice were irradiated with 2,500 Rads (X-rad irradiator)

and 3 x 10^5 cells were seeded into each well of a 96 well plate. Naïve T cells (CD90.2$^+$, CD4$^+$, CD62L$^+$,

CD44$^+$, CD25$^-$) were sorted from C57BL/6 background *WT* or *Rag1$^{-/-}$* mice using a FACSaria Fusion cell

sorter (BD Biosciences), and 3 x 10^4 sorted cells were added to each well containing irradiated feeder

cells. All cells were cultured with 10 μg/ml anti-CD3 (Biolegend) and 3 μg/ml anti-CD28 (Biolegend) in

RPMI with 10% FBS, 1% PSG, and 1x NEAA (Invitrogen), 1x Sodium Pyruvate (Invitrogen), and 0.001%

2-mercaptoethanol. rhIL-2 was used at 30 IU/mL, rmIL-12 and rmIL-4 at 10 ng/mL, rmIL-6 at 20 ng/mL,

mTGFβ1 at 1 ng/mL, anti-mIL-4, anti-mIFNγ and anti-mIL-12 at 10 μg/mL (from Biolegend except rhIL-

2, rmIL-12, and rmIL-6 from Prepotech). The following cytokines/blocking antibodies were used to skew

towards respective Th-subsets: Th0 condition: rhIL-2; Th1 condition: rhIL-2, rmIL-12 and anti-mIL-4;

Th2 condition: rhIL-2, rmIL-4, anti-mIFNγ and anti-mIL-12; Th17 condition: rhIL-2, mIL-6, mTGFβ1,

anti-mIL-4 and anti-mIL-12; Treg condition: rhIL-2, mTGFβ1, anti-mIL-4 and anti-mIL-12.

LCMV Infections

Mice were infected intraperitoneally with 2×10^5 plaque-forming units of LCMV-Armstrong and

euthanized at indicated timepoints. Virus and gp33-tetramer were kindly provided by E. John Wherry

(University of Pennsylvania).

T Cell Transfer Model of Colitis

This model was performed as described (31). Briefly, naïve CD4$^+$ T cells were sorted from respective

donor mice (CD90.2$^+$, CD4$^+$, CD62L$^+$, CD44$^+$, CD25$^-$) on a FACSAria Fusion cell sorter. 0.5×10^6 cells

were transferred in 100 μl of PBS into each of the donor mice via intraperitoneal injection. Non-transferred

mice were injected with PBS alone. Mice were weighed weekly for the first 4 weeks, and then twice

weekly for 4 weeks. Mice were euthanized 8 weeks post-transfer unless severe morbidity enforced

premature euthanasia (in accordance with CHOP IACUC approved guidelines) or spontaneous death

occurred.

Diabetes Onset in NOD Background Mice

NOD background *WT* and *PG* mice were aged and assessed for spontaneous development of diabetes.

Mice were aged and monitored for disease to a maximum of 30 weeks of age. Mice older than 10 weeks

of age were weekly tested for glycosuria (Uristix®, Siemens). For positive mice, glycosuria was assayed

by a second test 24 hours later. Capillary blood glucose of mice testing positive twice for glycosuria was

tested via facial vein bleeding. Mice with blood glucose ≥250 mg/dl (13.9 mmol/l) were considered as

diabetic and euthanized.

Statistical analysis

280 All data was analyzed in Graphpad Prism 8 using statistical tests indicated in the figure legends. Error

281 bars indicate mean +/- SEM unless otherwise stated. n.s.=$p>0.05$, *$p<0.05$, **$p<0.01$, ***$p<0.001$,

282 ****$p<0.0001$.

                                                    Results

Homozygous Rag1 P326G mutation diminishes TCR gene assembly and subsequent thymocyte

developmental stage transitions.

To determine potential *in vivo* functions of the RAG1 ubiquitin ligase, we created and analyzed C57BL/6

strain mice homozygous for the Rag1 P326G mutation (*Rag1$^{P326G/P326G}$* mice; see Materials and Methods)

that abrogates ubiquitin ligase activity but minimally destabilizes RAG1 protein (22). We initially

analyzed αβ T cell development and TCR gene rearrangements in *Rag1$^{P326G/P326G}$* (*PG*) mice and wild-

type (*WT*) mice. As compared to *WT* mice, *PG* mice have a ~2-fold higher percentage and number of DN3

thymocytes and correspondingly lower percentage and number of DN4 thymocytes (Fig. 1, A and B). *PG*

mice also exhibit ~10% reductions in the percentage and number of DP thymocytes and the ratio of CD4$^+$

SP thymocytes to DP thymocytes (Fig. 1, C-E). Despite these minor impairments of the DN-to-DN4 and

DP-to-SP thymocyte developmental transitions, *PG* mice have normal numbers of total thymocytes and

splenic αβ T cells (Fig. 1 F). We observed ~50% lower levels of rearrangement for nearly all individual

Vβ segments in DN3 thymocytes of *PG* mice (Fig. 2, A and B). *Rag1* or *Rag2* mutations that substantially

impair RAG endonuclease activity, thymocyte development, and thymic cellularity also alter the usage of

individual Vβ segments (32, 33). Yet, reflecting the minor effects of Rag1 P326G mutation, unbiased

high-throughput sequencing of *Tcrb* rearrangements in DN3 cells shows similar usage of each Vβ segment

in *PG* and *WT* mice (Fig. 2 C). Finally, we detected ~50% lower levels of rearrangements for many of the

Vα/Jα combinations assayed in DP thymocytes of *PG* mice (Fig. 2 D). Together, these data show that

homozygous Rag1^{P326G} mutation produces RAG endonuclease complexes that catalyze decreased levels

of *Tcrb* and *Tcra* rearrangements and support less efficient thymocyte developmental transitions that

depend on the functional assembly of a *Tcrb* or *Tcra* gene. Notably, the degrees to which TCR gene

rearrangements and αβ T cell development are reduced in *PG* mice are dramatically less than reported for

homozygous Rag1 C235Y mutation (21). For example, Rag1^{P236G} mice generate 90% of the normal

number of DP thymocytes, whereas homozygous Rag1^{C235Y} mice produce only 1% of the normal number

of DP thymocytes. The markedly different severities of these phenotypes indicate that abrogation of RAG1

ubiquitin ligase activity without substantial disruption of RAG1 protein structure only slightly impairs

V(D)J recombination and αβ T cell development.

Inactivation of the RAG1 ubiquitin ligase decreases expression of the signaling proteins Syk and

Zap70 in thymocytes.

The greater than normal number of DN3 thymocytes in *PG* mice is not typical of a mutation that impairs

V(D)J recombination. Instead, this phenotype is consistent with diminished pre-TCR signaling where

DN3 cells experiencing β-selection exhibit increased survival but are less efficient than normal at

transitioning into DN4 thymocytes. It is notable that thymocytes lacking the Syk pre-TCR signaling

protein display this phenotype (34, 35), and that RAG DSBs induced in pre-B cells trigger a genetic

program that transcriptionally represses expression of Syk to dampen pre-BCR signaling (36). Thus, we

determined whether inactivation of the RAG1 ubiquitin ligase reduces Syk expression in DN3 thymocytes.

We detected a slightly lower than normal level of Syk protein in *PG* mice (Fig. 3, A and B); however,

unexpectedly, this correlated with an increase in the level of Syk mRNA (Fig. 3 C). These results suggest

that RAG DSBs downregulate Syk gene transcription in DN thymocytes, as in pre-B cells, and fewer RAG

DSBs from less *Tcrb* recombination in *PG* DN cells leads to higher than normal Syk transcript levels.

Despite this, the lower Syk protein expression in *PG* DN thymocytes implies that the RAG1 ubiquitin

ligase normally increases translation of Syk mRNA and/or stability of Syk protein during αβ T cell

development. Regardless of the underlying mechanisms, these data open the possibility that Syk-mediated

pre-TCR signaling is impaired, underlies the accumulation of DN3 thymocytes, and contributes to

impaired DN3-to-DN4 thymocyte progression in mice lacking RAG1 ubiquitin ligase activity.

Given that Syk is related to the Zap70 protein that transduces αβ TCR signals, we determined whether

inactivation of the RAG1 ubiquitin ligase also reduces Zap70 expression. To this end, we quantified levels

of Zap70 protein and mRNA in DP cells at each of four stages of αβ TCR-signaled positive selection as

defined by expression of CD69 and CCR7 proteins (37)(Fig. 3 D). In *PG* mice, we observed lower than

normal Zap70 protein levels, but normal levels of TCRβ protein, at each stage of selection (Fig. 3, E-H).

We detected normal levels of Zap70 mRNA in cells at each stage assayed in *PG* mice (Fig. 3 I), implying

that lower Zap70 mRNA translation and/or Zap70 protein stability is the basis for lower Zap70 expression.

RAG DSBs induced during lymphocyte development signal gene expression changes that persist in mature

NK cells lacking RAG1 and RAG2 expression (38). We therefore considered whether the Zap70

alterations caused by mutant RAG1 during development would also persist in mature T cells after RAG1

is no longer expressed. However, we detected normal Zap70 protein levels in splenic CD4$^+$ and CD8$^+$ αβ

T cells of *PG* mice (Fig. 3, J and K), indicating that Rag1^{P326G} mutation only reduces Zap70 expression in

immature αβ T cells. These data imply that Zap70-mediated   TCR signaling might be weaker than

normal in developing αβ T cells and thereby contribute to the impaired DP-to-SP thymocyte progression

of mice lacking RAG1 ubiquitin ligase activity.

Homozygous Rag1 P326G mutation decreases efficiencies of positive and negative selection.

Considering the critical role of Zap70 for positive and negative selection, we investigated whether either

of these αβ TCR quality control processes is impaired by the Rag1^{P326G} mutation. As a proxy to establish

whether positive selection is impaired in *PG* mice, we determined the frequency of αβ TCR expressing

(TCRβ$^+$) Stage 1 DP thymocytes that initiate positive selection as indicated by progression to Stage 2 in

PG and *WT* mice (Fig. 4 A). This frequency is lower in *PG* mice (Fig. 4 B). We note that this is an

imperfect approach to quantify positive selection because it cannot measure differences in progression

through subsequent stages nor cellular fate decisions from specific TCR/pMHC contacts. Nevertheless, our data are consistent with less efficient positive selection in mice lacking RAG1 ubiquitin ligase activity.

To determine whether the $Rag1^{P326G}$ mutation impairs negative selection, we utilized a naturally occurring system of superantigen-mediated negative selection (39). Laboratory mice express remnant mouse mammary tumor virus (MMTV) retroviral products. Some of these products can bind the MHCII I-E^d protein and certain Vβ peptides, including Vβ5.1-5.2 and Vβ12, to trigger exceptionally strong αβ TCR signaling that promotes negative selection (39, 40). BALB/c, but not C57BL/6, strain mice express I-E^d protein. As a result, thymocytes expressing Vβ5.1-5.2 or Vβ12 are efficiently deleted in BALB/c, but not B6, genetic backgrounds. We bred BALB/c background $Rag1^{-/-}$ mice with C57BL/6 background Rag1 PG or WT mice to generate and analyze $Rag1^{PG/-}$ and $Rag1^{WT/-}$ mice expressing I-E^d protein. To ascertain whether these mice exhibit a difference in negative selection, we measured frequencies of pre-selected DP thymocytes and post-selection SP thymocytes expressing I-E^d:MMTV-reactive (Vβ5$^+$ or Vβ12$^+$) or - unreactive (Vβ6$^+$, Vβ8$^+$, or Vβ14$^+$) αβ TCRs (example staining shown in Fig. 4 C). We quantified the fractions of thymocytes expressing each Vβ that progresses through development (are positively selected) and normalized these to the frequencies of total pre- and post-selected cells. For each experiment, we normalized values to the fraction for $Rag1^{WT/-}$ mice so that we could combine data from independent experiments (Fig. 4 D). Our analysis indicates that greater fractions of I-E^d:MMTV-reactive thymocytes proceed through DP-to-SP development in $Rag1^{PG/-}$ mice as compared to $Rag1^{WT/-}$ mice (Fig. 4 D), consistent with impaired negative selection of highly self-reactive thymocytes. In contrast, we found that the types of un-reactive thymocytes assayed develop equivalently in each genotype (Fig. 4 D). While the latter finding does not reflect impaired positive selection, we note that this experiment monitors a small subset of thymocytes expressing specific Vβ peptides and thus is not as sensitive as determining the

frequencies of DP cells that initiate positive selection. Regardless, we conclude that inactivation of the

RAG1 ubiquitin ligase renders negative selection less efficient.

Mature αβ T cells that develop without RAG1 ubiquitin ligase activity exhibit normal responses to

activation.

RAG endonuclease activity during lymphocyte ontogeny triggers permanent changes in gene expression

that correlate with impaired immune responses of mature lymphocytes (38). Although we did not observe

any differences in Zap70 protein levels between splenic αβ T cells of *PG* and *WT* mice, we sought to

elucidate whether mature αβ T cells that develop without RAG1 ubiquitin ligase activity have normal or

altered response to activation. We first conducted *in vitro* analyses. Upon stimulation with PMA and

ionomycin, splenic CD44^{hi} αβ T cells from *PG* mice produce normal levels of IFNγ, TNFα, IL-2, and IL-

17A (Fig. 5 A). These cells also show normal proliferation upon anti-CD3/anti-CD28 stimulation (Fig. 5

B). Moreover, stimulated CD4$^+$ and CD8$^+$ αβ T cells of *PG* mice normally express their canonical

transcription factors (Fig. 5, C and D) and cytokines (Fig. 5, E and F) when cultured in respective skewing

conditions. We next monitored αβ T cell responses *in vivo* following infection of mice with an acute strain

of lymphocytic choriomeningitis virus (LCMV-Armstrong). We analyzed mice either seven days later to

monitor at the immune response peak or 35 days later to monitor memory cells. At each timepoint, we

detected similar frequencies and numbers of antigen-specific CD8$^+$ αβ T cells that stain with gp33:H-2Db

tetramer in *PG* and *WT* mice (Fig. 5 G), indicating that *PG* mice display normal numbers and dynamic

responses of αβ T cells that recognize LCMV-Armstrong. We also observed equivalent dynamic

expression of Tbet, Eomes, KLRG1, and CD127 proteins on αβ T cells from *PG* and *WT* mice (Fig. 5 H),

indicating normal contraction of short-lived effector cells (SLECs), and sustained memory/memory

precursor effector cells (MPECs) in *PG* mice. Finally, we detected similar numbers of antigen specific αβ

T cells able to respond to the gp33, NP396, or gp61 viral peptides within a peptide re-stimulation assay

(Fig. 5, I and J). Together, these data support the notion that the development of αβ T cells in the absence

of RAG1 ubiquitin ligase activity does not substantially alter responses of these mature immune cells to

activation.

Mature αβ T cells that develop in the absence of RAG1 ubiquitin ligase activity exhibit increased

predisposition to autoimmunity.

In a setting of less efficient negative selection as with homozygous $Rag1^{P326G}$ mutation, an increased

predisposition to αβ T cell mediated autoimmunity would be expected. However, we have not observed

overt autoimmune symptoms within *PG* mice, possibly because mice of the C57BL/6 background are

resistant to spontaneous autoimmune diseases (41, 42). Thus, to investigate whether the $Rag1^{P326G}$

mutation can predispose to autoimmunity, we utilized an induced model of autoimmune colitis in

C57BL/6 mice and a spontaneous model of autoimmune diabetes in NOD mice.

The colitis model involves transferring into $Rag1^{-/-}$ mice naïve $CD4^+$ effector αβ T cells, which become

activated and trigger cytokine-driven colitis and associated systemic pathologies including weight loss

due to the absence of Tregs (31, 43, 44). We observed accelerated wasting of $Rag1^{-/-}$ mice receiving *PG*

cells relative to *WT* cells or no transfer (Fig. 6 A). We analyzed mice at eight weeks after transfer. As

compared to *WT* recipients, we detected greater numbers of total splenic $CD4^+$ T cells and of $CD4^+$ T cells

producing IFNγ, TNFα, and/or GM-CSF cytokines (Fig. 6, B and C). Colon length is used as a measure

of disease severity in this model, and mice receiving transferred *PG* cells had shorter colons relative to

mice receiving *WT* cells, consistent with worse disease caused by *PG* cells (Fig. 6 D). Histologic

assessment and scoring in this colitis model has been formalized (44), and can be grossly broken down

into scores related to inflammatory cell infiltration and scores related to injury of tissue architecture. There

was no statistical difference in inflammation scores between mice receiving *WT* and *PG* cells. However,

tissue damage scores were higher in mice receiving *PG* cells, suggesting that, on a per cell basis, *PG* cells

induce greater tissue injury in this model (Fig. 6, E-F). Notably, specifically in *PG*-transferred animals,

we observed autoimmune pathologies in addition to colitis, including dermatitis, chylous ascites, severe

small intestine enlargement, and neurological stress (Fig. 6 H). Dermatitis was most common and involved

massive inflammatory infiltration and tissue damage (Fig. 6 I). Consistent with more widespread

autoimmune conditions in *PG*-transferred mice, we significant splenomegaly in these mice (Fig. 6 J).

Finally, while not statistically significant, we observed an increased rate of premature death in the mice

transferred with *PG* cells compared to *WT* cells (Fig. 6K), which was especially surprising because death

is an unexpected outcome in this model. Collectively, the data from this colitis model indicate that CD4$^+$

effector T cells from *PG* mice have greater than normal intrinsic autoimmune potential.

To determine whether inactivation of the Rag1 ubiquitin ligase enhances autoimmunity in another genetic

background, we utilized CRISPR/Cas9-mediated gene editing to establish NOD mice with the Rag1^{P326G}

mutation. NOD mice develop spontaneous $\alpha\beta$ T cell driven diabetes at an incidence of 60–80% in females

and 20–30% in males (45, 46). We bred together NOD:*Rag1$^{P326G/WT}$* mice to create cohorts of NOD *PG*

and *WT* females, which we aged and screened for diabetes by monitoring glycosuria and blood glucose

levels. We observed accelerated onset of diabetes in *PG* mice compared to *WT* mice (Fig. 6 L).

Collectively, our analyses of the colitis and diabetes models on mice of two different genetic backgrounds

indicate that homozygous Rag1^{P326G} mutation increases predisposition to $\alpha\beta$ T cell mediated

autoimmunity.

Discussion

Our analysis of homozygous Rag1^{P326G} mice reveals roles for the RAG1 RING domain ubiquitin ligase in stimulating TCR gene assembly, shaping thymocyte gene expression, and enhancing TCR quality control checkpoints. As expected from *in vitro* analyses, the Rag1^{P326G} mutant produces RAG endonuclease complexes that catalyze modestly decreased V(D)J recombination of *Tcrb* and *Tcra* loci in thymocytes. While a less efficient RAG endonuclease could account entirely for the reduced assembly of *Tcrb* and *Tcra* genes, additional experiments are necessary to investigate potential functions of the RAG1 ubiquitin ligase in modulating chromatin accessibility and/or RAG binding of these loci. The reduced assembly of *Tcrb* genes most likely contributes to the impaired DN3-to-DN4 thymocyte development of *PG* mice, as is the case in mice homozygous for other *Rag1* or *Rag2* mutations that lower RAG endonuclease activity (21, 33). To our knowledge, reduced V(D)J recombinase function has not been demonstrated to increase DN3 thymocyte numbers; however, Syk inactivation has because it impairs pre-TCR signals that drive differentiation of DN3 cells (34, 35). Accordingly, it is plausible that reduced Syk protein expression in DN3 thymocytes contributes to their accumulation in *PG* mice, highlighting a potential role of the RAG1 ubiquitin ligase in promoting early thymocyte development beyond stimulating *Tcrb* gene assembly. Yet, unequivocal conclusions require elucidating how Rag1^{P326G} mutation impairs Syk expression and specifically disrupting these mechanisms.

We next considered effects of Rag1 RING domain ubiquitin ligase inactivation on the DP-to-SP thymocyte developmental transition. Given that deletion of most Jα segments in mice impairs DP-to-SP thymocyte development due to fewer chances for each DP cell to produce a positively-selected TCR (47), it is possible that reduced *Tcra* recombination contributes to less efficient positive selection in *PG* mice. Accordingly, we cannot definitively attribute a non-V(D)J recombination effect of Rag1 ubiquitin ligase inactivation to positive selection. In contrast, our data indeed support a non-V(D)J recombination effect

of reduced Rag1 ubiquitin ligase activity on negative selection. In DP thymocytes with normal V(D)J

recombinase function, the expression of a highly self-reactive αβ TCR signals apoptosis before ongoing

Tcra recombination can generate a new αβ TCR (48). Consequently, it is hard to conceive how reduced

RAG endonuclease activity would impair negative selection. However, as reduced Zap70 activity impairs

αβ TCR signaling and negative selection (49-51), attenuated αβ TCR signaling from lower Zap70 protein

expression is a plausible explanation for less efficient superantigen-mediated negative selection in *PG*

mice. However, definitive conclusions necessitate determining how Rag1[P326G] mutation impairs Zap70

expression and specifically disrupting these mechanisms.

Based on our findings, we propose that RAG1 ubiquitin ligase activity upregulates expression of Syk and

Zap70 protein respectively, to enhance β-selection of positive and negative selection of thymocytes. We

acknowledge that Syk and Zap70 protein levels are only slightly lower than normal in *PG* mice and that

we have not proven that these differences underlie impaired β-selection, positive selection, and negative

selection. We did not observe differences in Syk or Zap70 activation or other aspects of TCR signaling

between *WT* and *PG* thymocytes following anti-CD3 stimulation (data not shown). However, this methods

of activating TCR signaling is not as sensitive for detecting differences as using MHC to present different

peptides to a fixed transgenic αβ TCR (9, 10, 16), which is beyond the scope of this study. Moreover, we

note that small reductions in Zap70 activity render negative selection less efficient (49-51). In addition,

we consider it likely that Rag1[P326G] mutation also might alter expression of other proteins, with the

cumulative changes having synergistic effects on TCR signaling and thymocyte selection. To rigorously

test our model, it will be necessary to elucidate the extent and mechanisms by which the RAG1 ubiquitin

ligase modulates gene expression, and then disrupt these pathways, preferably without also reducing RAG

endonuclease activity. Likewise, it will be important to determine whether Rag1[P326G] mutation has any

effects on the thymic microenvironment (i.e. defective maturation of thymic epithelial cells and/or

distribution of dendritic cells) as is observed in mouse models with impaired V(D)J recombination and a resulting dramatically reduced thymic cellularity (52). However, unlike these models, *PG* mice have relatively normal numbers of thymocytes, which should facilitate distinction of thymocyte intrinsic versus extrinsic effects on αβ TCR selection.

There are multiple not mutually exclusive mechanisms by which the RAG1 ubiquitin ligase domain could increase Syk and Zap70 protein levels within thymocytes. In pre-B cells, RAG DSBs signal transcriptional activation of a broad multifunctional genetic program, with one component being decreased Syk protein expression to attenuate pre-BCR signaling and facilitate assembly of Ig light chain genes (36, 53). RAG DSBs in thymocytes could trigger transcriptional and post-transcriptional up-regulation of Syk and Zap70 expression, where fewer RAG DSBs from $Rag1^{P326G}$ mutation specifically impairs post-transcriptional mechanisms. DSB-induced covalent modifications of histones in chromatin around breakage sites are critical for amplifying DNA damage response (54). The inability of $Rag1^{P326G}$ to ubiquitylate histone H3 and other potential target proteins at RAG DSBs might dampen or prevent signals that mediate post-transcriptional upregulation of Syk and Zap70. Alternatively, the RAG1 ubiquitin ligase domain might elevate Syk and Zap70 protein expression independent of RAG endonuclease activity, such as by binding and/or ubiquitylating factors that regulate translation or turnover of these TCR signaling proteins. Whatever mechanisms an intact RAG1 ubiquitin ligase domain employs to stimulate Syk and Zap70 protein expression, their effects would be temporary and terminated after cessation of RAG1 expression following positive selection as evidenced by normal Zap70 protein levels in mature αβ T cells as well as their normal function to infectious challenges.

The requirement for αβ TCR selection on pMHC presents major challenges for establishing an αβ T cell population that recognizes and responds to diverse foreign peptides, but not self-peptides. An inability to positively-select αβ TCRs with very low affinity/avidity for pMHC could eliminate T cells capable of

binding foreign peptides with no similarities to self-peptides, generating holes in adaptive immunity. At

the same time, any positively-selected TCR is able to recognize a self-peptide and thus could cause

autoimmunity. To meet these challenges, cell intrinsic mechanisms have evolved to render thymocytes

more sensitive to TCR stimulation than their mature αβ T cell counterparts (10-14). This "TCR signal

tuning" elevates the affinity/avidity thresholds for both positive and negative selection because any

functional TCR/pMHC interaction triggers a stronger TCR signal in thymocytes versus αβ T cells (6-9).

This lower sensitivity of mature αβ T cells prevents their activation when encountering an pMHC identical

to that on which they were positively-selected. Although several factors have been implicated in regulating

TCR signal tuning, each is expressed in both thymocytes and mature αβ T cells (16, 55-59). Our

observations of reduced thymic Zap70 expression and less efficient positive and negative selection in *PG*

mice suggest that the RAG1 protein might serve as a "TCR signal tuning" factor. The thymocyte specific

developmental timing of RAG1 expression makes it an excellent candidate to serve as one such tuning

factor. In this scenario, the ability of the RAG1 ubiquitin ligase domain to upregulate Zap70, and perhaps

modulate expression of additional factors, would raise TCR/pMHC affinity/avidity thresholds for positive

selection and negative selection. Therefore, we propose that the RAG1 ubiquitin ligase domain provides

developmental stage-specific means to stimulate positive selection of potentially beneficial TCRs and

negative selection of highly self-reactive TCRs with greatest autoimmune potential. Critically, our study

provides a phenotypic framework from which to elucidate the exact molecular mechanisms by which the

RAG1 ubiquitin ligase promotes TCR gene assembly and transcends this process to govern the proper

selection of developing αβ T cells.

Acknowledgements

549   We thank Kyutae Lee for designing the gene-editing strategy and Adele Harman of the CHOP Transgenic

550   Core to generate the NOD *PG* mice. The authors have no completing financial interests to declare.

References

1. Bassing, C. H., W. Swat, and F. W. Alt. 2002. The mechanism and regulation of chromosomal V(D)J recombination. *Cell* 109 Suppl: S45-55.

2. Schatz, D. G., and P. C. Swanson. 2011. V(D)J recombination: mechanisms of initiation. *Annu Rev Genet* 45: 167-202.

3. Bhandoola, A., H. von Boehmer, H. T. Petrie, and J. C. Zuniga-Pflucker. 2007. Commitment and developmental potential of extrathymic and intrathymic T cell precursors: plenty to choose from. *Immunity* 26: 678-689.

4. von Boehmer, H., and F. Melchers. 2010. Checkpoints in lymphocyte development and autoimmune disease. *Nat Immunol* 11: 14-20.

5. Carico, Z., and M. S. Krangel. 2015. Chromatin Dynamics and the Development of the TCRalpha and TCRdelta Repertoires. *Adv Immunol* 128: 307-361.

6. Hogquist, K. A., and S. C. Jameson. 2014. The self-obsession of T cells: how TCR signaling thresholds affect fate 'decisions' and effector function. *Nat Immunol* 15: 815-823.

7. Gascoigne, N. R., V. Rybakin, O. Acuto, and J. Brzostek. 2016. TCR Signal Strength and T Cell Development. *Annu Rev Cell Dev Biol* 32: 327-348.

8. Klein, L., B. Kyewski, P. M. Allen, and K. A. Hogquist. 2014. Positive and negative selection of the T cell repertoire: what thymocytes see (and don't see). *Nat Rev Immunol* 14: 377-391.

9. Fu, G., V. Rybakin, J. Brzostek, W. Paster, O. Acuto, and N. R. Gascoigne. 2014. Fine-tuning T cell receptor signaling to control T cell development. *Trends Immunol* 35: 311-318.

10. Davey, G. M., S. L. Schober, B. T. Endrizzi, A. K. Dutcher, S. C. Jameson, and K. A. Hogquist. 1998. Preselection thymocytes are more sensitive to T cell receptor stimulation than mature T cells. *J Exp Med* 188: 1867-1874.

11. Lucas, B., I. Stefanova, K. Yasutomo, N. Dautigny, and R. N. Germain. 1999. Divergent changes in the sensitivity of maturing T cells to structurally related ligands underlies formation of a useful T cell repertoire. *Immunity* 10: 367-376.

12. Yachi, P. P., C. Lotz, J. Ampudia, and N. R. Gascoigne. 2007. T cell activation enhancement by endogenous pMHC acts for both weak and strong agonists but varies with differentiation state. *J Exp Med* 204: 2747-2757.

13. Pircher, H., U. H. Rohrer, D. Moskophidis, R. M. Zinkernagel, and H. Hengartner. 1991. Lower receptor avidity required for thymic clonal deletion than for effector T-cell function. *Nature* 351: 482-485.

14. Sant'Angelo, D. B., and C. A. Janeway, Jr. 2002. Negative selection of thymocytes expressing the D10 TCR. *Proc Natl Acad Sci U S A* 99: 6931-6936.

15. Choi, S., R. Cornall, R. Lesourne, and P. E. Love. 2017. THEMIS: Two Models, Different Thresholds. *Trends Immunol* 38: 622-632.

16. Fu, G., J. Casas, S. Rigaud, V. Rybakin, F. Lambolez, J. Brzostek, J. A. Hoerter, W. Paster, O. Acuto, H. Cheroutre, K. Sauer, and N. R. Gascoigne. 2013. Themis sets the signal threshold for positive and negative selection in T-cell development. *Nature* 504: 441-445.

17. Jones, J. M., and M. Gellert. 2003. Autoubiquitylation of the V(D)J recombinase protein RAG1. *Proc Natl Acad Sci U S A* 100: 15446-15451.

18. Singh, S. K., and M. Gellert. 2015. Role of RAG1 autoubiquitination in V(D)J recombination. *Proc Natl Acad Sci U S A* 112: 8579-8583.

19. Jones, J. M., A. Bhattacharyya, C. Simkus, B. Vallieres, T. D. Veenstra, and M. Zhou. 2011. The RAG1 V(D)J recombinase/ubiquitin ligase promotes ubiquitylation of acetylated, phosphorylated histone 3.3. *Immunol Lett* 136: 156-162.

20.   Grazini, U., F. Zanardi, E. Citterio, S. Casola, C. R. Goding, and F. McBlane. 2010. The RING
domain of RAG1 ubiquitylates histone H3: a novel activity in chromatin-mediated regulation of
V(D)J joining. *Mol Cell* 37: 282-293.
21.   Deng, Z., H. Liu, and X. Liu. 2015. RAG1-mediated ubiquitylation of histone H3 is required for
chromosomal V(D)J recombination. *Cell Res* 25: 181-192.
22.   Simkus, C., P. Anand, A. Bhattacharyya, and J. M. Jones. 2007. Biochemical and folding defects
in a RAG1 variant associated with Omenn syndrome. *J Immunol* 179: 8332-8340.
23.   Villa, A., C. Sobacchi, L. D. Notarangelo, F. Bozzi, M. Abinun, T. G. Abrahamsen, P. D.
Arkwright, M. Baniyash, E. G. Brooks, M. E. Conley, P. Cortes, M. Duse, A. Fasth, A. M.
Filipovich, A. J. Infante, A. Jones, E. Mazzolari, S. M. Muller, S. Pasic, G. Rechavi, M. G. Sacco,
S. Santagata, M. L. Schroeder, R. Seger, D. Strina, A. Ugazio, J. Valiaho, M. Vihinen, L. B. Vogler,
H. Ochs, P. Vezzoni, W. Friedrich, and K. Schwarz. 2001. V(D)J recombination defects in
lymphocytes due to RAG mutations: severe immunodeficiency with a spectrum of clinical
presentations. *Blood* 97: 81-88.
24.   Zheng, N., P. Wang, P. D. Jeffrey, and N. P. Pavletich. 2000. Structure of a c-Cbl-UbcH7 complex:
RING domain function in ubiquitin-protein ligases. *Cell* 102: 533-539.
25.   Albert, T. K., H. Hanzawa, Y. I. Legtenberg, M. J. de Ruwe, F. A. van den Heuvel, M. A. Collart,
R. Boelens, and H. T. Timmers. 2002. Identification of a ubiquitin-protein ligase subunit within
the CCR4-NOT transcription repressor complex. *EMBO J* 21: 355-364.
26.   Schwenk, F., U. Baron, and K. Rajewsky. 1995. A cre-transgenic mouse strain for the ubiquitous
deletion of loxP-flanked gene segments including deletion in germ cells. *Nucleic Acids Res* 23:
5080-5081.
27.   Lee, K. D., and C. H. Bassing. 2020. Two Successive Inversional Vbeta Rearrangements on a
Single Tcrb Allele Can Contribute to the TCRbeta Repertoire. *J Immunol* 204: 78-86.
28.   Gopalakrishnan, S., K. Majumder, A. Predeus, Y. Huang, O. I. Koues, J. Verma-Gaur, S.
Loguercio, A. I. Su, A. J. Feeney, M. N. Artyomov, and E. M. Oltz. 2013. Unifying model for
molecular determinants of the preselection Vbeta repertoire. *Proc Natl Acad Sci U S A* 110: E3206-
3215.
29.   Shih, H. Y., J. Verma-Gaur, A. Torkamani, A. J. Feeney, N. Galjart, and M. S. Krangel. 2012.
Tcra gene recombination is supported by a Tcra enhancer- and CTCF-dependent chromatin hub.
Proc Natl Acad Sci U S A 109: E3493-3502.
30.   Chen, L., Z. Carico, H. Y. Shih, and M. S. Krangel. 2015. A discrete chromatin loop in the mouse
Tcra-Tcrd locus shapes the TCRdelta and TCRalpha repertoires. *Nat Immunol* 16: 1085-1093.
31.   Ostanin, D. V., J. Bao, I. Koboziev, L. Gray, S. A. Robinson-Jackson, M. Kosloski-Davidson, V.
H. Price, and M. B. Grisham. 2009. T cell transfer model of chronic colitis: concepts,
considerations, and tricks of the trade. *Am J Physiol Gastrointest Liver Physiol* 296: G135-146.
32.   Liang, H. E., L. Y. Hsu, D. Cado, L. G. Cowell, G. Kelsoe, and M. S. Schlissel. 2002. The
"dispensable" portion of RAG2 is necessary for efficient V-to-DJ rearrangement during B and T
cell development. *Immunity* 17: 639-651.
33.   Horowitz, J. E., and C. H. Bassing. 2014. Noncore RAG1 regions promote Vbeta rearrangements
and alphabeta T cell development by overcoming inherent inefficiency of Vbeta recombination
signal sequences. *J Immunol* 192: 1609-1619.
34.   Colucci, F., D. Guy-Grand, A. Wilson, M. Turner, E. Schweighoffer, V. L. Tybulewicz, and J. P.
Di Santo. 2000. A new look at Syk in alpha beta and gamma delta T cell development using
chimeric mice with a low competitive hematopoietic environment. *J Immunol* 164: 5140-5145.
35.   Palacios, E. H., and A. Weiss. 2007. Distinct roles for Syk and ZAP-70 during early thymocyte
development. *J Exp Med* 204: 1703-1715.
36.   Bednarski, J. J., R. Pandey, E. Schulte, L. S. White, B. R. Chen, G. J. Sandoval, M. Kohyama, M.
Haldar, A. Nickless, A. Trott, G. Cheng, K. M. Murphy, C. H. Bassing, J. E. Payton, and B. P.

Sleckman. 2016. RAG-mediated DNA double-strand breaks activate a cell type-specific
checkpoint to inhibit pre-B cell receptor signals. *J Exp Med* 213: 209-223.

37.   Kimura, M. Y., J. Thomas, X. Tai, T. I. Guinter, M. Shinzawa, R. Etzensperger, Z. Li, P. Love, T.
Nakayama, and A. Singer. 2016. Timing and duration of MHC I positive selection signals are
adjusted in the thymus to prevent lineage errors. *Nat Immunol* 17: 1415-1423.

38.   Karo, J. M., D. G. Schatz, and J. C. Sun. 2014. The RAG recombinase dictates functional
heterogeneity and cellular fitness in natural killer cells. *Cell* 159: 94-107.

39.   Jordan, M. S., J. E. Smith, J. C. Burns, J. E. Austin, K. E. Nichols, A. C. Aschenbrenner, and G.
A. Koretzky. 2008. Complementation in trans of altered thymocyte development in mice
expressing mutant forms of the adaptor molecule SLP76. *Immunity* 28: 359-369.

40.   Surh, C. D., and J. Sprent. 1994. T-cell apoptosis detected in situ during positive and negative
selection in the thymus. *Nature* 372: 100-103.

41.   Jiang, W., M. S. Anderson, R. Bronson, D. Mathis, and C. Benoist. 2005. Modifier loci condition
autoimmunity provoked by Aire deficiency. *J Exp Med* 202: 805-815.

42.   Jin, Y., A. Lee, J. H. Oh, H. W. Lee, and S. J. Ha. 2019. The R229Q mutation of Rag2 does not
characterize severe immunodeficiency in mice. *Sci Rep* 9: 4415.

43.   Mottet, C., H. H. Uhlig, and F. Powrie. 2003. Cutting edge: cure of colitis by CD4+CD25+
regulatory T cells. *J Immunol* 170: 3939-3943.

44.   Eri, R., M. A. McGuckin, and R. Wadley. 2012. T cell transfer model of colitis: a great tool to
assess the contribution of T cells in chronic intestinal inflammation. *Methods Mol Biol* 844: 261-
275.

45.   Anderson, M. S., and J. A. Bluestone. 2005. The NOD mouse: a model of immune dysregulation.
Annu Rev Immunol 23: 447-485.

46.   Delovitch, T. L., and B. Singh. 1997. The nonobese diabetic mouse as a model of autoimmune
diabetes: immune dysregulation gets the NOD. *Immunity* 7: 727-738.

47.   Huang, C. Y., B. P. Sleckman, and O. Kanagawa. 2005. Revision of T cell receptor {alpha} chain
genes is required for normal T lymphocyte development. *Proc Natl Acad Sci U S A* 102: 14356-
14361.

48.   Kreslavsky, T., H. J. Kim, S. B. Koralov, D. Ghitza, T. Buch, H. Cantor, K. Rajewsky, and H. von
Boehmer. 2013. Negative selection, not receptor editing, is a physiological response of
autoreactive thymocytes. *J Exp Med* 210: 1911-1918.

49.   Sakaguchi, N., T. Takahashi, H. Hata, T. Nomura, T. Tagami, S. Yamazaki, T. Sakihama, T.
Matsutani, I. Negishi, S. Nakatsuru, and S. Sakaguchi. 2003. Altered thymic T-cell selection due
to a mutation of the ZAP-70 gene causes autoimmune arthritis in mice. *Nature* 426: 454-460.

50.   Mallaun, M., G. Zenke, and E. Palmer. 2010. A discrete affinity-driven elevation of ZAP-70 kinase
activity initiates negative selection. *J Recept Signal Transduct Res* 30: 430-443.

51.   Hsu, L. Y., Y. X. Tan, Z. Xiao, M. Malissen, and A. Weiss. 2009. A hypomorphic allele of ZAP-
70 reveals a distinct thymic threshold for autoimmune disease versus autoimmune reactivity. *J Exp*
Med 206: 2527-2541.

52.   Marrella, V., P. L. Poliani, L. D. Notarangelo, F. Grassi, and A. Villa. 2014. Rag defects and
thymic stroma: lessons from animal models. *Front Immunol* 5: 259.

53.   Bredemeyer, A. L., B. A. Helmink, C. L. Innes, B. Calderon, L. M. McGinnis, G. K. Mahowald,
E. J. Gapud, L. M. Walker, J. B. Collins, B. K. Weaver, L. Mandik-Nayak, R. D. Schreiber, P. M.
Allen, M. J. May, R. S. Paules, C. H. Bassing, and B. P. Sleckman. 2008. DNA double-strand
breaks activate a multi-functional genetic program in developing lymphocytes. *Nature* 456: 819-
823.

54.   Van, H. T., and M. A. Santos. 2018. Histone modifications and the DNA double-strand break
response. *Cell Cycle* 17: 2399-2410.

700 55. Fu, G., S. Vallee, V. Rybakin, M. V. McGuire, J. Ampudia, C. Brockmeyer, M. Salek, P. R. Fallen,
701    J. A. Hoerter, A. Munshi, Y. H. Huang, J. Hu, H. S. Fox, K. Sauer, O. Acuto, and N. R. Gascoigne.
702    2009. Themis controls thymocyte selection through regulation of T cell antigen receptor-mediated
703    signaling. *Nat Immunol* 10: 848-856.
704 56. Lesourne, R., S. Uehara, J. Lee, K. D. Song, L. Li, J. Pinkhasov, Y. Zhang, N. P. Weng, K. F.
705    Wildt, L. Wang, R. Bosselut, and P. E. Love. 2009. Themis, a T cell-specific protein important for
706    late thymocyte development. *Nat Immunol* 10: 840-847.
707 57. Johnson, A. L., L. Aravind, N. Shulzhenko, A. Morgun, S. Y. Choi, T. L. Crockford, T. Lambe,
708    H. Domaschenz, E. M. Kucharska, L. Zheng, C. G. Vinuesa, M. J. Lenardo, C. C. Goodnow, R. J.
709    Cornall, and R. H. Schwartz. 2009. Themis is a member of a new metazoan gene family and is
710    required for the completion of thymocyte positive selection. *Nat Immunol* 10: 831-839.
711 58. Wang, D., M. Zheng, L. Lei, J. Ji, Y. Yao, Y. Qiu, L. Ma, J. Lou, C. Ouyang, X. Zhang, Y. He, J.
712    Chi, L. Wang, Y. Kuang, J. Wang, X. Cao, and L. Lu. 2012. Tespa1 is involved in late thymocyte
713    development through the regulation of TCR-mediated signaling. *Nat Immunol* 13: 560-568.
714 59. Li, Q. J., J. Chau, P. J. Ebert, G. Sylvester, H. Min, G. Liu, R. Braich, M. Manoharan, J. Soutschek,
715    P. Skare, L. O. Klein, M. M. Davis, and C. Z. Chen. 2007. miR-181a is an intrinsic modulator of
716    T cell sensitivity and selection. *Cell* 129: 147-161.

Footnotes

727 This work was supported by the Fondation ARC pour la recherche sur le cancer (C.M.), La Ligue contre

728 le cancer (C.M.), Fondation Pasteur Mutualite (C.M.), Nancy Taylor Foundation for Chronic Disease

729 (E.M.B.) and NIH grants R03 AI097611 (J.M.J.), R01 HL112836 (E.M.B.), R01 AI121250-A1 (E.M.B.),

730 R21 A1 135435 (C.H.B.), and RO1 AI 112621 (C.H.B.).

731
732
733

734
735
736

Figure 1. *PG* mice exhibit impaired thymocyte developmental transitions that require TCR gene

assembly. **(A)** Representative flow cytometry plots of DN stages of thymocyte development showing the

frequency of cells at each stage. **(B)** Quantification of numbers of cells in each DN stage. **(C)**

Representative flow cytometry plots of DN, DP, CD4$^+$ SP, and CD8$^+$ SP thymocytes. **(D)** Quantification

of numbers of cells in each stage. **(E)** Quantification of the ratio of CD4$^+$ SP or CD8$^+$ SP thymocytes to

DP thymocytes. **(F-G)** Quantification of the numbers of total thymocytes (F) and splenic αβ T cells (G).

(A-G) All data were collected from 6-10 weeks old mice and (B, D, E, F, and G) are combined from four

independent experiments including a total of 14-16 mice per genotype. Bars indicate mean +/- SEM. Stats:

2-way ANOVA with Sidak multiple comparison test. n.s. $p > 0.05$; * $p < 0.05$, **$p < 0.01$, *** $p < 0.001$, ****

$p < 0.0001$.

Figure 2. Thymocytes of *PG* mice support diminished levels of *Tcrb* and *Tcra* recombination. **(A-B)**

Taqman qPCR quantification of rearrangements of each Vβ segment to DJβ1.1 (A) or DJβ2.1 (B)

complexes conducted on genomic DNA from DN3 thymocytes. Signals from each assay were normalized

to values from an assay for the invariant CD19 gene. Shown are average values +/- SEM from three

independent DN cell preparations. **(C)** Adaptive Immunoseq quantification of the percentage usage of

each Vβ in *Tcrb* rearrangements of DN3 thymocytes. Shown are average values from two independent

DN cell preparations. **(D)** qPCR quantification of specific Vα/Jα rearrangements peformed on genomic

DNA from DP thymocytes. Signals from each assay were normalized to values from an assay for the

invariant β2m gene. Shown are the average values +/- SEM from three independent DP cell preparations.

For all graphs in figure: #, not detected, n.s. $p > 0.05$; * $p < 0.05$, **$p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$

Figure 3. Thymocytes of *PG* mice have reduced Syk and Zap70 protein levels. **(A-B)** Representative flow

cytometry histogram plots (A) and quantification (B) of Syk protein in DN3 thymocytes. (B) Shown are

all data points and average values +/- SEM from one of two independent experiments (n = 3-4 mice per

genotype, unpaired t-test). **(C)** qRT-PCR quantification of Syk mRNA in DN3 cells. Relative mRNA

levels were calculated using ddCt method with signals from each assay normalized to values from an assay

for the *Hprt* gene. Shown are all data points and average values +/- SEM from two independent

experiments (n = 8-9 mice per genotype, 2-way ANOVA with experiment an independent variable, stats

indicate difference as a result of genotype). **(D)** Gating strategy for Stages 1-4 of DP thymocyte selection.

Pre-gated on live singlets and $CD8^+$ and/or $CD4^+$. **(E-H)** Representative flow cytometry histogram plots

(E, G) and quantification (F, H) of Zap70 (E, F) and TCRβ (G, H) protein in Stages 1-4 of DP thymocyte

selection. (F, H) Shown are all data points and average values +/- SEM from one of three independent

experiments (n = 4-5 mice per genotype, 2-way ANOVA with Sidak's multiple comparison post-test) **(I)**

qRT-PCR quantification of Zap70 mRNA in DN3 cells or Stage 2 or 4 DP cells. Relative mRNA levels

were calculated using ddCt method with signals from each assay normalized to values from an assay for

the Hprt gene. Shown are all data points and average values +/- SEM from one experiment (n = 4-5 mice

per genotype, 2-way ANOVA with sidak multiple comparison post-test. **(J-K)** Representative flow

cytometry histogram plots (J) and quantification (K) of Zap70 protein in naïve $CD4^+$ or $CD8^+$ T cells.

Shown are all data points and average values +/- SEM from two independent experiments (n = 3 mice per

genotype, 2-way ANOVA with Sidak multiple comparison post-test). For all graphs in figure: n.s. p>0.05;

* p<0.05, **p<0.01, *** p<0.001, **** p<0.0001.

Figure 4. *PG* mice exhibit impaired positive and negative selection. **(A)** Gating strategy for measuring

positive selection of thymocytes showing Stage 1 and 2 gates. Pre-gated on live singlets and $CD4^+$ and/or

$CD8^+$. **(B)** Quantification of the ratio of the number of Stage 2 cells to the total number of Stage 1 and 2

cells. Shown are all data points and global mean from three independent experiments (n = 10-11 mice per

genotype, 2-way ANOVA indicating effect of genotype). **(C)** Gating strategy form measuring negative

selection of Vβ6$^+$ and Vβ8$^+$ thymocytes in F1 progeny of C57BL/6 background *PG* mice bred with

BALB/c background *Rag1$^{-/-}$* mice. Pre-gated on live singlets. **(D)** Quantification of ratios of indicated Vβ$^+$

cells that progress to post-selection versus those Vβ$^+$ cells that are produced at the pre-selection stage.

Normalized to total thymocytes in pre or post-selection population and *WT* controls. Shown are all data

points and average +/- SEM from two independent experiments (n = 9 mice per genotype, 2-way ANOVA

with Sidak multiple comparison post-test. For all graphs in figure: n.s. $p>0.05$; * $p<0.05$, **$p<0.01$, ***

$p<0.001$, **** $p<0.0001$.

Figure 5. Mature αβ T cells of PG mice exhibit normal immune response. **(A)** Representative plot of

cytokine expression on CD4$^+$CD44^{hi} αβ T cells following *ex vivo* stimulation of splenocytes with PMA

and ionomycin. **(B-F)** Representative analyses of naïve splenic CD4$^+$ or CD8$^+$ T cells after *ex vivo*

simulation with anti-CD3/anti-CD28 antibodies and additional antibodies to skew towards indicated

lineages. Shown is CFSE incorporation (B) or expression of relevant transcription factors (C, D) or

cytokines (E, F). **(G, H)** Representative flow data showing (G) frequencies of splenic αβ T cells that stain

with the gp33:H-2Db tetramer or (H) express indicated cytokines after LCMV infection of mice. Data are

representative of two independent experiments with at least four mice of each genotype. **(I, J)**

Representative plots (J) and (I) quantification of antigen-specific αβ T cells able to respond to indicated

viral peptides within a peptide re-stimulation assay. (A-F) Data are representative of two independent

experiments. (G-J) Data are combined from four independent experiments (two D7 and two D35

timepoints) with 8-11 mice of each genotype per time-point.

806
807

Figure 6. Mature αβ T cells of *PG* mice possess higher than normal autoimmune hazard**. (A)** Graph of body weights of *Rag1*[-/-] hosts over time after transfer of *WT* or *PG* CD4[+] T cells or no cells (NT). Data combined from three independent experiments with a total of 13-17 mice of each genotype. Lines indicate fitted segmented-linear regression lines +/- 95% confidence interval. Day 15 was fixed as the inflection point. Difference between gradients of second slope was determined by linear regression analysis. **(B)** Numbers of CD4[+] T cells in spleen eight weeks post-transfer. **(C)** Total number of IFNγ, TNFα, and/or GM-CSF producing cells per spleen. Shown are all data points from three independent experiments with 12-14 mice per group. Bar indicates global mean. Pre-gating on live singlets, CD90.2[+], CD4[+], and CD44[+]. **(D)** Colon length of mice sacrificed at eight weeks post-infection. **(E)** Representative hemotoxylin and eosin stained colon sections from respective mice at eitght weeks post-transfer. 10x magnification. **(F-G)** Total colitis score as assessed by blinded pathologist. Scored as per previously described guidelines for the five phenotypes (44): (F) Colon inflammation score (contribution of scores for crypt abcesses and inflammatory cell infiltrate). (G) Colon tissue damage score (contribition of scores for crypt architecture, tissue damage, goblet cell loss). **(H)** Non-colitis pathologies described for mice transferred with *WT* versus *PG* cells. **(I)** Hemotoxylin and eosin stained skin section from *PG*-transferred mouse with severe dermatitis at eight weeks post-transfer. 10x magnification. **(J)** Spleen weight at eight weeks post-transfer normalized to total mouse bodyweight prior to transfer. **(K)** Survival curves of mice transferred with *WT* or *PG* cells. Data combined from three independent experiments. Statistical analysis = Log-rank Mantel-Cox test. Overall: 0/13 NT mice died, 1/15 *WT*-transferred mice died, and 4/17 *PG*-transferred mice died. (B, C, F, G, and J) Shown are all data points from three independent experiments, 12-14 mice per group. Experiment indicated by shape of data point. Bar indicates global mean. Analyzed by 2-way ANOVA with Tukey HSD post-test. n.s. $p>0.05$; * $p<0.05$, **$p<0.01$, *** $p<0.001$, **** $p<0.0001$ **(L)** Curves showing diabetes free survival of female NOD *WT* and *PG* mice. Mice were generated by mating together NOD *Rag1*[WT/PG] mice and monitored for their first 30 weeks of life. The percent diabetes survival curves

832 include mice that have passed 20 weeks of age or gotten diabetes before 20 weeks of age. Analyzed by

833 Gehan-Breslow-Wilcoxon test. p=0.049.

857 **Figure 1**

Figure 2

868   **Figure 3**

880 **Figure 5**

883 **Figure 6**

885