Proton receptors regulate synapse-specific reconsolidation in the amygdala

Erin E Koffman[1,2][#], Charles M Kruse[1,2][#], Kritika Singh [2], FarzanehSadat Naghavi [2],

Jennifer Egbo [2], Sandra Boateng [2], Mark Houdi BA [2], Boren Lin [2], Jacek Debiec [3],

Jianyang Du [1, 4][*]

[1] Department of Anatomy and Neurobiology, University of Tennessee Health Science

Center, Memphis, TN 38163, USA.

[2] Department of Biological Sciences, University of Toledo, Toledo, OH 43606, USA.

[3] Molecular & Behavioral Neuroscience Institute and Department of Psychiatry,

University of Michigan, Ann Arbor, MI, USA.

[4] Neuroscience Institute, University of Tennessee Health Science Center, Memphis, TN,
United States.
12

[#] Equal contribution

* Correspondence:

Jianyang Du

Department of Anatomy and Neurobiology,

University of Tennessee Health Science Center,

Memphis, TN 38163, USA

TEL: 901-448-3463

E-MAIL: jdu15@uthsc.edu

SUMMARY

When an extinction procedure is performed within the reconsolidation window, the

original aversive memory can be replaced by one that is less traumatic. Recent studies

revealed that carbon dioxide (CO_2) inhalation during retrieval enhances memory lability.

However, the effects of CO_2 inhalation on the central nervous system can be extensive,

and there is lack of evidence suggesting that the effects of CO_2 are selective to a

reactivated memory. We discovered that CO_2 inhalation paired with memory retrieval

potentiates the specific aversive memory trace, resulting in greater memory lability. The

specific effects of CO_2 depend on acid-sensing ion channels (ASICs), the proton

receptors that are involved in synaptic transmission and plasticity in the amygdala. In

addition, CO_2 inhalation alters memory lability via synaptic plasticity at selectively

targeted synapses. Overall, our results suggest that inhaling CO_2 during the retrieval

event increases the lability of an aversive memory through a synapse-specific

reconsolidation process.

KEYWORDS: carbon dioxide (CO_2); acid-sensing ion channels; proton receptors;

aversive conditioning; aversive memory; memory retrieval; AMPA receptors;

reconsolidation; synaptic plasticity; spine structure.

INTRODUCTION

Recently, aversive memory research in both rodents and humans has focused on a

window of time after aversive memory retrieval, in which the memory is labile and

subject to intervention (Kida, 2020; Lee, 2009; Monfils et al., 2009; Nader et al., 2000;

Sara, 2000; Schiller et al., 2010; Tronson and Taylor, 2007). This window of time is

known as a reconsolidation window, and is thought to last up to six hours after initiation

(Clem and Huganir, 2010; Duvarci and Nader, 2004; Monfils et al., 2009; Schiller et al.,

2010). Several studies have demonstrated that interrupting the updating process

aroused by retrieval prevents memory restorage, generating selective amnesia (Agren

et al., 2012; Clem and Huganir, 2010; Mactutus et al., 1979; Monfils et al., 2009; Nader

et al., 2000; Schiller et al., 2010). Studies using rodent models have indicated that

pharmacological intervention within the reconsolidation window successfully erases

reactivated specific aversive memory (Nader et al., 2000; Sara and Hars, 2006; Schiller

et al., 2010). Despite the efficacy, ethical and practical concerns may prevent similar

pharmacological interventions from being used in humans (Agren et al., 2012; Schiller

et al., 2010). Recently, drug-free paradigms have been proposed that effectively prevent

the return of aversive memories in both rodents and humans (Bjorkstrand et al., 2016;

Clem and Huganir, 2010; Huang et al., 2020; Liu et al., 2014; Monfils et al., 2009;

Schiller et al., 2010). Monfils *et al.* developed a novel protocol in which an isolated

retrieval trial was followed by an extinction event within a specific time frame. This

resulted in the weakening of the original memory trace, thereby preventing reversion of

the original aversive memory by spontaneous recovery, renewal, or reinstatement

(Clem and Huganir, 2010; Jarome et al., 2015; Monfils et al., 2009; Quirk et al., 2010;

Wu et al., 2017). These studies suggest the mechanism by which retrieval changes the

lability of memory may become a target for novel clinical treatments for anxiety

disorders.

Currently, the outcomes of the combination of retrieval and extinction paradigms are

variable, depends on multiple factors, in both rodents (Auber et al., 2013; Chan et al.,

2010; Costanzi et al., 2011; Goode et al., 2017; Ishii et al., 2012) and humans (Golkar

et al., 2012; Klucken et al., 2016). Because it is difficult to completely erase the original

memory, a more reliable strategy for triggering memory erasure is necessary. In search

of this strategy, we studied the effects that CO_2 inhalation may have on the erasure of a

specific memory. As we showed previously, when mice receive a retrieval tone

supplemented by CO_2 inhalation, the memory becomes more labile than the

presentation of a retrieval tone alone (Du et al., 2017). Within the reconsolidation

window, the labile memory becomes more convertible, either weakened by extinction or

strengthened by reconditioning. Moreover, the effects of CO_2 inhalation on memory

lability were dependent on acid-sensing ion channels (ASICs), since disrupting ASICs in

the amygdala eliminated these effects of CO_2 (Du et al., 2017).

Recent studies have revealed that protons are potential neurotransmitters (Du et al.,

2014; Highstein et al., 2014; Kawasaki et al., 2009) and ASICs serve as postsynaptic

proton receptors that play key roles in neurotransmission and synaptic plasticity in the

amygdala, a brain region that is critical for the formation of aversive memories (Cohen

et al., 2017; Du et al., 2014; Farley et al., 2018; Ziemann et al., 2009). ASICs are

members of the Degenerin/ Epithelial sodium channel (DEG/ ENaC) family (Ben-

Shahar, 2011). To date, six family members have been identified (ASIC1a, ASIC1b,

ASIC2a, ASIC2b, ASIC3, and ASIC4) (Wemmie et al., 2013). These proteins assemble

as homo- or heterotrimers to form channels that are proton-gated, voltage-insensitive,

permeable to both Na^+ and Ca^{2+} and activated by extracellular protons (Waldmann et al.,

1997; Waldmann and Lazdunski, 1998). ASIC1a is widely expressed in many areas of

the brain, where it is associated with numerous brain functions and disorders, including

hippocampally-dependent learning and memory, anxiety, depression, stroke,

neurodegeneration, seizure, inflammation, and nerve injury (Chu and Xiong, 2012; Gao

et al., 2005; Ortega-Ramírez et al., 2017; Wang et al., 2018; Wemmie et al., 2002;

Wemmie et al., 2006). ASIC1a is highly expressed in the amygdala and its ion channel

activity is evoked by a reduction in extracellular pH within the physiological range

(Taugher et al., 2017; Wemmie et al., 2003). Disruption of ASIC1a affects synaptic

transmission and plasticity (Mango and Nisticò, 2019; Soto et al., 2018; Wemmie et al.,

2003; Wu et al., 2004). Loss of ASIC1a function also leads to impaired high-frequency

electrical stimulation-induced long-term potentiation (LTP) (Chiang et al., 2015; Du et

al., 2014; Liu et al., 2016; Wemmie et al., 2002). Moreover, we demonstrated that

presynaptic stimulation induces a transient synaptic drop in pH and activates ASIC-like

excitable postsynaptic currents (EPSCs) in pyramidal neurons (Du et al., 2014; Kreple

et al., 2014), suggesting that protons sufficiently activate postsynaptic ASIC1a in

synaptic transmission. In mice, disruption of ASIC1a activity reduces innate fear

learning and memory and alters neuronal activity in the fear circuit (Coryell et al., 2007),

whereas its overexpression has opposite effects (Wemmie et al., 2004). Also, reducing

pH by CO_2 inhalation or the injection of acid into the amygdala induced aversive

behavior response and enhanced aversive memory, and the CO_2 effects are ASIC1a

dependent (Ziemann et al., 2009).

Although our previous data suggested that CO_2 inhalation affects the aversive memory

retrieval and alters the memory lability, we also know that the CO_2 - affected memory

trace is associated with the retrieval cue (Du et al., 2017), inhalation of CO_2 may reduce

pH outside of the amygdala (Dulla et al., 2005; Zandbergen et al., 1989). Thus, whether

CO_2 and ASICs specifically regulate memory trace in retrieval is still unknown and this

question is outstanding. In this study, we found that CO_2 inhalation paired with memory

retrieval selectively potentiated memory lability in mice. Furthermore, electrorheological

and imaging studies in brain slices support the conclusion that the effects of CO_2 on

memory retrieval are specifically associated with a given memory. Our study proposes

that inhaling CO_2 within the reconsolidation window regulates aversive memory with

specificity, providing a unique angle to further study the mechanism by which memory is

modulated.

EXPERIMENTAL PROCEDURES

Mice

For our experiment, we used both male and female mice between 10-14 months of age.

Mice were derived from a congenic C57BL/6 background including wild-type, ASIC1a

knock out (ASIC1a$^{-/-}$), and TetTag-c-fos-tTA mice. TetTag-cFos-tTA mice were

obtained from Jackson Laboratory and crossed with C57BL/6J mice. Mice carrying

the Fos-tTA transgene were selected; Fos-tTA mice have a Fos promoter driving

expression of nuclear-localized, two-hour half-life EGFP (shEGFP) (Du et al., 2017;

Koffman and Du, 2017; Ramirez et al., 2013). The Fos promoter also drives the

expression of tetracycline transactivator (tTA), which bind to the tetracycline-responsive

element (TRE) site on an injected recombinant adeno-associated virus, $AAV_{2/9}$-TRE-

mCherry virus, resulting in the expression of mCherry (Du et al., 2017; Koffman and Du,

2017). The binding of the tTA to the TRE site is inhibited by doxycycline (DOX).

Inhibition of tTA binding prevents target gene expression (Das et al., 2016; Liu et al.,

2012; Ramirez et al., 2013).

Both male and female mice age10-14 weeks were randomly selected for the experiment

groups. Experimental mice were maintained on a standard 12-hour light-dark cycle and

received standard chow and water ad libitum. Animal care and procedures met the

National Institutes of Health standards. The University of Tennessee Health Science

Center Laboratory Animal Care Unit (Protocol #19-0112) and University of Toledo

Institutional Animal Care and Use Committee (Protocol #108791) approved all

procedures.

Aversive conditioning, retrieval, extinction, and memory test

The protocols for each experiment are detailed in the schematics of each figure. All

mice were handled by experimenters for 30 minutes on each of 3 days before aversive

conditioning. On day 1, mice were habituated to the aversive conditioning chamber

(Med Associates Inc.) for 7 minutes. Mice were then exposed to varying conditioning

protocols, as described below.

164 *Experiment 1: Standard conditioned stimulus (CS) auditory aversive conditioning,*

165 *retrieval, extinction, and memory tests.*

166 On day 1 in a curated environment (context A), the experimental mice were presented

167 with six pure tones (80 dB, 2 kHz, 20 seconds each) paired with 6 foot shocks - one

168 shock at the end of each tone (0.7 mA, 2 seconds). The interval between each tone was

169 100 seconds. On day 2, the mice were placed into a new environment (context B) and

170 habituated for 4 minutes. Mice then inhaled either unaltered air or air containing 10%

171 CO_2 for 7 minutes. Five minutes after inhalation of CO_2 or air began, mice were

172 presented with one 20 second pure tone to retrieve the memory. The mice were then

173 returned to their home cages. 30 minutes later, the mice returned to the retrieval

174 chamber (context B) and underwent two rounds of extinctions. In the first round of

175 extinction, mice were exposed to 20 pure tones with an interval between tones of 100

176 seconds. Mice were then returned to their home cage. 30 minutes later, the mice went

177 through the extinction protocol again with 20 pure tones. On day 7, the mice were tested

178 to see if their aversive response would recur via spontaneous recovery in context B with

179 4 pure tones. Thirty minutes after spontaneous recovery, the mice were returned to the

180 original context of the aversive memory, context A, in a recovery protocol with 4 pure

181 tones.

182

183 Freezing behavior in mice (the absence of movement beyond respiration) is used as a

184 measure of aversive response. To evaluate the outcomes of freezing behavior in mice,

185 the percentage of time during CS presentation spent in freezing was scored

186 automatically using VideoFreeze software (Med Associates Inc.). In the spontaneous

recovery and renewal tests, outcomes of the percentage of time freezing were averaged

from each of the 4 CSs.

Experiment 2: Two distinct CSs aversive conditioning, retrieval, extinction, and

memory tests.

This procedure was used to test the specificity of the effects of CO_2 on memory

retrieval. The context settings and parameters are similar to the above standard one CS

auditory aversive conditioning. In contrast to experiment 1, the mice were presented

with three pure tones (80 dB, 2 kHz, 20 seconds each) that alternated with three white

noises (60 dB, 2 kHz, 20 seconds each); all six stimuli were paired with foot shocks. On

day 2, the mice inhaled either unaltered air or air containing 10% CO_2 for 7 minutes.

Five minutes after inhalation of CO_2 or air began, the mice underwent retrieval with one

single pure tone followed by one white noise with or without CO_2. This was followed

thirty minutes later by two sections of extinctions with either pure tones or white noises.

On day 7, the mice were tested via spontaneous recovery and renewal protocols with 4

pure tones and 4 white noises respectively.

Experiment 3: Two distinct CSs aversive conditioning, retrieval, anisomycin, and

memory tests.

In a series of experiments, the standard extinction procedure was replaced with

amygdala infusion of anisomycin (detailed in the surgery procedure below). In brief, the

cannula was implanted on the amygdala 4-7 days before the behavioral experiments.

Starting on day 1, the mice were subjected to the aversive conditioning described in

experiment 2. On day 2, 30 minutes after retrieval, the mice were infused with 62.5

µg/µl anisomycin via the cannula in the lateral nuclei of the amygdala (LA) bilaterally

and returned to their home cage (Debiec et al., 2010). On day 7, the mice were tested

via spontaneous recovery and renewal as described in experiment 2.

Surgery and chemical infusion

For the cannula placement procedure, mice were anesthetized with isoflurane through

an anesthetic vaporizer, secured to the stereotaxic instrument and a cannula made from

a 25-gauge needle was inserted bilaterally into LA (relative to bregma: -1.2 mm

anterioposterior; ±3.5 mm mediolateral; -4.3 mm dorsoventral) (Du et al., 2017; Koffman

and Du, 2017). Dental cement secured the cannula and bone anchor screw in place.

Mice recovered for 4-5 days before any subsequent testing was carried out. A 10 µL

Hamilton syringe connected to a 30-gauge injector was inserted 1 mm past the cannula

tip to inject anisomycin (diluted in 1 µl artificial cerebrospinal fluid (ACSF), pH 7.3) over

5 sec. The injection sites were mapped post-mortem by sectioning the brain (10 µm

coronal) and performing cresyl violet staining.

Brain slice preparation and patch-clamp recording of amygdala neurons

Ten minutes after the memory retrieval experiment ended, mice were euthanized with

overdosed isoflurane and whole brains were dissected into pre-oxygenated (5% CO_2

and 95% O_2) ice-cold high sucrose dissection solution containing (in mM): 205 sucrose,

5 KCl, 1.25 NaH_2PO_4, 5 $MgSO_4$, 26 $NaHCO_3$, 1 $CaCl_2$, and 25 glucose (Du et al., 2017).

A vibratome sliced brains coronally into 300 µm sections that were maintained in normal

ACSF containing (in mM): 115 NaCl, 2.5 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 1.25 NaH_2PO_4, 11

glucose, 25 $NaHCO_3$ bubbled with 95% O_2/5% CO_2, pH 7.35 at 20°C-22°C. Slices were

incubated in the ACSF at least 1 hour before recording. For experiments, individual

slices were transferred to a submersion-recording chamber and were continuously

perfused with the 5% CO_2/95% O_2 solution (~3.0 ml/min) at room temperature (20°C -

22°C).

As we described previously (Du et al., 2017), pyramidal neurons in the lateral amygdala

were studied using whole-cell patch-clamp recordings. The pipette solution containing

(in mM): 135 KSO_3CH_3, 5 NaCl, 10 HEPES, 4 MgATP, 0.3 Na_3GTP, 0.5 K-EGTA

(mOsm=290, adjusted to pH 7.25 with KOH). The pipette resistance (measured in the

bath solution) was 3-5 MΩ. High-resistance (>1 GΩ) seals were formed in voltage-

clamp mode. Picrotoxin (100 µM) was added to the ACSF throughout the recordings to

yield excitatory responses. In AMPAR current rectification experiments, we applied D-

APV (100 µM) to block NMDAR-conducted EPSCs. The peak amplitude of ESPCs was

measured to determine current rectification. The amplitude was measured ranging from

-80 mV to +60 mV in 20 mV steps. The peak amplitude of EPSCs at -80 mV and +60

mV was measured for the rectification index. In EPSC ratio experiments, neurons were

measured at -80 mV to record AMPAR-EPSCs and were measured at +60 mV to record

NMDAR-EPSCs. To determine the AMPAR-to-NMDAR ratio, we measured the peak

amplitude of ESPCs at -80 mV as AMPAR-currents, and peak amplitude of EPSCs at

+60 mV at 70 ms as NMDAR-currents after onset. Data were acquired at 10 kHz using

Multiclamp 700B and pClamp 10.1. The mEPSCs events (>5pA) were analyzed in

Clampfit 10.1. The decay time (τ) of mEPSCs was fitted to an exponential using

Clampfit 10.1.

Immunohistochemistry and cell counting

The AAV-TRE-mCherry plasmid was obtained from the laboratory of Dr. Susumu

Tonegawa (Liu et al., 2012; Ramirez et al., 2013), and was used to produce $AAV_{2/9}$ by

the University of Iowa Gene Transfer Vector Core. For one week leading up to virus

microinjection, TetTag Fos-tTA mice were fed with food containing 40 mg/kg DOX. We

used a 10 μl Hamilton microsyringe and a WPI microsyringe pump to inject virus (0.5 μl

of 1.45E+12 viral genomes/ml of $AAV_{2/9}$-TRE-mCherry) bilaterally into the amygdala

(relative to bregma: -1.2 mm anterioposterior; ±3.5 mm mediolateral; -4.3 mm

dorsoventral), as described previously (Du et al., 2017; Koffman and Du, 2017). For

a two-week window between surgery and behavior training, mice were housed and fed

with a DOX-containing diet. The DOX-containing diet was ceased twenty-four hours

before aversive conditioning began on day one (replaced by a regular diet), then

immediately restarted afterward. Thirty minutes after retrieval on day two, the mice were

euthanized according to protocol. We used transcardial perfusion with 4%

paraformaldehyde (PFA) to fix whole brains, followed by continued fixation in 4% PFA at

4°C for 24 hours (Wright et al., 2020). Following perfusion, we used a vibratome (Leica

VT-1000S) to dissect 50 μm amygdala coronal slices, which were collected in ice-cold

PBS. To complete immunostaining, slices were placed in Superblock solution (Thermo

Fisher Scientific) plus 0.2% Triton X-100 for 1 hour and incubated with primary

antibodies (1:1000 dilution) at 4°C for 24 hours (Du et al., 2017). Primary antibodies we

used include: rabbit polyclonal IgG anti-RFP (Rockland Cat# 600-401-379); chicken IgY

anti-GFP (Thermo Fisher Scientific Cat# A10262) and mouse anti-NeuN (Millipore Cat#

MAB377X) (Liu et al., 2012; Ramirez et al., 2013). We then washed and incubated

slices for one hour with secondary antibodies (Alexa Fluor 488 goat anti-chicken IgG

(H+L) (Molecular Probes Cat# A-11039); Alexa Fluor 568 goat anti-rabbit IgG (H+L)

(Molecular Probes Cat# A-21429); Alexa Fluor 647 goat anti-mouse IgG (H+L) (Thermo

Fisher Scientific Cat# A-21235), 1:200 dilution). VectaShield H-1500 (Vector

Laboratories Cat# H-1500) was used to mount slices, while confocal microscopy was

used to view the slices. We used ImageJ software to analyze dendritic spine

morphology. Thin, mushroom and stubby spines were categorized based on the

following parameters: 1) mushroom spines: head-to-neck diameter ratio >1.1:1 and

spine head diameter >0.35 µm; 2); thin spines: head-to-neck diameter ratio >1.1:1 and

spine head diameter >0.35 µm or spine head-to-neck diameter ratios <1.1:1 and spine

length-to-neck diameter > 2.5 µm; 3); stubby spines: spine head-to-neck diameter ratios

<1.1:1 and spine length-to-neck diameter ≤ 2.5 µm (Kreple et al., 2014; Wright et al.,

2020).

Statistical analysis

One-way ANOVA and Tukey's post-hoc multiple comparison tests were used for

statistical comparison of groups. An unpaired Student's t-test was used to compare

results between two groups. P<0.05 was considered statistically significant, and we did

not exclude potential outliers from our data except the ones did not receive successful

aversive conditioning. The graphing and statistical analysis software Graphpad Prism 8

was used to analyze statistical data, which was presented as means ± SEM. Sample

sizes (n) are indicated in the figure legends, and data are reported as biological

replicates (data from different mice, different brain slices). Each group contained tissues

pooled from 4-5 mice. Due to variable behavior within groups, we used sample sizes of

10-16 mice per experimental group as we previously described in earlier experiments

(Du et al., 2017). In behavioral studies, we typically studied groups with four randomly

assigned animals per group, as our recording equipment allowed us to record four

separate animal cages simultaneously. The experiments were repeated with another set

of four animals until we reached the target number of experimental mice per group.

Experimentation groups were repeated in this manner so that each animal had the

same controlled environment-the same time of day and with similar handling,

habituation, and processes.

RESULTS

Our recent studies suggested that CO_2 inhalation throughout memory retrieval

enhances the lability of the memory and boosts the efficiency of the memory erasure

(Du et al., 2017). To further test whether the effects of CO_2 on memory retrieval are

synapse-specific, we designed a series of unique experiments by which we were able to

generate two distinct auditory aversive memories and identify the specificity of CO_2

effects on each of them.

CO_2 selectively enhances the lability of auditory aversive memory in the

amygdala

Previous studies have described an aversive conditioning paradigm in which memory

can be selectively reactivated and reconsolidated, suggesting synapse-specific

reconsolidation of distinct aversive memories in the amygdala (Debiec et al., 2010;

Doyere et al., 2007). We followed this paradigm albeit with modifications **(Fig.1A, Fig.**

s1A). On day 1, we trained the animals with two distinct conditioned stimuli: three pure

tones and three white noises paired with one foot-shock per stimuli as aversive

unconditioned stimuli (US) (see the detailed description in Materials and Methods). We

evaluated the outputs of the aversive conditioning through the percentage of the

freezing time within the time of CSs. The freezing was significantly increased after each

of the three conditioned stimuli, indicating the mice were trained sufficiently under the

designed condition **(Fig.1B, Fig. s1B)**.

On day 2, the animals were placed into a new context (context B) and presented with

one pure tone followed by a single noise (or vice versa) to retrieve the memory **(Fig. 1A,**

C, Fig. s1A, C). The animals were then returned to their home cages. 30 minutes later,

all mice underwent two blocks of extinctions in context B, each extinction contains 20

pure tones **(Fig.1D)** or 20 white noise **(Fig. s1D).** At the end of the extinction procedure,

the freezing dropped down to a low level **(Fig. 1D, Fig. s1D)**, suggesting that the

extinction procedure was sufficient to suppress the memory. Five days later, the mice

underwent spontaneous recovery (context B) and renewal (context A) respectively. Four

tones and four noises were presented throughout the memory test **(Fig.1A, Fig. s1A)**.

The group that underwent extinction with a specific CS showed specificity in which

freezing response was lowered after extinction **(Fig.1E, Fig. s1E)**. For example, when

the pure tone was presented during extinction, freezing in the pure tone group was

lower than freezing in the noise group **(Fig.1E)**, and vice versa **(Fig. s1E)**. When

retrieval was paired with 10% CO_2 inhalation, memory erasure effects were enhanced

(Fig.1 F-I, Fig. s1F-I), and there are statistical significances between the groups with or

without CO_2 in both spontaneous recovery and renewal (green columns, **Fig.1 J-K** and

red columns, **Fig. s1J-K**). To further evaluate the specificity of the effects of

reconsolidation on memory modifications, we designed another retrieval protocol in

which we presented pure tone and white noise, either of them paired with 10% CO_2

inhalation, followed by an unrelated extinction of CS, white noise, or pure tone

respectively **(Fig. s2)**. Consistent with our expectation, CO_2 did not boost the effects on

the retrieved memory in the absence of paired extinction, compared to the data in Fig. 1

and Fig. S1 **(Fig. s2B-I)**. When compared to the data in **Fig. 1F-I** and **Fig. s1F-I**, we

found the application of 10% CO_2 to the retrieval event failed to enhance the outcome

after extinction, indicating a specificity of the CO_2 effects. In all, our data suggest that

memory encoded in the amygdala can be distinct, and the effects of CO_2 on memory

are specific.

To focus on testing the specific effects of CO_2 on retrieval, we replaced the extinction

procedure with an injection of a protein synthesis inhibitor, anisomycin, to obliterate the

aversive memory **(Fig. 2A)**. Anisomycin, when injected bilaterally into the amygdala

after retrieval, causes memory erasure compared to the saline injection group (Debiec

et al., 2010). We conditioned the mice and retrieve the memory with pure tones

(Fig. 2B-C), followed by anisomycin/saline injection **(Fig. 2D)**. Our experiments show

that anisomycin disrupts the memory during reconsolidation **(Fig. 2E)**, and that memory

retrieval is required for memory erasure with anisomycin injection **(Fig. s3A-E)**.

Consistent with the extinction results in Fig. 1, when retrieval was paired with 10%

CO_2 inhalation, we found anisomycin reduced more aversive response-further

confirming that CO_2 enhances memory lability specifically **(Fig. 2F-I)**. To exclude the

possibility that anisomycin associates with one CS other the other, as rigorous controls,

we conditioned the mice with both pure tone and white noise and carried out memory

retrieval with both CSs and found anisomycin has equal effects on memory in both tone

and noise groups **(Fig. s3F-M)**. When 10% CO_2 was applied while the CSs were

presented, the retrieval group paired with CO_2 showed less freezing, regardless of the

type of CSs (pure tone or white noise) **(Fig.s4 B-I)**. As rigorous controls, we applied

CO_2 for both retrieval events together and anisomycin decreased the freezing level in

both groups, suggesting that CO_2 had equal effects on both tone and noise **(Fig. s4 J-**

Q). As another control, saline injection following retrieval and CO_2 did not cause

memory erasure, indicating that anisomycin was necessary to disrupt the reactivated

memory **(Fig. s5)**. Talking together, our data demonstrate that the effects of CO_2 are

specific to a distinct memory that is activated by a specific CS.

The specific effects of CO_2 on memory lability is ASIC-dependent.

We have previously found the effects of CO_2 on memory retrieval to be ASIC dependent

(Du et al., 2017). However, it is still unknown if CO_2 application to a specific memory

trace affects an ASIC-dependent mechanism. To answer this question, we first

performed distinct aversive conditioning in ASIC1a[-/-] mice with three pure tones and

white noises on day 1 **(Fig. 3A)**, followed by a pure tone and white noise for retrieval

on day 2. 30 minutes post-retrieval, we performed extinctions with pure tones. Five days

later, we tested spontaneous recovery and renewals with 4 pure tones and white

noises. Similar to the response we saw in WT mice, the freezing level in the pure tone

group of ASIC1a$^{-/-}$ mice were less than that in the white noise group (spontaneous

recovery, 46% decrease; renewal, 47.5% decrease) **(Fig. 3B-E)**. When 10% CO_2

inhalation was paired with pure tone in retrieval, we found that CO_2 did not have

additional effects on the memory with the specific CS in ASIC1a$^{-/-}$ mice (spontaneous

recovery, 43.4% decrease; renewal, 45.6% decrease) **(Fig. 3F-I)**, and there are no

statistical significances between the groups with or without CO_2 in both spontaneous

recovery and renewal (green columns, **Fig. 3J, K**). We had hypothesized that the

effects of CO_2 on memory retrieval would be ASIC dependent and our data supported

this prediction. We then replaced the extinction procedure with anisomycin, to obliterate

the aversive memory **(Fig. 4A)**. The ASIC1a$^{-/-}$ mice were conditioned with pure tones

(Fig. 4B) and followed by a single tone as retrieval **(Fig. 4C)**. We then apply anisomycin

infusions **(Fig. 4D)** and test the memory 5 days later **(Fig. 4E)**. Anisomycin dramatically

reduced freezing in memory tests that followed, whereas pairing with CO_2 in the CS did

not cause an additional reduction in the ASIC1a$^{-/-}$ mice, suggesting an ASIC

dependency **(Fig. 4E)**. We also presented two distinct CSs (pure tone and white noise)

during conditioning and then retrieval, with or without 10% CO_2, followed by anisomycin

infusions in the ASIC1a$^{-/-}$ mice **(Fig. 4F)**. Anisomycin dramatically reduced freezing in

memory tests that followed, whereas pairing with CO_2 in either CSs (pure tone or white

noise) did not cause an additional reduction in the ASIC1a$^{-/-}$ mice, suggesting an ASIC

dependency **(Fig. 4G-N)**.

To provide evidence that an acute ASIC1a blockage was able to eliminate the effects of

CO_2, we injected 100nM PcTX-1 into the lateral amygdala bilaterally 1 hour before the

application of CO_2 to the retrieval **(Fig. 5A)**. Our data suggest that compared to the

saline injection group **(Fig. 5 B-E)**, inhibiting ASIC1a by PcTX-1 significantly reduced

the CO_2 effects on memory retrieval **(Fig. 5 F-I)**, statistical analysis in the spontaneous

recovery and renewal groups supported this conclusion (green columns, **Fig. 5J, K**).

This pattern of findings suggests that the effects of CO_2 on specific memory traces are

ASIC dependent.

Activation of ASICs through CO_2 inhalation alters reconsolidation of distinct

memory through alteration of AMPARs.

AMPARs are glutamatergic receptors that have crucial roles in modulating memory

retrieval and destabilization (Auber et al., 2013; Chan et al., 2010; Clem and Huganir,

2010; Monfils et al., 2009; Quirk et al., 2010). Previous studies suggest that the

exchange of Ca^{2+}-impermeable AMPARs (CI-AMPARs) for Ca^{2+}-permeable AMPARs

(CP-AMPARs) occurs after retrieval (Clem and Huganir, 2010; Hong et al., 2013). 10%

CO_2 inhalation paired with retrieval induces a stronger current rectification of AMPARs

(the signature of CP-AMPARs) than in the retrieval alone group, indicating a greater

exchange of AMPARs (Du et al., 2017). Interestingly, no further enhancement was

observed in the ASIC1a$^{-/-}$ brain slices, indicating that the effect of CO_2 inhalation on

AMPAR exchange is ASIC dependent (Du et al., 2017). To further study whether

CO_2 specifically alters the AMPARs exchange in retrieval, we designed a unique

experiment to separate the aversive conditioning and retrieval and measure the

rectification of AMPARs **(Fig. 6A)**. To study this, we conditioned the mice with 6 pure

tones on day 1 **(Fig. 6B, left)**. On day 2, the mice were divided into 4 groups based on

retrieval conditions - the first group received pure tone only; the second group-pure tone

plus 10% CO_2 inhalation; the third group - white noise only; and the fourth group

received white noise+10% CO_2 inhalation **(Fig. 6B, right)**. Ten minutes after retrieval,

we dissected brain slices and AMPAR current was recorded in the pyramidal neurons in

the lateral amygdala through stimulation of thalamic inputs **(Fig. 6A)**. Rectification, a

signature of CP-AMPARs, was compared among all groups. Consistent with earlier

reports, pure tone retrieval increased current rectification (Clem and Huganir, 2010;

Hong et al., 2013) and CO_2 paired with pure tone retrieval caused stronger rectification

(Fig. 6C). However, when white noise was presented as the retrieval event, both white

noise and white noise plus CO_2 failed to cause a significant rectification compared to

the pure tone group **(Fig. 6C)**. This data supports our prediction that CO_2 was

associated with a specific memory trace that was reactivated. To control for the possible

effects of the order of presentation of CSs, we switched over the pure tone and white

noise in the aversive conditioning and retrieval. Similar results were observed,

confirming the effects of CO_2 were not artificial **(Fig. 6D, E)**.

We then asked whether synaptic strength changed with the application of an unrelated

retrieval CS and CO_2. The ratio of AMPAR-EPSCs to NMDAR-EPSCs might represent

the strength of the synapse (Rao and Finkbeiner, 2007). Previous studies reported that

the AMPAR/ NMDAR-EPSC ratio increased after aversive conditioning whereas

retrieval did not potentiate further increase, suggesting that memory retrieval did not

alter the synaptic strength (Clem and Huganir, 2010; Hong et al., 2013). Our previous

studies also indicated that CO_2 inhalation during memory retrieval did not strengthen the

synapse in the amygdala (Du et al., 2017). We further tested whether the pairing of CO_2

inhalation with the specific retrieval CS influenced the strength of a synapse. Currents

were recorded at -80mV for AMPAR-EPSCs and +60mV for NMDAR-EPSCs. Our data

suggest that retrieval plus CO_2 inhalation did not change the AMPAR/ NMDAR-EPSCs

ratio **(Fig. 6F)**. Moreover, the characteristics of miniature EPSCs (mEPSCs), including

amplitude, frequency, and decay times, were not altered **(Fig. 6G)**. In addition, the

pairing of CO_2 with another unrelated CS in retrieval did not change the strength of the

synapse **(Fig. 6F, G)**. This data suggests that the effects of CO_2 inhalation on specific

memory retrieval enhance the destabilization of the synapse without changing synaptic

strength.

The effects of CO_2 inhalation on distinct memory trace.

Our previous studies indicate that CO_2 enhances memory trace that is associated with

aversive conditioning (Du et al., 2017). In this experiment, we examined the mechanism

behind the specificity of CO_2 effects on memory traces. Using the TetTag-c-fos driven-

GFP mouse model, neurons in the amygdala involved in memory trace after aversive

conditioning can be labeled with a long-lasting mCherry fluorescent protein and the

neuron in the retrieval trace can be labeled with a 2-hour short half-life nuclear-localized

EGFP (shEGFP) **(Fig. 7A, B)** (see the details in Materials and Methods) (Du et al.,

2017; Koffman and Du, 2017). The overlapped labeling (yellow) represents the neurons

in the same memory trace of aversive and retrieval (Du et al., 2017; Liu et al., 2012;

Ramirez et al., 2013). In this experiment, the mice were first conditioned with pure tone,

activating the associated neurons that were labeled with mCherry **(Fig. 7C)**. Right

after aversive conditioning, the mice were immediately fed with a DOX diet thereby

preventing further mCherry labeling. On day 2, the mice were divided into two groups-

one group of mice were presented with a single pure tone to retrieve the memory,

another was presented with white noise. A temporary, 2-hour half-life, shEGFP was

labeled after the retrieval. Thirty minutes after the retrieval event, we sliced the

amygdala and imaged shEGFP- and mCherry-positive cells **(Fig. 7D)**. Compared to

pure tone aversive conditioning/pure tone retrieval group (same memory trace of

aversive conditioning and retrieval), inhalation of CO_2 in the pure tone aversive

conditioning/white noise retrieval group did not result in an increase of neurons positive

for expression of both mCherry-positive and shEGFP-positive neurons (overlapped

labeling, yellow) **(Fig. 7E)**. Control experiments to identify the efficiency of the aversive

conditioning on the expression of mCherry on the cells were performed **(Fig. 7F)**. These

findings indicate that CO_2, when paired with retrieval, only enhances the memory trace

that has been reactivated; CO_2 paired with unrelated retrieval cue does not affect the

original memory trace. These findings suggest a specific effect of CO_2 on the memory

trace.

We then examined the effects of CO_2 on dendritic spine morphology after memory

retrieval. Spine morphology has been widely indicated in the mechanism of synaptic

plasticity (Woolfrey and Srivastava, 2016; Wright et al., 2020; Yang et al., 2009).

Dendritic spines are the primary target of neurotransmission input in the central nervous

system (Bourne and Harris, 2008), and their density and structure provide the basis for

physiological changes in synaptic efficacy that underlie learning and memory (Bailey et

al., 2015). Spine formation and plasticity are regulated by many conditions, including

exterior stimulation and behavior (Gipson and Olive, 2017). We hypothesized that CO_2

inhalation during retrieval alters both structure and plasticity of dendritic spines. The

molecular mechanism by which CO_2 regulates spine plasticity may explain how CO_2

converts memory into the labile stage.

Using the TetTag-c-fos driven-GFP mouse model, we imaged spine structure and

assessed spine density and morphology in overlapping neurons of the amygdala in

each group (pure tone aversive conditioning, pure tone retrieval and pure tone aversive

conditioning, white noise retrieval) **(Fig. 7A, C)**. Mature spines - most of which display

"mushroom-like" morphology - have more stable postsynaptic structures enriched in

AMPARs. In contrast, immature spines with a "thin-like" morphology, are unstable

postsynaptic structures that have the transitional ability. Immature dendritic spines are

thought to be responsible for synaptic plasticity, as they have the potential for

strengthening (Berry and Nedivi, 2017). The categories of spines were identified based

on the parameters in the previous studies **(Fig. 7G)** (see the details in the Material and

Methods) (Kreple et al., 2014; Wright et al., 2020). The behavior procedure was

described in **Fig. 7C**, the animals underwent aversive conditioning with a tone as a

CS and followed by a retrieval on day 2 with tone or noise. We found increased spine

numbers after aversive conditioning, indicating that aversive conditioning increases

synaptic strength. There was no additional increase in the density of spines in all

groups, suggesting that retrieval and CO_2 inhalation did not change the synaptic

strength **(Fig. 7H, lower-right)**.

We further analyzed spine subtypes as described in the experimental procedures. Thin

spines are deemed to represent the immature structure of the synapses (Berry and

Nedivi, 2017). We examined the ratio of the number of thin spines to the total number of

spines. Interestingly, we found that there were more thin spines in the retrieval group,

which might suggest a greater potential for synaptic plasticity in this group **(Fig. 7H,**

lower-middle). Importantly, when the retrieval group (tone) was paired with CO_2, we

found an additional increase of thin spines compared to the retrieval group alone **(Fig.**

7H, lower-middle). This finding suggests that CO_2 paired with retrieval might boost

synaptic plasticity compared to memory retrieval alone. Consistently, the mushroom

spine numbers decreased in the tone and CO_2 paired retrieval groups, suggesting a

higher turnover rate after memory retrieval **(Fig. 7H, lower-left)**. However, when the

mice were trained with the pure tone and during the retrieval session were presented

with white noise (a generated unrelated CS), we found that in the noise group with or

without CO_2 inhalation, the thin spine number did not increase compared to the pure

tone/retrieval group **(Fig. 7H, lower-middle)**. This finding supports the specific effect of

CO_2 on the memory trace.

DISCUSSION

A newly acquired aversive memory is labile and can be easily disrupted before it is

transformed into a long-term stable state (Alberini, 2011). An existing memory, when

reactivated, may become labile again during a short post-reactivation period known are

reconsolidation window (Schiller et al., 2012). Previous studies using auditory aversive

conditioning found that a retrieval event utilizing a single tone CS renders the memory

labile during the reconsolidation window (Monfils et al., 2009). During this

reconsolidation window, memory is sensitive to the updating processes that may either

enhance or weaken the original memory (Bouton et al., 2020; Du et al., 2017;

Fukushima et al., 2014). The reconsolidation window offers an opportunity to determine

the mechanisms underlying the lability of an existing memory. We have recently found

that when mice inhale 10% CO_2 during retrieval, memory lability increases, and the

original memory may be replaced through the update mechanisms (Du et al., 2017).

Here, we show that the effects of CO_2 on memory are specific to the reactivated

conditioned cue (Debiec et al., 2010; Doyere et al., 2007). To address the specificity of

the effects of CO_2 on reactivated memories, we used a unique behavioral approach with

distinct CSs and US, which allowed us the creation of distinct memories. An application

of CO_2 to the reactivated but not no-reactivated memory enhanced its lability.

We then asked about the cellular mechanisms through which CO_2 enhances the lability

of aversive memory. Memory retrieval-induced lability is glutamate receptor-dependent

(Rose and Rankin, 2006; Milton, A.L., Everitt, B.J. 2013). We have shown that CO_2

activates ASIC1a by decreasing pH in the brain during retrieval and the activation of

ASIC1a increases postsynaptic intracellular calcium and increases AMPAR exchange

(Du et al., 2017). Previous research studying the mechanism of retrieval of aversive

memories have revealed the rapid and transient exchange from CI-AMPARs to CP-

AMPARs in the lateral amygdala synapses (Clem and Huganir, 2010; Hong et al., 2013)

after presenting the CS. In this study, we focused on the specificity of CO_2 effects on the

exchange of the AMPARs in the lateral amygdala's synapses. We conditioned the mice

with a pure tone CS and reactivated the memory with the pure tone with or without

CO_2 inhalation. Consistent with our earlier findings, we observed that CO_2 increases

AMPAR exchange when it is paired with retrieval. However, when mice were presented

with an unrelated CS during the retrieval stage, CO_2 did not alter the AMPAR exchange-

suggesting the effects of CO_2 on memory trace are memory specific. In addition,

synaptic strength (ratio of AMPAR/NMDAR and amplitude of the mEPSCs) was not

altered while applying CO_2 during retrieval, with or without combining with the memory

trace. The exchange of CI-AMPARs to CP-AMPARs indicates a synaptic plasticity

change. Interestingly, CO_2 did not increase the total number of spines compared to the

retrieval alone group, which suggests the strength of the synapse in a memory trace

does not change. In addition, thin spine density significantly increased when retrieval

was combined with CO_2 inhalation, suggesting that CO_2 application changes plasticity.

Using this model, we found when an unrelated CS was presented during the retrieval

session, no additional increase of immature spine density occurred. This finding

indicates no plasticity change in the memory trace. Thus, we can conclude from our

findings that the effects of CO_2 on memory trace are specific.

Although we have provided evidence that CO_2 indeed acts with specificity on a memory

trace, this study did not determine how CO_2 directly regulates a memory. We cannot

exclude the possibility that CO_2 might trigger specific effects on memory through other

targets. For instance, CO_2 inhalation increases cerebral blood flow and arterial blood

pressure and might affect brain functions, such as cognition. Although no direct

evidence supports the possibility that increased cerebral blood flow and arterial blood

pressure affect learning and memory, future studies will have the mechanisms

underlying the specificity of CO_2 on memory. Future studies will also have to examine

whether CO_2 has similar effects on synapses in other brain regions and on other learned

behaviors (*e.g.* appetitive behaviors and drug addictive behaviors).

In conclusion, the effects of CO_2 on the lability of aversive memory are found to be

specific under certain conditions. Our research tests the novel hypothesis that protons

are neurotransmitters that activate the postsynaptic proton receptors, ASICs, to

manipulate memory updates. This non-invasive, drug-free methodology is innovative,

efficacious, and safe for translation to clinical use. As a result, this research may lead to

an effective complementary treatment for many mental health-related disorders for

which efficient treatments are lacking. We hope this research will lead to new areas of

inquiry into CO_2-related mechanisms that underlie memory modification and lead to the

development of novel therapies for anxiety disorders such as PTSD.

ACKNOWLEDGMENTS

We thank Olivia Miller, Melissa Curtis, Nora Abdul-Aziz, Rida Naqvi, Caitlin Kilmurry,

Becca Sturges, Jen Page, Chase Carr, Jordan Jones for their assistance. We thank

Drs. Susumu Tonegawa for providing the TRE-mCherry plasmid. J.Du. is supported by

the National Institutes of Mental Health (1R01MH113986), the University of Toledo

start-up fund, and the University of Tennessee Health Science Center start-up fund.

AUTHOR CONTRIBUTIONS

J.DU., J.D., E.K., C.K. and conceived the project. J.DU., E.K., C.K. and J.D. designed

the experiments. E.K., C.K., J.E., S.B., M.H., B.L., performed the behavior experiments.

K.S., performed the spine morphology experiments and data analysis. F.N., and J.DU.

performed the patch-clamp experiments and data analysis. E.K., and J.DU., wrote the

manuscript. All authors reviewed and edited the manuscript.

DECLARATION OF INTERESTS: The authors declare no competing financial

interests.

FIGURE LEGENDS

Figure 1: CO_2 inhalation during a selective memory retrieval potentiates the effect

of the extinction. **(A)** Representative schematic of protocol for the aversive

conditioning (Av. Cond.), memory retrieval (Ret), extinction (Ext), and memory test-

spontaneous recovery (Spon Rec) and renewal. On day1, the mice were subjected to 3

pure tones and 3 white noises, paired with 6 foot shocks in context A. One day after, the

mice were placed in context B and were subjected to both tone and noise as retrieval

events. 30 mins after retrieval, the mice were treated with 2 blocks of extinctions with a

pure tone as the CS. On day 7, spontaneous recovery and renewal were tested

individually in context B and then context A. 4 pure tones and 4 white noises were

presented as CSs during each memory testing. **(B-E)** Data are presented by the

percentage of freezing time during the CSs (tone and noise) in aversive conditioning

(B), retrievals (tone and noise) **(C)**, two sections of extinction with tone **(D)**, memory test

of spontaneous recovery (Spon Rec) and renewal with tone and noise **(E)**. **(F-I)** Data

are presented by the percentage of freezing time during the CSs (tone and noise) in

aversive conditioning **(F)**, retrievals (tone plus CO_2 inhalation and noise) **(G)**, two

sections of extinction with tone **(H)**, memory test of spontaneous recovery (Spon Rec)

and renewal with tone and noise **(I)**. **(J-K),** comparison data based on spontaneous

recovery and renewal respectively from panels **E** and **I**. Data are mean ± SEM. n = 10-

12 mice in each group. 'n.s.' indicates not statistically significant. * indicates $p<0.05$, **

indicates $p<0.01$, *** indicates $p<0.001$, **** indicates $p<0.0001$, by unpaired Student's t-

test and one-way ANOVA with Tukey's post-hoc multiple comparisons.

See also Figure S1 and Figure S2.

Figure 2: CO_2 inhalation during memory retrieval potentiates the effect of the

anisomycin. (A) Representative schematic of the protocol for aversive conditioning,

memory retrieval, anisomycin injection, and memory test (spontaneous recovery and

renewal). On day1, the mice were subjected to 6 pure tones, paired with 6 foot shocks

in context A. One day after CS, the mice were placed in context B and were subjected

to one single tone as a retrieval event with or without CO_2 inhalation. 30 mins after

retrieval, the mice were then infused with 62.5 µg/µl anisomycin or saline and then

returned to their home cage. On day 7, spontaneous recovery and renewal were tested

individually in context B and then context A. 4 pure tones were presented as CSs during

each memory testing. **(B-E)** Data are presented by the percentage of freezing time

during the tone presentation in aversive conditioning **(B)**, retrieval (tone) **(C)**, saline or

anisomycin infusion in the amygdala **(D)**, spontaneous recovery (Spon Rec), and

renewal test with tones **(E)**. **(F-I)** Data are presented by the percentage of freezing time

during the tone presentation in aversive conditioning **(F)**, retrieval (tone) with or without

CO_2 **(G)**, anisomycin infusion **(H)**, spontaneous recovery (Spon Rec), and renewal test

with tones **(I)**. Data are mean ± SEM. n = 8-10 mice in each group. * indicates $p<0.05$,

*** indicates $p<0.001$, by unpaired Student's t-test. See also Fig. s3, s4, s5

Figure 3: The effect of CO_2 inhalation on selective memory retrieval is ASIC1a-

dependent. (A) Representative schematic of protocol for the aversive conditioning,

memory retrieval, extinction, and memory test (spontaneous recovery and renewal) in

ASIC1a[-/-] mice. On day1, the mice were subjected to 3 pure tones and 3 white noises,

paired with 6 foot shocks in context A. One day later, the mice were placed in context B

and subjected to both tone and noise as retrieval events. 30 mins after retrieval, the

mice were treated with 2 blocks of extinctions with a pure tone as the CS. On day 7,

spontaneous recovery and renewal were tested individually in context B and then

context A. 4 pure tones and 4 white noises were presented as CSs during each memory

testing. **(B-E)** Data in ASIC1a$^{-/-}$ mice are presented by the percentage of freezing time

during the CSs (tone and noise) in aversive conditioning **(B)**, retrievals (tone and noise)

(C), two sections of extinction with tone **(D)**, memory test of spontaneous recovery

(Spon Rec) and renewal with tone and noise **(E)**. **(F-I)** Data in ASIC1a$^{-/-}$ mice in

aversive conditioning **(F)**, retrievals (pure tone plus 10% CO_2 inhalation and white noise)

(G), two sections of extinction with pure tone **(H)**, memory test of spontaneous recovery

(Spon Rec) and renewal with tone and noise **(I)**. **(J-K),** comparison data based on

spontaneous recovery and renewal respectively from panels **E** and **I**. Data are mean ±

SEM. n = 12-16 mice in each group. * indicates $p<0.05$, ** indicates $p<0.01$, ***

indicates $p<0.001$, by unpaired Student's t-test and one-way ANOVA with Tukey's post-

hoc multiple comparisons.

Figure 4: The CO_2 potentiated anisomycin effects are ASIC1a-dependent.

(A) Representative schematic of protocol for the aversive conditioning, memory

retrieval, anisomycin injection, and memory test (spontaneous recovery and renewal).

On day 1, the mice were subjected to 6 pure tones, paired with 6 foot shocks in context

A. One day after, the mice were placed in context B and subjected to one single tone as

a retrieval event with or without CO_2 inhalation. 30 mins after retrieval, the mice were

infused with 62.5 µg/µl anisomycin or saline and then returned to their home cage. On

day 7, spontaneous recovery and renewal were tested individually in context B and then

context A. 4 pure tones were presented as CSs during each memory testing. **(B-E)** Data in ASIC1a[-/-] mice are presented by the percentage of freezing time during the tone presentation in aversive conditioning **(B)**, retrieval (tone) with or without CO_2 **(C)**, anisomycin infusion in the amygdala **(D)**, spontaneous recovery (Spon Rec) and renewal test with tones **(E)**. **(F)** Representative schematic of protocol for the aversive conditioning, memory retrieval, extinction, and memory test (spontaneous recovery and renewal). On day 1, ASIC1a[-/-] mice were subjected to 3 pure tones and 3 white noises, paired with 6 foot shocks in context A. One day later, the mice were placed in context B and subjected to both tone and noise as retrieval events. 30 mins after retrieval, the mice were infused with anisomycin or saline and returned to their home cage. On day 7, spontaneous recovery and renewal were tested individually in context B and then context A. 4 pure tones and 4 white noises were presented as CSs during each memory testing. **(G-J)** Data are presented by the percentage of freezing time during the tone presentation in aversive conditioning **(G)**, retrieval (noise plus CO_2 inhalation, then tone) **(H)**, anisomycin infusion in the amygdala **(I)**, spontaneous recovery (Spon Rec), and renewal test with tones **(J)**. **(K-N)** Data are presented by the percentage of freezing time during the tone presentation in aversive conditioning **(K)**, retrieval (tone) with or without CO_2 **(L)**, anisomycin infusion in the amygdala **(M)**, spontaneous recovery (Spon Rec), and renewal test with tones **(N)**. Data are mean ± SEM. n = 12-16 mice in each group. 'n.s.' indicates not statistically significant by unpaired Student's t-test between groups.

Figure 5: Blockage of ASIC1a in the amygdala reduces the CO_2 effects on selective memory retrieval. (A) Representative schematic of protocol for the aversive conditioning, PcTX-1 injection, memory retrieval, extinction, and memory test

(spontaneous recovery and renewal). On day1, the mice were subjected to 3 pure tones

and 3 white noises, paired with 6 foot shocks in context A. One day later, the mice were

injected with 100nM PcTX-1 or saline, then the mice were placed in context B and

subjected to both tone and noise as retrieval events with or without CO_2. 30 mins after

retrieval, the mice were treated with 2 blocks of extinctions with a pure tone as the CS.

On day 7, spontaneous recovery and renewal were tested individually in context B and

then context A. 4 pure tones and 4 white noises were presented as CSs during each

memory testing. **(B-E)** Data are presented by the percentage of freezing time during the

CSs (tone and noise) in aversive conditioning **(B)**, retrievals (tone plus CO_2 inhalation

and noise) after saline injection in the amygdala **(C)**, two sections of extinction with tone

(D), memory test of spontaneous recovery (Spon Rec) and renewal with tone and noise

(E). **(F-I)** Data are presented by the percentage of freezing time during the CSs (tone

and noise) in aversive conditioning **(F)**, retrievals (tone plus CO_2 inhalation and noise)

after PcTx-1 injection in the amygdala **(G)**, two sections of extinction with tone **(H)**,

memory test of spontaneous recovery (Spon Rec) and renewal with tone and noise **(I)**.

(J-K), comparison data based on spontaneous recovery and renewal respectively from

panels **E** and **I**. Data are mean ± SEM. n = 12 mice in each group. 'n.s.' indicates not

statistically significant. * indicates $p<0.05$, ** indicates $p<0.01$, *** indicates $p<0.001$, ****

indicates $p<0.0001$, by unpaired Student's t-test and one-way ANOVA with Tukey's

post-hoc multiple comparisons.

Figure 6: CO_2 inhalation during a selective memory retrieval enhances the

retrieval dependent AMPAR current rectification. (A) Representative schematic of

the protocol. On day 1, the animal underwent 6 CSs (tones or noses), paired with 6-foot

shocks in context A. On day 2, the mice were divided into 4 groups for the retrieval-pure

tone only; pure tone plus 10% CO_2 inhalation; white noise only; white noise+ 10% CO_2

inhalation. Ten minutes after retrieval, the brain slices were dissected and AMPAR

current was recorded in the pyramidal neurons in the lateral amygdala through

stimulation of thalamic input. **(B-E)** Mice underwent 6 pure tones **(B)** or 6 noises **(D)** in

aversive conditioning, Data are presented by the percentage of freezing time during the

tone presentation in aversive conditioning, retrieval (noise plus CO_2 inhalation, then

tone). **(C, E)** Left, AMPARs current-voltage relationships in the recorded neurons. Insets

show an example of the AMPAR-EPSCs in -80mV and + 60mV. Right, AMPAR

rectification index (I-80 mV / I+60 mV). Data are mean±SEM. n = 20-26 for each group.

(F) Left, examples of EPSC recordings of AMPAR-EPSCs (-80mV) and NMDAR-

EPSCs (+60mV). Right, AMPAR/NMDAR EPSC ratios. Current amplitudes were

measured 70 ms after onset. n = 20-26 for each group. **(G)** Miniature EPSCs recordings

from the neurons after retrieval. Upper, representative mEPSC traces from different

groups. Lower, cumulative distributions of mEPSC amplitudes, inter-event intervals, and

decay-times. n = 25-40 for each group. Data are mean ± SEM. 'n.s.' indicates not

statistically significant. * indicates $p < 0.05$, ** indicates $p < 0.01$, by ANOVA with Tukey's

post hoc multiple comparisons.

Figure 7: CO_2 inhalation during a selective memory retrieval enhances the

retrieval-related memory trace. (A) Schematic showing the c-Fos-tTA-GFP mouse

system combined with an $AAV_{2/9}$-mCherry to label a specific memory trace. The Fos

promoter in transgenic mice was activated by activities, followed by a transient, 2-hour

half-life, GFP expression in the cells. When the $AAV_{2/9}$-mCherry virus was injected into

the brain, the activation of c-FOS also induces the expression of mCherry. When the

mice were fed with DOX, the mCherry expression was interrupted. **(B)** an example

image showing the efficiency of the expression of GFP and mCherry in the amygdala.

(C) The procedure of aversive conditioning, memory retrieval, and memory trace

labeling using the system in A. Mice was fed with DOX for at least one week, followed

by an injection of $AAV_{2/9}$-mCherry in the amygdala. Two weeks later, DOX was removed

and mice were subjected to aversive conditioning with a pure tone as the CS. DOX has

added back again immediately after aversive conditioning. 24 hours later, the mice

underwent retrieval with pure tone or noise. The brain slices were collected 30 mins

after retrieval and then were stained for microscopy. **(D)** Left, representative images of

the neurons labeled by mCherry, GFP, and DAPI; Right, the enlarged area from the

"merge" image showing the overlapping expression of mCherry and GFP neurons. The

overlapping neurons indicate their "consanguinity" in the same memory trace. **(E)**

Summarized data are the percentage of the overlapping expression of mCherry and

GFP neurons in different behavior groups. All mice underwent aversive conditioning

with a tone as the CS. 24 hours later, the mice were separated into 4 groups. Tone

group: retrieval CS by tone; Tone+CO_2 group: retrieval CS by tone along with CO_2

inhalation; Noise group: retrieval CS by noise; Noise +CO_2 group: retrieval CS by noise

along with CO_2 inhalation. **(F)** Control experiment showing the expression of mCherry

with or without DOX as well as with or without aversive conditioning. **(G)** Left, a

representative image showing the spine morphology in the mCherry and GFP

colocalized neurons. The mature spines were categorified as "mushroom" spines and the immature spines were categorified as "thin" spines; Right, an enlarged image showing the details of mushroom and thin spines. **(H)** Upper, representative images of the spine structures in different animal groups showed in E; Lower, summarized data of the spine densities of mushroom spines, thin spines, and total spines in the different groups. Data are mean ± SEM. n =8-12 mice for each group. 'n.s.' indicates not statistically significant. * indicates $p<0.05$, *** indicates $p<0.001$, **** indicates $p<0.0001$,by unpaired Student's t-test and one-way ANOVA with Tukey's post-hoc multiple comparisons.

Figure s1: White noise was an appropriate successful CS to involve use in studying the effects of CO_2 effects on aversive conditioning and retrieval.

(A) Representative schematic of protocol for the aversive conditioning (pure tone and white noise), memory retrieval (pure tone and white noise), extinction (white noise), and memory test (spontaneous recovery and renewal). **(B-E)** Data are presented by the percentage of freezing time during the CSs (tone and noise) in aversive conditioning **(B)**, retrievals (tone and noise) **(C)**, two sections of extinction with white noise **(D)**, memory test of spontaneous recovery (Spon Rec) and renewal with tone and noise **(E)**. **(F-I)** Data are presented by the percentage of freezing time during the CSs (tone and noise) in aversive conditioning **(F)**, retrievals (pure tone and white noise plus CO_2 inhalation) **(G)**, two sections of extinction with white noise **(H)**, memory test of spontaneous recovery (Spon Rec) and renewal with tone and noise **(I)**. **(J-K)**, comparison data based on spontaneous recovery and renewal respectively from panels

E and **I**. Data are mean ± SEM. n = 12 mice in each group. 'n.s.' indicates not

statistically significant. * indicates p<0.05, ** indicates p<0.01, *** indicates p<0.001, ***

indicates p<0.0001, by unpaired Student's t-test and one-way ANOVA with Tukey's

post-hoc multiple comparisons.

Figure s2: CO_2 inhalation does not boost extinction when pairs with an unrelated

CS during memory retrieval. (A) Representative schematic of protocol for the aversive

conditioning (pure tone and white noise), memory retrieval (pure tone and white noise),

extinction (white noise), and memory test (spontaneous recovery and renewal). **(B-**

E) Data are presented by the percentage of freezing time during the CSs (tone and

noise) in aversive conditioning **(B)**, retrievals (pure tone and white noise plus CO_2

inhalation) **(C)**, two sections of extinction with pure tone **(D)**, memory test of

spontaneous recovery (Spon Rec) and renewal with tone and noise, Blue arrow and %

indicated the difference (decreases) between the tone and noise groups **(E)**. **(F-I)** Data

are presented by the percentage of freezing time during the CSs (tone and noise) in

aversive conditioning **(F)**, retrievals (pure tone plus CO_2 inhalation and white noise) **(G)**,

two sections of extinction with white noise **(H)**, memory test of spontaneous recovery

(Spon Rec) and renewal with tone and noise, Blue arrow and % indicated the difference

(decreases) between the tone and noise groups. **(I)**. Data are mean ± SEM. n = 12 mice

in each group. * indicates p<0.05 by unpaired Student's t-test.

Figure s3: Anisomycin disrupts the memory within the reconsolidation window.

(A) Representative schematic of protocol for the aversive conditioning (pure tone), with

or without memory retrieval (pure tone), anisomycin infusion, and memory test

(spontaneous recovery and renewal). **(B-E)** Data are presented by the percentage of freezing time during the tone presentation in aversive conditioning **(B)**, with or without retrieval (tone) **(C)**, anisomycin infusion in the amygdala **(D)**, spontaneous recovery (Spon Rec), and renewal test with tones **(E)**. **(F-I)** Data are presented by the percentage of freezing time during the CSs (tone and noise) in aversive conditioning **(F)**, retrievals (pure tone and white noise) **(G)**, anisomycin injection in the amygdala **(H)**, memory test of spontaneous recovery and renewal with tone and noise **(I)**. **(J-M)** Data are presented by the percentage of freezing time during the CSs (tone and noise) in aversive conditioning **(J)**, retrievals (pure tone plus CO_2 inhalation and white noise) **(K)**, saline injection in the amygdala **(L)**, memory test of spontaneous recovery and renewal with tone and noise, **(M)**. Data are mean ± SEM. n = 11-12 mice in each group. 'n.s.' indicates not statistically significant. * indicates $p < 0.05$ by unpaired Student's t-test.

Figure s4: CO_2 enhances memory retrieval and potentiates the effects of anisomycin on the reconsolidation. (A) Representative schematic of protocol for the aversive conditioning (pure tone and white noise), memory retrieval (pure tone and white noise) with or without CO_2 inhalation, anisomycin infusion, and memory test (spontaneous recovery and renewal). **(B-E)** Data are presented by the percentage of freezing time during the CSs (tone and noise) presentation in aversive conditioning **(B)**, retrieval (tone plus CO_2 and noise) **(C)**, anisomycin infusion in the amygdala **(D)**, spontaneous recovery (Spon Rec) and renewal test with tone and noise **(E)**. **(F-I)** Data are presented by the percentage of freezing time during the CSs (tone and noise) in aversive conditioning **(F)**, retrieval (noise plus CO_2 and tone) **(G)**, anisomycin infusion in the amygdala **(H)**, memory test of spontaneous recovery (Spon Rec) and renewal with

tone and noise **(I)**. **(J-M)** Data are presented by the percentage of freezing time during

the CSs (tone and noise) in aversive conditioning **(J)**, retrievals (tone and noise) plus

CO_2 **(K)**, saline infusion in the amygdala **(L)**, memory test of spontaneous recovery

(Spon Rec) and renewal with tone and noise **(M)**. **(N-Q)** Data are presented by the

percentage of freezing time during the CSs (tone and noise) in aversive conditioning

(N), retrievals (tone and noise) plus CO_2 **(O)**, anisomycin infusion in the amygdala **(P)**,

memory test of spontaneous recovery (Spon Rec) and renewal with tone and noise **(Q)**.

Data are mean ± SEM. n = 12 mice in each group. 'n.s.' indicates not statistically

significant. * indicate $p<0.05$ by unpaired Student's t-test.

Figure s5: CO_2 does not affect memory reconsolidation without anisomycin.

(A) Representative schematic of protocol for the aversive conditioning (pure tone and

white noise), memory retrieval (pure tone and white noise) with or without CO_2

inhalation, saline infusion, and memory test (spontaneous recovery and renewal). **(B-**

E) Data are presented by the percentage of freezing time during the CSs (tone and

noise) presentation in aversive conditioning **(B)**, retrieval (tone plus CO_2 and noise) **(C)**,

saline infusion in the amygdala **(D)**, spontaneous recovery (Spon Rec) and renewal test

with tone and noise **(E)**. **(F-I)** Data are presented by the percentage of freezing time

during the CSs (tone and noise) in aversive conditioning **(F)**, retrieval (noise plus CO_2

and tone) **(G)**, saline infusion in the amygdala **(H)**, memory test of spontaneous

recovery (Spon Rec) and renewal with tone and noise **(I)**. Data are mean ± SEM. n = 12

mice in each group. 'n.s.' indicates not statistically significant by unpaired Student's t-

test.

REFERENCES

Agren, T., Engman, J., Frick, A., Bjorkstrand, J., Larsson, E.M., Furmark, T., and Fredrikson, M. (2012). Disruption of reconsolidation erases a fear memory trace in the human amygdala. Science *337*, 1550-1552.

Alberini, C.M. (2011). The role of reconsolidation and the dynamic process of long-term memory formation and storage. Front Behav Neurosci *5*, 12.

Auber, A., Tedesco, V., Jones, C.E., Monfils, M.H., and Chiamulera, C. (2013). Post-retrieval extinction as reconsolidation interference: methodological issues or boundary conditions? Psychopharmacology (Berl) *226*, 631-647.

Bailey, C.H., Kandel, E.R., and Harris, K.M. (2015). Structural Components of Synaptic Plasticity and Memory Consolidation. Cold Spring Harb Perspect Biol *7*, a021758.

Ben-Shahar, Y. (2011). Sensory functions for degenerin/epithelial sodium channels (DEG/ENaC). Adv Genet *76*, 1-26.

Berry, K.P., and Nedivi, E. (2017). Spine Dynamics: Are They All the Same? Neuron *96*, 43-55.

Bjorkstrand, J., Agren, T., Ahs, F., Frick, A., Larsson, E.M., Hjorth, O., Furmark, T., and Fredrikson, M. (2016). Disrupting Reconsolidation Attenuates Long-Term Fear Memory in the Human Amygdala and Facilitates Approach Behavior. Curr Biol *26*, 2690-2695.

Bourne, J.N., and Harris, K.M. (2008). Balancing structure and function at hippocampal dendritic spines. Annu Rev Neurosci *31*, 47-67.

Bouton, M.E., Maren, S., and McNally, G.P. (2020). Behavioral and Neurobiological Mechanisms of Pavlovian and Instrumental Extinction Learning. Physiol Rev.

Chan, W.Y., Leung, H.T., Westbrook, R.F., and McNally, G.P. (2010). Effects of recent exposure to a conditioned stimulus on extinction of Pavlovian fear conditioning. Learn Mem *17*, 512-521.

Chiang, P.H., Chien, T.C., Chen, C.C., Yanagawa, Y., and Lien, C.C. (2015). ASIC-dependent LTP at multiple glutamatergic synapses in amygdala network is required for fear memory. Sci Rep *5*, 10143.

Chu, X.P., and Xiong, Z.G. (2012). Physiological and pathological functions of acid-sensing ion channels in the central nervous system. Curr Drug Targets *13*, 263-271.

Clem, R.L., and Huganir, R.L. (2010). Calcium-permeable AMPA receptor dynamics mediate fear memory erasure. Science *330*, 1108-1112.

957 Cohen, J.L., Jackson, N.L., Ballestas, M.E., Webb, W.M., Lubin, F.D., and Clinton, S.M.
958 (2017). Amygdalar expression of the microRNA miR-101a and its target Ezh2 contribute
959 to rodent anxiety-like behaviour. Eur J Neurosci *46*, 2241-2252.
960
961 Coryell, M.W., Ziemann, A.E., Westmoreland, P.J., Haenfler, J.M., Kurjakovic, Z., Zha,
962 X.M., Price, M., Schnizler, M.K., and Wemmie, J.A. (2007). Targeting ASIC1a reduces
963 innate fear and alters neuronal activity in the fear circuit. Biol Psychiatry *62*, 1140-1148.
964
965 Costanzi, M., Cannas, S., Saraulli, D., Rossi-Arnaud, C., and Cestari, V. (2011).
966 Extinction after retrieval: effects on the associative and nonassociative components of
967 remote contextual fear memory. Learn Mem *18*, 508-518.
968
969 Das, A.T., Tenenbaum, L., and Berkhout, B. (2016). Tet-On Systems For Doxycycline-
970 inducible Gene Expression. Curr Gene Ther *16*, 156-167.
971
972 Debiec, J., Diaz-Mataix, L., Bush, D.E., Doyere, V., and Ledoux, J.E. (2010). The
973 amygdala encodes specific sensory features of an aversive reinforcer. Nat Neurosci *13*,
974 536-537.
975
976 Doyere, V., Debiec, J., Monfils, M.H., Schafe, G.E., and LeDoux, J.E. (2007). Synapse-
977 specific reconsolidation of distinct fear memories in the lateral amygdala. Nat Neurosci
978 *10*, 414-416.
979
980 Du, J., Price, M.P., Taugher, R.J., Grigsby, D., Ash, J.J., Stark, A.C., Hossain Saad,
981 M.Z., Singh, K., Mandal, J., Wemmie, J.A.*, et al.* (2017). Transient acidosis while
982 retrieving a fear-related memory enhances its lability. Elife *6*.
983
984 Du, J., Reznikov, L.R., Price, M.P., Zha, X.M., Lu, Y., Moninger, T.O., Wemmie, J.A.,
985 and Welsh, M.J. (2014). Protons are a neurotransmitter that regulates synaptic plasticity
986 in the lateral amygdala. Proc Natl Acad Sci U S A *111*, 8961-8966.
987
988 Dulla, C.G., Dobelis, P., Pearson, T., Frenguelli, B.G., Staley, K.J., and Masino, S.A.
989 (2005). Adenosine and ATP link PCO2 to cortical excitability via pH. Neuron *48*, 1011-
990 1023.
991
992 Duvarci, S., and Nader, K. (2004). Characterization of fear memory reconsolidation. J
993 Neurosci *24*, 9269-9275.
994
995 Farley, S.J., Albazboz, H., De Corte, B.J., Radley, J.J., and Freeman, J.H. (2018).
996 Amygdala central nucleus modulation of cerebellar learning with a visual conditioned
997 stimulus. Neurobiol Learn Mem *150*, 84-92.
998
999 Fukushima, H., Zhang, Y., Archbold, G., Ishikawa, R., Nader, K., and Kida, S. (2014).
1000 Enhancement of fear memory by retrieval through reconsolidation. Elife *3*, e02736.

1001   Gao, J., Duan, B., Wang, D.G., Deng, X.H., Zhang, G.Y., Xu, L., and Xu, T.L. (2005).
1002   Coupling between NMDA receptor and acid-sensing ion channel contributes to ischemic
1003   neuronal death. Neuron *48*, 635-646.

1004
1005   Gipson, C.D., and Olive, M.F. (2017). Structural and functional plasticity of dendritic
1006   spines - root or result of behavior? Genes Brain Behav *16*, 101-117.

1007
1008   Golkar, A., Bellander, M., Olsson, A., and Ohman, A. (2012). Are fear memories
1009   erasable?-reconsolidation of learned fear with fear-relevant and fear-irrelevant stimuli.
1010   Front Behav Neurosci *6*, 80.

1011
1012   Goode, T.D., Holloway-Erickson, C.M., and Maren, S. (2017). Extinction after fear
1013   memory reactivation fails to eliminate renewal in rats. Neurobiol Learn Mem *142*, 41-47.

1014
1015   Highstein, S.M., Holstein, G.R., Mann, M.A., and Rabbitt, R.D. (2014). Evidence that
1016   protons act as neurotransmitters at vestibular hair cell-calyx afferent synapses. Proc
1017   Natl Acad Sci U S A *111*, 5421-5426.

1018
1019   Hong, I., Kim, J., Kim, J., Lee, S., Ko, H.G., Nader, K., Kaang, B.K., Tsien, R.W., and
1020   Choi, S. (2013). AMPA receptor exchange underlies transient memory destabilization
1021   on retrieval. Proc Natl Acad Sci U S A *110*, 8218-8223.

1022
1023   Huang, F., Zou, G., Li, C., Meng, H., Liu, X., and Yang, Z. (2020). A novelty-retrieval-
1024   extinction paradigm leads to persistent attenuation of remote fear memories. Sci Rep
1025   *10*, 3319.

1026
1027   Ishii, D., Matsuzawa, D., Matsuda, S., Tomizawa, H., Sutoh, C., and Shimizu, E. (2012).
1028   No erasure effect of retrieval-extinction trial on fear memory in the hippocampus-
1029   independent and dependent paradigms. Neurosci Lett *523*, 76-81.

1030
1031   Jarome, T.J., Ferrara, N.C., Kwapis, J.L., and Helmstetter, F.J. (2015). Contextual
1032   Information Drives the Reconsolidation-Dependent Updating of Retrieved Fear
1033   Memories. Neuropsychopharmacology *40*, 3044-3052.

1034
1035   Kawasaki, H., Eguchi, S., Miyashita, S., Chan, S., Hirai, K., Hobara, N., Yokomizo, A.,
1036   Fujiwara, H., Zamami, Y., Koyama, T.*, et al.* (2009). Proton acts as a neurotransmitter
1037   for nicotine-induced adrenergic and calcitonin gene-related peptide-containing nerve-
1038   mediated vasodilation in the rat mesenteric artery. J Pharmacol Exp Ther *330*, 745-755.

1039
1040   Kida, S. (2020). Function and mechanisms of memory destabilization and
1041   reconsolidation after retrieval. Proc Jpn Acad Ser B Phys Biol Sci *96*, 95-106.

1042
1043   Klucken, T., Kruse, O., Schweckendiek, J., Kuepper, Y., Mueller, E.M., Hennig, J., and
1044   Stark, R. (2016). No evidence for blocking the return of fear by disrupting
1045   reconsolidation prior to extinction learning. Cortex *79*, 112-122.

Koffman, E.E., and Du, J. (2017). Labeling Aversive Memory Trace in Mouse Using a Doxycycline-inducible Expression System. Bio-protocol *7*, e2578.

Kreple, C.J., Lu, Y., Taugher, R.J., Schwager-Gutman, A.L., Du, J., Stump, M., Wang, Y., Ghobbeh, A., Fan, R., Cosme, C.V.*, et al.* (2014). Acid-sensing ion channels contribute to synaptic transmission and inhibit cocaine-evoked plasticity. Nat Neurosci *17*, 1083-1091.

Lee, J.L. (2009). Reconsolidation: maintaining memory relevance. Trends Neurosci *32*, 413-420.

Liu, J., Zhao, L., Xue, Y., Shi, J., Suo, L., Luo, Y., Chai, B., Yang, C., Fang, Q., Zhang, Y.*, et al.* (2014). An unconditioned stimulus retrieval extinction procedure to prevent the return of fear memory. Biol Psychiatry *76*, 895-901.

Liu, M.G., Li, H.S., Li, W.G., Wu, Y.J., Deng, S.N., Huang, C., Maximyuk, O., Sukach, V., Krishtal, O., Zhu, M.X.*, et al.* (2016). Acid-sensing ion channel 1a contributes to hippocampal LTP inducibility through multiple mechanisms. Sci Rep *6*, 23350.

Liu, X., Ramirez, S., Pang, P.T., Puryear, C.B., Govindarajan, A., Deisseroth, K., and Tonegawa, S. (2012). Optogenetic stimulation of a hippocampal engram activates fear memory recall. Nature *484*, 381-385.

Mactutus, C.F., Riccio, D.C., and Ferek, J.M. (1979). Retrograde amnesia for old (reactivated) memory: some anomalous characteristics. Science *204*, 1319-1320.

Mango, D., and Nisticò, R. (2019). Acid-Sensing Ion Channel 1a Is Involved in N-Methyl D-Aspartate Receptor-Dependent Long-Term Depression in the Hippocampus. Front Pharmacol *10*, 555.

Monfils, M.H., Cowansage, K.K., Klann, E., and LeDoux, J.E. (2009). Extinction-reconsolidation boundaries: key to persistent attenuation of fear memories. Science *324*, 951-955.

Nader, K., Schafe, G.E., and Le Doux, J.E. (2000). Fear memories require protein synthesis in the amygdala for reconsolidation after retrieval. Nature *406*, 722-726.

Ortega-Ramírez, A., Vega, R., and Soto, E. (2017). Acid-Sensing Ion Channels as Potential Therapeutic Targets in Neurodegeneration and Neuroinflammation. Mediators Inflamm *2017*, 3728096.

Quirk, G.J., Paré, D., Richardson, R., Herry, C., Monfils, M.H., Schiller, D., and Vicentic, A. (2010). Erasing fear memories with extinction training. J Neurosci *30*, 14993-14997.

1090 Ramirez, S., Liu, X., Lin, P.A., Suh, J., Pignatelli, M., Redondo, R.L., Ryan, T.J., and
1091 Tonegawa, S. (2013). Creating a false memory in the hippocampus. Science *341*, 387-
1092 391.
1093
1094 Rao, V.R., and Finkbeiner, S. (2007). NMDA and AMPA receptors: old channels, new
1095 tricks. Trends Neurosci *30*, 284-291.
1096
1097 Rose, J.K., and Rankin, C.H. (2006). Blocking memory reconsolidation reverses
1098 memory-associated changes in glutamate receptor expression. J Neurosci *26*, 11582-
1099 11587.
1100
1101 Sara, S.J. (2000). Retrieval and reconsolidation: toward a neurobiology of remembering.
1102 Learn Mem *7*, 73-84.
1103
1104 Sara, S.J., and Hars, B. (2006). In memory of consolidation. Learn Mem *13*, 515-521.
1105 Schiller, D., Monfils, M.H., Raio, C.M., Johnson, D.C., Ledoux, J.E., and Phelps, E.A.
1106 (2010). Preventing the return of fear in humans using reconsolidation update
1107 mechanisms. Nature *463*, 49-53.
1108
1109 Schiller, D., Raio, C.M., and Phelps, E.A. (2012). Extinction training during the
1110 reconsolidation window prevents recovery of fear. J Vis Exp, e3893.
1111
1112 Soto, E., Ortega-Ramírez, A., and Vega, R. (2018). Protons as Messengers of
1113 Intercellular Communication in the Nervous System. Front Cell Neurosci *12*, 342.
1114
1115 Taugher, R.J., Lu, Y., Fan, R., Ghobbeh, A., Kreple, C.J., Faraci, F.M., and Wemmie,
1116 J.A. (2017). ASIC1A in neurons is critical for fear-related behaviors. Genes Brain Behav
1117 *16*, 745-755.
1118
1119 Tronson, N.C., and Taylor, J.R. (2007). Molecular mechanisms of memory
1120 reconsolidation. Nat Rev Neurosci *8*, 262-275.
1121
1122 Waldmann, R., Champigny, G., Bassilana, F., Heurteaux, C., and Lazdunski, M. (1997).
1123 A proton-gated cation channel involved in acid-sensing. Nature *386*, 173-177.
1124
1125 Waldmann, R., and Lazdunski, M. (1998). H(+)-gated cation channels: neuronal acid
1126 sensors in the NaC/DEG family of ion channels. Curr Opin Neurobiol *8*, 418-424.
1127
1128 Wang, Q., Wang, Q., Song, X.L., Jiang, Q., Wu, Y.J., Li, Y., Yuan, T.F., Zhang, S., Xu,
1129 N.J., Zhu, M.X.*, et al.* (2018). Fear extinction requires ASIC1a-dependent regulation of
1130 hippocampal-prefrontal correlates. Sci Adv *4*, eaau3075.
1131
Wemmie, J.A., Askwith, C.C., Lamani, E., Cassell, M.D., Freeman, J.H., Jr., and Welsh,
M.J. (2003). Acid-sensing ion channel 1 is localized in brain regions with high synaptic
density and contributes to fear conditioning. J Neurosci *23*, 5496-5502.

Wemmie, J.A., Chen, J., Askwith, C.C., Hruska-Hageman, A.M., Price, M.P., Nolan,
B.C., Yoder, P.G., Lamani, E., Hoshi, T., Freeman, J.H., Jr.*, et al.* (2002). The acid-
activated ion channel ASIC contributes to synaptic plasticity, learning, and memory.
Neuron *34*, 463-477.

1139

1140   Wemmie, J.A., Coryell, M.W., Askwith, C.C., Lamani, E., Leonard, A.S., Sigmund, C.D.,
1141   and Welsh, M.J. (2004). Overexpression of acid-sensing ion channel 1a in transgenic
1142   mice increases acquired fear-related behavior. Proc Natl Acad Sci U S A *101*, 3621-
1143   3626.

1144

1145   Wemmie, J.A., Price, M.P., and Welsh, M.J. (2006). Acid-sensing ion channels:
1146   advances, questions and therapeutic opportunities. Trends Neurosci *29*, 578-586.

1147

1148   Wemmie, J.A., Taugher, R.J., and Kreple, C.J. (2013). Acid-sensing ion channels in
1149   pain and disease. Nat Rev Neurosci *14*, 461-471.

1150

1151   Woolfrey, K.M., and Srivastava, D.P. (2016). Control of Dendritic Spine Morphological
1152   and Functional Plasticity by Small GTPases. Neural Plast *2016*, 3025948.

1153

1154   Wright, W.J., Graziane, N.M., Neumann, P.A., Hamilton, P.J., Cates, H.M., Fuerst, L.,
1155   Spenceley, A., MacKinnon-Booth, N., Iyer, K., Huang, Y.H.*, et al.* (2020). Silent
1156   synapses dictate cocaine memory destabilization and reconsolidation. Nat Neurosci *23*,
1157   32-46.

1158

1159   Wu, C.T., Haggerty, D., Kemere, C., and Ji, D. (2017). Hippocampal awake replay in
1160   fear memory retrieval. Nat Neurosci *20*, 571-580.

1161

1162   Wu, L.J., Duan, B., Mei, Y.D., Gao, J., Chen, J.G., Zhuo, M., Xu, L., Wu, M., and Xu,
1163   T.L. (2004). Characterization of acid-sensing ion channels in dorsal horn neurons of rat
1164   spinal cord. J Biol Chem *279*, 43716-43724.

1165

1166   Yang, G., Pan, F., and Gan, W.B. (2009). Stably maintained dendritic spines are
1167   associated with lifelong memories. Nature *462*, 920-924.

1168

1169   Zandbergen, J., Pols, H., De Loof, C., Lousberg, H., and Griez, E. (1989). Effect of
1170   hypercapnia and other disturbances in the acid-base-balance on panic disorder. Hillside
1171   J Clin Psychiatry *11*, 185-197.

1172

1173   Ziemann, A.E., Allen, J.E., Dahdaleh, N.S., Drebot, II, Coryell, M.W., Wunsch, A.M.,
1174   Lynch, C.M., Faraci, F.M., Howard, M.A., 3rd, Welsh, M.J.*, et al.* (2009). The amygdala
1175   is a chemosensor that detects carbon dioxide and acidosis to elicit fear behavior. Cell
1176   *139*, 1012-1021.

1177

Fig. 1

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. s1

Fig. s2

Fig. s4

A Av. Cond. — Context A, 24 hr — Tone, noise ± CO_2, Context B, 30 min — Anisomycin injection, Anisomycin/saline, 5 days — Memory tests: Spon Rec, Context B, 30 min — Renewal, Context A

B Av. Cond. — Tone+CO_2, Noise — % of time freezing — Tones+shocks

C Tone+CO_2, Noise

D Anisomycin injection — Mice return to home cage after injection

E Memory tests — Spon Rec, Renewal

F Av. Cond. — Tone, Noise+CO_2 — % of time freezing — Tones+shocks

G Noise+CO_2, Tone

H Anisomycin injection — Mice return to home cage after injection

I Memory tests — Spon Rec, Renewal

J Av. Cond. — CSt (tone), CSn (noise) — % of time freezing — Tones+shocks

K Tone+CO_2, Noise+CO_2

L Saline injection — Mice return to home cage after injection

M Memory tests — Spon Rec, Renewal

N Av. Cond. — CSt (tone), CSn (noise) — % of time freezing — Tones+shocks

O Noise+CO_2, Tone+CO_2

P Anisomycin injection — Mice return to home cage after injection

Q Memory tests — Spon Rec, Renewal

Fig. s5