Rapid selection of HIV envelopes that bind to neutralizing antibody B cell lineage members with functional improbable mutations

Olivia Swanson[1], Brianna Rhodes[1], Avivah Wang[1], Shi-Mao Xia[1], Cooper Melissa[1], Robert Parks[1], Aja Sanzone[1], Mark K. Louder[5], Bob C. Lin[5], Nicole A. Doria-Rose[5], Kevin O. Saunders[1,3], Mattia Bonsignori[1,2], Kevin Wiehe[1,2], Barton F. Haynes[1,2,4], Mihai L. Azoitei[1,2*]

[1]Duke Human Vaccine Institute, Duke University, Durham, NC

[2]Department of Medicine, Duke University, Durham, NC

[3]Department of Surgery, Duke University, Durham, NC

[4]Department of Immunology, Duke University, Durham, NC

[5] Vaccine Research Center, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD

* Correspondence: mihai.azoitei@duke.edu

Summary

Elicitation of broadly neutralizing antibodies (bnAbs) by an HIV vaccine will involve priming the immune system to activate antibody precursors, followed by boosting immunizations to select for antibodies with functional features required for neutralization breadth. The higher the number of mutations necessary for function, the more convoluted are the antibody developmental pathways. HIV bnAbs acquire a large number of somatic mutations, but not all mutations are functionally important. Here we identified a minimal subset of mutations sufficient for the function of to the naturally occurring, more mutated V3-glycan bnAb DH270.6. Using antibody library screening, candidate envelope immunogens that interacted with DH270.6-like antibodies containing this set of key mutations were identified and selected *in vitro*. Our results demonstrate that less complex B cell evolutionary pathways than those naturally observed exist for the induction of HIV bnAbs by vaccination, and establish rational approaches to identify boosting sequential envelope candidate immunogens.

Introduction

A major goal of HIV vaccine development is to elicit broadly neutralizing antibodies (bnAbs) (Haynes et al., 2019; Kwong and Mascola, 2018). High levels of bnAbs are rarely observed upon HIV infection (Doria-Rose et al., 2010; Gray et al., 2009) although ~50% of HIV-infected individuals make detectable levels of bnAbs over time (Hraber et al., 2014). When they do occur, bnAbs typically take years to develop and their maturation follows intricate evolutionary pathways that depend on a complex interplay between viral evolution and immune adaptation (Bonsignori et al., 2017b; Bonsignori et al., 2018; Bonsignori et al., 2016; Doria-Rose and Landais, 2019; Gao et al., 2014; Haynes et al., 2016; Liao et al., 2013b; Wu et al., 2015). One strategy of HIV vaccine development mimics the process of natural bnAb formation by targeting the unmutated common ancestor B cell receptors (BCRs) of naïve B cell precursors of bnAb B cells, and using sequential immunogens to guide otherwise disfavored B cell lineage maturation pathways rapidly acquire functional mutations required for to achieve broad neutralization (Haynes et al., 2019; Haynes et al., 2012; Kwong and Mascola, 2018).

Compared to neutralizing antibodies against other viruses such as influenza or SARS-CoV-2, HIV bnAbs acquire an unusually large number of somatic mutations, ranging from 11% to 42%, during their evolution from the unmutated common ancestor (UCA) to their mature form (Landais and Moore, 2018; Sok et al., 2013; Wiehe et al., 2018). However, only about a third of the sequence changes observed in the development of HIV bnAbs VRC01 and BG18 are sufficient to provide neutralization breadth and potency (Jardine et al., 2016; Steichen et al., 2019). Thus, for all bnAbs, it is unlikely that all acquired mutations are functionally important nor needed for vaccine-induced bnAb development (Georgiev et al., 2014; Wiehe et al., 2018).

Recently, it was shown that HIV bnAbs are enriched for improbable mutations acquired during affinity maturation (Shen et al., 2020; Wiehe et al., 2018). These amino acid changes are expected to occur with low frequency naturally, either because they occur in gene regions that

are not typically targeted by the Activation-Induced Cytidine Deaminase (AID) or due to the

number of nucleotide changes required to transform a respective UCA amino acid into the

mature one (Hwang et al., 2017; Yaari et al., 2013). Notably, improbable mutations typically play

a critical role in HIV bnAb function and represent developmental barriers that need to be

overcome for the development of neutralization breadth (Bonsignori et al., 2017a; Shen et al.,

2020). For example, in the evolution of the DH270.6 bnAb, the acquisition of an improbable

glycine to arginine mutation at position 57 in the heavy chain initiates the development of

heterologous neutralization breadth (Bonsignori et al., 2017a). Similarly, in the VRC34 lineage

the only branch that gives rise to bnAbs acquired a rare tyrosine to proline mutation that is

essential for broad neutralizing activity (Shen et al., 2020). Since improbable functional

mutations represent rare events that restrict bnAb development, antibodies with such sequence

changes will likely need to be explicitly selected by candidate immunogens in vaccinations.

Significant progress has been made in the design of immunogens that activate

precursors of HIV bnAbs such as VRC01, BG18, CH235 and DH270 (Havenar-Daughton et al.,

2018; LaBranche et al., 2019; Saunders et al., 2019; Steichen et al., 2016; Steichen et al.,

2019). Targeting immunogens against these antibodies have been shown to stimulate B cells

displaying the respective UCAs in knock-in mouse models and to promote the early

development of bnAb lineages (Dosenovic et al., 2015b; LaBranche et al., 2019; Saunders et

al., 2019; Steichen et al., 2019). However, how to further induce maturation of bnAb precursors

to attain significant neutralization breadth remains a major challenge. A series of boosting

immunizations will be required following germline BCR targeting, with immunogens needed to

select for bnAb lineage B cells with rare BCR features required for neutralization breadth, such

as CDR loop deletions or insertions, the ability to accommodate HIV Env glycans, or the

presence of functionally important improbable amino acids (Bonsignori et al., 2018; Kepler et al.,

2014; Klein et al., 2013; Wiehe et al., 2018).

The V3-glycan epitope on HIV Env is a region targeted by HIV bnAbs DH270, PGT121

and BG18 (Barnes et al., 2018; Freund et al., 2017; Julien et al., 2013; Mouquet et al., 2012).

Antibodies against this site have diverse immunogenetics, but all recognize the conserved base

of the V3 loop and one or both of the glycans present at positions 301 and 332 (Barnes et al.,

2018; Mouquet et al., 2012). DH270.6 neutralized 55% of the viruses from a representative

global panel of 208 pseudoviruses (IC_{50}=0.08ug/mL) and 77.4% of those viruses containing a

glycan at position 332 (Bonsignori et al., 2017a). The induction of DH270.6-like antibodies by

vaccination should therefore provide significant protection against HIV infections. Unlike other

bnAbs (Walker et al., 2011; Wu et al., 2010), DH270.6 does not acquire any deletions or

insertions in its CDR loops during development. Therefore, the elicitation of DH270.6-like bnAbs

should be in principle easier to attain, and require the selection and maturation of putative

DH270.6-like precursor antibodies with the key functional somatic mutations of DH270.6.

Recently, a germline targeting immunogen was described that robustly activates

DH270.6 precursors in mice (Saunders et al., 2019). This SOSIP Env trimer, named 10.17DT,

elicited DH270.6-like antibodies in DH270.6 UCA knock-in mice. Isolated antibodies from

animals vaccinated with 10.17DT bound the glycan moiety at position 332 and neutralized both

autologous and some heterologous viruses. Importantly, some of the 10.17DT-elicited

antibodies acquired improbable mutations from the UCA that are essential for DH270.6 function

including heavy chain residues Arg57 and Thr98. The 10.17DT immunogen therefore activates

DH270.6 precursors and promotes their acquisition of key functional mutations. In order to

mature these responses, additional boosting immunizations will be required to induce high titers

of antibodies containing other DH270.6 mutations essential for neutralization breadth and

potency. Given the complexity of designing immunogens to guide otherwise disfavored bnAb B

cell lineages to neutralization breadth, and the need to speed up immunogen design towards

rapid HIV vaccine development, a rapid iterative process is needed to rationally identify

sequential immunogens capable of selecting bnAb lineage BCRs containing functional

improbable mutations.

Here we describe a new approach that relies on computational modeling to identify the

functionally important somatic mutations present in HIV bnAbs and a strategy to identify and

validate candidate immunogens that select antibodies with these mutations *in vitro*. We illustrate

this approach by rapidly identifying the minimal functional mutations in V3-glycan bnAb B cell

lineage DH270.6. A subset of 12 out of the 42 acquired mutations accounted for most of the

DH270.6 breadth and potency. The resulting minimized DH270.6 (DH270min) antibody that

contained these functional mutations represented a bnAb evolutionary path that is shorter and

less complex to induce by vaccination. Since the majority of the acquired mutations necessary

for the function of DH270.6 were found to be improbable, antibodies containing these amino

acids are predicted to occur with low frequency *in vivo.* Consequently, immunogens will have to

explicitly target antibody improbable mutation enrichment in order to induce robust protection by

vaccination (Wiehe et al., 2018). To address this issue, we developed a new approach for

immunogen design that relies on high throughput screening of antibody libraries to identify

envelopes that interact with DH270.6-derived antibodies through the key acquired mutation.

Alternative recognition modes of DH270min that employ more probable amino acids at the key

functional sites were explored. This work illustrates a general approach to identify key functional

mutations in HIV bnAbs and describes a high throughput method to rationally discover

immunogens that target antibody functional mutations.

Results

Identification of a subset of DH270.6 acquired mutations sufficient for

neutralization function. DH270.6 acquired 42 mutations during its development from the UCA

to the mature, broadly neutralizing antibody form. Fourteen of these mutations are predicted to

be improbable, i.e. they are expected to occur with low frequency prior to antigenic selection in the germinal center, based on computational simulations with ARMADiLLO (Wiehe et al., 2018). Structural mapping revealed that 14 mutations interact directly with the viral envelope, while the other 28 are located far away from the paratope (Figure 1A). This finding suggests that, as previously shown for other HIV bnAbs (Jardine et al., 2016; Steichen et al., 2019), not all of the DH270.6 acquired mutations are required for broad and potent HIV neutralization. While sequence changes located in the paratope are likely important for function, mutations away from the binding interface could also play an important role in the stability and conformation of the antibody. We used computational modeling with Rosetta (Bender et al., 2016; Das and Baker, 2008; Leaver-Fay et al., 2011) to identify acquired mutations away from the paratope that could affect function. On the backbone of the DH270.6 scFv (PDBID: 6cbp), acquired mutations deemed nonessential were reversed to their UCA identity and the energetics of the new antibodies were assessed computationally (Supplementary Figure 1A). Upon analysis of different sets of acquired mutations, two "minimized" antibodies, DH270min1 and DH270min2, that contained only 17 and 20 of the 42 mutations acquired by DH270.6 respectively, were determined to have energies comparable to DH270.6 by Rosetta (Supplementary Figure 1B), and were subsequently selected for experimental characterization. Both DH270min1 and DH270min2 included acquired mutations located at the binding interface as well as improbable mutations located immediately adjacent to the paratope (Figure 1A, B). DH270min2 contained additional mutations at the interface of the heavy and light chains. Three other antibodies were also tested as reference: DH270min3, that contained only the DH270.6 acquired mutation located in the paratope, as well as DH270Improb and DH270Prob that contained only the acquired improbable and probable mutations respectively. The Rosetta energy of DH270min3 (Supplementary Figure 1B) was less favorable than that of DH270.6, DH270min1 or DH270min2, suggesting that acquired mutations located outside the paratope play an important structural role. Antibodies were expressed recombinantly, purified as IgGs, and tested for

neutralization on a representative panel of 20 DH270.6-sensitive HIV pseudoviruses (Figure

1C). Both DH270min1 and DH270min2 had the same neutralization breadth and potency on this

panel as WT DH270.6, revealing that less than half of the acquired mutations are sufficient for

DH270.6 function. In contrast, DH270min3 showed significantly reduced breadth (66%) and

potency (0.99ug/mL compared to 0.09ug/mL in DH270.6), confirming the Rosetta modeling

prediction that antibody residues outside the binding interface are functionally important.

Remarkably, even though there are twice as many probable mutations as improbable in

DH270.6, DH270Impr retained 80% of the DH270.6 breadth, while DH270Prob did not

neutralize any of the viruses in the panel. Both DH270min1 and DH270min2 contain 13 of the

total 14 improbable mutations acquired by DH270.6, showing that improbable mutations are

essential for bnAb function.

We next determined whether all of the 17 acquired mutations present in DH270min1

were necessary for its breadth, or if the antibody can be further "minimized" by reverting

additional amino acids to their UCA identity. Single mutant DH270min1 antibody variants, where

each of the acquired amino acids maintained from DH270.6 were changed to the corresponding

UCA residue, were expressed recombinantly and tested for neutralization as before (Figure 2A).

Changing five of the acquired amino acids (F32Y$_{HC}$, Q54N$_{HC}$, T60Y$_{HC,}$ K26S$_{LC}$, H32Y$_{LC}$) to their

DH270UCA identity had no effect on the breadth or potency of the resulting DH270min1

variants; five mutations to UCA (D31G$_{HC}$, M51I$_{HC}$, T55S$_{HC}$, G94S$_{LC}$, A97S$_{LC}$) had a moderate

effect on neutralization, while seven (R57G$_{HC}$, T98R$_{HC}$, Y110G$_{HC,}$ Y27S$_{LC}$, Y48L$_{LC}$, F93Y$_{LC}$,

C101F$_{LC}$) significantly affected the breadth and potency of the resulting DH270min1 mutants

(Figure 2A). Based on these results, two other minimized antibody variants, DH270min11, which

combined the 12 mutations that had any effect on antibody breadth, and DH270min12, which

preserved only the seven acquired mutations most critical for function, were developed and

experimentally characterized. Compared to DH270.6 and DH270min1, these antibodies were

less broad and potent on the same panel of 20 DH270.6-sensitive viruses used in the previous

experiments (Figure 2A). To further assess the minimized DH270.6 variants, neutralization was

also measured on a multi-clade panel of 208 viral isolates . DH270.6 was previously shown to

neutralize 114 viruses from the global panel with a mean IC_{50} of 0.136µg/ml (Bonsignori et al.,

2017a). In comparison, DH270min1 neutralized 100 viruses (mean IC_{50}=0.113µg/ml),

DH270min11 neutralized 99 viruses (mean IC_{50}=0.135µg/ml) and DH270min12 neutralized 68

viruses (mean IC_{50}=0.408µg/ml) (Figure 2B, Supplementary Figure 2). Therefore, DH270min11,

which contained only 12 of the 42 acquired mutations, recapitulated 90% of the breadth of

DH270.6, revealing that only a small subset of all DH270.6 acquired mutations is required for

the development of the antibody into its broadly neutralizing form. Nine of the 12 mutations in

DH270min11 ($G31D_{HC}$, $S55T_{HC}$, $G57R_{HC}$, $R98T_{HC}$, $G110Y_{HC}$, $S27Y_{LC}$, $L48Y_{LC}$, $Y93F_{LC}$, $S97A_{LC}$)

interact with the three components of the glycan-V3 epitope, the base of the V3 loop and the

glycans at positions 301 and 332, while three mutations are located away from the paratope and

likely contribute to the stability and conformation of the antibody (Figure 2C). While improbable

mutations make up 33% of the total mutations acquired by DH270.6, they represent 83% (10

out of 12) of those maintained in DH270min11. Taken together these results demonstrate that

only a small fraction (28%) of the mutations acquired by DH270.6 are required for its breadth

and potency. However, the majority of the functionally critical mutations are predicted to occur

with low probability during DH270.6 development and may restrict the development of this

lineage *in vivo*.

Identification of molecules that preferentially bind antibodies containing the

DH270min functional mutations. DH270min11 offers a significantly shorter evolutionary

pathway to elicit DH270.6-like antibodies. In order to identify boosting candidate immunogens to

elicit similar bnAbs, we aimed to find molecules that preferentially interact with the key acquired

mutations in DH270min11. We reasoned that molecules that engage residues critical for

DH270.6 breadth *in vitro* may elicit antibodies with those target residues by vaccination, as was

previously demonstrated for the induction of DH270.6-like precursor antibodies containing the

acquired mutations R57$_{HC}$ and T98$_{HC}$ by the DH270 lineage activating immunogen 10.17DT

(Saunders et al., 2019). To further mature these immune responses to breadth, boosting

immunogens will be required to select additional mutations contained in DH270min11.

We previously isolated multiple HIV quasispecies that developed in subject CH848 in

whom the DH270.6 V3-glycan B cell lineage arose (Bonsignori et al., 2017a). We measured the

ELISA binding of recombinantly expressed CH848 gp120s derived from 96 Envs in this panel to

DH270min1 and to mutated antibodies where each residue maintained from DH270.6 was

reverted to its UCA identity (Figure 3, Supplementary Figure 3). The "selection strength" of each

gp120 for an acquired mutation present in DH270min1 was determined by comparing its

DH270min1 binding to its interaction with a DH270min1 antibody containing the UCA mutation

at the site of interest. The calculated fold decrease in binding indicated the degree to which a

given potential immunogen engages a respective DH270min1 acquired mutation; we predict

that molecules that preferentially interact with a given antibody residue *in vitro* are promising

candidate immunogens to engage and expand antibodies that developed the same amino acid

mutation *in vivo*.

From this panel, we identified molecules that preferentially interacted with eight of the

twelve DH270.6 acquired mutations present in DH270min11, and six of the seven mutations

present in DH270min12 (Figure 3, Supplementary Figure 3). Multiple gp120s preferentially

bound the antibody through the key acquired mutations R57$_{HC}$, T98$_{HC}$, Y110$_{HC}$, Y27$_{LC}$ and

Y48$_{LC}$, while no gp120 selectively interacted with mutations at positions 31$_{HC}$, 51$_{HC}$, 55$_{HC}$ and

101$_{LC}$. Previously, a vaccination regimen made of sequential immunizations with Envs 10.17,

836.31, 358.06, 1432.41, and 526.02 was proposed for the elicitation of DH270.6-like antibodies

(Bonsignori et al., 2017a). These immunogens were chosen based on their predicted ability to

select for the G57R$_{HC}$ improbable mutation and because they have progressively longer V1

loops. These immunogens were included in this analysis and revealed that five out of the six

molecules showed strong selection for at least three mutations (Figure 3B). As a group, this

immunogen panel preferentially engaged 9 out of the 12 acquired mutations in DH270min11

(Figure 3B, Supplementary Figure 3)). However, our results indicate that more efficient

sequential vaccination regimens that use a subset of these previously proposed molecules

combined with newly identified ones could be developed based on our new criteria. In addition

to the engagement of specific mutations, the overall affinity of an immunogen for a target

antibody is also critical (Abbott et al., 2018). Both 836.31 and 1432.41 have a similar mutational

selection profile, but 1432.41 has a significantly higher binding for DH270min1 (Supplementary

Figures 3 and 4), which recommends its use over 836.31. While 526.02 showed strong

selection for a large number of the targeted mutations, its overall affinity for DH270min1 was

found to be low (Supplementary Figure 4). However, our analysis identified another molecule,

526.09, that had a comparable mutational profile, but a significantly higher affinity for

DH270min11 (Supplementary Figure 4). The "minimized" DH270 can therefore be leveraged to

identify molecules that preferentially interact with the key functional mutations of DH270.6.

Characterization of Envs that bind antibodies with multiple DH270min acquired

mutations. In the single mutant DH270min1 antibody screen, we evaluated gp120 molecules

for their ability to interact with one critical mutation at a time. However, in a vaccination scenario,

immunogens will interact with diverse BCRs that contain various amino acids at multiple sites in

the binding interface. To drive activated DH270.6 precursors to breadth, candidate immunogens

need to preferentially select from this diverse BCR repertoire the ones that contain the key

DH270.6 amino acids identified in DH270min1. One way to assess this at the vaccine design

stage is to ensure that selected immunogens preferentially interact with antibodies containing

the DH270.6 amino acids over the ones present in the unmutated common ancestor (UCA). To

measure the ability of our selected molecules to achieve this goal *in vivo*, we established an *in*

vitro approach that employs high throughput screening of single-chain variable fragment (scFv)

libraries displayed on the surface of yeast (Chao et al., 2006). A library was developed that

contains all the possible scFv variants of DH270min1 with either the amino acid present in the

UCA or in DH270.6 at each mutated amino acid position (Figure 4A). Therefore, this library

represents a subset of the possible evolutionary paths of the DH270UCA towards DH270min1.

The library was sorted by FACS for binding to five different envelopes: the germline targeting

immunogen 10.17DT and 1432.41, 836.31, 358.06, and 526.02 identified above (Figure 4B).

While gp120s were used for immunogen identification above due to their availability and ease of

production (Figure 3), candidate immunogens will be expressed as SOSIP Envs for

immunizations and were therefore used in this format for library screening. The sequences of

the selected scFvs bound by each envelope were isolated and analyzed by next generation

sequencing. The frequency of the "mature" DH270.6 amino acids across the selected clones at

each target site was compared to the frequency of that mutation in the naïve library in order to

determine the "enrichment levels" at a given site for the DH270.6 amino acid over the one from

the UCA (Figure 4C). To validate this approach, we first compared the enrichment of amino

acids selected from the scFv library by sorting with a given envelope to the presence of the

same mutations in antibodies elicited by vaccination with the respective molecule. The only

available data set for this analysis comes from 10.17DT, which was previously used to

immunize DH270UCA knock-in mice and the heavy chain diversity of the antibody repertoire

induced by vaccination was described in detail (Saunders et al., 2019). The enrichment of a

given amino acid at a site in the heavy chain of 10.17DT elicited antibodies was determined by

computing the ratio of the experimentally observed versus the computationally expected

frequency for that amino acid with ARMADiLLO (Wiehe et al., 2018). Enrichment of a given

DH270.6 amino acid from the scFv library correlated well with the observed frequency of that

same residue in vaccine elicited antibodies (R^2= 0.625 over 9 positions analyzed) (Figure 4D,

Supplementary Figure 5). Therefore, our scFv library screening approach can describe, at least

qualitatively, the ability of a given envelope to elicit DH270.6-derived antibodies with target

functional mutations.

Library selection and sequence analysis revealed that, taken as a group, 10.17DT,

1432.41 and 358.06 favored the mature over the UCA amino acids at all the key positions in

DH270min11, with the exception of 31_{HC}, 51_{HC} and 93_{LC} (enrichment levels>10%). The acquired

mutations at position 31_{HC} and 93_{LC} are probable and expected to occur with high frequency *in*

vivo, which should facilitate their selection; indeed, significant levels of $D31_{HC}$ are observed in

the repertoire of DH270UCA knock-in mice vaccinated with 10.17DT (Saunders et al., 2019).

The sequence of 10.17DT, 1432.41 and 358.06 envelopes showed an affinity gradient for the

selection of key amino acids at positions 55_{HC}, 57_{HC}, 98_{HC}, 110_{HC}, 94_{LC}, 97_{LC}, 101_{LC}, based on

the progressively higher enrichment levels observed for the target amino acids in the library

clones selected with these three Envs. However, for the robust induction of antibodies with the

full breadth of DH270.6, our results suggest that additional immunogens will likely be needed to

more strongly select for key improbable mutations $M51_{HC}$, $Y27_{LC}$ and $L48_{LC}$. The scFv library

was also labeled with 526.02, but only a small number (<0.1%) of clones showed binding above

background, which impeded sequence analysis (data not shown). These data, corroborated by

the weak binding of 526.02 to DH270min1 (Supplementary Figure 4), suggest that 526.02 only

interacts with DH270.6-derived antibodies that already contain the majority of the functional

mutations required for breadth, and further supports the use of the alternative envelope 526.09

as a final "boost". Taken together, the scFv library screens showed that the sequence of

envelopes composed of 1432.41, 358.06 and 526.09 can preferentially select for antibodies

containing the majority of the acquired mutations present in DH270.6-like bnAbs.

Identification of alternative Env recognition modes by minimized DH270.6

antibodies. The significant number of functional improbable mutations in DH270.6 likely

represents a barrier for the induction of such antibodies by vaccination. While envelopes can be

identified that engage DH270.6-derived antibodies with these rarely occurring amino acids as

shown above, it would be advantageous to discover alternative developmental pathways of

DH270.6-like bnAbs that may occur with higher frequency *in vivo* and may be easier to replicate

by vaccination. Combinations of more probable amino acids, not observed during natural

evolution, may exist at the key positions in DH270min11 resulting in antibodies with alternative

sequences that have similar breadth. In order to determine if such amino acids exist, we

performed high throughput screening of DH270min11 derived scFv libraries on the surface of

yeast. Three separate libraries were constructed that test alternative amino acids in the

DH270min11 paratope regions that contact the different components of the glycan-V3 epitope:

the base of the V3 loop, the N332 and the N301 glycans. Residues involved in the recognition of

the same epitope region were sampled together, such that the total diversity of the resulting

libraries (~10^6 variants) was in the range that can be analyzed by yeast display (Figure 5). The

V3 loop targeting library sampled positions 51_{HC}, 55_{HC}, 57_{HC}, and 97_{LC} from DH270min11, while

the 301 glycan library diversified sites 110_{HC}, 27_{LC}, 94_{LC}, and 101_{LC}; all these sites contained

improbable amino acids maintained in DH270min11. Acquired mutation $Y32F_{HC}$ was also shown

to be important for interactions with the 301 glycan in DH270min11 (Figure 2). However, this

mutation is expected to be probable and was therefore preserved in the 301 glycan libraries.

The N332 library samples all the possible combinations at positions 31_{HC}, 32_{HC}, 98_{HC}, and 48_{LC}.

While the acquired mutation present at position 31_{HC}, G31D, is probable, this site plays a

synergistic role in binding the N332 glycan together with residues at positions 32_{HC}, 98_{HC}, and

48_{LC}. Therefore, position 31_{HC} was also diversified in order to "unlock" potential alternative

recognition modes of this epitope region.

Libraries were sorted in succession for binding to SOSIP Envs that were identified above

(10.17DT, 358.06, 1432.41, and 526.02). Pseudoviruses with Envs from this panel are

progressively harder to neutralize by DH270.6 and by antibodies from the same lineage that

appeared earlier in the development (Supplementary Figure 6), with 10.17DT being the easiest

to neutralize, while 526.02 was the hardest (Bonsignori et al., 2017a). Therefore, successive

selections of the scFv libraries with these envelopes should identify antibody sequences with

increasing breadth for difficult to neutralize viruses (Supplementary Figure 7). Library sorts were

done sequentially with 10.17DT, 358.06, 1432.41 twice and 526.02. Clones selected by an

immunogen were collected, expanded and subjected to additional sorts with the next molecules

in the panel. At the final sort with 526.02 only the scFv variants that showed high affinity were

collected. The DNA of clones bound by 10.17DT, 1432.41 and 526.02 was isolated and

analyzed with PacBio long read sequencing technology to identify the full-length sequence of

selected scFv variants. As expected, the overall sequence diversity of the isolated clones was

reduced from one round of selection to the next, since fewer amino acid variants are expected

to be tolerated in antibodies that bind the Envs of harder to neutralize viruses. Of note, after the

last selection, the clones present with the highest frequency in the sorted libraries contained

combinations of amino acids at the key DH270min11 sites that are not present in either the

DH270.6 or its UCA (Figure 5). This indicates that antibodies with diverse sequences and

recognition modes not yet observed in nature may exist that have DH270.6-like breadth. Using

ARMADiLLO analysis (Wiehe et al., 2018), the expected probability of the newly identified

amino acids at the DH270min11 critical positions was determined (Supplementary Figure 8). At

11 out of the 12 critical positions in DH270min11 at least one other amino acid of similar or

higher probability to the one present in DH270.6 was identified. Notably, positions 27_{LC} and

48_{LC}, where the DH270.6 amino acids occur with expected probabilities of less than 0.1%, 7 and

3 alternate amino acids of higher probability were identified. Taken together, these results show

that diverse amino acids that occur with higher probability *in vivo* are functionally equivalent to

the native ones present at the key DH270.6 paratope sites. Therefore, multiple, more accessible

pathways than the one naturally observed may exist to induce DH270.6-like antibodies by

vaccination.

Discussion

A major task of current HIV vaccine development efforts is to design successful

sequential immunogen regimens capable of binding to B cell receptors of bnAb lineage

members that have functional improbable mutations required for potent bnAb activity. Here we

report a rapid strategy for identification of the minimal functional mutations needed for antibody

breadth and the identification of HIV Envs as sequential immunogen candidates that bind to

bnAb lineage BCRs. Our results show that only a small number of acquired mutations are

necessary for the development of DH270.6 V3-glycan bnAb precursors into broadly neutralizing

antibodies. Twelve amino acid changes from UCA were sufficient to recapitulate 90% of

DH270.6 neutralization. While the majority of these mutations were located in the paratope,

three of them did not directly contact HIV Env. Therefore, residues away from the binding site

also modulate function, likely by affecting the antibody conformation or stability (Klein et al.,

2013). Indeed, DH270min3, which contained only the acquired mutations in the paratope,

displayed only 66% of the breadth of the mature bnAb. These results are in line with those

reported for BG18, another HIV bnAb that targets the same glycan-V3 epitope, but uses a

different binding orientation and develops from other germline genes (Barnes et al., 2018).

Recently, a "minimized" BG18 was described that preserves 22 out of the 60 acquired mutations

and recapitulates 66% of the neutralization breadth of the mature bnAb (Steichen et al., 2019).

minBG18 contains only the acquired mutations located at the binding interface, while all the

amino acids located away from the paratope are reverted to their UCA identity. In comparison,

by considering acquired mutations outside the binding interface in our designs, DH270min11

almost fully recapitulates the breadth of DH270.6, while maintaining a lower percentage (28%)

of the total acquired mutations than minBG18 (36%). VRC01, a bnAb that targets the CD4

binding site on Env, was the first bnAb to be "minimized". minVRC01 retained 93% of the

mature VRC01 breadth by preserving 21 out of 65 acquired mutations together with a three

amino acid deletion in the CDR L1 loop of the inferred UCA (Jardine et al., 2016). minVRC01

was developed by screening 4 different scFv libraries that sampled UCA and mature amino

acids for over 20 selection rounds (Jardine et al., 2016). Compared to this experimentally

intensive approach, our strategy relied on computational and structural analysis in order to

determine the residues critical for the neutralization function of DH270.6 more rapidly and at a

significantly reduced cost. Amino acids that contact the HIV Env were identified from available

structural data (Fera et al., 2018; Saunders et al., 2019), while residues located away from the

interface critical for broad neutralizing function were determined by computational modeling.

This approach identified a subset of 17 out of 42 acquired mutations that recapitulated almost all

of the DH270.6 breadth; subsequent single site mutagenesis studies reduced this subset to 12

mutations in DH270min11. Our combined computational and experimental approach can

therefore efficiently identify functionally important residues that occur during antibody

development, and should be broadly applicable to the study of other heavily mutated bnAbs.

Of the 12 acquired mutations present in the minimized DH270.6, 10 were improbable

based on computational predictions that estimate the propensity of AID to mutate the antibody

gene regions where they are located (Wiehe et al., 2018). HIV bnAbs are enriched for

improbable mutations and such mutations have been shown to have important functional roles

across different antibodies (Bonsignori et al., 2016; Shen et al., 2020). Forty-three percent of the

acquired mutations maintained in the minVRC01 antibody and 27% of those in minBG18 are

improbable (Wiehe et al., 2018). In comparison, while improbable changes constitute 33% of the

acquired mutations in the DH270.6, 83% of the amino acids maintained in DH270min11 are

improbable, highlighting the functional importance of these rarely occurring sequence changes

for the breadth development. This conclusion is further supported by our data showing that a

DH270.6-derived antibody containing only the naturally acquired probable mutations had no

neutralization ability. Since only a small number of sequence changes, but no insertions or

deletions, are required for the development of DH270.6-like bnAbs from germline precursors,

these antibodies are an attractive target for elicitation by vaccination. DH270min11 is only 8%

mutated from the DH270UCA, and represents to our knowledge one of the least mutated HIV

antibodies with broadly neutralizing capabilities (Simonich et al., 2016). While DH270min11

should be easier to induce by vaccination than DH270.6, the presence of a large number of

improbable mutations may present a significant barrier for its elicitation. To address this

challenge, we aimed to identify non-native amino acids, predicted to occur with higher frequency

in vivo, at the key sites in DH270min11. High throughput library screening led to the

identification of other functionally equivalent amino acids that are expected to occur with higher

probability *in vivo*. At each of the 10 improbable sites in DH270min11 at least one other more

probable alternate amino acid was found, with 7 of 10 sites tolerating at least two such residues.

Thus, these results indicate that diverse developmental pathways exist for the elicitation of

DH270.6-like bnAbs, and that such pathways are more direct and expected to occur with higher

frequencies than those pathways naturally observed. Therefore, glycan-V3 targeting bnAbs that

resemble DH270.6/DH270min11 mAbs are an attractive HIV vaccine target.

Compared to previous HIV bnAb minimization studies, we leveraged our minimized

DH270 antibodies to find and characterize envelopes for the induction of similar antibodies by

vaccination. Our approach relied on the identification of proteins that preferentially interacted

with DH270.6-derived antibodies through mutations deemed essential for function. Diverse

molecular properties can control the ability of candidate immunogens to contact different

DH270min11 acquired mutations. The size and branching of glycans present at positions N301

and N332 can have a significant effect on antibody recognition. These glycans are variable

across different Envs and their processing is affected by the presence of neighboring

glycosylation sites, for example those that may exist at the tip of the V1 loop (Cao et al., 2017;

Saunders et al., 2019; Struwe et al., 2018). In addition, sequence variation in the V3 loop can

also affect molecular contacts with the acquired mutations. Envelopes were chosen from a

panel of molecules previously selected from the patient who gave rise to the DH270 lineage

(Bonsignori et al., 2017a), based on their ability to differentially bind to minimized antibodies that

either contain or lack a target mutation. Chosen envelopes were further tested in a high

throughput fashion for their ability to select antibodies containing multiple DH270min11 acquired

mutations *in vitro*. Characterization of diverse molecules in this fashion can rationally and rapidly

identify candidate immunogens towards the elicitation of antibodies containing key acquired

mutations *in vivo*. With this approach we identified Envs hypothesized to select for DH270.6-like

bnAbs that contain all but one of the acquired improbable mutations in DH270min11 (Figure 6).

In order to more robustly select for the improbable acquired mutations M51$_{HC}$, Y27$_{LC}$ and L48$_{LC}$,

additional boosting immunogens may be required. Based on one available data set (Saunders

et al., 2019), DH270min11 residues selected from the antibody libraries by the 10.17DT

immunogen correlated well with amino acids present at the same positions in antibodies elicited

by 10.17DT immunizations. Nevertheless, elicitation of antibodies *in vivo* depends on many

factors beyond binding affinity that are not captured in our *in vitro* characterization. For example,

precursor frequency and their ability to be recruited into germinal centers by different

immunizations together with the stimulation of T helper cells will be essential for the success of

this vaccination regimen. (Abbott et al., 2018; Lee et al., 2020; Mesin et al., 2020). Therefore,

the envelope discovery and characterization approach presented here is primarily meant to

provide a rational way to classify and prioritize candidate immunogens for *in vivo* testing. The

potential of the vaccination regimen identified here to induce DH270.6-like bnAbs that contain

the key target mutations *in vivo* will be assessed in animal models in upcoming work.

HIV vaccine boosting regimens have typically been determined empirically and usually

involve serial immunizations with molecules that progressively resemble the native HIV Env

(Dosenovic et al., 2015a; Escolano et al., 2016; Sanders et al., 2015; Steichen et al., 2016; Tian

et al., 2016). With the exception of one study that started with a later stage bnAb precursor and

used sera screening and B cell repertoire analysis to inform immunogen choice (Escolano et al.,

2016), such boosting regimens had limited success in maturing activated bnAb precursors

(Briney et al., 2016; Dosenovic et al., 2015a). Our approach aims to rationally identify and

validate Envs that specifically bind to bnAb antibody lineage members bearing functional

improbable mutations. Similar library platforms to the one described here for DH270.6 can be

developed for other HIV bnAbs, in order to identify sequential Envs that target bnAb lineages

that contain rare mutations necessary for HIV neutralization breadth. In conclusion, this study

describes a rational and efficient method to determine the key amino acids required for antibody

function, and establishes a generalizable approach to identify candidate envelope immunogens

for vaccine studies.

Methods

Computational identification of functionally critical acquired mutations.
Computational design of minimally mutated DH270.6 antibodies were based on the crystal
structure of the DH270.6 scFv in complex with Man9-V3 glycopeptide (PDB: 6CBP), a
previously described synthetic peptide that mimics the glycan-V3 epitope on Env (Alam et al.,
2017; Fera et al., 2018). Based on structural analysis, different subsets of amino acids were
identified that were deemed not important for DH270.6 stability or binding. The amino acids at
these sites were changed to their UCA identity and the Rosetta energy of the new antibodies
was computed. Upon reversion to UCA, 5000 Rosetta models were generated for each
"minimized" antibody decoys by sampling all amino acid side chains and allowing fine backbone
movement (Supplementary Figure 1) (Lauck et al., 2010). Antibodies with decoys that sampled
Rosetta free energy levels on par with those of the mature DH270.6 were selected for
experimental characterization (Supplementary Figure 1).

Plasmids and DNA synthesis. DNA for yeast display libraries was synthesized as a
pooled oligo library by BioXp (CodexDNA). Genes encoding the antibody heavy and light chains
were commercially synthesized and cloned into pcDNA3.1 vector (GenScript). DNA primers for
sequencing and insert amplification were ordered from IDT.

Development and screening on scFv libraries on the surface of yeast. *Library*
design and synthesis. DH270min1 sites with amino acids maintained from DH270.6 (heavy
chain: G31D, Y32F, I51M, N54Q, S55T, G57R, R98T, G110Y; light chain: S26K, S27Y, Y32H,
L48Y, Y60T, Y93F, A94G, S97A, F101C) were allowed to sample either the mature DH270.6 or
the germline DH270UCA3 residue in an all against all fashion (library size of 1.31×10^5 scFv
variants). Library DNA was synthesized as a pooled oligo library by BioXp and amplified with
High Fidelity Phusion polymerase (New England Biolabs). PCR products were gel extracted

(Qiagen Gel Extraction kit) to select full length genes that were further purified (Qiagen PCR

cleanup kit) as per the manufacturer's protocol. The three libraries that aimed to identify

alternative amino acids at the functional sites in DH270min11 maintained from DH270.6, tested

all the possible amino acids combinations as follows: V3 loop library (heavy chain: M51, T55,

R57; light chain: A97); 301 glycan library (heavy chain: Y110; light chain: Y27, G94, C101); 332

glycan library (heavy chain: D31, F32, T98; light chain: Y48).

Library transformation into S. Cerevisae. Libraries were generated to display DH270.6

scFv variants on the surface of yeast as previously described (Benatuil et al., 2010; Chao et al.,

2006). *S. cerevisiae* EBY100 cells were transformed by electroporation with a 3:1 ratio of 12µg

scFv library DNA and 4µg pCTCON2 plasmid digested with BamHI, SalI, NheI (New England

Biolabs). The typical sizes of the transformed libraries, determined by serial dilution on selective

plates, ranged from $1-5\times10^7$. Around 70-90% of the sequences recovered from the transformed

libraries were confirmed to contain full length, in-frame genes by Sanger sequencing. Yeast

Libraries were grown in SDCAA media (Teknova) with Pen-Strep at 30C and 225 rpm.

Library screening by FACS. scFv expression on the surface of yeast was induced by

culturing the libraries in SGCAA (Teknova) media at a density of 1x10^7 cells/mL for 24-36

hours. Cells were washed twice in ice cold PBSA (0.01M sodium phosphate, pH 7.4, 0.137M

sodium chloride, 1g/L bovine serum albumin) and labeled with biotinylated SOSIP Envs (10.17

and 10.17DT) at a concentration of 300nM and incubated for 1hr at 4C. Cells were then washed

twice with PBSA and resuspended in secondary labeling reagent 1:100 α c-myc:FITC (ICL) and

1:20 streptavidin:R-PE (Sigma-Aldrich) and incubated at 4C for 30 minutes. Cells were washed

twice with PBSA after incubation with the fluorescently labeled probes and sorted on a BD

FACS-DiVa. Double positive cells for PE and FITC were collected and expanded for one week

in SDCAA media supplemented with pen-strep before successive rounds of enrichment. Sorts

of the libraries with non-biotinylated SOSIPs (358.06, 1432.41, 836.31 and 0526.02) used 4-fold

excess PGT151 mAb and 1:60 α-hIgG:PE (Invitrogen) pre-incubated for 60 minutes as the

phycoerythrin signal. The *DH270min1 library* was screened separately for binding of SOSIPs

10.17DT, 10.17, 1432.41, 358.06, 526.02, 836.31. The *DH270min11 N301, N332, and V3 loop*

libraries were sorted sequentially with SOSIPS 1017.DT, 358.06, 1432.41 twice, and 052.02.

FACS data was analyzed with Flowjo_v10.6 software (Becton, Dickinson & Company). All

clones by FACS were expanded, and their DNA was extracted (Zymo Research) for analysis by

Next Generation Sequencing and Sanger sequencing analysis (Genewiz).

Sequence analysis of library clones. scFv encoding plasmids were recovered from

yeast cultures by yeast miniprep with the Zymoprep yeast plasmid miniprep II kit (Zymo

Research). Isolated DNA was transformed into NEB5α strain of *E. coli* (New England Biolabs)

and the DNA of individual bacterial colonies was isolated (Wizard Plus SV Minipreps, Promega)

and analyzed by Sanger sequencing to confirm the validity of the FACS selections. To prepare

plasmids for Next Generation Sequencing, the scFv containing region from isolated plasmids

was amplified by PCR using Q5 high fidelity PCR (NEB). Illumina NGS samples were prepped

and run using the Illumina MiSeq v3 reagent kit following manufacturer's protocols pooling four

library samples per lane. Illumina sequencing returned an average of 14.3 million reads per

sample, of which an average of 13.1 million mapped to the scFv amplicon.

PacBio long read NGS was completed by Genewiz pooling 3-4 library samples per SMRT cell.

Sequencing results by PacBio returned 35-47k reads per sample sequenced, of which 15-26k

aligned to the scFv amplicon. Sequencing data was processed using Geneious Prime and

programs developed in-house to compute the amino acid frequency and distribution.

Antibody expression and purification. Antibodies were expressed and purified as

previously described (Saunders et al., 2019). Briefly, 100mL cultures of Expi293F cells at a

density of 2.5×10^6 cells/mL were transiently transfected with 50µg each heavy and light chain

encoding plasmids and Expifectamine (Invitrogen) per manufacturer's protocol. Five days after

transfection, cell culture media was cleared of cells by centrifugation, and the surpernatant was

filtered with 0.8 micron filters. Clarified supernatant was incubated with Protein A beads

(ThermoFisher) over night at 4C, washed with 20mM Tris supplemented with 350mM NaCl

(pH=7), followed by elution with a 2.5% Glacial Acetic Acid Elution Buffer and subsequent buffer

exchange into 25mM Citric Acid supplemented with125mM NaCl (pH=6). IgGs expression was

confirmed by reducing SDS-PAGE analysis, and quantified by measuring absorbance at 280nM

(Nanodrop 2000)

Recombinant HIV Env production. SOSIP envelopes were produced recombinantly as

previously described (Saunders et al., 2019). Briefly, Freestyle 293 cells were transfected with

293Fectin complexed with envelope-expressing DNA and furin-expressing plasmid DNA. After 6

days, SOSIPS were purified via PGT145 affinity chromatography with subsequent size

exclusion chromatography. Trimeric HIV-1 Env fractions were pooled, snap-frozen, and stored

at -80C in 10mM Tris pH 8, 500mM NaCl buffer. Expression of gp120 proteins occurred as

previously described in 293F cells by transient transfection with 293Fectin (Invitrogen) and

purification by metal affinity and Size Exclusion Chromatography (Alam et al., 2013; Liao et

al., 2013a; Liao et al., 2006)

Pseudovirus neutralization assay. Neutralization of antibodies was measured in TZM-

bl cells in a 96 or 384-well plate format assay using either a 20-virus panel comprised of viruses

known to be sensitive to DH270.6, or on a 208 Env-pseudovirus panel representative of the

major genetic subtypes of circulating virus (Bonsignori et al., 2017a; Bonsignori et al., 2016).

Signal was calculated as a reduction in luminescence compared to control wells and reported as

IC50 in ug/mL.

ELISA Assay. Direct-binding ELISAs were performed as described on 96 gp120

molecules isolated previously (Bonsignori et al., 2017a; Bonsignori et al., 2016). 384-well plates

were blocked overnight at 4C. Starting concentrations of 100ug/mL purified antibody were

serially diluted 3-fold and incubated at room temperature for 1 hour. 1:30,000 HRP- conjugated

hIgG in assay diluent was added to plates, incubated for one hour, and developed using TMB

substrate. Plates were read at 450nm on a SpectraMax 384 PLUS reader (Molecular Devices).

(LogAUC).

Acknowledgements. We thank J. Baalwa, D. Ellenberger, F. Gao, B. Hahn, K. Hong, J. Kim, F.

McCutchan, D. Montefiori, L. Morris, E. Sanders-Buell, G. Shaw, R. Swanstrom, M. Thomson,

S. Tovanabutra, C. Williamson, and L. Zhang for contributing the HIV-1 envelope plasmids used

in the 208-strain panel. We thank K. McKee, C. Moore, S. O'Dell, G. Padilla, S.D. Schmidt, C.

Whittaker, and A.B. McDermott for assistance with neutralization assessments on the 208-strain

panel.

References

Abbott, R.K., Lee, J.H., Menis, S., Skog, P., Rossi, M., Ota, T., Kulp, D.W., Bhullar, D., Kalyuzhniy, O., Havenar-Daughton, C.*, et al.* (2018). Precursor Frequency and Affinity Determine B Cell Competitive Fitness in Germinal Centers, Tested with Germline-Targeting HIV Vaccine Immunogens. Immunity *48*, 133-146.e136.

Alam, S.M., Aussedat, B., Vohra, Y., Meyerhoff, R.R., Cale, E.M., Walkowicz, W.E., Radakovich, N.A., Anasti, K., Armand, L., Parks, R.*, et al.* (2017). Mimicry of an HIV broadly neutralizing antibody epitope with a synthetic glycopeptide. Science Translational Medicine *9*, eaai7521.

Alam, S.M., Liao, H.-X., Tomaras, G.D., Bonsignori, M., Tsao, C.-Y., Hwang, K.-K., Chen, H., Lloyd, K.E., Bowman, C., Sutherland, L.*, et al.* (2013). Antigenicity and Immunogenicity of RV144 Vaccine AIDSVAX Clade E Envelope Immunogen Is Enhanced by a gp120 N-Terminal Deletion. Journal of Virology *87*, 1554-1568.

Barnes, C.O., Gristick, H.B., Freund, N.T., Escolano, A., Lyubimov, A.Y., Hartweger, H., West, A.P., Cohen, A.E., Nussenzweig, M.C., and Bjorkman, P.J. (2018). Structural characterization of a highly-potent V3-glycan broadly neutralizing antibody bound to natively-glycosylated HIV-1 envelope. Nature Communications *9*, 1251.

Benatuil, L., Perez, J.M., Belk, J., and Hsieh, C.M. (2010). An improved yeast transformation method for the generation of very large human antibody libraries. Protein Eng Des Sel *23*, 155-159.

Bender, B.J., Cisneros, A., Duran, A.M., Finn, J.A., Fu, D., Lokits, A.D., Mueller, B.K., Sangha, A.K., Sauer, M.F., Sevy, A.M.*, et al.* (2016). Protocols for Molecular Modeling with Rosetta3 and RosettaScripts. Biochemistry *55*, 4748-4763.

Bonsignori, M., Kreider, E.F., Fera, D., Meyerhoff, R.R., Bradley, T., Wiehe, K., Alam, S.M., Aussedat, B., Walkowicz, W.E., Hwang, K.K.*, et al.* (2017a). Staged induction of HIV-1 glycan-dependent broadly neutralizing antibodies. Sci Transl Med *9*.

Bonsignori, M., Liao, H.X., Gao, F., Williams, W.B., Alam, S.M., Montefiori, D.C., and Haynes, B.F. (2017b). Antibody-virus co-evolution in HIV infection: paths for HIV vaccine development. Immunol Rev *275*, 145-160.

Bonsignori, M., Scott, E., Wiehe, K., Easterhoff, D., Alam, S.M., Hwang, K.K., Cooper, M., Xia, S.M., Zhang, R., Montefiori, D.C.*, et al.* (2018). Inference of the HIV-1 VRC01 Antibody Lineage Unmutated Common Ancestor Reveals Alternative Pathways to Overcome a Key Glycan Barrier. Immunity *49*, 1162-1174.e1168.

Bonsignori, M., Zhou, T., Sheng, Z., Chen, L., Gao, F., Joyce, M.G., Ozorowski, G., Chuang, G.-Y., Schramm, C.A., Wiehe, K.*, et al.* (2016). Maturation Pathway from Germline to Broad HIV-1 Neutralizer of a CD4-Mimic Antibody. Cell *165*, 449-463.

Briney, B., Sok, D., Jardine, J.G., Kulp, D.W., Skog, P., Menis, S., Jacak, R., Kalyuzhniy, O., de Val, N., Sesterhenn, F.*, et al.* (2016). Tailored Immunogens Direct Affinity Maturation toward HIV Neutralizing Antibodies. Cell *166*, 1459-1470.e1411.

Cao, L., Diedrich, J.K., Kulp, D.W., Pauthner, M., He, L., Park, S.-K.R., Sok, D., Su, C.Y., Delahunty, C.M., Menis, S.*, et al.* (2017). Global site-specific N-glycosylation analysis of HIV envelope glycoprotein. Nature Communications *8*, 14954.

Chao, G., Lau, W.L., Hackel, B.J., Sazinsky, S.L., Lippow, S.M., and Wittrup, K.D. (2006). Isolating and engineering human antibodies using yeast surface display. Nature Protocols *1*, 755-768.

Das, R., and Baker, D. (2008). Macromolecular Modeling with Rosetta. Annual Review of Biochemistry *77*, 363-382.

Doria-Rose, N.A., Klein, R.M., Daniels, M.G., Dell, S., Nason, M., Lapedes, A., Bhattacharya, T., Migueles, S.A., Wyatt, R.T., Korber, B.T.*, et al.* (2010). Breadth of Human Immunodeficiency Virus-Specific

Neutralizing Activity in Sera: Clustering Analysis and Association with Clinical Variables. Journal of
Virology *84*, 1631.
Doria-Rose, N.A., and Landais, E. (2019). Coevolution of HIV-1 and broadly neutralizing antibodies. Curr
Opin HIV AIDS *14*, 286-293.
Dosenovic, P., von Boehmer, L., Escolano, A., Jardine, J., Freund, N.T., Gitlin, A.D., McGuire, A.T., Kulp,
D.W., Oliveira, T., Scharf, L*., et al.* (2015a). Immunization for HIV-1 Broadly Neutralizing Antibodies in
Human Ig Knockin Mice. Cell *161*, 1505-1515.
Dosenovic, P., von Boehmer, L., Escolano, A., Jardine, J., Freund, Natalia T., Gitlin, Alexander D.,
McGuire, Andrew T., Kulp, Daniel W., Oliveira, T., Scharf, L*., et al.* (2015b). Immunization for HIV-1
Broadly Neutralizing Antibodies in Human Ig Knockin Mice. Cell *161*, 1505-1515.
Escolano, A., Steichen, J.M., Dosenovic, P., Kulp, D.W., Golijanin, J., Sok, D., Freund, N.T., Gitlin, A.D.,
Oliveira, T., Araki, T*., et al.* (2016). Sequential Immunization Elicits Broadly Neutralizing Anti-HIV-1
Antibodies in Ig Knockin Mice. Cell *166*, 1445-1458.e1412.
Fera, D., Lee, M.S., Wiehe, K., Meyerhoff, R.R., Piai, A., Bonsignori, M., Aussedat, B., Walkowicz, W.E.,
Ton, T., Zhou, J.O*., et al.* (2018). HIV envelope V3 region mimic embodies key features of a broadly
neutralizing antibody lineage epitope. Nature Communications *9*, 1111.
Freund, N.T., Wang, H., Scharf, L., Nogueira, L., Horwitz, J.A., Bar-On, Y., Golijanin, J., Sievers, S.A., Sok,
D., Cai, H*., et al.* (2017). Coexistence of potent HIV-1 broadly neutralizing antibodies and antibody-
sensitive viruses in a viremic controller. Sci Transl Med *9*.
Gao, F., Bonsignori, M., Liao, H.X., Kumar, A., Xia, S.M., Lu, X., Cai, F., Hwang, K.K., Song, H., Zhou, T*., et
al.* (2014). Cooperation of B cell lineages in induction of HIV-1-broadly neutralizing antibodies. Cell *158*,
481-491.
Georgiev, I.S., Rudicell, R.S., Saunders, K.O., Shi, W., Kirys, T., McKee, K., O'Dell, S., Chuang, G.-Y., Yang,
Z.-Y., Ofek, G*., et al.* (2014). Antibodies VRC01 and 10E8 Neutralize HIV-1 with High Breadth and Potency
Even with Ig-Framework Regions Substantially Reverted to Germline. The Journal of Immunology *192*,
1100.
Gray, E.S., Taylor, N., Wycuff, D., Moore, P.L., Tomaras, G.D., Wibmer, C.K., Puren, A., DeCamp, A.,
Gilbert, P.B., Wood, B*., et al.* (2009). Antibody Specificities Associated with Neutralization Breadth in
Plasma from Human Immunodeficiency Virus Type 1 Subtype C-Infected Blood Donors. Journal of
Virology *83*, 8925.
Havenar-Daughton, C., Sarkar, A., Kulp, D.W., Toy, L., Hu, X., Deresa, I., Kalyuzhniy, O., Kaushik, K.,
Upadhyay, A.A., Menis, S*., et al.* (2018). The human naive B cell repertoire contains distinct subclasses
for a germline-targeting HIV-1 vaccine immunogen. Science translational medicine *10*, eaat0381.
Haynes, B.F., Burton, D.R., and Mascola, J.R. (2019). Multiple roles for HIV broadly neutralizing
antibodies. Science Translational Medicine *11*, eaaz2686.
Haynes, B.F., Kelsoe, G., Harrison, S.C., and Kepler, T.B. (2012). B-cell-lineage immunogen design in
vaccine development with HIV-1 as a case study. Nat Biotechnol *30*, 423-433.
Haynes, B.F., Shaw, G.M., Korber, B., Kelsoe, G., Sodroski, J., Hahn, B.H., Borrow, P., and McMichael, A.J.
(2016). HIV-Host Interactions: Implications for Vaccine Design. Cell Host Microbe *19*, 292-303.
Hraber, P., Seaman, M.S., Bailer, R.T., Mascola, J.R., Montefiori, D.C., and Korber, B.T. (2014). Prevalence
of broadly neutralizing antibody responses during chronic HIV-1 infection. Aids *28*, 163-169.
Hwang, J.K., Wang, C., Du, Z., Meyers, R.M., Kepler, T.B., Neuberg, D., Kwong, P.D., Mascola, J.R., Joyce,
M.G., Bonsignori, M*., et al.* (2017). Sequence intrinsic somatic mutation mechanisms contribute to
affinity maturation of VRC01-class HIV-1 broadly neutralizing antibodies. Proc Natl Acad Sci U S A *114*,
8614-8619.
Jardine, J.G., Sok, D., Julien, J.P., Briney, B., Sarkar, A., Liang, C.H., Scherer, E.A., Henry Dunand, C.J.,
Adachi, Y., Diwanji, D*., et al.* (2016). Minimally Mutated HIV-1 Broadly Neutralizing Antibodies to Guide
Reductionist Vaccine Design. PLoS Pathog *12*, e1005815.

Julien, J.-P., Sok, D., Khayat, R., Lee, J.H., Doores, K.J., Walker, L.M., Ramos, A., Diwanji, D.C., Pejchal, R.,
Cupo, A*., et al.* (2013). Broadly neutralizing antibody PGT121 allosterically modulates CD4 binding via
recognition of the HIV-1 gp120 V3 base and multiple surrounding glycans. PLoS pathogens *9*, e1003342-
e1003342.
Kepler, T.B., Liao, H.-X., Alam, S.M., Bhaskarabhatla, R., Zhang, R., Yandava, C., Stewart, S., Anasti, K.,
Kelsoe, G., Parks, R*., et al.* (2014). Immunoglobulin gene insertions and deletions in the affinity
maturation of HIV-1 broadly reactive neutralizing antibodies. Cell host & microbe *16*, 304-313.
Klein, F., Diskin, R., Scheid, Johannes F., Gaebler, C., Mouquet, H., Georgiev, I.S., Pancera, M., Zhou, T.,
Incesu, R.-B., Fu, Brooks Z*., et al.* (2013). Somatic Mutations of the Immunoglobulin Framework Are
Generally Required for Broad and Potent HIV-1 Neutralization. Cell *153*, 126-138.
Kwong, P.D., and Mascola, J.R. (2018). HIV-1 Vaccines Based on Antibody Identification, B Cell Ontogeny,
and Epitope Structure. Immunity *48*, 855-871.
LaBranche, C.C., Henderson, R., Hsu, A., Behrens, S., Chen, X., Zhou, T., Wiehe, K., Saunders, K.O., Alam,
S.M., Bonsignori, M*., et al.* (2019). Neutralization-guided design of HIV-1 envelope trimers with high
affinity for the unmutated common ancestor of CH235 lineage CD4bs broadly neutralizing antibodies.
PLOS Pathogens *15*, e1008026.
Landais, E., and Moore, P.L. (2018). Development of broadly neutralizing antibodies in HIV-1 infected
elite neutralizers. Retrovirology *15*, 61.
Lauck, F., Smith, C.A., Friedland, G.F., Humphris, E.L., and Kortemme, T. (2010). RosettaBackrub—a web
server for flexible backbone protein structure modeling and design. Nucleic Acids Research *38*, W569-
W575.
Leaver-Fay, A., Tyka, M., Lewis, S.M., Lange, O.F., Thompson, J., Jacak, R., Kaufman, K., Renfrew, P.D.,
Smith, C.A., Sheffler, W*., et al.* (2011). ROSETTA3: an object-oriented software suite for the simulation
and design of macromolecules. Methods Enzymol *487*, 545-574.
Lee, J.H., Nakao, C., Moyer, T.J., Irvine, D.J., Schief, W.R., and Crotty, S. (2020). Exploration of roles of
CD4 T cell help in recruitment of rare B cells to germinal centers. The Journal of Immunology *204*,
167.165.
Liao, H.-X., Bonsignori, M., Alam, S.M., McLellan, Jason S., Tomaras, Georgia D., Moody, M.A., Kozink,
Daniel M., Hwang, K.-K., Chen, X., Tsao, C.-Y*., et al.* (2013a). Vaccine Induction of Antibodies against a
Structurally Heterogeneous Site of Immune Pressure within HIV-1 Envelope Protein Variable Regions 1
and 2. Immunity *38*, 176-186.
Liao, H.-X., Sutherland, L.L., Xia, S.-M., Brock, M.E., Scearce, R.M., Vanleeuwen, S., Alam, S.M.,
McAdams, M., Weaver, E.A., Camacho, Z.T*., et al.* (2006). A group M consensus envelope glycoprotein
induces antibodies that neutralize subsets of subtype B and C HIV-1 primary viruses. Virology *353*, 268-
282.
Liao, H.X., Lynch, R., Zhou, T., Gao, F., Alam, S.M., Boyd, S.D., Fire, A.Z., Roskin, K.M., Schramm, C.A.,
Zhang, Z*., et al.* (2013b). Co-evolution of a broadly neutralizing HIV-1 antibody and founder virus. Nature
496, 469-476.
Mesin, L., Schiepers, A., Ersching, J., Barbulescu, A., Cavazzoni, C.B., Angelini, A., Okada, T., Kurosaki, T.,
and Victora, G.D. (2020). Restricted Clonality and Limited Germinal Center Reentry Characterize Memory
B Cell Reactivation by Boosting. Cell *180*, 92-106.e111.
Mouquet, H., Scharf, L., Euler, Z., Liu, Y., Eden, C., Scheid, J.F., Halper-Stromberg, A., Gnanapragasam,
P.N.P., Spencer, D.I.R., Seaman, M.S*., et al.* (2012). Complex-type N-glycan recognition by potent broadly
neutralizing HIV antibodies. Proceedings of the National Academy of Sciences of the United States of
America *109*, E3268-E3277.
Sanders, R.W., van Gils, M.J., Derking, R., Sok, D., Ketas, T.J., Burger, J.A., Ozorowski, G., Cupo, A.,
Simonich, C., Goo, L*., et al.* (2015). HIV-1 neutralizing antibodies induced by native-like envelope trimers.
Science *349*, aac4223.

Saunders, K.O., Wiehe, K., Tian, M., Acharya, P., Bradley, T., Alam, S.M., Go, E.P., Scearce, R., Sutherland,
L., Henderson, R.*, et al.* (2019). Targeted selection of HIV-specific antibody mutations by engineering B
cell maturation. Science *366*.
Shen, C.H., DeKosky, B.J., Guo, Y., Xu, K., Gu, Y., Kilam, D., Ko, S.H., Kong, R., Liu, K., Louder, M.K.*, et al.*
(2020). VRC34-Antibody Lineage Development Reveals How a Required Rare Mutation Shapes the
Maturation of a Broad HIV-Neutralizing Lineage. Cell Host Microbe *27*, 531-543 e536.
Simonich, C.A., Williams, K.L., Verkerke, H.P., Williams, J.A., Nduati, R., Lee, K.K., and Overbaugh, J.
(2016). HIV-1 Neutralizing Antibodies with Limited Hypermutation from an Infant. Cell *166*, 77-87.
Sok, D., Laserson, U., Laserson, J., Liu, Y., Vigneault, F., Julien, J.P., Briney, B., Ramos, A., Saye, K.F., Le, K.*,*
et al. (2013). The effects of somatic hypermutation on neutralization and binding in the PGT121 family
of broadly neutralizing HIV antibodies. PLoS Pathog *9*, e1003754.
Steichen, J.M., Kulp, D.W., Tokatlian, T., Escolano, A., Dosenovic, P., Stanfield, R.L., McCoy, L.E.,
Ozorowski, G., Hu, X., Kalyuzhniy, O.*, et al.* (2016). HIV Vaccine Design to Target Germline Precursors of
Glycan-Dependent Broadly Neutralizing Antibodies. Immunity *45*, 483-496.
Steichen, J.M., Lin, Y.C., Havenar-Daughton, C., Pecetta, S., Ozorowski, G., Willis, J.R., Toy, L., Sok, D.,
Liguori, A., Kratochvil, S.*, et al.* (2019). A generalized HIV vaccine design strategy for priming of broadly
neutralizing antibody responses. Science *366*.
Struwe, W.B., Chertova, E., Allen, J.D., Seabright, G.E., Watanabe, Y., Harvey, D.J., Medina-Ramirez, M.,
Roser, J.D., Smith, R., Westcott, D.*, et al.* (2018). Site-Specific Glycosylation of Virion-Derived HIV-1 Env Is
Mimicked by a Soluble Trimeric Immunogen. Cell Rep *24*, 1958-1966.e1955.
Tian, M., Cheng, C., Chen, X., Duan, H., Cheng, H.-L., Dao, M., Sheng, Z., Kimble, M., Wang, L., Lin, S.*, et*
al. (2016). Induction of HIV Neutralizing Antibody Lineages in Mice with Diverse Precursor Repertoires.
Cell *166*, 1471-1484.e1418.
Walker, L.M., Huber, M., Doores, K.J., Falkowska, E., Pejchal, R., Julien, J.P., Wang, S.K., Ramos, A., Chan-
Hui, P.Y., Moyle, M.*, et al.* (2011). Broad neutralization coverage of HIV by multiple highly potent
antibodies. Nature *477*, 466-470.
Wiehe, K., Bradley, T., Meyerhoff, R.R., Hart, C., Williams, W.B., Easterhoff, D., Faison, W.J., Kepler, T.B.,
Saunders, K.O., Alam, S.M.*, et al.* (2018). Functional Relevance of Improbable Antibody Mutations for
HIV Broadly Neutralizing Antibody Development. Cell Host Microbe *23*, 759-765 e756.
Wu, X., Yang, Z.Y., Li, Y., Hogerkorp, C.M., Schief, W.R., Seaman, M.S., Zhou, T., Schmidt, S.D., Wu, L., Xu,
L.*, et al.* (2010). Rational design of envelope identifies broadly neutralizing human monoclonal
antibodies to HIV-1. Science *329*, 856-861.
Wu, X., Zhang, Z., Schramm, C.A., Joyce, M.G., Kwon, Y.D., Zhou, T., Sheng, Z., Zhang, B., O'Dell, S.,
McKee, K.*, et al.* (2015). Maturation and Diversity of the VRC01-Antibody Lineage over 15 Years of
Chronic HIV-1 Infection. Cell *161*, 470-485.
Yaari, G., Vander Heiden, J.A., Uduman, M., Gadala-Maria, D., Gupta, N., Stern, J.N.H., O'Connor, K.C.,
Hafler, D.A., Laserson, U., Vigneault, F.*, et al.* (2013). Models of somatic hypermutation targeting and
substitution based on synonymous mutations from high-throughput immunoglobulin sequencing data.
Front Immunol *4*, 358-358.

Figure 1. Design of first generation minimized DH270.6 antibodies. A. Structural mapping of DH270.6 somatic mutations maintained in different DH270min variants. Mutations (*magenta*, sticks) were mapped on the structure of DH270.6 scFv (*grey*) in complex with an HIV Env derived V3-loop glycopeptide (*cyan*) (PDBID: 6cbp). Glycans at positions N301 and N332 are shown in sticks. **B.** Venn diagram of the acquired mutations present in the DH270min antibodies colored by their predicted *in vivo* probability (high=*blue*; low=*red)*. **C.** Neutralization breadth and potency of DH270.6, DH270min antibodies, and DH270.6 variants that contain only the "probable" (DH270Prob) or "improbable" mutations (DH270Improb) on a panel of DH270.6 sensitive pseudoviruses. #AA Muts=the number of DH270.6 acquired mutations maintained in the respective antibodies.

Figure 2. Additional minimization of DH270min1 by single site mutagenesis.
A. The effect on neutralization of every DH270min1 acquired mutation assessed by single site mutagenesis to the corresponding UCA residue. Second generation minimized antibodies DH270min11 and DH270min12 contained only the acquired mutations important for DH270min1 neutralization. **B.** Neutralization breadth of minDH270 antibodies on a global panel of 208 pseudoviruses. **C.** Location and binding interactions of the DH270.6 acquired mutations (*magenta*, sticks) maintained in DH270min11.

A

B

Figure 3. Identification of immunogen candidates that preferentially interact with DH270min1 through its acquired mutations. A. The number of tested gp120 immunogen candidates that preferentially interact with each of the acquired mutations in DH270min1. Only the acquired mutations present in DH270min11 are displayed. The top 50% gp120 molecules by DH270min1 affinity are shown. **B.** The selection strength of individual candidate immunogens for each of the acquired mutations present in DH270min11. NB=not binding.

Figure 4. In vitro characterization of candidate immunogens towards selection of DH270.6-derived antibodies that contain the acquired mutations present in DH270min1.
A. Design of a scFv library that sampled in combination each acquired mutation site in DH270min1. At each sampled position the amino acid present in the mature DH270.6 mAb (*blue*) or that found in the UCA (*red*) was randomly assorted. **B.** Selection of the scFv library for binding to different candidate immunogens by FACS. **C.** Enrichment levels of the acquired mutations in the clones isolated by the different immunogen candidates in **B**, computed relative to the frequency of the mutations in the unsorted library. Only the key acquired mutations present in DH270min11 are shown. **D.** The enrichment of mutations by vaccination, averaged from 5 DH270UCA knock-in mice immunized with 10.17DT (x-axis), was plotted against the enrichment of the same mutations in library clones selected by 10.17DT from **C** (y-axis). Data was fitted with a linear equation (R^2=0.625). Immunization enrichment was determined by calculating the ratio between the frequency of the observed mutations in the antibody repertoire of vaccinated animals analyzed by next generation sequencing, and the expected frequency of the mutations in the absence of immunogen selection predicted with ARMADiLLO (Wiehe et al., 2018). Immunization data is available only for the heavy chain sequences (Saunders et al., 2019).

Figure 5. Identification of alternative amino acids at the acquired mutation sites in DH270min11. Structural mapping of the DH270min11 residues simultaneously sampled in the three libraries (*top*). Sequence diversity in the antibody library variants selected by immunogen candidates 10.17DT_SOSIP, 1432.41_SOSIP, and 526.02_SOSIP. Logo plots display the amino acid diversity present at each library site in the selected clones, with the mature DH270.6 amino acid shown in *blue*. Mutations more probable than the DH270.6 amino acid are marked in *green*. Sequence tables show the frequency of the 5 most common co-occurring DH270min11 variants selected by the respective immunogen candidates.

Figure 6. Vaccination regimen for the elicitation of DH270.6-like antibodies that contain the key functional mutations. Priming with the germline targeting immunogen 10.17DT will be followed by successive boosts with molecules identified in this study. Arrows point to the DH270min11 acquired mutations predicted to be selected by the immunogen depicted in the same color. Mutations are colored by probability (*red*=improbable; *blue*=probable).

Supplementary Figures

A

B

Rosetta scores of minDH270 models

min1   UCA+ binding interface + 2nd shell improbable mutants
min2   UCA+ binding interface + 2nd shell improbable mutants + H/L chain interface muts
min3   UCA+ binding interface muts

Supplementary Figure 1. Computational design of minimized DH270.6 antibodies.
A. Rosetta protocol to evaluate the effect of acquired mutations on antibody packing and stability. **B.** Rosetta energy landscapes of designed mins (5000 decoys) described in **A** and chosen for experimental characterization.

High Throughput Antibody Screen - Panel 38

Assay - Luc/TZM-bl
values represent IC in ug/ml

Legend: <0.001 | .001-.01 | .01-.100 | .100-1.00 | 1.00-10.0 | >10.0

IC50 — Data from Bonsignori

Virus ID	Clade	Exp. Code	DH270.min1	DH270.min11	DH270.min12	DH270	DH270_IA1	DH270_IA2	DH270_IA4
0260.v5.c36	A	CVL2139_001-05	0.351	0.433	1.37	2.32	0.295	2.17	>50
0330.v4.c3	A	CVL2139_001-06	>50	>50	>50	>50	>50	>50	>50
0439.v5.c1	A	CVL2139_001-07	>50	>50	>50	>50	>50	>50	>50
3365.v2.c20	A	CVL2139_003-07	>50	>50	>50	>50	>50	>50	>50
3415.v1.c1	A	CVL2139_003-08	>50	>50	>50	>50	>50	>50	>50
3718.v3.c11	A	CVL2139_003-12	>50	>50	>50	>50	>50	>50	>50
398-F1.F6.20	A	CVL2139_004-03	0.016	0.015	0.030	0.038	0.010	0.026	1.14
BB201.B42	A	CVL2139_006-05	>50	>50	>50	>50	>50	>50	>50
BB539.2B13	A	CVL2139_006-06	>50	>50	>50	>50	>50	>50	>50
BG505.W6M.C2	A	CVL2139_006-08	>50	>50	>50	>50	>50	>50	>50
BI369.9A	A	CVL2139_006-09	0.103	0.111	0.325	0.407	0.086	0.579	>50
BS208.B1	A	CVL2139_007-05	>50	>50	>50	>50	>50	>50	>50
KER2008.12	A	CVL2139_011-04	0.076	0.214	>50	14.6	0.080	>50	>50
KER2018.11	A	CVL2139_011-05	>50	>50	>50	>50	>50	>50	>50
KNH1209.18	A	CVL2139_011-06	0.021	0.025	0.026	0.085	0.032	0.082	2.74
MB201.A1	A	CVL2139_011-08	>50	>50	>50	>50	>50	>50	>50
MB539.2B7	A	CVL2139_011-09	>50	>50	>50	>50	>50	>50	>50
MI369.A5	A	CVL2139_011-10	0.129	0.196	0.146	0.501	0.221	0.620	>50
MS208.A1	A	CVL2139_011-12	>50	>50	>50	>50	>50	>50	>50
Q23.17	A	CVL2139_012-06	0.020	0.024	0.024	0.058	0.022	0.074	16.3
Q259.17	A	CVL2139_012-07	>50	>50	>50	>50	>50	>50	>50
Q769.d22	A	CVL2139_012-09	>50	>50	>50	>50	>50	>50	>50
Q769.h5	A	CVL2139_012-10	>50	>50	>50	>50	>50	>50	>50
Q842.d12	A	CVL2139_012-11	>50	>50	>50	>50	>50	>50	>50
QH209.14M.A2	A	CVL2139_013-01	>50	>50	>50	>50	>50	>50	>50
RW020.2	A	CVL2139_013-07	0.012	0.006	0.007	0.051	0.010	0.023	0.176
UG037.8	A	CVL2139_015-06	0.101	0.219	>50	1.53	0.317	>50	>50
246-F3.C10.2	AC	CVL2139_002-04	>50	>50	>50	>50	>50	>50	>50
3301.v1.c24	AC	CVL2139_003-04	0.080	0.115	5.53	0.587	0.109	>50	>50
3589.v1.c4	AC	CVL2139_003-10	>50	>50	>50	>50	>50	>50	>50
6540.v4.c1	AC	CVL2139_005-02	>50	>50	>50	>50	>50	>50	>50
6545.v4.c1	AC	CVL2139_005-03	>50	>50	>50	>50	>50	>50	>50
0815.v3.c3	ACD	CVL2139_001-08	0.150	0.139	0.136	0.635	0.127	0.380	3.50
6095.v1.c10	ACD	CVL2139_004-08	>50	>50	>50	>50	>50	>50	>50
3468.v1.c12	AD	CVL2139_003-09	>50	>50	>50	>50	>50	>50	>50
Q168.a2	AD	CVL2139_012-05	>50	>50	>50	>50	>50	>50	>50
Q461.e2	AD	CVL2139_012-08	>50	>50	>50	>50	>50	>50	>50
620345.c1	AE	CVL2139_004-10	>50	>50	>50	>50	>50	>50	>50
BJOX009000.02.4	AE	CVL2139_006-11	>50	>50	>50	>50	>50	>50	>50
BJOX010000.06.2	AE	CVL2139_006-12	>50	>50	>50	>50	>50	>50	>50
BJOX025000.01.1	AE	CVL2139_006-13	>50	>50	>50	>50	>50	>50	>50
BJOX028000.10.3	AE	CVL2139_007-01	>50	>50	>50	>50	>50	>50	>50
C1080.c3	AE	CVL2139_007-07	>50	>50	>50	>50	>50	>50	>50
C2101.c1	AE	CVL2139_007-08	>50	>50	>50	>50	>50	>50	>50
C3347.c11	AE	CVL2139_007-09	>50	>50	>50	>50	>50	>50	>50
C4118.09	AE	CVL2139_007-10	>50	>50	>50	>50	>50	>50	>50
CM244.ec1	AE	CVL2139_008-10	>50	>50	>50	>50	>50	>50	>50
CNE3	AE	CVL2139_009-05	>50	>50	>50	>50	>50	>50	>50
CNE5	AE	CVL2139_009-10	>50	>50	>50	>50	>50	>50	>50
CNE55	AE	CVL2139_009-12	>50	>50	>50	>50	>50	>50	>50
CNE56	AE	CVL2139_009-13	>50	>50	>50	>50	>50	>50	>50
CNE59	AE	CVL2139_010-03	>50	>50	>50	>50	>50	>50	>50
CNE8	AE	CVL2139_010-05	>50	>50	>50	>50	>50	>50	>50
M02138	AE	CVL2139_011-07	>50	>50	>50	>50	>50	>50	>50
R1166.c1	AE	CVL2139_013-02	>50	>50	>50	>50	>50	>50	>50
R2184.c4	AE	CVL2139_013-03	>50	>50	>50	>50	>50	>50	>50
R3265.c6	AE	CVL2139_013-04	>50	>50	>50	>50	>50	>50	>50
TH023.6	AE	CVL2139_014-10	>50	>50	>50	>50	>50	>50	>50
TH966.8	AE	CVL2139_014-11	>50	>50	>50	>50	>50	>50	>50
TH976.17	AE	CVL2139_014-12	>50	>50	>50	>50	>50	>50	>50
235-47	AG	CVL2139_002-02	0.185	0.240	11.5	3.11	0.118	>50	>50
242-14	AG	CVL2139_002-03	>50	>50	>50	>50	>50	>50	>50
263-8	AG	CVL2139_002-10	1.12	5.06	>50	>50	9.03	>50	>50
269-12	AG	CVL2139_002-11	0.089	0.139	>50	2.74	0.096	>50	>50
271-11	AG	CVL2139_002-12	>50	>50	>50	>50	>50	>50	>50
928-28	AG	CVL2139_005-10	23.0	21.7	>50	>50	8.00	>50	>50
DJ263.8	AG	CVL2139_010-06	0.083	0.102	1.56	0.201	0.072	0.598	>50
T250-4	AG	CVL2139_014-01	>50	>50	>50	>50	>50	>50	>50
T251-18	AG	CVL2139_014-02	>50	>50	>50	>50	>50	>50	>50
T253-11	AG	CVL2139_014-03	>50	>50	>50	>50	>50	>50	>50
T255-34	AG	CVL2139_014-04	>50	>50	>50	>50	>50	>50	>50
T257-31	AG	CVL2139_014-05	>50	>50	>50	>50	>50	>50	>50
T266-60	AG	CVL2139_014-06	0.952	2.74	>50	6.21	0.787	>50	>50
T278-50	AG	CVL2139_014-07	>50	>50	>50	>50	>50	>50	>50
T280-5	AG	CVL2139_014-08	0.015	0.018	0.020	0.044	0.013	0.067	39.5
T33-7	AG	CVL2139_014-09	>50	>50	>50	>50	>50	>50	>50
3988.25	B	CVL2139_004-02	0.017	0.020	1.07	0.090	0.021	0.463	>50
5768.04	B	CVL2139_004-07	0.314	0.927	>50	>50	0.079	>50	>50
6101.10	B	CVL2139_004-09	0.035	0.039	0.023	0.124	0.027	0.116	1.45
6535.3	B	CVL2139_005-01	0.025	0.029	0.022	0.102	0.023	0.069	0.432
7165.18	B	CVL2139_005-08	0.056	0.068	0.710	0.281	0.043	1.11	>50
45_01dG5	B	CVL2139_004-05	0.037	0.057	0.063	0.099	0.074	0.177	16.3
89.6.DG	B	CVL2139_005-09	0.039	0.044	44.7	0.267	0.056	>50	>50
AC10.29	B	CVL2139_006-01	0.019	0.022	>50	1.18	0.012	>50	>50
ADA.DG	B	CVL2139_006-02	0.060	0.076	0.892	0.463	0.047	1.21	>50
Bal.01	B	CVL2139_006-03	0.054	0.055	0.331	0.176	0.032	0.300	>50
BaL.26	B	CVL2139_006-04	0.034	0.039	0.105	0.221	0.027	0.226	>50
BG1168.01	B	CVL2139_006-07	>50	>50	>50	>50	>50	>50	>50
BL01.DG	B	CVL2139_007-02	20.0	>50	>50	>50	>50	>50	>50
BR07.DG	B	CVL2139_007-04	>50	>50	>50	>50	>50	>50	>50
BX08.16	B	CVL2139_007-06	0.015	0.019	0.030	0.074	0.006	0.053	9.78
CAAN.A2	B	CVL2139_007-11	0.018	0.035	2.53	0.088	0.030	8.69	>50
CNE10	B	CVL2139_008-11	>50	>50	>50	>50	>50	>50	>50
CNE12	B	CVL2139_008-12	0.052	0.057	0.051	0.128	0.052	0.111	13.1
CNE14	B	CVL2139_008-13	0.022	0.025	0.023	0.066	0.019	0.049	0.761
CNE4	B	CVL2139_009-08	>50	>50	>50	>50	>50	>50	>50
CNE57	B	CVL2139_010-01	0.056	0.059	0.530	0.135	0.068	1.15	>50
HO86.8	B	CVL2139_010-12	>50	>50	>50	>50	>50	>50	>50
HT593.1	B	CVL2139_010-13	>50	>50	>50	>50	>50	>50	>50
HXB2.DG	B	CVL2139_011-01	>50	>50	>50	>50	>50	>50	>50
JRCSF.JB	B	CVL2139_011-02	0.049	0.077	47.7	1.17	0.069	>50	>50
JRFL.JB	B	CVL2139_011-03	0.028	0.025	0.182	0.075	0.018	0.396	>50
MN.3	B	CVL2139_011-11	>50	>50	>50	>50	>50	>50	>50
PVO.04	B	CVL2139_012-04	0.082	0.103	>50	0.447	0.146	>50	>50
QH0515.01	B	CVL2139_012-12	4.94	21.2	>50	>50	45.1	>50	>50
QH0692.42	B	CVL2139_012-13	0.216	0.288	>50	3.96	0.437	>50	>50
REJO.67	B	CVL2139_013-05	>50	>50	>50	>50	>50	>50	>50
RHPA.7	B	CVL2139_013-06	0.048	0.055	22.5	0.285	0.092	40.6	>50
SC422.8	B	CVL2139_013-08	0.087	0.085	49.3	0.646	0.091	>50	>50
SF162.LS	B	CVL2139_013-09	0.012	0.016	0.031	0.036	0.006	0.026	1.53
SS1196.01	B	CVL2139_013-12	0.036	0.040	0.645	0.080	0.046	0.633	>50
THRO.18	B	CVL2139_014-13	>50	>50	>50	>50	>50	>50	>50
TRJO.58	B	CVL2139_015-01	0.241	0.440	>50	>50	0.099	>50	>50
TRO.11	B	CVL2139_015-02	0.049	0.064	0.094	0.189	0.039	0.091	25.6
WITO.33	B	CVL2139_015-07	>50	>50	>50	>50	>50	>50	>50
X2278.C2.B6	B	CVL2139_015-13	0.047	0.069	>50	0.855	0.030	>50	>50
YU2.DG	B	CVL2139_016-01	0.111	0.147	0.857	0.634	0.070	0.742	>50
BJOX002000.03.2	BC	CVL2139_006-10	0.019	0.021	0.026	0.096	0.015	0.181	>50
CH038.12	BC	CVL2139_008-05	0.034	0.038	>50	0.217	0.033	>50	>50
CH070.1	BC	CVL2139_008-06	0.119	0.126	1.25	0.621	0.058	22.1	>50
CH117.4	BC	CVL2139_008-07	>50	>50	>50	>50	>50	>50	>50
CH119.10	BC	CVL2139_008-08	0.058	0.056	0.064	0.235	0.028	0.159	>50
CH181.12	BC	CVL2139_008-09	0.086	0.145	44.2	0.773	0.349	>50	>50
CNE15	BC	CVL2139_009-01	>50	>50	>50	>50	>50	>50	>50
CNE19	BC	CVL2139_009-02	0.048	0.140	>50	>50	0.104	>50	>50
CNE20	BC	CVL2139_009-03	>50	>50	>50	>50	>50	>50	>50
CNE21	BC	CVL2139_009-04	0.038	0.050	40.4	0.535	0.049	>50	>50

IC80 — Data from Bonsignori

Virus ID	Clade	Exp. Code	DH270.min1	DH270.min11	DH270.min12	DH270	DH270_IA1	DH270_IA2	DH270_IA4
0260.v5.c36	A	CVL2139_001-05	0.699	0.999	3.54	5.38	0.995	8.09	>50
0330.v4.c3	A	CVL2139_001-06	>50	>50	>50	>50	>50	>50	>50
0439.v5.c1	A	CVL2139_001-07	>50	>50	>50	>50	>50	>50	>50
3365.v2.c20	A	CVL2139_003-07	>50	>50	>50	>50	>50	>50	>50
3415.v1.c1	A	CVL2139_003-08	>50	>50	>50	>50	>50	>50	>50
3718.v3.c11	A	CVL2139_003-12	>50	>50	>50	>50	>50	>50	>50
398-F1.F6.20	A	CVL2139_004-03	0.040	0.038	0.064	0.101	0.033	0.086	7.39
BB201.B42	A	CVL2139_006-05	>50	>50	>50	>50	>50	>50	>50
BB539.2B13	A	CVL2139_006-06	>50	>50	>50	>50	>50	>50	>50
BG505.W6M.C2	A	CVL2139_006-08	>50	>50	>50	>50	>50	>50	>50
BI369.9A	A	CVL2139_006-09	0.230	0.309	1.48	1.01	0.299	3.64	>50
BS208.B1	A	CVL2139_007-05	>50	>50	>50	>50	>50	>50	>50
KER2008.12	A	CVL2139_011-04	0.585	1.89	>50	>50	0.654	>50	>50
KER2018.11	A	CVL2139_011-05	>50	>50	>50	>50	>50	>50	>50
KNH1209.18	A	CVL2139_011-06	0.050	0.067	0.081	0.205	0.077	0.177	11.3
MB201.A1	A	CVL2139_011-08	>50	>50	>50	>50	>50	>50	>50
MB539.2B7	A	CVL2139_011-09	>50	>50	>50	>50	>50	>50	>50
MI369.A5	A	CVL2139_011-10	0.316	0.408	0.459	1.32	0.554	1.41	>50
MS208.A1	A	CVL2139_011-12	>50	>50	>50	>50	>50	>50	>50
Q23.17	A	CVL2139_012-06	0.046	0.066	0.075	0.138	0.061	0.160	>50
Q259.17	A	CVL2139_012-07	>50	>50	>50	>50	>50	>50	>50
Q769.d22	A	CVL2139_012-09	>50	>50	>50	>50	>50	>50	>50
Q769.h5	A	CVL2139_012-10	>50	>50	>50	>50	>50	>50	>50
Q842.d12	A	CVL2139_012-11	>50	>50	>50	>50	>50	>50	>50
QH209.14M.A2	A	CVL2139_013-01	>50	>50	>50	>50	>50	>50	>50
RW020.2	A	CVL2139_013-07	0.026	0.027	0.025	0.116	0.035	0.062	0.910
UG037.8	A	CVL2139_015-06	0.353	1.11	>50	7.61	2.13	>50	>50
246-F3.C10.2	AC	CVL2139_002-04	>50	>50	>50	>50	>50	>50	>50
3301.v1.c24	AC	CVL2139_003-04	0.213	0.297	>50	1.50	0.362	>50	>50
3589.v1.c4	AC	CVL2139_003-10	>50	>50	>50	>50	>50	>50	>50
6540.v4.c1	AC	CVL2139_005-02	>50	>50	>50	>50	>50	>50	>50
6545.v4.c1	AC	CVL2139_005-03	>50	>50	>50	>50	>50	>50	>50
0815.v3.c3	ACD	CVL2139_001-08	0.375	0.358	0.441	1.64	0.571	1.03	24.1
6095.v1.c10	ACD	CVL2139_004-08	>50	>50	>50	>50	>50	>50	>50
3468.v1.c12	AD	CVL2139_003-09	>50	>50	>50	>50	>50	>50	>50
Q168.a2	AD	CVL2139_012-05	>50	>50	>50	>50	>50	>50	>50
Q461.e2	AD	CVL2139_012-08	>50	>50	>50	>50	>50	>50	>50
620345.c1	AE	CVL2139_004-10	>50	>50	>50	>50	>50	>50	>50
BJOX009000.02.4	AE	CVL2139_006-11	>50	>50	>50	>50	>50	>50	>50
BJOX010000.06.2	AE	CVL2139_006-12	>50	>50	>50	>50	>50	>50	>50
BJOX025000.01.1	AE	CVL2139_006-13	>50	>50	>50	>50	>50	>50	>50
BJOX028000.10.3	AE	CVL2139_007-01	>50	>50	>50	>50	>50	>50	>50
C1080.c3	AE	CVL2139_007-07	>50	>50	>50	>50	>50	>50	>50
C2101.c1	AE	CVL2139_007-08	>50	>50	>50	>50	>50	>50	>50
C3347.c11	AE	CVL2139_007-09	>50	>50	>50	>50	>50	>50	>50
C4118.09	AE	CVL2139_007-10	>50	>50	>50	>50	>50	>50	>50
CM244.ec1	AE	CVL2139_008-10	>50	>50	>50	>50	>50	>50	>50
CNE3	AE	CVL2139_009-05	>50	>50	>50	>50	>50	>50	>50
CNE5	AE	CVL2139_009-10	>50	>50	>50	>50	>50	>50	>50
CNE55	AE	CVL2139_009-12	>50	>50	>50	>50	>50	>50	>50
CNE56	AE	CVL2139_009-13	>50	>50	>50	>50	>50	>50	>50
CNE59	AE	CVL2139_010-03	>50	>50	>50	>50	>50	>50	>50
CNE8	AE	CVL2139_010-05	>50	>50	>50	>50	>50	>50	>50
M02138	AE	CVL2139_011-07	>50	>50	>50	>50	>50	>50	>50
R1166.c1	AE	CVL2139_013-02	>50	>50	>50	>50	>50	>50	>50
R2184.c4	AE	CVL2139_013-03	>50	>50	>50	>50	>50	>50	>50
R3265.c6	AE	CVL2139_013-04	>50	>50	>50	>50	>50	>50	>50
TH023.6	AE	CVL2139_014-10	>50	>50	>50	>50	>50	>50	>50
TH966.8	AE	CVL2139_014-11	>50	>50	>50	>50	>50	>50	>50
TH976.17	AE	CVL2139_014-12	>50	>50	>50	>50	>50	>50	>50
235-47	AG	CVL2139_002-02	0.490	0.492	>50	12.1	0.564	>50	>50
242-14	AG	CVL2139_002-03	>50	>50	>50	>50	>50	>50	>50
263-8	AG	CVL2139_002-10	10.8	>50	>50	>50	>50	>50	>50
269-12	AG	CVL2139_002-11	0.611	0.987	>50	9.15	1.56	>50	>50
271-11	AG	CVL2139_002-12	>50	>50	>50	>50	>50	>50	>50
928-28	AG	CVL2139_005-10	>50	>50	>50	>50	>50	>50	>50
DJ263.8	AG	CVL2139_010-06	0.233	0.315	>50	0.865	0.184	3.32	>50
T250-4	AG	CVL2139_014-01	>50	>50	>50	>50	>50	>50	>50
T251-18	AG	CVL2139_014-02	>50	>50	>50	>50	>50	>50	>50
T253-11	AG	CVL2139_014-03	>50	>50	>50	>50	>50	>50	>50
T255-34	AG	CVL2139_014-04	>50	>50	>50	>50	>50	>50	>50
T257-31	AG	CVL2139_014-05	>50	>50	>50	>50	>50	>50	>50
T266-60	AG	CVL2139_014-06	7.59	24.1	>50	30.9	19.0	>50	>50
T278-50	AG	CVL2139_014-07	>50	>50	>50	>50	>50	>50	>50
T280-5	AG	CVL2139_014-08	0.039	0.042	0.053	0.169	0.048	0.244	>50
T33-7	AG	CVL2139_014-09	>50	>50	>50	>50	>50	>50	>50
3988.25	B	CVL2139_004-02	0.047	0.053	7.43	0.184	0.056	2.81	>50
5768.04	B	CVL2139_004-07	5.41	21.4	>50	>50	0.799	>50	>50
6101.10	B	CVL2139_004-09	0.093	0.080	0.076	0.327	0.083	0.287	8.57
6535.3	B	CVL2139_005-01	0.067	0.074	0.069	0.276	0.066	0.171	1.61
7165.18	B	CVL2139_005-08	0.138	0.188	9.08	0.815	0.140	18.6	>50
45_01dG5	B	CVL2139_004-05	0.093	0.152	0.185	0.301	0.241	0.568	>50
89.6.DG	B	CVL2139_005-09	0.120	0.151	>50	0.628	0.265	>50	>50
AC10.29	B	CVL2139_006-01	0.049	0.056	>50	4.47	0.057	>50	>50
ADA.DG	B	CVL2139_006-02	0.177	0.244	>50	1.97	0.221	>50	>50
Bal.01	B	CVL2139_006-03	0.156	0.154	>50	0.636	0.120	>50	>50
BaL.26	B	CVL2139_006-04	0.097	0.133	3.21	0.541	0.112	0.516	>50
BG1168.01	B	CVL2139_006-07	>50	>50	>50	>50	>50	>50	>50
BL01.DG	B	CVL2139_007-02	>50	>50	>50	>50	>50	>50	>50
BR07.DG	B	CVL2139_007-04	>50	>50	>50	>50	>50	>50	>50
BX08.16	B	CVL2139_007-06	0.042	0.048	0.091	0.198	0.047	0.156	>50
CAAN.A2	B	CVL2139_007-11	0.052	0.090	>50	0.267	0.092	>50	>50
CNE10	B	CVL2139_008-11	>50	>50	>50	>50	>50	>50	>50
CNE12	B	CVL2139_008-12	0.085	0.090	0.162	0.325	0.133	0.292	>50
CNE14	B	CVL2139_008-13	0.050	0.063	0.071	0.153	0.065	0.121	6.23
CNE4	B	CVL2139_009-08	>50	>50	>50	>50	>50	>50	>50
CNE57	B	CVL2139_010-01	0.120	0.130	3.87	0.291	0.193	6.31	>50
HO86.8	B	CVL2139_010-12	>50	>50	>50	>50	>50	>50	>50
HT593.1	B	CVL2139_010-13	>50	>50	>50	>50	>50	>50	>50
HXB2.DG	B	CVL2139_011-01	>50	>50	>50	>50	>50	>50	>50
JRCSF.JB	B	CVL2139_011-02	0.159	0.306	>50	4.55	0.307	>50	>50
JRFL.JB	B	CVL2139_011-03	0.064	0.063	1.25	0.202	0.059	3.61	>50
MN.3	B	CVL2139_011-11	>50	>50	>50	>50	>50	>50	>50
PVO.04	B	CVL2139_012-04	0.232	0.303	>50	1.39	0.633	>50	>50
QH0515.01	B	CVL2139_012-12	>50	>50	>50	>50	>50	>50	>50
QH0692.42	B	CVL2139_012-13	0.737	1.60	>50	24.1	2.27	>50	>50
REJO.67	B	CVL2139_013-05	>50	>50	>50	>50	>50	>50	>50
RHPA.7	B	CVL2139_013-06	0.138	0.235	>50	1.20	0.260	>50	>50
SC422.8	B	CVL2139_013-08	0.198	0.223	>50	1.82	0.337	>50	>50
SF162.LS	B	CVL2139_013-09	0.025	0.037	0.096	0.096	0.044	0.068	>50
SS1196.01	B	CVL2139_013-12	0.078	0.089	5.91	0.214	0.106	5.39	>50
THRO.18	B	CVL2139_014-13	>50	>50	>50	>50	>50	>50	>50
TRJO.58	B	CVL2139_015-01	0.667	1.73	>50	>50	0.470	>50	>50
TRO.11	B	CVL2139_015-02	0.126	0.148	0.309	0.510	0.120	0.291	>50
WITO.33	B	CVL2139_015-07	>50	>50	>50	>50	>50	>50	>50
X2278.C2.B6	B	CVL2139_015-13	0.120	0.187	>50	4.58	0.137	>50	>50
YU2.DG	B	CVL2139_016-01	0.321	0.387	25.4	1.81	0.195	3.83	>50
BJOX002000.03.2	BC	CVL2139_006-10	0.048	0.059	0.074	0.236	0.048	1.05	>50
CH038.12	BC	CVL2139_008-05	0.109	0.113	>50	0.674	0.123	>50	>50
CH070.1	BC	CVL2139_008-06	0.291	0.351	11.3	1.70	0.237	>50	>50
CH117.4	BC	CVL2139_008-07	>50	>50	>50	>50	>50	>50	>50
CH119.10	BC	CVL2139_008-08	0.091	0.140	0.185	0.412	0.084	0.381	>50
CH181.12	BC	CVL2139_008-09	0.230	0.376	>50	2.52	1.14	>50	>50
CNE15	BC	CVL2139_009-01	>50	>50	>50	>50	>50	>50	>50
CNE19	BC	CVL2139_009-02	>50	>50	>50	>50	>50	>50	>50
CNE20	BC	CVL2139_009-03	>50	>50	>50	>50	>50	>50	>50
CNE21	BC	CVL2139_009-04	0.112	0.138	>50	1.61	0.153	>50	>50

IC50 Table (left)

Virus	Clade	CVL ID	DH270.min1	DH270.min11	DH270.min12	DH270	DH270_IA1	DH270_IA2	DH270_IA4
CNE40	BC	CVL2139_009-09	>50	>50	>50	>50	>50	>50	>50
CNE7	BC	CVL2139_010-04	>50	>50	>50	>50	>50	>50	>50
286.36	C	CVL2139_018-01	0.038	0.057	0.120	0.119	0.041	0.085	16.6
288.38	C	CVL2139_003-01	0.028	0.028	0.057	0.148	0.049	0.104	7.76
0013095-2.11	C	CVL2139_001-01	>50	>50	>50	>50	>50	>50	>50
001428-2.42	C	CVL2139_001-02	>50	>50	>50	>50	>50	>50	>50
0077.v1.c16	C	CVL2139_001-03	>50	>50	>50	>50	>50	>50	>50
00836-2.5	C	CVL2139_001-04	>50	>50	>50	>50	>50	>50	>50
0921.v2.c14	C	CVL2139_001-09	>50	>50	>50	>50	>50	>50	>50
16055-2.3	C	CVL2139_001-10	>50	>50	>50	>50	>50	>50	>50
16845-2.22	C	CVL2139_001-11	2.20	1.80	>50	>50	2.17	>50	>50
16936-2.21	C	CVL2139_001-12	0.051	0.066	0.404	0.725	0.073	32.1	>50
25710-2.43	C	CVL2139_002-06	0.112	0.153	32.4	2.07	0.170	>50	>50
25711-2.4	C	CVL2139_002-07	0.018	0.019	0.571	0.200	0.023	>50	>50
25925-2.22	C	CVL2139_002-08	8.27	7.12	>50	>50	11.2	>50	>50
26191-2.48	C	CVL2139_002-09	>50	>50	>50	>50	>50	>50	>50
3168.v4.c10	C	CVL2139_003-03	0.218	0.167	28.8	0.856	0.396	>50	>50
3637.v5.c3	C	CVL2139_003-11	>50	>50	>50	>50	>50	>50	>50
3873.v1.c24	C	CVL2139_004-01	0.056	0.052	8.95	0.374	0.091	>50	>50
426c	C	CVL2139_004-04	0.981	2.09	>50	>50	1.97	>50	>50
6322.v4.c1	C	CVL2139_004-11	>50	>50	>50	>50	>50	>50	>50
6471.v1.c16	C	CVL2139_004-13	6.82	5.04	>50	>50	1.26	>50	>50
6631.v3.c10	C	CVL2139_005-04	>50	>50	>50	>50	>50	>50	>50
6644.v2.c33	C	CVL2139_005-05	0.025	0.031	0.174	0.128	0.031	0.148	>50
6785.v5.c14	C	CVL2139_005-06	0.050	0.054	0.174	0.247	0.033	0.462	>50
6838.v1.c35	C	CVL2139_005-07	>50	>50	>50	>50	>50	>50	>50
96ZM651.02	C	CVL2139_005-11	0.022	0.025	5.65	0.660	0.016	18.1	>50
BR025.9	C	CVL2139_007-03	0.011	0.013	0.016	0.060	0.011	0.032	0.500
CAP210.E8	C	CVL2139_007-12	>50	>50	>50	>50	>50	>50	>50
CAP244.D3	C	CVL2139_007-13	>50	>50	>50	>50	14.4	>50	>50
CAP256.206.C9	C	CVL2139_008-01	0.053	0.049	0.043	0.128	0.040	0.125	>50
CAP45.G3	C	CVL2139_008-02	>50	>50	>50	>50	>50	>50	>50
Ce1176.A3	C	CVL2139_008-03	0.066	0.075	0.115	0.294	0.076	0.204	>50
CE703010217.B6	C	CVL2139_008-04	0.033	0.041	0.144	0.457	0.023	>50	>50
CNE30	C	CVL2139_009-06	1.02	0.765	4.54	3.02	0.801	9.12	>50
CNE31	C	CVL2139_009-07	>50	>50	>50	>50	>50	>50	>50
CNE53	C	CVL2139_009-11	0.103	0.102	0.100	0.406	0.107	0.513	>50
CNE58	C	CVL2139_010-02	0.208	0.270	43.7	2.20	0.284	>50	>50
DU123.06	C	CVL2139_010-07	0.102	0.126	0.193	0.531	0.112	0.591	>50
DU151.02	C	CVL2139_010-08	>50	>50	>50	>50	>50	>50	>50
DU156.12	C	CVL2139_010-09	0.077	0.088	0.087	0.234	0.055	0.323	>50
DU172.17	C	CVL2139_010-10	1.29	1.81	>50	13.8	0.428	>50	>50
DU422.01	C	CVL2139_010-11	0.144	0.232	0.211	0.554	0.147	28.0	>50
MW965.26	C	CVL2139_011-13	0.027	0.026	>50	0.135	0.036	>50	>50
SO18.18	C	CVL2139_013-11	0.023	0.030	0.141	0.073	0.025	0.374	>50
TV1.29	C	CVL2139_018-03	1.06	0.828	16.9	8.94	0.742	15.8	>50
TZA125.17	C	CVL2139_015-04	>50	>50	>50	>50	>50	>50	>50
TZBD.02	C	CVL2139_015-05	0.053	0.060	0.212	1.82	0.078	>50	>50
ZA012.29	C	CVL2139_016-02	0.040	0.054	2.71	9.33	0.055	>50	>50
ZM106.9	C	CVL2139_016-03	0.034	0.044	0.054	0.127	0.057	0.381	>50
ZM109.4	C	CVL2139_016-04	>50	>50	>50	>50	>50	>50	>50
ZM135.10a	C	CVL2139_016-05	0.110	0.240	>50	2.12	0.142	>50	>50
ZM176.66	C	CVL2139_016-06	2.65	2.77	>50	>50	14.7	>50	>50
ZM197.7	C	CVL2139_016-07	>50	>50	>50	>50	>50	>50	>50
ZM214.15	C	CVL2139_016-08	1.58	2.26	>50	46.9	1.32	>50	>50
ZM215.8	C	CVL2139_016-09	1.76	1.52	>50	5.13	0.547	>50	>50
ZM233.6	C	CVL2139_016-10	>50	>50	>50	>50	>50	>50	>50
ZM249.1	C	CVL2139_016-11	>50	>50	>50	>50	>50	>50	>50
ZM53.12	C	CVL2139_016-12	>50	>50	>50	>50	>50	>50	>50
ZM55.28a	C	CVL2139_016-13	0.030	0.044	0.057	0.145	0.036	0.072	43.7
3326.v4.c3	CD	CVL2139_003-05	>50	>50	>50	>50	>50	>50	>50
3337.v2.c6	CD	CVL2139_003-06	31.3	35.4	>50	34.0	>50	>50	>50
3817.v2.c59	CD	CVL2139_003-13	0.221	0.327	>50	0.736	0.222	>50	>50
191821.E6.1	D	CVL2139_001-13	>50	>50	>50	>50	>50	>50	>50
231965.c01	D	CVL2139_002-01	>50	>50	>50	>50	>50	>50	>50
247-23	D	CVL2139_002-05	>50	>50	>50	>50	>50	>50	>50
3016.v5.c45	D	CVL2139_003-02	>50	>50	>50	>50	>50	>50	>50
57128.vrc15	D	CVL2139_017-04	26.6	25.6	>50	17.4	>50	>50	>50
6405.v4.c34	D	CVL2139_004-12	>50	>50	>50	>50	>50	>50	>50
A03349M1.vrc4a	D	CVL2139_005-12	>50	>50	>50	>50	>50	>50	>50
A07412M1.vrc12	D	CVL2139_005-13	>50	>50	>50	>50	>50	>50	>50
NKU3006.ec1	D	CVL2139_012-01	>50	>50	>50	>50	>50	>50	>50
UG021.16	D	CVL2139_019-02	>50	>50	>50	>50	>50	>50	>50
UG024.2	D	CVL2139_019-01	>50	>50	>50	>50	>50	>50	>50
P0402.c2.11	G	CVL2139_012-02	3.83	3.53	>50	14.8	5.20	>50	>50
P1981.C5.3	G	CVL2139_012-03	0.013	0.013	0.015	0.047	0.011	0.031	11.6
X1193.c1	G	CVL2139_015-08	1.07	1.70	>50	31.3	0.372	>50	>50
X1254.c3	G	CVL2139_015-09	0.067	0.078	2.08	0.259	0.082	17.3	>50
X1632.S2.B10	G	CVL2139_015-10	>50	>50	>50	>50	>50	>50	>50
X2088.c9	G	CVL2139_015-11	0.020	0.029	0.065	0.158	0.027	6.68	>50
X2131.C1.B5	G	CVL2139_015-12	0.169	0.214	>50	0.972	0.079	>50	>50
IVmac251.30.SG3	NA	CVL2139_013-10	>50	>50	>50	>50	>50	>50	>50
SVA.MLV	NA	CVL2139_013-13	>50	>50	>50	>50	>50	>50	>50

	DH270.min1	DH270.min11	DH270.min12	DH270	DH270_IA1	DH270_IA2	DH270_IA4
# Viruses	208	208	208	208	208	208	208
Total VS Neutralized							
IC50 <50ug/ml	100	99	68	88	98	53	19
IC50 <10ug/ml	96	95	57	81	94	46	11
IC50 <1.0ug/ml	83	83	46	65	87	39	4
IC50 <0.1ug/ml	61	54	23	17	63	14	0
IC50 <0.01ug/ml	0	1	1	0	2	0	0
% VS Neutralized							
IC50 <50ug/ml	48	48	33	42	47	25	9
IC50 <10ug/ml	46	46	27	39	45	22	5
IC50 <1.0ug/ml	40	40	22	31	42	19	2
IC50 <0.1ug/ml	29	26	11	8	30	7	0
IC50 <0.01ug/ml	0	0	0	0	1	0	0
Median IC50	0.059	0.076	0.188	0.390	0.073	0.374	7.76
Geometric Mean	0.113	0.135	0.408	0.482	0.103	0.449	4.41

IC80 Table (right)

Virus	Clade	CVL ID	DH270.min1	DH270.min11	DH270.min12	DH270	DH270_IA1	DH270_IA2	DH270_IA4
CNE40	BC	CVL2139_009-09	>50	>50	>50	>50	>50	>50	>50
CNE7	BC	CVL2139_010-04	>50	>50	>50	>50	>50	>50	>50
286.36	C	CVL2139_018-01	0.106	0.147	0.328	0.342	0.088	0.200	>50
288.38	C	CVL2139_003-01	0.062	0.090	0.166	0.390	0.128	0.299	44.5
0013095-2.11	C	CVL2139_001-01	>50	>50	>50	>50	>50	>50	>50
001428-2.42	C	CVL2139_001-02	>50	>50	>50	>50	>50	>50	>50
0077.v1.c16	C	CVL2139_001-03	>50	>50	>50	>50	>50	>50	>50
00836-2.5	C	CVL2139_001-04	>50	>50	>50	>50	>50	>50	>50
0921.v2.c14	C	CVL2139_001-09	>50	>50	>50	>50	>50	>50	>50
16055-2.3	C	CVL2139_001-10	>50	>50	>50	>50	>50	>50	>50
16845-2.22	C	CVL2139_001-11	8.38	10.5	>50	>50	9.28	>50	>50
16936-2.21	C	CVL2139_001-12	0.136	0.243	2.19	2.05	0.234	>50	>50
25710-2.43	C	CVL2139_002-06	0.301	0.383	>50	8.80	0.450	>50	>50
25711-2.4	C	CVL2139_002-07	0.042	0.051	3.36	1.50	0.051	>50	>50
25925-2.22	C	CVL2139_002-08	>50	>50	>50	>50	>50	>50	>50
26191-2.48	C	CVL2139_002-09	>50	>50	>50	>50	>50	>50	>50
3168.v4.c10	C	CVL2139_003-03	0.654	0.839	>50	4.14	1.38	>50	>50
3637.v5.c3	C	CVL2139_003-11	>50	>50	>50	>50	>50	>50	>50
3873.v1.c24	C	CVL2139_004-01	0.125	0.159	>50	0.976	0.258	>50	>50
426c	C	CVL2139_004-04	24.2	>50	>50	>50	>50	>50	>50
6322.v4.c1	C	CVL2139_004-11	>50	>50	>50	>50	>50	>50	>50
6471.v1.c16	C	CVL2139_004-13	>50	>50	>50	>50	10.1	>50	>50
6631.v3.c10	C	CVL2139_005-04	>50	>50	>50	>50	>50	>50	>50
6644.v2.c33	C	CVL2139_005-05	0.095	0.106	>50	0.328	0.122	0.660	>50
6785.v5.c14	C	CVL2139_005-06	0.118	0.140	1.09	0.525	0.115	1.66	>50
6838.v1.c35	C	CVL2139_005-07	>50	>50	>50	>50	>50	>50	>50
96ZM651.02	C	CVL2139_005-11	0.102	0.348	>50	>50	0.429	>50	>50
BR025.9	C	CVL2139_007-03	0.032	0.045	0.042	0.127	0.039	0.078	1.97
CAP210.E8	C	CVL2139_007-12	>50	>50	>50	>50	>50	>50	>50
CAP244.D3	C	CVL2139_007-13	>50	>50	>50	>50	>50	>50	>50
CAP256.206.C9	C	CVL2139_008-01	0.114	0.129	0.119	0.325	0.105	0.328	>50
CAP45.G3	C	CVL2139_008-02	>50	>50	>50	>50	>50	>50	>50
Ce1176.A3	C	CVL2139_008-03	0.188	0.190	0.349	0.755	0.238	0.499	>50
CE703010217.B6	C	CVL2139_008-04	0.087	0.110	0.660	1.81	0.076	>50	>50
CNE30	C	CVL2139_009-06	2.03	2.30	16.6	6.49	2.18	40.6	>50
CNE31	C	CVL2139_009-07	>50	>50	>50	>50	>50	>50	>50
CNE53	C	CVL2139_009-11	0.297	0.323	0.330	1.14	0.306	1.25	>50
CNE58	C	CVL2139_010-02	0.659	0.650	>50	10.3	0.865	>50	>50
DU123.06	C	CVL2139_010-07	0.280	0.377	1.15	1.72	0.362	4.46	>50
DU151.02	C	CVL2139_010-08	>50	>50	>50	>50	>50	>50	>50
DU156.12	C	CVL2139_010-09	0.155	0.216	0.264	0.710	0.159	1.53	>50
DU172.17	C	CVL2139_010-10	12.2	23.3	>50	>50	5.12	>50	>50
DU422.01	C	CVL2139_010-11	0.380	0.481	0.539	1.30	0.391	>50	>50
MW965.26	C	CVL2139_011-13	0.119	0.104	>50	0.773	0.121	>50	>50
SO18.18	C	CVL2139_013-11	0.059	0.070	0.695	0.167	0.081	2.32	>50
TV1.29	C	CVL2139_018-03	3.86	3.15	>50	26.1	2.96	>50	>50
TZA125.17	C	CVL2139_015-04	>50	>50	>50	>50	>50	>50	>50
TZBD.02	C	CVL2139_015-05	0.125	0.186	2.13	10.4	0.256	>50	>50
ZA012.29	C	CVL2139_016-02	0.107	0.158	>50	>50	0.187	>50	>50
ZM106.9	C	CVL2139_016-03	0.088	0.122	0.117	0.316	0.136	2.13	>50
ZM109.4	C	CVL2139_016-04	>50	>50	>50	>50	>50	>50	>50
ZM135.10a	C	CVL2139_016-05	0.391	1.02	>50	18.7	0.559	>50	>50
ZM176.66	C	CVL2139_016-06	15.6	21.6	>50	>50	>50	>50	>50
ZM197.7	C	CVL2139_016-07	>50	>50	>50	>50	>50	>50	>50
ZM214.15	C	CVL2139_016-08	4.38	9.36	>50	>50	15.0	>50	>50
ZM215.8	C	CVL2139_016-09	25.7	15.0	>50	31.1	8.29	>50	>50
ZM233.6	C	CVL2139_016-10	>50	>50	>50	>50	>50	>50	>50
ZM249.1	C	CVL2139_016-11	>50	>50	>50	>50	>50	>50	>50
ZM53.12	C	CVL2139_016-12	>50	>50	>50	>50	>50	>50	>50
ZM55.28a	C	CVL2139_016-13	0.072	0.118	0.147	0.329	0.113	0.206	>50
3326.v4.c3	CD	CVL2139_003-05	>50	>50	>50	>50	>50	>50	>50
3337.v2.c6	CD	CVL2139_003-06	>50	>50	>50	>50	>50	>50	>50
3817.v2.c59	CD	CVL2139_003-13	0.856	1.39	>50	2.76	1.02	>50	>50
191821.E6.1	D	CVL2139_001-13	>50	>50	>50	>50	>50	>50	>50
231965.c01	D	CVL2139_002-01	>50	>50	>50	>50	>50	>50	>50
247-23	D	CVL2139_002-05	>50	>50	>50	>50	>50	>50	>50
3016.v5.c45	D	CVL2139_003-02	>50	>50	>50	>50	>50	>50	>50
57128.vrc15	D	CVL2139_017-04	>50	>50	>50	>50	>50	>50	>50
6405.v4.c34	D	CVL2139_004-12	>50	>50	>50	>50	>50	>50	>50
A03349M1.vrc4a	D	CVL2139_005-12	>50	>50	>50	>50	>50	>50	>50
A07412M1.vrc12	D	CVL2139_005-13	>50	>50	>50	>50	>50	>50	>50
NKU3006.ec1	D	CVL2139_012-01	>50	>50	>50	>50	>50	>50	>50
UG021.16	D	CVL2139_019-02	>50	>50	>50	>50	>50	>50	>50
UG024.2	D	CVL2139_019-01	>50	>50	>50	>50	>50	>50	>50
P0402.c2.11	G	CVL2139_012-02	49.5	45.4	>50	>50	>50	>50	>50
P1981.C5.3	G	CVL2139_012-03	0.029	0.031	0.058	0.108	0.028	0.125	>50
X1193.c1	G	CVL2139_015-08	5.39	8.05	>50	>50	1.90	>50	>50
X1254.c3	G	CVL2139_015-09	0.143	0.183	22.9	0.727	0.157	>50	>50
X1632.S2.B10	G	CVL2139_015-10	>50	>50	>50	>50	>50	>50	>50
X2088.c9	G	CVL2139_015-11	0.064	0.095	0.304	0.406	0.084	>50	>50
X2131.C1.B5	G	CVL2139_015-12	0.506	0.435	>50	5.13	0.416	>50	>50
SIVmac251.30.SG3	NA	CVL2139_013-10	>50	>50	>50	>50	>50	>50	>50
SVA.MLV	NA	CVL2139_013-13	>50	>50	>50	>50	>50	>50	>50

	DH270.min1	DH270.min11	DH270.min12	DH270	DH270_IA1	DH270_IA2	DH270_IA4
# Viruses	208	208	208	208	208	208	208
Total VS Neutralized							
IC80 <50ug/ml	92	90	48	79	89	42	9
IC80 <10ug/ml	86	83	44	71	86	40	6
IC80 <1.0ug/ml	79	73	31	42	74	23	1
IC80 <0.1ug/ml	30	23	13	1	23	4	0
IC80 <0.01ug/ml	0	0	0	0	0	0	0
% VS Neutralized							
IC80 <50ug/ml	44	43	23	38	43	20	4
IC80 <10ug/ml	41	40	21	34	41	19	3
IC80 <1.0ug/ml	38	35	15	20	36	11	0
IC80 <0.1ug/ml	14	11	6	0	11	2	0
IC80 <0.01ug/ml	0	0	0	0	0	0	0
Median IC80	0.138	0.188	0.329	0.815	0.195	0.542	7.39
Geometric Mean	0.246	0.303	0.493	1.02	0.267	0.739	6.21

Note: Median and Geometric Mean titers are calculated only for samples with IC50 <50ug/ml

Supplementary Figure 2. Neutralization profile of minimized DH270 variants on global panel. of 208 isolates. Values indicated calculated IC$_{50}$ and IC$_{80}$ of minDH270 against the individual viruses in the panel. DH270.6 neutralized 114 viruses from this panel with a mean IC$_{50}$ of 0.136 µg/ml. DH270min1 neutralized 100 viruses (mean IC$_{50}$=0.113 µg/ml), DH270min11 neutralized 99 viruses (mean IC$_{50}$=0.135 µg/ml) and DH270min12 neutralized 68 viruses (mean IC$_{50}$=0.408 µg/ml).

Fold binding decrease of DH270.min1 single residue reversion to UCA

	LogAUC DH270.6	HC.D31G	HC.M51I	HC.T55S	HC.R57G	HC.T98R	HC.Y110G	LC.Y27S	LC.Y48L	LC.F93Y	LC.G94A	LC.A97S	LC.C101F
10.17 [CH0848d0949.10.17D11gp120/293F/Mon]	13.175	1.0	1.0	1.0	1.1	1.2	1.3	1.1	1.1	1.2	1.0	1.0	1.1
1432.41 [CH848.3.d1432.5.41D11gp120/293F/Mon]	12.831	1.0	1.0	1.0	1.2	17.9	21.7	1.4	3.1	1.4	1.2	1.3	1.1
358.06 [CH848.3.d0358.80.06D11gp120]	10.718	1.2	1.4	1.2	3.1	NB	NB	2.6	53.9	3.2	1.7	1.6	1.4
526.02 [CH848.3.d0526.25.02D11gp120/293F]	7.015	1.5	1.2	1.3	8.5	NB	NB	13.5	NB	26.4	3.2	2.6	2.2
836.31 [CH0848d0836.10.31D11gp120/293F]	9.845	1.2	1.0	1.1	2.2	9.4	29.7	2.2	4.3	2.7	1.4	1.9	1.3
526.09 [CH848.3.d0526.25.09D11gp120/293F]	8.924	1.3	1.3	1.2	4.1	NB	NB	6.5	NB	11.0	2.4	2.4	1.7
CH848.3.d1120.10.41D11gp120/293F/PEI	13.274	1.0	1.0	1.0	1.0	1.1	1.2	1.1	1.1	1.0	1.0	1.0	1.0
CH848.3d0864.7.26D11gp120	12.808	1.0	1.0	1.1	1.2	2.2	3.3	1.2	1.7	1.2	1.1	1.2	1.1
CH848.3d0808.15.25D11gp120	12.515	1.1	1.0	1.1	1.9	17.7	100.1	1.6	3.7	1.7	1.2	1.4	1.1
CH0848d0808.15.27D11gp120/293F	12.409	1.0	1.0	1.0	1.3	2.7	3.3	1.3	1.9	1.3	1.1	1.2	1.1
CH848.3.d1120.10.32D11gp120/293F/PEI	12.397	1.0	1.0	1.0	1.1	1.8	1.9	1.2	1.5	1.1	1.0	1.1	1.0
CH848.3.d0135.27.06D11gp120	12.091	1.1	1.5	1.1	2.1	NB	19.6	1.6	16.1	1.6	1.5	1.4	1.3
CH0848d0808.15.15D11gp120/293F	12.04	1.1	1.1	1.1	1.5	3.2	3.3	1.5	2.3	1.5	1.2	1.2	1.1
CH0848d1432.5.48D11gp120/293F	11.961	1.0	1.0	1.0	2.1	6.7	23.4	1.5	3.0	1.6	1.2	1.4	1.1
CH0848d0780.15.29D11gp120/293F	11.937	1.1	1.0	1.1	1.5	2.8	2.9	1.5	2.1	1.5	1.2	1.2	1.1
CH848.3.d0611.20.28D11gp120	11.897	1.1	1.0	1.1	1.3	15.9	40.7	1.7	5.0	1.8	1.3	1.2	1.2
CH848.3.d0864.7.39D11gp120	11.794	1.1	1.0	1.1	1.5	3.6	6.7	1.6	2.4	1.7	1.2	1.4	1.2
CH848.3.d0794.5.41D11gp120/293F	11.788	1.0	1.0	1.0	1.4	2.3	2.7	1.4	1.9	1.4	1.1	1.2	1.1
CH848.3.d0611.9.02D11gp120	11.76	1.1	1.0	1.1	1.2	11.3	84.6	1.7	3.8	1.9	1.3	1.2	1.2
CH0848d1432.5.06D11gp120/293F	11.751	1.0	1.1	1.0	2.3	8.4	29.9	1.6	3.4	1.7	1.2	1.4	1.1
CH0848d0780.25.05D11gp120	11.207	1.1	1.2	1.1	3.2	NB	NB	2.8	45.4	3.5	1.8	1.6	1.4
CH0848d0808.15.43D11gp120/293F	11.649	1.1	1.0	1.1	2.1	6.8	8.1	1.7	3.9	1.9	1.3	1.4	1.2
CH848.3.d1621.4.44D11gp120/293F	11.346	1.1	1.1	1.0	2.0	NB	27.4	1.7	28.9	1.7	1.5	1.7	1.3
CH848.3.d0358.80.44D11gp120	11.275	1.2	1.1	1.2	2.5	NB	78.3	2.0	32.8	2.2	1.6	1.5	1.3
CH848.3.d0526.25.21D11gp120	11.116	1.1	1.0	1.1	2.6	NB	NB	2.1	17.4	2.5	1.6	1.6	1.3
CH848.3.d0274.30.09D11gp120/293F	10.95	1.2	1.4	1.2	2.8	NB	121.7	2.2	33.8	2.5	1.7	1.6	1.4
CH848.3.d0700.15.29D11gp120/293F	10.942	1.1	1.0	1.1	3.2	NB	NB	2.9	43.4	3.8	1.8	1.7	1.4
CH848.3.d0358.80.17D11gp120/293F	10.918	1.2	1.1	1.2	3.0	NB	NB	2.4	37.8	2.8	1.7	1.6	1.4
CH848.3.d1120.10.24D11gp120/293i/PEI	10.849	1.1	1.0	1.1	1.8	5.2	10.8	1.7	3.0	1.9	1.3	1.6	1.2
CH848.3.d0893.10.06D11gp120/293F	10.793	1.1	1.0	1.1	1.4	7.0	29.7	1.7	3.5	2.0	1.3	1.6	1.2
CH848.3.d0445.25.26D11gp120/293F	10.782	1.2	1.1	1.2	3.3	NB	136.5	2.7	113.5	3.3	1.8	1.6	1.5
CH0848d0780.15.22D11gp120/293VRC	10.781	1.1	1.0	1.1	2.1	3.8	4.8	2.0	3.1	2.3	1.5	1.5	1.3
CH848.3.d0107.30.27D11gp120/293F	10.704	1.2	1.7	1.2	3.2	NB	65.7	2.2	127.4	2.5	1.6	1.7	1.4
CH848.3.d0700.15.15D11gp120	10.6	1.2	1.4	1.2	3.7	NB	NB	3.5	108.2	4.9	2.0	1.7	1.4
CH848.3.d0836.10.36D11gp120/293F	10.547	1.1	1.2	1.1	1.7	4.8	11.3	1.8	2.9	2.0	1.3	1.8	1.2
CH0848d1120.10.05D11gp120/293F	10.483	1.1	1.0	1.0	1.5	6.5	9.7	1.6	2.9	1.7	1.1	1.4	1.1
CH848.3.d0445.30.42D11gp120	10.474	1.2	1.1	1.2	4.0	NB	NB	3.2	45.0	3.9	1.9	1.8	1.5
CH848.3.d1432.5.18D11gp120/293F	10.33	1.1	1.2	1.0	2.6	NB	69.3	2.1	100.3	2.1	1.7	2.0	1.3
CH848.3.d0135.60.05D11gp120	10.327	1.2	1.2	1.2	3.5	NB	77.1	2.4	NB	2.5	1.8	1.6	1.4
CH0848d0794.3.03D11gp120/293F	10.326	1.0	1.1	1.1	1.1	24.7	109.9	2.1	3.5	3.1	1.4	1.1	1.4
CH848.3.d0526.25.10D11gp120/293F	10.064	1.2	1.2	1.2	4.4	NB	NB	3.8	NB	5.5	2.0	1.8	1.5
CH848.3.d1432.5.27D11gp120/293F	9.881	1.1	1.0	1.1	1.5	6.3	4.2	1.6	2.5	1.8	1.2	0.9	1.3
CH848.3.d0194.25.17D11gp120	9.785	1.2	1.1	1.2	4.7	NB	NB	4.0	NB	5.5	1.9	1.8	1.4
CH848.3.d0135.60.20D11gp120/293F	9.757	1.2	1.3	1.2	3.7	NB	NB	2.8	NB	3.2	1.9	1.7	1.4
CH848.3.d0107.30.31D11gp120/293F	9.647	1.3	2.0	1.3	4.4	NB	NB	3.4	NB	3.9	2.0	1.9	1.5
CH848.3.d0611.20.12D11gp120/293F	9.641	1.2	1.1	1.1	1.6	NB	NB	2.7	15.2	3.7	1.7	1.6	1.4
CH848.3 TF gp120	9.615	1.1	1.1	1.2	3.4	NB	NB	2.8	NB	2.9	1.8	1.6	1.3
CH848.3.d0078.30.42D11gp120/293F	9.515	1.2	1.2	1.2	3.8	NB	NB	2.8	NB	3.1	1.9	1.7	1.4
CH848.3.d1651.10.04D11gp120/293F	9.341	1.2	1.1	1.1	2.9	NB	NB	3.8	NB	4.5	2.0	3.0	1.5
CH848.3.d0107.30.12D11gp120	9.12	1.3	1.5	1.2	4.4	NB	NB	4.0	NB	4.6	2.1	1.9	1.5
CH848.3.d0445.25.04D11gp120/293F	9.026	1.3	1.1	1.2	4.9	NB	NB	4.7	NB	8.3	2.2	1.8	1.6
CH848.3.d1432.5.56D11gp120/293F	8.83	1.3	1.5	1.1	3.2	NB	NB	2.8	NB	2.8	1.9	2.6	1.4
CH848.3.d1621.4.15D11gp120/293F	8.805	1.3	1.2	1.2	5.9	NB	NB	7.8	NB	11.2	2.5	2.1	1.6
CH848.3.d0078.30.02D11gp120	8.763	1.3	0.9	1.2	4.1	NB	NB	3.8	NB	4.5	2.1	1.9	1.4
CH848.3.d1677.521.D11gp120/293F	8.713	1.2	1.2	1.1	2.5	NB	NB	3.4	NB	3.3	1.9	2.8	1.4
CH848.3.d1120.10.21D11gp120/293F/PEI	8.698	1.2	1.0	1.2	2.6	NB	NB	3.2	22.9	3.2	1.9	2.8	1.6
CH848.3.d0274.30.07D11gp120/293F	8.693	1.2	1.1	1.2	5.0	NB	NB	4.7	NB	5.1	2.3	2.1	1.7
CH0848d0949.10.18D11gp120/293F	8.648	1.1	1.1	1.1	1.6	9.1	9.7	1.9	2.8	2.5	1.4	1.2	1.4
CH848.3.d0526.25.32D11gp120	8.101	1.3	1.4	1.2	7.5	NB	NB	10.3	NB	19.2	2.8	2.6	2.0
CH0848d0526.25.26D11gp120/293F	8.083	1.3	1.0	1.2	6.1	NB	NB	6.9	NB	10.8	2.6	2.3	1.8
CH848.3.d1651.7.34D11gp120/293F	7.959	1.2	1.2	1.1	3.1	NB	NB	4.2	NB	4.9	2.1	3.9	1.5
CH848.3.d0135.60.32D11gp120/293F	7.94	1.3	1.3	1.3	6.9	NB	NB	7.9	NB	10.1	2.7	2.5	1.7
CH848.3.d0358.80.03D11gp120/293F	7.573	1.3	1.2	1.3	7.9	NB	NB	6.6	NB	11.8	2.6	1.9	1.7
CH0848d0780.15.22D11gp120/293F	7.567	1.2	1.0	1.2	2.8	5.9	8.8	2.7	4.4	3.2	1.8	2.0	1.6
CH848.3.d0526.25.39D11gp120	7.567	1.4	1.5	1.3	7.8	NB	NB	10.1	NB	15.7	2.9	2.3	1.9
CH848.3.d1621.4.31D11gp120/293F	7.39	1.4	1.2	1.1	3.8	NB	NB	7.1	NB	7.9	2.4	4.4	1.6
CH848.3.d0700.27.06D11gp120/293F	7.219	1.2	1.3	1.1	0.7	NB	NB	11.2	NB	26.1	3.1	1.3	2.6
CH848.3.d0135.60.34D11gp120/293F	6.983	1.5	1.4	1.3	6.8	NB	NB	8.6	NB	10.9	2.9	2.6	1.9
CH0848d0864.3.03D11gp120/293F	6.867	1.1	1.2	1.2	1.1	NB	NB	3.9	9.0	8.5	2.0	1.2	1.9
CH848.3.d0526.25.11D11gp120/293F	6.793	1.3	0.9	1.3	11.9	NB	NB	10.0	NB	15.7	2.7	4.7	2.0
CH848.3.d1621.4.12D11gp120/293F	6.676	1.8	1.5	1.3	5.4	NB	NB	48.4	NB	63.0	3.7	2.9	3.1
CH848.3.d1720.5D11gp120/293F	6.366	1.4	0.9	1.5	24.6	NB	NB	11.9	NB	14.7	3.2	5.2	2.3
CH848.3.d0274.30.14D11gp120/293F	6.191	1.5	1.1	1.5	16.6	NB	NB	23.5	NB	63.2	3.8	3.5	2.9
CH848.3.d0274.30.02D11gp120/293F	4.89	1.6	1.1	1.5	16.9	NB	NB	20.4	NB	26.3	4.2	3.6	3.0
CH848.3.d0445.25.18D11gp120	4.764	1.5	1.4	1.5	20.3	NB	NB	36.1	NB	NB	4.6	3.6	2.9
CH848.3.d0135.60.19D11gp120/293F	4.562	1.7	1.1	1.4	11.9	NB	NB	31.9	NB	40.4	3.7	3.9	2.8
CH0848d0700.15.05D11gp120/293F	4.512	1.4	1.3	1.4	12.6	NB	NB	NB	NB		5.7	5.1	3.1
CH0848d0949.10.10D11gp120/293F	3.471	1.2	1.1	1.3	1.4	NB	NB	6.6	11.1	46.3	2.9	1.8	2.4
CH848.3.d0445.30.41D11gp120	2.985	1.7	1.1	1.5	26.9	NB	NB	NB	NB		4.9	5.2	3.6
CH848.3.d0611.20.14D11gp120	2.949	1.7	1.1	1.3	2.1	NB	NB	18.4	NB	NB	4.0	2.7	2.4
CH848.3.d1120.10.13D11gp120/293F	2.528	1.1	1.1	1.1	1.4	NB	NB	6.5	18.5	19.0	2.4	1.5	2.1
CH848.3.d0194.25.24D11gp120/293F	2.23	2.2	1.1	1.9	NB	NB	NB	NB	NB		8.7	5.2	5.6
CH848.3.d0194.25.21D11gp120	2.192	1.7	2.5	1.6	NB	NB	NB	NB	NB		8.5	9.7	4.6
CH0848d0893.10.05D11gp120/293F	2.174	1.1	1.3	1.3	1.0	NB	NB	7.3	11.4	NB	3.0	1.8	2.4
CH848.3.d0135.60.14D11gp120/293F	1.464	2.1	1.0	2.1	NB	NB	NB	NB	NB		20.1	NB	8.3
CH848.3.d0194.25.48D11gp120/293F	0.584	6.3	1.1	3.2	NB	NB	NB	NB	NB	NB	NB	5.4	NB
CH848.3.d1305.10.35D11gp120/293F	0.092	NB	NB	0.4	0.1	NB	NB	NB	NB	NB	NB	NB	NB
CH0848d1432.5.35D11gp120/293F	0	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
CH848.3.d1305.10.13D11gp120/293F	0	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
CH848.3.d1305.10.21D11gp120/293F	0	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
CH848.3.d1305.10.30D11gp120/293F	0	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
CH848.3.d1432.5.50D11gp120	0	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
CH848.3.d1621.4.25D11gp120	0	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
CH848.3.d1621.4.46D11gp120	0	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB
CH848.3.d1651.7.50D11gp120/293F	0	NB	NB	NB		NB	NB	NB	NB	NB	NB	NB	NB
CH848.d1651.10.07D11gp120/293F	0	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB	NB

Legend:
	<1 fold
	1-2 fold
	2-5 fold
	5-10 fold
	10-25 fold
	25-50 fold
	>50 fold
NB	no binding

Supplementary Figure 3. Interactions of candidate gp120 immunogens with DH270min1 antibody variants.
ELISA binding of mutated DH270min1 antibodies with each acquired mutation changed to the corresponding UCA residue to a panel of gp120 immunogen candidates. Values represent fold decrease in binding of a given gp120 to a respective DH270min1 mutated antibody relative to its binding to the unmutated DH270min1 mAb. Highlighted molecules at the top of the table were considered for further analysis. Binding values (logAUC) to DH270min1 are shown as reference. NB=not binding.

Supplementary Figure 4. ELISA binding of candidate immunogen gp120s to DH270min1. The binding of DH270min1 was measured to immunogen candidates by ELISA suggests an affinity gradient exists amongst the immunogen candidates. This correlates with previous data that these viruses are progressively more difficult to neutralize described in *Supplementary Figure 6* and supports the use of these immunogens in sequential prime/boost strategy.

Supplementary Figure 5. Correlation between the amino acids selected from the scFv library by 10.17DT and the acquired mutations in the antibodies elicited by immunizations with 10.17DT. The same values as in *Figure 4D*, individually plotted. Negative library enrichment values are plotted as 0. The enrichment of mutations by 10.17DT vaccination in mice and the enrichment of yeast library clones selected by 10.17DT (y-axis) plotted against heavy chain mutation. Library enrichment values were calculated as described in Figure 4C. For immunization enrichment, both the data from individual animals (*gray, dashed*) and the calculated average (*blue*) is displayed.

Virus	V1 loop length	DH270.UCA	DH270.IA4	DH270.IA3	DH270.IA2	DH270.IA1	DH270.5	DH270.min11
CH848.0949.10.17	17	>50	0.64	0.2	0.14	<0.02	0.05	0.42
CH848.0358.80.06	24	>50	>50	>50	>50	0.02	<0.02	0.026
CH848.1432.5.41	30	>50	>50	>50	>50	0.33	0.21	<0.023
CH848.0526.25.02	34	>50	>50	>50	>50	1.42	1.56	0.026

IC50 ug/ml
<0.1
0.1 to <1
1 to <10
10 to 50
>50

Supplementary Figure 6. DH270min11 neutralization of pseudoviruses that containin the Envs used for the analysis of scFv libraries. Values indicate IC_{50} neutralization values of DH270 lineage precursor antibodies and DH270min11 mAb against pseudoviruses containing Envs that were used to develop the immunogens described in *Figures 3, 4 and 5.* DH270.IA4, DH270.IA3, DH270.IA2, DH270IA1 and DH270.5 are DH270 lineage members that are progressively closer in sequence to DH270.6 as previously described (Bonsignori et al., 2017). Neutralization data for these mAbs comes from (Bonsignori et al., 2017).

Supplementary Figure 7. Selection of DH270min11 libraries from *Figure 5* by FACS. A. FACS sorting work-flow and sequence analysis work-flow of the three libraries that tested all the possible amino acid combinations at the acquired mutation sites in DH270min11. **B.** Facsograms of libraries stained and sorted with the respective SOSIPs. Residues that interacted with the different glycan-V3 epitope regions (V3 loop, N301 glycan, and N332 glycan respectively) were sampled together in the three libraries. Double positive clones were selected, expanded and subjected to additional selection rounds as shown.

Probability of Selected Heavy Chain Residues						
D31$_{HC}$	F32$_{HC}$	M51$_{HC}$	T55$_{HC}$	R57$_{HC}$	T98$_{HC}$	Y110$_{HC}$
D=0.14351	F=0.05742	M=0.01914	T=0.04783	R=0.01415	T=0.02094	Y=0.00235
R=0.00536	C=0.0957	V= 0.04787	R=0.07251	A=0.04853	S=0.04702	R=0.01718
T=0.00125	R=0.00415	L=0.03062	G=0.04371	S=0.00659	E=0.00334	L=0.00127
P=0.00048	G=0.00159	R=0.00581	A=0.00209	T=0.00125	A=0.00269	Q=0.0001
	I=0.001	C=0.00019			P=0.00119	
	M=0.00006				L=0.00037	

Probability of Selected Light Chain Residues				
Y27$_{LC}$	Y48$_{LC}$	G94$_{LC}$	A97$_{LC}$	C101$_{LC}$
Y=0.00194	Y=0.00061	G=0.00498	A=0.00431	C=0.02239
R=0.0702	P=0.02995	V=0.00688	R=0.26982	L=0.17787
G=0.04776	A=0.00347	R=0.00307	G=0.06867	S=0.02623
N=0.04296	G=0.00161	H=0.00005	E=0.00113	V=0.01593
T=0.04036		W=0.00004		
P=0.00293				
A=0.00248				
L=0.00117				
E=0.00019				
Q=0.00012				

DH270.6
more probable than DH270.6
similar probability to DH270.6

Supplementary Figure 8. The predicted probability of DH270min11 alternate amino acids to occur *in vivo*. ARMADiLLO calculated probabilities of alternate amino acids identified at the key functional sites of DH270min11 by the library screening approach described in *Figure 5 and Supplementary Figure 7*. Residues present in the mature DH270.6 together with their expected probability to occur *in vivo* are shown in blue.