Medium-throughput zebrafish optogenetic platform identifies deficits in subsequent neural

activity following brief early exposure to cannabidiol and Δ-9-tetrahydrocannabinol

Richard Kanyo[1,2], Md Ruhul Amin[2], Laszlo F. Locskai[1,2], Danika D. Bouvier [2],

Alexandria M. Olthuis[2], W. Ted Allison[1,2,3], and Declan W. Ali [2, *]

[1] Centre for Prions & Protein Folding Disease, University of Alberta, Edmonton AB, T6G 2M8

[2] Department of Biological Sciences, University of Alberta, Edmonton AB, T6G 2E9

[3] Department of Medical Genetics, University of Alberta, Edmonton AB, T6G 2H7, Canada

23 * Author for correspondence (phone: (780) 492-6094; fax: (780) 492-9234; email:

24 declan.ali@ualberta.ca)

Abstract

In the light of legislative changes and the widespread use of cannabis as a recreational and medicinal drug, delayed effects of cannabis upon brief exposure during embryonic development are of high interest as early pregnancies often go undetected. Here, zebrafish embryos were exposed to cannabidiol (CBD) and Δ-9-tetrahydrocannabinol l (THC) until the end of gastrulation (1-10 hours post-fertilization) and analyzed later in development (4-5 days post-fertilization). In order to measure neural activity, we implemented CaMPARI (Calcium-Modulated Photoactivatable Ratiometric Integrator) and optimized the protocol for a 96-well format complemented by locomotor analysis. Our results revealed that neural activity was decreased by CBD more than THC. At higher doses, both cannabinoids could dramatically reduce neural activity and locomotor activity. Interestingly, the decrease was more pronounced when CBD and THC were combined. At the receptor level, CBD-mediated reduction of locomotor activity was partially prevented using cannabinoid type 1 and 2 receptor inhibitors. Overall, we report that CBD toxicity occurs via two cannabinoid receptors and is synergistically enhanced by THC exposure to negatively impact neural activity late in larval development. Future studies are warranted to reveal other cannabinoids and receptors involved in this pathway to understand the subsequent health implications of cannabis consumption on fetal development.

Introduction

Cannabis is consumed most commonly as a recreational drug that is often portrayed as harmless, yet the health implications are not fully understood[1]. The positive association of this drug in the public eye is not a surprise considering extracts of the *Cannabis sativa* plant have been used for medical purposes for almost 5000 years, especially for pain treatment[2]. However, scientific knowledge is limited and despite controversy, some countries, more recently Canada, continue to legalize cannabis for recreational use. Cannabis has been reported to be one of the most illicitly used drugs during pregnancy, with increase in consumption over the years, and key compounds, Δ-9-tetrahydrocannabinol (THC) and cannabidiol (CBD), can readily cross the placenta[3-8]. We were especially interested in CBD due to the reported health benefits and availability in many natural products including oils and food[9]. In the past, the negative impacts of cannabis in connection to embryonic development have been principally associated with the psychoactive THC[10]. In humans, epidemiological and clinical studies associate maternal cannabis exposure to behavioural disturbances in the offspring linked to increased risk for neuropsychiatric disorders[11]. In rats, maternal exposure of THC changed a series of behaviours in the offspring, including water-induced grooming, increased light sensitivity and altered exploratory behaviour[12]. Recently, the negative impacts of cannabis have expanded and also include the non-psychotropic CBD, which disrupts motor-neuron development in zebrafish[13]. This study is in contrast to reports that suggest positive health benefits of CBD, by treating nausea during pregnancy[14,15].

The mechanistic pathway(s) by which THC and CBD are toxic are elusive. A new study linked these two cannabinoids to the sonic hedgehog signaling pathway in mice and zebrafish, and appears to involve cannabinoid type 1 receptor (CB1R)[16]. It is well established that THC

binds and activates as a partial agonist two distinct classes of G-coupled protein receptors, CB1R

and CB2R[17]. Both receptors are highly expressed in neuronal tissue where CB1R is localized

mostly in the CNS[18-20] and CB2R in the peripheral nervous system[21], although CB2R has been

also linked to the CNS[22-24]. CBD has been reported to interact with CB1R and CB2R, but with a

lower affinity relative to THC, and rather antagonizes cannabinoid-induced effects indirectly

through other receptors[25-29]. Additional data to illuminate signaling pathways of THC and CBD

in vivo would highlight functional important mechanisms.

Zebrafish embryos have several advantages that complement mammalian models.

Experiments are economical because many embryos are available, and exposure studies are

simple as compounds can be easily added and removed. Further, embryos develop outside the

female and therefore not confounded by maternal physiology and variable transport to fetus.

Translucent larvae provide opportunities to implement cutting-edge fluorescing calcium sensors

and measure neural activity. Past studies suggest that the endocannabinoid system plays a role in

zebrafish development[30-32]. In addition, both cannabinoid receptors investigated here have

similar expression profiles in the CNS compared to mammals, with CB1R sharing a 70 % protein

sequence identity with the human homolog[33]. Therefore, the zebrafish is an excellent model

organism to further illuminate biological mechanisms. Considering the anticipated effects of

THC and CBD, high-throughput quantification of neural activity following early exposures

would be of great interest. In this study, the calcium modulated photoactivatable ratiometric

integrator (CaMPARI)[34], was implemented. Green fluorescing CaMPARI undergoes

photoconversion (PC) to a red fluorescing protein only in the presence of intense 405 nm light

and high Ca^{2+} levels. This conversion is irreversible, thus creating a temporal snapshot in a form

of a ratiometric red/green output. CaMPARI is expressed exclusively in neuronal tissue due to a

pan-neural promoter (*elavl3*, a.k.a. HuC) and is a direct read-out of relative neural activity.

Initially, this innovation was not suitable for a screening tool as the photoconversion was not

efficient enough. Thus, we have optimized the use of CaMPARI, such that the photoconversion

is more efficient and practical for high throughput well-plate-formats. This allows our novel *in*

vivo assay to measure neural activity and potentially screen large libraries of compounds. Our

results were supplemented with a behavioral assay obtained from the same larvae, which offers

an independent proxy measure that is related to neural activity.

Here, we investigated the effects of the two most abundant cannabinoids found in the

plant, THC and CBD, on neural activity and characterized CB1R and CB2R *in vivo* on the CBD

effects in zebrafish. The effects of early THC and CBD exposure from immediately after egg

fertilization until the end of gastrulation, at 10 hour post-fertilization (hpf), were monitored later

in embryonic development, at 4 and 5 days post-fertilization (dpf). The concentrations used in

this study mimic plasma levels of human subjects with high cannabis consumption[35-37], but we

do have to consider that the molecules have to pass from the waterbath through the chorion into

the plasma of the zebrafish embryo. THC and CBD were studied individually and then together

to understand the impact of individual compounds by themselves and to partially mimic the

cannabis available for recreational purposes. Further, we characterized the cannabinoid receptors

involved using established receptor inhibitors for CB1R and CB2R. This was important, because

with many receptors being linked to cannabinoids *in vitro*, additional functional *in vivo* evidence

would be of great value.

Results

High-throughput assessment of CaMPARI is a reliable metric of neural activity.

In this study the objective was to gain insight in the effects of THC and CBD on neural activity during early embryonic development. We measured neural activity with CaMPARI. CaMPARI photoconverts irreversibly from a green to a red fluorescing protein only if user-applied 405 nm light application coincides with high calcium levels (Fig. 1a)[34]. Red and green fluorescing CaMPARI are quantified as ratio (referred here as "CaMPARI activity"). We expressed CaMPARI through a pan-neural promoter, *elavl3* exclusively in the CNS and thus as a read-out of neural activity (Fig. 1b). Here, we optimized the use of CaMPARI zebrafish for an automated high-content INCell2000 plate-reader (Fig.1c). CaMPARI transgene expression was most suitable for this platform at 4 dpf, which is a suitable balance of fluorescing intensity and a sufficiently developed brain.

CaMPARI activity was quantified and displayed as a heatmap consisting of the red/green ratio in fluorescing intensities. At 4 dpf, optic tectum and hindbrain regions of larvae showed a clear measureable output, whereas without photoconversion CaMPARI activity was nearly zero. To establish a baseline of near-zero neural activity, larvae were anesthetized with MS-222, and a significant reduction ($p<0.01$) in red/green ratio was obtained compared to freely swimming larvae. In contrast, drug-induced neural activity, with the established convulsants pentylenetetrazole (PTZ) or 4-aminopyridine (4-AP), showed a consistent increase in CaMPARI activity, even at concentrations considered to be minimal when inducing seizure (Fig. 1d,e)[38].

We benchmarked our CaMPARI outputs against locomotor activity of zebrafish larvae, which is an established metric of neural activity[39]. We chose 5 dpf as an optimal developmental

timepoint for analyzing locomotor activity, because younger larvae are largely inactive. Larvae

were left in the same 96-well plate for assessing CaMPARI and locomotion. Exposing larvae to

photoconversion with LED light the day before had no impact on the mean locomotor activity

(Fig. 1f). Larvae anesthetized with MS-222 did not display any detectable swimbouts (Fig. 1f).

The level of neural activity correlated with locomotor activity and will be presented later.

Overall, these results show that CaMPARI, deployed in zebrafish larvae is a reliable high-

throughput tool for measuring neural activity.

CBD and THC reduce subsequent neural activity.

We sought to assess how embryonic exposure to cannabinoids impacts upon subsequent

neural activity later in development. Here, the doses of CBD and THC aligned with our previous

work[13], reflecting high cannabis consumption in humans. Comparisons of our dosage to humans

requires various considerations: (i) blood plasma concentrations of THC can peak up to 0.25

mg/l while smoking a single cigarette[35]; (ii) the content of THC has increased in the past 20

years; and (iii) doses of intraperitoneally administered medical CBD can vary greatly, from 5-

100 mg/kg, and daily maximum of 1500 mg/kg[36,37]. The current study uses up to 6 µg/mL of

THC and 3 µg/mL of CBD. Absorption studies using Liquid Chromatography-Tandem Mass

Spectrometry suggest that typically an estimated 0.1-10 % of toxic compounds will pass through

the chorion to reach the embryo[40,41].

We wanted to gain insight into the effects of CBD and THC separately (Fig. 2a), and

therefore each was applied in a dose-response format. This validation was necessary as

CaMPARI has not been used in studying cannabinoids. Compounds were added to the bath early

in animal development at (0.5 hpf) and then washed out towards the end of gastrulation (10 hpf).

CaMPARI was imaged at 4 dpf and locomotor activity was assessed at 5 dpf (Fig. 2b). Animals

that were exposed to CBD at concentrations of 2 µg/ml and 3 µg/ml exhibited a dose-dependent

reduction in CaMPARI activity (Fig. 2c,d). These reductions in CaMPARI output are substantial

when compared against the ~50 % reduction we observed in anesthetized larvae, where little

neural activity is expected. Coordinated with this, locomotion was also reduced starting at 1.5

µg/ml and was significant ($p<0.01$) at 3 µg/ml (Fig. 2e). THC had a similar effect as CBD and

also reduced neural activity at higher doses, 4 µg/ml and 6 µg/ml, (Fig. 2f,g). Locomotor activity

exhibited a more extensive reduction at 6 µg/ml (Fig. 2h). We compared neural activity and

locomotion in the same individuals exposed to effective doses of cannabinoids and found

significant correlations (Fig. S2; CBD r=0.52 ($p<0.01$) and THC r=0.71 ($p<0.01$)). Indeed, most

larvae, that displayed reduced neural activity also showed reduced locomotion. Together, our

findings show that both CBD and THC reduced neural activity when exposed early in

development.

Antagonistic effects of CBD on neural activity is enhanced when combined with sub-

effective doses of THC. Cannabis consumption during pregnancy exposes the fetus to THC and

CBD in concert. In order to investigate whether CBD and THC have a combined effect that is

different from either compound on its own, neural activity was measured when larvae were

exposed to both cannabinoids, using the same timeline as the previous experiment (Fig. 2b).

First, various sub-effective doses of CBD were applied from 0.5 to 1.5 µg/ml in the presence of 2

µg/ml THC (identified as sub-effective in Fig. 2). Preliminary dosage attempts revealed small,

but significant differences with $p<0.01$ and $p<0.05$, respectively, on neural activity when

combining 0.5 or 1.5 µg/ml CBD with 2.0 µg/ml THC (Fig. S3a,b). Locomotor activity also

trended towards a dose-dependent reduction during concerted application of CBD and THC and

was significant at 1.5 µg/ml with p<0.01 (Fig. S3c). Next, CBD and THC were added in a 1:1

ratio mixture at 1.0 and 2.0 µg/ml each. An enhanced antagonistic effect was revealed on neural

activity when both CBD and THC were added at 2.0 µg/ml. CBD by itself at 2.0 µg/ml

significantly (p<0.05) reduced neural activity (Fig. 3d,e) consistent with results in Fig. 2d, but

CBD and THC together, further (p<0.01) reduced neural activity compared to CBD or THC

alone (Fig. 3d,e). The combined effects of CBD and THC seemed synergistic in that CBD by

itself at 2.0 µg/ml, although significant, had a relative small effect and THC by itself at 2.0

µg/ml had no effect. However, when CBD and THC at the same concentration were combined

neural activity was reduced close to the mean of MS-222 anesthetized samples suggesting no to

very little real neural activity. Locomotor activity was mainly affected by CBD with no

additional reduction when compared to CBD and THC combined (Fig. 3f). No significant effect

on neural activity or locomotor activity was obtained when 1.0 µg/ml of CBD and THC was

applied (Fig. 3a-c). Due to the effect on locomotion, we also wanted to assess the integrity of

neurons in connection to motor axons. Reticulospinal neurons in the hindbrain were stained with

RMO44 antibody targeting NEFM, an established marker for neuronal damage[42,43]. Indeed,

RMO44 immunostaining was further reduced when CBD and THC were combined at 2.0 µg/ml

suggesting that both CBD and THC affect neuronal health (Fig. 4S). Together, these results

suggest that CBD and THC are more potent in reducing neural activity when applied in

combination.

CB1R and CB2R are both required for CBD-induced reduction of locomotor

activity. Some reports suggest that CBD mediates its actions through CB1- and CB2-

receptors[25]. To test whether these two receptors are involved in the neuronal toxicity mediated by

CBD *in vivo*, we deployed corresponding receptor inverse agonist AM251 and antagonist

AM630. This pharmacological approach allowed the temporal specificity required for our

experimental design. Both inhibitors appear to act with good specificity in zebrafish. Previous

works support the specificity of CB1R and CB2R inhibitors used in this study[44,45]. In connection

to CB2R, a recent study has shown that CB2 receptors are required for the action of AM630 on

zebrafish behaviour. The impact AM630 had on wildtype larvae was not detectable when applied

to *cnr2*[-/-] mutant larvae, supporting drug specificity. Moreover, AM630 treated zebrafish showed

very similar photo-dependent response compared to the CB2 knockout[44]. Similar in regards to

CB1R, because knocking out CB1R and inhibiting CB1R with AM251 caused a similar response

in rescueing locomotor activity when supressed with cannbinoid-receptor agonist, WIN55,212-

2[45]. All three metrics measured (acclimatization, dark-adaption and recovery) for locomotion

displayed arguably the same rescue-pattern in *cnr1*[-/-] mutant and AM251 treated larvae[45].

Further, both inhibtors were used at nanomolar range in our study, which is considered modest

and ensuring a level of specificity towards their respective receptors (AM251, IC50= 8 nM

towards CB1R; AM630, IC50= 31.2 nM towards CB2R). As described previously, the

experimental timeline is as depicted in figure 2b with CBD being added immediately after

AM251 and AM630. When analyzing CaMPARI activity, our results revealed that the CBD-

mediated reduction in neural activity could not be completely prevented with concentrations

from 0.1 to 10 nM AM251 or AM630 in the presence of 3 µg/ml of CBD (Fig. 4a-d). In

contrast, CBD-mediated reduction of locomotor activity was prevented with a concentration of

AM251 as low as 1 nM (Fig. 4e) or with 10 nM of AM630 (Fig. 4f). In the absence of CBD,

when using the maximum dose (10 nM) of AM251 or AM630, neither inhibitor had an effect

(Fig. 4g,h). This finding excludes the possibility that the effects of AM251 or AM630 on

locomotion were independent of CBD and rather specific towards CB1R and CB2R,

respectively. Overall, this suggests a role for both CB1R and CB2R receptors in CBD-mediated

effects on locomotor activity. In sum, inhibiting CB1R and CB2R can ameliorate the negative

impacts of early CBD on late CNS function, at least with respect to locomotion.

Discussion

In some jurisdictions, including Canada, cannabis has been legalized despite the lack of

mechanistic understanding and health implications. Cannabis consumption has increased in the

past decade[46,47] and reports emerged associating prenatal cannabis consumption with stillbirths

and autism spectrum disorder, a neurodevelopmental syndrome[48,49]. Yet, cannabis and CBD oil

have been used by consumers to treat labor pain[50]. In animal models, such as rats and chicken,

cannabis has been shown to have detrimental effects on developing embryos[10, 51]. Intriguingly,

more recently, new evidence revealed that exposure to not only the more widely studied THC,

but also CBD, during gastrulation impacts the overall development in zebrafish[13]. Interestingly,

the effect of the two key cannabinoids, CBD and THC, were delayed, and observed much later in

the development after the exposure had ended.

In vitro, CBD and THC can function through CB1R and CB2R, which are expressed in

neuronal tissue[17-25]. Our goal was to specifically investigate the effects of CBD, how it might

interact with the more widely studied THC during early development and characterize the impact

on neural activity *in vivo* on a mechanistic level. For our study, we implemented CaMPARI

transgenic zebrafish as a read-out of neural activity and supplemented this approach by

measuring the locomotor activity of the same larvae. CaMPARI has proven to be a simple and

economical neural integrator that can be optimized in a well-format using our animal model. We

found that neural activity was reduced later in zebrafish development upon brief exposure of not

only THC, but also CBD early in development. Surprisingly, CBD had similar impacts on neural

activity at lower concentrations compared to THC. Notably, the impact of CBD was increased by

sub-effective doses of THC. Our data quantifying locomotion, suggested that the CBD-mediated

effect involves CB1R and CB2Rs.

We optimized CaMPARI such that this innovation can be used for large number of

samples to screen compounds in future studies. Zebrafish larvae can be deployed in a 96-well

set-up quantifying locomotion[39,52-56]. Nevertheless, locomotor activity is not a direct

measurement of neural activity. Calcium sensors, such as the GCaMP series[57,58], the more

recently introduced GECO[59,60], and CaMPARI, are more accurate. CaMPARI provides a

temporal-snapshot of neural activity in the freely swimming animal, which can be anesthetized

and imaged after the experiment[34]. Furthermore, the ratiometric red/green signals are imaged

together and mitigate inter-individual differences in expression levels or focal plane. We

optimized conditions for a well-plate set-up analyzing neural activity in the optic tectum and

hindbrain. Unfortunately, less defined areas or specific neurons could not be quantified. Effects

on motor axons will be detected by our behavioural assay. In general, there seems to be a good

correlation between neural activity measured by CaMPARI and locomotor activity[61,62]. However,

some findings may be more sensitive towards one assay vs the other, as described above.

Furthermore, when interpreting our data as neural activity, it is important to consider that when

anesthetized a clear red to green CaMPARI ratio can be still measured. This is unlikely due to

autofluoresence because larvae that were not photoconverted yielded zero CaMPARI activity.

We cannot exclude the possibility that due to longer photoconversion some CaMPARI has

converted into the red fluorescence version even when anesthetized. It is well established that

when anesthetized with MS-222, neurons are barely active[63,64]. Indeed, our behavioural assay

shows no locomotion in the presence of MS-222 and we used this assay as a supplement in this

study. We believe this method is a suitable technique to measure substantial changes in neural

activity in the larger parts of the zebrafish CNS. For future studies in connection to epilepsy this

innovation may be particularly helpful as some studies use convulstants, PTZ or 4-AP, to

simulate seizure[62,65]. We found that both, PTZ and 4-AP, induce CaMPARI activity. Considering

the advantages, this method outweighs its limitation and could serve for future studies, especially

in connection with epilepsy.

We have considered several competing technologies prior to designing this study, in

hopes that we can build upon and complement these potent approaches. Other than behavioural

assays described above, the GCaMP series has been of interest[57,58]. However, the larvae have to

be analyzed with high-resolution live microscopy during the experiment and movement of the

larvae would add considerable variability. The assays would be data-intensive and technically

challenging, where large sample sizes and different experimental groups are difficult to obtain

(especially *in vivo*). Additionally, we were still interested in measuring outputs from freely

behaving animals as constraining them for imaging likely impacts the results and is time-

consuming. Other widely used markers for neural activity include the immediate-early gene *c-fos*

and phosphorylation of ERK[65,66]. However, CaMPARI is more temporal specific due to

immediate photoconversion. In addition, both markers are not only less tissue specific, but also

respond to stress and other factors. The *elavl3* promoter that drives CaMPARI ensures CNS-

specific expression. Overall, CaMPARI is a practical direct output of neural activity and thus the

chosen innovation for this study.

Both CBD and THC have been implicated in embryonic development including abnormal

changes to neurons. Delayed abnormalities after early exposure of key cannabinoids have only

recently been investigated[13]. Here, early exposure of CBD and THC during the first 10 hours of

embryonic development reduced neural activity when measured 4 days later, in a dose-dependent

manner. These later effects may be due to abnormal neurodevelopment and/or the slow release of

the cannabinoids from the tissue due to their lipophilic nature. Consistent with this finding,

zebrafish larvae with a similar exposure (5-10 hpf) to cannabinoids have shown effects that could

be due abnormal development (at 5 dpf). This was not exclusive to neurons, but included other

changes such as morphological abnormalities, decreased survival rate, decreased mEPCs activity

(may relate to muscle development), decreased heart rate and delayed hatching[13]. Considering

the specific time-window of exposure (first 10 hours in development), an abrupt effect on the

developmental program including neurons seems plausible. Interestingly, CBD seemed to impact

neural activity at a lower concentration than THC, which also aligns with our previous study[13]. It

is important to acknowledge that CBD consumption has been documented to help with seizures

and cancer by reducing pain[67,68]. Nevertheless, our study suggests caution during pregnancies.

Considering that cannabis contains large proportions of both CBD and THC, we looked

at the effects of these two compounds in combination and found a more potent effect in

decreasing neural activity. Reticulospinal neurons in the hindbrain of the zebrafish control motor

neurons during swimming[42,69]. In line with a reduction in locomotion, we observed reduced

staining when exposed to both cannabinoids, suggesting that the reticulospinal neuronal integrity

was compromised. However, this finding did not explain why locomotor activity was mainly

influenced by CBD. This could be due in part that CBD has a greater effect at lower

concentration than THC. The effect of CBD may be amplified by effecting motor neurons and

potentially muscle development. Overall, these findings were surprising, because CBD was

found to negate the effects of THC on CB1R[27]. One possibility is that at higher concentrations

both, CBD and THC, affect the developmental program. This could explain the abnormal

morphologies reported previously[13]. We have observed similar morphologies using higher

concentrations of CBD and/or THC. Thus, it is of no surprise that abnormal morphologies

impacted neural activity and locomotion. Nevertheless, because at 2 µg/mL we found a

significant reduction in neural activity, but no morphological change was observed previously at

that concentration[13], it is unlikely that neural activity is affected by morphological changes alone.

In short, we believe that although in physiological range, high concentrations of CBD and THC

during embryonic development disrupt proper growth and neuron function with similar

consequences and thus the effects of CBD and THC are same. This could result in overriding the

negating effects of CBD. In contrast, at lower concentrations and in fully developed brains the

impact of CBD and THC is different. Indeed, several receptors have been shown to have

different affinities to THC and CBD[25]. A second possibility, therefore, is that at high

concentration CBD and THC affect receptors other than CB1R and CB2R. This could explain

why inhibiting CB1R and CB2R did not fully prevent CBD-mediated reduction of neural

activity, as further discussed below.

At this stage, it is unclear what mechanism(s) account for the reduction in neural activity

later in development when embryos are exposed briefly to CBD and THC in the first hours after

fertilization. In this study, we were especially interested in the less widely studied CBD. Using

HEK cells several receptor candidates have been identified showing to interact with CBD[25]. A

recent study supports a functional role of CB1R and CB2R in zebrafish[32]. Here, by inhibiting

CB1R, and CB2R, we found that the reduction in locomotor activity mediated by CBD was

partially prevented at nanomolar concentrations of the inhibitors. This suggests that,

mechanistically, CBD acts through both receptors. Previous studies have implied that CBD may

bind to both CB1R and CB2R *in vitro*[25] and our study is in support that these interactions may be

important *in vivo*. In line with our findings, CBD has been shown to cause abnormal craniofacial

and brain development in mice and zebrafish, which seems to be mediated through CB1R and

the hedgehog signaling pathway[16]. Whether CB2R can mediate its effects through the same

pathway remains to be determined. An explanation to why inhibiting CB1R and CB2R did

prevent the maximum effect of CBD, which is delayed, could relate to the fact that the CB1R

and CB2R are expressed very early in development and found on stem cell progenitor cells[61]. As

mentioned above, an early effect may impact the developmental program such that changes

persist until later, even if cannabinoids are no longer present in the circulation. Such changes

may not be specific to the CNS as CB receptors are not only expressed in neuronal progenitor

cells, but in stem cells forming other types of tissues[70]. This could explain our previous findings

where CBD affected heart-rate and overall morphology after 5 dpf when exposed only briefly

during gastrulation[11]. The CBD-mediated reduction in neural activity was not prevented when

either CB1 or CB2Rs were inhibited. While the reason for this is unclear, it could relate that the

effects of CBD are larger outside the optic tectum and the hindbrain, thus more visible when

quantifying locomotion. As described above, our CaMPARI assay in well-format is not suitable

to resolve motor axons in detail, which have been shown to be affected previously[13]. It is also

possible that other receptors such as TRPV1 are involved[26], but this needs further investigation.

Overall, both CBD and THC exposure during embryonic stages had a negative impact on

later neural activity, which was additive when combined. Together, this could mean that the

impact of cannabis on early development is higher than using isolated compounds and profound.

Our findings also support an *in vivo* mechanism of CBD functioning through CB1R and CB2R.

We believe this study opens up a new path for investigating detailed mechanisms, including

other receptors, in specific neurons that are affected by cannabinoid toxicity early in

development and persist. The accumulating evidence for prenatal cannabis consumption having

negative consequences on neurodevelopmental disorders in the offspring[3] has to be considered.

As the recreational use is becoming legalized and socially acceptable, more studies are warranted

to not only fully understand the impact on human development, but also the mechanisms of the

diseases.

METHODS

Animal Ethics and exposure to drugs

Zebrafish maintenance was approved by the Animal Care & Use Committee: Biosciences

at the University of Alberta and operated under the guidelines of the Canadian Council of

Animal Care. The fish were maintained in the University of Alberta fish facility at 28°C under a

14/10 light/dark cycle as previously described[71] and all authors complied with ARRIVE (Animal

Research: Reporting of In Vivo Experiments) guidelines[72].

CaMPARI transgenic zebrafish were outcrossed with *Casper* strains not carrying the

transgene. For experimental set up, embryos were randomly collected and placed in egg water

(60 µg/ml Instant Ocean) in groups of 20 embryos as early as practically possible within 0.5 hpf.

Stock solutions of CBD and THC were obtained from Sigma at 1 mg/ml dissolved in methanol.

Receptor inhibitors AM251 (Selleck Chemicals, Houston, TX, USA) and AM630 (Adooq

Bioscience, Irvine, CA, USA) were dissolved in DMSO. Accordingly, zebrafish embryos were

treated with vehicle or compounds as close to 1 hpf as practically possible and removed towards

the end of gastrulation (10 hpf). Within experiments all treatments contained the same amount of

methanol or DMSO, specific percentages are indicated in the figure legends. Compounds were

removed with several washes and egg water was replaced every day until CaMPARI analysis.

Not all larvae carried the transgene and were screened for green fluorescence. An increase in

mortality and abnormal morphology was observed when treated with higher doses of CBD or

THC, as reported previously[13].

Engineering CaMPARI transgenic zebrafish for integrative calcium imaging.

Zebrafish expressing CaMPARI, i.e. the Tg[*elavl3*:CaMPARI (W391F+V398L)][ua3144]

line were generated as described previously[61,62]. We re-derived CaMPARI fish in response to a

federal moratorium limiting zebrafish import into Canada[73].

CaMPARI photoconversion

For photoconversion, bright green Tg[*elavl3*:CaMPARI (W391F+V398L)][ua3144] larvae at

4 dpf were placed in a 96-well plate containing 150 µl egg water (made as previously decribed[71])

and acclimatized for 2 h. The central 48 wells of the plate were exposed to 405 nm LED array

(Loctite) for 300 sec at a distance of 10 cm. LED array illuminated the plates entirely and evenly.

The plates were floating in water at room temperature to ensure that the larvae did not overheat.

For imaging, the larvae were anesthetized in 0.24 mg/mL tricaine methanesulfonate (MS-222,

Sigma) following photoconversion. MS-222 was washed out and zebrafish were rested for a day

prior analysis for locomotor activity.

CaMPARI analysis

The ratio of red to green fluorescent emissions (red photoconverted CaMPARI in ratio to

green CAMPARI) was interpreted as relative neural activity, as previously defined[34]. Images

were objectively obtained with an InCell 2000 microscope (GE Healthcare, US). 96-well plates

were run using FITC (1 second exposure) and dsRed (2 second exposure) channels. Blinded

experimentalists used ImageJ to quantify the fluorescence mean of the red (dsRed) and green

(FITC) channels. The area of the optic and hindbrain area were quantified as those regions were

most consistent in the view. A mask was applied to exclude noise outside of the CNS.

Quantifying Locomotor activity

The same zebrafish larvae being placed in the 96-well plate the day prior for CaMPARI

analysis were objectively quantified for locomotion using Basler GenICaM (Basler acA 1300-

60) scanning camera (75-mm f2.8 C-mount lens) and EthoVision® XT-11.5 software by Noldus

(Wageningen, Netherlands) as described previously[74]. Briefly, larvae were acclimatized for 3 h

and one hour of movement was recorded from each individual larvae in each well. Beneath the

plate an infrared backlight source was located and the scanning camera above. Activity was

defined here as % pixel change within well between frames (recordings were at 25 frames per

second) and quantified with EthoVision® XT-11.5 software. This set-up allowed to see whether

there is a correlation between level of neural activity and the the mean activity % within a

swimbout (swimbout size). Swimbout size was shown to increase in response to the convulsant

PTZ previously[52] and here it can be a useful marker for neural activity. Transparent *Casper* strain

zebrafish[75] were unable to be tracked as ususal with Noldus. Therefore, quantification of

swimbouts provided an additional advantage. Clear movement was detected above the noise

treshold due to movement of the pigmented eyes. Although the % swimbout is small, reliable

measurements were obtained. Only clear swimbouts were quantified, which were set above a

treshold of 0.1 % of activity. Figure 1f shows that measuring the mean activity of a swimbout is

a useful tool in obtaining output that is clearly diminished in presence of MS-222. In addition,

the output also shows sensitivity in freely swimming larvae.

RMO44 staining

The zebrafish used in this experiment were of the Tuebingen Longfin strain. Set-up was

similar as described above. Following the experiment, larvae were stained with anti-RMO44

(Thermo Fisher Scientific, Waltham, MA, USA) and imaged using a Zeiss LSM 710 confocal

microscope (Oberkochen, Germany) as described elsewhere[76].

Statistics

GraphPad Prism Software (Version 7, GraphPad, San Diego, CA) was used to analyze statistics

from data obtained from our CaMPARI assay or tracking locomotor activity. All the data were

presented as mean ±SEM (standard error of the mean). Statistical significance of $p < 0.05$ was

determined using a non-parametric t-test between two groups followed by a Mann-Whitney

analysis, if approriate. Multiple groups were compared using One-Way ANOVA with Dunnett's

multiple comparisons test. Pearson r from correlation analysis was also determined where

indicated.

References

1. Degenhardt L, Chiu W-T, Sampson N, Kessler RC, Anthony JC, Angermeyer M, et al. Toward a global view of alcohol, tobacco, cannabis, and cocaine use: findings from the WHO World Mental Health Surveys. PLoS Med. 2008;5:e141.

2. Winek CL. Some historical aspects of marijuana. Clin Toxicol. 1977;10:243–53.

3. Bérard A. The importance of generating more data on cannabis use in pregnancy. Nat Med. 2020;26:1515–6.

4. Fergusson DM, Horwood LJ, Northstone K, ALSPAC Study Team. Avon Longitudinal Study of Pregnancy and Childhood. Maternal use of cannabis and pregnancy outcome. BJOG. 2002;109:21–7.

5. Gómez M, Hernández M, Johansson B, de Miguel R, Ramos JA, Fernández-Ruiz J. Prenatal cannabinoid and gene expression for neural adhesion molecule L1 in the fetal rat brain. Brain Res Dev Brain Res. 2003;147:201–7.

6. Hutchings DE, Martin BR, Gamagaris Z, Miller N, Fico T. Plasma concentrations of delta-9-tetrahydrocannabinol in dams and fetuses following acute or multiple prenatal dosing in rats. Life Sci. 1989;44:697–701.

7. Young-Wolff KC, Tucker L-Y, Alexeeff S, Armstrong MA, Conway A, Weisner C, et al. Trends in Self-reported and Biochemically Tested Marijuana Use Among Pregnant Females in California From 2009-2016. JAMA. 2017;318:2490.

8. van Gelder MMHJ, Reefhuis J, Caton AR, Werler MM, Druschel CM, Roeleveld N. Characteristics of pregnant illicit drug users and associations between cannabis use and perinatal outcome in a population-based study. Drug and Alcohol Dependence. 2010;109:243–7.

9. Singer L, Tokish H, Park F, Campisi C, Milanaik RL. The cannabidiol conundrum: potential benefits and risks of cannabidiol products for children. Curr Opin Pediatr. 2020;32:198–205.

10. Friedrich J, Khatib D, Parsa K, Santopietro A, Gallicano GI. The grass isn't always greener: The effects of cannabis on embryological development. BMC Pharmacol Toxicol. 2016 Sep 29;17:45.

11. Morris CV, DiNieri JA, Szutorisz H, Hurd YL. Molecular mechanisms of maternal cannabis and cigarette use on human neurodevelopment. Eur J Neurosci. 2011;34:1574–83.

12. Navarro M, Rubio P, de Fonseca FR. Behavioural consequences of maternal exposure to natural cannabinoids in rats. Psychopharmacology (Berl). 1995;122:1–14.

539 13. Ahmed KT, Amin MR, Shah P, Ali DW. Motor neuron development in zebrafish is altered
540   by brief (5-hr) exposures to THC (Δ9-tetrahydrocannabinol) or CBD (cannabidiol) during
541   gastrulation. Sci Rep. 2018;8:10518.
542
543 14. Parker LA, Rock EM, Limebeer CL. Regulation of nausea and vomiting by cannabinoids. Br
544   J Pharmacol. 2011;163:1411–22.
545
546 15. Roberson EK, Patrick WK, Hurwitz EL. Marijuana use and maternal experiences of severe
547   nausea during pregnancy in Hawai'i. Hawaii J Med Public Health. 2014;73:283–7.
548
549 16. Fish EW, Murdaugh LB, Zhang C, Boschen KE, Boa-Amponsem O, Mendoza-Romero HN,
550   et al. Cannabinoids Exacerbate Alcohol Teratogenesis by a CB1-Hedgehog Interaction. Sci
551   Rep. 2019;9:16057.
552
553 17. Pertwee RG. Ligands that target cannabinoid receptors in the brain: from THC to
554   anandamide and beyond. Addict Biol. 2008;13:147–59.
555
556 18. Herkenham M, Lynn AB, Little MD, Johnson MR, Melvin LS, de Costa BR, et al.
557   Cannabinoid receptor localization in brain. Proc Natl Acad Sci U S A. 1990;87:1932–6.
558
559 19. Watson S, Chambers D, Hobbs C, Doherty P, Graham A. The endocannabinoid receptor,
560   CB1, is required for normal axonal growth and fasciculation. Mol Cell Neurosci.
561   2008;38:89–97.
562
563 20. Kano M, Ohno-Shosaku T, Hashimotodani Y, Uchigashima M, Watanabe M.
564   Endocannabinoid-mediated control of synaptic transmission. Physiol Rev. 2009;89:309–80.
565
566 21. Smita K, Sushil Kumar V, Premendran JS. Anandamide: an update. Fundam Clin Pharmacol.
567   2007;21:1–8.
568
569 22. Benarroch E. Endocannabinoids in basal ganglia circuits: implications for Parkinson disease.
570   Neurology. 2007;69:306–9.
571
572 23. Stempel AV, Stumpf A, Zhang H-Y, Özdoğan T, Pannasch U, Theis A-K, et al. Cannabinoid
573   Type 2 Receptors Mediate a Cell Type-Specific Plasticity in the Hippocampus. Neuron.
574   2016;90:795–809.
575
576 24. Liu Q-R, Canseco-Alba A, Zhang H-Y, Tagliaferro P, Chung M, Dennis E, et al.
577   Cannabinoid type 2 receptors in dopamine neurons inhibits psychomotor behaviors, alters
578   anxiety, depression and alcohol preference. Sci Rep. 2017;7:17410.
579
580 25. McPartland JM, Duncan M, Di Marzo V, Pertwee RG. Are cannabidiol and Δ(9) -
581   tetrahydrocannabivarin negative modulators of the endocannabinoid system? A systematic
582   review. Br J Pharmacol. 2015;172:737–53.
583

26. Gray RA, Whalley BJ. The proposed mechanisms of action of CBD in epilepsy. Epileptic
Disorders. 2020;22:S10–5.

27. Laprairie RB, Bagher AM, Kelly MEM, Denovan☐Wright EM. Cannabidiol is a negative
allosteric modulator of the cannabinoid CB1 receptor. Br J Pharmacol. 2015;172:4790–805.

28. Ibeas Bih C, Chen T, Nunn AVW, Bazelot M, Dallas M, Whalley BJ. Molecular Targets of
Cannabidiol in Neurological Disorders. Neurotherapeutics. 2015;12:699–730.

29. Ryberg E, Larsson N, Sjögren S, Hjorth S, Hermansson N-O, Leonova J, et al. The orphan
receptor GPR55 is a novel cannabinoid receptor. Br J Pharmacol. 2007;152:1092–101.

30. Akhtar MT, Ali S, Rashidi H, van der Kooy F, Verpoorte R, Richardson MK. Developmental
effects of cannabinoids on zebrafish larvae. Zebrafish. 2013;10:283–93.

31. Oltrabella F, Melgoza A, Nguyen B, Guo S. Role of the endocannabinoid system in
vertebrates: Emphasis on the zebrafish model. Dev Growth Differ. 2017;59:194–210.

32. Sufian MS, Amin MR, Kanyo R, Allison WT, Ali DW. CB1 and CB2 receptors play
differential roles in early zebrafish locomotor development. J Exp Biol. 2019;222(Pt 16).

33. Lam CS, Rastegar S, Strähle U. Distribution of cannabinoid receptor 1 in the CNS of
zebrafish. Neuroscience. 2006;138:83–95.

34. Fosque BF, Sun Y, Dana H, Yang C-T, Ohyama T, Tadross MR, et al. Neural circuits.
Labeling of active neural circuits in vivo with designed calcium integrators. Science.
2015;347:755–60.

35. Huestis MA. Human cannabinoid pharmacokinetics. Chem Biodivers. 2007;4:1770–804.

36. Rohleder C, Müller JK, Lange B, Leweke FM. Cannabidiol as a Potential New Type of an
Antipsychotic. A Critical Review of the Evidence. Front Pharmacol. 2016;7:422.

37. Iffland K, Grotenhermen F. An Update on Safety and Side Effects of Cannabidiol: A Review
of Clinical Data and Relevant Animal Studies. Cannabis Cannabinoid Res. 2017;2:139

38. Ellis LD, Seibert J, Soanes KH. Distinct models of induced hyperactivity in zebrafish larvae.
Brain Research. 2012;1449:46–59.

39. Rihel J, Prober DA, Arvanites A, Lam K, Zimmerman S, Jang S, et al. Zebrafish behavioral
profiling links drugs to biological targets and rest/wake regulation. Science. 2010;327:348–
51.

40. Zhang F, Qin W, Zhang J-P, Hu C-Q. Antibiotic Toxicity and Absorption in Zebrafish Using
Liquid Chromatography-Tandem Mass Spectrometry. PLOS ONE. 2015;10:e0124805.

41. Brox S, Ritter AP, Küster E, Reemtsma T. A quantitative HPLC–MS/MS method for
studying internal concentrations and toxicokinetics of 34 polar analytes in zebrafish (Danio
rerio) embryos. Anal Bioanal Chem. 2014;406:4831–40.

42. Banote RK, Edling M, Eliassen F, Kettunen P, Zetterberg H, Abramsson A. β-Amyloid
precursor protein-b is essential for Mauthner cell development in the zebrafish in a Notch-
dependent manner. Developmental Biology. 2016;413:26–38.

43. Braissant O, Henry H, Villard A-M, Zurich M-G, Loup M, Eilers B, et al. Ammonium-
Induced Impairment of Axonal Growth Is Prevented through Glial Creatine. J Neurosci.
2002;22:9810–20.

44. Acevedo-Canabal A, Colón-Cruz L, Rodriguez-Morales R, Varshney GK, Burgess S,
González-Sepúlveda L, et al. Altered Swimming Behaviors in Zebrafish Larvae Lacking
Cannabinoid Receptor 2. Cannabis and Cannabinoid Research. 2019;4:88–101.

45. Luchtenburg FJ, Schaaf MJM, Richardson MK. Functional characterization of the
cannabinoid receptors 1 and 2 in zebrafish larvae using behavioral analysis.
Psychopharmacology (Berl). 2019;236:2049–58.

46. Canada Health. Canadian Tobacco Alcohol and Drugs (CTADS) Survey: 2017 summary
2018 https://www.canada.ca/en/health-canada/services/canadian-tobacco-alcohol-drugs-
survey/2017-summary.html#n3 Accessed 30 November 2020

47. Statistics Canada. National Cannabis Survey, first quarter 2019. 2019;(11):15.

48. Varner MW, Silver RM, Rowland Hogue CJ, Willinger M, Parker CB, Thorsten VR, et al.
Association between stillbirth and illicit drug use and smoking during pregnancy. Obstet
Gynecol. 2014;123:113–25.

49. Corsi DJ, Donelle J, Sucha E, Hawken S, Hsu H, El-Chaâr D, et al. Maternal cannabis use in
pregnancy and child neurodevelopmental outcomes. Nat Med. 2020;26:1536–40.

50. Sarrafpour S, Urits I, Powell J, Nguyen D, Callan J, Orhurhu V, et al. Considerations and
Implications of Cannabidiol Use During Pregnancy. Curr Pain Headache Rep. 2020;24:38.

51. Gustafsson SB, Jacobsson SOP. Effects of cannabinoids on the development of chick
embryos in ovo. Scientific Reports. 2019;9:13486.

52. Hoffman EJ, Turner KJ, Fernandez JM, Cifuentes D, Ghosh M, Ijaz S, et al. Estrogens
Suppress a Behavioral Phenotype in Zebrafish Mutants of the Autism Risk Gene, CNTNAP2.
Neuron. 2016;89:725–33.

53. MacRae CA, Peterson RT. Zebrafish as tools for drug discovery. Nat Rev Drug Discov.
2015;14:721–31.
675

676 54. Baraban SC, Dinday MT, Hortopan GA. Drug screening in Scn1a zebrafish mutant identifies
677   clemizole as a potential Dravet syndrome treatment. Nature Communications. 2013;4:2410.
678

679 55. Bose P, Tremblay E, Maois C, Narasimhan V, Armstrong GAB, Liao M, et al. The Novel
680   Small Molecule TRVA242 Stabilizes Neuromuscular Junction Defects in Multiple Animal
681   Models of Amyotrophic Lateral Sclerosis. Neurotherapeutics. 2019;16:1149–66.
682

683 56. Ibhazehiebo K, Gavrilovici C, de la Hoz CL, Ma S-C, Rehak R, Kaushik G, et al. A novel
684   metabolism-based phenotypic drug discovery platform in zebrafish uncovers HDACs 1 and 3
685   as a potential combined anti-seizure drug target. Brain. 2018;141:744–61.
686

687 57. Mank M, Griesbeck O. Genetically Encoded Calcium Indicators. Chem Rev.
688   2008;108:1550–64.
689

690 58. Akerboom J, Chen T-W, Wardill TJ, Tian L, Marvin JS, Mutlu S, et al. Optimization of a
691   GCaMP calcium indicator for neural activity imaging. J Neurosci. 2012;32:13819–40.
692

693 59. Shaner NC, Lambert GG, Chammas A, Ni Y, Cranfill PJ, Baird MA, et al. A bright
694   monomeric green fluorescent protein derived from Branchiostoma lanceolatum. Nat
695   Methods. 2013;10:407–9.
696

697 60. Zarowny L, Aggarwal A, Rutten VMS, Kolb I, GENIE Project, Patel R, et al. Bright and
698   High-Performance Genetically Encoded Ca2+ Indicator Based on mNeonGreen Fluorescent
699   Protein. ACS Sens. 2020;5:1959–68.
700

701 61. Alyenbaawi H, Kanyo R, Locskai LF, Kamali-Jamil R, DuVal MG, Bai Q, et al. Seizures are
702   a druggable mechanistic link between TBI and subsequent tauopathy. bioRxiv. 14 May 2020
703   (accepted in eLife). https://doi.org/10.1101/2020.05.12.091819.
704

705 62. Kanyo R, Wang CK, Locskai LF, Li J, Allison WT, Kurata HT. Functional and behavioral
706   signatures of Kv7 activator drug subtypes. Epilepsia. 2020;61:1678-1690
707

708 63. Medler S. Anesthetic MS-222 eliminates nerve and muscle activity in frogs used for
709   physiology teaching laboratories. Adv Physiol Educ. 2019;43:69–75.
710

711 64. Hedrick MS, Winmill RE. Excitatory and inhibitory effects of tricaine (MS-222) on fictive
712   breathing in isolated bullfrog brain stem. American Journal of Physiology-Regulatory,
713   Integrative and Comparative Physiology. 2003;284:R405–12.
714

715 65. Leighton PLA, Kanyo R, Neil GJ, Pollock NM, Allison WT. Prion gene paralogs are
716   dispensable for early zebrafish development and have nonadditive roles in seizure
717   susceptibility. J Biol Chem. 2018;293:12576–92.

718
719 66. Özcan, Güliz Gürel, Sumi Lim, Patricia LA Leighton, W. Ted Allison, and Jason Rihel.
720   "Sleep Is Bi-Directionally Modified by Amyloid Beta Oligomers." ELife 9 (14 2020).
721   https://doi.org/10.7554/eLife.53995.
722
723 67. Perucca E. Cannabinoids in the Treatment of Epilepsy: Hard Evidence at Last? J Epilepsy
724   Res. 2017;7:61–76.
725
726 68. Russo EB. Cannabinoids in the management of difficult to treat pain. Ther Clin Risk Manag.
727   2008;4:245–59.
728
729 69. Kohashi T, Nakata N, Oda Y. Effective Sensory Modality Activating an Escape Triggering
730   Neuron Switches during Early Development in Zebrafish. J Neurosci. 2012;32:5810–20.
731
732 70. Galve-Roperh I, Chiurchiù V, Díaz-Alonso J, Bari M, Guzmán M, Maccarrone M.
733   Cannabinoid receptor signaling in progenitor/stem cell proliferation and differentiation.
734   Progress in Lipid Research. 2013;52:633–50.
735
736 71. Westerfield, M. The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio
737   rerio). University of Oregon Press:Eugene, OR ;2000.
738
739 72. Sert NP du, Hurst V, Ahluwalia A, Alam S, Avey MT, Baker M, et al. The ARRIVE
740   guidelines 2.0: Updated guidelines for reporting animal research. PLOS Biology.
741   2020;18:e3000410.
742
743 73. Hanwell D, Hutchinson SA, Collymore C, Bruce AE, Louis R, Ghalami A, et al. Restrictions
744   on the Importation of Zebrafish into Canada Associated with Spring Viremia of Carp Virus.
745   Zebrafish. 2016;13 Suppl 1:S153-163.
746
747 74. Kanyo R, Leighton PLA, Neil GJ, Locskai LF, Allison WT. Amyloid-β precursor protein
748   mutant zebrafish exhibit seizure susceptibility that depends on prion protein. Exp Neurol.
749   2020;328:113283.
750
751 75. White RM, Sessa A, Burke C, Bowman T, LeBlanc J, Ceol C, et al. Transparent adult
752   zebrafish as a tool for in vivo transplantation analysis. Cell Stem Cell. 2008;2:183–9.
753
754 76. Amin MR, Ahmed KT, Ali DW. Early Exposure to THC Alters M-Cell Development in
755   Zebrafish Embryos. Biomedicines. 4 January 2020;8.
756   https://doi.org/10.3390/biomedicines8010005
757
758
759
760

ACKNOWLEDGEMENTS

This study was supported by grants to W.T. Allison from Alberta Prion Research Institute-Alberta Innovates BioSolutions and the Alzheimer Society of Alberta and the Northwest Territories, and to D.W. Ali from the Natural Sciences and Engineering Research Council of Canada and Alberta Innovates mCannabis. L.F. Locskai was supported by undergraduate studentships from Natural Sciences and Engineering Research Council of Canada and from Alberta Innovates Health Solutions. R. Kanyo was supported by a SynAD postdoctoral fellowship funded via Alzheimer Society of Alberta and Northwest Territories through their Hope for Tomorrow program and the University Hospital Foundation. We would like to also extend our thanks to members of the Allison and Ali lab. In addition, we would also like to acknowledge the Sciences Animal Support Services at the Unversity of Alberta.

AUTHOR'S CONTRIBUTIONS

RK, MRA, WTA, DWA contributed to the experimental design. RK, MRA, LFL, DDB, AMO, performed the experiments and analyzed the data. RK, MRA and DWA interpreted the results. RK, WTA and DWA wrote the manuscript. All authors read and approved the manuscript.

Additional Information

Competing Interests: The authors declare no competing interests.

Figure 1: High-throughput quantification of neural activity in freely-swimming zebrafish

larvae. (**a**) CaMPARI photoconverts from green to red fluorescing versions in the neuron only in

the presence of both high intracellular calcium concentrations and a bright 405 nm light source.

(**b**) Lateral view of green fluorescing CaMPARI merged with brightfield image shows exclusive

expression in the CNS due to a pan-neural promoter (*elavl3*). (**c**) Larvae were transferred to 48

wells in the centre of a 96-well plate to ensure that the 405 nm LED Flood Array covers all

larvae entirely. To minimize overheating, the plate was floating in a waterbath at 10 cm distant to

the LED while CaMPARI was photoconverted (PC) by the LED Flood Array. (**d**) Lateral view of

zebrafish with exemplar "CaMPARI Activity" heat maps. Heat maps show ratio of red/green

(R/G) fluorescent output as indicated by the calibration bar and can be interpreted as relative

neural activity. CNS regions with higher levels of neural activity translate to "hotter" pixels.

Images were acquired using an automated INCell 2000 high-content microscope and reveal a

reduction in neural activity when the larvae were anesthetized with MS-222 or an increase with

convulsants, PTZ and 4-AP. Top, Enlarged lateral view from freely swimming larvae illustrating

the optic tectum and hindbrain areas (dashed line) from which the R/G ratio was obtained. (**e**)

Neural activity, inferred from the mean R/G ratio in optic tectum and hindbrain. CaMPARI

activity is reduced to baseline in MS222-anaesthetized fish and photoconversion is undetectable

when CaMPARI photoconverting light is omitted. Dashed green line represents mean for MS-

222-anesthetized samples, which show a clear reduction in signal compared to freely swimming

larvae and provides a baseline "near-zero" activity level for reference in subsequent experiments.

Drivers of neural activity, PTZ and 4-AP induce a significant increase in CaMPARI activity. (**f**)

Locomotor activity was measured using behavioural tracking software. PC light from the day

prior at 4 days post-fertilization (dpf) did not affect locomotion at 5 dpf. MS-222 treatment

abolished any swimbouts. Biological replicates are individual larvae = n. ** Significantly

different from freely swimming samples, p<0.01.

Figure 2: Early application of CBD and THC reduce subsequent neural activity and

locomotor activity. (**a**) The chemical structures of key cannabinoids used in this study: CBD

and THC. (**b**), Timeline of experimental set up where zebrafish embryos were treated with drugs

between 0.5 and 1 hour post-fertilization (hpf) and washed out at 10 hpf towards the end of

gastrulation. CaMPARI imaging was obtained at 4 days post-fertilization (dpf) and locomotor

activity was tracked at 5 dpf. (**c**) and (**f**), Representative CaMPARI activity maps obtained from

treated larvae and corresponding quantification and statistics shown in (**d**) and (**g**), respectively.

(**e**) and (**h**), Locomotor activity of the same larvae. (**c**) CBD reduces neural activity and

locomotor activity as shown in (**d**) and (**e**). (**f**) THC requires a higher does than CBD to reduce

neural activity and locomotor activity as shown in (**g**) and (**h**). CaMPARI activity heat maps

show ratio of R/G channels as indicated by the calibration bar, with higher ratios (hotter colours)

representing greater neural activity. Green-dashed lines depict mean baseline (zero) value for

MS-222 anesthetized samples (From Fig 1). Biological replicates are n=8-19. * is p<0.05; **

p<0.01 compared to the vehicle control (0.3 % MeOH in experiments with CBD and 0.6 %

MeOH for THC).

Figure 3: Antagonistic effect of CBD on neural activity is enhanced when combined with

sub-effective doses of THC. Zebrafish larvae were exposed to a series of CBD and THC

concentrations by themselves or in combination, which are mostly sub-effective, or in the case of

CBD, were minimally effective. (**a**) and (**d**), Exemplar CaMPARI activity heat maps show an

additive effect (minimizing neural activity) when exposed to 2 µg/ml (**d**) of each CBD and THC

compared to CBD or THC by themselves (as plotted in (**e**)). When 1 µg/ml of 1:1 CBD and THC

was applied, ratios show no additive effect as illustrated by quantifications and statistics in (**b**).

(**c**) and (**f**), Locomotor activity from the same well at 5 dpf shows a clear reduction when CBD

and THC is combined at 2 µg/ml each, but reduction mediated by CBD alone is almost as low as

when combined with THC suggesting that CBD is the main component affecting locomotor

activity. R/G is indicated by the calibration bar. Green-dashed lines depict mean values for MS-

222-anesthetized samples (from Figure 1). Biological replicates are n=15-25. * compared to

vehicle control (equal amount of MeOH in all experiments); [#] compared to CBD plus THC. One

symbol is p<0.05; Two symbols is p<0.01.

838
Figure 4: Cannabinoid receptor 1 (CB1R) and CB2R are both involved in CBD-mediated

reduction of locomotor activity. Zebrafish larvae were exposed to CBD (3 µg/ml) with CB1R-

and CB2R inhibitors, AM251 and AM630, respectively. (**a**) and (**b**), Exemplar CaMPARI heat

maps showing R/G ratios obtained at 4 dpf. (**c**) and (**d**), Corresponding quantifications showing

CBD mediated-reduction of neural activity is not fully rescued when CB1R is inhibited with

AM251 or CB2R with AM630 in the nanomolar (nM) range. (**e**) and (**f**), Locomotor activity at 5

dpf shows a clear rescue when inhibiting either CB1R or CB2R with 0.1 nM AM251, or 10 nM

AM630, respectively. (**g**) and (**h**), shows that applying AM251 or AM630 without CBD did not

affect CaMPARI activity or locomotion. R/G is indicated by the calibration bar. All samples

contained the same amount of vehicle DMSO (0.1 %) and MeOH (0.3 %) including the vehicle

control (Veh.). Green-dashed lines depict mean values for MS-222 anesthetized samples.

Biological replicates are n=13-30. * is p<0.05; ** p<0.01 compared to CBD.

Figure S1: Supplementary figure accompanying Fig.2. Re-plotting data to illustrate that CBD

shows an effect at lower concentration than THC. Quantifications of CaMPARI activity (left) or

locomotor activity (right) showing a statistical difference between CBD and THC at 4 or 5 days

post-fertilization when adding 3.0 µg/ml of either cannabinoids. Green-dashed lines depict mean

values for MS-222 anesthetized samples. Biological replicates are n=14-28. Statistics using

unpaired t-test shows ** p<0.01 compared to CBD.

Figure S2: Supplementary figure accompanying Fig.2. Re-plotting data from groups that were

treated with vehicle, CBD (3 µg/ml) or THC (6 µg/ml) to assess correlation between CaMPARI

activity vs. Locomotor activity within the same larvae. There is a significant positive linear

correlation between neural activity and locomotor activity when corresponding vehicle controls

(black circles) were plotted with either CBD (**a**; red circles) or THC (b; blue circles). Green-

dashed lines depict mean values for MS-222 anesthetized samples.

Figure S3: Supplementary figure accompanying Fig.3. Additive effects in reducing neural

activity are also obtained at lower CBD concentrations while THC concentration is kept constant

at 2 µg/ml. (**a**) Exemplar CaMPARI activity heat maps obtained at 4 days post-fertilization show

an effect at 0.5 and 1.5 µg/ml of CBD when combined with 2 µg/ml THC (blue) compared to

CBD alone (red), as illustrated by quantifications in (**b**). (**c**) Locomotor activity from the same

well at 5 dpf. R/G as indicated by the calibration bar. Green-dashed lines depict mean values for

MS-222 anesthetized samples. Biological replicates are n=14-28. * is compared to CBD; [#] is

compared to Vehicle. One symbol is p<0.05; two symbols are p<0.01.

Figure S4: Supplementary figure accompanying Fig.3. RMO44 staining of reticulospinal

neurons in the hindbrain shows a decrease in fluorescence intensities when combining CBD and

THC at 2 µg/ml at 5 dpf compared to when CBD or THC was added alone. Images show

biological replicates (n=3), left to right, from corresponding treatment group, top to bottom.

a

405 nm PC Light

↑[Ca²⁺]

↓[Ca²⁺]

b

400 µm

c

405 nm PC Light

d

no photoconversion

Relative Neural Activity (R/G)

2.8

2.1

1.4

0.7

0.0

MS-222 anesthetized

freely swimming

PTZ (2.5 mM)

4-AP (200 µM)

400 µM

e

CaMPARI Activity

no photoconversion
MS-222 anesthetized
freely swimming
PTZ (2.5 mM)
4-AP (200 µM)

R/G ±SEM

2.0

1.5

1.0

0.5

0.0

**

**

**

**

**

MS-222 anesthetized

n=13   n=19   n=40   n=11   n=12

Locomotor Activity

no photoconversion
MS-222 anesthetized
freely swimming

Δ % Activity within Swimbouts ±SEM

0.6

0.4

0.2

0.0

**

MS-222 anesthetized

n=16   n=5   n=32