Embryonic exposure to cannabidiol disrupts active neural circuits, an effect increased by

Δ☐-tetrahydrocannabinol and involving CB1R and CB2R in zebrafish

Richard Kanyo[a,b], Md Ruhul Amin [b], Laszlo F. Locskai[a,b], Danika D. Bouvier [b],

Alexandria M. Olthuis [b], W. Ted Allison[a, b, c], and Declan W. Ali [b, *]

[a] Centre for Prions & Protein Folding Disease, University of Alberta, Edmonton AB, T6G 2M8

[b] Department of Biological Sciences, University of Alberta, Edmonton AB, T6G 2E9, Canada

[c] Department of Medical Genetics, University of Alberta, Edmonton AB, T6G 2H7

23 * Author for correspondence (phone: (780) 492-6094; fax: (780) 492-9234; email:

24 declan.ali@ualberta.ca)

Abstract

Considering the wide spread use of cannabis as a recreational and medicinal drug, a knowledge gap exists regarding biological mechanisms and health implication. In the light of legislative changes, delayed effects of cannabis upon brief exposure during embryonic development are of high interest as early pregnancies often go undetected. We hypothesized that brief early cannabinoid exposure impacts neural activity by affecting embryonic brain development through cannabinoid-1 (CB1R) and -2 (CB2R). Zebrafish larvae were exposed to cannabidiol (CBD) and Δ-tetrahydrocannabinol (THC) until the end of gastrulation (1-10 hours post-fertilization) and analyzed later in development (4-5 days post-fertilization). In order to measure neural activity, we implemented the fluorescence based calcium detector CaMPARI (Calcium-Modulated Photoactivatable Ratiometric Integrator) and optimized the protocol such that a high number of samples can be economically used in a 96-well format complemented by locomotor analysis. Our results revealed that neural activity was decreased by CBD more than THC. At higher doses both cannabinoids could reduce neural activity close to a level comparable to anesthetized samples and locomotor activity was abolished. Interestingly, the decrease was more pronounced when CBD and THC were combined. At the receptor level, CBD-mediated reduction of locomotor activity was partially prevented using CB1R or CB2R inhibitors. Overall, we provided a mechanistic link to perturbed brain development *in vivo*, which seems to involve CBD and receptors, CB1R and CB2R. Future studies are warranted to reveal other cannabinoids and receptors involved in this pathway to understand the full health implications of cannabis consumption on fetal development.

Introduction

Publicly, cannabis is used as a recreational drug [1] that is often portrayed as harmless, yet the health implications are not fully understood. Interestingly, extracts of the *Cannabis sativa* plant have been used for medical purposes for almost 5000 years [2]. However, scientific knowledge is limited and despite controversy, some countries continue to legalize cannabis for recreational use. Cannabis has been reported to be one of the most used drugs during pregnancy and key compounds, Δ⊡-tetrahydrocannabinol (THC) and cannabidiol (CBD), can readily cross the placenta [3-6]. We were especially interested in CBD due to the reported health benefits and availability in many natural products including oils and food [7]. In the past, the negative impacts of cannabis in connection to embryonic development have been principally associated with the psychoactive THC [8]. In humans, epidemiological and clinical studies associate maternal cannabis exposure to behavioural disturbances in the offspring linked to increased risk for neuropsychiatric disorders [9]. In rats, maternal exposure of THC changed a series of behaviours in the offspring, including water-induced grooming, increased light sensitivity and altered exploratory behaviour [10]. Recently, the negative impacts of cannabis have expanded and also include the non-psychotropic CBD, which disrupts motor-neuron development in zebrafish [11]. This study is in contrast to reports that suggest positive health benefits of CBD, by treating nausea during pregnancy [12, 13].

The mechanistic pathway(s) by which THC and CBD are toxic are elusive. Recently, these two cannabinoids were linked to the sonic hedgehog signaling pathway in mice and zebrafish, and appears to involve cannabinoid type 1 receptor (CB1R) [14, 15]. It is well established that THC binds and activates as a partial agonist two distinct classes of G-coupled protein receptors, CB1R and CB2R [16]. Both receptors are highly expressed in neuronal tissue

where CB1R is localized mostly in the CNS [17-19] and CB2R in the peripheral nervous system

[20], although CB2R has been also linked to the CNS [21-23]. CBD has been reported to interact

with a lower affinity to CB1R and CB2R relative to THC, and rather antagonizes cannabinoid-

induced effects indirectly through other receptors [24-28]. Additional data to illuminate signaling

pathways of THC and CBD *in vivo* would be of great value.

Zebrafish embryos have several advantages that complement mammalian models.

Experiments are economical because many embryos are available, and exposure studies are

simple. Translucent larvae provide opportunities to implement cutting-edge fluorescing calcium

sensors and measure neural activity. Further, past studies suggest that the endocannabinoid

system plays a role in zebrafish development [29-31]. Both cannabinoid receptors investigated

here have similar expression profiles in the CNS compared to mammals, with CB1R sharing a

70% protein sequence identity with the human homolog [18, 31]. Considering the anticipated

effects of THC and CBD, high-throughput quantification of neural activity following early

exposures would be of great interest. In this study, the calcium modulated photoactivatable

ratiometric integrator (CaMPARI) [32], was implemented. Green fluorescing CaMPARI

undergoes photoconversion (PC) to a red fluorescing protein only in the presence of intense 405

nm light and high Ca^{2+} levels. This conversion is irreversible, thus creating a temporal snapshot

in a form of a ratiometric red/green output. CaMPARI is expressed exclusively in neuronal tissue

due to a pan-neural promoter (*elavl3*, a.k.a. HuC) and is a direct read-out of relative neural

activity. Initially, the photoconversion was not efficient, but our optimized protocol allows for

high-throughput well-plate-formats. Locomotion was also quantified from the same larvae as an

independent proxy. Here, we investigated the effects of the two most abundant cannabinoids

found in the plant, THC and CBD, on neural activity and characterized CB1R and CB2R *in vivo*

on the CBD effects using zebrafish.

METHODS

Animal Ethics and zebrafish husbandry

Zebrafish maintenance was approved by the Animal Care & Use Committee: Biosciences at the

University of Alberta and operated under the guidelines of the Canadian Council of Animal

Care. The fish were maintained in the University of Alberta fish facility at 28°C under a 14/10

light/dark cycle as previously described [33].

Engineering CaMPARI transgenic zebrafish for integrative calcium imaging.

Zebrafish expressing CaMPARI, i.e. the Tg[*elavl3*:CaMPARI (W391F+V398L)][ua3144]

line were generated as described previously [34, 35]. We re-derived CaMPARI fish in response

to a federal moratorium limiting zebrafish import into Canada [36].

CaMPARI photoconversion

For photoconversion, bright green Tg[*elavl3*:CaMPARI (W391F+V398L)][ua3144] larvae at

4 dpf were placed in a 96-well plate containing 150 µl egg water (made as previously decribed

[33]) and acclimatized for 2 h. The central 48 wells of the plate were exposed to 405 nm LED

array (Loctite) for 300 sec at a distance of 10 cm. LED array illuminated the plates entirely and

evenly. The plates were floating in water at room temperature to ensure that the larvae did not

overheat. For imaging, the larvae were anesthetized in 0.24 mg/mL tricaine methanesulfonate

(MS-222, Sigma) following photoconversion. MS-222 was washed out and zebrafish were rested

for a day prior locomotor activity assay.

CaMPARI analysis

The ratio of fluorescent emissions, taken as red photoconverted CaMPARI in ratio to

green CAMPARI, was interpreted as relative neural activity, as previously defined [32]. Images

were obtained with an InCell 2000 microscope (GE Healthcare, US). 96-well plates were run

using FITC (1 second exposure) and dsRed (2 second exposure) channels. ImageJ was used to

quantify the fluorescence mean of the red (dsRed) and green (FITC) channels. The area of the

optic and hind brain area were quantified as those regions were most consistent in the view. A

mask was applied to exclude noise outside of the CNS. Statistics used one-way ANOVA and

Dunnett's multiple comparison test with GraphPad Prism Software (Version 7, GraphPad, San

Diego, CA).

Quantifying Locomotor activity

The same zebrafish larvae being placed in the 96-well plate the day prior for CaMPARI

analysis were quantified for locomotion using Basler GenICaM (Basler acA 1300-60) scanning

camera (75-mm f2.8 C-mount lens) and EthoVision® XT-11.5 software by Noldus (Wageningen,

Netherlands) as described previously [37, 38]. This set-up allowed to see whether there is a

correlation between level of neural activity and the the mean activity % within a swimbout

(swimbout size). Swimbout size was shown to increase in response to the convulsant PTZ

previously [39] and here it can be a useful marker for neural activity. Transparent *Casper* strain

zebrafish [40] were unable to be tracked as ususal with Noldus. Therefore, quantification of

swimbouts provided an additional advantage. Clear movement was detected above the noise

treshold due to movement of the pigmented eyes. Although the % swimbout is small, reliable

measurements were obtained. Only clear swimbouts were quantified, which were set above a

treshold of 0.1 % of activity. Figure 1F shows that measuring the mean activity of a swimbout is

a useful tool in obtaining output that is clearly diminished in presence of MS-222. In addition,

the output also shows sensitivity in freely swimming larvae. Statistics used here were as

described above.

RMO44 staining

The zebrafish used in this experiment were of the Tuebingen Longfin strain. Set-up was

similar as described above. Following the experiment, larvae were stained with anti-RMO44

(Thermo Fisher Scientific, Waltham, MA, USA) and imaged using a Zeiss LSM 710 confocal

microscope (Oberkochen, Germany) as described elsewhere [41].

Results

High-throughput assessment of CaMPARI is a reliable metric of neural activity.

In this study the objective was to gain insight in the effects of THC and CBD on neural

activity during early embryonic development. We measured neural activity with CaMPARI.

CaMPARI photoconverts irreversibly from a green to a red fluorescing protein only if user-

applied 405 nm light application coincides with high calcium levels (Fig. 1A) [32]. Red and

green fluorescing CaMPARI are quantified as ratio (referred here as "CaMPARI activity"). We

expressed CaMPARI through a pan-neural promoter, *elavl3* (a.k.a HuC) exclusively in the CNS

and thus as a read-out of neural activity (Fig. 1B). Here, we optimized the use of CaMPARI

zebrafish for an automated high-content INCell2000 plate-reader (Fig.1C). CaMPARI transgene

expression was most suitable for this platform at 4 dpf, which is a suitable balance of fluorescing

intensity and a sufficiently developed brain.

CaMPARI activity was quantified and displayed as a heatmap consisting of the red/green

ratio in fluorescing intensities. At 4 dpf, optic tectum and hindbrain regions of larvae showed a

clear measureable output, whereas without photoconversion CaMPARI activity was nearly zero

(Fig. 1D and E). To establish a baseline of near-zero neural activity, larvae were anesthetized

with MS-222, and a significant reduction in red/green ratio was obtained compared to freely

swimming larvae (Fig. 1D and E). In contrast, drug-induced neural activity, with the established

convulsants pentylenetetrazole (PTZ) or 4-aminopyridine (4-AP), showed a consistent increase

in CaMPARI activity, even at concentrations considered to be minimal when inducing seizure

(Fig. 1D, E)[34,35].

We benchmarked our CaMPARI outputs against locomotor activity of zebrafish larvae,

which is an established metric of neural activity [43]. We chose 5 dpf as an optimal

developmental timepoint for analyzing locomotor activity, because younger larvae are largely

inactive. Larvae were left in the same 96-well plate for assessing CaMPARI and locomotion.

Exposing larvae to photoconversion with LED light the day before had no impact on the mean

locomotor activity (Fig. 1F). Larvae anesthetized with MS-222 did not display any detectable

swimbouts (Fig. 1F). The level of neural activity correlated with locomotor activity and will be

presented later. Overall, these results show that CaMPARI, deployed in zebrafish larvae is a

reliable high-throughput tool for measuring neural activity.

CBD and THC reduce subsequent neural activity.

We sought to assess how embryonic exposure to cannabinoids impacts upon subsequent

neural activity later in development. Here, the doses of CBD and THC aligned with our previous

work [11], reflecting high cannabis consumption in humans. Comparisons of our dosage to

humans requires various considerations: (i) blood plasma concentrations can peak up to 0.25

mg/l while smoking a single cigarette [44]; (ii) the content of THC has increased in the past 20

years; and (iii) doses of intraperitoneally administered medical CBD can vary greatly, from 5-

100mg/kg, and daily maximum of 1500mg/kg [45,46]. The current study uses up 6 µg/mL of

THC and 3 µg/mL CBD with an estimated 0.1-10% of this passing through the chorion to reach

the embryo.

We wanted to gain insight into the effects of CBD and THC separately (Fig. 2A), and

therefore each was applied in a dose-response format. This validation was necessary as

CaMPARI has not been used in studying cannabinoids. Compounds were added to the bath early

in animal development at (0.5 hpf) and then washed out towards the end of gastrulation (10 hpf).

CaMPARI was imaged at 4 dpf and locomotor activity was assessed at 5 dpf (Fig. 2B). Animals

that were exposed to CBD at concentrations of 2 µg/ml and 3 µg/ml exhibited a dose-dependent

reduction in CaMPARI activity (Fig. 2C, F). These reductions in CaMPARI output are

substantial when compared against the ~50% reduction we observed in anesthetized larvae,

where little neural activity is expected. Coordinate with this, locomotion was also reduced

starting at 1.5 µg/ml and was significant at 3 µg/ml (Fig. 2F). THC had a similar effect than

CBD and also reduced neural activity at higher doses, 4 µg/ml and 6 µg/ml, (Fig. 2F and G).

Locomotor activity exhibited a more extensive reduction at 6 µg/ml. We compared neural

activity and locomotion in the same individuals exposed to effective doses of cannabinoids and

found significant correlations (Fig. S2; CBD r=0.52 and THC r=0.71). Indeed, most larvae, that

displayed reduced neural activity also showed reduced locomotion. Together, our findings show

that both CBD and THC reduced neural activity when exposed early in development.

Antagonistic effects of CBD on neural activity is enhanced when combined with sub-

effective doses of THC. Cannabis consumption during pregnancy exposes the fetus to THC and

CBD in concert. In order to investigate whether CBD and THC have a combined effect that is

different from either compound on its own, neural activity was measured when larvae were

exposed to both cannabinoids, using the same timeline as the previous experiment (Fig. 2B).

First, various sub-effective doses of CBD were applied from 0.5 to 1.5 µg/ml in the presence of 2

µg/ml THC (identified as sub-effective in Fig. 2). Preliminary dosage attempts revealed small,

but significant differences on neural activity when combining 0.5 or 1.5 µg/ml CBD with 2.0

µg/ml THC (Fig. S3A and B). Locomotor activity also trended towards a dose-dependent

reduction during concerted application of CBD and THC, but was statistically non-significant

(Fig. S3C). Next, CBD and THC were added in a 1:1 ratio mixture at 1.0 and 2.0 µg/ml each. An

enhanced antagonistic effect was revealed on neural activity when both CBD and THC were

added at 2.0 µg/ml. CBD by itself at 2.0 µg/ml significantly reduced neural activity (Fig.3D)

consistent with results in Fig.2D, but CBD and THC together, further reduced neural activity

compared to CBD or THC alone (Fig. 3D and E). Locomotor activity was mainly affected by

CBD with no additional reduction when compared to CBD and THC combined (Fig. 3F). No

significant effect on neural activity or locomotor activity was obtained when 1.0 µg/ml of CBD

and THC was applied (Fig. 3A-C). Due to the effect on locomotion, we also wanted to assess the

integrity of neurons in connection to motor axons. Reticulospinal neurons in the hindbrain were

stained with RMO44 antibody targeting NEFM, an established marker for neuronal damage [47,

48]. Indeed, RMO44 immunostaining was further reduced when CBD and THC were combined

at 2.0 µg/ml suggesting that both CBD and THC affect neuronal health (Fig. 4S). Together, these

results suggest that CBD and THC are more potent in reducing neural activity when applied in

combination.

CB1R and CB2R are both required for CBD-induced reduction of locomotor

activity. Some reports suggest that CBD mediates its actions through CB1- and CB2- receptors

[24]. To test whether these two receptors are involved in the neuronal toxicity mediated by CBD

in vivo, we deployed corresponding receptor inverse agonist AM251 and antagonist AM630. As

described previously, the experimental timeline is as depicted in figure 2B with CBD being

added immediately after AM251 and AM630. When analyzing CaMPARI activity, our results

revealed that the CBD-mediated reduction in neural activity could not be completely prevented

with concentrations from 0.1 to 10 nM AM251 or AM630 in the presence of 3 µg/ml of CBD

(Fig. 4A-C, D). In contrast, CBD-mediated reduction of locomotor activity was prevented with a

concentration of AM251 as low as 1 nM (Fig. 4D) or with 10 nM of AM630. In the absence of

CBD, when using the maximum dose (10 nM) of AM251 or AM630, neither antagonist had an

effect. This finding excludes the possibility that the effect of AM251 or AM630 on locomotion

were independent of CBD (Fig. 4G and H). Overall, this suggests a role for both CB1R and

CB2R receptors in CBD-mediated effects on locomotor activity. In sum, inhibiting CB1R and

CB2R can ameliorate the negative impacts of early CBD on late CNS function, at least with

respect to locomotion.

Discussion

In some jurisdictions, including Canada, cannabis has been legalized despite the lack of

mechanistic understanding and health implications. Cannabis consumption has increased in the

past decade [49, 50] and reports emerged associating prenatal cannabis consumption with

stillbirths[51] and autism spectrum disorder, a neurodevelopmental syndrome [52, 53]. Yet,

cannabis and CBD oil have been used by consumers to treat labor pain [54]. Cannabis has been

shown to have detrimental effects on developing embryos in rats and chicken [10, 55]. More

recently, new evidence revealed that exposure to not only the more widely studied THC, but also

CBD, during gastrulation impacts the overall development in zebrafish [11]. Interestingly, the

effect of the two key cannabinoids, CBD and THC, were delayed, and observed after the

exposure had ended.

In vitro, CBD and THC can function through CB1R and CB2R, which are expressed in

neuronal tissue [16-22, 24]. Our goal was specifically to investigate the effects of CBD, and how

it might interact with the more widely studied THC, during early development and characterize

the impact on neural activity *in vivo* using our CaMPARI assay. We found that neural activity

was reduced later in zebrafish development upon brief exposure of not only THC, but also CBD

early in development. Surprisingly, CBD had similar impacts on neural activity at lower

concentrations compared to THC. Notably, the impact of CBD was increased by sub-effective

doses of THC. Our data quantifying locomotion, suggested that the CBD-mediated effect

involves CB1R and CB2Rs.

CaMPARI Assay

We optimized CaMPARI such that this innovation can be used for large number of

samples to screen compounds in future studies. Zebrafish larvae can be deployed in a 96-well

set-up quantifying locomotion [39, 42, 56-61]. Nevertheless, locomotor activity is not a direct

measurement of neural activity. Calcium sensors, such as the GCaMP series [62, 63], or more

recently GECO [64, 65], are more accurate, but CAMPARI became more practical once we

optimized the assay. CaMPARI provides a temporal-snapshot of neural activity in the freely

swimming animal, which can be anesthetized and imaged after the experiment [32].

Furthermore, the ratiometric red/green signals are imaged together and mitigate inter-individual

differences in expression levels or focal plane. We optimized conditions for a well-plate set-up

analyzing neural activity in the optic tectum and hindbrain. Unfortunately, less defined areas or

specific neurons could not be quantified. Furthermore, when interpreting our data as neural

activity, it is important to consider that when anesthetized a clear red to green CaMPARI ratio

can be still measured. This is unlikely due to autofluoresence because larvae that were not

photoconverted yielded zero CaMPARI activity. We cannot exclude the possibility that due to

longer photoconversion some CaMPARI has converted into the red fluorescence version even

when anesthetized. It is well established that when anesthetized with MS-222, neurons are barely

active [69, 70]. Indeed, our behavioural assay shows no locomotion in the presence of MS-222.

We believe this method is a suitable technique to measure substantial changes in neural activity

in the larger parts of the zebrafish CNS. Considering the advantages, this method outweighs its

limitation and could serve for future studies, especially in connection with epilepsy.

Delayed implications after early embryonic exposure of CBD and THC

Both CBD and THC have been implicated in embryonic development including abnormal

changes to neurons. Delayed abnormalities after early exposure of key cannabinoids have only

recently been investigated [11]. Here, early exposure of CBD and THC during the first 10 hours

of embryonic development reduced neural activity when measured days later, in a dose-

dependent manner. These later effects may be due to abnormal neurodevelopment and/or the

slow release of the cannabinoids from the tissue due to their lipophilic nature. Consistent with

this finding, zebrafish larvae with a similar exposure (5-10 hpf) to cannabinoids have shown

effects that could be due abnormal development (at 5 dpf). This was not exclusive to neurons,

but included other changes such as morphological abnormalities, decreased survival rate,

decreased mEPCs activity (may relate to muscle development), decreased heart rate and delayed

hatching [11]. Considering the specific time-window of exposure (first 10 hours in

development), an abrupt effect on the developmental program including neurons seems

plausible. Interestingly, CBD seemed to impact neural activity at a lower concentration than

THC, which also aligns with our previous study [11]. It is important to acknowledge that CBD

consumption has been documented to help with seizures and cancer by reducing pain [71, 72].

Nevertheless, our study suggests caution during pregnancies.

Considering that cannabis contains large proportions of both CBD and THC, we looked

at the effects of these two compounds in combination and found a more potent effect in

decreasing neural activity. Reticulospinal neurons in the hindbrain of the zebrafish control motor

neurons during swimming [47, 73]. In line with a reduction in locomotion, we observed reduced

staining when exposed to both cannabinoids, suggesting that the reticulospinal neuronal integrity

was indeed compromised. However, this finding did not explain why locomotor activity was

mainly influenced by CBD. This could be due in part that CBD has a greater effect at lower

concentration than THC. The effect of CBD may be amplified by effecting motor neurons and

potentially muscle development. Overall, these findings were surprising, because CBD was

found to negate the effects of THC on CB1R [26]. One possibility is that at higher concentrations

both, CBD and THC, affect the developmental program. In contrast, at lower concentrations and

in fully developed brains the impact of CBD and THC is different. This could result in

overriding the negating effects of CBD. Indeed, several receptors have been shown to have

different affinities to THC and CBD [24]. A second possibility, therefore, is that at high

concentration CBD and THC affect receptors other than CB1Rs and CB2Rs. This could explain

why inhibiting CB1R and CB2R did not fully prevent CBD-mediated reduction of neural

activity, as further discussed below.

Potential mechanism involved in the delayed effect of CBD and THC

At this stage, it is unclear what mechanism(s) account for the reduction in neural activity

later in development when embryos are exposed briefly to CBD and THC in the first hours after

fertilization. In this study, we were especially interested in the less widely studied CBD. Using

HEK cells several receptor candidates have been identified showing to interact with CBD [24]. A

recent study supports a functional role of CB1R and CB2R role in zebrafish [33]. Here, by

inhibiting CB1R, and CB2R, we found that the reduction in locomotor activity mediated by CBD

was partially prevented at nanomolar concentrations of the inhibitors suggesting that,

mechanistically, CBD acts through both receptors. Previous studies have implied that CBD may

bind to both CB1R and CB2R *in vitro* [24] and our study is in support that these interactions are

important *in vivo*. In line with this, CBD can lead to abnormal craniofacial and brain

development in mice and zebrafish, which is mediated through CB1R and the hedgehog

signaling pathway [15]. Whether CB2R can mediate its effects through the same pathway

remains to be determined. An explanation to why inhibiting CB1R and CB2R did prevent the

maximum effect of CBD, which is delayed, could relate to the fact that the CB1R and CB2R

receptors are expressed very early in development and found on stem cell progenitor cells [74].

As mentioned above an early effect may impact the developmental program such that changes

persist until later even if cannabinoids are no longer present in the circulation. Changes may not

be specific to the CNS as CB receptors are not only expressed in neuronal progenitor cells, but in

stem cells forming other types of tissues [74]. This could explain our previous study CBD

affected heart-rate and overall morphology after 5 dpf when exposed only briefly during

gastrulation [11]. The CBD-mediated reduction in neural activity was not prevented when either

CB1 or CB2Rs were inhibited. While the reason for this is unclear, it could relate that the effects

of CBD are larger outside the optic tectum and the hindbrain and more visible when quantifying

locomotion. It is possible that other receptors such as TRPV1 are involved [27, 75, 76], but this

needs further investigation.

Conclusion

Overall, both CBD and THC exposure during embryonic stages had a negative impact on

later neural activity, which was additive when combined. Together, this could mean that the

impact of cannabis on early development is higher than using isolated compounds and profound.

Our findings also support an *in vivo* mechanism of CBD functioning through CB1R and CB2R.

We believe this study opens up a new path for investigating detailed mechanisms, including

other receptors, in specific neurons that are affected by cannabinoid toxicity early in

development and persist. The accumulating evidence for prenatal cannabis consumption having

negative consequences on neurodevelopmental disorders in the offspring [3] has to be

considered. As the recreational use is becoming legalized and socially acceptable, more studies

are warranted to not only fully understand the impact on human development, but also the

mechanisms of the diseases.

FUNDING

This study was supported by grants to W.T. Allison from Alberta Prion Research Institute-

Alberta Innovates BioSolutions and the Alzheimer Society of Alberta and the Northwest

Territories, and to D.W. Ali from the Natural Sciences and Engineering Research Council of

Canada and Alberta Innovates mCannabis. L.F. Locskai was supported by undergraduate

studentships from Natural Sciences and Engineering Research Council of Canada and from

Alberta Innovates Health Solutions. R. Kanyo was supported by a SynAD postdoctoral

fellowship funded via Alzheimer Society of Alberta and Northwest Territories through their

Hope for Tomorrow program and the University Hospital Foundation.

DISCLOSURE

None of the authors have any conflicts of interest to disclose.

ACKNOWLEDGEMENTS

We would like to extend our thanks to members of the Allison and Ali lab. In adition, we would also like to acknowledge the Sciences Animal Support Services at the Unversity of Alberta.

AUTHOR'S CONTRIBUTIONS

RK, MRA, WTA, DWA contributed to the experimental design. RK, MRA, LFL, DDB, AMO, performed the experiments and analyzed the data. RK, MRA and DWA interpreted the results. RK, WTA and DWA wrote the manuscript. All authors read and approved the manuscript.

References

1. Degenhardt L, Chiu W-T, Sampson N, Kessler RC, Anthony JC, Angermeyer M, et al. Toward a global view of alcohol, tobacco, cannabis, and cocaine use: findings from the WHO World Mental Health Surveys. PLoS Med. 2008;5:e141.

2. Winek CL. Some historical aspects of marijuana. Clin Toxicol. 1977;10:243–53.

3. Bérard A. The importance of generating more data on cannabis use in pregnancy. Nat Med. 2020;26:1515–6.

4. Fergusson DM, Horwood LJ, Northstone K, ALSPAC Study Team. Avon Longitudinal Study of Pregnancy and Childhood. Maternal use of cannabis and pregnancy outcome. BJOG. 2002;109:21–7.

5. Gómez M, Hernández M, Johansson B, de Miguel R, Ramos JA, Fernández-Ruiz J. Prenatal cannabinoid and gene expression for neural adhesion molecule L1 in the fetal rat brain. Brain Res Dev Brain Res. 2003;147:201–7.

6. Hutchings DE, Martin BR, Gamagaris Z, Miller N, Fico T. Plasma concentrations of delta-9-tetrahydrocannabinol in dams and fetuses following acute or multiple prenatal dosing in rats. Life Sci. 1989;44:697–701.

7. Singer L, Tokish H, Park F, Campisi C, Milanaik RL. The cannabidiol conundrum: potential benefits and risks of cannabidiol products for children. Curr Opin Pediatr. 2020;32:198–205.

8. Friedrich J, Khatib D, Parsa K, Santopietro A, Gallicano GI. The grass isn't always greener: The effects of cannabis on embryological development. BMC Pharmacol Toxicol. 2016 Sep 29;17:45.

9. Morris CV, DiNieri JA, Szutorisz H, Hurd YL. Molecular mechanisms of maternal cannabis and cigarette use on human neurodevelopment. Eur J Neurosci. 2011;34:1574–83.

10. Navarro M, Rubio P, de Fonseca FR. Behavioural consequences of maternal exposure to natural cannabinoids in rats. Psychopharmacology (Berl). 1995;122:1–14.

11. Ahmed KT, Amin MR, Shah P, Ali DW. Motor neuron development in zebrafish is altered by brief (5-hr) exposures to THC (Δ9-tetrahydrocannabinol) or CBD (cannabidiol) during gastrulation. Sci Rep. 2018;8:10518.

12. Parker LA, Rock EM, Limebeer CL. Regulation of nausea and vomiting by cannabinoids. Br J Pharmacol. 2011;163:1411–22.

13. Roberson EK, Patrick WK, Hurwitz EL. Marijuana use and maternal experiences of severe nausea during pregnancy in Hawai'i. Hawaii J Med Public Health. 2014;73:283–7.

14. Subbaraman MS, Kerr WC. Simultaneous Versus Concurrent Use of Alcohol and Cannabis in the National Alcohol Survey. Alcoholism: Clinical and Experimental Research. 2015;39:872–9.

15. Fish EW, Murdaugh LB, Zhang C, Boschen KE, Boa-Amponsem O, Mendoza-Romero HN, et al. Cannabinoids Exacerbate Alcohol Teratogenesis by a CB1-Hedgehog Interaction. Sci Rep. 2019;9:16057.

16. Pertwee RG. Ligands that target cannabinoid receptors in the brain: from THC to anandamide and beyond. Addict Biol. 2008;13:147–59.

17. Herkenham M, Lynn AB, Little MD, Johnson MR, Melvin LS, de Costa BR, et al. Cannabinoid receptor localization in brain. Proc Natl Acad Sci U S A. 1990;87:1932–6.

18. Watson S, Chambers D, Hobbs C, Doherty P, Graham A. The endocannabinoid receptor, CB1, is required for normal axonal growth and fasciculation. Mol Cell Neurosci. 2008;38:89–97.

19. Kano M, Ohno-Shosaku T, Hashimotodani Y, Uchigashima M, Watanabe M. Endocannabinoid-mediated control of synaptic transmission. Physiol Rev. 2009;89:309–80.

20. Smita K, Sushil Kumar V, Premendran JS. Anandamide: an update. Fundam Clin Pharmacol. 2007;21:1–8.

21. Benarroch E. Endocannabinoids in basal ganglia circuits: implications for Parkinson disease. Neurology. 2007;69:306–9.

22. Stempel AV, Stumpf A, Zhang H-Y, Özdoğan T, Pannasch U, Theis A-K, et al. Cannabinoid Type 2 Receptors Mediate a Cell Type-Specific Plasticity in the Hippocampus. Neuron. 2016;90:795–809.

23. Liu Q-R, Canseco-Alba A, Zhang H-Y, Tagliaferro P, Chung M, Dennis E, et al. Cannabinoid type 2 receptors in dopamine neurons inhibits psychomotor behaviors, alters anxiety, depression and alcohol preference. Sci Rep. 2017;7:17410.

24. McPartland JM, Duncan M, Di Marzo V, Pertwee RG. Are cannabidiol and Δ(9)-tetrahydrocannabivarin negative modulators of the endocannabinoid system? A systematic review. Br J Pharmacol. 2015;172:737–53.

25. Gray RA, Whalley BJ. The proposed mechanisms of action of CBD in epilepsy. Epileptic Disorders. 2020;22:S10–5.

26. Laprairie RB, Bagher AM, Kelly MEM, Denovan⬜Wright EM. Cannabidiol is a negative allosteric modulator of the cannabinoid CB1 receptor. Br J Pharmacol. 2015;172:4790–805.

27. Ibeas Bih C, Chen T, Nunn AVW, Bazelot M, Dallas M, Whalley BJ. Molecular Targets of Cannabidiol in Neurological Disorders. Neurotherapeutics. 2015;12:699–730.

28. Ryberg E, Larsson N, Sjögren S, Hjorth S, Hermansson N-O, Leonova J, et al. The orphan receptor GPR55 is a novel cannabinoid receptor. Br J Pharmacol. 2007;152:1092–101.

29. Akhtar MT, Ali S, Rashidi H, van der Kooy F, Verpoorte R, Richardson MK. Developmental effects of cannabinoids on zebrafish larvae. Zebrafish. 2013;10:283–93.

30. Oltrabella F, Melgoza A, Nguyen B, Guo S. Role of the endocannabinoid system in vertebrates: Emphasis on the zebrafish model. Dev Growth Differ. 2017;59:194–210.

31. Sufian MS, Amin MR, Kanyo R, Allison WT, Ali DW. CB1 and CB2 receptors play differential roles in early zebrafish locomotor development. J Exp Biol. 2019;222(Pt 16).

32. Fosque BF, Sun Y, Dana H, Yang C-T, Ohyama T, Tadross MR, et al. Neural circuits. Labeling of active neural circuits in vivo with designed calcium integrators. Science. 2015;347:755–60.

33. Westerfield, M. The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio rerio). University of Oregon Press:Eugene, OR ;2000.

34. Kanyo R, Wang CK, Locskai LF, Li J, Allison WT, Kurata HT. Functional and behavioral signatures of Kv7 activator drug subtypes. Epilepsia. 2020;61:1678-1690

35. Alyenbaawi H, Kanyo R, Locskai LF, Kamali-Jamil R, DuVal MG, Bai Q, et al. Seizures are a druggable mechanistic link between TBI and subsequent tauopathy. bioRxiv. 14 May 2020 (accepted in *eLife*). https://doi.org/10.1101/2020.05.12.091819.

36. Hanwell D, Hutchinson SA, Collymore C, Bruce AE, Louis R, Ghalami A, et al. Restrictions on the Importation of Zebrafish into Canada Associated with Spring Viremia of Carp Virus. Zebrafish. 2016;13 Suppl 1:S153-163.

37. Leighton PLA, Kanyo R, Neil GJ, Pollock NM, Allison WT. Prion gene paralogs are dispensable for early zebrafish development and have nonadditive roles in seizure susceptibility. J Biol Chem. 2018;293:12576–92.

38. Kanyo R, Leighton PLA, Neil GJ, Locskai LF, Allison WT. Amyloid-β precursor protein mutant zebrafish exhibit seizure susceptibility that depends on prion protein. Exp Neurol. 2020;328:113283.

531 39. Hoffman EJ, Turner KJ, Fernandez JM, Cifuentes D, Ghosh M, Ijaz S, et al. Estrogens
532   Suppress a Behavioral Phenotype in Zebrafish Mutants of the Autism Risk Gene, CNTNAP2.
533   Neuron. 2016;89:725–33.
534
535 40. White RM, Sessa A, Burke C, Bowman T, LeBlanc J, Ceol C, et al. Transparent adult
536   zebrafish as a tool for in vivo transplantation analysis. Cell Stem Cell. 2008;2:183–9.
537
538 41. Amin MR, Ahmed KT, Ali DW. Early Exposure to THC Alters M-Cell Development in
539   Zebrafish Embryos. Biomedicines. 4 January 2020;8.
540   https://doi.org/10.3390/biomedicines8010005
541
542 42. Ellis LD, Seibert J, Soanes KH. Distinct models of induced hyperactivity in zebrafish larvae.
543   Brain Research. 2012;1449:46–59.
544
545 43. Rihel J, Prober DA, Arvanites A, Lam K, Zimmerman S, Jang S, et al. Zebrafish behavioral
546   profiling links drugs to biological targets and rest/wake regulation. Science. 2010;327:348–
547   51.
548
549 44. Huestis MA. Human cannabinoid pharmacokinetics. Chem Biodivers. 2007;4:1770–804.
550
551 45. Rohleder C, Müller JK, Lange B, Leweke FM. Cannabidiol as a Potential New Type of an
552   Antipsychotic. A Critical Review of the Evidence. Front Pharmacol. 2016;7:422.
553
554 46. Iffland K, Grotenhermen F. An Update on Safety and Side Effects of Cannabidiol: A Review
555   of Clinical Data and Relevant Animal Studies. Cannabis Cannabinoid Res. 2017;2:139–54.
556
557 47. Banote RK, Edling M, Eliassen F, Kettunen P, Zetterberg H, Abramsson A. β-Amyloid
558   precursor protein-b is essential for Mauthner cell development in the zebrafish in a Notch-
559   dependent manner. Developmental Biology. 2016;413:26–38.
560
561 48. Braissant O, Henry H, Villard A-M, Zurich M-G, Loup M, Eilers B, et al. Ammonium-
562   Induced Impairment of Axonal Growth Is Prevented through Glial Creatine. J Neurosci.
563   2002;22:9810–20.
564
565 49. Canada Health. Canadian Tobacco Alcohol and Drugs (CTADS) Survey: 2017 summary
566   2018 https://www.canada.ca/en/health-canada/services/canadian-tobacco-alcohol-drugs-
567   survey/2017-summary.html#n3 Accessed 30 November 2020
568
569 50. Statistics Canada. National Cannabis Survey, first quarter 2019. 2019;(11):15.
570
571 51. Varner MW, Silver RM, Rowland Hogue CJ, Willinger M, Parker CB, Thorsten VR, et al.
572   Association between stillbirth and illicit drug use and smoking during pregnancy. Obstet
573   Gynecol. 2014;123:113–25.
574
575 52. Fombonne E. Is there an epidemic of autism? Pediatrics. 2001;107:411–2.

53. Corsi DJ, Donelle J, Sucha E, Hawken S, Hsu H, El-Chaâr D, et al. Maternal cannabis use in pregnancy and child neurodevelopmental outcomes. Nat Med. 2020;26:1536–40.

54. Sarrafpour S, Urits I, Powell J, Nguyen D, Callan J, Orhurhu V, et al. Considerations and Implications of Cannabidiol Use During Pregnancy. Curr Pain Headache Rep. 2020;24:38.

55. Gustafsson SB, Jacobsson SOP. Effects of cannabinoids on the development of chick embryos in ovo. Scientific Reports. 2019;9:13486.

56. MacRae CA, Peterson RT. Zebrafish as tools for drug discovery. Nat Rev Drug Discov. 2015;14:721–31.

57. Rennekamp AJ, Peterson RT. 15 years of zebrafish chemical screening. Curr Opin Chem Biol. 2015;24:58–70.

58. Kokel D, Bryan J, Laggner C, White R, Cheung CYJ, Mateus R, et al. Rapid behavior—based identification of neuroactive small molecules in the zebrafish. Nat Chem Biol. 2010;6:231–7.

59. Baraban SC, Dinday MT, Hortopan GA. Drug screening in Scn1a zebrafish mutant identifies clemizole as a potential Dravet syndrome treatment. Nature Communications. 2013;4:2410.

60. Bose P, Tremblay E, Maois C, Narasimhan V, Armstrong GAB, Liao M, et al. The Novel Small Molecule TRVA242 Stabilizes Neuromuscular Junction Defects in Multiple Animal Models of Amyotrophic Lateral Sclerosis. Neurotherapeutics. 2019;16:1149–66.

61. Ibhazehiebo K, Gavrilovici C, de la Hoz CL, Ma S-C, Rehak R, Kaushik G, et al. A novel metabolism-based phenotypic drug discovery platform in zebrafish uncovers HDACs 1 and 3 as a potential combined anti-seizure drug target. Brain. 2018;141:744–61.

62. Mank M, Griesbeck O. Genetically Encoded Calcium Indicators. Chem Rev. 2008;108:1550–64.

63. Akerboom J, Chen T-W, Wardill TJ, Tian L, Marvin JS, Mutlu S, et al. Optimization of a GCaMP calcium indicator for neural activity imaging. J Neurosci. 2012;32:13819–40.

64. Shaner NC, Lambert GG, Chammas A, Ni Y, Cranfill PJ, Baird MA, et al. A bright monomeric green fluorescent protein derived from Branchiostoma lanceolatum. Nat Methods. 2013;10:407–9.

65. Zarowny L, Aggarwal A, Rutten VMS, Kolb I, GENIE Project, Patel R, et al. Bright and High-Performance Genetically Encoded Ca2+ Indicator Based on mNeonGreen Fluorescent Protein. ACS Sens. 2020;5:1959–68.

66. Sidik SM, Hortua Triana MA, Paul AS, El Bakkouri M, Hackett CG, Tran F, et al. Using a Genetically Encoded Sensor to Identify Inhibitors of Toxoplasma gondii Ca2+ Signaling. J Biol Chem. 2016;291:9566–80.

67. Streit AK, Fan YN, Masullo L, Baines RA. Calcium Imaging of Neuronal Activity in Drosophila Can Identify Anticonvulsive Compounds. PLoS One. 2016;11:e0148461.

68. Seshadri S, Hoeppner DJ, Tajinda K. Calcium Imaging in Drug Discovery for Psychiatric Disorders. Front Psychiatry. 2020;11:713.

69. Medler S. Anesthetic MS-222 eliminates nerve and muscle activity in frogs used for physiology teaching laboratories. Adv Physiol Educ. 2019;43:69–75.

70. Hedrick MS, Winmill RE. Excitatory and inhibitory effects of tricaine (MS-222) on fictive breathing in isolated bullfrog brain stem. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 2003;284:R405–12.

71. Perucca E. Cannabinoids in the Treatment of Epilepsy: Hard Evidence at Last? J Epilepsy Res. 2017;7:61–76.

72. Russo EB. Cannabinoids in the management of difficult to treat pain. Ther Clin Risk Manag. 2008;4:245–59.

73. Kohashi T, Nakata N, Oda Y. Effective Sensory Modality Activating an Escape Triggering Neuron Switches during Early Development in Zebrafish. J Neurosci. 2012;32:5810–20.

74. Galve-Roperh I, Chiurchiù V, Díaz-Alonso J, Bari M, Guzmán M, Maccarrone M. Cannabinoid receptor signaling in progenitor/stem cell proliferation and differentiation. Progress in Lipid Research. 2013;52:633–50.

75. McKemy DD, Neuhausser WM, Julius D. Identification of a cold receptor reveals a general role for TRP channels in thermosensation. Nature. 2002;416:52–8.

76. Moran MM. TRP Channels as Potential Drug Targets. Annu Rev Pharmacol Toxicol. 2018;58:309–30.

Figure 1: High-throughput quantification of neural activity in freely-swimming zebrafish

larvae. A, Calcium-modulated photoactivatable ratiometric integrator (CaMPARI)

photoconverts from green fluorescent emission to red emission in presence of both high calcium

concentrations and a bright 405 nm light source. This photoconversion is irreversible and thus

creates a temporal snapshot, integrated over the time window of light application, of increased

calcium levels (neural activity) as a red-to-green ratio of fluorescing CaMPARI. **B**, Lateral view

of green fluorescing CaMPARI merged with brightfield image shows expression of this neural

activity biosensor in the CNS due to a pan-neural promoter (*elavl3* aka HuC) allowing the red-to-

green ratio of CaMPARI to be interpreted as a direct output of relative neural activity of freely-

swimming larvae. **C**, Larvae were transferred to 48 wells in the centre of a 96-well plate to

ensure that the 405 nm LED Flood Array covers all larvae entirely. To minimize overheating, the

plate was floating in a waterbath at 10 cm distant to the LED while CaMPARI was

photoconverted (PC) by the LED Flood Array. Variations in fluorescence intensity of a sample

due to variable larvae positioning in different wells are normalized due to the ratiometric feature

of the CaMPARI output. **D**, Lateral view of zebrafish akin to panel B with Exemplar "CaMPARI

Activity" heat maps that were acquired in the 96 well plate using an automated INCell 2000

high-content microscope and reveal a consistent 2-fold reduction in neural activity when the

larvae were anesthetized with MS-222 and an expected increase when neural activity was

stimulated with convulsants, pentylenetetrazole (PTZ) and 4-aminopyridine (4-AP). Top,

Enlarged lateral view from freely swimming larvae illustrating the optic tectum and hindbrain

areas (dotted line) from which the R/G ratio was obtained. Heat maps show ratio of red/green

(R/G) fluorescent output as indicated by the calibration bar and can be interpreted as relative

calcium levels or neural activity. CNS regions with higher levels of neural activity translate to

higher R/G values and are reported here as "hotter" pixels in image. **E**, Neural activity, inferred

from the mean R/G ratio in optic tectum and hindbrain, increases and decreases following

application of stimulants or anesthetics, respectively. CaMAPRI is reduced to baseline in

MS222-anaesthetized fish and photoconversion is undetectable when CaMPARI photoconverting

light is omitted. Dashed green line represents mean for MS-222 anesthetized samples, which

show a clear reduction in signal compared to freely swimming larvae and provides a baseline

"near-zero" activity level for reference in subsequent experiments. Drivers of neural activity,

such as the convulsants PTZ and 4-AP induce a significant increase in CaMPARI signal **F**,

Locomotor activity was measured using behavioural tracking software. PC light from the day

prior at 4 days post-fertilization did not affect locomotion at 5 dpf. Anesthetic (MS-222)

treatment abolished any swimbouts. Biological replicates are individual larvae = n. **

Significantly different from freely swimming samples, $p<0.01$.

Figure 2: Early application of Cannabidiol (CBD) and Δ-tetrahydrocannabinol (THC)

reduce subsequent neural activity and locomotor activity. A, The chemical structures of key

cannabinoids used in this study: CBD and THC. **B**, Timeline of experimental set up where

zebrafish embryos were treated with drugs between 0.5 and 1 hour post-fertilization (hpf) and

washed out at 10 hpf towards the end of gastrulation. CaMPARI imaging was obtained at 4 days

post-fertilization (dpf) and locomotor activity was tracked at 5 dpf. **C** and **F**, Representative

CaMPARI activity maps obtained from treated larvae and corresponding quantification and

statistics shown in **D** and **G**, respectively. **E** and **H**, Locomotor activity of the same larvae. **C**,

CBD reduces neural activity and locomotor activity as shown in **D** and **E**. **F**, THC requires a

higher does than CBD to reduce neural activity and locomotor activity as shown in **G** and **H**.

CaMPARI heat maps show ratio of R/G channels as indicated by the calibration bar., with higher

ratios (hotter colours) representing greater neural activity. Green-dashed lines depict mean

baseline (zero) value for MS-222 anesthetized samples (From Fig 1). Biological replicates are

n=8-19. * is $p<0.05$; ** $p<0.01$ compared to the vehicle control (0.3 % MeOH in experiments

with CBD and 0.6 % MeOH for THC).

Figure 3: Antagonistic effect of CBD on neural activity is enhanced when combined with

sub-effective doses of THC. Zebrafish larvae were exposed to a series of CBD and THC

concentrations by themselves or in combination, which are mostly sub-effective, or in the case of

CBD, were minimally effective. **A** and **D**, Exemplar CaMPARI activity heat maps show an

additive effect (minimizing neural activity) when exposed to 2 µg/ml (**D**) of each CBD and THC

compared to CBD or THC by themselves (as plotted in **E**). When 1 µg/ml of 1:1 CBD and THC

was applied, ratios shows no additive effect as illustrated by quantifications and statistics in **B**. **C**

and **F**, Locomotor activity from the same well at 5 dpf shows a clear reduction when CBD and

THC is combined at 2 µg/ml each, but reduction mediated by CBD alone is almost as low as

when combined with THC suggesting that CBD is the main component affecting locomotor

activity. R/G is indicated by the calibration bar. Green-dashed lines depict mean values for MS-

222 anesthetized samples (from Figure 1). Biological replicates are n=15-25. * compared to

vehicle control (equal amount of MeOH in all experiments); [#] compared to CBD plus THC. One

symbol is $p<0.05$; Two symbols is $p<0.01$.

728
Figure 4: Cannabinoid receptor 1 (CB1R) and CB2R are both involved in CBD-mediated

reduction of locomotor activity. Zebrafish larvae were exposed to CBD (3 µg/ml) with CB1R-

and CB2R antagonists, AM251 and AM630, respectively. **A** and **B**, Exemplar CaMPARI heat

maps showing R/G ratios obtained at 4 dpf. **C** and **D**, Corresponding quantifications showing

CBD mediated-reduction of neural activity is not fully rescued when CB1R with AM251, but

shows significant rescue when blocking CB2R with AM630 in the nanomolar (nM) range. **E** and

F, Locomotor activity at 5 dpf shows a clear rescue when inhibiting either CB1R or CB2R with

0.1nM AM251, or 10 nM AM630, respectively. **G** and **H**, shows that applying AM251 or

AM630 without CBD did not affect CaMPARI activity or locomotion. All samples contained the

same amount of vehicle DMSO (0.1 %) and methanol (0.3 %) and the experiment included a

corresponding vehicle control (Veh.). Green-dashed lines depict mean values for MS-222

anesthetized samples. Biological replicates are n=13-30. * is $p<0.05$; ** $p<0.01$ compared to

CBD.

Figure S1: Supplementary figure accompanying Fig.2. Re-plotting data to illustrate that CBD

shows an effect at lower concentration than THC. Quantifications of CaMPARI activity (left) or

locomotor activity (right) showing a statistical difference between CBD and THC at 4 or 5 days

post-fertilization when adding 3.0 µg/ml of either cannabinoids. Green-dashed lines depict mean

values for MS-222 anesthetized samples. Biological replicates are n=14-28. Statistics using

unpaired t-test shows ** $p<0.01$ compared to CBD.

Figure S2: Supplementary figure accompanying Fig.2. Re-plotting data from groups that were

untreated, treated with vehicle, CBD (3 µg/ml) or THC (6 µg/ml) to assess CaMPARI activity

vs. Locomotor activity within the same individual larvae. There is a significant positive linear

correlation between neural activity and locomotor activity when corresponding vehicle controls

(black circles) were plotted with either CBD (A, red circles) or THC (B, blue circles).

Figure S3: Supplementary figure accompanying Fig.3. Additive effects in reducing neural

activity are also obtained at lower CBD concentrations while THC concentration is kept

constant. **A**, Exemplar CaMPARI activity heat maps obtained at 4 days post-fertilization show an

effect at 0.5 and 1.5 µg/ml of CBD when combined with 2 µg/ml THC compared to CBD as

illustrated by quantifications in **B**. **C**, Locomotor activity from the same well at 5 dpf. R/G as

indicated by the calibration bar. Green-dotted lines depict mean values for MS-222 anesthetized

samples. Biological replicates are n=14-28. * is $p<0.05$; ** $p<0.01$ compared to CBD.

Figure S4: Supplementary figure accompanying Fig.3. RMO44 staining of reticulospinal

neurons in the hindbrain shows a decrease in fluorescence intensities when combining CBD and

THC at 2 µg/ml at 5 dpf compared to when CBD or THC was added alone. Images show

exemplar biological replicates, left to right, from corresponding treatment group, top to bottom.

A 405 nm PC Light

↑[Ca²⁺]

↓[Ca²⁺]

B 400 µm

C 405 nm PC Light

D

Relative Neural Activity (R/G)

no photoconversion

MS-222 anesthetized

freely swimming

PTZ (2.5 mM)

4-AP (200 µM)

2.8
2.1
1.4
0.7
0.0

400 µM

E CaMPARI Activity

no photoconversion
MS-222 anesthetized
freely swimming
PTZ (2.5mM)
4-AP (200µM)

R/G ±SEM

2.0
1.5
1.0
0.5
0.0

**

**

**

**

**

MS-222 anesthetized

n=13   n=19   n=40   n=11   n=12

no photoconversion
MS-222 anesthetized
freely swimming

Δ % Activity within Swimbout ±SEM

0.4

0.2

0.0

**

MS-222 anesthetized

n=16   n=5   n=32