Title: Determinants of genetic diversity and species richness of North
American amphibians

Running title:

Authors: Chloé Schmidt[1*], Jason Munshi-South[2], Stéphane Dray[3], Colin J Garroway[1*]

Affiliations:

[1]Department of Biological Sciences, University of Manitoba, Winnipeg, MB, Canada

[2]Department of Biological Sciences and Louis Calder Center, Fordham University, Armonk, NY,
USA

[3]Univ Lyon, Université Claude Bernard Lyon 1, CNRS, Laboratoire de Biométrie et Biologie
Evolutive, F-69100, Villeurbanne, France

*Correspondence to:

Chloé Schmidt
Department of Biological Sciences
50 Sifton Rd
University of Manitoba
Winnipeg, MB R3T 2N2
email: schmid46@myumanitoba.ca
19
Colin J Garroway
Department of Biological Sciences
50 Sifton Rd
University of Manitoba
Winnipeg, MB R3T 2N2
email: colin.garroway@umanitoba.ca

Abstract: Ecological limits on the population sizes and number of species a region is capable of

supporting are thought to simultaneously produce spatial patterns in genetic diversity and species

richness. However, we do not know the extent to which resource-based environmental limits

jointly determine these patterns of biodiversity in ectotherms because of their low energy

requirements compared to endotherms. Here, we adapt a framework for the ways ecological

limits may shape genetic diversity and species richness previously tested in mammals for

amphibians to determine whether similar processes produce continental patterns of biodiversity

across both taxa. Repurposing open, raw microsatellite data from 19 species sampled at 554 sites

in North America we found that spatial patterns of genetic diversity run opposite to patterns of

species richness and genetic differentiation. However, while measures of resource availability

and niche heterogeneity predict 89% of the variation in species richness, these landscape metrics

are poor predictors of genetic diversity. Although heterogeneity appears to be an important driver

of genetic and species biodiversity patterns in both amphibians and mammals, our results suggest

that variation in genetic diversity both within and across species makes it difficult to infer

general processes producing spatial patterns of amphibian genetic diversity.

Keywords: latitudinal diversity gradient, biogeography, heterozygosity, frogs, salamanders,

more individuals hypothesis, macrogenetics

Introduction

Although species richness is higher in the tropics for most taxa, the details of diversity patterns

differ among species groups. In North America for instance, vertebrate richness generally

increases with resource availability, but mammals and birds tend to have higher species richness

in dry, mountainous areas, and reptiles and amphibians are more diverse in wet, lower elevation

regions (Currie 1991). This pattern suggests that while richness increases with resource

availability, taxon-specific traits may cause richness patterns to diverge from a strictly latitudinal

gradient. Broad-scale patterns of biodiversity at the genetic level have only recently begun to be

mapped thanks to the accumulation of open data in public repositories (Miraldo et al. 2016;

Manel et al. 2020; Schmidt et al. 2020; Theodoridis et al. 2020). Genetic diversity is typically

thought of as the most fundamental level of biodiversity because it bears on populations'

capacities to evolve adaptively in response to environmental change (Frankham 1995). Recent

analyses of mammals suggest that environments can simultaneously shape species richness and

genetic diversity on continental scales (Schmidt et al. 2020). Whether this is also true in

ectothermic taxa is unknown. Understanding whether common processes underlie variation in

biogeographic patterns of diversity across taxa with different environmental requirements can

help us move toward a general understanding of the drivers of biodiversity at multiple levels.

In mammals, continental scale multi-species patterns of nuclear genetic diversity and species

richness could be inferred with estimates of resource and niche availability, or ecological

opportunity (Schmidt et al. 2020). This suggested that ecological limits placed on the number of

individuals and species an environment can support are important drivers of broad-scale

biodiversity patterns. Resource-rich environments supported larger populations with higher

genetic diversity and species richness, while niche availability in heterogeneous habitats

promoted species coexistence but reduced population sizes and genetic diversity because

specialization restricts available resources. These processes are related to two well-supported

hypotheses for the production of biogeographic patterns of species richness: the more-individuals

hypothesis (Wright 1983), and the effects of environmental and resource heterogeneity (Allouche

et al. 2012; Stein et al. 2014). The more individuals hypothesis posits that resource-rich regions

near the equator are capable of supporting larger populations and communities, and thus more

species than temperate regions. Heterogeneity is thought to increase species richness due to

greater niche availability in more complex heterogeneous environments which allows more

species to coexist, but with smaller population sizes because resources are partitioned.

The effects of resource heterogeneity on species richness and genetic diversity seem likely to be

generally applicable across taxa (Stein et al. 2014; Schmidt et al. 2020), however the general

relevance of the more-individuals hypothesis for ectotherms is uncertain (Buckley and Jetz

2010). This is because compared to endotherms, ectotherms have lower energy requirements and

can behaviorally thermoregulate, meaning their abundances are less likely to be limited by

resource-related ecological limits (Pough 1980; Buckley and Jetz 2010). Instead, ectotherm

distributions, and therefore species richness, seem more directly constrained by environmental

temperature because fewer species have evolved thermal adaptations required for expanding into

cooler regions (Buckley and Jetz 2007, 2010). Further, the evolution of traits associated with

better survival in temperate regions may have additional effects on speciation dynamics. For

example, species turnover tends to be higher among viviparous squamate reptiles, which

typically occupy cooler regions (Pyron and Burbrink 2014). Reaching a more comprehensive

understanding of how biodiversity patterns emerge requires knowledge about whether

relationships between genetic diversity, species richness, and environments are consistent across

endothermic and ectothermic taxa.

The determinants of species richness across all terrestrial vertebrates are generally related to

resource availability as estimated by energy (e.g., potential evapotranspiration, primary

productivity), water-energy balance (e.g., actual evapotranspiration, precipitation), and

heterogeneity (e.g., elevation variability, land cover diversity) (Currie 1991; Kerr and Packer

1997; Hawkins et al. 2003; Rodríguez et al. 2005; Buckley and Jetz 2007; Stein et al. 2014;

Jiménez-Alfaro et al. 2016). Amphibians are interesting because they are constrained both by

water availability and temperature and water availability is consistently identified as an

important driver of diversity in amphibians (Rodríguez et al. 2005; Buckley and Jetz 2007).

Indeed in Europe, the best predictors of species richness in mammals and birds shift from energy

to water availability at decreasing latitudes, but amphibian species richness remains strongly

related to water-energy balance regardless of latitude (Whittaker et al. 2007).

The causes of population genetic diversity are rarely studied at the same time or scale as patterns

of species richness (but see Marshall and Camp 2006; Schmidt et al. 2020), yet the presumed

mechanisms related to the more-individuals hypothesis and heterogeneity are closely related to

carrying capacity and population-level processes (Schmidt et al. 2020). The more-individuals

mechanism predicts a positive relationship between species richness and population genetic

diversity because bigger populations and communities tend to have higher levels of genetic and

species diversity (Kimura 1983; Hubbell 2001; Schmidt et al. 2020). With higher carrying

capacities, more species persist because they can reach minimal viable population sizes. On the

other hand, heterogeneity is predicted to cause negative correlations between genetic diversity

and species richness by increasing the number of species a given area can support which in turn

reduces population size and limits gene flow due to increased niche specialization.

Heterogeneous environments also facilitate population differentiation due to spatially varying

selection. In mammals, evolutionary processes acting on the population level scaled up and

interacted with resource availability and heterogeneity to produce genetic diversity and species

richness patterns (Schmidt et al. 2020).

Whether mechanisms related to ecological limits and niche availability predict patterns of

species richness and genetic diversity in ectotherms is unclear. To test this prediction, we

analyzed previously-published microsatellite genotype data from 19 North American amphibian

species (8 frogs, 11 salamanders), with >13000 individuals sampled at 554 sites. Our first

objective was to identify existing spatial patterns in genetic diversity and differentiation and

quantify the extent to which genetic diversity and species richness covary. We then tested

whether limits on resources and niche availability jointly determined genetic diversity and

species richness using structural equation models, which allowed us to evaluate multiple

hypotheses at genetic and species level biodiversity simultaneously. We based our conceptual

framework on previous findings in mammals (Schmidt et al. 2020) and adapted it for

amphibians. First, we excluded human presence because previous investigation shows it did not

have a clear effect on amphibian genetic diversity (Schmidt and Garroway 2021a). Second, we

included a measure of water availability as an additional indicator of resource availability. Third,

body size was used as a proxy for whole species census size in mammals, however we do not

know the extent to which body size is correlated with effective population size or genetic

diversity within amphibians. We therefore first tested for a relationship between body size and

genetic diversity for species in our sample to determine whether it should be incorporated into

the model. Finally, we compare our results to previous results in mammals (Schmidt et al. 2020)

to infer whether similar environmental features contribute to diversity gradients across

endothermic and ectothermic taxa in North America.

Methods

Biodiversity data

Genetic diversity and differentiation. We used raw genotypes (i.e. called allele sizes) of North

American amphibians compiled by (Schmidt and Garroway 2021a). This data set was assembled

from raw microsatellite datasets publicly archived in Dryad (DataDryad.org). To identify data

sets we conducted a systematic search of the Dryad data repository with the following keywords:

species name (e.g. *Plethodon cinereus*), "microsat*", "short tandem*", and "single tandem*".

We used the IUCN Red List database to obtain a list of amphibian species native to North

America for the search. We excluded datasets that lacked spatial reference, were not located in

North America, did not sample neutral microsatellite loci, or had study designs that may have

affected genetic diversity (including sampling island populations, or captive or managed

populations). In total the data sets include genotypes for 13680 individuals spanning 19 species

sampled at 554 locations in the contiguous United States and Canada. We used gene diversity as

a measure of genetic diversity because it is minimally affected by sample size (Nei 1973;

Charlesworth and Charlesworth 2010). Our measure of differentiation was population-specific

F_{ST} (Weir and Goudet 2017). Population-specific F_{ST} differs from pairwise F_{ST} in that it

measures how far single populations in a sample have diverged from a common ancestor—this

means it is comparable across studies and species. Population-specific F_{ST} was not estimable at 2

sites because it requires a minimum of 2 sample sites in the original dataset ($n = 552$).

Species richness. We estimated species richness at each of our genetic diversity sample sites

using amphibian range extent data from the IUCN RedList (IUCN 2019), applying filters for

native, extant species ranges. We measured species richness as the number of species' ranges

overlapping each genetic sample site.

Diversity maps and spatial variation partitioning

We used distance-based Moran's eigenvector maps (MEMs) to detect spatial patterns in genetic

diversity and differentiation and compare these to patterns of species richness. MEMs are

orthogonal spatial eigenvectors with eigenvalues that are directly proportional to Moran's *I*.

They measure spatial autocorrelation at all scales present in the data. We computed MEMs in the

R package adespatial (Dray et al. 2017). We used the forward selection procedure described in

Blanchet et al. (2008) to select two sets of MEMs describing important patterns in genetic

diversity, differentiation, and species richness. We then selected MEMs which explained the

broadest spatial patterns in these data (Moran's I > 0.25). To create maps of genetic diversity,

differentiation, and species richness (Fig. 1), we used the fitted values for gene diversity,

population-specific F_{ST}, and species richness regressed on broad-scale MEMs. This approach

allowed us to visualize broad purely spatial patterns in genetic diversity, differentiation, and

species richness without variation due to other sources, such as local environments, species

identity, or population history. Maps of raw values are presented in Fig. S1.

Next, we determined the extent to which spatial patterns in genetic diversity and species richness

were shared using variation partitioning. Because our MEM analysis for both levels of

biodiversity had the same input distance matrix, the resulting spatial MEMs were directly

comparable. This was not the case for genetic differentiation, which had fewer sample sites. We

therefore did not partition variation in genetic differentiation because these MEMs are not the

same as those computed for genetic diversity and species richness.

We determined the fraction of total variation explained by spatial structure, shared spatial

structure, and non-spatial variation using variation partitioning as follows. We ran a series of

linear regressions with either species richness (y_{SR}) or gene diversity (y_{GD}) as the response

variable using all MEMs selected for that variable (Equations 1 and 2), or only MEMs shared by

both variables as predictors (Equations 3 and 4):

$y_{SR} \sim \alpha + \beta_{1S}(\mathrm{MEM}_{1S}) + \beta_{2S}(\mathrm{MEM}_{2S}) + \cdots + \beta_{8S}(\mathrm{MEM}_{8S}) + \epsilon$ Eq. 1

$y_{GD} \sim \alpha + \beta_{1G}(\mathrm{MEM}_{1G}) + \beta_{2G}(\mathrm{MEM}_{2G}) + \cdots + \beta_{5G}(\mathrm{MEM}_{5G}) + \epsilon$ Eq. 2

$y_{SR} \sim \alpha + \beta_{1SG}(\mathrm{MEM}_{1SG}) + \beta_{2SG}(\mathrm{MEM}_{2SG}) + \cdots + \beta_{5SG}(\mathrm{MEM}_{5SG}) + \epsilon$ Eq. 3

$y_{GD} \sim \alpha + \beta_{1SG}(\mathrm{MEM}_{1SG}) + \beta_{2SG}(\mathrm{MEM}_{2SG}) + \cdots + \beta_{5SG}(\mathrm{MEM}_{5SG}) + \epsilon$ Eq. 4

where α is the grand mean, and MEM_{iS} and MEM_{iG} are the set of MEMs selected for species

richness (8 MEMs) and genetic diversity (5 MEMs), respectively. The coefficients of variation

(R^2) from Eqs. 1 and 2 give the total amount of variation explained by spatial patterns for species

richness and genetic diversity. Subtracting these values from 1 gives the amount of non-spatial

variation. MEM_{iSG} represents the set of MEMs shared by both species richness and genetic

diversity (5 MEMs). R^2 values from Eqs. 3 and 4 tell us the amount of variation in each response

variable which can be explained by spatial variation shared at both levels of diversity. When

subtracted from the total spatial variation in genetic diversity or species richness (Eqs. 1 and 2),

we get the proportion of non-shared spatial variation.

Structural equation modeling

Environmental data. We measured resource and niche heterogeneity by computing Simpson's

Diversity Index for landcover categories within buffers at each site. We obtained a 30 m

resolution map of North American landcover data from the Commission for Environmental

Cooperation (CEC et al. 2015). This map is based on 2015 satellite imagery and has 19 standard

land cover classifications which includes forests, shrubland, grassland, wetlands, cropland,

barren land, and built-up land. Because this variable is scale-dependent, we recorded

heterogeneity within 4 buffer sizes (10, 25, 40, and 80 km) around each site.

We used two measures of resource availability because amphibians are habitat-limited by both

temperature and water availability. Water availability can be measured by evapotranspiration, or

the amount of water removed from the Earth's surface through soil or open water evaporation

and plant transpiration processes. Potential evapotranspiration (PET) measures the atmospheric

demand for water, depending on factors such as temperature and wind (Peng et al. 2019). It is

strongly correlated with temperature. PET is the maximum amount of water that would be

removed in the absence of biophysical limitations (Peng et al. 2019). The amount of water

actually removed, actual evapotranspiration (AET), reflects water availability and soil moisture

levels. Actual evapotranspiration has also been shown to be one of the strongest predictors of

amphibian species richness (Buckley and Jetz 2007). PET can be viewed as a measure of energy

availability, and AET one of water-energy balance (see Currie 1991; Buckley and Jetz 2007;

Kreft and Jetz 2007). Together these variables represent total ecosystem resource availability at

sites. We measured mean PET and AET (mm/yr) at sites within 10, 25, 40, and 80 km buffers

using data from the CGIAR Consortium for Spatial Information (Trabucco and Zomer 2019).

Population size. Body size is a readily measured trait that is typically negatively correlated with

long-term effective population size across a diversity of species (Frankham 1996; Romiguier et

al. 2014; Mackintosh et al. 2019). To explore whether body size was a useful substitute for long-

term effective population size in amphibians, we obtained body length data from the AmphiBIO

v1 database (Oliveira et al. 2017). We then regressed it on gene diversity in a hierarchical model

with species as a random effect allowing intercepts to vary and predicted a negative relationship.

We found no relationship between body size and genetic diversity in our data (Fig. S2), thus we

excluded population size from our proposed causal framework. Without population size as an

intermediate variable, we predicted that resource availability would have direct positive effects

on both genetic diversity and species richness, and that heterogeneity would have direct negative

and positive effects on genetic diversity and species richness, respectively. We hypothesize that

these direct effects would act through population size were such data available.

Analysis. We used structural equation modeling (SEM) to determine whether genetic diversity

and species richness are shaped by differential ecological limits due to limits on resources and

niche availability. Structural equation modeling begins with a causal diagram, or conceptual

model, where paths between variables represent hypothesized causal relationships (Fig. 2a).

Hypotheses are envisioned as a network where variables are nodes, and paths connecting them

represent causal relationships. In SEM, the effects of multiple predictors are simultaneously

assessed for multiple response variables (Shipley 2016). We implemented structural equation

models using the piecewiseSEM package (version 2.0.2), which uses a local estimation approach

for models in the hypothesis network allowing for the incorporation of more complex model

types (Lefcheck et al. 2019). Model fit is evaluated using tests of directed separation (Shipley

2016), which determine whether an association exists between two variables in the network

conditional on each of their causes. If two variables are not conditionally independent, the model

is updated by adding a path between them to make the model more consistent with the data. *P*-

values from tests of directed separation are used to calculate Fisher's *C,* which is then used to

calculate an overall *p*-value for the model network. Models are a good fit to the data when *p* >

0.05, indicating the null hypothesis—the proposed hypothesis network—is not rejected.

Our model network consisted of two models with genetic diversity and species richness as

response variables. Our genetic data is hierarchical with multiple sites nested within species. To

account for variation in mean genetic diversity across species we modeled random intercepts for

species. Additionally, we allowed the responses of genetic diversity to resource availability and

heterogeneity to vary across species by including a random slope term for each environmental

predictor (PET, AET, and land cover diversity). With this random effect structure, we thus allow

relationships to vary among species and do not assume effects will be of the same magnitude or

direction for all species. We fit the hierarchical model for genetic diversity using the lme4

package (Bates et al. 2015) within piecewiseSEM. We scaled and centered all variables before

analysis so path coefficients were comparable. Finally, we checked model residuals for spatial

autocorrelation using Moran tests. We conducted SEM analyses in parallel for each of our 4

buffer sizes.

Effect of heterogeneity on population differentiation

Finally, we tested whether landscape heterogeneity was related to increased population

differentiation. We regressed heterogeneity on population-specific F_{ST} using a hierarchical model

with a random effect for species accounting for differences in mean F_{ST} (intercepts) while

allowing the strength and direction of the effect of heterogeneity (slopes) to vary across species.

To account for spatial variation we included MEMs describing spatial patterns in F_{ST} as

covariates. We performed these analyses in parallel across all four heterogeneity buffers.

Results

Spatial patterns in genetic variation. We detected spatial patterns across genetic diversity,

genetic differentiation, and species richness (Fig. 1). The major axis of broad-scale variation is

longitudinal, and across all three biodiversity metrics the western sample sites show little

variation whereas spatial patterns were more complex in the east and appear to vary latitudinally.

This broad longitudinal pattern is consistent with environmental variation in North America (Fig.

1) as well as genetic diversity and species richness patterns in mammals (Schmidt et al. 2020).

We recovered known patterns of amphibian species richness with MEMs, where richness was

highest in the southeastern United States (Fig. 1) (Currie 1991). Western USA, which is hotter

and drier, had a comparatively low number of species. In eastern sample sites, genetic diversity

increased with latitude, while differentiation and species richness increase towards the tropics.

Genetic diversity and differentiation in the western samples were in the mid-range of values

across sample sites (Fig. 1).

In general, species richness was more spatially structured than genetic diversity, with 85% and

23% of variation explained by spatial patterns, respectively (Fig. 1). We detected shared spatial

patterns between both levels of biodiversity, however, while shared patterns accounted for the

entirety of the spatial variation in genetic diversity, they explained less of the variation in species

richness (18%).

Common causes of genetic diversity and species richness. Our conceptual model (Fig 2a) fit the

data well (Fisher's C = 0.49, *p* = 0.78, 2 degrees of freedom) with no additional links suggested

at any scale. Note that for SEM, *p* > 0.05 means that our conceptual model is not rejected. We

present results from the 40 km buffer in the main text; results from all models can be found in

tables S1 – 4. Species richness was well explained (R^2 = 0.89) and increased with water

availability, environmental heterogeneity, and species body size (Fig 2, Table S3). Water

availability had the strongest effect on species richness. Genetic diversity was not well predicted

by any variables in our model (R^2 = 0.04; Fig. 2). Residuals from genetic diversity models did

not exhibit spatial autocorrelation. Species richness residuals were spatially autocorrelated at

local scales (Moran's *I* = 0.06). In general, the environmental covariates in our models captured

broad spatial patterns well, and we did not incorporate fine-scale spatial structure into our

models as this was likely due to clustered sampling of some species. Lastly, genetic

differentiation within species decreased with heterogeneity (β = -0.32 ± 0.10 SE) at the most

local spatial scale we tested (10 km buffer), but this relationship disappeared at larger buffer

sizes.

Discussion

We found spatial variation shared across genetic diversity, differentiation, and species richness at

broad spatial scales (Fig. 1). In general, areas with high species richness tended to have

genetically differentiated populations with relatively low genetic diversity. There was a

latitudinal gradient in genetic diversity across eastern sites which varied in the opposite direction

of the gradient in species richness and genetic differentiation. These patterns are consistent with

our predicted effects of heterogeneity, however, these relationships were not well-reflected by

our structural equation model (Fig. 2). Our variation partitioning suggests that all spatial

variation in genetic diversity was shared with species richness, but the environments that

predicted species richness did not predict genetic diversity well. This finding suggests that other

processes drive the genetic diversity patterns we detect. These processes are likely species-

specific, and not generalizable at continental scales as they were in mammals (Schmidt et al.

2020).

We suspect the general lack of relationship between genetic diversity and species richness and

climate we find in our structural equation model may come down to amphibian population

dynamics. Ecological limits hypotheses assume that communities are in equilibrium with respect

to speciation, colonization, and extinction dynamics (Storch et al. 2018)— extending this to the

genetic level, we also assume populations are in an equilibrium state with regards to gene flow,

mutation, and genetic drift. However, amphibians have variable local population sizes which can

sometimes fluctuate by orders of magnitude from year to year (Collins et al. 2009), high rates of

species turnover at local and regional scales (Werner et al. 2007; Buckley and Jetz 2008), and

relatively low occupancy within potential distributions (Munguía et al. 2012). Species turnover at

sites is associated with environmental heterogeneity in freshwater habitats, especially

hydroperiod variation in temporary ponds (Urban 2004). Varying population dynamics could

obscure general relationships between population genetic diversity and population size, species

richness, and the climatic factors we explore here.

Interestingly, although genetic diversity was not affected by environmental heterogeneity at any

scale, genetic differentiation decreased with heterogeneity at the most local scale we tested;

genetic differentiation was relatively low in the northeast where landscape heterogeneity was

higher (Fig. 1). If heterogeneity increases niche availability and creates opportunities for

specialization and divergence, we predicted that it would increase genetic differentiation.

However, the pattern we detect here may be expected if species that were capable of recolonizing

northern regions following glaciation tend to be widely distributed generalists that maintain

population connectivity over relatively large geographic distances (Smith et al. 2005; Zeisset and

Beebee 2008). Niche partitioning in amphibians may also occur at finer scales within suitable

habitats (Karlin et al. 1984; Cloyed and Eason 2017)—for example by modifying microhabitat

usage, diets, foraging strategies, or behaviors across species. Thus, the landscape level metric we

use here may not capture the varying ways that environmental heterogeneity affects genetic

diversity and population differentiation if species respond to environments in different ways.

These patterns point to the importance of species-specific responses to environmental conditions

and environmental instability in generating biogeographic patterns of genetic diversity and

species richness in amphibians (Urban 2004).

Previous exploration of the relationships between nuclear genetic diversity, species richness, and

environments in plethodontid salamanders produced similarly mixed results (Marshall and Camp

2006). Genetic diversity was positively associated with temperature and rainfall across all eight

species studied, but topographic heterogeneity had both positive and negative effects on genetic

diversity depending on species. Allelic richness was only correlated with species richness for

Desmognathus species and *Plethodon jordani*, but relationships were positive and negative

(Marshall and Camp 2006). In another example, Karlin et al. (1984) report a negative

relationship between species richness and genetic diversity across populations of *D. fuscus*.

Similar to our findings, it appears resource availability and heterogeneity simultaneously affect

biodiversity on genetic and species levels, but genetic diversity in general is less well predicted

by environments alone. Variation between species may reduce our ability to detect general

relationships across species.

Although we did not detect clear latitudinal or longitudinal gradients in genetic diversity across

North America, previous findings suggest mitochondrial genetic diversity in amphibians and

other ectotherms varies latitudinally and mirrors species richness patterns (Miraldo et al. 2016;

Manel et al. 2020). Miraldo et al. (2016) found that amphibian mitochondrial genetic diversity in

North America was highest in the species-rich southeastern United States. They suggested this

pattern may be related to the evolutionary speed hypothesis, where presumably high

environmental temperature increases rates of population divergence and speciation through its

effects on mutation rate and generation time (Miraldo et al. 2016). In addition to a negative

correlation between patterns of nuclear genetic diversity and species richness, we detected no

effect of temperature on nuclear genetic diversity in our SEM, casting doubt on this hypothesis

for amphibians. Furthermore, a lack of latitudinal gradient indicates that nuclear genetic diversity

is not related to temperature in a straightforward way. Environmental temperature has varied and

complex effects on metabolism and mitochondrial processes (Munro and Treberg 2017; Zhang

and Wong 2021), making relationships with nuclear and mitochondrial mutation rates unlikely to

be generalizable (Lanfear et al. 2007; Schmidt and Garroway 2021b). More generally

mitochondrial DNA alone is not a reliable marker for detecting intraspecific demographic

patterns (Bazin et al. 2006; Galtier et al. 2009; Schmidt and Garroway 2021b). Thus, marker

choice is very likely responsible for the diverging patterns we find here. This also appears to be

true in mammals, where patterns of genetic diversity measured using mitochondrial DNA and

nuclear DNA also trended in opposite directions (Schmidt et al. 2020).

Despite limitations modeling population size, it appears that spatial patterns in genetic diversity

and species richness in amphibians are driven by processes similar to mammals. Interestingly,

similar proportions of variation in genetic diversity (~ 25%) could be attributed to spatial

processes across both taxa. Additionally, it was the case in both groups that spatial variation in

genetic diversity was primarily due to factors that also shaped species richness, but other factors

contribute to unique spatial variation in species richness. The overall negative correlation

between spatial patterns of species richness and genetic diversity in amphibians and mammals

(Schmidt et al. 2020) suggests heterogeneity and niche partitioning are major contributors to

diversity across genetic and species levels in endo- and ectothermic vertebrates. Heterogeneity

has previously been put forth as a universal driver of species richness (Stein et al. 2014). In

general, similar environmental factors seem capable of generating an overall species richness

gradient across taxa, but slight deviations from this general pattern are mediated by differential

interactions between environments, species traits, and population processes.

Genetic diversity and species richness are two important metrics for biodiversity conservation

because they contribute to the resilience of populations and communities in rapidly changing

environments (Oliver et al. 2015). Neutral genetic diversity is indicative of population mean

fitness and the efficiency of selection in response to environmental change—although it is only

weakly correlated with additive genetic diversity, it is nevertheless informative for conservation

purposes because it reflects levels of inbreeding and the efficiency of selection due to its

relationship to the effective population size (Frankham 1995; Mittell et al. 2015). Amphibians

are among the most imperiled vertebrates (Stuart et al. 2004) and are especially susceptible to

environmental change. Macrogenetics approaches mapping multispecies patterns of genetic

diversity at broad scales have great potential for incorporation into conservation policies

targeting regional conservation of genetic diversity. However, complex ecophysiological

requirements, life histories, and population dynamics may render this approach impractical for

amphibians because the environmental factors affecting genetic diversity may differ depending

on species (Schmidt and Garroway 2021a). Species-specific measures of environmental

heterogeneity, resource availability, and habitat suitability may prove to be more reliable

predictors of genetic diversity, but may be less relevant for species richness. We are only

beginning to explore broad scale patterns of intraspecific genetic diversity across several species,

but it is already apparent that they are not as consistently clear as gradients in species richness

(Miraldo et al. 2016; Manel et al. 2020; Schmidt et al. 2020; Theodoridis et al. 2020). We look

forward to the continued exploration of these patterns in other taxonomic groups to help build a

comprehensive picture of the distribution of genetic biodiversity across the globe.

Author contributions: C.J.G., and C.S. conceived of the study. C.S., J.M.S, S.D. and C.J.G.

designed the study and C.S. conducted the analyses with input from S.D. and C.J.G. All authors

contributed to data interpretation. C.S. wrote the first draft of the manuscript and all authors

participated in editing subsequent manuscript drafts.

Acknowledgements: We would like to thank Mitchell Green for assistance with data

acquisition, as well as the authors of the original datasets for making their data public. C.S. and

C.J.G. were supported by a Natural Sciences and Engineering Research Council of Canada

Discovery Grant to C.J.G. C.S. was also supported by a U. Manitoba Graduate Fellowship, and a

U. Manitoba Graduate Enhancement of Tri-council funding grant to C.J.G.

Data availability: Synthesized genetic data is available from the Dryad Data Repository (DOI:

10.5061/dryad.qv9s4mwf0). Species range boundary files and environmental data are available

from open online sources (see Methods).

References

Allouche, O., Kalyuzhny, M., Moreno-Rueda, G., Pizarro, M., and Kadmon, R. 2012. Area-
heterogeneity tradeoff and the diversity of ecological communities. Proc. Natl. Acad. Sci.
U. S. A. **109**(43): 17495–17500. doi:10.1073/pnas.1208652109.

Bates, D., Mächler, M., Bolker, B., and Walker, S. 2015. Fitting linear mixed-effects models
using lme4. J. Stat. Softw. **67**(1). doi:10.18637/jss.v067.i01.

Bazin, E., Glémin, S., and Galtier, N. 2006. Population size does not influence mitochondrial
genetic diversity in animals. Science **312**(5773): 570–572. doi:10.1126/science.1122033.

Blanchet, G.F., Legendre, P., and Borcard. 2008. Forward selection of explanatory variables.
Ecology **89**(9): 2623–2632.

Buckley, L.B., and Jetz, W. 2007. Environmental and historical constraints on global patterns of
amphibian richness. Proc. R. Soc. B Biol. Sci. **274**(1614): 1167–1173.
doi:10.1098/rspb.2006.0436.

Buckley, L.B., and Jetz, W. 2008. Linking global turnover of species and environments. Proc.
Natl. Acad. Sci. U. S. A. **105**(46): 17836–17841. doi:10.1073/pnas.0803524105.

Buckley, L.B., and Jetz, W. 2010. Lizard community structure along environmental gradients. J.
Anim. Ecol. **79**(2): 358–365. doi:10.1111/j.1365-2656.2009.01612.x.

CEC, NRCan/CCMEO, USGS, INEGI, CONABIO, and CONAFOR. 2015. 2015 North
American Land Cover at 30 m spatial resolution. Available from http://www.cec.org/north-
american-environmental-atlas/land-cover-30m-2015-landsat-and-rapideye/.

Charlesworth, B., and Charlesworth, D. 2010. Elements of evolutionary genetics. Roberts &
Company Publishers, Greenwood Village, Colorado, USA.

Cloyed, C.S., and Eason, P.K. 2017. Niche partitioning and the role of intraspecific niche
variation in structuring a guild of generalist anurans. R. Soc. Open Sci. **4**(3).
doi:10.1098/rsos.170060.

Collins, J.P., Crump, M.L., and Lovejoy III, T.E. 2009. Extinction in Our Times: Global
Amphibian Decline. Oxford University Press, Oxford.

Currie, D.J. 1991. Energy and large-scale patterns of animal- and plant-species richness. Am.
Nat. **137**(1): 27–49.

Dray, S., Blanchet, G., Borcard, D., Clappe, S., Guenard, G., Jombart, T., Larocque, G.,
Legendre, P., Madi, N., and Wagner, H.H. 2017. adespatial: Multivariate Multiscale Spatial

Analysis. Available from https://cran.r-project.org/package=adespatial.

Frankham, R. 1995. Conservation Genetics. Annu. Rev. Genet. **29**: 305–332.

Frankham, R. 1996. Relationship of Genetic Variation to Population Size in Wildlife. Conserv.
Biol. **10**(6): 1500–1508. doi:10.1046/j.1523-1739.1996.10061500.x.

Galtier, N., Nabholz, B., Glémin, S., and Hurst, G.D.D. 2009. Mitochondrial DNA as a marker
of molecular diversity: A reappraisal. Mol. Ecol. **18**(22): 4541–4550. doi:10.1111/j.1365-
294X.2009.04380.x.

Hawkins, B.A., Field, R., Cornell, H. V., Currie, D.J., Guégan, J.-F, Kaufman, D.M., Kerr, J.T.,
Mittelbach, G.G., Oberdorff, T., O'Brien, E.M., Porter, E.E., and Turner, J.R.G. 2003.
Energy, water, and broad-scale geographic patterns of species richness. Ecology **84**(12):
3105–3117. doi:10.1890/03-8006.

Hubbell, S.P. 2001. The Unified Neutral Theory of Biodiversity and Biogeography. Princeton
University Press, Princeton NJ.

IUCN. 2019. The IUCN Red List of Threatened Species. Version 2019-1. Available from
https://www.iucnredlist.org.

Jiménez-Alfaro, B., Chytrý, M., Mucina, L., Grace, J.B., and Rejmánek, M. 2016. Disentangling
vegetation diversity from climate-energy and habitat heterogeneity for explaining animal
geographic patterns. Ecol. Evol. **6**(5): 1515–1526. doi:10.1002/ece3.1972.

Karlin, A.A., Guttman, S.I., Rathbun, S.L., and Rathbun, S.L. 1984. Spatial autocorrelation
analysis of heterozygosity and geographic distribution in populations of Desmognathus
fuscus (Amphibia: Plethodontidae). Copeia **1984**(2): 343–356.

Kerr, J.T., and Packer, L. 1997. Habitat heterogeneity as a determinant of mammal species
richness. Nature **385**: 253–254.

Kimura, M. 1983. The Neutral Theory of Molecular Evolution. Cambridge University Press,
Cambridge.

Kreft, H., and Jetz, W. 2007. Global patterns and determinants of vascular plant diversity. Proc.
Natl. Acad. Sci. **104**(14): 5925–5930. doi:10.1073/pnas.0608361104.

Lanfear, R., Thomas, J.A., Welch, J.J., Brey, T., and Bromham, L. 2007. Metabolic rate does not
calibrate the molecular clock. Proc. Natl. Acad. Sci. U. S. A. **104**(39): 15388–15393.
doi:10.1073/pnas.0703359104.

Lefcheck, J., Byrnes, J., and Grace, J. 2019. piecewiseSEM: Piecewise Structural Equation
Modeling. Available from https://cran.r-project.org/package=piecewiseSEM.

Mackintosh, A., Laetsch, D.R., Hayward, A., Charlesworth, B., Waterfall, M., Vila, R., and
Lohse, K. 2019. The determinants of genetic diversity in butterflies. Nat. Commun. **10**(1):
1–9. doi:10.1038/s41467-019-11308-4.

Manel, S., Guerin, P.E., Mouillot, D., Blanchet, S., Velez, L., Albouy, C., and Pellissier, L. 2020.
Global determinants of freshwater and marine fish genetic diversity. Nat. Commun. **11**(1):
1–9. Springer US. doi:10.1038/s41467-020-14409-7.

Marshall, J.L., and Camp, C.D. 2006. Environmental correlates of species and genetic richness in lungless salamanders (family plethodontidae). Acta Oecologica **29**(1): 33–44. doi:10.1016/j.actao.2005.07.008.

Miraldo, A., Li, S., Borregaard, M.K., Florez-Rodriguez, A., Gopalakrishnan, S., Rizvanovic, M., Wang, Z., Rahbek, C., Marske, K.A., and Nogues-Bravo, D. 2016. An Anthropocene map of genetic diversity. Science **353**(6307): 1532–1535. doi:10.1126/science.aaf4381.

Mittell, E.A., Nakagawa, S., and Hadfield, J.D. 2015. Are molecular markers useful predictors of adaptive potential? Ecol. Lett. **18**(8): 772–778. doi:10.1111/ele.12454.

Munguía, M., Rahbek, C., Rangel, T.F., Diniz-Filho, J.A.F., and Araújo, M.B. 2012. Equilibrium of global amphibian species distributions with climate. PLoS One **7**(4): e34420. doi:10.1371/journal.pone.0034420.

Munro, D., and Treberg, J.R. 2017. A radical shift in perspective: Mitochondria as regulators of reactive oxygen species. J. Exp. Biol. **220**(7): 1170–1180. doi:10.1242/jeb.132142.

Nei, M. 1973. Analysis of gene diversity in subdivided populations. Proc. Natl. Acad. Sci. U. S. A. **70**(12): 3321–3323. doi:10.1073/pnas.70.12.3321.

Oliveira, B.F., São-Pedro, V.A., Santos-Barrera, G., Penone, C., and Costa, G.C. 2017. AmphiBIO, a global database for amphibian ecological traits. Sci. Data **4**: 1–7. doi:10.1038/sdata.2017.123.

Oliver, T.H., Heard, M.S., Isaac, N.J.B., Roy, D.B., Procter, D., Eigenbrod, F., Freckleton, R., Hector, A., Orme, C.D.L., Petchey, O.L., Proença, V., Raffaelli, D., Suttle, K.B., Mace, G.M., Martín-López, B., Woodcock, B.A., and Bullock, J.M. 2015. Biodiversity and resilience of ecosystem functions. Trends Ecol. Evol. **30**(11): 673–684. doi:10.1016/j.tree.2015.08.009.

Peng, L., Zeng, Z., Wei, Z., Chen, A., Wood, E.F., and Sheffield, J. 2019. Determinants of the ratio of actual to potential evapotranspiration. Glob. Chang. Biol. **25**(4): 1326–1343. doi:10.1111/gcb.14577.

Pough, F.H. 1980. The Advantages of Ectothermy for Tetrapods. Am. Nat. **115**(1): 92–112.

Pyron, R.A., and Burbrink, F.T. 2014. Early origin of viviparity and multiple reversions to oviparity in squamate reptiles. Ecol. Lett. **17**(1): 13–21. doi:10.1111/ele.12168.

Rodríguez, M.Á., Belmontes, J.A., and Hawkins, B.A. 2005. Energy, water and large-scale patterns of reptile and amphibian species richness in Europe. Acta Oecologica **28**(1): 65–70. doi:10.1016/j.actao.2005.02.006.

Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., Chiari, Y., Dernat, R., Duret, L., Faivre, N., Loire, E., Lourenco, J.M., Nabholz, B., Roux, C., Tsagkogeorga, G., Weber, A.A.T., Weinert, L.A., Belkhir, K., Bierne, N., Glémin, S., and Galtier, N. 2014. Comparative population genomics in animals uncovers the determinants of genetic diversity. Nature **515**(7526): 261–263. doi:10.1038/nature13685.

Schmidt, C., Dray, S., and Garroway, C.J. 2020. Genetic and species-level biodiversity patterns are linked by demography and ecological opportunity. bioRxiv.

539   doi:10.1101/2020.06.03.132092.

540 Schmidt, C., and Garroway, C.J. 2021a. The population genetics of urban and rural amphibians
541   in North America. Mol. Ecol. doi:10.1111/mec.16005.

542 Schmidt, C., and Garroway, C.J. 2021b. The conservation utility of mitochondrial genetic
543   diversity in macrogenetic research. Conserv. Genet. **22**(3): 323–327. doi:10.1007/s10592-
544   021-01333-6.

545 Shipley, B. 2016. Cause and Correlation in Biology, 2nd edition. Cambridge University Press,
546   Cambridge.

547 Smith, S.A., Stephens, P.R., and Wiens, J.J. 2005. Replicate patterns of species richness,
548   historical biogeography, and phylogeny in holarctic treefrogs. Evolution **59**(11): 2433–
549   2450. doi:10.1111/j.0014-3820.2005.tb00953.x.

550 Stein, A., Gerstner, K., and Kreft, H. 2014. Environmental heterogeneity as a universal driver of
551   species richness across taxa, biomes and spatial scales. Ecol. Lett. **17**(7): 866–880.
552   doi:10.1111/ele.12277.

553 Storch, D., Bohdalková, E., and Okie, J. 2018. The more-individuals hypothesis revisited: the
554   role of community abundance in species richness regulation and the productivity–diversity
555   relationship. Ecol. Lett. **21**(6): 920–937. doi:10.1111/ele.12941.

556 Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L., and
557   Waller, R.W. 2004. Status and trends of amphibian declines and extinctions worldwide.
558   Science **306**(5702): 1783–1786. doi:10.1126/science.1103538.

559 Theodoridis, S., Fordham, D.A., Brown, S.C., Li, S., Rahbek, C., and Nogues-Bravo, D. 2020.
560   Evolutionary history and past climate change shape the distribution of genetic diversity in
561   terrestrial mammals. Nat. Commun. **11**(1): 2557. doi:10.1038/s41467-020-16449-5.

562 Trabucco, A., and Zomer, R. 2019. Global Aridity Index and Potential Evapotranspiration (ET0)
563   Climate Database v2. doi:10.6084/m9.figshare.7504448.v3.

564 Urban, M.C. 2004. Disturbance heterogeneity determines freshwater metacommunity structure.
565   Ecology **85**(11): 2971–2978. doi:10.1890/03-0631.

566 Weir, B.S., and Goudet, J. 2017. A Unified Characterization of Population Structure and
567   Relatedness. Genetics **206**(4): 2085–2103. doi:10.1534/genetics.116.198424.

568 Werner, E.E., Yurewicz, K.L., Skelly, D.K., and Relyea, R.A. 2007. Turnover in an amphibian
569   metacommunity: the role of local and regional factors. Oikos **116**(10): 1713–1725.
570   doi:10.1111/j.2007.0030-1299.16039.x.

571 Whittaker, R.J., Nogués-Bravo, D., and Araújo, M.B. 2007. Geographical gradients of species
572   richness: a test of the water-energy conjecture of Hawkins et al. (2003) using European data
573   for five taxa. Glob. Ecol. Biogeogr. **16**: 76–89. doi:10.1111/j.1466-822x.2006.00268.x.

574 Wright, D.H. 1983. Species-Energy Theory: An Extension of Species-Area Theory. Oikos **41**(3):
575   496–506.

576 Zeisset, I., and Beebee, T.J.C. 2008. Amphibian phylogeography: a model for understanding

historical aspects of species distributions. Heredity. **101**: 109–119.
doi:10.1038/hdy.2008.30.

Zhang, Y., and Wong, H.S. 2021. Are mitochondria the main contributor of reactive oxygen
species in cells? J. Exp. Biol. **224**(5). doi:10.1242/jeb.221606.

Table 1. Data summary. Summary of aggregated raw genetic data: mean gene diversity, mean number of loci, median number of
individuals at sites per species. Species richness is the mean species richness at sites. Environmental variables (40 km
buffer): potential evapotranspiration (PET); actual evapotranspiration (AET); heterogeneity is the mean land cover diversity measured
with Simpson's Index. Ranges of values are given in parentheses where applicable.

Species (# sites)	Gene diversity	Loci	Individuals	Species richness	PET	AET	Heterogeneity
Anura							
Ascaphus montanus (100)	0.74 (0.41–0.87)	13	20 (10–41)	7.35 (3–11)	1097.84 (1006.52–1327.35)	506.86 (341.66–601.4)	0.54 (0.31–0.72)
Lithobates pipiens (5)	0.86 (0.83–0.9)	9	40 (20–46)	18.4 (16–21)	961.02 (924.69–1041.02)	656.24 (614.39–711.78)	0.74 (0.69–0.78)
Lithobates sylvaticus (90)	0.8 (0.7–0.9)	10.57 (7–15)	24 (5–36)	21.4 (11–38)	1062.37 (843.19–1437.5)	759.82 (669.33–1050.57)	0.76 (0.63–0.84)
Pseudacris crucifer (11)	0.75 (0.65–0.81)	11	28 (5–114)	16.18 (15–17)	984.96 (913.86–1122.62)	656.36 (569.9–689.03)	0.53 (0.27–0.8)
Pseudacris streckeri (17)	0.59 (0.38–0.82)	14	8 (5–22)	24.71 (15–34)	1479.33 (1257.25–1925.78)	878.45 (521.88–1001.36)	0.63 (0.3–0.86)
Rana draytonii (17)	0.51 (0.27–0.66)	15	13 (7–83)	9.41 (7–12)	1776.46 (1586.25–2179.73)	405.28 (331.77–493.63)	0.71 (0.55–0.8)
Rana luteiventris (25)	0.55 (0.37–0.69)	8	15 (6–205)	9.16 (5–11)	1200.7 (1077.99–1563.67)	494.77 (254–579.38)	0.66 (0.41–0.74)
Rana pretiosa (23)	0.32 (0.16–0.55)	10.39 (7–13)	34 (26–299)	9.35 (5–18)	1195.73 (924.34–1341.85)	515.32 (424.91–631.07)	0.55 (0.37–0.81)
Caudata							
Ambystoma barbouri (76)	0.78 (0.54–0.88)	11	22 (8–43)	28.28 (21–34)	1274.78 (1189.33–1337.51)	889.87 (811.77–967.52)	0.68 (0.55–0.75)
Ambystoma maculatum (97)	0.69 (0.45–0.75)	9.16 (7–14)	24 (8–36)	20.74 (16–35)	1074.81 (961.91–1406.61)	752.9 (682.96–931.33)	0.74 (0.37–0.8)
Desmognathus fuscus (5)	0.39 (0.15–0.65)	5	26 (22–35)	26.4 (26–27)	1258.69 (1224.99–1271.15)	827 (813.24–849.54)	0.62 (0.58–0.67)
Dicamptodon aterrimus (3)	0.48 (0.43–0.53)	9	91 (90–180)	6.33 (5–7)	1039.28 (1029.16–1046.55)	527.8 (469.46–564.46)	0.36 (0.32–0.41)
Dicamptodon copei (29)	0.69 (0.48–0.87)	11	18 (6–81)	13.86 (13–15)	905.66 (826.03–1074.59)	652.43 (628.31–678.36)	0.57 (0.48–0.74)
Ensatina eschscholtzii (4)	0.8 (0.75–0.84)	10	11.5 (5–19)	10 (9–11)	2140.32 (2039.14–2308.24)	368.69 (341.99–380.48)	0.63 (0.6–0.72)
Hydromantes brunus (6)	0.52 (0.38–0.61)	10	10 (7–18)	9.5 (8–11)	2021.39 (2008.05–2037.4)	347.91 (291.31–424.01)	0.7 (0.63–0.75)
Hydromantes platycephalus (15)	0.44 (0.25–0.64)	10	13 (6–21)	6.93 (4–9)	1786.55 (1523.97–2159.97)	466.68 (326.23–576.76)	0.65 (0.55–0.73)
Plethodon albagula (21)	0.48 (0.44–0.55)	20	16 (10–24)	31 (31–31)	1349.69 (1348.91–1350.48)	833.13 (832.48–833.85)	0.66 (0.66–0.66)
Plethodon cinereus (1)	0.6	7	122	22	1348.41 (1348.41–1348.41)	843.96	0.77
Taricha granulosa (9)	0.4 (0.26–0.77)	6	16 (6–32)	4 (2–13)	746.76 (614.95–1612.18)	522.65 (480.12–609.21)	0.73 (0.69–0.82)

Figure 1. *(Top row)* Maps of predicted genetic diversity, species richness, and genetic differentiation at genetic sample sites (points) based on spatial MEMs. MEMs were able to recover known patterns of species richness, which are negatively correlated with spatial patterns of genetic diversity. Patterns of genetic differentiation mirror those of species richness. *(Bottom row)* Maps depicting the environmental variables predicted to have simultaneous effects on genetic diversity and species richness, and variation partitioning results. Note land cover heterogeneity is a categorical variable and this map represents different land cover classes.

Figure 2. Structural equation model results. (A) Our hypothesized conceptual model. Line color denotes predicted directions of effect. (B) Results for model using environments within 40 km of sites (results for all spatial scales are in Tables S1-S4). Regression coefficients with standard errors are shown along each path. Paths between variables where no effect was detected are colored in gray (see Table S3 for a complete summary of all paths). R^2, the proportion of variation explained by the model, is given for genetic diversity and species richness. For genetic diversity, R^2_m is the variation explained by fixed effects and R^2_c is the variation explained by both fixed effects and the random species effect.

Determinants of genetic diversity and species richness of North American amphibians

Supplementary Information

Figures S1-3

Tables S1-4

Genetic diversity

Species richness

Genetic differentiation

Figure S1. Maps of raw data for genetic diversity, species richness, and genetic differentiation. Genetic diversity is Nei's gene diversity; species richness is the number of amphibian species' ranges overlapping each site; genetic differentiation is site-specific F_{ST}.

Figure S2. Relationship between genetic diversity and body size (snout-vent length). We detected no relationship using a mixed-effects model including a random intercept for species (β = -0.03 ± 0.2 SE).

Figure S3. Spatial patterns (MEMs) for genetic diversity and species richness are correlated with environments: energy availability (potential evapotranspiration; PET), water availability (actual evapotranspiration; AET), and heterogeneity (land cover diversity). MEMs are ordered along the x-axis according to spatial scale explained, from broadest (MEM1) to finest (MEM18). MEMs to the left of the dashed lines indicate the broadest-scale patterns with Moran's I > 0.25 used to produce maps of genetic diversity and species richness. Light blue bars are MEMs explaining spatial patterns of species richness, and dark blue bars are MEMs explaining spatial patterns shared by genetic diversity and species richness. No MEMs explained spatial patterns in genetic diversity only.

Table S1. Path coefficients and standard errors for SEM model with heterogeneity measured within a 10 km buffer around sites (Fisher's C = 0.28, p = 0.87, 2 degrees of freedom). Genetic diversity R^2_m = 0.04 (variance explained by fixed effects); R^2_c = 0.92 (variance explained by fixed and random effects); species richness R^2 = 0.87.

Response	Predictor	Estimate ± SE
gene diversity	potential evapotranspiration	0.16 ± 0.16
gene diversity	actual evapotranspiration	0.13 ± 0.14
gene diversity	heterogeneity	0.16 ± 0.09
species richness	potential evapotranspiration	0.18 ± 0.02
species richness	actual evapotranspiration	0.90 ± 0.02
species richness	heterogeneity	0.09 ± 0.02

7   **Table S2.** Path coefficients and standard errors for SEM model with heterogeneity measured
8   within a 25 km buffer around sites (Fisher's C = 0.59, p = 0.74, 2 degrees of freedom). Genetic
9   diversity R^2_m = 0.04 (variance explained by fixed effects); R^2_c = 0.92 (variance explained by
10   fixed and random effects); species richness R^2 = 0.88.

Response	Predictor	Estimate ± SE
gene diversity	potential evapotranspiration	0.20 ± 0.16
gene diversity	actual evapotranspiration	0.07 ± 0.16
gene diversity	heterogeneity	0.19 ± 0.12
species richness	potential evapotranspiration	0.19 ± 0.01
species richness	actual evapotranspiration	0.90 ± 0.02
species richness	heterogeneity	0.10 ± 0.02

13 **Table S3.** Path coefficients and standard errors for SEM model with heterogeneity measured
14 within a 40 km buffer around sites (Fisher's C = 0.49, $p = 0.78$, 2 degrees of freedom). Genetic
15 diversity $R^2_m = 0.04$ (variance explained by fixed effects); $R^2_c = 0.93$ (variance explained by
16 fixed and random effects); species richness $R^2 = 0.89$.

Response	Predictor	Estimate ± SE
gene diversity	potential evapotranspiration	0.22 ± 0.17
gene diversity	actual evapotranspiration	0.10 ± 0.17
gene diversity	heterogeneity	0.15 ± 0.11
species richness	potential evapotranspiration	0.20 ± 0.01
species richness	actual evapotranspiration	0.91 ± 0.01
species richness	heterogeneity	0.09 ± 0.01

20 **Table S4.** Path coefficients and standard errors for SEM model with heterogeneity measured
21 within an 80 km buffer around sites (Fisher's C = 0.40, p = 0.82, 2 degrees of freedom). Genetic
22 diversity R^2_m = 0.03 (variance explained by fixed effects); R^2_c = 0.92 (variance explained by
23 fixed and random effects); species richness R^2 = 0.90.

Response	Predictor	Estimate ± SE
gene diversity	potential evapotranspiration	0.16 ± 0.19
gene diversity	actual evapotranspiration	0.24 ± 0.20
gene diversity	heterogeneity	0.09 ± 0.09
species richness	potential evapotranspiration	0.22 ± 0.01
species richness	actual evapotranspiration	0.95 ± 0.01
species richness	heterogeneity	0.07 ± 0.01