Deletion of *Serf2* shifts amyloid conformation in an Aβ amyloid mouse model

E. Stroo[1*], L. Janssen[1*], O. Sin[1], W. Hogewerf[1], M. Koster[2], L. Harkema[2], S.A. Youssef[2,3], N. Beschorner[4], A.H.G. Wolters[5], B. Bakker[1], A. Thathiah[6,7], F. Foijer[1], B. van de Sluis[2], J. van Deursen[8], M. Jucker[4], A. de Bruin[2,3], E.A.A. Nollen[1]

[1]European Research Institute for the Biology of Ageing, University of Groningen, University Medical Centre Groningen, The Netherlands

[2]Department of Biomolecular Health Sciences, Faculty of Veterinary Medicine, Utrecht University, The Netherlands

[3]Department of Pediatrics, Molecular Genetics Section, University of Groningen, University Medical Centre Groningen, The Netherlands

[4]Department of Cellular Neurology, Hertie-Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany.

[5]Department of Biomedical Sciences of Cells and Systems, University Medical Centre Groningen, The Netherlands

[6]VIB Center for the Biology of Disease, KU Leuven Center for Human Genetics, University of Leuven, Leuven, Belgium

[7] Department of Neurobiology, University of Pittsburgh Brain Institute, University of Pittsburgh School of Medicine, Pittsburgh, PA USA

[8]Mayo Clinic, United States of America

*These authors contributed equally

[#]Correspondence: e.a.a.nollen@umcg.nl

Keywords: Small EDRK-rich factor 2, SERF, protein aggregation, amyloid-beta, amyloid polymorphisms

Highlights

- Loss of SERF2 slows embryonic development and causes perinatal lethality
- SERF2 affects proliferation in a cell-autonomous fashion
- Brain-specific *Serf2* knockout does not affect viability or Aβ production
- Brain deletion of *Serf2* shifts the amyloid conformation of Aβ

Abstract

Neurodegenerative diseases like Alzheimer, Parkinson and Huntington disease are characterized by

aggregation-prone proteins that form amyloid fibrils through a nucleation process. Despite the shared

β-sheet structure, recent research has shown that structurally different polymorphs exist within fibrils

of the same protein. These polymorphs are associated with varying levels of toxicity and different

disease phenotypes. MOAG-4 and its human orthologs SERF1 and SERF2 have previously been

shown to modify the nucleation and drive amyloid formation and protein toxicity *in vitro* and in *C.*

elegans. To further explore these findings, we generated a *Serf2* knockout (KO) mouse model and

crossed it with the *APPPS1* mouse model for Aβ amyloid pathology. Full-body KO of *Serf2* resulted

in a developmental delay and perinatal lethality due to insufficient lung maturation. Therefore, we

proceeded with a brain-specific *Serf2* KO, which was found to be viable. We examined the Aβ

pathology at 1 and 3 months of age, which is before and after the start of amyloid deposition. We

show that SERF2 deficiency does not affect the production and overall Aβ levels. *Serf2* KO-*APPPS1*

mice displayed an increased intracellular Aβ accumulation at 1 month and a higher number of Aβ

deposits compared to *APPPS1* mice with similar Aβ levels. Moreover, conformation-specific dyes and

electron microscopy revealed a difference in the structure and amyloid content of these Aβ deposits.

Together, our results reveal that SERF2 causes a structural shift in Aβ aggregation in a mammalian

brain. These findings indicate that a single endogenous factor may contribute to amyloid

polymorphisms, allowing for new insights into this phenomenon's contribution to disease

manifestation.

Introduction

Protein aggregation is a pathological hallmark shared by several age-related neurodegenerative diseases, such as Alzheimer (AD), Parkinson and Huntington disease. The amyloid-like aggregates that accumulate in each of these diseases are composed of disease-specific proteins, i.e., amyloid-beta (Aβ) and tau in AD(Glenner & Wong, 1984; Masters *et al*, 1985; Grundke-Iqbal *et al*, 1986a, 1986b), alpha-synuclein (α-Syn) in Parkinson disease(Langley & Charlesworth, 1997) and Huntingtin (HTT) in Huntington disease(Scherzinger *et al*, 1999). While the exact molecular mechanisms underlying the disease pathology remain to be elucidated, genetic evidence indicates that these aggregation-prone proteins play a key role in the disease processes. In AD multiple causative mutations have been found in three genes: the amyloid precursor protein (*APP*)(Goate *et al*, 1991; Chartier-Harlin *et al*, 1991; Citron *et al*, 1992; Eckman *et al*, 1997; Murrell *et al*, 2000; Kumar-Singh *et al*, 2000; Cruts *et al*, 2003; Kumar-Singh *et al*, 2006; Tomiyama *et al*, 2008), presenilin-1 (*PSEN1*)(Schellenberg *et al*, 1992; Sherrington *et al*, 1995; Borchelt *et al*, 1996; Lemere *et al*, 1996; Guerreiro *et al*, 2010) and presenilin-2 (*PSEN2*)(Rogaev *et al*, 1995; Citron *et al*, 1997; Müller *et al*, 2014; Guerreiro *et al*, 2010) (for a full list of known AD-linked mutations see https://www.alzforum.org/mutations). The *APP* gene encodes the precursor protein from which Aβ is produced through sequential cleavage by a β- and γ-secretase. The *PSEN* genes encode the main catalytic subunit of the γ-secretase complex. The fact that all the causative mutations affect the production, length and/or structure of the Aβ peptides and are sufficient to cause the disease(Chartier-Harlin *et al*, 1991; Citron *et al*, 1997; Cruts *et al*, 2003; Eckman *et al*, 1997; Guerreiro *et al*, 2010; Goate *et al*, 1991; Kumar-Singh *et al*, 2000, 2006; Lemere *et al*, 1996; Müller *et al*, 2014; Murrell *et al*, 2000; Rogaev *et al*, 1995; Schellenberg *et al*, 1992; Sherrington *et al*, 1995; Tomiyama *et al*, 2008; Borchelt *et al*, 1996; Citron *et al*, 1992), is arguably the strongest piece of evidence for the involvement of Aβ in the disease process. Causative mutations generally affect the Aβ peptide in one of three ways: (i) an overall increase in Aβ production(Haass *et al*, 1995; Citron *et al*, 1992), (ii) a shift in the production of shorter variants, like $A\beta_{40,}$ to more aggregation-prone, longer variants, like $A\beta_{42}$ (or $A\beta_{43}$)(Suzuki *et al*, 1994; Duff *et al*, 1996; Murayama *et al*, 1999; Murrell *et al*, 2000; Borchelt *et al*, 1996), (iii) Aβ peptides with a modified core region,

resulting in an increased propensity for aggregation(Nilsberth *et al*, 2001; Miravalle *et al*, 2000; Ni *et*

al, 2011; Tomiyama *et al*, 2008). Taken together, these mutations do not necessarily increase the

amount of Aβ, but rather increase the likelihood of Aβ aggregation. Similarly, several mutations

in the *SNCA* gene that have been linked to Parkinson disease have been shown to alter α-Syn

aggregation(Conway *et al*, 1998; Narhi *et al*, 1999; Conway *et al*, 2000; Ghosh *et al*, 2013, 2014; Choi

et al, 2004). In Huntington disease, any expansion of a polyglutamine stretch in the *HTT* gene that

exceeds 36 repeats has been shown to facilitate HTT aggregation and cause the disease(Cooper *et al*,

1998). Hence, changes in the folding and aggregation of these disease proteins are believed to be at the

basis of the observed pathology. In addition, there is increasing evidence that distinct structural

variants exist between aggregates of the same disease protein and that these polymorphs are associated

with varying degrees of toxicity(Nekooki-Machida *et al*, 2009; Sun *et al*, 2015; Cohen *et al*, 2015;

Qiang *et al*, 2017; Bousset *et al*, 2013; Heise *et al*, 2008; Li *et al*, 2018; Guerrero-Ferreira *et al*, 2019;

Gath *et al*, 2014; Cohen *et al*, 2009) . For instance, recent studies with brain samples from human AD

patients have shown that variations in the disease type and disease progression correlate with the

presence of structurally different aggregates and fibrils(Lu *et al*, 2013; Rasmussen *et al*, 2017; Cohen

et al, 2015; Qiang *et al*, 2017). The knowledge that small changes in the structural conformation of

these aggregates can have far-reaching effects on the disease manifestation could provide a new

avenue for the development of therapeutics. Until now, most research into potential treatments of AD

has focused on either reducing the overall amount of aggregating proteins or preventing aggregation,

not modifying the aggregation process.

In a previous study, our group performed a genetic screen in *Caenorhabditis elegans* (*C. elegans*) and

identified a small, intrinsically disordered protein of unknown function with the capacity to modify

protein aggregation and to enhance the resulting toxicity of multiple disease-related, aggregation-

prone proteins(van Ham *et al*, 2010). This peptide was dubbed *mo*difier of *ag*gregation-4 (MOAG-4)

and was found to be an ortholog of two human genes: *Serf1A* and *Serf2*. Both SERF1A and SERF2

were shown to exacerbate protein aggregation and toxicity in human neuroblastoma and HEK293 cell

lines as well(van Ham *et al*, 2010). *In vitro* assays with MOAG-4 and SERF1A have demonstrated

that these proteins can specifically accelerate the aggregation kinetics of amyloidogenic proteins, but

do not affect aggregation of non-amyloidogenic proteins(Falsone *et al*, 2012; Yoshimura *et al*, 2017).

For α-Syn, this amyloid-promoting effect has been shown to depend on the transient interaction of

these positively-charged peptides with negatively-charged regions and the disruption intramolecular

electrostatic interactions within the protein(Merle *et al*, 2019; Falsone *et al*, 2012; Yoshimura *et al*,

2017). Removal of MOAG-4 or SERF from a biological system did not change the overall levels of

the aggregation-prone proteins or activate chaperone pathways known to suppress aggregation(van

Ham *et al*, 2010). Therefore, the reduced toxicity would appear to rely solely on the structural shift in

the aggregation process(van Ham *et al*, 2010; Falsone *et al*, 2012). In this study, we investigate

whether the removal of SERF2 modifies the aggregation of Aβ in the biologically more complex

environment of the mammalian brain.

Results

Full-body Serf2 KO results in developmental delay and perinatal lethality in mice

To establish the role of SERF2 *in vivo*, full-body *Serf2* knockout (KO) mice were generated (Figure

S1A-C). During an initial cross, using the four homozygous *Serf2* KO animal (*Serf2^{-/-}*) and eight

heterozygous *Serf2* KO animals (*Serf2^{+/-}*) we obtained, only one *Serf2^{-/-}* survived the day of birth

compared to 27 *Serf2^{+/-}* and 96 *Serf2* wild-type (*Serf2^{+/+}*) mice (Supplemental Table 1). After

backcrossing into C57BL/6J background, interbreeding of *Serf2^{+/-}* animals did not result in any viable

Serf2^{-/-} mice at the time of weaning (Figure 1A,1B). To determine whether *Serf2^{-/-}* mice died during

fetal development, we examined the offspring at multiple developmental stages: embryonic day 13.5,

15.5 and 17.5. In all of the examined embryonic stages we observed the expected Mendelian ratio of

genotypes, around 25 % *Serf2^{+/+}* and *Serf2^{-/-}* embryos and around 50 % *Serf2^{+/-}* (Chi Square: E13.5

p=0.9672; E15.5 p= 0.4432; E17.5 p=0.7674) (Figure 1B). At birth (P0), all *Serf2^{-/-}* displayed

respiratory distress and the majority (5/7) died within 30 minutes of being born. At the time of

weaning, the Mendelian ratio was maintained between the *Serf2^{+/+}* and *Serf2^{+/-}* animals, but no *Serf2^{-/-}*

mice survived until this stage (Chi Square: P22 p<0.0001) . Examination of *Serf2* mRNA expression at

E13.5 showed a 50 % reduction for *Serf2^{+/-}* and complete KO for *Serf2^{-/-}* compared to *Serf2^{+/+}* (One-

way ANOVA: *Serf2^{+/-}* and *Serf2^{-/-}* vs *Serf2^{+/+}* p_{Bon}<0.0001) (Figure 1C). A similar ratio in SERF2

expression was also observed on the protein level (Figure 1D). The *Serf1* mRNA levels were not

affected by the change in *Serf2* levels, indicating there is no compensatory mechanism between these

genes (One-way ANOVA: p=0.2403) (Figure 1C). Further analysis of the *Serf2^{-/-}* embryonic

phenotype revealed a reduction in embryo size of *Serf2^{-/-}* compared to *Serf2^{+/+}* and *Serf2^{+/-}* from E15.5

until birth (Two-way ANOVA: E17.5 WT vs KO p_{bon}<0.05; P0 WT vs ko p_{Bon}<0.0001) (Figure 1E,

S1D-E). No differences in size could be observed between *Serf2^{+/+}* and *Serf2^{+/-}* embryos during the

embryonic stage leading up to birth (P0). Histological examination of various tissues revealed a

developmental delay of approximately one to two days in the organs of E17.5 *Serf2^{-/-}* embryos

compared to the previous time points and standard embryonic development(Kaufman, 1992; Kaufman

& Bard, 1999). This was most prominently in the lungs, through reduced expansion of alveoli and

augmented tissue density, and in the kidneys, by the increased amount of mesenchyme between the

tubules and glomeruli (Figure S1F). At the time of birth, the lungs of $Serf2^{-/-}$ pups displayed a

significantly reduced saccular space (t-test: p= 0.0088) (Figure 1F), reflecting insufficient lung

expansion and maturation. This partial fetal atelectasis seems to explain the observed respiratory

distress and a likely cause of the perinatal lethality in $Serf2^{-/-}$ pups. Placental abnormalities have been

shown to cause impaired transfer of nutrients and oxygen to the embryo, resulting in growth

delays(Ward *et al*, 2012). To exclude this as a potential cause, we examined the placenta at all three

embryonic stages for microscopic lesions in the placental labyrinth or other changes. No genotype-

related pathology could be found, indicating that absence of SERF2 in the embryo itself is responsible

for the observed growth phenotype. This seems to be supported by the fact that CRISPR-induced

$Serf2^{-/-}$ clones of HEK293T cells display a delay in proliferation (Figure S2A). Moreover, RNA

sequencing analysis of mouse embryonic fibroblasts (MEF) isolated from E13.5 embryos showed a

clear enrichment of GO-terms clustered around three closely related biological functions: cell cycle,

cell proliferation and cell adhesion (Supplemental Figure S2A-B). Interestingly, in the cell cycle

cluster we observe an overall upregulation of factors driving cell division and downregulation of

growth arrest factors in $Serf2^{-/-}$ MEFs (Supplemental Table 2). This observation seems in line with a

developmental delay phenotype in $Serf2^{-/-}$ mice, because it is known that cell proliferation is favored

over differentiation in earlier stages of embryonic development compared to later stages(Ciemerych &

Sicinski, 2005). Overall, the cell culture results point towards a cell-autonomous effect of SERF2 and

suggest that loss of *Serf2* results in delayed growth. While our previous study also demonstrated a cell-

autonomous function for moag-4, it did not show any effects of moag-4 deletion on the viability, life

span or size of *C. elegans*(van Ham *et al*, 2010). All together, our results suggest that *Serf2* KO causes

delayed maturation of certain organs at the time of birth resulting in death. Given the cell-autonomous

nature of the effects of SERF2, we aimed to avoid these adverse effects by creating an organ-specific

conditional KO model.

Brain-specific KO of Serf2 does not affect viability or the Aβ production pathway

To study the effect of *Serf2* KO on amyloid pathology in the brain, we circumvented the perinatal

lethality of full-body *Serf2* KO mice by generating a brain-specific *Serf2* KO mouse model (*Serf2*$^{br-/-}$).

The conditional KO was obtained by combining our *Serf2*$^{flox/flox}$ mice with a *Sox1-Cre*

model(Takashima *et al*, 2007) (Figure S3A). SOX1 is one of the earliest transcription factors

expressed in neuronal precursors of the ectoderm from embryonic day 7.5(Pevny *et al*, 1998; Wood &

Episkopou, 1999). Consequently, the *Sox1* promotor was selected as a pan-neuronal driver of

Recombinase cre expression to restrict homologous recombination to the central nervous

system(Takashima *et al*, 2007). Analysis of *Serf2* expression on the mRNA and protein level in

various organs confirmed brain-specific ablation of *Serf2* by *Sox1*-mediated *Cre* expression in our

Serf2$^{flox/flox}$ mice (t-test WT vs *Serf2*$^{br-/-}$ brain: 1m p=0.0022; 3m p=0.00527) (Figure S3B-C). This

conditional KO was crossed with the *APPPS1-21* amyloid (AM) model(Radde *et al*, 2006), which

contains human transgenes for both APP with the Swedish KM670/671NL mutation and PSEN1 with

the L66P mutation. From this cross we obtained four experimental groups: WT, *Serf2*$^{br-/-}$, AM and

AM;*Serf2*$^{br-/-}$ (Figure 2A). Upon crossing of the *Serf2*$^{br-/-}$ with the amyloid model, we observed the

expected Mendelian ratio for all the genotypes at P22, indicating that the viability is not significantly

affected by brain-specific KO of *Serf2* (Chi Square: p= 0.37)(Supplemental Table 3). Based on the

known progression of Aβ plaque pathology in the AM mice, we selected two age groups for further

analysis: before (1 month) and after (3 months) Aβ plaque deposition (Figure 2B). Analysis of the

Serf2 mRNA levels in both age groups showed that *Serf2* expression was not altered by the plaque

pathology and confirmed the KO in the *Serf2*$^{br-/-}$ and AM;*Serf2*$^{br-/-}$ mice. Some residual *Serf2*

expression could be detected in the brains of the KO groups likely due to the presence of cells from

non-neuronal lineage in the samples, e.g. cells from circulatory system (Two-way ANOVA: *Serf2*$^{br-/-}$

and AM;*Serf2*$^{br-/-}$ vs WT 1m p$_{bon}$ <0.001; 3m p$_{bon}$ <0.0001) (Figure 2C). These results were also

confirmed on the protein level using western blot (Figure 2D). In accordance with the effects observed

in the full-body *Serf2*$^{-/-}$ mice, the *Serf2*$^{br-/-}$ and the AM;*Serf2*$^{br-/-}$ mice did show a reduction in brain

weight compared to WT in both age groups (Two-way ANOVA: *Serf2*$^{br-/-}$ vs WT both ages p$_{bon}$<0.01;

AM;*Serf2*$^{br-/-}$ vs WT 1m p$_{bon}$<0.0001; 3m p$_{bon}$<0.001) (Figure 2E). While the weight reduction seems

to be slightly more pronounced in AM;*Serf2*$^{br-/-}$ compared to *Serf2*$^{br-/-}$, AM by itself displayed no

significant change in brain weight. This would indicate that *Serf2* ablation, *Cre* expression or a

combination of both is most likely responsible for the difference in weight. However, AM mice may

be more susceptible to this effect (Figure 2E). To distinguish between the effects of *Cre* expression

and *Serf2* KO, we also examined the *Sox1-Cre* mice(Takashima *et al*, 2007) without floxed *Serf2*

(*Cre^{br}*) and crossed them with *APPPS1-21*(Radde *et al*, 2006) without floxed *Serf2*. The *Cre*^{br} mice

also showed a reduction in brain weight compared to the WT controls, which was more pronounced at

1 month (One-Way ANOVA: *Cre^{br}* vs WT 1m $p_{bon}<0.05$) (Figure S3D), but was overall less extensive

than the reduction in *Serf2*^{br-/-} animals (One-way ANOVA: *Cre^{br}* vs WT $p_{bon}<0.01$; AM;*Cre^{br}* vs WT

$p_{bon}<0.001$) (Figure 2F). Together, this suggests that *Cre* expression is partially responsible for the

observed decrease in brain size, particularly at 1 month of age. Due to ongoing brain maturation

between 1 and 3 months, all groups show a significant increase in brain weight (Two-way Anova: Age

$p<0.0001$), but this maturation appeared unaffected by *Serf2* KO. Moreover, the difference in brain

weight between WT and *Serf2*^{br-/-}remained constant beyond the age of 3 months up to 11 months of

age (Figure S3E). This suggests that the reduction in brain size takes place during specific stages of

brain development, but that the continued development is not necessarily altered by these changes.

Additional histological examination of the brains with hematoxylin-eosin staining revealed no

difference in the cell density (Figure 2G), nor did we find any evidence of degeneration, apoptosis or

necrosis. Apart from the overall reduction in weight, the brains showed no structural abnormalities and

the tissue appeared healthy.

Next, we examined whether the brain KO of *Serf2* affected any of the key components in the Aβ

aggregation process, like altering the levels of APP or the production of its cleavage products, Aβ$_{40}$

and Aβ$_{42}$. We determined the expression of human *APP* in WT, *Serf2*^{br-/-}, AM and AM;*Serf2*^{br-/-} mice

at both 1 and 3 months of age and observed no difference due to brain specific *Serf2* KO (t-test: 1m

p=0.9170; 3m p=0.9963) (Figure 2H, data from WT, *Serf2*^{br-/-} not included due to lack of APP

construct and absence of signal). Western blot analysis confirmed there was also no difference in APP

protein levels between AM and AM;*Serf2*^{br-/-} (t-test: 1m p= 0.1157;3m p=0.5908) (Figure 2I and J).

Further analysis of Aβ$_{40}$ (Two-way ANOVA: Genotype p=0.8841; Age $p<0.0001$;Interaction

p=0.8841) (Figure 2K) and Aβ_{42} (Two-way ANOVA: Genotype p=0.7006; Age p<0.0001 ;Interaction

p=0.6522) (Figure 2L) showed an increase in concentration between 1 and 3 months, but this was the

same in both AM and AM;*Serf2*$^{br-/-}$. Given the variability in the Aβ concentrations at 3 months and the

fact that the ratio between these two peptides is known to affect the aggregation process, we also

investigated the correlation between these two Aβ levels (Figure 2M). This analysis showed that mice

with a high Aβ_{42} concentration display a similarly high Aβ_{40} level, maintaining a comparable Aβ_{40}

/Aβ_{42} ratio for both AM and AM;*Serf2*$^{br-/-}$ animals. Taken together, this data suggests that SERF2 does

not affect the levels of APP and its cleavage products, Aβ_{40} and Aβ_{42}. These findings are in accordance

with the previously observed effects of MOAG-4 on the aggregation of polyglutamine, Aβ and αSyn

in *C. elegans*, which modifies the aggregation process without changing the overall levels of these

aggregation-prone proteins(van Ham *et al*, 2010). While the difference in brain size is definitely a

factor to be considered during further analysis, given the fact that we observed no obvious changes in

the overall health, continued development and APP processing pathway of these brain-specific *Serf2*

KO mice, we proceeded with our analysis of Aβ pathology in these animals.

Serf2 KO alters the amount of Aβ deposition in the brain

We next investigated whether SERF2 affects the Aβ aggregation by performing immunohistological

analysis of the Aβ plaque pathology. Initially, we performed a general Aβ staining using the 6E10

antibody. In accordance with the known progression of Aβ plaque pathology in the amyloid model, we

only found extracellular deposits in the 3-month-old AM and AM;*Serf2*$^{br-/-}$ mice. As expected, most

plaques were found in the cortex, with some pathology beginning to spread to the hippocampus as

well (Figure S4). Microscopic inspection of the Aβ deposits revealed a broad range of sizes with a

wide variety in staining intensity and patterns, but we found no clear morphological differences

between the plaque population found in AM and AM;*Serf2*$^{br-/-}$ (Figure 3A, S4). Quantification of the

6E10-positive deposits showed a slight increase in AM;*Serf2*$^{br-/-}$ compared to AM, but this just failed

to reach statistical significance in the Bonferroni corrected post-hoc comparison (Two-way ANOVA:

all 3m non-AM vs AM groups p$_{bon}$ <0.0001; AM vs AM;*Serf2*$^{br-/-}$ p$_{bon}$ =0.063)(Figure 3B). Given the

high variability in plaque load between animals within the same experimental group, we examined

whether this variability reflected the previously observed variation in Aβ levels. Unsurprisingly,

animals with higher levels of $A\beta_{40}$ (Figure 3C) and $A\beta_{42}$ (Figure 3D) displayed a higher overall plaque

load in both groups. More interestingly, our linear regression analysis revealed that AM;$Serf2^{br-/-}$ mice

tend to have a significantly higher plaque load than AM mice with similar Aβ levels (Supplemental

Table 4). A second general Aβ stain with the W0-2 antibody displayed a similar slight increase in the

amount of plaque deposits in AM;$Serf2^{br-/-}$ at 3 months (Two-way ANOVA: all 3m non-AM vs AM

groups p_{bon} <0.0001; AM vs AM;$Serf2^{br-/-}$ p_{bon} =0.101) (Figure 3E). In fact, a comparison between the

6E10 and the W0-2 staining in individual animals confirmed the similarity between both staining

patterns at 3 months (Figure 3F, S5, Supplemental Table S3). Interestingly, our detection algorithm

also picked up a small increase in the counts of W0-2-positive foci in the 1 month AM;$Serf2^{br-/-}$ group

(Two-way ANOVA: 1m WT vs AM;$Serf2^{br-/-}$ p_{bon} =0.088 ; $Serf2^{br-/-}$ vs AM;$Serf2^{br-/-}$ p_{bon} =0.081; AM

vs AM;$Serf2^{br-/-}$ p_{bon} =0.127; all other comparisons p_{bon} >0.999) (Figure 3E). Closer examination of the

microscopic images revealed the identified spots were not extracellular plaques, but were the result of

increased levels of W0-2-positive intracellular staining in AM;$Serf2^{br-/-}$ (Figure S7A). Some low level

intracellular W0-2 and 6E10 staining could also be observed in the other 1-month-old AM mice, but

not to the extent that it was picked up by the detection algorithm (Figure S7A-B). This would suggest

that the deletion of *Serf2* already alters the intracellular accumulation of Aβ prior to plaque deposition.

Overall, these results indicate that the presence of SERF2 is indeed able to modify Aβ aggregation in

the mouse brain.

SERF2 deficiency changes the proportion of ThS-positive cores, but not the immunoreactivity
of deposits

To investigate if SERF2 also affected the amount of Thioflavin-positive fibrils, we performed

Thioflavin-S (ThS) staining. Here, we again observed a large variety in the amount of ThS-positive

plaques between individual animals, but we found no mean difference between AM and AM;$Serf2^{br-/-}$

mice (Two-way ANOVA: all 3m non-AM vs AM groups p_{bon} <0.001) (Figure 3G). However, further

analysis of the correlation between Aβ levels and the amount of ThS-positive plaques revealed that

AM mice showed a positive correlation similar to the general Aβ stainings, while AM;$Serf2^{br-/-}$ mice

displayed an inverse relationship (Figure 3H, 3I, Supplemental Table S3). Consequently, a general

increase in Aβ deposits does not equate to an increase in amyloid deposition in AM;*Serf2*[br-/-] mice, as

it does in AM mice (Figure 3J), further demonstrating a change in the amyloid formation in *Serf2* KO

mice. In addition, we investigated microglia activation and whether we could detect a change in the

gliosis associated with Aβ plaques. Staining for Iba1 showed a slight increase in the AM;*Serf2*[br-/-] mice

compared to AM (Two-way ANOVA: 3m non-AM groups vs AM p_{bon} <0.01; non-AM groups vs

AM;*Serf2*[br-/-] p_{bon} <0.0001; AM vs AM;*Serf2*[br-/-] p_{bon} <0.01) (Figure 3K), similar to what was observed

with the 6E10 staining (Figure 3B). We therefore examined if this increase was simply related to the

increase in Aβ deposits or if AM;*Serf2*[br-/-] also displayed a different immune response to plaques

compared to AM. Our analysis revealed a similar correlation between 6E10-positive deposits and

immune response for AM and AM;*Serf2*[br-/-] mice, suggesting the deposits evoked a similar immune

response in both groups(Figure S6A, Supplemental Table S3).

Knockout of Serf2 changes the plaque composition and fibril structure

Given the differences in the reactivity of the Aβ deposits between the various stains, we decided to

explore the composition within the individual plaques in greater detail. To this end, we made use of

two luminescent conjugated oligothiophenes (LCO), qFTAA (quadro-formylthiophene acetic acid)

and hFTAA (hepta-formylthiophene acetic acid). These LCOs are small hydrophobic ligands with a

high affinity for binding distinct types of amyloids and can easily be distinguished from each other

based on their spectral properties(Klingstedt *et al*, 2011). The smallest of the two molecules, qFTAA,

is characterized by a peak in its emission spectrum around 500 nm and has been shown to stain

compact, multifilament fibril structures, coinciding with ThT and Congo red staining(Psonka-

Antonczyk *et al*, 2016; Nyström *et al*, 2013; Klingstedt *et al*, 2011). Whereas hFTAA emits in the

yellow to red range with a peak around 600 nm and is capable of staining prefibrillar (non-

thioflavinophilic) and single fibril aggregates found during the lag-phase of in vitro ThT kinetics

(Figure 4A). The prefibrillar aggregates were shown to be stained with 6E10, but not Congo red, ThT

or qFTAA(Klingstedt *et al*, 2011; Nyström *et al*, 2013; Psonka-Antonczyk *et al*, 2016). Previous

studies have shown that by combining LCO dyes it is possible to uncover structural differences in

plaque composition(Lord *et al*, 2011; Nyström *et al*, 2013; Rasmussen *et al*, 2017). We performed

confocal spectral imaging of randomly selected plaques in 3-month-old AM and AM;*Serf2*$^{br-/-}$ mice

after double staining with both LCOs and determined the ratio of the light intensity emitted at 502nm

and 588nm as a measure for the relative amounts of fibrillar and prefibrillar aggregates within the

individual plaques. An exploratory frequency analysis of the intensity ratios found in both groups

revealed a three-peak pattern in which the middle peak was similarly represented in both groups

(Figure 4B, green area). The first peak, however, was predominantly found in AM mice (Figure 4B,

red area), while the third peak was mainly detected in AM;*Serf2*$^{br-/-}$ (Figure 4B, blue area). This shift

in ratio was also reflected by a significant difference in the average intensity ratio between AM and

AM;*Serf2*$^{br-/-}$ (t-test p<0.001) (Figure 4C). This increase in intensity ratio could not be observed in

AM;*Cre*^{br} mice (t-test p=0.6047) (Figure 4C, insert), indicating that this effect is caused by the *Serf2*

KO and unrelated to *Cre* expression. Given the fact that AM;*Cre*^{br} animals also show a decrease in

brain weight without a change in LCO staining, this would also suggest that the weight reduction is

not a factor in these results. Although, as the change in brain weight is more pronounced in *Serf2*$^{br-/-}$

animals, we cannot completely exclude this as a potential contributing factor. Next, we examined if

the change in the intensity ratio correlated with the slight increase in 6E10-positive deposits we

observed in AM;*Serf2*$^{br-/-}$ (Figure 4D). However, the higher average intensity ratio in AM;*Serf2*$^{br-/-}$

mice was not related to this general plaque load nor the number of W0-2- or ThS-positive deposits

(Figure S6B, S6C). Further microscopic analysis of low intensity ratio deposits from AM mice showed

a plaque with a qFTAA- and hFTAA-positive core and a border that was only stained by hFTAA

(Figure 4E, left panels). The high intensity ratio deposits from AM;*Serf2*$^{br-/-}$ mice, on the other hand,

revealed a qFTAA-positive core, but virtually no hFTAA-staining (Figure 4E, right panels),

demonstrating the difference in plaque composition reflected by the shift in LCO intensity ratios.

Additionally, we visualized the global fibrillar structure of the plaques on a nanometer scale using

high resolution, scanning transmission electron microscopy (STEM). Overall, STEM images of

plaques from AM;*Serf2*$^{br-/-}$ tended to show more condensed plaques composed of short, thick and

densely packed bundles of fibers with little space in between (Figure 4F, right panels). In the AM

mice, however, the plaques we observed displayed more loosely packed fibrils (Figure 4F, left panels).

Taken together, these findings suggest *Serf2* KO in mice leads to a shift in the Aβ aggregation process,

resulting in an altered structural composition of the plaques.

Discussion

Previous studies identified MOAG-4 and its human orthologs SERF1A and SERF2 and their potential

to modify the aggregation of amyloidogenic proteins. While the exact endogenous biological function

of these peptides remains unclear, a recent study has provided evidence for RNA-binding properties of

SERF1A through a region which is highly conserved in MOAG-4 and SERF2(Meyer *et al*, 2019). In

this study, we have demonstrated in cells and mice that SERF2 provides a growth advantage during

development. The absence of SERF2 or MOAG-4 in itself does not appear to be lethal in cells, worms

or the brain-specific KO mice. Therefore, we postulate that the observed perinatal lethality in the full-

body KO mice is a secondary effect of the delay in growth due to insufficient maturation of certain

organs at birth. This appears to be supported by the fact that some pups survived the initial minutes

after birth and incomplete penetrance of the lethality in the earliest generations with mixed genetic

background. The 129SV genetic background present in these mice is known to have a gestation period

that is on average around one day longer than C57/Bl6J mice(Murray *et al*, 2010), which could result

in improved lung maturation. While the exact function of SERF2 remains unknown, we propose its

effects on growth and development are most likely caused through one of two mechanisms: either

SERF2 facilitates cell proliferation or it protects cells from stressors that create an unfavorable

environment for cell division. In the latter case, we would expect to see an upregulation of stress

pathways or cell cycle arrest proteins inhibiting the cell cycle in $Serf2^{-/-}$ cells. Interestingly, the RNA

seq analysis of the MEFs from our $Serf2^{-/-}$ mice showed the exact opposite. Cell cycle driving factors

were upregulated (e.g. cyclins). Whereas, stress signaling and cell cycle inhibiting factors were

downregulated (e.g. GADD45 and GAS1). Thus, the $Serf2^{-/-}$ cells actually displayed an increased

stimulation of cell proliferation mechanisms, which would fit with cells from earlier stages of

embryonic development. This would suggest the first mechanism, with a beneficial role for SERF2 in

the efficient execution of cell proliferation, as the most likely explanation for the observed growth

effect. Furthermore, an increase in cellular stress would most likely result in decreased health and

fitness. However, *moag-4* deletion worms showed no changes in life span compared to controls(van

Ham *et al*, 2010) and, apart from the initial growth deficit, the tissues of *Serf2* KO mice appeared

healthy and continued to develop similarly to controls after birth. A recent study showed SERF1a

might play a role as an RNA-organizing protein, which localizes to membraneless nuclear organelles,

like the nucleolus, under normal physiological conditions. However, under stress conditions, it was

shown to migrate to the cytoplasm, where it could drive amyloid toxicity.(Meyer *et al*, 2019) While

there was no compensatory upregulation of SERF1 in response to SERF2 KO and a similar mode of

action remains to be demonstrated for SERF2, the structural homology of the two proteins would

suggest a comparable function. Given the importance of nucleolar disassembly and reassembly in cell

cycle control(Visintin & Amon, 2000; Leung *et al*, 2004), an RNA-organizing function could explain

how SERF2 facilitates cell proliferation. This developmental role would have limited biological

importance in non-proliferating, differentiated cells, like the brain's neurons or the cells of adult *C.*

elegans, and might also explain why we saw no adverse effects in the *moag-4* deletion worms or our

adult *Serf2*$^{br-/-}$ mice. While a reduced biological importance of SERF2 later in life could prove

interesting with regard to the treatment of age-related neurodegenerative disorders, further exploration

of SERF2's endogenous function and how it evolves with ageing will be needed to fully assess this

mechanism's therapeutic potential.

In this study, we confirmed the ability of SERF2 to modify the Aβ pathology *in vivo* in a mouse model

without changing the overall Aβ levels. We showed that mice lacking SERF2 were more prone to

form Aβ deposits and that the composition of these deposits was structurally different. In AM mice

higher levels of Aβ result in an increase in the numbers of Aβ deposits as well as an increase in the

number of ThS-positive deposits. AM;*Serf2*$^{br-/-}$ mice, on the other hand, also show an increase in Aβ

deposits, but this did not lead to higher numbers of ThS-positive plaques, indicating an altered

dynamic for amyloid formation. These findings were further corroborated by the LCO spectra, which

revealed that the plaques in AM;*Serf2*$^{br-/-}$ mice have a different conformation of pre-fibrillar and

fibrillar Aβ compared to AM mice. Finally, STEM imaging also confirmed a globally altered structure

of the amyloid fibrils in the plaques. Amyloid fibrils are formed through a nucleated self-assembly

process characterized by a structural conformation to a β-strand secondary structure and the formation

of a critical nucleus during the initial oligomerization. The nuclei act as seeds from which the fibrils

grow and have been shown to propagate a specific fibril structure(Petkova *et al*, 2005; Qiang *et al*,

2017). In vitro kinetic assays with SERF and MOAG-4 have shown that these peptides accelerate

aggregation by acting on the nucleation phase(Meinen *et al*, 2019; Falsone *et al*, 2012; Yoshimura *et*

al, 2017; Merle *et al*, 2019). Recently, we have shown that SERF2 is equally capable to drive amyloid

formation of α-Syn and Aβ *in vitro* through a conserved, positively charged region(Pras *et al*,

2020). Meanwhile, another study has demonstrated that binding of the intrinsically disordered protein

yeast SERF to α-Syn and Aβ results in fuzzy complexes with heterogeneous structural

conformations that are more extended in nature(Meinen *et al*, 2019). For α-Syn, the disruption of

intra-molecular electrostatic interactions by the positively charged region of MOAG-4/SERF1 was

found to expose an amyloid nucleation site(Merle *et al*, 2019; Yoshimura *et al*, 2017). Intra-molecular

electrostatic interactions have also been proven to play a part in the dynamics and structure of the Aβ

monomer folding, which is at the basis of nucleus formation. In fact, several of the familial AD

mutations located within the Aβ sequence and reported not to affect the Aβ levels, appear to exert their

effects by modifying the intra-molecular interactions and monomer folding(Grant *et al*, 2007; Lazo *et*

al, 2009; Elkins *et al*, 2016; Ni *et al*, 2011). These single amino acid modifications in the Aβ

sequences also result in alterations in the aggregation kinetics, fibril structure and aggregate toxicity in

in vitro studies(Ovchinnikova *et al*, 2011; Van Nostrand *et al*, 2001; Miravalle *et al*, 2000). Moreover,

patients and mouse models with these mutations show differences in the localization of Aβ

accumulation and deposition, as well as differences in its structure, giving rise to specific disease

phenotypes. For instance, while some mutations result in increased Aβ deposition and altered plaque

structure, other mutations at the same location result in cerebral amyloid angiopathy with Aβ

deposition in blood vessels rather than the brain parenchyma or even an absence of extracellular Aβ

deposition(Inayathullah & Teplow, 2011; Bugiani *et al*, 2010; Grabowski *et al*, 2001; Philipson *et al*,

2009; Shimada *et al*, 2011; Tomiyama *et al*, 2010; Grant *et al*, 2007; Ni *et al*, 2011; Nilsberth *et al*,

2001; Lord *et al*, 2011; Tomiyama *et al*, 2008; Van Nostrand *et al*, 2001). In addition, some of these

mutants showed increased intracellular Aβ accumulation which differed in structure-specific staining

characteristics(Lord *et al*, 2006; Kulic *et al*, 2012; Tomiyama *et al*, 2010). The differences in Aβ

pathology between these mutants seem to display some similarities to the changes we observed

between our AM and AM;*Serf2*^{br-/-} mice. Together with the *in vitro* findings about the mechanism of

SERF2 interaction with aggregation-prone proteins, this would appear to suggest modulation of intra-

molecular interactions and altered nucleation as the mechanism for SERF2's effect on aggregation and

amyloid formation in our mice. Interestingly, there is increasing evidence that qualitative, structural

properties may be more related to toxicity than the quantity of aggregates and deposits. Recent studies

have provided increasing evidence that structural polymorphs are associated with differences in

toxicity and different clinical phenotypes in sporadic and familial cases(Lu *et al*, 2013; Cohen *et al*,

2015; Rasmussen *et al*, 2017; Qiang *et al*, 2017; Cohen *et al*, 2009). One recent study demonstrated

that these structural conformations of distinct disease phenotypes could also be detected by differences

in the LCO spectra of the plaques and that these spectral properties could, at least partially, be

transmitted to a mouse model through seeding(Rasmussen *et al*, 2017). While the effect of familial

mutations on the formation of distinct polymorphs has already been explored by others(Philipson *et al*,

2012; Lord *et al*, 2006; Kulic *et al*, 2012; Rasmussen *et al*, 2017), our study provides the first evidence

of a single endogenous factor, separate from Aβ and its production pathway, contributing to a

structural shift of amyloid pathology in a mammalian system. Further research will be needed to

elucidate the exact structural changes at an atomic level and if they affect toxicity and disease

progression similarly to what was previously observed in *C. elegans*. This will provide new insights

into the structural properties and diversity of disease-protein aggregation, contributing to a better

understanding of the variability in disease manifestation and open up previously unexplored avenues

for therapeutic research.

Materials and methods

Animals

Serf2 knockout mice were generated by introducing *loxP* sites in the *Serf2* gene on either side of exon

2, in accordance with the 'knockout-first' allele design as described by Skarnes and

colleagues(Skarnes *et al*, 2011). Full-body *Serf2* knockout mice were obtained by crossing these

floxed mice with 129SV mice expressing *Cre* under the *Hprt* promoter, resulting in the removal of

exon 2 by *Cre*-mediated recombination (Figure S1A). Subsequently, these mice with mixed

background were backcrossed at least 6 times to a C57Bl/6J background. Homozygous and

heterozygous full-body *Serf2* knockout mice were examined with their wild-type littermates at

embryonic day 13.5, 15.5 and 17.5, as well as at the day of birth (P0) and day of weaning (P22).

Heterozygous *Sox1-Cre* mice(Takashima *et al*, 2007) were backcrossed at least 8 times to a C57BL/6J

background. These animals were also used as controls to differentiate between effects of *Cre*

expression and Serf2 knockout. To generate the conditional *Serf2* knockout mice, the floxed mice

were first crossed with Tg(ACTFLPe)9205Dym (Jackson#003800) mice to flip the FRT site and

remove the lacZ and neo cassette. Subsequently, these mice were backcrossed at least 6 times to a

C57BL/6J background. Finally, the resulting homozygous *Serf2*$^{\text{flox/flox}}$ mice were crossed with the

backcrossed *Sox1-Cre* mice, to obtain the brain-specific *Serf2* knockout mice (Figure S3A). The

conditional *Serf2* knockout mice were crossed with the APPPS1-21 transgenic mice (APP

(KM670/671NL)/PS1(L166P), Rottenburg, Germany) (Radde *et al*, 2006). A final cross between

heterozygous APPPS1-21; *Serf2*$^{\text{flox/flox}}$ and heterozygous *Sox1-Cre*; *Serf2*$^{\text{flox/flox}}$ resulted in four

experimental groups of *Serf2*$^{\text{flox/flox}}$ mice: *Sox1-Cre*$^{-/-}$; APPPS1-21$^{-/-}$ (WT), *Sox1-Cre*$^{+/-}$; APPPS1-21$^{-/-}$

(*Serf2*$^{\text{br-/-}}$), *Sox1-Cre*$^{-/-}$; APPPS1-21$^{+/-}$ (AM) and *Sox1-Cre*$^{+/-}$; APPPS1-21$^{+/-}$ (AM;*Serf2*$^{\text{br-/-}}$) (Figure 2A).

All experimental groups were tested at 1 month or 3 months of age.

All experiments were approved by the Institutional Animal Care and Use Committee of the University

of Groningen (Groningen, The Netherlands). All mice were maintained on a C57BL/6J background

and housed in a 12:12 hour light/dark cycle and the animals had ad libitum access to food and water.

Genotyping of embryos was performed using tail and yolk sac biopsies. Otherwise, ear biopsies were

used. DNA was purified using prepGEM® Tissue kit according to a protocol adapted from the

manufacturer (ZYGEPTI0500, ZyGEM, VWR International BV) and subjected to PCR using the

primers listed in Supplemental Table S5. Mice were terminated through CO_2-inhalation and cervical

dislocation for subsequent tissue collection.

Embryo processing and histochemistry

Embryos were fixed in 4 % formalin (Kinipath) for a minimum of 24 hours at room temperature. For

the pathological analysis, the embryos and their placenta were bisected longitudinally and embedded

in paraffin. Using the microm HM 340E (Thermo Scientific) 4 µm sections were cut for the

hematoxylin-eosin (HE) staining. The HE sections were incubated at 60 °C for 15 minutes. Next,

sections were deparaffinized and rehydrated in xylene (2x), 100 % alcohol (2x), 96 % alcohol, 70 %

alcohol and water. HE staining was performed by incubation with hematoxylin for 4 minutes, water

for 10 minutes, eosin for 1 minute and water for 10 seconds. After staining all sections were

dehydrated in 70 % alcohol, 96 % alcohol, 100 % alcohol (2x) and xylene (2x).

Brain processing

Upon termination, all brains were collected and weighed. From each experimental group three whole

brains were collected for HE staining according to the protocol described under embryo processing.

HE-stained sections were scanned with the TissueFAXs microscope using 20X objective. Images were

processed for nuclei counts using Histoquest software. The remaining brains were divided sagittally.

The left hemibrain was prepared for histochemistry and fixed in 4 % paraformaldehyde (PFA) for 48

hours at room temperature on a shaker. Next, the brains were placed in 30 % sucrose for

approximately 12 hours. Afterwards, the excess sucrose was removed and hemibrains were frozen in

molds with Tissue Tek O.C.T. compound (Sakura, The Netherlands) on dry ice. The right hemibrain

was snap frozen in liquid nitrogen and stored at -80°C. This tissue was homogenized using a liquid

nitrogen cooled mortar for subsequent protein and RNA analysis.

Quantitative RT-PCR

Total RNA was extracted from snap frozen tissue using TRIzol Reagent (Life Technologies) according

to the manufacturers' description. Total RNA quality and concentration were assessed using a

NanoDrop 2000 spectrophotometer (Thermo Scientific/Isogen Life Science). cDNA was made from

1,5 µg total RNA with a RevertAid H Minus First Strand cDNA Synthesis kit (Thermo Scientific)

using random hexamer primers. Quantitative real-time PCR was performed using a Roche LightCycler

480 Instrument II (Roche Diagnostics) with SYBR green dye (Bio-Rad Laboratories) to detect DNA

amplification. Relative transcript levels were quantitated using a standard curve of pooled cDNA

solutions. Expression levels were normalized to β-Actin or 18S mRNA levels. The primers for

Quantitative PCR used are listed in Supplemental Table S5.

Western Blot analysis

For SERF2 analysis tissues were homogenized in RIPA buffer (50 mM Tris pH 8, 150 mM NaCl,

5mM EDTA, 0.5 % SDS, 0.5 % SDO, 1 % NP-40) with protease inhibitor cocktail (Roche) and

incubated on ice for 1 hour, spun down at 13,300 rpm for 30 minutes at 4 °C and the supernatant was

collected. Protein measurements were performed using a BCA kit (Pierce) and 150 µg was loaded on a

10-20 % tris/tricine SDS-page gels (Bio-Rad Laboratories) and blotted onto 0,2 µm nitrocellulose

membrane (Bio-Rad Laboratories). Membranes were incubated overnight with SERF2 (1/1500,

Protein tech) or Actin (1/10 000, MP biomedicals) antibody. Next, the membranes were incubated

with anti-mouse or –rabbit secondary antibodies tagged with horseradish peroxidase (1/10 000, Bio-

Rad Laboratories) for one hour at room temperature and visualized by chemiluminescence (Amersham

ECL prime western blotting detection reagent, VWR).

RNA sequencing

RNA sequencing analysis was performed on three MEF cell lines from *Serf2*$^{-/-}$ mice and four *Serf2*$^{+/+}$

littermate controls. Total RNA was isolated from MEFs using the Qiagen RNeasy isolation kit.

Integrity of the RNA based on RIN scores as determined by a Bioanalyzer (Agilent). RNA-sequencing

libraries were prepared using TruSeq Stranded Total RNA with Ribo-Zero Human/Mouse/Rat (RS-

122-2201; Illumina, CA, USA) according to manufacturers' protocol. Pooled libraries were sequenced

on an Illumina HiSeq 2500 (single-end 50 bp). Reads were aligned to the mouse reference genome (mm10) using a splicing-aware aligner (StarAligner). Aligned reads were normalized to fragments per million (FPM), excluding low abundance genes (mean FPM>1 in at least two samples). The raw count data were pre-processed using the programming language R (3.4.0) [R Core Team, 2016, available online at: www.r-project.org], the program RStudio (1.0.143) [RStudio Team, 2016, available online at: http://www.rstudio.com/] and the EdgeR package (3.18.0)(Robinson et al., 2010). Genes that displayed FPM value >1 in at least two libraries were retained, resulting in a list of 12,808 genes for differential analysis. Differentially expressed (DE) genes between the $Serf2^{+/+}$ and $Serf2^{-/-}$ MEFs were identified using the EdgeR general linear model approach. After statistical correction for multiple comparisons with the "false discovery rate" (FDR) method (FDR<0.05), a list of 738 DE genes was obtained. DAVID (6.8) was used to perform functional annotation analyses on this gene list and to identify significantly enriched gene ontology (GO) terms [using GOTERM_BP_DIRECT] (Huang *et al*, 2009b, 2009a). The connectivity between the enriched GO terms was further examined by determining the amount of associated genes found (AGF) that were shared between two GO terms. The most significant relationships (where a GO term shared > 25 % of the AGF) were mapped in a network using the igraph package (1.0.1). The color of the edges reflects the major node, with the largest amount of AGF, in each relationship. The weight of the edges was determined by the percentage of AGF shared by the minor node in the relationship. We defined three clusters of GO-terms, where multiple minor GO-terms connected to one or two central, major GO-terms . Minor GO-terms were always clustered with the major GO-term with whom they shared the strongest connection. One GO term, positive regulation of the apoptotic process, showed no direct relation with one of the major GO-terms and was therefore assigned to the only cluster it had a significant connection with (Figure S2A-B). RNA-seq data have been deposited in the ArrayExpress database at EMBL-EBI (www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-10083. (http://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-10083)

[to be deleted] Upon acceptance of this work the data will be open access available, without password. Pending acceptance, the website can only be accessed by collaborators and reviewers

ELISA

The ELISA experiments were performed as described previously(Huang *et al*, 2015). Briefly, the

frozen brain samples were homogenized in tissue protein extraction reagent (Pierce) supplemented

with complete protease inhibitors (Roche), and centrifuged for 1 hour at 430 000 g at 4 °C. The

supernatant was used for ELISA. The $A\beta_{40}$ and $A\beta_{42}$ levels were determined by standard sandwich

ELISAs using end-specific antibodies (Janssen Pharmaceutical), and the monoclonal JRFcAβ_{40}/28 and

JRFcAβ_{42}/26 antibodies as the capture antibodies. Horseradish peroxidase conjugated JRFAβN/25 or

JRFAβN/25 were used as the detection antibodies for, respectively, human Aβ or murine Aβ. Samples

were loaded in triplicate for each ELISA. The ELISAs were performed in triplicate for the 3-month-

old animals and in duplicate for the 1-month-old animals.

Brain immunohistochemistry

To evaluate Aβ-related pathology, the brain was cut in 30 μm thick serial cryostat sections. A series of

sections was selected with 300 μm spacing between individual sections and blind-coded, 6 per mouse

for the 6E10 and Iba1 antibody and 3 per mouse for W02. Sections were fixed in 4 % PFA for 10

minutes, followed by 20 minutes antigen retrieval in citrate acid (pH 6.0) and 30 minutes in 1 % H_2O_2

in methanol. Next the sections were incubated for 1 hour in 10 % goat or donkey serum, depending on

the secondary antibody, and immunolabeled overnight with antibodies against Aβ, and Iba1

(Supplemental Table S6). The sections were washed and incubated with the complementary

biotinylated secondary antibody (1/500, Jackson Immunoresearch) for 2 hours. Next, sections were

incubated in ABC complex and reacted with diamino-benzidine (0.5 mg/ml H2O with 0.01 % H_2O_2).

The stained sections were scanned with the TissueFAXs microscope using 20X objective. All images

were analyzed using the Fiji platform of the ImageJ software(Schindelin *et al*, 2012). For unbiased

feature detection we used a single automated script with optimized threshold method and parameters

for the complete image set of each staining method.

Thioflavin-S Staining

A 300 μm-spaced series of three blind-coded 30 μm sections were fixed in 4 % PFA for 10 min,

followed by incubation in 0.25 % potassium permanganate solution for 15 min, and a bleaching step

with 1 % potassium metabisulfite/1 % oxalic acid for 5 min, incubation with 0.02 % Thioflavin-S

solution in 50 % ethanol (Sigma T1892) for 8 minutes, rinsed with water between every step. Finally

the sections were incubated with 1 μl/ml DAPI. The sections were analyzed as described for the

immunostained sections. Stained sections were scanned with the TissueFAXs microscope using 20X

objective. Images were processed using Tissuequest software, selecting the cortex for automated

analyses.

Luminescent conjugated oligothiophene (LCO) staining

For the Luminescent conjugated oligothiophenes (LCO) stainings, two different LCO variants,

qFTAA (quadro-formyl thiophene acetic acid) and hFTAA (hepta-formyl thiophene acetic acid) were

used. Blind-coded 30 μm sections were double-stained with qFTAA and hFTAA (2.4 μM qFTAA and

0.77 μM hFTAA in PBS) similar to a previous description (Nyström et al., 2013). Sections were

incubated for 30 minutes in the dark at room temperature. Stainings were analyzed on the Zeiss LSM

510 META confocal microscope equipped with an argon 458 nm laser (Carl Zeiss MicroImaging

GmbH, Jena, Germany). An 40x objective (oil-immersion, 1.3 NA, Zeiss) was used for spectral

imaging of the Aβ deposits. Stacked images were acquired every 10 nm in the emission spectrum

between 470 to 695 nm. Locations were selected randomly from the plaque containing regions of the

temporal, frontal and occipital cortex. Images were analyzed using Fiji (Schindelin et al., 2012, 2015)

and standard computer algorithms to provide a reproducible, unbiased detection of the plaques. First,

the stacks of images across the different wavelengths of the spectrum were combined into one image

using the Z-projection sum. A threshold was applied to the resulting image using the "Triangle dark"

algorithm to automatically identify and delineate the plaques. Partial plaques on the edge of the images

and particles smaller than 25 μm^2 were excluded. These settings allowed us to detect over 98 % of all

the plaques in the images. The missed plaques were either too small or too low in intensity to be

accurately distinguished from the background. Incorrectly identified particles due to (lipofuscin) auto

fluorescence were readily identified based on their wavelength intensity profile and visual inspection

and were also excluded from the analysis. We identified between 17-25 plaques in each animal for the

spectral intensity measurements. The ratio of the intensity of emitted light at the blue-shifted portion

(502 nm) and red-shifted peak (588 nm) was used as a parameter for spectral distinction of different

Aβ deposits. These peaks of the spectra were selected to maximize the spectral distinction.

Large-scale STEM (nanotomy)

The scanning transmission electron microscopy (STEM) experiments were adapted from previously

described experiments (Kuipers et al., 2015). Briefly, paraffin embedded cortical sections of AM and

AM;*Serf2$^{br-/-}$* were deparaffinized and postfixed with 1 % osmium tetroxide/1.5 % potassium

ferrocyanide in 0.1 M sodium cacodylate, dehydrated through ethanol and embedded in EPON (Serva)

using a tissue processor (Leica EM TP 709202). Ultrathin sections (80 nm) were cut using the Leica

uc7 ultramicrotome and collected on formvar coated cupper grids (electron microscopy sciences,

Hatfield, Pennsylvania). A large area scan using Scanning Transmission Detection was made using a

Zeiss supra55 SEM with Atlas. STEM detection with a four quadrant STEM detector was used in

inverted darkfield mode, at 28 kV with 30 μm aperture at 3.5 mm working distance. All images were

recorded at the same scan speed (cycle time 1.5 minute at 3072×2304 pixels). Contrast and brightness

were set based on a live histogram. High resolution large-scale STEM images at 2.5 nm pixel size

were generated with the external scan generator ATLAS (Fibics, Canada), individual tiles were

stitched in VE viewer (Fibics, Canada), exported as a html-file and uploaded to the website

www.nanotomy.org. Nanotomy datasets are open access available via the repository website

www.nanotomy.org/PW/temp02/EllenNollen.

[to be deleted] Upon acceptance of this work the website will be open access available, without

password. Pending acceptance, the website can only be accessed by collaborators and reviewers

Cell culture

Mouse embryonic fibroblasts (MEFs) were isolated from *Serf2$^{+/+}$, Serf2$^{+/-}$* and *Serf2$^{-/-}$* E13.5 embryos.

MEFs were cultured in T75 culture flasks (Greiner Bio-One, 658175), high–glucose Dulbecco's

modified Eagle's medium (Gibco), supplemented with 10% fetal bovine serum (Sigma 12133C), 1 %

penicillin/streptomycin (Gibco), non-essential amino acids (Gibco) and β-mercaptoethanol at 37°C, 5

% CO_2 and 3 % O_2. E13.5 embryos. Wild-type HEK293T and two independent clonal lines of

CRISPR *Serf2* KO mutant HEK 293T cells were cultured in Dulbecco's Modified Eagle Medium

(DMEM, Gibco; high glucose, pyruvate, 41966052), supplemented with 10 % Bovine Cow Serum

(BCS; Sigma 12133C) and 1 % Penicillin/Streptomycin (Gibco; 10 000 U/mL, 15140122), at 37 °C, 5

% CO_2. For passaging of cells, 0.05 % Trypsin-EDTA (1X), Phenol Red (Invitrogen; 25300-054) was

used. Regular mycoplasma tests were performed.

Cell proliferation assay

For the proliferation assay, 200 000 cells of each cell line were plated in triplicate in a standard 12-

well plate and incubated inside the IncuCyte Zoom Live-Cell Imaging System (Essen Bioscience) for

over 100 hours. The Incucyte system captured phase contrast images with a 10X magnification every 2

hours at nine distinct locations in each well to determine average confluency as a measure of cell

growth. The quantification was performed using the IncuCyte Zoom software. The growth experiment

was repeated three times. The confluency data of each experiment was normalized by min-max scaling

and the average of the three experiments was plotted (solid line) with the standard deviation (dashed

lines).

Statistical analysis

Unless specified otherwise, statistical analysis was performed using Graphpad v7.02. The exact

statistical test used for each comparison is specified in the main body of the text, together with the

resulting p-values. In the case of multiple comparisons, a post-hoc analysis with Bonferroni correction

was performed. Any corrected p-values derived from this post-hoc analysis are indicated in the text as

p_{bon}.

Acknowledgments

All LCOs were kindly provided by K. Peter R. Nilsson, Linköping University, Sweden. Part of the

work has been performed in the UMCG Microscopy and Imaging Center (UMIC), sponsored by

ZonMW grant 91111.006 and NWO 40-00506-98-9021. We thank Ben Giepmans for advise on EM

studies. This project was funded by a Meervoud Grant from NWO (836.09.001) (to E.A.A.N.), a

European Research Council (ERC) starting grant (281622 PDControl) (to E.A.A.N.), the Alumni

chapter Gooische Groningers facilitated by the Ubbo Emmius Fonds (to E.A.A.N), an Aspasia

fellowship from NWO (015.014.005) (to E.A.A.N.).

Conflict of Interests

The authors declare that they have no conflict of interest.

References

Borchelt DR, Thinakaran G, Eckman CB, Lee MK, Davenport F, Ratovitsky T, Prada CM, Kim G, Seekins S, Yager D, *et al* (1996) Familial Alzheimer's disease-linked presenilin I variants elevate aβ1- 42/1-40 ratio in vitro and in vivo. *Neuron*

Bousset L, Pieri L, Ruiz-Arlandis G, Gath J, Jensen PH, Habenstein B, Madiona K, Olieric V, Böckmann A, Meier BH, *et al* (2013) Structural and functional characterization of two alpha-synuclein strains. *Nat Commun* 4: 2575

Bugiani O, Giaccone G, Rossi G, Mangieri M, Capobianco R, Morbin M, Mazzoleni G, Cupidi C, Marcon G, Giovagnoli A, *et al* (2010) Hereditary Cerebral Hemorrhage With Amyloidosis Associated With the E693K Mutation of APP. *Arch Neurol* 67: 987–995

Chartier-Harlin M-C, Crawford F, Houlden H, Warren A, Hughes D, Fidani L, Goate A, Rossor M, Roques P, Hardy J, *et al* (1991) Early-onset Alzheimer's disease caused by mutations at codon 717 of the β-amyloid precursor protein gene. *Nature* 353: 844–846

Choi W, Zibaee S, Jakes R, Serpell LC, Davletov B, Anthony Crowther R & Goedert M (2004) Mutation E46K increases phospholipid binding and assembly into filaments of human α-synuclein. *FEBS Lett* 576: 363–368

Ciemerych MA & Sicinski P (2005) Cell cycle in mouse development. *Oncogene* doi:10.1038/sj.onc.1208608 [PREPRINT]

Citron M, Oltersdorf T, Haass C, McConlogue L, Hung AY, Seubert P, Vigo-Pelfrey C, Lieberburg I & Selkoe DJ (1992) Mutation of the β-amyloid precursor protein in familial Alzheimer's disease increases β-protein production. *Nature*

Citron M, Westaway D, Xia W, Carlson G, Diehl T, Levesque G, Johnson-Wood K, Lee M, Seubert P, Davis A, *et al* (1997) Mutant presenilins of Alzheimer's disease increase production of 42-

residue amyloid beta-protein in both transfected cells and transgenic mice. *Nat Med* 3: 67–72

Cohen E, Paulsson JF, Blinder P, Burstyn-Cohen T, Du D, Estepa G, Adame A, Pham HM,

Holzenberger M, Kelly JW, *et al* (2009) Reduced IGF-1 Signaling Delays Age-Associated

Proteotoxicity in Mice. *Cell* 139: 1157–1169

Cohen ML, Kim C, Haldiman T, ElHag M, Mehndiratta P, Pichet T, Lissemore F, Shea M, Cohen Y,

Chen W, *et al* (2015) Rapidly progressive Alzheimer's disease features distinct structures of

amyloid-β. *Brain* 138: 1009–1022

Conway KA, Harper JD & Lansbury PT (1998) Accelerated in vitro fibril formation by a mutant α-

synuclein linked to early-onset Parkinson disease. *Nat Med*

Conway KA, Lee SJ, Rochet JC, Ding TT, Williamson RE & Lansbury PT (2000) Acceleration of

oligomerization, not fibrillization, is a shared property of both α-synuclein mutations linked to

early-onset Parkinson's disease: Implications for pathogenesis and therapy. *Proc Natl Acad Sci U*

S A

Cooper JK, Schilling G, Peters MF, Herring WJ, Sharp AH, Kaminsky Z, Masone J, Khan FA,

Delanoy M, Borchelt DR, *et al* (1998) Truncated N-terminal fragments of huntingtin with

expanded glutamine repeats form nuclear and cytoplasmic aggregates in cell culture. *Hum Mol*

Genet

Cruts M, Dermaut B, Rademakers R, Van Den Broeck M, Stögbauer F & Van Broeckhoven C (2003)

Novel APP mutation V715A associated with presenile Alzheimer's disease in a German family.

J Neurol 250: 1374–1375

Duff K, Eckman C, Zehr C, Yu X, Prada CM, Perez-Tur J, Hutton M, Buee L, Harigaya Y, Yager D,

et al (1996) Increased amyloid-β42(43) in brains of mice expressing mutant presenilin 1. *Nature*

Eckman CB, Mehta ND, Crook R, Perez-tur J, Prihar G, Pfeiffer E, Graff-Radford N, Hinder P, Yager

D, Zenk B, *et al* (1997) A new pathogenic mutation in the APP gene (I716V) increases the
relative proportion of A beta 42(43). *Hum Mol Genet* 6: 2087–9

Elkins MR, Wang T, Nick M, Jo H, Lemmin T, Prusiner SB, Degrado WF, Stö Hr J & Hong M (2016)
Structural Polymorphism of Alzheimer's β-Amyloid Fibrils as Controlled by an E22 Switch: A
Solid-State NMR Study.

Falsone SF, Meyer NH, Schrank E, Leitinger G, Pham CLL, Fodero-Tavoletti MT, Holmberg M,
Dulle M, Scicluna B, Gesslbauer B, *et al* (2012) SERF Protein Is a Direct Modifier of Amyloid
Fiber Assembly. *Cell Rep* 2: 358–371

Gath J, Bousset L, Habenstein B, Melki R, Böckmann A & Meier BH (2014) Unlike Twins: An NMR
Comparison of Two α-Synuclein Polymorphs Featuring Different Toxicity. *PLoS One* 9: e90659

Ghosh D, Mondal M, Mohite GM, Singh PK, Ranjan P, Anoop A, Ghosh S, Jha NN, Kumar A & Maji
SK (2013) The Parkinson's Disease-Associated H50Q Mutation Accelerates α-Synuclein
Aggregation *in Vitro*. *Biochemistry* 52: 6925–6927

Ghosh D, Sahay S, Ranjan P, Salot S, Mohite GM, Singh PK, Dwivedi S, Carvalho E, Banerjee R,
Kumar A, *et al* (2014) The Newly Discovered Parkinson's Disease Associated Finnish Mutation
(A53E) Attenuates α-Synuclein Aggregation and Membrane Binding. *Biochemistry* 53: 6419–
6421

Glenner GG & Wong CW (1984) Alzheimer's disease: Initial report of the purification and
characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun*

Goate A, Chartier-Harlin MC, Mullan M, Brown J, Crawford F, Fidani L, Giuffra L, Haynes A, Irving
N, James L, *et al* (1991) Segregation of a missense mutation in the amyloid precursor protein
gene with familial Alzheimer's disease. *Nature* 349: 704–706

Grabowski TJ, Cho HS, Vonsattel JPG, Rebeck GW & Greenberg SM (2001) Novel amyloid

720 precursor protein mutation in an Iowa family with dementia and severe cerebral amyloid
721 angiopathy. *Ann Neurol* 49: 697–705

722 Grant MA, Lazo ND, Lomakin A, Condron MM, Arai H, Yamin G, Rigby AC & Teplow DB (2007)
723 Familial Alzheimer's disease mutations alter the stability of the amyloid-protein monomer
724 folding nucleus

725 Grundke-Iqbal I, Iqbal K, Quinlan M, Tung YC, Zaidi MS & Wisniewski HM (1986a) Microtubule-
726 associated protein tau. A component of Alzheimer paired helical filaments. *J Biol Chem* 261:
727 6084–6089

728 Grundke-Iqbal I, Iqbal K, Tung Y-C, Quinlan M, Wisniewski HM & Bindert LI (1986b) Abnormal
729 phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal
730 pathology. *Proc Natl Acad Sci U S A* 83: 4913–4917

731 Guerreiro RJ, Baquero M, Blesa R, Boada M, Brás JM, Bullido MJ, Calado A, Crook R, Ferreira C,
732 Frank A, *et al* (2010) Genetic screening of Alzheimer's disease genes in Iberian and African
733 samples yields novel mutations in presenilins and APP. *Neurobiol Aging* 31: 725–731

734 Guerrero-Ferreira R, Taylor NMI, Arteni A-A, Kumari P, Mona D, Ringler P, Britschgi M, Lauer ME,
735 Verasdock J, Riek R, *et al* (2019) Two new polymorphic structures of alpha-synuclein solved by
736 cryo-electron microscopy. *bioRxiv*: 654582

737 Haass C, Lemere CA, Capell A, Citron M, Seubert P, Schenk D, Lannfelt L & Selkoe DJ (1995) The
738 Swedish mutation causes early-onset Alzheimer's disease by beta-secretase cleavage within the
739 secretory pathway. *Nat Med* 1: 1291–6

740 van Ham TJ, Holmberg MA, van der Goot AT, Teuling E, Garcia-Arencibia M, Kim H eui, Du D,
741 Thijssen KL, Wiersma M, Burggraaff R, *et al* (2010) Identification of MOAG-4/SERF as a
742 regulator of age-related proteotoxicity. *Cell* 142: 601–612

Heise H, Celej MS, Becker S, Riedel D, Pelah A, Kumar A, Jovin TM & Baldus M (2008) Solid-State

NMR Reveals Structural Differences between Fibrils of Wild-Type and Disease-Related A53T

Mutant α-Synuclein. *J Mol Biol* 380: 444–450

Huang DW, Sherman BT & Lempicki RA (2009a) Bioinformatics enrichment tools: paths toward the

comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 37: 1–13

Huang DW, Sherman BT & Lempicki RA (2009b) Systematic and integrative analysis of large gene

lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44–57

Huang Y, Skwarek-Maruszewska A, Horré K, Vandewyer E, Wolfs L, Snellinx A, Saito T, Radaelli E,

Corthout N, Colombelli J, *et al* (2015) Loss of GPR3 reduces the amyloid plaque burden and

improves memory in Alzheimer's disease mouse models. *Sci Transl Med* 7: 309ra164

Inayathullah M & Teplow DB (2011) Structural dynamics of the Δe22 (Osaka) familial Alzheimer's

disease-linked amyloid β-protein. *Amyloid* 18: 98–107

Kaufman MH (1992) The Atlas of Mouse Development Elsevier Science

Kaufman MH & Bard JBL (1999) The Anatomical Basis of Mouse Development Elsevier Science

Klingstedt T, Aslund A, Simon RA, Johansson LBG, Mason JJ, Nyström S, Hammarström P &

Nilsson KPR (2011) Synthesis of a library of oligothiophenes and their utilization as fluorescent

ligands for spectral assignment of protein aggregates. *Org Biomol Chem* 9: 8356–70

Kulic L, McAfoose J, Welt T, Tackenberg C, Späni C, Wirth F, Finder V, Konietzko U, Giese M,

Eckert A, *et al* (2012) Early accumulation of intracellular fibrillar oligomers and late congophilic

amyloid angiopathy in mice expressing the Osaka intra-Aβ APP mutation. *Transl Psychiatry* 2:

e183–e183

Kumar-Singh S, De Jonghe C, Cruts M, Kleinert R, Wang R, Mercken M, De Strooper B,

Vanderstichele H, Löfgren A, Vanderhoeven I, *et al* (2000) Nonfibrillar diffuse amyloid
deposition due to a gamma(42)-secretase site mutation points to an essential role for N-truncated
A beta(42) in Alzheimer's disease. *Hum Mol Genet* 9: 2589–98

Kumar-Singh S, Theuns J, Van Broeck B, Pirici D, Vennekens K, Corsmit E, Cruts M, Dermaut B,
Wang R & Van Broeckhoven C (2006) Mean age-of-onset of familial Alzheimer disease caused
by presenilin mutations correlates with both increased Aβ42 and decreased Aβ40. *Hum Mutat*

Langley CH & Charlesworth B (1997) Nature © Macmillan Publishers Ltd 1997 expansion of
xenotransplantation

Lazo ND, Grant MA, Condron MC, Rigby AC & Teplow DB (2009) On the nucleation of amyloid β-
protein monomer folding. *Protein Sci* 14: 1581–1596

Lemere CA, Lopera F, Kosik KS, Lendon CL, Ossa J, Saido TC, Yamaguchi H, Ruiz A, Martinez A,
Madrigal L, *et al* (1996) The E280A presenilin 1 Alzheimer mutation produces increased A beta
42 deposition and severe cerebellar pathology. *Nat Med* 2: 1146–50

Leung AKL, Gerlich D, Miller G, Lyon C, Lam YW, Lleres D, Daigle N, Zomerdijk J, Ellenberg J &
Lamond AI (2004) Quantitative kinetic analysis of nucleolar breakdown and reassembly during
mitosis in live human cells. *J Cell Biol* 166: 787–800

Li B, Ge P, Murray KA, Sheth P, Zhang M, Nair G, Sawaya MR, Shin WS, Boyer DR, Ye S, *et al*
(2018) Cryo-EM of full-length α-synuclein reveals fibril polymorphs with a common structural
kernel. *Nat Commun* 9: 3609

Lord A, Kalimo H, Eckman C, Zhang X-Q, Lannfelt L & Nilsson LNG (2006) The Arctic Alzheimer
mutation facilitates early intraneuronal Aβ aggregation and senile plaque formation in transgenic
mice. *Neurobiol Aging* 27: 67–77

Lord A, Philipson O, Klingstedt T, Westermark G, Hammarström P, Nilsson KPR & Nilsson LNG

(2011) Observations in APP bitransgenic mice suggest that diffuse and compact plaques form via

independent processes in Alzheimer's disease. *Am J Pathol* 178: 2286–2298

Lu JX, Qiang W, Yau WM, Schwieters CD, Meredith SC & Tycko R (2013) Molecular structure of β-

amyloid fibrils in alzheimer's disease brain tissue. *Cell*

Masters CL, Simms G, Weinman NA, Multhaupt G, Mcdonald BL & Beyreuthert K (1985) Amyloid

plaque core protein in Alzheimer disease and Down syndrome. *Proc Natl Acad Sci U S A* 82:

4245–4249

Meinen BA, Gadkari V V., Stull F, Ruotolo BT & Bardwell JCA (2019) SERF engages in a fuzzy

complex that accelerates primary nucleation of amyloid proteins. *Proc Natl Acad Sci*

Merle DA, Witternigg A, Tam-Amersdorfer C, Hartlmüller C, Spreitzer E, Schrank E, Wagner-

Lichtenegger S, Werzer O, Zangger K, Kungl AJ, *et al* (2019) Increased Aggregation Tendency

of Alpha-Synuclein in a Fully Disordered Protein Complex. *J Mol Biol* 431: 2581–2598

Meyer NH, Dellago H, Tam-Amersdorfer C, Merle DA, Parlato R, Gesslbauer B, Almer J,

Gschwandtner M, Leon A, Franzmann TM, *et al* (2019) Structural fuzziness of the RNA-

organizing protein SERF1a determines a toxic gain-of-interaction. *bioRxiv*: 713511

Miravalle L, Tokuda T, Chiarle R, Giaccone G, Bugiani O, Tagliavini F, Frangione B & Ghiso J

(2000) Substitutions at Codon 22 of Alzheimer's A Peptide Induce Diverse Conformational

Changes and Apoptotic Effects in Human Cerebral Endothelial Cells*.

Müller U, Winter P, Bolender C & Nolte D (2014) Previously Unrecognized Missense Mutation

E126K of PSEN2 Segregates with Early Onset Alzheimer's Disease in a Family. *J Alzheimer's*

Dis 42: 109–113

Murayama O, Tomita T, Nihonmatsu N, Murayama M, Sun X, Honda T, Iwatsubo T & Takashima A

(1999) Enhancement of amyloid β 42 secretion by 28 different presenilin 1 mutations of familial

811  Alzheimer's disease. *Neurosci Lett* 265: 61–63

812 Murray SA, Morgan JL, Kane C, Sharma Y, Heffner CS, Lake J & Donahue LR (2010) Mouse
813  Gestation Length Is Genetically Determined. *PLoS One* 5: e12418

814 Murrell JR, Hake AM, Quaid KA, Farlow MR & Ghetti B (2000) Early-Onset Alzheimer Disease
815  Caused by a New Mutation (V717L) in the Amyloid Precursor Protein Gene. *Arch Neurol* 57:
816  885

817 Narhi L, Wood SJ, Steavenson S, Jiang Y, Wu GM, Anafi D, Kaufman SA, Martin F, Sitney K, Denis
818  P, *et al* (1999) Both Familial Parkinson's Disease Mutations Accelerate α-Synuclein
819  Aggregation. *J Biol Chem* 274: 9843–9846

820 Nekooki-Machida Y, Kurosawa M, Nukina N, Ito K, Oda T & Tanaka M (2009) Distinct
821  conformations of in vitro and in vivo amyloids of huntingtin-exon1 show different cytotoxicity.
822  *Proc Natl Acad Sci U S A* 106: 9679–84

823 Ni C-L, Shi H-P, Yu H-M, Chang Y-C & Chen Y-R (2011) Folding stability of amyloid-β 40
824  monomer is an important determinant of the nucleation kinetics in fibrillization. *FASEB J* 25:
825  1390–1401

826 Nilsberth C, Westlind-Danielsson A, Eckman CB, Condron MM, Axelman K, Forsell C, Stenh C,
827  Luthman J, Teplow DB, Younkin SG, *et al* (2001) The 'Arctic' APP mutation (E693G) causes
828  Alzheimer's disease by enhanced Aβ protofibril formation. *Nat Neurosci* 4: 887–893

829 Van Nostrand WE, Melchor JP, Cho HS, Greenberg SM & Rebeck GW (2001) Pathogenic effects of
830  D23N Iowa mutant amyloid beta -protein. *J Biol Chem* 276: 32860–6

831 Nyström S, Psonka-Antonczyk KM, Ellingsen PG, Johansson LBG, Reitan N, Handrick S, Prokop S,
832  Heppner FL, Wegenast-Braun BM, Jucker M, *et al* (2013) Evidence for Age-Dependent in Vivo
833  Conformational Rearrangement within Aβ Amyloid Deposits. *ACS Chem Biol* 8: 1128–1133

Ovchinnikova OY, Finder VH, Vodopivec I, Nitsch RM & Glockshuber R (2011) The Osaka FAD

Mutation E22Δ Leads to the Formation of a Previously Unknown Type of Amyloid β Fibrils and

Modulates Aβ Neurotoxicity. *J Mol Biol* 408: 780–791

Petkova AT, Leapman RD, Guo Z, Yau WM, Mattson MP & Tycko R (2005) Self-propagating,

molecular-level polymorphism in Alzheimer's β-amyloid fibrils. *Science (80-)*

Pevny LH, Sockanathan S, Placzek M & Lovell-Badge R (1998) A role for SOX1 in neural

determination. *Development*

Philipson O, Hammarström P, Nilsson KPR, Portelius E, Olofsson T, Ingelsson M, Hyman BT,

Blennow K, Lannfelt L, Kalimo H, *et al* (2009) A highly insoluble state of Aβ similar to that of

Alzheimer's disease brain is found in Arctic APP transgenic mice. *Neurobiol Aging* 30: 1393–

1405

Philipson O, Lord A, Lalowski M, Soliymani R, Baumann M, Thyberg J, Bogdanovic N, Olofsson T,

Tjernberg LO, Ingelsson M, *et al* (2012) The Arctic amyloid-β precursor protein (AβPP)

mutation results in distinct plaques and accumulation of N- and C-truncated Aβ. *Neurobiol Aging*

33: 1010.e1-1010.e13

Pras A, Houben B, Aprile FA, Seinstra R, Gallardo R, Janssen L, Hogewerf W, De Vleeschouwer M,

Mata-Cabana A, Koopman M, *et al* (2020) The cellular modifier MOAG-4/SERF drives amyloid

formation through charge complementation. *bioRxiv*: 2020.12.09.417709

Psonka-Antonczyk KM, Hammarström P, Johansson LBG, Lindgren M, Stokke BT, Nilsson KPR &

Nyström S (2016) Nanoscale Structure and Spectroscopic Probing of Aβ1-40 Fibril Bundle

Formation. *Front Chem*

Qiang W, Yau W-M, Lu J-X, Collinge J & Tycko R (2017) Structural variation in amyloid-β fibrils

from Alzheimer's disease clinical subtypes. *Nature* 541: 217–221

Radde R, Bolmont T, Kaeser S a, Coomaraswamy J, Lindau D, Stoltze L, Calhoun ME, Jäggi F,

Wolburg H, Gengler S, *et al* (2006) Abeta42-driven cerebral amyloidosis in transgenic mice

reveals early and robust pathology. *EMBO Rep* 7: 940–6

Rasmussen J, Mahler J, Beschorner N, Kaeser SA, Häsler LM, Baumann F, Nyström S, Portelius E,

Blennow K, Lashley T, *et al* (2017) Amyloid polymorphisms constitute distinct clouds of

conformational variants in different etiological subtypes of Alzheimer's disease. *Proc Natl Acad*

Sci 114: 13018–13023

Rogaev EI, Sherrington R, Rogaeva EA, Levesque G, Ikeda M, Liang Y, Chi H, Lin C, Holman K,

Tsuda T, *et al* (1995) Familial Alzheimer's disease in kindreds with missense mutations in a gene

on chromosome 1 related to the Alzheimer's disease type 3 gene. *Nature* 376: 775–778

Schellenberg G, Bird T, Wijsman E, Orr H, Anderson L, Nemens E, White J, Bonnycastle L, Weber J,

Alonso M, *et al* (1992) Genetic linkage evidence for a familial Alzheimer's disease locus on

chromosome 14. *Science (80-)* 258: 668–671

Scherzinger E, Sittler A, Schweiger K, Heiser V, Lurz R, Hasenbank R, Bates GP, Lehrach H &

Wanker EE (1999) Self-assembly of polyglutamine-containing huntingtin fragments into

amyloid-like fibrils: Implications for Huntington's disease pathology (huntingtinglutamine

repeataggregation)

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

Saalfeld S, Schmid B, *et al* (2012) Fiji: an open-source platform for biological-image analysis.

Nat Methods 9: 676–682

Sherrington R, Rogaev EI, Liang Y, Rogaeva EA, Levesque G, Ikeda M, Chi H, Lin C, Li G, Holman

K, *et al* (1995) Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's

disease. *Nature* 375: 754–760

Shimada H, Ataka S, Tomiyama T, Takechi H, Mori H & Miki T (2011) Clinical course of patients

with familial early-onset Alzheimer's disease potentially lacking senile plaques bearing the

E693Δ mutation in amyloid precursor protein. *Dement Geriatr Cogn Disord* 32: 45–54

Skarnes WC, Rosen B, West AP, Koutsourakis M, Bushell W, Iyer V, Mujica AO, Thomas M,

Harrow J, Cox T, *et al* (2011) A conditional knockout resource for the genome-wide study of

mouse gene function. *Nature* 474: 337–342

Sun C-S, Lee C-C, Li Y-N, Yao-Chen Yang S, Lin C-H, Chang Y-C, Liu P-F, He R-Y, Wang C-H,

Chen W, *et al* (2015) Conformational switch of polyglutamine-expanded huntingtin into benign

aggregates leads to neuroprotective effect. *Sci Rep* 5: 14992

Suzuki N, Cheung TT, Cai XD, Odaka A, Otvos L, Eckman C, Golde TE & Younkin SG (1994) An

increased percentage of long amyloid β protein secreted by familial amyloid β protein precursor

(βAPP717) mutants. *Science (80-)*

Takashima Y, Era T, Nakao K, Kondo S, Kasuga M, Smith AG & Nishikawa S-I (2007)

Neuroepithelial Cells Supply an Initial Transient Wave of MSC Differentiation. *Cell* 129: 1377–

1388

Tomiyama T, Matsuyama S, Iso H, Umeda T, Takuma H, Ohnishi K, Ishibashi K, Teraoka R, Sakama

N, Yamashita T, *et al* (2010) A Mouse Model of Amyloid β Oligomers: Their Contribution to

Synaptic Alteration, Abnormal Tau Phosphorylation, Glial Activation, and Neuronal Loss In

Vivo. *J Neurosci* 30: 4845–4856

Tomiyama T, Nagata T, Shimada H, Teraoka R, Fukushima A, Kanemitsu H, Takuma H, Kuwano R,

Imagawa M, Ataka S, *et al* (2008) A new amyloid β variant favoring oligomerization in

Alzheimer's-type dementia. *Ann Neurol* 63: 377–387

Visintin R & Amon A (2000) The nucleolus: the magician's hat for cell cycle tricks. *Curr Opin Cell*

Biol 12: 372–377

Ward JM, Elmore SA & Foley JF (2012) Pathology Methods for the Evaluation of Embryonic and
Perinatal Developmental Defects and Lethality in Genetically Engineered Mice. *Vet Pathol* 49:
71–84

Wood HB & Episkopou V (1999) Comparative expression of the mouse Sox1, Sox2 and Sox3 genes
from pre-gastrulation to early somite stages. *Mech Dev*

Yoshimura Y, Holmberg MA, Kukic P, Andersen CB, Mata-Cabana A, Falsone SF, Vendruscolo M,
Nollen EAA & Mulder FAA (2017) MOAG-4 promotes the aggregation of α-synuclein by
competing with self-protective electrostatic interactions. *J Biol Chem* 292: 8269–8278

Figure legends

Figure 1. Full-body *Serf2*$^{-/-}$ mice display a developmental delay and perinatal lethality. (A)

Schematic overview of the crossing scheme with full-body *Serf2*$^{+/-}$ mice and the resulting genotypic

composition of the offspring at embryonic and postnatal stage. **(B)** Absolute animal counts for the

three genotypes in the offspring at various days of embryonic development(E13.5, E15.5 and E17.5),

at birth (P0) and at time of weaning (P22). **(C)** Real time RT-PCR analyses of *Serf1* and *Serf2* RNA

levels in E13.5 heads normalized to housekeeping gene beta-actin (n=4/group, mean±SD, One-way

ANOVA for each gene, Bonferroni corrected post-hoc comparison between genotypes ***p<0.001).

(D) Western blot of SERF2 and actin in *Serf2*$^{+/+}$, *Serf2*$^{+/-}$, *Serf2*$^{-/-}$embryos at E13.5 (black arrow

indicates SERF2 band). **(E)** Length measurements of *Serf2*$^{+/+}$, *Serf2*$^{+/-}$, *Serf2*$^{-/-}$ embryos at E15.5, E17.5

and P0 (black line = mean, Two-way ANOVA with factors age and genotype, Bonferroni corrected

post-hoc comparison for *Serf2*$^{+/-}$ and *Serf2*$^{-/-}$ compared to *Serf2*$^{-/-}$ *p<0.05; ***p<0.001). **(F)**

Hematoxylin and eosin stained lung tissue from *Serf2*$^{+/+}$ and *Serf2*$^{-/-}$ pups at P0 (top: *Serf2*$^{-/-}$, bottom:

Serf2$^{+/+}$, scale bar=500μm, rectangle =5x magnification in right picture, **(G)** Quantification of the

saccolar space in lung tissue from *Serf2*$^{+/+}$ and *Serf2*$^{-/-}$ pups at P0 (black line = mean, t-test **p<0.01).

For all mouse data panels: *Serf2*$^{+/+}$= black, *Serf2*$^{+/-}$ = dark grey, *Serf2*$^{-/-}$ = light grey.

Figure 2. Crossing a conditional *Serf2* KO with an AM background does not affect the overall

levels of APP and Aβ. (A) Schematic overview of the cross between brain-specific *Serf2* KO mice

and the amyloid model resulting in four experimental groups: WT, *Serf2*$^{br-/-}$, AM and AM;*Serf2*$^{br-/-}$.

(B) Timeline for the Aβ pathology in the amyloid model and the selected time points for this study.

(C) Real time RT-PCR analyses of *Serf2* RNA levels in the brain tissue of all experimental groups at 1

and 3 months of age normalized to housekeeping gene beta-actin (n=6/group, mean±SD, Two-way

ANOVA with factors age and genotype, Bonferroni corrected post-hoc for individual groups

compared to WT). **(D)** Western blot of SERF2 and actin in brain tissue of all experimental groups at 1

and 3 months of age. **(E)** Brain weight of all experimental groups at 1 and 3 months of age (black line

= mean, Two-way ANOVA with factors age and genotype, Bonferroni corrected post-hoc for

individual groups compared to WT). **(F)** Brain weight of unfloxed control groups at 3 months of age

(black line = mean, One-way ANOVA, Bonferroni corrected post-hoc for individual groups compared

to WT). **(G)** Number of cortical nuclei per mm^2 in hematoxylin and eosin stained brain tissue of all

experimental groups at 3 months of age. (n=3/group, mean±SD, Kruskal-Wallis p=0.5566). **(H)** Real

time RT-PCR analyses of APP RNA levels in the brain tissue of AM and AM;*Serf2$^{br-/-}$* mice at 1 and 3

months of age normalized to housekeeping gene beta-actin (n=6/group, mean±SD, t-test between

genotypes at both ages not significant). **(I)** Western blot of APP and beta-actin in brain tissue of AM

and AM;*Serf2$^{br-/-}$* mice at 3 months of age versus 1-month-old non-AM controls. **(J)** Quantification of

total APP protein levels in brains of AM and AM;*Serf2$^{br-/-}$* mice at 1 and 3 months of age normalized

to beta-actin levels. (black line = mean, t-test between genotypes at both ages not significant). **(K)**

Aβ$_{40}$ and **(L)** Aβ$_{42}$ levels in brain lysate from AM and AM;*Serf2$^{br-/-}$* mice at 1 and 3 months of age as

determined by ELISA normalized to total protein content. (black line = mean, t-test between

genotypes at both ages not significant). **(M)** Correlation plot depicting the relationship between Aβ$_{40}$

and Aβ$_{42}$ levels in AM (black) and AM;*Serf2$^{br-/-}$* (grey) mice at 3 months of age (detailed statistics in

Supplemental Table 3). In panels A-L: WT = black, *Serf2$^{br-/-}$* = light grey , AM = dark grey and

AM;*Serf2$^{br-/-}$* = white; ***p<0.001; **p<0.01.

Figure 3. Brain-specific *Serf2* KO display increased Aβ deposition and amyloid .(A) Detailed light

microscope images of the 6E10-stained cortical region marked on the sagittal section in Supplemental

Figure 5 from 3-month-old AM and AM;*Serf2$^{br-/-}$* mice (scale bar = 100µm). **(B)** Quantification of the

Aβ deposits in the region of interest (ROI) of 6E10 immunostained sagittal brain sections from 1- and

3-month-old AM and AM;*Serf2$^{br-/-}$* mice (black line = mean, One-way ANOVA between genotypes at

both ages, Bonferroni corrected post-hoc between AM and AM;*Serf2$^{br-/-}$* not significant). **(C)**

Correlation plot depicting the relationship between Aβ$_{40}$ levels and 6E10 plaque density in AM (black)

and AM;*Serf2$^{br-/-}$* (grey) mice at 3 months of age. **(D)** Correlation plot depicting the relationship

between Aβ$_{42}$ levels and 6E10 plaque density in AM (black) and AM;*Serf2$^{br-/-}$* (grey) mice at 3 months

of age. **(E)** Quantification of the Aβ deposits in the ROI of W0-2 immunostained sagittal brain

sections from 1- and 3-month-old AM and AM;*Serf2$^{br-/-}$* mice (black line = mean, One-way ANOVA

between genotypes at both ages, Bonferroni corrected post-hoc between AM and AM;*Serf2*[br-/-] not

significant). **(F)** Correlation plot depicting the relationship between 6E10 deposits and W0-2 deposits

in AM (black) and AM;*Serf2*[br-/-] (grey) mice at 3 months of age. **(G)** Quantification of the plaque

density in the ROI of Thioflavin-S stained sagittal brain sections from 1- and 3-month-old AM and

AM;*Serf2*[br-/-] mice (black line = mean, One-way ANOVA between genotypes at both ages, Bonferroni

corrected post-hoc between AM and AM;*Serf2*[br-/-] not significant). **(H)** Correlation plot depicting the

relationship between $A\beta_{40}$ levels and ThS plaque density in AM (black) and AM;*Serf2*[br-/-] (grey) mice

at 3 months of age. **(I)** Correlation plot depicting the relationship between $A\beta_{42}$ levels and ThS plaque

density in AM (black) and AM;*Serf2*[br-/-] (grey) mice at 3 months of age. **(I)** Correlation plot depicting

the relationship between 6E10 deposits and ThS plaques in AM (black) and AM;*Serf2*[br-/-] (grey) mice

at 3 months of age. In all panels ROI = cortical and hippocampal area; In panels B,E and G: WT =

black, *Serf2*[br-/-] = light grey , AM = dark grey and AM;*Serf2*[br-/-] = white; for correlation plots: detailed

statistics in Supplemental Table 3; ***p<0.001; **p<0.01; *p <0.05.

Figure 4. Brain-specific *Serf2* KO alters the structural composition of amyloid plaques. (A)

Graph showing the theoretical LCO dye emission spectra for qFTAA, which binds mature $A\beta$ fibrils

(blue), and hFTAA, which binds both mature fibrils and prefibrillar $A\beta$ species (red). **(B)** Frequency

distribution of the ratio in fluorescence intensity at 502 nm and 588 nm for all amyloid plaques

identified by confocal microscopy in AM (blue) and AM;*Serf2*[br-/-] (red) mice at 3 months of age. Three

peak frequency categories were identified: low IR(red), medium IR (green) and high IR (blue) (n=6

mice/genotype). **(C)** Dot plot representing the intensity ratio of all amyloid plaques identified by

confocal microscopy for AM and AM;*Serf2*[br-/-] mice at 3 months of age (colors represent the IR

categories defined in panel B, black circles correspond to the plaques depicted in panel E, black line =

mean, t-test ***p<0.001) (n=6 mice/genotype). **(D)** Correlation plot depicting the relationship between

6E10 plaque density in the ROI and the LCO average intensity ratio in AM (black) and AM;*Serf2*[br-/-]

(grey) mice at 3 months of age (ROI = cortical and hippocampal area, detailed statistics in

Supplemental Table 3). **(E)** Spectral confocal microscopy images of amyloid plaques in AM and

AM;*Serf2*[br-/-] mice at 3 months of age of double stained with qFTAA and hFTAA (top= fluorescence

at 502 nm, middle= fluorescence at 588 nm, bottom=merged image of 502 nm and 588 nm

fluorescence, scale bar = 5 μm). **(F)** Scanning transmission electron microscopy pictures of amyloid

plaques in the cortex of AM and AM;*Serf2*$^{br-/-}$ mice at 3 months of age (top scale bar 5μm, rectangle =

zoomed region in bottom picture, bottom scale bar = 1μm). High-resolution EM images are available

via http://www.nanotomy.org.

Figure 1

Figure 2

Figure 3

Figure 4

Supplemental figure legends

Supplemental Figure 1. Additional full-body *Serf2$^{-/-}$* mice validation and phenotype confirmation. (A) Targeting strategy used to delete exon 2 of *Serf2* by using the 'knockout-first'-allele (adjusted from Skarnes et al. 2011). Top panel: exon/intron structure of *Serf2* including the positions of the LacZ and Neo cassettes, as well as loxP sites. Bottom panel: the *Serf2* allele after recombination events as a result of crosses with *Hprt-Cre* mice. **(B)** Western blot analysis of liver, heart and brain tissue of a *Serf2$^{+/+}$*, *Serf2$^{+/-}$* and *Serf2$^{-/-}$* adult mice to detect Serf2 and actin. **(C)** Real time RT-PCR analyses of *Serf2* in different organs of adult mice with the following genotypes *Serf2$^{+/+}$* (n=3), *Serf2$^{+/-}$* (n=8) and *Serf2$^{-/-}$* (n=1). *Serf2* levels were normalized to the housekeeping gene 18S (mean ± SEM). **(D)** Lateral view of a *Serf2$^{-/-}$* and *Serf2$^{+/-}$* embryo and placenta at embryonic day 15.5 and 17.5. **(E)** Weight measurements of *Serf2$^{+/+}$*, *Serf2$^{+/-}$*, *Serf2$^{-/-}$* embryos at E17.5 and P0 (black line = mean, Two-way ANOVA with factors age and genotype, Bonferroni corrected post-hoc comparison for *Serf2$^{+/-}$* and *Serf2$^{-/-}$* compared to *Serf2$^{-/-}$* *p<0.05; ***p<0.001). **(F)** Hematoxylin and Eosin staining on *Serf2$^{-/-}$* and *Serf2$^{+/+}$* lung and kidney tissue at E17.5 (10 times magnification). For all panels: *Serf2$^{+/+}$*= black, *Serf2$^{+/-}$* = dark grey, *Serf2$^{-/-}$* = light grey.

Supplemental Figure 2. Cell culture data of *Serf2* KO and control HEK cells and RNA sequencing analysis of mouse embryonic fibroblasts (MEFs) isolated from full-body *Serf2$^{-/-}$* and control mice. (A) Growth curve of two CRISPR-induced *Serf2$^{-/-}$* clones of HEK293T versus wild-type HEK293T. (Three replicates measured,full line=replicate average, dashed lines=standard deviation on average) **(B)** Network showing the interconnectivity of enriched GO-terms in RNA sequencing data from *Serf2$^{-/-}$* and control MEFs. Circle size indicates amount of differentially expressed genes found in each GO-category. Edges were drawn when the minor category shared more than 25 % of its found genes with the major category. Edge color indicates the major category in the relationship. Edge thickness indicates the percentage of found genes shared by the minor category. **(C)** Bar chart indicating the fold change for each GO-term as determined by DAVID. Brackets indicate the three main GO-term clusters defined based on the network in (B).

Supplemental Figure 3. Additional validation of the brain-specific knock-out of *Serf2*. (A)

Modified targeting strategy used to delete the exon 2 of *Serf2* specifically in the central nervous

system. **(B)** Real time RT-PCR analyses of *Serf2* RNA expression in different organs *Serf2*$^{+/+}$ and

Serf2$^{br-/-}$ female mice at 1 month and 3 months of age. *Serf2* expression was normalized to

housekeeping gene beta-actin (all groups n=6, mean ± SEM). **(C)** Western blot analysis and

quantification of *Serf2* and beta-actin in brain lysates of WT and *Serf2*$^{br-/-}$ female mice at 1 month

(both groups n=6, mean ± SEM, t-test **p<0.01). **(D)** Brain weight of unfloxed WT and CRE^{br} mice at

1 and 3 months of age (black line = mean, Two-way ANOVA with factors age and genotype,

Bonferroni corrected post-hoc for CRE^{br} compared to WT *p$_{bon}$<0.05). **(E)** Evolution of brain weight

in WT and *Serf2*$^{br-/-}$ female mice between 1 and 11 months of age (black line = mean).

Supplemental Figure 4. Overview of distribution and general morphological parameters of

6E10-stained Aβ deposits in AM and AM;*Serf2*$^{br-/-}$ mice at 3 months of age. (A) Light microscope

images of the 6E10 immunostained sagittal section of 3-month-old AM and AM;*Serf2*$^{br-/-}$ female mice

(scale bar = 1mm, black rectangle = magnified region in main Figure 3). **(B)** Boxplots showing the

distribution of the surface area of the individual plaques measured in AM and AM;*Serf2*$^{br-/-}$ mice at 3

months of age. **(C)** Boxplots showing the distribution of the circularity of the individual plaques

measured in AM and AM;*Serf2*$^{br-/-}$ mice at 3 months of age. **(D)** Boxplots showing the distribution of

the solidity of the individual plaques measured in AM and AM;*Serf2*$^{br-/-}$ mice at 3 months of age. For

all boxplots: black line = median, box = 25th-75th percentile, whiskers = 1.5 x interquartile range

(IQR), circles = outlier value > 1.5x IQR, asterisks = extreme outlier value > 3x IQR.

Supplemental Figure 5. Overview of distribution and detailed W0-2-stained Aβ deposits in AM

and AM;*Serf2*$^{br-/-}$ mice at 3 months of age. Light microscope images of the W0-2 immunostained

sagittal section of 3-month-old AM and AM;*Serf2*$^{br-/-}$ female mice (scale bar = 1mm, black rectangle =

magnified in insert on the right, insert scale bar = 100μm).

Supplemental Figure 6. Correlation plots illustrating the relationship of different stains in

individual mice. (A) Correlation plot depicting the relationship between the density of 6E10 deposits

and Iba1-positive spots in AM (black) and AM;*Serf2*$^{br-/-}$ (grey) mice at 3 months of age. **(B)**

Correlation plot depicting the relationship between W0-2 plaque density and the LCO average

intensity ratio in AM (black) and AM;*Serf2*$^{br-/-}$ (grey) mice at 3 months of age. **(C)** Correlation plot

depicting the relationship between ThS plaque density and the LCO average intensity ratio in AM

(black) and AM;*Serf2*$^{br-/-}$ (grey) mice at 3 months of age. For all plots: detailed statistics in

Supplemental Table 3; *p <0.05.

Supplemental Figure 7. Overview and detailed view of Aβ staining pattern after W0-2 and 6E10

staining in AM and AM;*Serf2*$^{br-/-}$ mice at 1 month of age. (A) Light microscope images of the W0-2

immunostained sagittal section of 1-month-old AM and AM;*Serf2*$^{br-/-}$ female mice. **(B)** Light

microscope images of the 6E10 immunostained sagittal section of 1-month-old AM and AM;*Serf2*$^{br-/-}$

female mice. All pictures: main scale bar = 1mm, black rectangle = magnified in insert on the right,

insert scale bar = 100μm.