Cellular feedback to organotelluranes displays modulation of antioxidant proteins gene expression

Felipe S. Pessoto[1,2], César H. Yokomizo[1], Rodrigo L. O. R. Cunha[1,]* and Iseli L. Nantes-Cardoso[1,]*

[1] Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, Santo André, SP, Brazil.

[2] INFAR, Universidade Federal de São Paulo, São Paulo, SP, Brazil.

KEYWORDS

Telluranes, redox modulation, thioredoxin, thiol reagents, gene expression, cellular feedback.

ABSTRACT

Organotelluranes, RT3 and RT4, are thiol reagents that induce mitochondrial transition pore (MTP) opening in the manner sensitive and insensitive to cyclosporin A. Although RT3 and RT4 promote glutathione depletion, paradoxically, they are also efficient antioxidant for membrane lipids. These antagonistic effects of these compounds elicited the challenging question of how the gene expression of antioxidant enzymes would respond to treatment with these compounds. The influence of RT3 and RT4 on the expression of antioxidant enzymes was investigated in cultured aortic smooth muscle cells (ASMC). RT-3 and RT4 promoted disruption of Ca^{2+} homeostasis, mitochondrial transmembrane potential ($\Delta\Psi$), and cell death in a dose-dependent manner. The cell death mechanisms responded qualitatively to the increase of the organotellurane concentration and changed from the predominance of apoptosis to necrosis. RT3 and RT4 increased significantly the expression of thioredoxin. RT3 increased also the expression of glutaredoxin and glutathione peroxidase, slightly the expression of catalase without significant

effects on SOD expression. The results are consistent with GSH and protein thiol depletion and discussed on basis of the mechanism of cell toxicity exhibited by these compounds.

Introduction

Tellurium forms inorganic and organic derivatives, both with a multiplicity of biological effects that are attractive due to the potential therapeutic applications and technological applications.[1–10] The organic derivatives of tellurium, are compounds with at least, one tellurium-carbon ligation in their structures.[2–4,11] Organotellurium comprise the groups of the divalent derivatives tellurols, diorganoditellurides, and diorganotellurides, and the hypervalent derivatives that are the organotellurium trihalides, diorganotellurium dihalides, organotellurium oxides, organotellurates, and organopertelluranes.[2] Most of the biological effects of tellurium compounds are related to the reactivity with thiol groups.[2,12–15] The inorganic tellurium compounds AS-101 and SAS form stable $Te-Cys_4$ complexes by replacing of labile ligands by four equivalents of cysteine.[2,16] Silberman et al demonstrated that distinct of inorganic tellurium compounds, the diaromatic organotelluranes of the type $Te^{IV}Ar_2X_n$ (X = Cl, O and CH_3COO) do not form stable complexes with cysteine.[3,16] The reaction mechanism of diaromatic organotelluranes forms complexes with two cysteines that are intermediates of the final products, the $Te^{II}Ar_2$ and cystine.[16] The effects of trichlorotelluro-dypnones on the bioenergetics of cells and their relationship with thiol reactivity were also investigated.[14] For these compounds, it was proposed a reaction mechanism like to SN2-type and SN1-type that are associative and dissociative mechanism.[14] All the postulated mechanisms implicate in the depletion of thiol groups with repercussions on the cell redox balance.[17–21] GSH is considered the most abundant redox buffer in cells and acts in conjunction with proteins, particularly the TRX system (TRX, TRX reductase, and NADPH) in maintaining the intracellular redox homeostasis.[17,22–27] The cells have redox-sensitive transcription factors that regulate gene expression in response to changes in the cellular redox status. The pathway of redox-sensitive gene expression involves agents acting

as sensors, transducers and effectors.[28–31] The redox state of cysteine residues is a key signal for a cellular gene expression in response to loss of redox homeostasis. In this regard, distinct oxidative states of cysteine residues elicit specific responses in cells. The reactivity of thiol groups depends on their protonation state. The thiolate group ($-S^-$) is more reactive with its conjugate acid form (-SH).[32,33] Also, the oxidation of protein bound thiol groups depends on the accessibility to the oxidative agents and different derivatives can be formed, i.e. sulfenic (-SOH), sulfinic (SO_2H) and sulfonic (SO_3H) acids.[34,35] The sulfenic and sulfinic groups are reversible oxidized states of thiol group while sulfonic acids represent irreversible modification of proteins. The thiol group of proteins are also targeted by the nitrosative species nitric oxide ($NO^\bullet$) and peroxynitrite ($ONOO^-$) that is formed by the reaction of $NO^\bullet$ with superoxide ion ($O_2^{-\bullet}$). The toxicity of nitrosative species results of covalent binding of $NO^\bullet$ to the thiol group of proteins leading to the formation of S-nitrosothiols (SNOs)[36] and the occurrence of protein nitration by the appending of nitro group ($-NO_2$) to a tyrosine residue.[37] A diversity of enzymes and transcription factors responds to the oxidative stress such as peroxiredoxins,[38–42] the transcription factors NF-κB and AP-1,[43–46] the negative regulator of Nrf2, protein 1 (Keap1),[47,48] thioredoxin[49–53], tyrosine phosphatases (PTP),[54,55] and the kinase MEKK1.[56–58] As an example, the transcription factor Nrf2 controls hundreds of genes involved in the combat of the oxidative stress. Nrf2 is repressed by association with the homodimeric form of the Keap1 protein. Human Keap1 has several of its 25 cysteine residues with low pK_a and consequently, they are highly reactive in the cell conditions. Therefore, Keap1 protein regulates Nrf2 in response to the cell redox state.[59–62] The redox-sensitive gene expression is also regulated by S-glutathionylation, that is the reversible formation of disulfide bonds between GSH and cysteinyl residues.[59,63,64] The S-glutathionylation is a negative regulatory process for tyrosine hydroxylase,[65] PKC-α,[66]

MEKK1 kinase,[56] PTP1B,[67,68] and inhibitor of kappa B kinase-β, which inhibits NF-κB[69,70] that in turn is reversed by glutaredoxin (Grx).[71,72] TRX is a low molecular weight disulfide oxido-reductase enzyme with 10-12 kDa. TRX has two reactive cysteine residues that forms a disulfide bond upon oxidation and that is recycled the action of TRX reductase and NADPH.[73] TRX exerts its antioxidant activity by the enzymatic action on other enzymes and transcription factors as well as the scavenging of reactive oxygen species.[2573] TRX is overexpressed in conditions in which the redox-active cysteines are inactivated by heavy metals, and reactive species produced by ischemia–reperfusion and viral infection.[74–76] In the present study, the toxicologic effects of RT-03 and RT-04 (Fig. 1) on aortic smooth muscle cells (ASMC) and the repercussions on the expression of thioredoxins (TRx), thioredoxin reductase, glutaredoxin, SOD, catalase, and glutathione peroxidase were investigated.

Figure 1. Molecular structures of RT-03 and RT-04.

Materials and Methods

Synthesis of organotelluranes

The organotelluranes used (RT-03 and RT-04) were synthesized by the electrophilic addition of p-methoxyphenyl tellurium trichloride to alkynes, called 1-ethynyl-1-cyclohexanol and 3-methyl -3-hydroxy-butyne, respectively.[77]

Animal handling and euthanasia

Rats of the Wistar lineage, maintained and euthanized using procedures approved by the internal ethics committee registered at COBEA, which supervises all norms and ethical principles in maintenance of animals, such as: space, ventilation, temperature, nutrition, hydration and proper handling. Only males of the Wistar lineage were used in this research, the euthanasia method used was the cervical dislocation for rats up to 200 grams, and cranial trauma for animals over 200 grams, these methods are described in Guide to the care and use of experimental animals, Canadian Council on Animal Care (CCAC).[78] The chemical method of euthanasia cannot be used because in this type of study it is contraindicated due to the interaction of the lethal chemical agents with membranes, interfering in the transport of ions, among other effects.

Isolation of hepatic mitochondria

Mitochondria were isolated from the liver of male Wistar rats of approximately 200 g using the differential centrifugation technique.[79] The liver was removed after the animal's death and washed in a solution containing 250 mM sucrose, 10 mM HEPES-KOH buffer and 1 mM EGTA at pH 7.4 at 4 ° C, chopped into small fragments and homogenized in Potter-Elvehjen. The

suspension was centrifuged at 580g for 5 minutes and the supernatant purchased was centrifuged at 10,300g for 10 minutes. The resulting supernatant was discarded along with the upper lipid phase with Pasteur pipette, and the resulting pellet resuspended in approximately 25 ml of medium containing 250 mM sucrose, 10 mM HEPES-KOH buffer and 0.3 mM EGTA at pH 7.4 at 4 °C and centrifuged at 3400g for 15 minutes. The final pellet or mitochondrial fraction was resuspended in a solution containing 250 mM sucrose, 10 mM HEPES-KOH buffer at pH 7.4 at 4 ° C, at a concentration of approximately 100 mg of protein per ml.

Dosage of proteins

The protein concentration was quantified by the Biuret method, where an aliquot (10 mL) of the sample to be determined was mixed with 100 µL of a solution of 5% deoxycholic acid (m / v) and water q.s. 1.5 mL. To this mixture was added 1.5 mL of the biuret reagent, composed of 0.15% (m/v) cupric sulfate, 0.6% (w/v) sodium and potassium tartrate and 0.75 M NaOH. After 10 min, the absorbance was determined at 540 nm against a blank of reagents and a 1% BSA solution was used as standard.[80][81]

ANT extraction of rat liver mitochondria

Isolation of ANT was done according to Rück et al, 1998.[82] All steps were done at 4°C. Isolated liver mitochondria (20 mg) in 1 ml of isolation medium were incubated for 5 min with an equal volume of extraction buffer consisting of 40 mM KH_2PO_4, 40 mM KCl, 2 mM EDTA and 6% Triton X-100, at pH 6.0. The suspension was centrifuged for 30 min at 24,000 g and the supernatant was loaded onto a column packed with 1g of dry hydroxyapatite (Bio-Rad). After elution with extraction buffer the fractions collected containing proteins were diluted in 20 mM MES buffer, 0.2 mM EDTA and 0.5% Triton X-100 at pH 6.0 in the ratio 1: 1. This sample was

applied to a 1-ml cation exchange HiTrapSP column (Pharmacia), connected with a FPLC system and eluted by a NaCl gradient (0.1 M NaCl in column buffer). Fractions were analyzed by 12% SDS-PAGE and by immunodetection by anti-ANT, porin, CyPD, and Bax antibodies after transfer to nitrocellulose sheets.

MALDI-TOF Spectrometry

The mass spectra were acquired in a MALDI-TOF Pro mass spectrometer device that can operate in linear or reflectron mode using harmonic reflecton which increases the resolution and narrows the focusing time. The samples were mixed with saturated solution (5 mg / ml) of 3,5-dimethoxy-4-hydroxycinnamic acid in 50% acetonitrile and 0.5% TFA, and 0.5 ml of the mixture was loaded onto the MALDI stainless steel films for analysis.[83]

Culture of rabbit aortic smooth muscle cells

Maintenance of rabbit aortic smooth muscle cell (ASMC) culture was performed every 3 days. The removal of the cells from the flasks, when confluent, was performed by the trypsinization process. After determining cell viability, 1 x 10^6 cells were distributed in flasks of 75 cm^3 and grown in DMEM medium (Sigma Chemical Co.) supplemented with 10% fetal bovine serum (BFS, Gibco, BRL-USA), and kept in a 5% of CO_2 at 37ºC. The trypsinization for cell handling was done by discarding of the culture medium followed by cell washing, at least 2 times with PBS solution. In sequence, a 2.5% trypsin solution containing 0.25% EDTA was added to the flask and held for 2 minutes at 37 °C to inactivate the reaction and then culture medium supplemented with BFS was added in a volume 2 times greater than that of trypsin used. After cells were detached from the flask, the solution was centrifuged at 450 g for 10 min at 20 °C and cell viability was determined.[84]

Cell viability assays

The yellow tetrazole MTT (3- (4,5-dimethylthiazolyl-2) -2, 5-diphenyltetrazol bromide) is metabolically reduced by living cells, in part by the action of dehydrogenase enzymes, to generate reduced equivalents such as NADH and NADPH. In the intracellular medium of viable cells, the reduction of the yellow tetrazole salt gives the purple formazan salt, which can be solubilized and quantified by spectroscopic measurements. The assay for EC50 determination was performed in 96-well plate, in each well, 2×10^5 cells were added, at least 12 hours after cell distribution that is the minimum time for cells to adhere on the plate. After the incubation time had elapsed, 10 μl of the MTT solution (5mg/ml) (Invitrogen Co., USA) was added to each well and the plate incubated for an additional 4 hours. The whole procedure was performed in the dark due to photosensitivity of the MTT reagent. After the incubation time, 100 μl of a 10% SDS solution in 0.01 M HCL was added to each well, and after incubation for 9 to 12 hours the absorbances at 570 and 650 nm of each well was read using a reader of (Biotek ELX800). Blue trypan assay was performed on 25 x 13 mm petri dishes containing 5 x 10 5 cells. After incubation, 200 μl of the 0.4% tripan blue solution was added and nine fields / plate were analyzed, 50 cells / field counted, in a total of three plates per concentration. The percentage of dead cells was calculated in relation to the total cells present in the field. For the V-FITC/7-AAD assay, the cells were trypsinized and centrifuged 500 g for 5 min. The binding medium containing 10 mM of HEPES, 140 mM NaCl, 2.5 mM $CaCl_2$, pH 7.4, and 1 μg/mL annexin V were added to the precipitate. Cells were incubated in light-protected propylene tubes and kept at room temperature for 15 min. Detection of the percentage of cells in apoptosis and necrosis was determined on a flow cytometer (Beckman Coulter, Lab Cell Quanta SC MPL model).

Intracellular Ca^{2+} measurements

Rabbit aortic cells were loaded with 10 μM of Fura-2AM. Before starting the experiments, the cells were washed with Ca^{2+}-free buffer to avoid interference of the external Ca^{2+}. Ca^{2+} levels were measured on isolated intact cells using a high resolution inverted digital microscope that was coupled to a CCD camera controlled by a software.

Intra-cell mitochondrial membrane potential

Rabbit aortic cells were loaded with 1 μM TMRM (tetramethyl rhodamine) for 30 min. Before starting the experiments, the cells were washed with fluorescence buffer. Fluorescence was monitored in isolated intact cells using a high-resolution digital microscope with an inverted microscope coupled to a CCD camera.

Oxidation of total thiol groups

After induction of thiol group oxidation by the organotelluranes or 0.6 mM t-BOOH, the thiol content of cells was determined using 5,5'dithiobis (2-nitrobenzoic acid) (DTNB). The samples were centrifuged 2 min at 10,000 rpm and the pellet was treated twice with 200 μL of 6.5% trichloroacetic acid (TCA). After the addition of TCA, the samples were submitted to centrifugation at 10,000 g during 5 min to precipitate the protein content. The final pellet was resuspended in 1 mL of the cell medium in which 100 μM 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB), 0.5 mM EGTA, and 0.5 M Tris, pH 8.3 were present. The colorimetric determination of SH was measured by the absorbance intensity at 412 nm, using cysteine for calibration.[85]

Results and Discussion

Effects of RT-03 and RT-04 on ASMC viability and bioenergetics.

The ASMCs were incubated with organotelluranes RT-03 and RT-04 at the concentrations of 0, 10, 25, 50, 75, and 100 μM for one hour and the cell viability and death mechanisms were determined by flow cytometry using the annexin V-FITC/7-AAD kit. (Figure 2A and B).

Figure 2. *Flow cytometry of MLAC cells incubated with organotelluranes.* MLAC cells (1x10^5 cells) were incubated for 1h at 37 °C under 5% CO_2 atmosphere, with RT-03 (panel A) and RT-04 (panel B). The insets show the curves for viable cells EC_{50} for each compound treatment. The percentages of apoptotic and necrotic cells were determined by flow cytometry using double labelling with annexin V-FITC and 7-AAD.

In viable cells, phosphatidylserine (PS) is restricted to the inner leaflet of the plasma membrane at expenses of ATP and it is not accessible for the specific staining with fluorescein-labeled annexin V. In the early step of apoptosis, the asymmetric PS distribution is randomized and makes the phospholipid accessible for labeling. During this phase, the membrane of dying cells is stained by fluorescein-labeled annexin V before the occurrence of DNA hydrolysis and lose of cellular morphology. Therefore, the annexin V staining indicates that cells are in the early apoptosis. The progression of apoptosis allows the staining of cells by the DNA-specific viability

dye 7-AAD and characterizes late apoptotic cells.[868788] Otherwise, the staining only with 7-AAD characterizes necrotic cells. Figures 2(a), and 2(b), lower panels, show the results of cell death mechanisms promoted by RT-03 and RT-04, respectively. The data of cell viability were used for the calculation of EC_{50} (Figures 2, upper panels). Figure 2A and B, upper panels, show the percentage of viable cells as a function of the logarithm of organotellurane concentration. In this condition, the EC_{50} for cell viability was calculated by fitting the data by equation 1.

$$\% \; cell \; death = \; A_1 + \frac{(A_2 - A_1)}{(1 + 10^{(logx_0 - x)p})} \qquad \textbf{(1)}$$

Where, A_1 and, A_2 are respectively the upper and lower asymptotic of the curve, p (Hill slope) the tangent slope to the curve and EC_{50} calculation from the curve turning point $(logx_0)$. The EC_{50} determined for RT-03 e RT-04, were, respectively = 34±4 and 50±5 μM. Cell viability was also determined using Trypan blue assay (not shown). The same EC_{50} for RT-03 (34±3 μM) was obtained using Trypan blue. The EC50 of RT-04 determined by Trypan blue assay was 64±3 μM and is close of the value obtained from flux cytometry data (50±5 μM). These results show the correlation between the events measured by both techniques. Therefore, the uptake of telluranes by the cells produces a toxic effect that triggers the mechanisms of programmed cell death whose events that signal its occurrence were measured by Annexin V / 7-AAD. At the same time these events lead to permeabilization of the plasma membrane which is measured by the uptake of Tripan blue. It is important to note that the significantly lower EC_{50} of RT-03 in relation to RT-04 (32-47%, considering Annexin V / 7-AAD and trypan blue, respectively) is consistent with the results previously obtained in isolated rat liver mitochondria (IRLM).[14] In IRLM, RT-03 was also more efficient than RT-04 to promote damages in the organelle. The treatment of ASMCs with RT-03 at concentrations of 10-25 μM increased the percentage of

apoptosis and necrosis by 31% and 44% respectively over the control. The use of 10-25 μM of RT-04 increased apoptosis up to 50% and necrosis up to 164%. Although the percentage of apoptosis had increased significantly after the treatment with 25 μM of the organotelluranes, around 75% of ASMCs remained viable. Bellow EC50 apoptosis responded predominantly for the cell death and above the EC_{50}, necrosis was the predominant cell death mechanism promoted by the organotelluranes. The cell viability EC_{50} values of RT-03 and RT-04 were close to the EC_{50} of SH depletion that were, respectively, 60.2 ± 3 and 68.7 ± 3 μM (Figure 3). Figure 3 shows also that RT-03 and RT-04 did not deplete GSH of ASMCs. These results suggest that cell death promoted by the organotelluranes are related to the depletion of protein thiol groups but not to the oxidative stress. In IRLM, RT-03 and RT-04 promoted depletion of both protein thiol and GSH. The concentration of mitochondrial and cytosolic GSH are both in the range of 10-15 mM.[89] However, the enzymatic machinery to synthesize GSH is present only in cytosol as well as the pentose-phosphate pathway that is an important source of NADPH (48). The organotelluranes, RT-03 and RT-03 react as Lewis acids leading to protein SH depletion according to two possible mechanisms that were previously postulated: bis alkylation and redox mechanism.[12]

Figure 3. Effect of RT-03 and RT-04 on the SH and GSH contents of ASMC.

Despite, different effects of the organotelluranes on GSH content of IRLM and ASMCs, loss of $\Delta\Psi$ and Ca^{2+} homeostasis were observed for ASMCs (Figure 4). Figure 4 shows the effects of RT-03 on the cytosolic Ca^{2+} concentration (black line). The cytosolic Ca^{2+} concentration was labelled by Fura-2AM fluorescence that shifts the spectrum peak from 380 to 340 nm after Ca^{2+} binding. The 340/380 nm intensity ratio increased progressively to the maximum within 5 min in response to the increase of cytosolic Ca^{2+} concentration. In the following, cytosolic Ca^{2+}

concentration decreased due to mitochondrial uptake and attained the minimal around 25 min after the organotellurane addition. In the same conditions, the mitochondrial transmembrane potential ($\Delta\Psi$) was measured for ASMC after the addition of the organotellurane. The loss of $\Delta\Psi$ promoted by RT-03 was accompanied by decay of TMRM (gray line) fluorescence at 340 nm. The halftime (t_1) of TMRM fluorescence decay was determined as 7.8 min by fitting the data using equation 2,

$$IRF = y_0 + A_1 \times e^{(-x/t_1)} \qquad \textbf{(2)}$$

Where y_0 is the offset, A_1 the decay amplitude and t_1, the halftime of the decay. The halftime of $\Delta\Psi$ leakage was coincident with the maximum of the cytosolic Ca^{2+} overload and indicated by the vertical red dotted line in Figure 4. The cytosolic Ca^{2+} buffering promoted by the mitochondrial uptake contributes to the organelle permeabilization. The results obtained with RT-04 was very similar (not shown). The inset of Figure 4 shows snapshot of ASMC immediately and 13 min after the addition of RT-03 as indicated by the letters *a* and *b*. Loss of $\Delta\Psi$ promoted by RT-03 and RT-04 was also observed using JC-1 as marker (not shown).

Figure 4. RT-03-promoted changes in Ca^{2+} homeostasis and mitochondrial permeabilization. ASM cells were grown on glass coverslips and incubated with TMRM and Fura-2AM in the presence of 50 μM of organotelluranes. The black line represents the 340/380 nm ratio of Fura-2AM fluorescence that increases proportionally to the cytosolic Ca^{2+} concentration. The gray line represents the ΔΨ dissipation curve that was fitted by equation 2 (green dotted line). The vertical red line marks the half-life of the ΔΨ decay that is coincident with the maximum of Ca^{2+} overload in cytosol. The inset shows the snapshot of ASMC labeled by TMRM at the times indicated by the green arrows.

Considering the effect of the organotelluranes on mitochondrial bioenergetic, it was investigated the capacity of RT-03 and RT-04 to promote the thiol depletion of isolated adenine nucleotide translocase was investigated (Figure 5). The organotelluranes decreased drastically the thiol content of isolated ANT that is consistent with mitochondria permeabilization promoted by these compounds.

Figure 5. Effect of organotelluranes on the thiol content of ANT. ANT was isolated and incubated for 15 min in the presence of RT-03 e RT-04 and the thiol content determined colorimetrically at 412 nm after the reaction with DTNB.

The effects of organotelluranes (50 $\square$M) on the $\Delta\Psi$, SH content and Ca^{2+} homeostasis of ASMC culminated with cell death that was evidenced by changes in the nuclear morphology (Figure 6).

Figure 6. Effect of RT-03 on ASMC morphology. Snapshots show ASMCs nucleus after 15 h of incubation in the absence (left panel) and presence (right panel) of 50 µM of RT-03. The cells were incubated during 15 and the nucleus labeled with Hoeschst 33342.

Figure 6 shows the snapshots representative of cell morphology after 15 h of incubation with RT-03. After incubation, cells were treated with aqueous solution of Hoechst 33342 (1µg/ml), for 15 min. The images show the normal morphology of control cells (left panel) and the typical apoptotic nucleus with the characteristics of condensation and pycnoses. RT-03 promoted also loss of ASMC adherence and even the cells that remained stuck on the plaque presented pycnotic nucleus. In the following, the effect of RT-03 and RT-04 on the expression of proteins related to redox balance of cells was determined by real time PCR (Figure 7A-L).

Figure 7. Effect of RT-03 and RT-04 concentrations on the cell death mechanisms and the expression of antioxidant enzymes.

In Figure 7A-L the data of RNAm production was shown overlapped with the percentage of apoptotic and necrotic ASMCs that were treated with the corresponding concentration of RT-03 and RT-04. Thioredoxin (TRX) had the expression extremely enhanced by the organotelluranes, particularly, RT-03. The results obtained with 100 μM of RT-03 suggests a total uncontrol of the TRX RNAm production. Contrarily, the organotellurane RT-03 did not affect significantly the expression of thioredoxin reductase (TRXr) while RT-04 promoted an inhibition its expression. In the case of GRX expression, only 100 μM of RT-03 promoted a significant increase of the RNAm production. The expression of superoxide dismutase (SOD) increased more significantly in ASMCs treated with 20 μM of organotelluranes and in this condition, the more severe effect was observed for RT-04. The concentration of 20 μM produced the greatest increase of apoptosis as the preponderant mechanism of cell death. In concentrations >20 μM, the expression of SOD was inhibited by both the organotelluranes, and more significantly by RT-04. The expression of catalase was also decreased significantly in ASMCs treated with 50 mM of RT-04 but not by RT-03 that duplicate the enzyme expression only at 100 μM. RT-03 also promoted increase of

glutathione peroxidase (GPx) expression in a concentration-dependent manner. RT-04 also increased the expression of GPx, except at the concentration of 50 μM in which a significant decrease of the enzyme expression was observed.

Discussion

Organotelluranes have a particular reactivity with the thiol groups herewith their biological activities are rationalized. The reaction of a thiol with organotelluranes is still a matter of debate since the rate of such processes difficult to access in experimental model systems; an early attempt was conducted by Albeck and coworkers who proposed two mechanistic pathways by which these reactions would occur, an associative and dissociative mechanisms (Scheme 1). [2,14]

Scheme 1. Proposed ligand exchange reactions of dichloro organotelluranes.

The difference of both mechanisms lies in the nature of the reaction intermediate as an anionic pentavalent intermediate is formed by addition of the nucleophile in the former mechanism whereas a cationic trivalent specie is formed in the later prior to the reaction with the thiol. Interestingly, orgatelluranes RT-04 and a dichloride derivative of RT-03 shows a remarkable stability in aqueous dimethylsulfoxide solutions, even in acidic media, showing that DMSO solutions of organotelluranes are very stable remaining in the form of dichlorides.[91] In general, the most commonly exploited organotelluranes in biological studies have a limited solubility in aqueous media thus theoretical studies have been carried out to investigate the fate of such species in water media since the hydrolysis can be anticipated. These studies indicate that after a

series of ligand-exchange reactions, a thermodynamically stable telluroxide may be formed which is reactive towards thiols, nonetheless[3,92]. The spontaneous formation of telluroxides from dihalo diorganotelluranes possessing an intramolecular coordination ether moiety further corroborated these theoretical studies.[92] Telluroxides may likely react with thiols but through a nucleophilic addition mechanism forming an monothiolate that can react with another thiol leading to a dithiolate that can disproportionate forming a telluride and the disulfide (Scheme 2).

Scheme 2. Reactions of thiols with organotelluroxide.

Organotellurane monothiolates originated in reactions with macromolecular thiols are likely to be kinetically more stable than those formed in reactions with low molecular weight thiols as cysteine, whose formation of the dithiol is more favorable due to lower steric hindrance. Whatever the mechanism, an intermediate protein-tellurane complex might form and become a target for the reaction with GSH and TRX. Also, the direct reaction of cysteine residues of TRX with organotelluranes cannot be discarded as it possesses more nucleophilic thiols than those of GSH and TRX. The protein-tellurane complexes can react with antioxidant proteins as TRX, GRX, and GSH by bis-thiolation leading to a redox pathway. In the redox pathway, glutathionylated protein or a mixed disulfide complex with TRX could be formed and subsequently recycled by TRXr and GRXr. However, inhibition of TRXr, previously described for some of organotelluranes, could prevent the recovery of protein thiol groups.[90] In the case of TRX, cells responded to the impaired recycling of TRX with its overexpression. In this regard,

RT-03 could more strongly inhibit TRXr and form more stable complexes with proteins leading to more drastic effects of TRX and GRX overexpression. The structure-dependent reactivity of RT-03-protein complexes via bis-thiolation (i.e., the formation of dithiolates) or redox mechanisms remains uncovered.

Before analyzing of the concentration-dependent effect of RT-03 and RT-04 on the expression of antioxidant enzymes, it is important to consider that for RT-03, but not for RT-04, the concentrations of 50 and 100 µM lead to a similar percentage of cell death. Also, RT-03 promoted overexpression of Grx, catalase, and GPx only at the concentration of 100 µM. Therefore, it is reasonable to consider that 100 µM of RT-03 promoted a severe disruption of cellular redox control and their effects are not comparative with the corresponding concentration of RT-04. Thioredoxin was the protein with the highest expression increase in response to treatment with RT-03 and RT-04. Overexpression of Trx was observed even below the EC_{50} of RT-03 and RT-04. Apart from the effect of 100 µM of RT-03, Grx was not overexpressed by ASMCs treated with 20 and 50 µM of RT-03 and had a decreased expression in the presence of 50 µM of RT-04. The expression of TrxR was not significantly affected by RT-03 and was inhibited by RT-04, more significantly above the EC_{50}.

These results can be understood based on the following reasoning. The reaction mechanisms of telluranes with thiols have been extensively studied and can comprise two stages: the formation of the tellurium-sulfur bond forming a monothiolate and possible secondary reactions of the Te-protein complexes with nucleophiles. The formation of organotellurane adducts with a protein thiol group may occur through associative and dissociative mechanisms described previously. The protein complexed with the organotellurane may be the target of nucleophilic attack by

glutathione, another protein, or another cysteine residue of its chain. Thus, the tellurane-protein adduct can be glutathionylated or form intra- or inter-molecular disulfides (RS-SR).

In any of these conditions, the protein may be recycled to its native form by the action of the Grx and Trx systems. Glutathionylated proteins can be reduced by the action o Grx and other disulfides by the Grx and Trx systems. Organotelluranes can inhibit TrxR impairing the recycling of oxidized Trx to the reduced form. The high consumption of reduced Trx without recycling by TrxR could lead cells to respond with the overexpression of the enzyme.

The absence of glutathione depletion suggests that the balance of oxidation and recycling of the peptide was not affected by RT-03 and RT-04 at the concentration range of 20-100 µM. Notwithstanding, GSH was not the preferential nucleophile that reacted with the Te-protein adducts. The reaction with GSH should be unfavorable because the pKa of its thiol group is 8.3, and most of the thiol is therefore protonated. Also, the organotelluranes are highly hydrophobic and should partition preferentially inside cell membranes. Taken together, these results suggest that membrane thiol groups are the principal nucleophiles reacting with RT-03 and RT-04 in ASMCs. Previously, it was demonstrated that RT-03 and RT-04 are also an efficient antioxidant for membrane lipids. In low concentrations (nanomolar scale), these organotelluranes acted as efficient antioxidants without promoting organelle damage. In high concentrations, RT-03 and RT-04 promoted the opening of the MPTP associated with thiol depletion but without lipid oxidation.

Consistently, 20 and 50 µM of RT-03 did not significantly affect the expression of SOD and catalase. Above the EC_{50}, RT-03 increased the expression of GPx. However, GPx-1 is a selenocysteine-containing enzyme, and its expression has unique forms of regulation that

involves the trace mineral selenium. Therefore, it is not surprising that tellurium-containing compounds could affect the expression of GPx. Further, GPx controls the cellular levels of hydrogen peroxide and consequently the signal transduction involving redox-sensitive cysteines. Several proteins are redox regulated by TRX, such as NF-κB subunit p50, hypoxia-inducible factor 1-α, the tumor suppressor p53, c-Fos/c-Jun complexes, and others. Overexpression of thioredoxin in cancer cells is associated with apoptosis inhibition. In pancreatic cancer, the complexation of TRX with apoptosis signal-regulating kinase 1 (ASK1) inhibits apoptosis. TRX binds to the N-terminal noncatalytic region of ASK1 that encompasses the amino acid residues 1-655 leading to the enzyme inhibition. Consistently, necrosis rather than apoptosis contributed to cell death promoted by RT-03 and RT-04.

Conclusions

The organotelluranes RT-03 and RT-04 have peculiar effects on the redox balance by attacking specifically the thiol groups of proteins with concomitant protection of membrane lipids. The attack of protein thiol groups and particularly the capacity to inhibit TRXr affected mitochondrial bioenergetic and consistently lead to a significant increase in TRx, GRx, and glutathione expression levels peroxidase. Three levels of damage previously observed in RLM treated with the organotelluranes were reproduced in ASMCs. In RLM, RT-03 and RT-04 promoted mitochondrial transition pore opening at the concentration range of 5-10 mM, alterations in the bilayer fluidity, phosphorylation impairment, uncoupling, transmembrane potential disruption, and depletion of mitochondrial reduced thiol groups at the range of 15-30 mM promoted, and complete RLM respiratory inhibition at 100 mM. Consistently, ASMCs exhibited a maximum of a 2-fold increase in apoptosis rate after the treatment with 25 mM of the organotelluranes without the significant occurrence of necrosis that increased fifty times at the

concentration of 75 mM of RT-03 and RT-04 and total uncontrol of the RNAm production at 100 mM. The damaging effects of RT-03 and RT-04, both in rat liver mitochondria and cells, are related to the reactivity with thiol groups of proteins. These findings are essential for comprehending how organotellurane affects cells, essential for further therapeutic applications of these compounds.

Acknowledgments

All acknowledgments (if any) should be included at the very end of the paper before the references and may include supporting grants, presentations, and so forth.

References

Corresponding Author

Iseli Lourenço Nantes-Cardoso, e-mail ilnantes@ufabc.edu.br, and ilnantes@gmail.com, Laboratory of Nanostructures for Biology and Advance Materials, NanoBioMAv, Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, Santo André, SP, Brasil, Av. dos Estados, 5001, Bairro Bangu, CEP 09210-580.

Rodrigo L. O. R. Cunha, e-mail rodrigo.cunha@ufabc.edu.br, Laboratory of Chemical Biology, Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, Santo André, SP, Brasil, Av. dos Estados, 5001, Bairro Bangu, CEP 09210-580.

Author Contributions

The manuscript was written through the contributions of all authors. All authors have approved the final version of the manuscript.

Funding Sources

FAPESP (2015/17688-0, 2017/02317-2), CAPES, finance code-001 and CNPq (309247/2017-9 and 305818/2018-0). National System of Nanotechnology Laboratories – SisNANO, Proc. CNPq 402289/2013-7.

ACKNOWLEDGMENT

Authors thank FAPESP, CNPq, CAPES, and the UFABC´s Strategic Research Units NBB and NuTS for the financial support and David da Mata Lopes for the technical assistance.

REFERENCES

(1)   Comasseto, J. V.; Cunha, R. L. O. R.; Clososki, G. C. Tellurium. In *Comprehensive Organometallic Chemistry III*; Elsevier, 2007; pp 587–648. https://doi.org/10.1016/B0-08-045047-4/00121-7.

(2)   Cunha, R. L. O. R.; Gouvea, I. E.; Juliano, L. A Glimpse on Biological Activities of Tellurium Compounds. *An. Acad. Bras. Cienc.* **2009**, *81* (3), 393–407. https://doi.org/10.1590/S0001-37652009000300006.

(4)   Pietrasiak, E.; Togni, A. Synthesis and Characterization of Fluorinated Hypervalent Tellurium Derivatives. *Organometallics* **2017**, *36* (19), 3750–3757. https://doi.org/10.1021/acs.organomet.7b00535.

(5)   Srivastava, K.; Panda, A.; Sharma, S.; Singh, H. B. Telluroxanes: Synthesis, Structure and Applications. *J. Organomet. Chem.* **2018**, *861*, 174–206. https://doi.org/10.1016/J.JORGANCHEM.2018.02.036.

(6)   Heimann, S.; Schulz, S.; Bläser, D.; Wölper, C. Syntheses and Solid-State Structures of Bis(Dialkylstibanyl)Sulfanes and -Telluranes. *Eur. J. Inorg. Chem.* **2013**. https://doi.org/10.1002/ejic.201300767.

(7)   Medina-Cruz, D.; Tien-Street, W.; Vernet-Crua, A.; Zhang, B.; Huang, X.; Murali, A.; Chen, J.; Liu, Y.; Garcia-Martin, J. M.; Cholula-Díaz, J. L.; Webster, T. Tellurium, the Forgotten Element: A Review of the Properties, Processes, and Biomedical Applications of the Bulk and Nanoscale Metalloid. In *Racing for the Surface: Antimicrobial and Interface Tissue Engineering*; 2020. https://doi.org/10.1007/978-3-030-34471-9_26.

(8)   Wang, H.; Chai, L.; Xie, Z.; Zhang, H. Recent Advance of Tellurium for Biomedical
        Applications. *Chemical Research in Chinese Universities*. 2020, pp 551–559.
        https://doi.org/10.1007/s40242-020-0193-0.

(9)   Hagar, M.; Roman, G.; Eitan, O.; Noam, B. Y.; Abrham, Z.; Benjamin, S. A Tellurium-
        Based Small Immunomodulatory Molecule Ameliorates Depression-Like Behavior in
        Two Distinct Rat Models. *NeuroMolecular Med.* **2020**. https://doi.org/10.1007/s12017-
        020-08603-5.

(10)   Machado, T. M.; Pereira da Silva, M. A. Preparation and Properties of Antimony Sodium
        Polyphosphate Glasses Containing Tellurium Particles. *Mater. Chem. Phys.* **2020**.
        https://doi.org/10.1016/j.matchemphys.2020.122731.

(11)   Nogueira, C. W.; Zeni, G.; Rocha, J. B. T. Organoselenium and Organotellurium
        Compounds: Toxicology and Pharmacology. *Chem. Rev.* **2004**, *104* (12), 6255–6286.
        https://doi.org/10.1021/cr0406559.

(12)   Pessoto, F. S.; Faria, P. A.; Cunha, R. L. O. R.; Comasseto, J. V; Rodrigues, T.; Nantes, I.
        L. Organotellurane-Promoted Mitochondrial Permeability Transition Concomitant with
        Membrane Lipid Protection against Oxidation. *Chem. Res. Toxicol.* **2007**.
        https://doi.org/10.1021/tx700092r.

(13)   Nantes, I. L.; Rodrigues, T.; Caires, A. C. F.; Cunha, R. L. O. R.; Pessoto, F. S.;
        Yokomizo, C. H.; Araujo-Chaves, J. C.; Faria, P. A.; Santana, D. P.; Dos Santos, C. G.
        Specific Effects of Reactive Thiol Drugs on Mitochondrial Bioenergetics. *J. Bioenerg.
        Biomembr.* **2011**. https://doi.org/10.1007/s10863-011-9328-9.

(14) Yokomizo, C. H.; Pessoto, F. S.; Prieto, T.; Cunha, R. L. O. R.; Nantes, I. L. Effects of Trichlorotelluro-Dypnones on Mitochondrial Bioenergetics and Their Relationship to the Reactivity with Protein Thiols. *Chem. Res. Toxicol.* **2015**, *28* (6), 1167–1175. https://doi.org/10.1021/tx5005166.

(15) Kheirabadi, R.; Izadyar, M. Computational Modeling of the Kinetics and Mechanism of Tellurium-Based Glutathione Peroxidase Mimic. *Int. J. Quantum Chem.* **2020**. https://doi.org/10.1002/qua.26201.

(16) Silberman, A.; Albeck, M.; Sredni, B.; Albeck, A. Ligand-Substitution Reactions of the Tellurium Compound AS-101 in Physiological Aqueous and Alcoholic Solutions. *Inorg. Chem.* **2016**, *55* (21), 10847–10850. https://doi.org/10.1021/acs.inorgchem.6b02138.

(17) Jones, D. P. Redefining Oxidative Stress. *Antioxid. Redox Signal.* **2006**, *8* (9–10), 1865–1879. https://doi.org/10.1089/ars.2006.8.1865.

(18) McBean, G. J.; Aslan, M.; Griffiths, H. R.; Torrão, R. C. Thiol Redox Homeostasis in Neurodegenerative Disease. *Redox Biology*. 2015. https://doi.org/10.1016/j.redox.2015.04.004.

(19) Paul, B. D.; Sbodio, J. I.; Snyder, S. H. Cysteine Metabolism in Neuronal Redox Homeostasis. *Trends in Pharmacological Sciences*. 2018. https://doi.org/10.1016/j.tips.2018.02.007.

(20) Benhar, M. Oxidants, Antioxidants and Thiol Redox Switches in the Control of Regulated Cell Death Pathways. *Antioxidants*. 2020. https://doi.org/10.3390/antiox9040309.

(21)   Allen, E. M. G.; Mieyal, J. J. Protein-Thiol Oxidation and Cell Death: Regulatory Role of Glutaredoxins. *Antioxidants and Redox Signaling*. 2012. https://doi.org/10.1089/ars.2012.4644.

(22)   McBean, G. Cysteine, Glutathione, and Thiol Redox Balance in Astrocytes. *Antioxidants* **2017**, *6* (3), 62. https://doi.org/10.3390/antiox6030062.

(23)   Munoz, L. E.; Janko, C.; Schorn, C.; Schett, G. Orum Eview Rticle. **2014**, *20* (6). https://doi.org/10.1089/ars.2013.5179.

(24)   Arnér, E. S. J.; Holmgren, A. Physiological Functions of Thioredoxin and Thioredoxin Reductase. *Eur. J. Biochem.* **2000**, *267* (20), 6102–6109. https://doi.org/10.1046/j.1432-1327.2000.01701.x.

(25)   Berndt, C.; Lillig, C. H.; Holmgren, A. Thiol-Based Mechanisms of the Thioredoxin and Glutaredoxin Systems: Implications for Diseases in the Cardiovascular System. *AJP Hear. Circ. Physiol.* **2006**, *292* (3), H1227–H1236. https://doi.org/10.1152/ajpheart.01162.2006.

(26)   Panday, S.; Talreja, R.; Kavdia, M. The Role of Glutathione and Glutathione Peroxidase in Regulating Cellular Level of Reactive Oxygen and Nitrogen Species. *Microvasc. Res.* **2020**, *131*, 104010. https://doi.org/10.1016/j.mvr.2020.104010.

(27)   Bjørklund, G.; Tinkov, A. A.; Hosnedlová, B.; Kizek, R.; Ajsuvakova, O. P.; Chirumbolo, S.; Skalnaya, M. G.; Peana, M.; Dadar, M.; El-Ansary, A.; Qasem, H.; Adams, J. B.; Aaseth, J.; Skalny, A. V. The Role of Glutathione Redox Imbalance in Autism Spectrum Disorder: A Review. *Free Radical Biology and Medicine*. 2020. https://doi.org/10.1016/j.freeradbiomed.2020.07.017.

(28)   Lens, T.; Cells, E.; Spector, A.; Yan, G.; Huang, C.; Mcdermott, M. J.; Gascoyneg, P. R. C.; Pigietll, V. The Effect of H202. *Biochemistry* **1988**, *263* (10), 4984–4990.

(29)   Hirota, K.; Matsui, M.; Murata, M.; Takashima, Y.; Cheng, F. S.; Itoh, T.; Fukuda, K.; Junji, Y. Nucleoredoxin, Glutaredoxin, and Thioredoxin Differentially Regulate NF-KB, AP-1, and CREB Activation in HEK293 Cells. *Biochem. Biophys. Res. Commun.* **2000**, *274* (1), 177–182. https://doi.org/10.1006/bbrc.2000.3106.

(30)   Gong, T.; Hu, Q.; Nie, X.; Liu, T.; Wang, H. Periodic Dynamic Regulation of MSCs Differentiation on Redox-Sensitive Elastic Switched Substrates. *ACS Appl. Bio Mater.* **2020**. https://doi.org/10.1021/acsabm.0c00245.

(31)   Messina, S.; De Simone, G.; Ascenzi, P. Cysteine-Based Regulation of Redox-Sensitive Ras Small GTPases. *Redox Biology*. 2019. https://doi.org/10.1016/j.redox.2019.101282.

(32)   Wall, S. B.; Oh, J. Y.; Diers, A. R.; Landar, A. Oxidative Modification of Proteins: An Emerging Mechanism of Cell Signaling. *Front. Physiol.* **2012**, *3 SEP* (September). https://doi.org/10.3389/fphys.2012.00369.

(33)   Castranova, V.; Asgharian, B.; Sayre, P.; Virginia, W.; Carolina, N. HHS Public Access. **2016**, 1922–2013. https://doi.org/10.1080/10937404.2015.1051611.INHALATION.

(34)   Jeong, J.; Jung, Y.; Na, S.; Jeong, J.; Lee, E.; Kim, M.-S.; Choi, S.; Shin, D.-H.; Paek, E.; Lee, H.-Y.; Lee, K.-J. Novel Oxidative Modifications in Redox-Active Cysteine Residues. *Mol. Cell. Proteomics* **2011**, *10* (3), M110.000513. https://doi.org/10.1074/mcp.M110.000513.

(35) Icimoto, M. Y.; Ferreira, J. C.; Yokomizo, C. H.; Bim, L. V; Marem, A.; Gilio, J. M.; Oliveira, V.; Nantes, I. L. Redox Modulation of Thimet Oligopeptidase Activity by Hydrogen Peroxide. *FEBS Open Bio* **2017**, *7* (7), 1037–1050. https://doi.org/10.1002/2211-5463.12245.

(36) Sun, J.; Steenbergen, C.; Murphy, E. S-Nitrosylation: NO-Related Redox Signaling to Protect against Oxidative Stress. *Antioxid. Redox Signal.* **2006**, *8* (9–10), 1693–1705. https://doi.org/10.1089/ars.2006.8.1693.

(37) Paes de Barros, M.; Casares Araujo-Chaves, J.; Marlise Mendes Brito, A.; Lourenço Nantes-Cardoso, I. Oxidative/Nitrative Mechanism of Molsidomine Mitotoxicity Assayed by the Cytochrome c Reaction with SIN-1 in Models of Biological Membranes. *Chem. Res. Toxicol.* **2020**. https://doi.org/10.1021/acs.chemrestox.0c00122.

(38) Kang, S. W.; Chang, T.-S.; Lee, T.-H.; Kim, E. S.; Yu, D.-Y.; Rhee, S. G. Cytosolic Peroxiredoxin Attenuates The Activation Of Jnk And P38 But Potentiates That Of Erk In Hela Cells Stimulated With Tumor Necrosis Factor-α. *J. Biol. Chem.* **2004**, *279* (4), 2535–2543. https://doi.org/10.1074/jbc.M307698200.

(39) Rhee, S. G.; Chae, H. Z.; Kim, K. Peroxiredoxins: A Historical Overview and Speculative Preview of Novel Mechanisms and Emerging Concepts in Cell Signaling. *Free Radic. Biol. Med.* **2005**, *38* (12), 1543–1552. https://doi.org/10.1016/j.freeradbiomed.2005.02.026.

(40) Jeong, W.; Park, S. J.; Chang, T.-S.; Lee, D.-Y.; Rhee, S. G. Molecular Mechanism of the Reduction of Cysteine Sulfinic Acid of Peroxiredoxin to Cysteine by Mammalian

Sulfiredoxin. *J. Biol. Chem.* **2006**, *281* (20), 14400–14407. https://doi.org/10.1074/jbc.M511082200.

(41)　Jiang, L.; Gong, Y.; Hu, Y.; You, Y.; Wang, J.; Zhang, Z.; Wei, Z.; Tang, C. Peroxiredoxin-1 Overexpression Attenuates Doxorubicin-Induced Cardiotoxicity by Inhibiting Oxidative Stress and Cardiomyocyte Apoptosis. *Oxid. Med. Cell. Longev.* **2020**. https://doi.org/10.1155/2020/2405135.

(42)　P., H.; J., Z.; L., S.; M., L.; Y., H.; Z., D.; Y., L. Brain-Derived Neurotrophic Factor Attenuates Doxorubicin-Induced Cardiac Dysfunction through Activating Akt Signalling in Rats. *J. Cell. Mol. Med.* **2017**. https://doi.org/10.1111/jcmm.13012 LK - http://resolver.ebscohost.com/openurl?sid=EMBASE&issn=15821838&id=doi:10.1111%2Fjcmm.13012&atitle=Brain-derived+neurotrophic+factor+attenuates+doxorubicin-induced+cardiac+dysfunction+through+activating+Akt+signalling+in+rats&stitle=J.+Cell.+Mol.+Med.&title=Journal+of+Cellular+and+Molecular+Medicine&volume=21&issue=4&spage=685&epage=696&aulast=Hang&aufirst=Pengzhou&auinit=P.&aufull=Hang+P.&coden=&isbn=&pages=685-696&date=2017&auinit1=P&auinitm=.

(43)　KLATT, P.; MOLINA, E. P.; DE LACOBA, M. G.; PADILLA, C. A.; MARTÍNEZ-GALISTEO, E.; BÁRCENA, J. A.; LAMAS, S. Redox Regulation of C-Jun DNA Binding by Reversible S-Glutathiolation. *FASEB J.* **1999**, *13* (12), 1481–1490. https://doi.org/10.1096/fasebj.13.12.1481.

(44)　Nishi, T.; Shimizu, N.; Hiramoto, M.; Sato, I.; Yamaguchi, Y.; Hasegawa, M.; Aizawa, S.; Tanaka, H.; Kataoka, K.; Watanabe, H.; Handa, H. Spatial Redox Regulation of a Critical Cysteine Residue of NF-KB *in Vivo*. *J. Biol. Chem.* **2002**, *277* (46), 44548–44556.

https://doi.org/10.1074/jbc.M202970200.

(45)   Gius, D.; Botero, A.; Shah, S.; Curry, H. A. Intracellular Oxidation/Reduction Status in the Regulation of Transcription Factors NF-KB and AP-1. *Toxicol. Lett.* **1999**, *106* (2–3), 93–106. https://doi.org/10.1016/S0378-4274(99)00024-7.

(46)   Attafi, I. M.; Bakheet, S. A.; Korashy, H. M. The Role of NF-KB and AhR Transcription Factors in Lead-Induced Lung Toxicity in Human Lung Cancer A549 Cells. *Toxicol. Mech. Methods* **2020**. https://doi.org/10.1080/15376516.2019.1687629.

(47)   Wakabayashi, N.; Dinkova-Kostova, A. T.; Holtzclaw, W. D.; Kang, M.-I.; Kobayashi, A.; Yamamoto, M.; Kensler, T. W.; Talalay, P. Protection against Electrophile and Oxidant Stress by Induction of the Phase 2 Response: Fate of Cysteines of the Keap1 Sensor Modified by Inducers. *Proc. Natl. Acad. Sci.* **2004**, *101* (7), 2040–2045. https://doi.org/10.1073/pnas.0307301101.

(48)   Priya Dharshini, L. C.; Vishnupriya, S.; Sakthivel, K. M.; Rasmi, R. R. Oxidative Stress Responsive Transcription Factors in Cellular Signalling Transduction Mechanisms. *Cell. Signal.* **2020**, *72*, 109670. https://doi.org/10.1016/j.cellsig.2020.109670.

(49)   Holmgren, A. Thioredoxin Structure and Mechanism: Conformational Changes on Oxidation of the Active-Site Sulfhydryls to a Disulfide. *Structure* **1995**, *3* (3), 239–243.

(50)   Sumbayev, V. V. S-Nitrosylation of Thioredoxin Mediates Activation of Apoptosis Signal-Regulating Kinase 1. *Arch. Biochem. Biophys.* **2003**, *415* (1), 133–136.

(51)   Shao, R.; Yang, Y.; Zhang, Y.; Zhao, S.; Zheng, Z.; Chen, G. The Expression of

Thioredoxin-1 and Inflammatory Cytokines in Patients with Sepsis. *Immunopharmacol. Immunotoxicol.* **2020**, *42* (3), 280–285. https://doi.org/10.1080/08923973.2020.1755309.

(52) Zeng, X. S.; Geng, W. S.; Wang, Z. Q.; Jia, J. J. Morphine Addiction and Oxidative Stress: The Potential Effects of Thioredoxin-1. *Frontiers in Pharmacology*. 2020. https://doi.org/10.3389/fphar.2020.00082.

(53) Rashidi, Z.; Aleyasin, A.; Eslami, M.; Nekoonam, S.; Zendedel, A.; Bahramrezaie, M.; Amidi, F. Quercetin Protects Human Granulosa Cells against Oxidative Stress via Thioredoxin System. *Reprod. Biol.* **2019**. https://doi.org/10.1016/j.repbio.2019.07.002.

(54) Denu, J. M.; Tanner, K. G. Specific and Reversible Inactivation of Protein Tyrosine Phosphatases by Hydrogen Peroxide: Evidence for a Sulfenic Acid Intermediate and Implications for Redox Regulation [†]. *Biochemistry* **1998**, *37* (16), 5633–5642. https://doi.org/10.1021/bi973035t.

(55) Hsu, M. F.; Koike, S.; Mello, A.; Nagy, L. E.; Haj, F. G. Hepatic Protein-Tyrosine Phosphatase 1B Disruption and Pharmacological Inhibition Attenuate Ethanol-Induced Oxidative Stress and Ameliorate Alcoholic Liver Disease in Mice. *Redox Biol.* **2020**. https://doi.org/10.1016/j.redox.2020.101658.

(56) Cross, J. V; Foss, F. W.; Rady, J. M.; Macdonald, T. L.; Templeton, D. J. The Isothiocyanate Class of Bioactive Nutrients Covalently Inhibit the MEKK1 Protein Kinase. *BMC Cancer* **2007**, *7* (1), 183. https://doi.org/10.1186/1471-2407-7-183.

(57) CROSS, J. V.; TEMPLETON, D. J. Oxidative Stress Inhibits MEKK1 by Site-Specific Glutathionylation in the ATP-Binding Domain. *Biochem. J.* **2004**, *381* (3), 675–683.

https://doi.org/10.1042/BJ20040591.

(58)   Matsushita, M.; Nakamura, T.; Moriizumi, H.; Miki, H.; Takekawa, M. Stress-Responsive
        MTK1 SAPKKK Serves as a Redox Sensor That Mediates Delayed and Sustained
        Activation of SAPKs by Oxidative Stress. *Sci. Adv.* **2020**.
        https://doi.org/10.1126/sciadv.aay9778.

(59)   Limón-Pacheco, J.; Gonsebatt, M. E. The Role of Antioxidants and Antioxidant-Related
        Enzymes in Protective Responses to Environmentally Induced Oxidative Stress. *Mutat.
        Res. - Genet. Toxicol. Environ. Mutagen.* **2009**, *674* (1–2), 137–147.
        https://doi.org/10.1016/j.mrgentox.2008.09.015.

(60)   Janssen-Heininger, Y. M.; Poynter, M. E.; Baeuerle, P. A. Recent Advances towards
        Understanding Redox Mechanisms in the Activation of Nuclear Factor KappaB. *Free
        Radic. Biol. Med.* **2000**, *28* (9), 1317–1327.

(61)   Krajka-Kuźniak, V.; Paluszczak, J.; Baer-Dubowska, W. The Nrf2-ARE Signaling
        Pathway: An Update on Its Regulation and Possible Role in Cancer Prevention and
        Treatment. *Pharmacological Reports*. 2017. https://doi.org/10.1016/j.pharep.2016.12.011.

(62)   Baird, L.; Yamamoto, M. The Molecular Mechanisms Regulating the KEAP1-NRF2
        Pathway. *Mol. Cell. Biol.* **2020**, *40* (13). https://doi.org/10.1128/mcb.00099-20.

(63)   Matsui, R.; Ferran, B.; Oh, A.; Croteau, D.; Shao, D.; Han, J.; Pimentel, D. R.;
        Bachschmid, M. M. Redox Regulation via Glutaredoxin-1 and Protein S-
        Glutathionylation. *Antioxidants and Redox Signaling*. Mary Ann Liebert Inc. April 1,
        2020, pp 677–700. https://doi.org/10.1089/ars.2019.7963.

(64) Mailloux, R. J.; Gill, R.; Young, A. Protein S-Glutathionylation and the Regulation of Cellular Functions. In *Oxidative Stress*; Elsevier, 2020; pp 217–247. https://doi.org/10.1016/b978-0-12-818606-0.00013-4.

(65) Borges, C. R.; Geddes, T.; Watson, J. T.; Kuhn, D. M. Dopamine Biosynthesis Is Regulated by *S* -Glutathionylation. *J. Biol. Chem.* **2002**, *277* (50), 48295–48302. https://doi.org/10.1074/jbc.M209042200.

(66) Ward, N. E.; Stewart, J. R.; Ioannides, C. G.; O'Brian, C. A. Oxidant-Induced *S* - Glutathiolation Inactivates Protein Kinase C-α (PKC-α): A Potential Mechanism of PKC Isozyme Regulation [†]. *Biochemistry* **2000**, *39* (33), 10319–10329. https://doi.org/10.1021/bi000781g.

(67) Lee, S. R.; Kwon, K. S.; Kim, S. R.; Rhee, S. G. Reversible Inactivation of Protein-Tyrosine Phosphatase 1B in A431 Cells Stimulated with Epidermal Growth Factor. *J. Biol. Chem.* **1998**, *273* (25), 15366–15372.

(68) William C. Barrett, ‡; Jon P. DeGnore, §; Simone König, §; Henry M. Fales, §; Yen-Fang Keng, □; Zhong-Yin Zhang, □; Moon B. Yim, ‡ and; P. Boon Chock*, ‡. Regulation of PTP1B via Glutathionylation of the Active Site Cysteine 215. **1999**. https://doi.org/10.1021/BI990240V.

(69) Reynaert, N. L.; van der Vliet, A.; Guala, A. S.; McGovern, T.; Hristova, M.; Pantano, C.; Heintz, N. H.; Heim, J.; Ho, Y.-S.; Matthews, D. E.; Wouters, E. F. M.; Janssen-Heininger, Y. M. W. Dynamic Redox Control of NF- B through Glutaredoxin-Regulated S-Glutathionylation of Inhibitory B Kinase Beta. *Proc. Natl. Acad. Sci.* **2006**, *103* (35),

13086–13091. https://doi.org/10.1073/pnas.0603290103.

(70)   Checconi, P.; Limongi, D.; Baldelli, S.; Ciriolo, M. R.; Nencioni, L.; Palamara, A. T. Role
        of   Glutathionylation   in   Infection   and   Inflammation.   *Nutrients*   **2019**.
        https://doi.org/10.3390/nu11081952.

(71)   Gallogly, M. M.; Mieyal, J. J. Mechanisms of Reversible Protein Glutathionylation in
        Redox Signaling and Oxidative Stress. *Curr. Opin. Pharmacol.* **2007**, *7* (4), 381–391.
        https://doi.org/10.1016/j.coph.2007.06.003.

(72)   Wen, J.; Li, X.; Zheng, S.; Xiao, Y. Upregulation of Glutaredoxin 2 Alleviates Oxygen-
        Glucose Deprivation/Reoxygenation-Induced Apoptosis and ROS Production in Neurons
        by Enhancing Nrf2 Signaling via Modulation of GSK-3β. *Brain Res.* **2020**, *1745*, 146946.
        https://doi.org/10.1016/j.brainres.2020.146946.

(73)   Collet, J.-F.; Messens, J. Structure, Function, and Mechanism of Thioredoxin Proteins.
        Antioxid. Redox Signal. **2010**, *13* (8), 1205–1216. https://doi.org/10.1089/ars.2010.3114.

(74)   Kasuno, K.; Nakamura, H.; Ono, T.; Muso, E.; Yodoi, J. Protective Roles of Thioredoxin,
        a Redox-Regulating Protein, in Renal Ischemia/Reperfusion Injury. *Kidney Int.* **2003**, *64*
        (4), 1273–1282. https://doi.org/10.1046/j.1523-1755.2003.00224.x.

(75)   Branco, V.; Pimentel, J.; Brito, M. A.; Carvalho, C. Thioredoxin, Glutathione and Related
        Molecules in Tumors of the Nervous System. *Curr. Med. Chem.* **2019**, *27* (12), 1878–
        1900. https://doi.org/10.2174/0929867326666190201113004.

(76)   Zhou, J.; Wang, C.; Wu, J.; Fukunaga, A.; Cheng, Z.; Wang, J.; Yamauchi, A.; Yodoi, J.;

Tian, H. Anti-Allergic and Anti-Inflammatory Effects and Molecular Mechanisms of Thioredoxin on Respiratory System Diseases. *Antioxid. Redox Signal.* **2020**, *32* (11), 785–801. https://doi.org/10.1089/ars.2019.7807.

(77)   Gilson Zeni, †; André Chieffi, †; Rodrigo L. O. R. Cunha, †; Julio Zukerman-Schpector, †; Hélio A. Stefani, ‡ and; João V. Comasseto*, †. Addition Reaction of P-Methoxyphenyltellurium Trichloride to 3-Hydroxy Alkynes. **1999**. https://doi.org/10.1021/OM980738D.

(78)   Cross, B. M.; Mcwilliam, A. A. *Guide To the Care and Use of Experimental*; 1993; Vol. 1.

(79)   Schneider, C. Research 1951. *Cancer Res.* **1951**, *11* (1), 23.

(80)   Layne, E. [73] Spectrophotometric and Turbidimetric Methods for Measuring Proteins. *Methods Enzymol.* **1957**, *3*, 447–454. https://doi.org/10.1016/S0076-6879(57)03413-8.

(81)   Noble, J. E.; Bailey, M. J. A. Chapter 8 Quantitation of Protein. *Methods Enzymol.* **2009**, *463*, 73–95. https://doi.org/10.1016/S0076-6879(09)63008-1.

(82)   Rück, A.; Dolder, M.; Wallimann, T.; Brdiczka, D. Reconstituted Adenine Nucleotide Translocase Forms a Channel for Small Molecules Comparable to the Mitochondrial Permeability Transition Pore. *FEBS Lett.* **1998**, *426* (1), 97–101. https://doi.org/10.1016/S0014-5793(98)00317-2.

(83)   Kawai, C.; Prado, F. M.; Nunes, G. L. C.; Di Mascio, P.; Carmona-Ribeiro, A. M.; Nantes, I. L. PH-Dependent Interaction of Cytochrome c with Mitochondrial Mimetic

Membranes: The Role of an Array of Positively Charged Amino Acids. *J. Biol. Chem.* **2005**, *280* (41), 34709–34717. https://doi.org/10.1074/jbc.M412532200.

(84) Rodrigues, T.; De França, L. P.; Kawai, C.; De Faria, P. A.; Mugnol, K. C. U.; Braga, F. M.; Tersariol, I. L. S.; Smaili, S. S.; Nantes, I. L. Protective Role of Mitochondrial Unsaturated Lipids on the Preservation of the Apoptotic Ability of Cytochrome c Exposed to Singlet Oxygen. *J. Biol. Chem.* **2007**, *282* (35), 25577–25587. https://doi.org/10.1074/jbc.M700009200.

(85) Castilho, R. F.; Kowaltowski, A. J.; Vercesi, A. E. The Irreversibility of Inner Mitochondrial Membrane Permeabilization by Ca2+ plus Prooxidants Is Determined by the Extent of Membrane Protein Thiol Cross-Linking. *J Bioenerg Biomembr* **1996**, *28* (6), 523–529.

(86) Pessoto, F. S.; Yokomizo, C. H.; Prieto, T.; Fernandes, C. S.; Silva, A. P.; Kaiser, C. R.; Basso, E. A.; Nantes, I. L. Thiosemicarbazone P-Substituted Acetophenone Derivatives Promote the Loss of Mitochondrial Δ ψ, GSH Depletion, and Death in K562 Cells. *Oxid. Med. Cell. Longev.* **2015**, *2015*. https://doi.org/10.1155/2015/394367.

(87) Schlegel, R. A.; Williamson, P. Phosphatidylserine, a Death Knell. *Cell Death Differ.* **2001**, *8* (6), 551–563. https://doi.org/10.1038/sj.cdd.4400817.

(88) Zachowski, A.; Henry, J.-P.; Devaux, P. F. Control of Transmembrane Lipid Asymmetry in Chromaffin Granules by an ATP-Dependent Protein. *Nature* **1989**, *340* (6228), 75–76. https://doi.org/10.1038/340075a0.

(89) Marí, M.; Morales, A.; Colell, A.; García-Ruiz, C.; Fernández-Checa, J. C. Mitochondrial

Glutathione, a Key Survival Antioxidant. *Antioxid. Redox Signal.* **2009**, *11* (11), 2685–2700. https://doi.org/10.1089/ARS.2009.2695.

(90)   Albeck, A.; Weitman, H.; Sredni, B.; Albeck, M. Tellurium Compounds: Selective Inhibition of Cysteine Proteases and Model Reaction with Thiols. *Inorg. Chem.* **1998**, *37* (8), 1704–1712. https://doi.org/10.1021/ic971456t.

(91)   Princival, C. R.; Archilha, M. V. L. R.; Dos Santos, A. A.; Franco, M. P.; Braga, A. A. C.; Rodrigues-Oliveira, A. F.; Correra, T. C.; Cunha, R. L. O. R.; Comasseto, J. V. Stability Study of Hypervalent Tellurium Compounds in Aqueous Solutions. *ACS Omega* **2017**, *2* (8), 4431–4439. https://doi.org/10.1021/acsomega.7b00628.

(92)   Silva, G. D.; Cunha, R. L. O. R.; Domingues, M.; Neto, C. Equilibrium Between Tri- and Tetra-Coordinate Chalcogenuranes Is Critical for Cysteine Protease Inhibition. **2020**, No. 4. https://doi.org/10.26434/CHEMRXIV.13148825.V1.