Diverse spatial representations in primary auditory cortex during active localization simultaneously code source location and identity

Diana Amaro[1,2], Dardo N. Ferreiro[1,3], Benedikt Grothe[1,2,4], Michael Pecka[1,+]

[1] Division of Neurobiology, Department Biology II, Ludwig-Maximilians-Universität München, Munich, Germany.

[2] Graduate School of Systemic Neurosciences, Ludwig-Maximilians-Universität München, Munich, Germany.

10 [3] Department of General Psychology and Education, Ludwig-Maximilians-Universität München, Munich, Germany.

[4] Max Planck Institute of Neurobiology, Martinsried, Germany

[+] pecka@bio.lmu.de

ABSTRACT:

Localizing and identifying sensory objects during active navigation are fundamental brain functions. However, how individual objects are neuronally represented during self-motion is mostly unexplored. Here we show that active localization during
20 unrestricted navigation promotes previously unreported spatial representations in primary auditory cortex. Spatial tuning differs between sources with distinct behavioral outcome associations, revealing a simultaneous population coding of egocentric source locations and angle-independent identification of individual sources during active sensing.

MAIN:

Information about the position of sensory objects is vital to navigate the environment. In the auditory system, neurons compute azimuthal spatial information via angle-specific differences of sound features between the ears. From brainstem to primary
30 auditory cortex (A1), a predominance of broad neuronal tuning to contralateral sound-source locations has been reported, with a smaller subset tuned to ipsilateral or frontal positions[1,2,3]. Accordingly, spatial tuning is assumed to be egocentric (i.e., based on the position of the sound-source relative to the observer). Nevertheless, most studies on spatial tuning to date were conducted in either anesthetized or head-
35 fixed and passively listening animals, thus lacking important aspects of real-life localization behavior that may crucially impact the nature of spatial coding.

Movements and active exploration are fundamental components of natural sensing[4]. Notably, while self-movement constantly alters the egocentric sound-source location, the perception of source position remains stable relative to the world coordinates, i.e.

40 is allocentric[5]. Recent results from one study with freely moving ferrets suggest the existence of allocentric representation components in a minority of neurons already in A1[6]. However, the stimuli and associated sources in this study were task-irrelevant since subjects were merely exposed to sounds while searching for water. Yet active sensing and task engagement critically modulate neuronal coding in A1[7,8,9,10,11], and

45 thus it remains unclear how sound-sources are represented during unrestricted movement and active localization.

We hypothesized that training freely exploring animals to actively localize sounds from sources with distinct task identities (i.e., associated behavioral outcome, Fig. 1a) that only differ in their allocentric locations would promote revealing potentially

50 undiscovered spatial representations. We took advantage of the recently developed Sensory Island Task (SIT) paradigm[12] and trained Mongolian gerbils (*Meriones unguiculatus*) to report whenever the presentation of a pulsed harmonic stack switched from a "background" loudspeaker to a "target" loudspeaker (separated by 180° in a circular arena, Fig. 1a). The source change was triggered whenever gerbils

55 entered a specific target area in the arena (the "island"), whose position was randomized across trials. The animals were trained to report the detection of the change by remaining within the island for 6 seconds to receive a food reward. Thus, activity of the target loudspeaker was associated with imminent reward delivery. As the sound presented from both loudspeakers was identical (Extended Data Fig. 1),

60 task completion was achievable only by determining the location of the active sound-source. Importantly, because the egocentric sound-source location can be the same for both loudspeakers, this information was rendered insufficient for task completion (Fig. 1a and Suppl. Video 1). Since the animals reliably exhibited highly significant performance levels in reporting the activity of the target loudspeaker (Fig. 1b,c), their

65 behavior demonstrates allocentric localization (e.g. "loudspeaker to the west of the initiation platform"). Note that switching the rewarded loudspeaker in catch-trials resulted in performances significantly below chance level (Fig. 1c, left column), corroborating a task-specific identification of the loudspeakers by the animals. This allowed studying how listening to two behaviorally meaningful sound-sources that are

70 defined by their world-based position influences the spatial representation in A1.

To investigate the neuronal processing during task performance, expert animals were implanted with custom-made tetrodes to allow chronic recordings of action potentials from neurons in A1 (Fig. 1d). Overall, we acquired responses during task performance from 364 single neurons and 246 multi-units from 5 gerbils. Activity in

75 A1 was strongly correlated with the subjects' goal-specific behavior, as response rates to the target loudspeaker were significantly lower when animals correctly finished the trial as opposed to when they wrongly left the island (Fig. 1e and Extended Data Fig. 2).

Neuronal firing rates exhibited significant modulation (see below and Methods) as a

80 function of the active loudspeaker's angle relative to the animal's body-axis (Fig. 1f see Methods) at specific time periods relative to sound presentation (Fig. 1g, "onset period": from the determined response latency + stimulus duration of 57 ms, 118 neurons; "offset period": end of onset period + 57 ms, 58 neurons; "late response period": remaining time between the periodic stimuli, 42 neurons, see Methods).

85 Construction of polar plots from these responses displayed a large variety of

egocentric spatial sensitivity (Fig. 1h). Alongside a fraction of "canonical" neurons that were similarly tuned to either loudspeaker with contralateral or ipsilateral preference ("i" and "iii" in Fig. 1h; onset period: 37/118 = 31.3 %; offset period: 7/58; late response period: 0/42), we observed neurons whose tuning to either loudspeaker can be described as orientation-sensitive (onset period: 16/118 = 13.6 %; offset period: 1/58; late period: 2/42), with an apparent bias to the front/back orientation ("ii" in Fig. 1h, comparable to reports in awake primates[14]).

Remarkably, the spatial tuning of a large fraction of the neurons was source-sensitive, inclusively during the onset period (65/118 = 55.1 %), as their responses differed for the two loudspeakers (Figs. 1h & 2a, and Extended Data Fig. 3). These neurons were either only significantly spatially tuned to one of the two sound-sources ("iv" in Figs. 1h & 2a; 47/65 = 72.3 %) or exhibited a large tuning difference between the two sources (difference in preferred egocentric angle > 90°, "v" in Figs. 1h & 2a; 18/65 = 27.7 %, see Methods). This variety in tuning types was evident during all three response epochs and also on the level of multi-unit responses (Extended Data Fig. 3c). Quantification of spatial selectivity exposed population-wide source-specific tuning, as vector strength (see Methods) was significantly higher for the target loudspeaker (Fig. 2b). While we did not notice a dependency of tuning preferences on the putative neuronal type (regular or fast spiking, Fig. 2a, Extended data Fig. 4), putative regular spiking neurons exhibited sharper tuning to either sound-source (Extended Data Fig. 4f,g). Notably, in a sizable fraction of neurons that classified as spatially un-tuned to either loudspeaker, the response magnitude differed significantly between the two loudspeakers (16/40, 40 %, Fig. 2a, "vi" in Fig. 1h), indicative of "purely" allocentric, i.e. sound-source angle independent coding.

Response timing relative to stimulus duration is known to provide additional information about sound-source location[3,15]. We found that the spatial representations in A1 of the actively localizing gerbils also depended on the relative timing of responses, as the egocentric tuning of individual neurons frequently varied considerably across response epochs (Fig. 2c,d). Additionally, often this change of preferred egocentric location across response time occurred only for one loudspeaker.

Together, our data suggests that active localization during self-motion in SIT reveals a variety of previously unreported characteristics of spatial sensitivity. However, movement can modulate cortical activity unspecifically[16,17]. We therefore performed additional control analyses with data restricted to neuronal responses from periods when the animals were not moving. This resulted in qualitatively similar observations (see Extended Data Fig. 5), demonstrating that our findings are not caused by unspecific movement modulation.

To test to what extent the diverse activity patterns that we observed may aid task performance, we implemented an artificial neural network model with one hidden layer (multilayer perceptron classifier, see Methods). The algorithm was trained on 224 units (141 single neurons and 83 multi-units) from two animals from a total of 21 sessions and the training classes were a combination of egocentric spatial information and allocentric sound-source identity (using random under-sampling to prevent class imbalance), in which the 360° around the animal were divided in 8

classes for each loudspeaker. The neural network was then presented with test data (see Methods) for decoding of specific spatial information. Initially, the data from both loudspeakers was merged to calculate the accuracy of the decoder on the egocentric location of the active loudspeaker. The neural network model classified the

135 egocentric locations with high accuracy (Fig. 3a). Next, we determined the capability of the algorithm in identifying the active loudspeaker irrespective of the egocentric location. Remarkably, the decoding accuracy for identifying the sound-source was also highly significant (Fig. 3b), demonstrating the coexistence of both subject-based and world-based reference frames (identity) in the neuronal representations (Fig. 3c).

140 Temporal analysis of model performance indicates that the egocentric position of the active loudspeaker was largely encoded within the first 100 ms after sound onset, whereas the information about the identity of the loudspeaker increased monotonically throughout the stimulation period (Fig. 3d) and reached behavioral performance levels (Fig. 1c).

145 To explore to what extent the diverse response patterns across the population of neurons contribute to the egocentric and identity decoding accuracy, we selectively eliminated units with specific tuning characteristics in the test data and re-assessed the decoding accuracy (Fig. 3e). Elimination of either all units with differential spatial tuning for the two sound-sources (18 units, compare Fig. 1h, type "v") or exclusion of

150 the same number of units with either diverse response profiles (randomly selected) or exclusively canonic hemispherically tuned units (compare Fig. 1h, type "i" and "iii") resulted in only mild reductions of accuracy for location and identity, respectively. Overall, the neural network algorithm demonstrated remarkable robustness towards these interventions, as decoding accuracy for both reference frames remained far

155 above chance level regardless of the eliminated neuronal tuning class (Extended Data Fig. 6). This robustness suggests that distributed interactions in population responses underlie an interlaced coding of egocentric and allocentric information.

Together, our data reveal spatial representations in A1 that were so far unreported. By introducing differential behavioral meaning to two loudspeakers at distinct

160 locations, we show that the spatial tuning of A1 neurons exceeds the canonical hemispheric preference and exhibits temporal complexity. Maybe most intriguingly, we observed differential spatial tuning to individual loudspeakers, especially at later response periods, which may result from top-down modulation[5,18]. These findings strengthen the hypotheses for the existence of identity coding[19] and allocentric

165 representations at an early cortical level[6], and further demonstrate an intricate coexistence with egocentric representations on the population level. More generally, our findings may contribute to identifying and tracking of a specific sound-source in the environment during self-motion and active sensing and thus may also help to better understand the cocktail party phenomenon[20].

Fig. 1: Sound localization task and non-canonical spatial tuning. a, General scheme of the task: the animal starts a trial in the initiation platform and hears a pulsed harmonic complex from the background loudspeaker until it enters the island, which triggers a switch to the target loudspeaker; to report the switch the animal must stay within the island for 6 seconds. Egocentric sound-source angles are ambiguous and hence insufficient for task completion. Pairing of the target loudspeaker with reward delivery allows the formation of source-specific identity associations **b**, Task performance of one animal in comparison to chance during a particular session as a function of time spent in the trial (40 trials, shaded areas correspond to 95% confidence interval). **c**, Left panel: task performance at maximum trial length (60s) in comparison to chance (11 animals, Wilcoxon signed-rank test, horizontal lines represent the median, previously reported[12]). Right panel: task performance at maximum trial length (60s) in comparison to chance in catch-trials with reversal of the reward contingency (5 animals, Wilcoxon signed-rank test, horizontal lines represent the median, previously reported[12]) **d**, Reconstruction of tetrode position from histological information, superimposed with the location of A1[13]. **e**, Left: Firing rates were significantly lower in the last 2s when the animal finished a trial correctly compared to when it wrongly abandons the island (321 neurons, Wilcoxon signed-rank test, mean firing rate). Right: This decrease in firing rate was enhanced with more time spent in the island (Mann-Whitney U-test, first 2s: 346 neurons, last 2s: 321 neurons). Violin plots surrounding boxplots (the median is represented by a white

dot) depict the distribution of the ratios between the mean firing rate when the animal correctly finished a trial and when it wrongly left the island. **f**, Trajectory of the animal during a successful trial. Colored dots indicate the position of the animal at the moment of sound presentation and corresponding egocentric locations of the active loudspeaker. The color code is the same as in (g). **g**, Raster plots of neuronal activity in two A1 neurons during task performance in the session represented in (b). Periods of stimulation by the target loudspeaker are highlighted by gray areas. Egocentric location of the active loudspeaker at the moment of the sound occurrence is color-coded. **h**, Spatial tuning for six representative neurons for the target and background loudspeakers. The shaded area corresponds to the 95% confidence interval and the colored lines indicate the preferred tuning angle, with the length scaling with the vector strength. Neurons i and ii correspond to the ones depicted in (g).

Fig. 2: Complexity of neuronal responses in a context-dependent manner. a, Correspondence of preferred egocentric locations between the target and the background loudspeakers. Color-code depicts putative neuronal type; FS: fast-spiking, RS: regular spiking (118 neurons, 71 of which tuned for both loudspeakers). Inset in lower left corner shows percentage of neurons that were not significantly (n.s.) tuned to either loudspeaker, but exhibited a significant difference in overall response magnitude (16/40 neurons, 5 FS, 11 RS). **b,** Vector strength was larger for the target than the background loudspeaker (Wilcoxon signed-rank test). Yellow data points represent neurons spatially tuned to both loudspeakers (71 neurons) and data from neurons that were not significantly spatially modulated for at least one of the loudspeakers are shown in gray (133 neurons). **c,** Peristimulus time histograms (PSTH) for two representative neurons with temporally complex spatial tunings. The

220 shaded area corresponds to the 95% confidence interval, the red line represents the sound stimulation period and the vertical lines indicate the time period considered for the construction of the spatial tuning (i.e. onset, offset and late response periods). The violet/green PSTHs correspond to responses acquired only for contra- or ipsilateral loudspeaker locations (in a 90° angular bin). **d**, Spatial tuning comparison

225 between the onset and the offset time periods (upper panel, 23 neurons) and between the onset and the late response (lower panel, 21 neurons). The dark blue lines connect background and target spatial tuning angles that correspond to the same neuron.

Fig. 3: Artificial neural network decoding of population activity. Confusion matrix for decoding **a,** the egocentric loudspeaker location; **b,** the identity of the active loudspeaker; **c,** the combination of the loudspeaker location and identity. F: front, L: left, B: back, R: right. For the construction of all the confusion matrices, 38980 values were used across 20 sampling cycles (1949 values per cycle). **d**, Relationship between the accuracy of the decoder and the time after sound onset used for decoding. Inset: Accuracy of decoding the egocentric location when using exactly one 25 ms bin. The normalized accuracy drops linearly from over 30% until 50 ms to almost chance level after 125ms after sound onset. All error bars correspond to the standard deviation of the mean normalized accuracy across 20 sampling cycles. **e**, Eliminating 18 units of different tuning types from the test dataset resulted in a minor drop in the decoding accuracy and remained highly significant above chance (differentially tuned: -3.6±2.2 % and -4.2±1.8 % for location and identity, respectively; randomly selected, -1.4±1.8 % and -1.2±1.7 %, hemispherically tuned: -4.8±2.4 % and -2.9±2.4 %). All error bars correspond to the standard deviation of the mean accuracy across 100 sampling cycles.

Methods

Animals

All procedures were approved in accordance with the stipulations of the German animal welfare law (AZ 55.2-1-54-2532-74-2016). Experiments were conducted on 11 male Mongolian gerbils (*Meriones unguiculatus*) from the breeding colony of the Biocenter of the Ludwig-Maximilians-Universität München. Animals were housed in groups of 3 to 4 individuals at a temperature of 22.4∘C and 66% humidity with 12 h light/dark cycles. The experiments were conducted during the light phase of the cycle.

Behavioral conditioning and stimulus control

Behavioral data was collected from eleven gerbils. Animals were required to be at least 8 weeks old at the beginning of training and underwent a general habituation period in the SIT setup for 15 minutes per day for 5 days. Gerbils had unrestricted access to water in their home cages. Food was provided as pellets ad libitum until the training phase started, after which animals were only allowed food as rewards for correct trials (half of a sunflower seed or 20 mg, TestDiet LabTab AIN-76A). The weight of every gerbil was measured daily and additional food was provided if needed to ensure the weight not to drop more than 5% between consecutive training days and to maintain it within the desired range of 60–80 g.

Animals were trained using the SIT paradigm[12] for probing sensory perception in unrestrained and actively engaged animals. In SIT, a freely moving animal was trained to search for an unknown area in the arena (target island), which prompted a change in the identity of the loudspeaker (from "background" to "target") from which a stimulation is played back, and to report the detection of the change by remaining in the target island for a determined time interval ("sit-time" = 6 s). The position of the target island was random and changed every trial (pseudo-randomly chosen from a uniform distribution), making the loudspeaker change the only useful cue to find the island. After correctly reporting the detection of the target stimulus, the animal was rewarded with food that automatically dropped in the arena from an overhead food dispenser. Trials had a time limit of 60 s. If the animal did not correctly report the target island within the time limit, a low-pass filtered noise was presented to the animal for 10 s, during which no new trial could be initiated.

Details of the setup and stimulation control were reported elsewhere[12] and are summarized here: The setup consisted of a circular arena (diameter = 92 cm) within a sound attenuated chamber (Fig. 1). Stimuli were computer generated and transmitted through an amplifier (AVR 445 Harman/Kardon, Germany) and small loudspeakers (Aurasound NSW1-205-8A 1" Extended Range) mounted externally of the arena (~5 cm distance to the perforated metal walls of the arena). Stimuli were 57 ms long harmonic complex sounds with a fundamental frequency of 147±4 Hz and low-pass filtered below 1.5 kHz. Animals were trained to initiate a trial by remaining on an initiation platform (approx. 1cm in height, 12 cm diameter) for 1 s. Trial initiation triggered the playback from the background loudspeaker (located either east or west of the initiation platform), and animal entrance into the island triggered the switch of the playback to the target loudspeaker (separated by 180° from the background

loudspeaker). Stimuli were played at a repetition rate of 4 Hz and their amplitude was 70 dB SPL roved ±5 dB. The animal's position was tracked via images captured every 250 ms with a Flea3 camera (FL3-U3-13Y3M-C, Point Grey Research Inc.). Stimulation parameters (loudspeaker identity) were updated online according to the animals' position within the arena. Custom-made software for animal tracking, stimuli generation and food reward delivery was developed in MATLAB.

No more than two training sessions were carried out per day, lasting up to 90 minutes. The training of the animals was performed by gradually reducing island size (starting at diameter = 42 cm, ~21% of the arena surface) and increasing sit-time (starting with 2 s) over the course of the training sessions. The final parameters of the island size were diameter = 25 cm (~7% of the arena surface) and sit-time = 6 s.

Video tracking

Custom-made online tracking scripts were developed in MATLAB. In the animals' tracking algorithm, an ellipse was fitted to the animal after background subtraction and the centroid of this ellipse was taken as the position of the gerbil. The recorded videos were offline analyzed in Python (with OpenCV library[21]) where the ellipse and its centroid position was re-analyzed and the orientation of the fitted ellipse determined. The median orientation error was assessed blindly to be 6°, with 75% of the orientation errors being below 15°.

Behavioral data analysis

Data analyses were performed in MATLAB (Mathworks) and Python using custom scripts. To test the performance of the animals, we compared the percentage of correct trials in each session with surrogate runs based on random target island shuffling. That is, for each trial (offline, a posteriori), 1000 surrogate (non-real) islands, non-overlapping with the target one, were randomly set and the real trajectory of the animal was used to calculate in how many of these islands the trial would have been correct at each time point given the required sit-time. The median chance performance and 95% confidence interval was calculated based on bootstrapping (random sampling with replacement from all the trials of the session). This method allows obtaining an estimate of the proportion of correct trials the animal would have gotten just by chance given their locomotion trajectory and dynamics.

Surgery

Electrophysiological data was collected from five trained gerbils. Animals were anesthetized with an intraperitoneal injection of a mixture of metedomidin (0.15 mg/kg), midazolam (7.5 mg/kg), and fentanyl (0.03 mg/kg). To maintain it at a constant level, the same mixture was subcutaneously re-injected every 90 min. After shaving and disinfecting the head, a local anesthetic (50 µl, 2% xylocaine) was injected under the scalp skin and below the skin near the ears. For protection and to prevent dehydration, the eyes were covered with an ophthalmic gel (Thilo-Tears SE, Alcon Pharma Gmbh). The animal was then transferred to the stereotactic apparatus, where its head was securely fixed via a bite and ear bars. Its internal temperature was monitored with a rectal thermometer and kept constant at 37°C throughout the experiment by a feedback controlled electric heating pad (Harvard Apparatus). After disinfection, a midline scalp incision was performed to expose the skull.

Subsequently, the connective tissue on the skull was removed with a bone curette and the skull was treated with 35% phosphoric acid (iBOND etch gel, Kulzer), which was promptly washed away. Structural screws were placed on top of the left frontal and right parietal bones and the ground screw on the occipital bone, so that it gently touched the brain. After stereotactic alignment, a 3x3 mm craniotomy and durotomy were performed on top of the left auditory cortex, followed by a very slow lowering (2 µm/s) of a tetrode bundle to a maximum depth of 0.9 mm into the cortex, using a micromanipulator (Scientifica). The craniotomy was carefully filled with KY-jelly and immediately sealed with dental cement (Paladur, Kulzer), which also fixated the bottom of the microdrive and the outer cannula that protected the tetrodes. 1 ml of Ringer's solution was subcutaneously injected at the end of the surgery and the anesthesia was reversed via subcutaneous injection of the antagonist mixture composed of naloxone (0.5 mg/kg), flumazenil (0.4 mg/kg), and atipamezol (0.375 mg/kg). Analgesics (0.2 mg/kg, meloxicam) and antibiotics (7.5 mg/kg, enrofloxacin) were orally administered post surgically for five subsequent recovery days. During this time, the animals had food and water ad libitum and were not trained.

The implant used in this experiment was a tetrode bundle consisting of four tetrodes glued together, which, on their turn, consisted of four insulated tungsten wires (12.7 µm diameter each, tungsten 99.95%, California Fine Wire) twisted around each other. Each wire was connected to a custom-made printed circuit board with Omnetics connector (Axona), which was attached to a lightweight microdrive (0.25 mm/turn, Axona). The tetrodes were glued together and protected by an inner and outer cannula that could slide by each other. On the day prior to the surgery, the tip of all electrodes were cut with sharp scissors and gold plated (Non-Cyanide Gold Plating Solution, Neuralynx) to reach a desired impedance of 100-150 kOhm (at 1 kHz). The tetrode bundle was implanted vertically in the following coordinates from lambda: 6.2 mm lateral, 2.6 mm anterior.

Electrophysiological recordings during task performance

Recordings were made via a wireless headstage (W2100-HS16, Multichannel Systems). The physiological signals were at first amplified between 1 Hz and 5 kHz and digitized (16-bit resolution) in the headstage, then wirelessly transmitted to the receiver (W2100-RE-AO, Multichannel Systems) at a sampling rate of 25 kHz and recorded by a PC via an interface board (MCS-IFB 3.0 Multi-boot, Multichannel Systems), and commercial software (Multi Channel Experimenter, Multichannel Systems). Simultaneous with the onset of sound presentation, a short signal was sent via sound card to the analog input of the interface board, allowing the synchronization of the physiological recordings with the sound presentation, and a digital signal was transmitted to the interface board indicating the beginning and end of the trial, which was later used to align the video information.

Spike sorting and neurophysiological data analysis

Initially, the raw electrophysiological signals were high-pass filtered above 300Hz and a common median referencing[22] was performed for the whitening among all channels and removal of large artifacts. The signals were low-pass filtered below 5 kHz and fed to a spike sorting algorithm based on template matching (Kilosort[23]). Afterwards, the automatically sorted spikes were manually inspected and the corresponding

clusters refined with the graphical user interface phy[24]. Only units with an isolation distance larger than 20, more than 200 spikes and less than 2% of the spikes within the 2 ms refractory period were considered single cells[25,26]. All units that passed the manual curation but not the isolation distance or refractory period requirements were considered multiunits.

The peristimulus time histogram (PSTH) was calculated based on bootstrapping data. An algorithm sampled 500 times (with replacement) from the total number of recordings of a given sound. For each cycle, a histogram was calculated with binning width 8 ms and the median and 95% confidence intervals were determined for the total of bootstrapping cycles normalized by the total number of analyzed sounds. For the calculation of neuronal latencies without contamination by baseline activity, the Bayesian blocks method[27] was used, which identifies statistically significant variation in a time series to optimally segment the data. If at least one new edge is created, the unit is considered modulated and therefore auditory responsive. The latency of a given unit was defined as the first created edge. To determine whether the onset response corresponds to an increase in the firing rate, the maximum displacement of the median values of the firing rate in the 15 ms after latency was compared to the baseline (last 50 ms before latency). If it was lower than the mean of the lower bound of the baseline's 95% confidence interval, the onset firing rate was considered to be decreasing and if it was higher than the mean of the upper bound of the baseline's 95% confidence interval, the onset firing rate was considered to be increasing. The same Bayesian blocks method was employed to calculate whether a unit had a significant offset response within the interval [latency+50:latency+70] ms. If so, the offset period was defined as the time point of the new edge creation, otherwise it was defined as latency + sound duration (57 ms). The start of the late response period was defined as the start of the offset period + sound duration (57 ms).

Mean firing rates were calculated as the mean of the PSTH across all time bins.

For the comparison between the first and last 2 s inside the island and whenever the animal wrongly left it, only situations in which the animal stayed at least 2 s in the island were considered. Furthermore, if the last 2 s overlapped in time with the first 2 s (the animal stayed less than 4 s in the island), only the non-overlapping part was used for the calculation of the last 2 s.

For the spatial tuning analysis, the 360° angular space around the animal was divided in 8 bins. A minimum of 10 sound presentations per binned angle was a primary condition for the spatial tuning analysis, as well as a maximum difference of 10 between the bins with the most and the least number of sound presentations. A 1000 cycle bootstrapping method with replacement was implemented to calculate the spatial tuning, in which an angular histogram was calculated for the corresponding spikes and normalized by the number of sounds at each egocentric sound-source angle bin for the chosen loudspeaker. The vector strength and corresponding preferred egocentric sound-source angle were then calculated for each bootstrapping cycle. As some units revealed a more complex spatial tuning (with two peaks in opposite directions), the folded vector strength[28] (in which responses to angles opposite to each other are summed) was additionally calculated together with the corresponding orientation angle.

$$\rho = \frac{\sum_{j=1}^{8} \alpha_j . e^{i\omega_j}}{\sum_{j=1}^{8} \alpha_j}$$

where $i = \sqrt{-1}$, $\alpha_j = \frac{\#spikes\ in\ angular\ bin_j}{\#sound\ presentations\ in\ angular\ bin_j}$ and $\omega_j = \frac{2\pi.middle\ of\ angular\ bin_j}{T}$.

430   The vector strength is $VS = |\rho|$ with T=360 and the corresponding preferred egocentric sound-source angle $ang_{dir} = \arg(\rho)$. The folded vector strength is $VS_{folded} = |\rho|$ with T=180 and the corresponding preferred egocentric sound-source angle angle $ang_{ori} = \arg(\rho)/2$. From the bootstrapped data, the 95% confidence interval for each angle bin was calculated, as well as the 68% confidence interval of

435   the vector strength and orientation vector. A unit was considered spatially tuned at a particular temporal period (onset, offset or late) if the vector strength (or folded vector strength, whichever is larger) was larger than 0.2 and the lower bound of 68% confidence interval of the vector strength was larger than 0.15.

Units were considered as un-tuned if the median vector strength was smaller than

440   0.15 in all three phases of the PSTH for both loudspeakers. The firing rate in response to the two loudspeakers was considered significantly different if there was no overlap in the 95% confidence interval of the calculation of the mean firing rate for each loudspeaker.

For the decoder analysis, the decoder was trained on the recorded responses of 224

445   units (141 single neurons and 83 multi-units) from two animals in a total of 21 sessions. Each sound presentation was categorized with respect to the active loudspeaker and the egocentric sound-source angle at the time of sound onset (in 8 angular bins). Therefore, each sound presentation corresponded to one of the 16 possible loudspeaker-angular bin classes. Spike counts, for each recorded unit, in

450   each of the 250 ms response periods were determined in 10 bins, each with a 25 ms duration, and then normalized between 0 and 1 (using the mix-max scaler algorithm) to avoid that units that are inherently more active dominate the results. These unit responses were then pooled across sessions for each of the angular-bin classes, creating 16 across-sessions class-specific population-response profile, which were

455   then used to train the decoder. To prevent class imbalance, due to non-uniform occurrence of angular-bin classes during the task, random under-sampling (without replacement) was implemented to feed the decoder with the same amount of data per class. The test dataset consisted of a minimum of 25% of the population responses per class.

460   The training data was fitted by a multi-layer perceptron classifier[29] (MLPC), a feedforward artificial neural network which uses backpropagation during training for optimization of the parameters in a supervised learning manner, and therefore was interpreted as a more biologically inspired decoder. In the MLPC, one hidden layer was implemented with the number of nodes as the mean between the number of

465   features used and the total number of classes (16). The optimization algorithm used was the "lbfgs" from the family of quasi-Newton methods and the biologically-inspired rectified linear unit was the activation function for the hidden layer.

The whole process was repeated 20 times to estimate errors (100 times in the case of the unit-elimination analysis). The accuracy of identifying the active loudspeaker

was determined per loudspeaker for each sampling repetition, later the accuracy per repetition was considered as the mean between the two loudspeakers and the total accuracy and standard deviation calculated across all 20 sampling turns. The accuracy of predicting the right class and egocentric sound-source location was similarly calculated. For the construction of the confusion matrices, all the predictions from all the sampling repetitions were simultaneously used.

To evaluate the influence of units with different spatial tuning, units were eliminated in the test data by setting all the bins corresponding to the chosen units to zero. The accuracy of the decoder (trained on the training data from all the units) was then compared to when the test data was complete. In the elimination of the random units, different units were chosen to be eliminated in each sampling turn. The units with differential spatial tuning for both loudspeakers which were eliminated corresponded to all the units in the training set which at some temporal response period had a difference in preferred egocentric sound-source angle between loudspeakers larger than 80°. The eliminated canonical units were randomly selected from the neurons which showed clear ipsi or contralateral tuning during the onset period.

Histology

After a 350 μl intraperitoneal injection of pentobarbital sodium, the animal was perfused with 4% paraformaldehyde (PFA) and the brain carefully removed and stored in PFA in the fridge. Afterwards the brain was twice washed in PBS and the frontal part cut with a vibrotome in 70μm thick slices. The slices were stained with green fluorescent Nissl (NeuroTrace™ 500/525) and then compared to the gerbil brain atlas[13] for determining the location of the recording sites.

Statistics and reproducibility

All error bars correspond to the standard deviation unless stated otherwise. All shaded areas correspond to the 95% confidence interval as calculated via bootstrapping. For comparisons of central tendencies on the group level, we used two-tailed non-parametric tests: Wilcoxon signed-rank tests for paired samples and Mann-Whitney U-test for independent samples. All hypotheses were tested at an alpha level of 0.05.

Data collection and analysis were not performed blind to the conditions of the experiments. Our experimental design provided a within-animal control, and because comparisons were not required between different groups, blinding was not necessary. All analyses were based on automated scripts applied across animals and thus were not subject to any experimenter bias.

References

1.Benson, D. A., Hienz, R. D. & Goldstein, M. H. Single-unit activity in the auditory cortex of monkeys actively localizing sound sources: Spatial tuning and behavioral dependency. *Brain Res.* **219**, 249–267 (1981).

2.Grothe, B., Pecka, M. & McAlpine, D. Mechanisms of Sound Localization in Mammals. *Physiol. Rev.* **90**, 983–1012 (2010).

3.Mickey, B. J. & Middlebrooks, J. C. Representation of Auditory Space by Cortical Neurons in Awake Cats. *J. Neurosci.* **23**, 8649–8663 (2003).

4.Schroeder, C. E., Wilson, D. A., Radman, T., Scharfman, H. & Lakatos, P. Dynamics of Active Sensing and perceptual selection. *Curr. Opin. Neurobiol.* **20**, 172–176 (2010).

5.Burgess, N. Spatial Cognition and the Brain. *Ann. N. Y. Acad. Sci.* **1124**, 77–97 (2008).

6.Town, S. M., Brimijoin, W. O. & Bizley, J. K. Egocentric and allocentric representations in auditory cortex. *PLOS Biol.***15**, e2001878 (2017).

7.Fritz, J., Shamma, S., Elhilali, M. & Klein, D. Rapid task-related plasticity of spectrotemporal receptive fields in primary auditory cortex. *Nat. Neurosci.* **6**, 1216–1223 (2003).

8.David, S. V., Fritz, J. B. & Shamma, S. A. Task reward structure shapes rapid receptive field plasticity in auditory cortex. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 2144–2149 (2012).

9.Lin, P.-A., Asinof, S. K., Edwards, N. J. & Isaacson, J. S. Arousal regulates frequency tuning in primary auditory cortex. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 25304–25310 (2019).

10.von Trapp, G., Buran, B. N., Sen, K., Semple, M. N. & Sanes, D. H. A Decline in Response Variability Improves Neural Signal Detection during Auditory Task Performance. *J. Neurosci.* **36**, 11097–11106 (2016).

11.Lee, C.-C. & Middlebrooks, J. C. Auditory cortex spatial sensitivity sharpens during task performance. *Nat. Neurosci.* **14**, 108–114 (2011).

12.Ferreiro, D. N. et al. Sensory Island Task (SIT): A New Behavioral Paradigm to Study Sensory Perception and Neural Processing in Freely Moving Animals. *Front. Behav. Neurosci.* **14**, 576154 (2020).

13.Radtke-Schuller, S. et al. Brain atlas of the Mongolian gerbil (*Meriones unguiculatus*) in CT/MRI-aided stereotaxic coordinates. *Brain Struct. Funct.* **221**, 1–272 (2016).

14.Remington, E. D. & Wang, X. Neural Representations of the Full Spatial Field in Auditory Cortex of Awake Marmoset (*Callithrix jacchus*). *Cereb. Cortex* **29**, 1199–1216 (2019).

15. Middlebrooks, J., Clock, A., Xu, L. & Green, D. A panoramic code for sound location by cortical neurons. *Science* **264**, 842–844 (1994).

16. Zhou, M. et al. Scaling down of balanced excitation and inhibition by active behavioral states in auditory cortex. *Nat. Neurosci.* **17**, 841–850 (2014).

17. Schneider, D. M., Nelson, A. & Mooney, R. A synaptic and circuit basis for corollary discharge in the auditory cortex. *Nature* **513**, 189–194 (2014).

18. Yin, P., Strait, D. L., Radtke-Schuller, S., Fritz, J. B. & Shamma, S. A. Dynamics and Hierarchical Encoding of Non-compact Acoustic Categories in Auditory and Frontal Cortex. *Curr. Biol.* **30**, 1649-1663.e5 (2020).

19. Town, S. M., Wood, K. C. & Bizley, J. K. Sound identity is represented robustly in auditory cortex during perceptual constancy. *Nat. Commun.* **9**, 4786 (2018).

20. Bizley, J. K. & Cohen, Y. E. The what, where and how of auditory-object perception. *Nat. Rev. Neurosci.* **14**, 693–707 (2013).

21. Bradski, G. The OpenCV Library. *Dr. Dobb's Journal of Software Tools* (2000).

22. Rolston, J. D., Gross, R. E. & Potter, S. M. Common median referencing for improved action potential detection with multielectrode arrays. in *2009 Annual International Conference of the IEEE Engineering in Medicine and Biology Society* 1604–1607 (IEEE, 2009). doi:10.1109/IEMBS.2009.5333230.

23. Pachitariu, M., Steinmetz, N., Kadir, S., Carandini, M. & Kenneth D., H. Kilosort: realtime spike-sorting for extracellular electrophysiology with hundreds of channels. Preprint at: http://biorxiv.org/lookup/doi/10.1101/061481 (2016).

24. Rossant, C. et al. Spike sorting for large, dense electrode arrays. *Nat. Neurosci*. **19**, 634–641 (2016).

25. Schmitzer-Torbert, N., Jackson, J., Henze, D., Harris, K. & Redish, A. D. Quantitative measures of cluster quality for use in extracellular recordings. *Neuroscience* **131**, 1–11 (2005).

26. Hill, D. N., Mehta, S. B. & Kleinfeld, D. Quality Metrics to Accompany Spike Sorting of Extracellular Signals. *J. Neurosci.* **31**, 8699–8705 (2011).

27. Scargle, J. D., Norris, J. P., Jackson, B. & Chiang, J. Studies in Astronomical Time Series Analysis. VI. Bayesian Block Representations. *Astrophys. J.* **764**, 167 (2013).

28. Mazurek, M., Kager, M. & Van Hooser, S. D. Robust quantification of orientation selectivity and direction selectivity. *Front. Neural Circuits* **8**, (2014).

29. Pedregosa, F. et al. Scikit-learn: Machine Learning in Python. Journal of Machine Learning Research 12, 2825–2830 (2011).

Acknowledgements

This study was supported by the Deutsche Forschungsgemeinschaft DFG (PE2251/2-1 to MP, and SFB 870, project B02 to MP and BG), and by the International Max Planck Research School for Molecular Life Sciences (to DA). We thank N. Lesica and P. Alcami for valuable comments on earlier versions of the manuscript. We are grateful to L. Wiegrebe for help on the behavioral paradigm.

Author Contributions

MP, DA, and BG conceived the study. MP and DA designed the experiments and DNF and BG contributed to paradigm refinement. DA performed the experiments and analyzed the data. DA, MP, and DNF designed the data presentation and wrote the manuscript. All authors provided comments and approved the manuscript.

Competing interests

The authors declare no competing interests.

Extended Data

Extended Data Fig. 1: No influence of swapping loudspeakers on task performance. The performance of three animals in the task before and after
625 physically swapping the loudspeakers (but not their task-specific identity) is similar, showing that animals were not using specific spectral cues from the loudspeakers to solve the task. The data is presented as median performance relative to chance at maximum trial length (60s). The error bars correspond to the 95% confidence interval. Number of trials before swapping: $N_{Gerbil\ 1}$=206 $N_{Gerbil\ 2}$=163 $N_{Gerbil\ 3}$=177
630 After swapping: $N_{Gerbil\ 1}$=237 $N_{Gerbil\ 2}$=234 $N_{Gerbil\ 3}$=172.

Extended Data Fig. 2: Changes in firing rate are specific to trial-outcome. The decrease in mean firing rate when the animal finishes the trial is not due to **a,** unspecific neuronal rate adaptation, as no firing rate decrease is observed in the initial 6 s of background stimulation (360 neurons); **b,** movement, since the reduction in firing rate persists even when calculated exclusively for periods without movement (267 neurons); **c,** a spatial bias in the orientation of the animal, because the decrease persists even when calculated exclusively for periods when the active loudspeaker was exclusively ipsilaterally located (137 neurons). The Wilcoxon signed-rank test was used in all analysis.

Extended Data Fig. 3: Preferred egocentric sound-source location correspondence between the target and the background loudspeakers. a, during the onset (left), offset (center) and late response periods (right) color-coded with the type of spatial tuning for each loudspeaker; D: direction-sensitive O: orientation-sensitive **b,** during the onset period, color-coded with the latency determined for each neuron; **c,** during the onset period, for the multi-unit activity.

Extended data figure 4: Different response properties of neuronal types. a, Response properties of putative regular spiking (RS) and fast spiking (FS) neurons were separated based on spike-waveform properties by k-means clustering analysis. The clustering was performed on the full width at half maximum of the peak (FWHP) and of the through, and on the through to peak time. (364 neurons: 102 FS and 262 RS). **b** to **e**, On the population level, FS neurons exhibited shorter latencies (364 neurons) (b) and higher mean firing rates, both for spontaneous activity (364 neurons) (c) as well as in response to the target (244 neurons) (d) or background loudspeaker (333 neurons) (e). **f** and **g**, Spatial tuning was sharper in RS neurons (204 and 354 neurons for target and background, respectively). All p-values based on the Mann-Whitney U-test.

During sound presentation in moments without movement

a Preferred egocentric sound-source location during the onset period

b Vector Strength

Extended Data Fig. 5: Tuning complexity and loudspeaker specificity are also evident when considering only stationary episodes. a, Equivalent to Fig. 2a for moments without movement. **b,** Equivalent to Fig. 2b for moments without movement.

Extended Data Fig. 6: Influence of eliminating neuronal tuning classes on the decoding accuracy. Difference in the confusion matrix relative to the original test dataset when 18 randomly selected units (left), 18 units with differential spatial tuning (middle) or 18 canonic hemispherically tuned units (right) are excluded from the test dataset (see also Fig. 3e). Number of sampling cycles: 100.

Video 1: Example trial of an animal performing in the task. Left: Allocentric perspective: The animal starts a trial by staying 1 s on top of the initiation platform (yellow circle) while its position and orientation is tracked (green ellipse and arrow). The moments of sound stimulation are represented by the blinking blue circle (background loudspeaker) or the orange circle (target loudspeaker. After staying 6 s in the island (orange circle), a reward is dropped. Right: Egocentric perspective: Representation of the location of the active loudspeaker relative to the animal - blue if the background loudspeaker is active and orange if the target loudspeaker is active.