Article Summary Line: Molecular detection of West Nile virus in horses in

Northeast and Southeast Brazil

Running Title: Genomic monitoring of West Nile virus in Brazil

West Nile virus detection in horses in three Brazilian states.

Erica Azevedo Costa[1^], Marta Giovanetti[2,3^], Lilian Silva Catenacci[4^], Vagner

Fonseca[3,5,6^], Flavia Aburjaile[2,3^], Felipe Melo Campos Iani[7], Marcelo Adriano da

Cunha e Silva Vieira[8], Maria Isabel Maldonado Coelho Guedes[1], Beatriz Senra

Álvares da Silva Ramos[1], Aila Solimar Gonçalves Silva[1], Tulio de Oliveira[5], Noely

Fabiana Oliveira de Moura[9], Alessandro Pecego Martins Romano[9], Lauro César

Soares Feitosa[10], José Joffre Martins Bayeux[11], Raffaella Bertoni Cavalcanti

Teixeira[12], Osmaikon Lisboa Lobato[13], Silvokleio da Costa Silva[13], José Lourenço[14],

Luiz Carlos Junior Alcantara[2,3]*.

Affiliations

[1]Departamento de Medicina Veterinária Preventiva, Escola de Veterinaria,

Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil;

[2]Laboratório de Flavivírus, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de

Janeiro, Brazil; [3]Laboratório de Genética Celular e Molecular, Universidade Federal

de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil; [4]Departamento De

Morfofisiologia Veterinária, Universidade Federal do Piauí; [5]KwaZulu-Natal

Research Innovation and Sequencing Platform (KRISP), School of Laboratory

Medicine and Medical Sciences, College of Health Sciences, University of KwaZulu-

Natal, Durban, South Africa; [6]Coordenação Geral dos Laboratórios de Saúde

Pública/Secretaria de Vigilância em Saúde, Ministério da Saúde, (CGLAB/SVS-MS)

Brasília, Distrito Federal 70719-040, Brazil; [7]Laboratório Central de Saúde Pública,

Fundação Ezequiel Dias, Belo Horizonte, Brazil; [8]Piaui State Health Department,

Terezina, Brazil; [9]Coordenacao Geral das Arboviroses, Secretaria de Vigilância em

Saúde/Ministério da Saúde, Brasília, Distrito Federal, Brazil; [10]Departamento de

Clínica e Cirurgia Veterinária, Centro de Ciências Agrárias, Universidade Federal do

Piauí, Teresina, Piauí, Brazil; [11]UNIVAP- Universidade Vale do Paraíba, Faculdade

de Ciências da Saúde, Medicina Veterinária, Urbanova, São José Dos Campos, SP;

[12]Departamento de Clinica e Cirurgia Veterinárias, Escola de Veterinaria,

Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil;

[13]Laboratório de Genética e Conservação de Germoplasma, Campus Prof.ª

Cinobelina Elvas, Universidade Federal do Piauí, Bom Jesus, Piauí, Brazil;

[14]Department of Zoology, University of Oxford, Oxford OX1 3PS, UK.

Keywords: West Nile virus, genomic monitoring, molecular detection, Brazil

^Denotes equal contribution

*Correspondence to: luiz.alcantara@ioc.fiocruz.br

Dr Luiz Carlos Júnior Alcantara - Laboratório de Flavivírus, Instituto Oswaldo Cruz,

Fundação Oswaldo Cruz Avenida Brasil, 4365, Manguinhos - 21040-360 Rio de

Janeiro, Brazil; +552125621707/+5531993971962 luiz.alcantara@ioc.fiocruz.br

Abstract

We report genetic evidence of WNV circulation in three Brazilian states, isolated from equine red blood cells. Our analyses demonstrate that much is still unknown on the virus' local epidemiology. We advocate for a shift to active surveillance, to ensure adequate control for future epidemics with spill-over potential to humans.

Text

West Nile virus (WNV), a member of the *Flaviviridae* family, was first identified in the West Nile district of Uganda in 1937, but is nowadays commonly found in Africa, Europe, North America, the Middle East, and Asia (*1, 2, 3*).

WNV transmission is maintained in a mosquito-bird cycle, for which the genus *Culex*, in particular *Cx. pipiens*, are considered the principal vectors (*4*). WNV can infect humans, equines and other mammals, but these are considered "dead-end" hosts, given their weak potential to function as amplifying hosts to spread infection onwards (*5, 6*). Around 80% of WNV infections in humans are asymptomatic while the rest may develop mild or severe disease. Mild disease includes fever, headache, tiredness and vomiting (*7, 8*), while severe disease (neuroinvasive) is characterized by high fever, coma, convulsions and paralysis (*7, 8*). Equine infections can occasionally cause neurological disease and death (*7, 8*), such that equines typically serve as sentinel species for WNV outbreaks with potential for spill-over into human populations.

Genome detection of WNV in South America were originally reported in horses (Argentina in 2006) and captive flamingos (Colombia, in 2012) (*9, 10*). The first ever

sequenced genome in Brazil was in 2019, when the virus was isolated from a horse

sicked with severe neurological disease in the Espírito Santo state (*11*). Despite

multiple studies reporting serological evidence suggestive of WNV circulation in

Brazil (*11-13*) and reports human WNV disease confirmed cases in the Piauí state

(*13*), much is unknown about genomic diversity, evolution and transmission

dynamics across the country.

Here, we report genetic evidence of WNV circulation in three Brazilian states

extracted from equine red blood cells (RBCs) with neurological or ophthalmic

disease and use a computational approach to explore the theoretical transmission

potential of WNV within one of those states.

Samples (RBCs) from three horses with suspected WNV infection obtained from

southern (Minas Gerais and São Paulo) and northeastern (Piauí) Brazilian states

were sent for molecular diagnosis at the *Departamento de Medicina Veterinária*

Preventiva at the Federal University of Minas Gerais (UFMG) (**for details see**

Appendix).

RNAs were extracted from red blood cells and tested using an in-house PCR assay

(**see Technical Appendix for details**). WNV-specific RT-PCR amplification

products were obtained using a combination of nested and multiplex PCR scheme

(**Figure 1 panel C, D**) (**see Technical Appendix for details**). A multiplex PCR

primer scheme was then designed (**Appendix Table 1**) to generate complete

genomes sequences by means of portable nanopore sequencing. The published

WNV genome from Brazil (MH643887) was used to generate a mean 98.4%

consensus sequences that formed the target for primer design. New genomes were

deposited in the GenBank with accession numbers MW420987, MW420988 and

MW420989 (**Table 1).**

We constructed phylogenetic trees to explore the relationship of the sequenced

genomes to those from elsewhere globally. We retrieved 2321 WNV genome

sequences with associated lineage date and country of collection from GenBank,

from which we generated a subset that included the highly supported (>0.9) clade

containing the newly WNV strains obtained in this study plus 29 sequences

(randomly sampled) from all lineages and performed phylogenetic analysis (**see**

Technical Appendix for detail). An automated online phylogenetic tool to identify

and classify WNV sequences was developed (available at

http://krisp.ukzn.ac.za/app/typingtool/wnv/job/9b40f631-51c4-419c-9edf-

2206e7cd8d9c/interactive-tree/phylo-WNV.xml).

Phylogenies estimated by the newly developed WNV typing tool, along with

maximum likelihood methods (**Figure 1 panel A),** consistently placed the Brazilian

genomes in a single clade within the 1a lineage with maximum statistical support

(bootstrap = 100%) (**Supplementary Figure 1**).

Time-resolved maximum likelihood tree appeared to be consistent with previous

estimates (*11*) and showed that the new genomes clustered with strong bootstrap

support (97%) with a WNV strain isolated from an *Aedes albopictus* mosquito in

Washington DC, USA in 2019 (**Figure 1 panel B**). Interestingly, these new isolates

did not group with the previously sequenced genome in 2019 from the Espirito Santo

state, suggesting that inter-continental introduction events might be frequent in
Brazil.

We also explored the dynamics of suspected human WNV cases (**see Technical**
Appendix for detail) in the three Brazilian states for which we had sequence data.
Between 2014 and 2019, all states had suspected cases, suggesting continuous
circulation or recent importation from other regions of the country or elsewhere. The
state of Piauí presented sufficient reports (N=227) for an exploration of the geo-
temporal dynamics of WNV spread (**Figure 2A**), while the other two states had a
much smaller number of total reports (N=5 for MG,24 for SP).

To estimate the transmission potential of WNV we calculated the index P, a
computational approach informed by climatic variables from Lourenço et al. recently
applied in Israel (*14*) (**see Technical Appendix for detail**). Between 2015 and 2016,
cases in Piauí presented a seasonal signal typical of WNV and other mosquito-borne
viruses, with a peak between February and June (late summer and autumn). We
compared suspected cases per month to the monthly mean transmission potential
across the state (**Figure 2B**). The index captured the seasonal variation in reporting,
albeit with a lag of 1 month. The latter could have resulted from the suspected nature
of the cases (rather than confirmed), and from averaging climatic variables across
the vast spatial dimension of the state, thus missing key geo-temporal resolution in
the association of the estimated transmission potential and cases. The Spearman's
correlation between the two variables was 0.4 (non-significant), but increased to 0.7
(significant, p-value=0.003) when shifting the index by a month into the future.
Similarly, to the recent work on WNV in Israel (*14*), the area under the index P curve

(AUC) was positively related to the total suspected cases per year: with

AUC2015>AUC2016 (AUC=3.71, 3.18, respectively), and cases2015>cases2016

(N=118, 76 respectively).

We also estimated the index P in space for 2015, which highlighted some geo-

patterns (**Figures 2C-F**). In summer, when reports were high, estimated

transmission potential was mainly highest in the center at the –42 longitude axis

(**Figure 2C).** Into autumn, the southern region was the first to present lower

transmission potential (**Figure 2D**), followed by the northern region in the winter

(**Figure 2E**). In spring, when the trough of reporting is observed (2015-2016), the

lowest transmission potential was estimated in the same central region presenting

highest potential in the summer and autumn (**Figure 2F**). Notably, these geo-

temporal snapshots highlight a possible wave of seasonal transmission potential,

starting in the south-west in spring when the rest of the state remains at lowest

potential (**Figure 2F**), moving to the north-east in summer when many regions

remain at highest potential (**Figure 2C**), and ending in the north during autumn

(**Figure 2D**).

Conclusions

Our analyses indicate that additional data is required to better identify routes of WNV

importation into Brazil, and to understand its local transmission dynamics. Critically,

it is still uncertain where in the country WNV is endemic. Indeed, if under reporting is

frequent, the data presented in this article is compatible with both sporadic or

endemic local transmission. Furthermore, the detection of WNV RNA from equine

whole blood presented in this study proved to be an effective diagnostic method in

horses. Shifting from passive to active WNV screening and sequencing in equines

and birds in Brazil must be implemented to better understand the virus' local

epidemiology, and to be able to act accordingly to prevent any future epidemics with

significant spill-over to humans.

Biographical sketch

Dr. Costa is a veterinarian and a Professor at the Federal University of Minas Gerais,

Minas Gerais state, Southeast Brazil. Her research focuses on the improvement of

diagnostic techniques for a prompt equine encephalitis diagnosis and managements.

Acknowledgements

The authors thank the important contributions of the Municipal and Piaui State

Health Department (SESAPI, FMS), Municipal and Piaui State Animal Health

Department (ADAPI), Laboratório de Saúde Pública do Piauí (LACEN-PI) and the

colleague Thiago dos Santos Silva. We also thank the sponsoring institutions: Saint

Louis Zoo WildCare Institute and Institute for Conservation Medicine (USA),

Universidade Federal do Piauí (UFPI), Fundação de Amparo a Pesquisa do Estado

do Piauí (FAPEPI). Authors also thank the Municipal and State Health Department of

São Paulo and Minas Gerais state.

Financial Disclosure

This work was supported by Decit/SCTIE/BrMoH/CNPq (440685/2016-8), by CAPES

(88887.130716/2016-00) and by the European Union's Horizon 2020 Research and

Innovation Programme under ZIKAlliance Grant Agreement no. 734548. MG and

LCJA is supported by Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro

(FAPERJ). JL is supported by a lectureship from the Department of Zoology,
University of Oxford.

References

1. Fall G, Di Paola N, Faye M, et al. Biological and phylogenetic characteristics
of West African lineages of West Nile virus (DWC Beasley, Ed.). PLoS Negl
Trop Dis. 2017;11: 1-23.

2. Smithburn KC, Hughes TP, Burke AW, et al. A neutrotropic virus isolated from
the blood of a native of Uganda. Am J Trop Med Hyg. 1940; 20: 471-2.

3. Murgue B, Zeller H, Deubel V. The ecology and epidemiology of West Nile
virus in Africa, Europe and Asia. Curr Top Microbiol Immunol. 2002;267:195-
221.

4. Campbell GL, Marfin AA, Lanciotti RS, et al. West Nile virus. Lancet Infect
Dis. 2002;2:519-29.

5. Gamino V, Ho□fle U. Pathology and tissue tropism of natural West Nile virus
infection in birds: a review. Vet Res. 2013;44:46-89.

6. Bunning ML, Bowen RA, Cropp CB, et al. Experimental infection of horses
with West Nile virus. Emerg Infect Dis. 2002;8:380-6.

7. Hayes EB, Sejvar JJ, Zaki SR, et al. Virology, pathology, and clinical
manifestations of West Nile virus disease. Emerg Infect Dis. 2005;11:1174–
1179.

8. Kramer LD, Li J, Shi P-Y. West Nile virus. Lancet Neurol. 2007;6:171–181.

9. Morales MA, Barrandeguy M, Fabbri C, et al. West Nile virus isolation from
equines in Argentina. Emerg Infect Dis. 2006;12:1559-61.

10. Osorio JE, Ciuoderis KA, Lopera JG, et al. Characterization of West Nile
viruses isolated from captive American flamingoes (Phoenicopterus ruber) in
Medellin, Colombia. Am J Trop Med Hyg. 2012;87:565-72.

11. Martins LC, Silva EVPD, Casseb LMN, et al. First isolation of West Nile virus
in Brazil. Mem Inst Oswaldo Cruz. 2019 Jan 17;114:e180332.

12. Pauvolid-Corrêa A, Morales MA, Levis S, Figueiredo LTM, Cou- to-Lima D,
Campos Z, et al. Neutralising antibodies for West Nile virus in horses from
Brazilian Pantanal. Mem Inst Oswaldo Cruz. 2011;106:467-74.

13. Vieira MA, Romano APM, Borba AS, et al. West Nile Virus Encephalitis: The
First Human Case Recorded in Brazil. Am J Trop Med Hyg. 2015; 93(2):377-
9.

14. Lourenço J, Thompson RN, Thézé J, et al. Characterising West Nile virus
epidemiology in Israel using a transmission suitability index. Euro Surveill.
2020;2:5-41.

Table 1. Epidemiological information and sequencing statistics of the 3 sequenced samples
of WNV sampled in Minas Gerais, São Paulo and Piauí Brazilian states.

ID	Sample	Collection date	Age	Sex	State	Municipality	Reads	Coverage (%)	Depth of coverage	Lineage Assignment	Acession Number	Clinical sign
BC02_07	RBCs	11/07/2018	9 months	F	MG	Sabara	343743	97.9	6527.6	Lineage 1a	MW420989	Chorioretinitis
BC03_04	RBCs	30/07/2019	13 years-old	M	SP	São Bernardo do Campo	170980	97.9	3189.7	Lineage 1a	MW420988	Muscle stiffness, tremor retinal and flaccid paralysis

												Motor incoordination, paddling movements, loss of sensitivity
BC05_06	RBCs	21/08/2020	5 years-old	F	PI	Parnaíba	222516	99.4	4121.4	Lineage 1a	MW420987		

ID=study identifier; Collection date=Sample collection date; Municipality=Municipality of residence; State=

MG-Minas Gerais; SP-São Paulo; PI-Piauí; Sex: M=Male; F=Female; Accession Number=NCBI accession

number.

Figure 1. Investigation of WNV infections in Brazil, between July 2018 - September 2020. A)

Midpoint rooted maximum-likelihood phylogeny of WNV genomes, showing major lineages. The scale

bar is in units of substitutions per site (s/s). Support for branching structure is shown by bootstrap

values at nodes. B) Time-resolved maximum likelihood tree showing the WNV strains belonged to the

1a lineage. Colors indicate geographic location of sampling. The new Brazilian WNV strains are

shown in bold. C-D) Agarose gel electrophoresis of amplicons from assay for WNV. (C) nested RT-

PCR. MW-Molecular weight marker; 1- plasm of horse from São Paulo; 2- buffy coat of horse from

São Paulo; 3- washed RBC of horse from São Paulo; 4- without sample; 5 and 6-positive control

(synthetic gene); NTC, no template control. (D) Multiplex PCR. MW-Molecular weight marker; 1-

horse form Minas Gerais (pair primers); 2- horse form Minas Gerais (impair primers); 3- horse form

Sao Paulo (pair primers); 4- horse form Sao Paulo (impair primers); 5-horse form Piaui (pair primers);

6-horse form Piaui (impair primers); NTC, no template control.

Figure 2 - WNV reporting of suspected cases and estimated transmission potential. (A) Time

series of suspected human WNV cases in the Brazilian states of Minas Gerais (MG, green), São

Paulo (SP, pink) and Piauí (PI, gold). (B) WNV human suspected cases and estimated monthly mean

transmission potential (index P, grey) for the state of Piauí (gold). The shaded area includes the

standard deviation. (C-F) Maps of estimated transmission potential (index P) per geo-location for four

different months in the year of 2015 (from C to F, March, May, August, November, respectively). Color

legend on the right applies to all maps. The city of Parnaíba from where the Piauí sequence is

available is marked by a black, full point in the north of the state.

Table 1. Epidemiological information and sequencing statistics of the 3 sequenced samples of WNV sampled in Minas Gerais, Sao Paulo, Piaui Brazilian states.

ID	Sample	Collection date	Age	Sex	State	Municipality	Reads	Coverage (%)	Depth of coverage	Lineage Assignment	Acession Number	Clinical sign
BC02_07	RBCs	11/07/2018	9 months	F	MG	Sabara	343743	97.9	6527.6	Lineage 1a	MW420989	Chorioretinitis
BC03_04	RBCs	30/07/2019	13 years-old	M	SP	São Bernardo do Campo	170980	97.9	3189.7	Lineage 1a	MW420988	Muscle stiffness, tremor retinal and flaccid paralysis
BC05_06	RBCs	21/08/2020	5 years-old	F	PI	Parnaíba	222516	99.4	4121.4	Lineage 1a	MW420987	Neurological complications

ID=study identifier; RBCs=Red Blood Cells; Collection date=Sample collection date; Municipality=Municipality of residence; State= MG-Minas Gerais; SP-Sao Paulo; PI-Piaui; Sex: M=Male; F=Female; Accession Number=NCBI accession number

A WNV in all states

B WNV and transmission potential in Piauí state

C Piauí, Mar. 2015

D Piauí, May 2015

E Piauí, Aug. 2015

F Piauí, Nov. 2015