Title

Review and performance evaluation of trait-based between-community dissimilarity measures

Author details

Attila Lengyel[1*] & Zoltán Botta-Dukát[2*]

*Centre for Ecological Research, Institute of Ecology and Botany, Alkotmány u. 2-4., H-2163

Vácrátót, Hungary

[1] corresponding author, lengyel.attila@ecolres.hu

[2] botta-dukat.zoltan@ecolres.hu

Abstract

1. In the recent years a variety of indices have been proposed with the aim of quantifying functional dissimilarity between communities. These indices follow different approaches to account for between-species similarities in the calculation of community dissimilarity, yet they all have been proposed as straightforward tools.

2. In this paper we reviewed the trait-based dissimilarity indices available in the literature, contrasted the approaches they follow, and evaluated their performance in terms of correlation with an underlying environmental gradient using individual-based community simulations with different gradient lengths. We tested how strongly dissimilarities calculated by different indices correlate with environmental distances. Using random forest models we tested the importance of gradient length, the choice of data type (abundance vs. presence/absence), the transformation of between-species similarities (linear vs. exponential), and the dissimilarity index in the predicting correlation value.

3. We found that many indices behave very similarly and reach high correlation with environmental distances. There were only a few indices (e.g. Rao's D_Q, and representatives of the nearest neighbour approach) which performed regularly poorer than the others. By far the strongest determinant of correlation with environmental distance was the gradient length, followed by the data type. The dissimilarity index and the transformation method seemed not crucial decisions when correlation with an underlying gradient is to be maximized.

4. *Synthesis*: We provide a framework of functional dissimilarity indices and discuss the approaches they follow. Although, these indices are formulated in different ways and follow different approaches, most of them perform similarly well. At the same time,

35   sample properties (e.g. gradient length) determine the correlation between trait-based

36   dissimilarity and environmental distance more fundamentally.

Keywords

39   beta diversity, dissimilarity index, distance metric, community ecology, functional traits

Abbreviations

42   CDF = cumulative distribution function, CWM = community-weighted mean, FDissim =

43   functional dissimilarity, VIS = variable importance score

Introduction

Understanding and explaining the variation of living communities along dimensions of space

and time have been in the focus of ecological research ever since. The widely applied scheme

by Whittaker (1960, 1972) to tackle questions of different aspects of community variation

divides community diversity into alpha (within-community), beta (between-community) and

gamma (across-community) components. It is no exaggeration to say that among these three,

beta diversity sparked the most controversy due to the multitude of ways how it can be

formulated (Tuomisto 2010a,b, Anderson et al. 2011, Podani & Schmera 2011, Baselga &

Leprieur 2015). One of the most popular approaches to beta diversity builds upon

quantification of variation between pairs of communities using dissimilarity indices

(Anderson et al. 2006, Legendre & De Cáceres 2013, Ricotta 2017). A broad spectrum of

such dissimilarity indices are available for many specific purposes providing elementary tools

for different fields of ecology and beyond (see reviews by Legendre & Legendre 1998, Podani

2000). Nevertheless, choosing from such many options requires a more or less subjective

decision from the researcher which may affect the final result of the analysis. Comparative

reviews of dissimilarity indices (Faith et al. 1987, Koleff et al. 2003) and evaluations of

effects of methodological decisions (Lengyel & Podani 2015) are inevitably helpful in making

these decisions.

The most popular, yet not exclusive, interpretations of diversity for long time considered

species as variables which are unrelated with each other. In the last two decades, however, the

functional approach to ecological questions gained unprecedented attention (Díaz & Cabido

2001, McGill et al. 2006). This approach relies on the fact that species are not all maximally

different from each other, rather they can be considered related with respect to similarities in

their traits thought to represent their roles in ecosystems (Violle et al. 2007). The need for

explicitly accounting for between-species relatedness generated a wave of methodological

improvements that introduced new methods in the calculation of diversity. Next to a lively

scientific discussion on how functional alpha diversity can be appropriately quantified (Mason

et al. 2005, Petchey & Gaston 2006, Villéger et al. 2008, Mouchet et al. 2010), suggestions

were made also for the expression of functional beta diversity (Swenson 2011, Botta-Dukát

2018, Chao et al. 2019). Among them, a large variety of indices for calculating dissimilarity

between pairs of communities on the basis of the traits of their species have been proposed

(e.g. Ricotta & Burrascano 2008, Cardoso et al. 2014, Ricotta & Pavoine 2015). Although

these indices have been introduced as straightforward measures for revealing between-

community dissimilarity on the basis of traits, they have very different concepts behind, and

we still lack a comparative review of them.

In this paper we aim to provide an overview and a conceptual framework for the pairwise

functional dissimilarity (hereafter called FDissim) measures available in the literature to our

best knowledge. We start with a (1) short overview of the concept and indices of ecological

83   (dis-)similarity without accounting for relatedness of species, then (2) we review and classify

84   FDissim indices according to their conceptual basis, and (3) we test the performance of

85   FDissim indices.

Short overview of taxon-based (dis-)similarity methods

Most FDissim measures are generalizations of simple indices which were originally designed

for expressing dissimilarity based on species composition (that is, omitting similarities

between species). We start the review of trait-based (dis-)similarity measures with a brief

summary of these species-based indices. Then, we present a framework of approaches

including several families of trait-based dissimilarity indices.

Species-based indices

Most indices can be written in either similarity (s) or dissimilarity ($d=1-s$) form but when we

do not see necessary to specify the form, we call them 'resemblances'. In the case of

presence/absence data, these indices are based on the well-known 2×2 contingency table

whose cells represent the number of species shared (denoted by a), as well as the number of

species occurring only in one of the communities (b and c). The fourth cell of the contingency

table quantifying the number of shared absences is disregarded by these indices and rarely

used in ecological analyses (but see Tamás et al. 2001). All these indices agree that they

express similarity as the proportion of shared diversity to total diversity. Hence, all of them

range between 0 and 1. In the case of presence/absence data the number of shared species, a,

in the numerator stands for shared diversity for all indices, while the denominators are

different. In the Sørensen index (s_S) the denominator is the arithmetic mean of the species

numbers of the two communities, in Ochiai index (s_O) it is their geometric mean, in

Kulczynski (s_K) it is their harmonic mean, while in Simpson index (s_{Si}) it is the richness of the

species poorer community. If the two communities are equally species-rich, then these indices

are equal, otherwise $s_S < s_O < s_K < s_{Si}$. In the Jaccard index (s_J), the denominator is the total

number of species in the two communities, while in Sokal & Sneath index (s_{SS}) species

occurring in a single community are taken into account with double weight. There is a direct

and monotonic relationship between Jaccard, Sørensen, and Sokal & Sneath indices (see

Appendix S1). Table 1 summarizes the similarity and dissimilarity forms of the above indices.

For abundance data, the resemblance of two communities is derived from the summation of

species-wise differences, with the simplest interpretation being the Euclidean and the

Manhattan distances, respectively:

Eq. 1. $$d_{Euclidean} = \sqrt{\sum_{i=1}^{S_{jk}} \left(x_{ij} - x_{ik} \right)^2}$$

Eq. 2. $$d_{Manhattan} = \sum_{i=1}^{S_{jk}} \left| x_{ij} - x_{ik} \right|$$

where x_{ij} and x_{ik} are the abundance of species i in communities j and k, S_{jk} is the total number

of species in j and k. For both indices, the minimum is 0 but the maximum of Euclidean

distance is the square-root of the sum of squared abundances, while for Manhattan distance

the maximum is the sum of abundances. Obviously, their dependence on total abundance

makes these index values difficult to compare across samples; therefore, indices including a

standardization have become more popular in ecological studies. The standardization is

possible in several ways. The first option is to standardize raw species contributions to

between-community dissimilarity (x_{ij}-x_{ik}), and then to sum them. Therefore, each species-level

difference in abundance should be divided by a scaling factor in a way that maximal species-

level difference is 1 and this difference is maximal if species present only one of the

compared communities. Summing x_{ij} and x_{ik} in the denominator satisfies this requirement and

gives a well-known distance measure, the Canberra index:

Eq. 3.
$$d_{Canberra} = \sum_{i=1}^{S_{jk}} \frac{|x_{ij}-x_{ik}|}{(x_{ij}+x_{ik})}$$

131   However, Canberra index still ranges between 0 and S_{jk}. According to Ricotta & Podani

132   (2017), the normalized Canberra index can be derived by unweighted averaging of species

133   contributions:

134   Eq. 4.
$$d_{NCanberra} = \frac{1}{S_{jk}}\sum_{i=1}^{S_{jk}} \frac{|x_{ij}-x_{ik}|}{(x_{ij}+x_{ik})}$$

135   Alternatively, species-level differences can be divided by max(x_{ij}, x_{ik}). It also results unity, if

136   species occur only either of the plots. Ricotta & Podani (2017) called this modified Canberra

137   index, whose normalized version follows:

138   Eq. 5.
$$d_{NMCanberra} = \frac{1}{S_{jk}}\sum_{i=1}^{S_{jk}} \frac{|x_{ij}-x_{ik}|}{\max{(x_{ij},x_{ik})}}$$

Calculating from binary data, both normalized Canberra and normalized modified Canberra

result in Jaccard dissimilarity.

A different way of standardization is possible if raw species-level differences are summed and

divided by the sum of their theoretical maxima. In this case, the denominator can follow the

logic of Canberra index, thus leading to the Bray-Curtis index:

Eq. 6.
$$d_{BC} = \frac{\sum_{i=1}^{S_{jk}}|x_{ij}-x_{ik}|}{\sum_{i=1}^{S_{jk}}(x_{ij}+x_{ik})}$$

145   Analogously with the normalized modified Canberra index, instead of the sum, the

146   denominator may contain the maximum of abundance, resulting in the formula known as

147   Marczewski-Steinhaus index:

148   Eq. 7.
$$d_{MS} = \frac{\sum_{i=1}^{S_{jk}}|x_{ij}-x_{ik}|}{\sum_{i=1}^{S_{jk}}\max{(x_{ij},x_{ik})}}$$

Worth to note that Bray-Curtis and Marczewski-Steinhaus indices calculated on

presence/absence data return the values of Sørensen index and Jaccard index in dissimilarity

form, respectively. Moreover, several abundance-based indices can be expressed if we

generalize *a*, *b*, and *c* quantities used during the definition of indices for presence/absence

data (Tamás et al. 2001).

Eq. 8. $$a' = \sum_{i=1}^{S_{jk}} \min\ (x_{ij}, x_{ik})$$

Eq. 9. $$b' = \sum_{i=1}^{S_{jk}} (\max\ (x_{ij}, x_{ik}) - x_{ij})$$

Eq. 10. $$c' = \sum_{i=1}^{S_{jk}} (\max\ (x_{ij}, x_{ik}) - x_{ik})$$

Substituting *a*, *b* and *c* with *a'*, *b'* and *c'* into the formula of Sørensen index gives Bray-

Curtis, and doing so with Jaccard index results in the Marczewski-Steinhaus. Abundance

versions of all other presence/absence indices can be created in the same manner.

A classification of FDissim indices

FDissim indices incorporate trait information into the calculation of dissimilarity in different

ways. The simplest solution is when summary statistics or distributions are calculated for the

two communities and a measure of distance or segregation is calculated between them. We

call this the *summary-based class*, and in our review, we include two approaches within this,

the *typical value approach* and the *distribution-based approach*. In the second class we

include indices which utilize a symmetrical species by species (dis-)similarity matrix and link

it directly through matrix operations with the compositional matrix. We call this the

dissimilarity-based class which includes the *probabilistic*, the *ordinariness-based*, the

diversity partitioning, and the *nearest neighbour approaches*. The third class includes

methods which make use of between-species (dis-)similarities for classification of species;

therefore, we call it the *classification-based class*. The classification either transforms the

original structure of the dissimilarity matrix into discrete groups of species which can be used

as functional types, or expresses dissimilarities in a form of a tree-graph where between-

species dissimilarities are organized in an inclusive hierarchy. This is a widespread approach

for accounting for phylogenetic relatedness, since phylogenies are commonly summarized in

the form of cladograms. Such methods heavily rely on the algorithm chosen for the

classification, including the decisions about the number of clusters and the method for

breaking tied values. Examples are provided by Hérault & Honnay (2007), Nipperess et al.

(2010), and Cardoso et al. (2014), while a review is available by Pavoine (2016). As there is

no general recommendation for the classification method, we omit this class from the

framework detailed below and the comparative test. The classification of trait-based

dissimilarity indices and their main properties are summarized on Table 2.

Typical value approach

Indices following this approach represent each community with a typical trait value, and

calculate a distance metric between them. The most commonly applied typical trait value is

the community weighted mean (CWM; Garnier et al. 2004). The rationale behind the CWM

can be linked with the mass ratio hypothesis (Grime 1998) stating that the effect of species on

ecosystem functioning is proportional to their relative abundances. Although, several issues

emerged regarding its limited applicability in statistical inference (Hawkins et al. 2017, Peres-

Neto et al. 2017, Zeleny 2018) and its negligence of within-community variation (Muscarella

& Uriarte 2016), difference in CWM is still considered a reliable indicator of robust changes

in trait composition induced by selective forces like environmental matching or succession

(De Bello et al. 2007, 2013, Kleyer et al. 2012). Ricotta et al. (2015) investigated the

relatedness of the distance between CWMs with the probabilistic approach (see therein) and

showed its applicability on phylogenetic data. Due to its tolerable requirements for

computational capacity, Lengyel et al. (2020) used the Euclidean distance between trait

CWMs of phytosociological relevés for the trait-based numerical classification of grasslands

of Poland with a sample size of 6985 sites and 885 species. Another advantage of this method

is its Euclidean property. Besides the community-weighted mean, other typical values, e.g. the

median or the mode, might be considered depending on the scaling of the trait variable and on

specific research aims.

Distribution-based approach

Instead of typical values, the distribution of trait values is considered a more reliable

representative of the trait composition and variability of a community. Continuous

distributions can be defined by a density function, while discrete distributions by the

probabilities of the possible values, while both types can be characterized by a cumulative

distribution function (CDF). A useful analogue of the distance between typical values might

be distance between discrete distributions, density functions or CDFs.

If data is available on intraspecific trait variation, trait values forms a continuous distribution.

First, separate density functions have to be fitted within each species. Then, density function

of this community-level distribution can be calculated as weighted sum of species level

density functions (Carmona, de Bello, Mason, & Lepš, 2016). If such data is not available, we

can use relative abundances as estimates of probabilities of the corresponding trait values.

Pairs of trait values and their probability form a discrete distribution.

Similarity of density functions can be measured by their overlap (see Appendix S2 for

overview of overlap measures). Overlap functions between within-species trait distributions

has already been proved useful in the quantification of between-species niche segregation

(MacArthur & Levins 1967, Mouillot et al. 2005) or trait-based dissimilarity of species (Lepš

et al. 2006, De Bello et al. 2013). Nevertheless, they are perfectly applicable to the

community level as well.

Gregorius et al. (2003) proposed an index called *delta* for the quantification differences

between discrete trait distributions. Delta is the minimal sum of frequencies shifted from one

trait state to another trait state, weighted by the differences between the respective states.

Minimizing the sum of shifted frequencies is known in linear programming as the

transportation problem (Hitchcock 1941). Due to its relatively high computational demand, it

is unfeasible for large compositional and trait data matrices typically used in ecological

research, therefore, we exclude this index from our comparison.

Difference between two CDFs can be calculated at each possible trait values (i.e. not only the

observed ones), then the sum of them can be used as a trait-based dissimilarity measure. In

Appendix S3 we introduce the distance between CDFs in more detail.

Maximally distinct communities

Species-based dissimilarities, except Euclidean, Manhattan and (non-normalized) Canberra

distances, equal unity, which is their maximum, when the two compared communities do not

share any species. In this context, we could call such communities maximally distinct.

However, when traits are considered, two communities can be similar, even if they do not

share any species. For example, if all species of community *A* is replaced by a similar species

in community *B*, the two communities have no shared species, but from functional point of

view, they are similar. In this context, two communities are maximally distinct, when

similarity of any species from the first community is zero to any species in the other

community. It is a desirable property for a functional similarity index to take the value 0 if

and only if the two compared communities are maximally distinct.

Probabilistic approach

This approach can be traced back to the diversity framework proposed by Rao (1982), and recently extended by Pavoine & Ricotta (2014). Rao's within community diversity is defined as the expected dissimilarity between two randomly drawn individuals from a single community:

Eq. 11. $$Q(\mathbf{p}) = \sum_i \sum_j p_i p_j \delta_{ij}$$

where p_i is the relative abundance of the *i*th species in the community and δ_{ij} is the dissimilarity between species *i* and *j*. This has become a widely used index of functional alpha diversity (Botta-Dukát 2005). Likewise, a between-community component of diversity, $Q(\mathbf{p},\mathbf{q})$, can be defined as the dissimilarity between two random individuals, each selected from different communities:

Eq. 12. $$Q(\mathbf{p}, \mathbf{q}) = \sum_i \sum_j p_i q_j \delta_{ij}$$

Between community diversity can be expressed using within community diversity of the two original communities ($Q(\mathbf{p})$ and $Q(\mathbf{q})$) and the community with mean relative abundances; $Q\left(\frac{\mathbf{p}+\mathbf{q}}{2}\right)$.

Eq. 13. $$2Q\left(\frac{\mathbf{p}+\mathbf{q}}{2}\right) = 2\sum_i \sum_j \frac{p_i+q_i}{2}\frac{p_j+q_j}{2}\delta_{ij} = \frac{1}{2}\sum_i \sum_j [p_i p_j + q_i q_j +$$

$$2p_i q_j]\,\delta_{ij} = \frac{Q(\mathbf{p})+Q(\mathbf{q})}{2} + Q(\mathbf{p}, \mathbf{q})$$

Subtracting mean within community diversity from the between community diversity leads to Rao's dissimilarity (also called *DISC*):

Eq. 14. $D_Q = \sum_i \sum_j p_i q_j \delta_{ij} - \frac{\sum_i \sum_j p_i p_j \delta_{ij} + \sum_i \sum_j q_i q_j \delta_{ij}}{2} = \sum_i \sum_j p_i q_j \delta_{ij} - \frac{Q(\mathbf{p}) + Q(\mathbf{q})}{2} =$

$2Q\left(\frac{\mathbf{p}+\mathbf{q}}{2}\right) - Q(\mathbf{p}) - Q(\mathbf{q})$

where p_i and q_i are the relative abundances of species i in the two communities. Champely

and Chessel (2002) proved that if δ has squared Euclidean property, Rao quadratic entropy is

concave function, i.e. $Q\left(\frac{\mathbf{p}+\mathbf{q}}{2}\right)$ is higher than or equal to mean of $Q(\mathbf{p})$ and $Q(\mathbf{q})$. Thus under

this condition, $D_Q \geq 0$. If $0 \leq \delta_{ij} \leq 1$, $\sum_i \sum_j p_i q_j \delta_{ij}$, which is the weighted average of

between-species distances, also has to be within this range. Therefore, $0 \leq D_Q \leq 1$. However,

D_Q may be much less than 1, even if the two communities are completely distinct, when $Q(\mathbf{p})$

and $Q(\mathbf{q})$ are high. Therefore, Pavoine & Ricotta (2014) suggested dividing D_Q by its

theoretical maximum (see equations 3 and 4 in Pavoine & Ricotta 2014). They recognized

that the resulting indices are representatives of a broader family of indices, hereafter called

dsimcom, which are actually the implementations of Rao's between-community and within-

community components of diversity into the similarity formulae designed for

presence/absence data. For this index, it is necessary to introduce the similarity between

species, $\varepsilon_{ij}=1- \delta_{ij}$. The expected similarity between individuals of different communities,

$A = \sum_i \sum_j p_i q_j \varepsilon_{ij}$ is taken analogous with the shared diversity, *a*, according to the parameters

of the similarity indices for presence/absence data disregarding species properties, while the

expected similarities within communities ($A + B = \sum_i \sum_j p_i p_j \varepsilon_{ij}$ and $A + C = \sum_i \sum_j q_i q_j \varepsilon_{ij}$) are

analogous with the species numbers (*a+b*, *a+c*). In this way, Pavoine & Ricotta (2014)

presented formulae following the Sokal & Sneath, Jaccard, Sørensen, and Ochiai indices.

Additionally, a formula analogous with Whittaker's effective species turnover ($\beta=\gamma/\alpha-1$;

Whittaker 1972, Tuomisto 2010a) is suggested for two communities, which in similarity form

is shown to be identical with the overlap index of Chiu et al. (2014). In this formulation

$\gamma=A+B+C$ and $\alpha=(2A+B+ C)/2$. Pavoine & Ricotta (2014) showed that members of the

dsimcom family provide meaningful values also if absolute abundances, percentage values or

binary occurrences are used instead of relative abundances.

When ε_{ij} contains taxonomical similarities, its off-diagonal elements are 0, and $A=a$, $B=b$, and

$C=c$.

Worth to note the inherent link between D_Q and *CWMdis* on the basis of the geometric

interpretation by Pavoine (2012) and Ricotta et al. (2015). Pavoine (2012) showed that if

between-species dissimilarities are in the form $\delta_{ij}=(d_{ij}^2)/2$ and d_{ij} is Euclidean embeddable, D_Q

is half the squared Euclidean distance between the centroids of two communities – a function

monotonically related with *CWMdis*, the simple Euclidean distance between centroids of

communities. As Ricotta et al. (2015) argue, if species relatedness is only described by a

dissimilarity matrix, which is the common case in phylogenetic analyses, species can be

mapped into a principal coordinate analysis ordination using d_{ij}. Given the Euclidean

embeddable property of d_{ij}, this ordination should produce S-1 or fewer ordination axes, all

with positive eigenvalues. Ordination scores for species can be used as traits, and therefore,

centroids of communities, and (squared) Euclidean distances between communities can be

calculated. In the special case when between-species dissimilarities are Euclidean distances,

D_Q must be equal with the Euclidean distance between the weighted averages of traits, that is,

CWMdis.

It is also notable that Swenson et al. (2011) and Swenson (2011) use the quantity $Q(\mathbf{p}, \mathbf{q})$ as a

standalone index of pairwise beta diversity and call it D_{pw} or "Rao's D". The latter name is

misleading since Rao (1982) himself noted with D_{ij} the *DISC* (or D_Q) index. $Q(\mathbf{p}, \mathbf{q})$ measures

dissimilarity between two communities but the dissimilarity of a community from itself is not

zero. Swenson (2011) also presents a standardized version of $Q(\mathbf{p}, \mathbf{q})$ under the name Rao's

H. With this formula the dissimilarity of a community to itself is scaled to 1, however, its

transformation to a meaningful scale where each community has dissimilarity value zero

towards itself is not elaborated. Due to this drawback, we do not consider these indices in our

review of functional dissimilarity measures.

Schmidt et al. (2017) proposed probabilistic indices with weighted and unweighted versions

for expressing community similarity on the basis of taxa interaction networks (called *TINA*,

taxa interaction-adjusted) and phylogenetic relatedness (*PINA*, phylogenetic interaction-

adjusted). *TINA* and *PINA* differ only in what type of data the interaction matrix contains.

Notably, the functional formula of weighted *TINA* is identical with the Ochiai version of

dsimcom. However, the unweighted *TINA*, abbreviated *TU*, is not a special case of *TINA*,

which we consider an inconsistency. Therefore, we did not include *TU* as a separate index.

Ordinariness-based approach

With respect to functional alpha diversity, Leinster & Cobbold (2012) introduced the concept

of species ordinariness defined as the weighted sum of relative abundances of species similar

to a focal species within the same community, or in other words, the expected similarity of an

individual of the focal species and an individual chosen randomly from the same community.

According to Ricotta & Pavoine (2015) it is straightforward to replace abundances with

ordinariness values in the species-based (dis-)similarity indices. Following this concept,

Ricotta & Pavoine (2015) introduced a new family of trait-based similarity measures called

dissABC. *dissABC* applies the schemes of Jaccard, Sørensen, Ochiai, Kulczynski, Sokal &

Sneath, and Simpson indices. Either relative or absolute abundances can be chosen as input

values. Species ordinariness values can be calculated either with respect to the pooled species

list of the two communities under comparison, or to the total species list of the data matrix.

For species-based analyses, Ricotta & Podani (2017) suggested a general formula of distance

measures in which community dissimilarity is calculated by the weighted averaging of

species-level differences in abundance. From this formula, a normalized Canberra distance,

Bray-Curtis distance, Marczewski-Steinhaus index, and an evenness-based dissimilarity index

(Ricotta 2018) can be derived. According to Pavoine & Ricotta (2019), replacing species

abundances with species ordinariness values, a meaningful dissimilarity index can be

designed, which is called *generalized_Tradidiss*. Additionally, this index contains a factor

which weights the contribution of each species to the overall dissimilarity between the two

communities. This weight can be set to give even weight to all species or to weigh them

proportionally to their relative abundance in the pooled communities.

Diversity partitioning approach

Following the work of Hill (1973), a community with *diversity of order q*, ^qD, is as diverse as

a theoretical community containing ^qD equally abundant species. The order of diversity, q,

expresses the weight given to differences in species abundance, $q = 0$ representing the

presence/absence case, $q = \infty$ considering only the relative abundance of the most abundant

species in the community. Without accounting for interspecific similarities, there is emerging

consensus that using effective numbers (also called number of equivalents) is a

straightforward way for partitioning diversity into within-community (alpha), between-

community (beta) and across-community (gamma) components (Jost 2007). Of these three,

the between-community component, beta diversity, can be interpreted as a form of

dissimilarity when applied for two communities (Ricotta 2017). Beta diversity can be derived

from alpha and gamma diversity in a multiplicative (beta = gamma/alpha) or an additive way

(beta = gamma – alpha). Jost (2007) and Chao et al. (2012) argued that multiplicative beta

diversity is a useful way for quantifying community differentiation; however, due to its

scaling between 1 and N (N being the number of communities) it is not comparable across

samples containing different numbers of communities. To remove this dependence, they offer

three solutions with which the value of multiplicative beta can be normed. Although, for

pairwise comparisons, N is always 2, it seems straightforward to follow these

recommendations, since the scaling between 0 and 1 has several advantages, and most other

indices also share this property. The rescaling formulae of Chao et al. (2012) embody

different concepts of community (dis-)similarity, which together we call the family of

multiplicative beta indices. The first formula is the relative turnover rate per community,

which is a linear transformation of beta to the normed scale.

Eq. 15. $$\beta_{turnover}\langle q \rangle = (^q\beta - 1)/(N - 1)$$

Here 0 means identical species composition, while 1 indicates totally distinct communities. In

the pairwise comparison (N = 2), $\beta_{turnover}(q) = {}^q\beta - 1$.

The second index measures homogeneity, and is a linear transformation of the inverse of beta.

With respect to the fact that the complement term of homogeneity is heterogeneity, we call its

dissimilarity form $\beta_{heterogeneity}$:

Eq. 16. $$\beta_{heterogeneity}\langle q \rangle = 1 - \left(\frac{1}{{}^q\beta} - \frac{1}{N}\right)\Big/\left(1 - \frac{1}{N}\right)$$

When N = 2, $\beta_{het}(q) = 2 - 2/{}^q\beta$. With $q = 0$ (presence/absence case) the index is identical with

Jaccard index, while with $q = \infty$ (abundance case) it is the Morisita & Horn index.

The third index measures overlap between communities, whose counterpart is segregation,

thus we call it β_{segr}:

Eq. 17. $$\beta_{segregation}\langle q \rangle = 1 - \left[\left(\frac{1}{{}^q\beta}\right)^{q-1} - \left(\frac{1}{N}\right)^{q-1}\right]\Big/\left[1 - \left(\frac{1}{N}\right)^{q-1}\right]$$

With $q = 0$, $\beta_{segregation}\langle q \rangle = \beta_{turnover}\langle q \rangle$, and both gives the Sørensen index.

According to Leinster & Cobbold (2012), it is possible to implement species similarities in the

calculation of effective numbers. This way, the meaning of $^qD^{\mathbf{Z}}$, is the diversity of a

theoretical community with $^qD^{\mathbf{Z}}$ *equally abundant* and *maximally different* species. Hence,

both unevenness in the abundance structure and the between-species similarities decrease the

value of effective species number. Due to measuring diversity in effective numbers, it is

possible to partition diversity into alpha, beta, and gamma fractions (Leinster & Cobbold

2012; Botta-Dukát 2018) in the multiplicative way. Then, this multiplicative beta can be

rescaled using the formulae proposed by Chao et al. (2012). These indices behave consistently

only if abundances are taken into account as relative abundances.

Nearest neighbour approach

The earliest representatives of this family were shown by Clarke & Warwick (1998) and Izsák

& Prince (2001), then Ricotta & Burrascano (2008), and Ricotta & Bacaro (2010; see D_{CW}

and D_{IP} indices). Later Ricotta et al. (2016) introduced a new, general family called *PADDis*.

All these indices were primarily defined for presence-absence data type. The approach is

based on a re-definition of the *b* and *c* quantities of the 2×2 contingency table. Looking at

species as maximally different, and taking X and Y the two communities under comparison, *b*

can be viewed as the total uniqueness of community X. The uniqueness of a single species in

X is 1 if it is absent in *Y*, otherwise it is 0. Therefore, *b* is the sum of species uniqueness

values. However, from a functional perspective, the uniqueness of a species present only in *X*

should be between 0 and 1 if it is absent in *Y* but a similar species present there. Therefore, it

is possible to define the analogue of *b* which accounts for similarities between species:

Eq. 18. $\qquad\qquad B = \sum_{i \in X}\left(1 - \max_{j \in Y} \varepsilon_{ij}\right) = S_x - \sum_{i \in X} \max_{j \in Y} \varepsilon_{ij}$

The same logic applies for c, which is the uniqueness of community Y, where C expresses the

degree of uniqueness:

Eq. 19. $\qquad\qquad C = \sum_{i \in Y}\left(1 - \max_{j \in X} \varepsilon_{ij}\right) = S_y - \sum_{i \in Y} \max_{j \in X} \varepsilon_{ij}$

Ricotta et al. (2016) define the A quantify as follows:

Eq. 20. $\qquad\qquad A = a + (b - B) + (c - C)$

Having A, B, and C defined as analogues of a, b, and c, it is now possible to design trait-based

similarity measures following the logics of Jaccard, Sørensen, Sokal & Sneath, Kulczynski,

Ochiai and Simpson indices. It is notable that Ricotta et al. (2016) define A as a quantity that

ensures the components B and C to add up to $a + b + c$ but with no explicit biological

interpretation. Notably, D_{IP} and D_{CW} are identical with the Sørensen and Kulczynski forms of

PADDis. The generalization of D_{IP} and D_{CW} to relative abundances, $D_{CW}(Q)$, was also derived

by Ricotta & Bacaro (2010). For these two versions, it is not necessary to explicitly define the

A component. Using the relationships between Jaccard, Sørensen, Kulczynski, Ochiai and

Sokal & Sneath indices, from $D_{CW}(Q)$ it is theoretically possible to derive the extension of

PADDis to relative abundances; however, the biological interpretation of A remains dubious

in this framework.

Methods

The performance of FDissim indices can be reliably tested on data sets with known

background processes driving community assembly which is hardly possible to satisfy with

real data. Therefore, we compared the performance of FDissim indices using simulated data

sets. The data sets were generated using the *comm.simul* function of the comsimitv R package

(Botta-Dukát & Czúcz 2016, Botta-Dukát 2020). This function follows an individual-based

model for a meta-community comprising *N* communities and a regional pool of *S* species.

Local communities include *J* individuals, and are distributed equidistantly along a continuous

environmental gradient (with gradient values between 0 and 1). Each individual possesses

three traits: an 'environmental', a 'competitive' trait, and a neutral trait, all ranging on [0; 1].

Intraspecific variation in trait values is neglected in the simulation, that is, individuals

belonging to the same species are identical. The environmental trait defines the optimum of

the species along the environmental gradient. The closer the position of a community along

the environmental gradient to the environmental trait value of a species, the more suitable it is

for that species:

Eq. 21. $$suitability = \frac{(environment - environmental\ trait)^2}{\sigma}.$$

where σ (sigma) is adjustable so as to change the niche width of the species, and hence, the

length of the gradient (see later). The competitive trait represents the resource acquisition

strategy of the individual. The more similar the latter value between two individuals, the

higher the competition is between them, which means that intraspecific competition is the

strongest. The neutral trait has no effect on community assembly, thus it is not considered in

our study. The simulation starts with the random assignment of all individuals of all

communities to species. The second step is a 'disturbance' event, when one individual 'dies'

in each community. This individual is to be replaced by an offspring of other individuals

within the same community or those of other communities. Each individual produces one

offspring or does not reproduce. Probability of reproduction depends on the strength of

competition. The offspring remains in the same community or randomly disperses into any of

the other communities. Finally, the dead individual is replaced by one new individual from

the seeds produced and dispersed. The probability that an individual of a certain species

replaces the dead individual is defined by the number of seeds of that species and the

suitability of the habitat. Steps between the disturbance event and the establishment of a new

individual constitute a single 'generation'. Community composition is evaluated after lot of

generations. The strength of the environmental filtering can be adjusted by the sigma

parameter, respectively. When sigma is 0, all species are maximally specialist, which means

that they can occur only at the optimum point of the gradient (that is, at the exact value for the

environmental trait). If sigma is infinity, species are maximally generalist and all points along

the environmental gradient are equally suitable for them. Therefore, sigma is the parameter

which defines the suitability of each point of the gradient for each species based on its

distance from the respective optima. We generated data sets with sigma values 0.01, 0.1, 0.25,

0.5, 1 and 5 in order to simulate situations with different strength of environmental filtering.

The number of communities was 30, each community comprised 200 individuals, the number

of species in the species pool was 300, the simulation iterated for 100 generations, and we

allowed no intraspecific trait variation. For all the other parameters, we used the default

options.

However, it needed further explanation what real situations the six simulated levels of

environmental filtering represent. To provide a reference and assist interpretation, we

calculated two species-based beta-diversity measures, the multiplicative beta (Whittaker

1960) and the gradient length of the first axis of a detrended correspondence analysis (DCA)

ordination (Hill & Gauch 1980; Appendix S5, Fig. S5.1). The former gives the number of

distinct communities present in the total species pool of the gradient, while the latter is

minimal number of average niche breadths (also called turnover units) necessary for covering

the total gradient length. Moreover, we plotted the abundance of species in the sample units

along the gradient as a visual tool for assessing gradient length (Appendix S5, Fig. S5.2). All

these methods indicated that with sigma = 0.01 the gradient is extremely long: there are more

than 10 distinct communities and near 20 turnover units along the gradient. Samples with such

high beta diversity are very rare and special in real ecological research; therefore, findings

from simulations with sigma = 0.01 are mostly of theoretical importance. Beta diversity

values from sigma = 0.1 to sigma = 1 are more similar to real study situations, hence they

should be more relevant for practice. At sigma = 5, environmental filtering is practically not

operating, between-community variation is driven by interspecific relations and chance.

We calculated between-species dissimilarities as the Gower distance between their

environmental trait values which in this case equals the Euclidean distance scaled to [0; 1].

These distances had to be transformed to similarities according to the requirements of the

FDissim indices. Several formulae are available with which it is possible; however, they may

assume different functional relationships between similarity and distance. One of such

formulae we used is the linear transformation according to *Similarity = 1-Distance*. Besides

this, we also used *Similarity = $e^{-u \times Distance}$* which supposes a curvilinear function between

similarity and distance (Leinster & Cobbold 2012). With this exponential formula, it is

possible to weight the importance of small Gower distances between species relative to large

distances. With changing the parameter *u* it is possible to adjust how steeply similarity

decreases with increasing distance. We set *u* = 10 which leads to a relatively steep decline.

Although, after this transformation the minimal value for similarity is higher than zero, we

considered it negligibly low ($e^{-10} \approx 0.000045$) so we did not apply the transformation proposed

by Botta-Dukát (2018). For all FDissim indices where it was necessary we used the similarity

matrix or a dissimilarity matrix calculated as *Dissimilarity = 1-Similarity* as input. The

dissimilarity matrix is identical with the Gower distance matrix if the similarities were

calculated in a linear way, but in the other case, it keeps the exponential relationship between

distance and (dis-)similarity.

Dissimilarity matrices were calculated for the four community data sets with different sigma

values, with the two functions transforming Gower distances, and across a broad range of

available FDissim indices. For indices where absolute or relative abundances could have been

taken into account, we opted for relative abundance for the sake of better comparability. With

generalized_Tradidiss, we calculated the 'even' and the 'uneven' weighting versions. The

entire analysis was run with abundance and presence/absence data. Some FDissim indices are

only suitable for binary data, thus the number of indices applied for relative abundance and

binary data were 25 and 31, respectively. In cases of indices handling both data types, we

used exactly the same version of the index as with abundance data, hence communities with

different numbers of species were given equal weight due to division by community totals.

Additionally, dissimilarity matrices were also calculated using the Bray-Curtis index (for

binary data: Sørensen index in dissimilarity form) to provide a contrast against the case

disregarding between-species dissimilarities.

Then for each dissimilarity matrices, we conducted two types of analyses. Firstly, we

compared how strongly the dissimilarity indices correlate with the environmental distance

using Kendall tau rank correlation. This gives an estimate of how well a dissimilarity index

reveals the monotonic relationship between trait composition of local communities and the

environmental gradient. We visually assessed the shape of relationship between dissimilarity

and environmental distance in the case of lowest sigma (i.e., longest gradient) when the

distortion of linear relationship between the two is supposed to be the strongest. Then, to

disentangle the effects of different methodological decisions and the sigma parameter on the

correlation between FDissim indices and environmental distance we calculated a random

forest model. In this model the dependent variable was the Kendall tau correlation coefficient,

while the independent variables were the sigma, the data type (abundance vs.

presence/absence), the transformation method for Gower distances (linear vs. exponential),

and the FDissim method. Within approaches FDissim methods often strongly correlated that

resulted in very similar Kendall's tau values. Therefore, only the Sørensen/Bray-Curtis

versions of *dsimcom*, *dissABC*, *PADDis*/*D_CW*, *generalized_Tradidiss* with uneven weights, as

well as $\beta_{turnover}$, *CWMdis*, and the *CDFdis* were included into this analysis. Variable

importance scores (VIS) in the random forest were estimated by the permutation approach

based on mean decrease in log-likelihood using the *varimp* function of the *partykit* package.

The effects of the model terms were also illustrated by heat-maps.

All statistical analyses were done in R (R Core Team 2019) using the *FD* (Laliberté &

Legendre 2010, Laliberté et al. 2014), *adiv* (Pavoine 2020a,b), *comsimitv* (Botta-Dukát 2020,)

vegan (Oksanen et al. 2019), *DescTools* (Signorell et al. 2020), *partykit* (Hothorn et al. 2006,

Strobl et al. 2007, Strobl et al. 2008, Hothorn & Zeileis 2015) packages.

Results

Kendall tau correlation coefficients decreased as the strength of environmental filtering

decreased (that is, with increasing sigma) in all examined cases. For FDissim indices which

handled both data types, presence/absence data resulted in lower correlations than abundance

data for all indices. For most indices, this difference was highest at intermediate values for

sigma. These trends were consistent between the linear and the exponential transformations.

Correlations for all indices at all sigma values with linear transformation are shown in Table 3

for abundances data and in Table 4 for presence/absence data.

In most simulation scenarios, the FDissim indices correlated more strongly with the

environmental gradient than the species-based Bray-Curtis index. However, in several

occasions, indices belonging to the nearest neighbour family performed poorer than the

species-based dissimilarity. Notably, at the highest sigma and with presence/absence data, all

indices showed correlation near to zero but among them the Bray-Curtis index had the highest

correlation with environmental distance.

As expected, we found perfect rank correlations among Jaccard, Sørensen, Sokal-Sneath and

Whittaker's beta versions of *dsimcom*, among Jaccard, Sørensen and Sokal-Sneath forms of

dissABC, between D_{IP} and Sørensen form of *PADDis* (only for presence-absence data),

between D_{CW} and Kulczynski form of *PADDis* (only for presence-absence data), and between

D_{IP} and D_{CW} (for abundance data type).

Dissimilarity indices showed various shapes of relationship with environmental distance

(Appendix S4). At strongest environmental filtering, all FDissim indices had dissimilarity

values near zero at minimal environmental distance, only the species-based Bray-Curtis which

had dissimilarity was near 0.4 at the smallest environmental distances. In case of linear

transformation of Gower distances and presence/absence data, approximately linear

relationship was found for *CWMdis*, *CDFdis*, D_Q, Sørensen and Ochiai forms of *dsimcom*,

Jaccard form of *dissABC*, Marczewski-Steinhaus form of *generalized_Tradidiss* with both

weighting versions, $\beta_{heterogeneity}$ and $\beta_{segregation}$; although, most other indices showed only a

small degree of distortion of linear function (Figure S4.1). Exponential relationship was found

for the evenness-based (PE) form of *generalized_Tradidiss*. Notably, the taxon-based Bray-

Curtis index had the steepest asymptotic function among all. In case of exponential

transformation all other indices relying on between-species dissimilarities showed an

asymptotic curve (Figure S4.2).

In the random forest, niche width (that is, sigma) acquired by far the highest variable

importance score (VIS=0.114). The less important variables were the data type (VIS=0.0176),

the dissimilarity method (VIS=0.0037) and the transformation (VIS=-0.00001). The heat map

(Figure 1) also revealed a strong decrease in correlation along increasing sigma. It is also

clearly shown that in most cases abundance data resulted in significantly higher correlation

than presence/absence. The difference between linear and exponential transformation methods

was not always visible. Regarding variation between dissimilarity indices, the most striking

patterns were the relatively poor performance of the *PADDis*/D_{CW} indices. All but the latter

index combined with abundance data and linear transformation of dissimilarities lead to the

highest correlation with environmental distance.

Discussion

General patterns in the correlation with environmental distance

We ran different simulation scenarios with varying strength of environmental filtering. We

expected that the correlation between FDissim indices and environmental distance to be the

highest when the environmental filtering is the strongest, and the correlation to become

neutral when environmental filtering is not effective. When environmental filtering was

strongest (that is, minimal overlap of species niches along the environmental gradient), all

FDissim indices correlated highly with the environmental gradient. As expected, correlation

between trait dissimilarity and environmental distance decreased as filtering weakened,

moreover, differences between families of indices became more apparent. This result suggests

that all tested methods are able to reveal the strong environmental filtering processes.

As the contribution of competitive exclusion and stochastic processes approach or override

environmental filtering, the correlation between FDissim indices and the background gradient

becomes weaker. This decrease itself is not a drawback of the FDissim methods, rather it is a

consequence of our study design, since we applied a series of scenarios where the effect of

niche filtering was decaying. However, we think that the degree of the decrease reflects the

sensitivity of the FDissim indices to the underlying trait-environmental relationship. Indices,

which showed high correlation with environmental distance, could be capable of revealing the

environmental signal even when it is weak. Actually, in our tests, most indices reached

similarly high correlation, and there were only a few combinations of simulation parameters

which resulted in a decreased correlation with environmental distance for some dissimilarity

indices.

Determinants of the correlation based on the random forest model

The random forest model revealed that the effect of gradient length is the most important

determinant of the correlation between dissimilarity and environmental distance, while

methodological decisions had much lower variable importance. These observations suggest

that the absolute value of the correlation between dissimilarity and environmental distance is

primarily dependent on the sample in hand, and can be influenced by methodological

decisions to a limited extent.

Correlations were stronger with abundance than with presence/absence data. This finding is at

least partly attributable to our simulation design where community composition was driven by

individual-based processes: birth, fitness difference, reproduction, and death. As a result,

species relative abundances had to be proportional with their environmental suitability in the

local community. Transforming such data to binary scale loses meaningful information and

weakens the correlation between dissimilarity in trait composition and environmental

background. In cases when presences and absences of species respond more robustly to the

main environmental gradient, while relative abundances change stochastically, or abundance

estimations are inaccurate, the binary data type might be more straightforward.

Transforming between-species dissimilarities has a potential to conform distributional

requirements, to approximate expert intuitions about relatedness of species or to customize

sensitivity to functional difference with respect to specific research aims. For most indices

across the tested range of gradient length and data type, the exponential transformation

resulted a somewhat lower correlation than with linear transformation. More insight is

provided by examining the shape of the relationships besides the pure correlation value. After

linear transformation of Gower distances, most dissimilarity indices showed a linear or

slightly curved function along environmental distance; although the scatter of the evenness-

based *generalized_Tradidiss* differed considerably from the straight line towards an

exponentially increasing one. After exponential transformation of between-species trait

dissimilarities, all indices in the direct dissimilarity-based class showed a rather steeply

increasing asymptotic function. This result suggests that with the exponential transformation

of between-species dissimilarities, it is possible to make FDissim indices more sensitive to

smaller differences in functional composition. Certainly, summary-based indices (*CWMdis*,

CDFdis) are not affected by this transformation, since they are not based on between-species

dissimilarities.

Comparison of taxon-based vs. trait-based dissimilarity

The basic assumption of functional ecology is that the traits of individuals should be in closer

relationship with ecological properties than their taxonomical status. Following this argument,

we expected that trait-based dissimilarity measures correlate more strongly with the

environmental background than species-based indices. In contrast, higher correlation of

species-based dissimilarity than trait-based dissimilarity indicates loss of information with the

introduction of between-species similarity – which is non-sensual since our data was

simulated in a way to possess a strong pattern in trait-environment relationship. We used the

Sørensen/Bray-Curtis index in a dissimilarity form as a reference method representing

species-based dissimilarity calculations disregarding traits. Our expectation was fulfilled by

all indices with the exception of the members of the nearest neighbour family (D_{IP}, D_{CW} and

PADDis). We suspect two potential reasons behind the low performance of these latter groups

of indices. The first one is the improper scaling factor used for standardizing the 'operational

part' of the indices (see the description in of the *PADDis* family and the discussion about it

under the paragraph "Within-family variation of indices"). Second, these indices rely on the

quantities of minimally different species in the two communities under comparison. However,

the minimum is a less robust descriptor of any sample distribution because of its dependency

on sampling error; therefore, it might provide a poor representation of total community

dissimilarity.

Although, we did not include dissimilarity values at exactly zero distance, the y-intercept (also

called 'nugget') of the dissimilarity vs. environmental distance functions can be extrapolated

with negligible error (Fortin & Dale 2005). Brownstein et al. (2012) argued that the nugget of

the distance decay relationship is a direct estimate of the amount of chance in the variation

between local communities. In this respect worth noting is that the nugget with species-based

Bray-Curtis index was near 0.4, while with all trait-based indices the nugget was near zero.

This suggests that without accounting for species similarities, environmental distance between

communities can be overestimated due to similar species replacing each other.

Within-family variation of indices

The perfect correlation between Jaccard, Sørensen and Sokal-Sneath forms of *dsimcom* and

dissABC families was expected, since the original, taxon-based Jaccard, Sørensen and Sokal-

Sneath indices are algebraically related, too (Janson & Vegelius 1981). However, for *PADDis*

Jaccard, Sørensen and Sokal-Sneath forms showed correlation below 1. At this family, the *B*

and *C* components of the 2×2 contingency table are defined as measurable quantities with

clear interpretation: the sum of species uniqueness values within each community. The total

diversity ($A+B+C$) is defined to be equal with the species richness of the pooled pair of

communities ($a+b+c$), and the quantity A is derived by subtracting ($B+C$) from it. With this

definition, A remains a virtual quantity with no biological interpretation. In PADDis indices,

trait-based quantities *B* and *C* appear in the numerator (the 'operational part' sensu Ricotta et

al. 2016) of the indices, while in the denominators (i.e., in the 'scaling factor') the taxon-

based quantities, *a*, *b* and *c* are used. We argue that the inconsistent behaviour of *PADDis* is

due to the application of taxon-based quantities for scaling factors of trait-based operational

parts. At the same time, we acknowledge that we either see no obvious solution to define total

diversity or shared diversity according to the uniqueness-based idea behind *PADDis* in a more

realistic way. In the *generalized_Tradidiss* family, the trait-based analogue of Bray-Curtis

index can be achieved by calculating generalized Canberra distance with uneven weighting of

species. We expected this to be perfectly correlated with Marczewski-Steinhaus form of

generalized_Tradidiss index with uneven weighting, since Bray-Curtis and Marczewski-

Steinhaus indices are the abundance forms of Sørensen and Jaccard indices, respectively.

However, the correlation between them was lower. In the *generalized_Tradidiss* family,

between-community dissimilarity is calculated as weighted sum a standardized differences in

species ordinariness values. Species ordinariness is calculated on the basis of species

abundance and trait values; however, weights used for adjusting species-level contributions

are derived solely from abundances. Therefore, *generalized_Tradidiss* also follows a 'hybrid'

approach in accounting for taxon-based vs. trait-based information. We argue that this is the

reason why the algebraic relationships between the original Sørensen and Jaccard indices does

not apply to its Sørensen/Bray-Curtis-type and Jaccard/Marczewski-Steinhaus-type forms. To

sum up, we point to our observation that Jaccard, Sørensen and Sokal-Sneath forms of certain

families of indices do not satisfy the algebraic relationships they supposed to, opening space

for potential confusion. These algebraic relations hold only if A, B and C quantities are

explicitly and consistently defined.

Families of FDissim indices combine abundance difference of species between plots and

interspecific trait differences in a unique way, while indices belonging to the same family

differ in how they relate this amount of 'unshared' variation (summarized as the b and c

portions of the contingency table) to the shared (a) variation. Some indices are able to handle

abundances either as absolute or relative abundance (e.g. *dsimcom*, *generalized_Tradidiss*,

dissABC), while others divide absolute abundances by their sum over the respective

community, thus they work only with relative abundances. When indices in the former group

are set to consider absolute abundances, they become sensitive to variation in the summed

abundances of the communities under comparison. To place our tests on a common ground,

we simulated communities with equal total number of individuals, and set all indices, where

relevant, to work with relative abundances. Hence, we removed the effect of differences in

total abundance. The constant number of individuals might have increased the similarity

between FDissim indices belonging to the same family and the correlation with the

environmental gradient. The sum of abundances, let them be measured on any quantitative

scale, may vary considerably in real study situations due to aggregated distribution of

individuals or uneven sampling effort. Therefore, our findings are more likely valid for

settings when the sum of abundances are relatively stable, e.g. when sampling effort is

controlled and individuals are dispersed evenly, or when abundances are recorded on

percentage scale.

Limitations of our study

In our study, we simulated a research situation in a simplistic way. We applied only one

environmental gradient which operated as an environmental filter driving convergence on a

single trait. Besides this, we applied another trait which was constantly affected by a low level

of competitive exclusion. These two traits were uncorrelated. Nevertheless, there was some

effect of random drift on community composition due to the probabilistic components of the

simulation algorithm. We varied the strength of environmental filtering thus it had different

relative contribution compared with competitive exclusion and stochasticity. In real research

situations local trait composition is influenced by a wide range of processes, including several

abiotic and biotic filters acting simultaneously. Unless they are manipulated as parts of an

experimental system, the full set of such filters are usually unknown for the researchers. The

multiplicity of filters may reduce the ability of FDissim indices in recovering trait-

environment relationships. Further research should clarify how increasing complexity of the

sample affects the behaviour of FDissim indices.

Conclusions

Considering the diversity of concepts they are built upon, FDissim indices showed

unexpectedly low variation in performance. CWMdis, dsimcom, generalized_Tradidiss

acquired the highest correlation with environmental distance in all simulation scenarios,

therefore they seem to be equally suitable for quantifying pairwise beta diversity based on

traits. Nevertheless, the most important determinant of the matching between trait-based

dissimilarity and environmental distance is the length of the trait gradient. Besides this, the

data type (presence/absence vs. abundance) also affected the correlation more strongly than

the choice of FDissim method. Extending the comparative tests of FDissim measure to more

complex gradients and real data sets could offer further insight into their behaviour.

Data availability

Simulated data was generated using the comsimitv R package. Own functions for functional

dissimilarity indices are made available through the Zenodo public repository:

10.5281/zenodo.4323590.

Author contributions

A.L. designed and carried out the analysis, lead writing, Z.B.D. discussed the concept and the

results, wrote parts of and commented on the manuscript.

References

Anderson, M. J., Crist, T. O., Chase, J. M., Vellend, M., Inouye, B. D., Freestone, A. L.,

Sanders, N. J., Cornell, H. V., Comita, L. S., Davies, K. F., Harrison, S. P., Kraft, N. J. B.,

Stegen, J. C. & Swenson, N. G. (2011). Navigating the multiple meanings of β diversity: a

roadmap for the practicing ecologist. Ecology Letters, 14(1), 19-28. doi:10.1111/j.1461-

0248.2010.01552.x

Anderson, M. J., Ellingsen, K. E. & McArdle, B. H. (2006). Multivariate dispersion as a

measure of beta diversity. Ecology Letters, 9(6), 683-693. doi:10.1111/j.1461-

0248.2006.00926.x

Baselga, A. & Leprieur, F. (2015). Comparing methods to separate components of beta

diversity. Methods in Ecology and Evolution, 6: 1069-1079. doi:10.1111/2041-210X.12388

Botta Dukát, Z. & Czúcz, B. (2016). Testing the ability of functional diversity indices to

detect trait convergence and divergence using individual based simulation. Methods in

Ecology and Evolution, 7, 114-126. https://doi.org/10.1111/2041-210X.12450

Botta Dukát, Z. (2005). Rao's quadratic entropy as a measure of functional diversity based

on multiple traits. Journal of Vegetation Science, 16, 533-540. https://doi.org/10.1111/j.1654-

1103.2005.tb02393.x

Botta Dukát, Z. (2018). The generalized replication principle and the partitioning of

functional diversity into independent alpha and beta components. Ecography, 41: 40-50.

doi:10.1111/ecog.02009

Botta-Dukat, Z. (2020). comsimitv: Flexible Framework for Simulating Community

Assembly. R package version 0.1.4. https://CRAN.R-project.org/package=comsimitv

Brownstein, G., Steel, J.B., Porter, S., Gray, A., Wilson, C., Wilson, P.G. & Wilson, J. B.

(2012). Chance in plant communities: a new approach to its measurement using the nugget

from spatial autocorrelation. Journal of Ecology, 100, 987-996.

https://doi.org/10.1111/j.1365-2745.2012.01973.x

Cardoso, P., Rigal, F., Carvalho, J.C., Fortelius, M., Borges, P.A.V., Podani, J. & Schmera, D.

(2014). Partitioning taxon, phylogenetic and functional beta diversity into replacement and

richness difference components. Journal of Biogeography, 41, 749-761.

doi:10.1111/jbi.12239

Carmona, C. P., de Bello, F., Mason, N. W. H., Lepš, J. (2016). Traits without borders:

Integrating functional diversity across scales. Trends in Ecology and Evolution 31(5), 382-

394. doi: 10.1016/j.tree.2016.02.003

Champely, S., Chessel, D. (2002). Measuring biological diversity using Euclidean metrics.

Environmental and Ecological Statistics 9, 167–177.

https://doi.org/10.1023/A:1015170104476

Chao, A., Chiu, C. and Hsieh, T.C. (2012). Proposing a resolution to debates on diversity

partitioning. Ecology, 93, 2037-2051. https://doi.org/10.1890/11-1817.1

Chao, A., Chiu, C. H., Villéger, S., Sun, I F., Thorn, S., Lin, Y. C., Chiang, J. M., &

Sherwin, W. B. (2019). An attribute diversity approach to functional diversity, functional

beta diversity, and related (dis)similarity measures. Ecological Monographs, 89(2), e01343.

10.1002/ecm.1343

Chiu, C.-H., Jost, L. & Chao, A. (2014). Phylogenetic beta diversity, similarity, and

differentiation measures based on Hill numbers. Ecological Monographs, 84(1), 21-44.

Clarke, K.R. & Warwick, R.M. (1993). Quantifying structural redundancy in ecological

communities. Oecologia, 113(2), 278-289.

De Bello, F., Carmona, C.P., Mason, N.W.H., Sebastià, M. T. and Lepš, J. (2013). Which

trait dissimilarity for functional diversity: trait means or trait overlap? Journal of Vegetation

Science, 24, 807-819. doi:10.1111/jvs.12008

De Bello, F., Lepš, J., Lavorel, S., & Moretti, M. (2007). Importance of species abundance for

assessment of trait composition: an example based on pollinator communities. Community

Ecology, 8(2), 163–170. https://doi.org/10.1556/ComEc.8.2007.2.3

Díaz, S., & Cabido, M. (2001). Vive la différence: plant functional diversity matters to

ecosystem processes. Trends in Ecology and Evolution, 16(11), 646–655.

https://doi.org/10.1016/S0169-5347(01)02283-2

Faith, D. P., Minchin, P. R. & Belbin, L. (1987). Compositional dissimilarity as a robust

measure of ecological distance. Vegetatio 69, 57-68.

Fortin, M. J. & Dale, M.R.T. (2005). Spatial Data Analysis: a Guide for Ecologists.

Cambridge University Press, Cambridge.

Garnier, E., Cortez, J., Billès, G., Navas, M., Roumet, C., Debussche, M., Laurent, G.,

Blanchard, A., Aubry, D., Bellmann, A., Neill, C. & Toussaint, J. (2004). Plant functional

markers capture ecosystem properties during secondary succession. Ecology, 85, 2630-2637.

doi:10.1890/03-0799

Gregorius, H. R., Gillet, E.M. & Ziehe, M. (2003). Measuring Differences of Trait

Distributions Between Populations. Biometrical Journal, 45, 959-973.

https://doi.org/10.1002/bimj.200390063

Grime, J. P. (1998). Benefits of plant diversity to ecosystems: immediate, filter and founder

effects. Journal of Ecology, 86, 902–910.

Hawkins, B.A., Leroy, B., Rodríguez, M.Á., Singer, A., Vilela, B., Villalobos, F., Wang, X.

& Zelený, D. (2017). Structural bias in aggregated species☐level variables driven by repeated

species co☐occurrences: a pervasive problem in community and assemblage data. Journal of

Biogeography, 44, 1199-1211.

Hérault, B., & Honnay, O. (2007). Using life-history traits to achieve a functional

classification of habitats. Applied Vegetation Science, 10(1), 73–80.

https://doi.org/10.1111/j.1654-109X.2007.tb00505.x

Hill, M. O. & Gauch, H. G. (1980). Detrended Correspondence Analysis: An Improved

Ordination Technique. Vegetatio, 42, 47–58.

Hill, M. O. (1973). Diversity and evenness: a unifying notation and its consequences.

Ecology, 54(2), 427–432.

Hitchcock, F.L. (1941). Distribution of a product from several sources to numerous localities.

Journal of Mathematical Physics, 20: 224-230.

Hothorn, T., Hornik, K., Van de Wiel, M. A. & Zeileis, A. (2006). A Lego System for

Conditional Inference. The American Statistician, 60(3), 257–263.

Hothorn, T., Zeileis, A. (2015). partykit: A Modular Toolkit for Recursive Partytioning in R.

Journal of Machine Learning Research, 16, 3905-3909. URL

http://jmlr.org/papers/v16/hothorn15a.html

Izsák, C., & Price. R. G. (2001). Measuring b-diversity using a taxonomic similarity index,

and its relation to spatial scale. Marine Ecology Progress Series 215, 69–77.

Janson, S. & J. Vegelius (1981). Measures of ecological association. Oecologia, 49(3), 371-

376.

Jost, L. (2007). Partitioning diversity into independent alpha and beta components. Ecology,

88, 2427–2439.

Kleyer, M., Dray, S., Bello, F., Lepš, J., Pakeman, R.J., Strauss, B., Thuiller, W. & Lavorel,

S. (2012). Assessing species and community functional responses to environmental gradients:

which multivariate methods? Journal of Vegetation Science, 23, 805-821. doi:10.1111/j.1654-

1103.2012.01402.x: 1199–1211.

Koleff, P., Gaston, K. J. & Lennon, J. J. (2003). Measuring beta diversity for presence–

absence data. Journal of Animal Ecology, 72, 367-382. doi:10.1046/j.1365-

2656.2003.00710.x

Laliberté, E. & P. Legendre (2010). A distance-based framework for measuring functional

diversity from multiple traits. Ecology, 91, 299-305.

Laliberté, E., Legendre, P., & Shipley, B. (2014). FD: measuring functional diversity from

multiple traits, and other tools for functional ecology. R package version 1.0-12.

Legendre, P. & Legendre, L. (1998) Numerical ecology. Elsevier, Amsterdam, NL

Legendre, P., De Cáceres, M. (2013). Beta diversity as the variance of community data:

dissimilarity coefficients and partitioning. Ecology Letters 16, 951–963

Leinster, T. & Cobbold, C.A. (2012). Measuring diversity: the importance of species

similarity. Ecology, 93, 477-489. doi:10.1890/10-2402.1

Lengyel, A. & Podani, J. (2015). Assessing the relative importance of methodological

decisions in classifications of vegetation data. Journal of Vegetation Science, 26, 804-815.

doi:10.1111/jvs.12268

Lengyel, A., Swacha, G., Botta-Dukát, Z. & Kacki, Z. (2020). Trait-based numerical

classification of mesic and wet grasslands in Poland. Journal of Vegetation Science, 31, 319–

330. https://doi.org/10.1111/jvs.12850

Lepš, J., de Bello, F., Lavorel, S. & Berman, S. (2006). Quantifying and interpreting

functional diversity of natural communities: practical considerations matter. Preslia, 78, 481–

501.

MacArthur, R., Levins, R. (1967). Limiting similarity convergence and divergence of

coexisting species. American Naturalist, 101, 377–387.

Mason, N. W. H., Mouillot, D., Lee, W. G. & Wilson, J. B. (2005). Functional richness,

functional evenness and functional divergence: the primary components of functional

diversity. Oikos, 111, 112-118. doi:10.1111/j.0030-1299.2005.13886.x

McGill, B., Enquist, B. J., Weiher, E., Westoby, M. (2006). Rebuilding community ecology

from functional traits. Trends in Ecology and Evolution 21(4), 178-185.

Mouchet, M.A., Villéger, S., Mason, N.W.H. and Mouillot, D. (2010). Functional diversity

measures: an overview of their redundancy and their ability to discriminate community

assembly rules. Functional Ecology, 24, 867-876. doi:10.1111/j.1365-2435.2010.01695.x

Mouillot, D., Stubbs, W., Faure, M., Dumay, O., Tomasini, J.A., Wilson, J.B. & Chi, T.D.

(2005). Niche overlap estimates based on quantitative functional traits: a new family of

non□parametric indices. Oecologia, 145, 345–353.

Muscarella, R. & Uriarte, M. (2016). Do community-weighted mean functional traits reflect

optimal strategies? Proceedings of the Royal Society B, 283, 20152434.

https://doi.org/10.1098/rspb.2015.2434

Nipperess, D.A., Faith, D.P. & Barton, K. (2010), Resemblance in phylogenetic diversity

among ecological assemblages. Journal of Vegetation Science, 21, 809-820.

doi:10.1111/j.1654-1103.2010.01192.x

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Peter R.

Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. M., Szoecs, E. &

Wagner, H. (2019). vegan: Community Ecology Package. R package version 2.5-6.

https://CRAN.R-project.org/package=vegan

Pavoine, S. & Ricotta, C. (2014). Functional and phylogenetic similarity among communities.

Methods in Ecology and Evolution, 5, 666--675.

Pavoine, S. & Ricotta, C. (2019). Measuring functional dissimilarity among plots: Adapting

old methods to new questions. Ecological Indicators, 97, 67-72.

Pavoine, S. (2012). Clarifying and developing analyses of biodiversity: towards a

generalisation of current approaches. Methods in Ecology and Evolution, 3, 509-518.

doi:10.1111/j.2041-210X.2011.00181.x

Pavoine, S. (2016). A guide through a family of phylogenetic dissimilarity measures among

sites. Oikos, 125, 1719-1732. doi:10.1111/oik.03262

Pavoine, S. (2020). adiv: An R package to analyse biodiversity in ecology. Methods in

Ecology and Evolution, 11, 1106– 1112. https://doi.org/10.1111/2041-210X.

Peres-Neto, P.R., Dray, S. & ter Braak, C.J.F. (2017). Linking trait variation to the

environment: critical issues with community☐weighted mean correlation resolved by the

fourth☐corner approach. Ecography, 40, 806-816.

Petchey, O. L. & Gaston, K. J. (2006). Functional diversity: back to basics and looking

forward. Ecology Letters, 9, 741-758. doi:10.1111/j.1461-0248.2006.00924.x

Podani, J. & Schmera, D. (2011). A new conceptual and methodological framework for

exploring and explaining pattern in presence – absence data. Oikos, 120, 1625-1638.

doi:10.1111/j.1600-0706.2011.19451.x

Podani, J. (2000). Introduction to the exploration of multivariate biological data. Backhuys,

Leiden, NL.

R Core Team (2019). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Rao, C. R. (1982). Diversity and dissimilarity coefficients: a unified approach. Theoretical

Population Biology, 21, 24-43.

Ricotta C. & Burrascano S. (2008). Beta diversity for functional ecology. Preslia, 80, 61–71.

Ricotta, C. & G. Bacaro. (2010). On plot-to-plot dissimilarity measures based on species

functional traits. Community Ecology, 11, 113–119.

Ricotta, C. & J. Podani. (2017). On some properties of the Bray-Curtis dissimilarity and their

ecological meaning. Ecological Complexity, 31, 201–205.

Ricotta, C. & Pavoine, S. (2015). Measuring similarity among plots including similarity

among species: an extension of traditional approaches. Journal of Vegetation Science, 26,

1061-1067. doi:10.1111/jvs.12329

Ricotta, C. (2017). Of beta diversity, variance, evenness, and dissimilarity. Ecology and

Evolution 7, 4835– 4843. https://doi.org/10.1002/ece3.2980

Ricotta, C. (2018). A family of (dis)similarity measures based on evenness and its relationship

with beta diversity. Ecological Complexity, 34, 69-73. DOI: 10.1016/j.ecocom.2018.03.002

Ricotta, C., Bacaro, G., Caccianiga, M., Cerabolini, B.E.L. & Moretti, M. (2015). A classical

measure of phylogenetic dissimilarity and its relationship with beta diversity. Basic and

Applied Ecology 16(1), 10-18. https://doi.org/10.1016/j.baae.2014.10.003

Ricotta, C., Podani, J., Pavoine, S. (2016). A family of functional dissimilarity measures for

presence and absence data. Ecology and Evolution, 6, 5383–5389. DOI: 10.1002/ece3.2214

Schmidt, T., Matias Rodrigues, J. & von Mering, C. (2017). A family of interaction-adjusted

indices of community similarity. ISME Journal 11, 791–807.

https://doi.org/10.1038/ismej.2016.139

Signorell, A. et mult. al. (2020). DescTools: Tools for descriptive statistics. R package version

0.99.38.

Strobl, C., Boulesteix, A.L., Kneib, T., Augustin, T. & Zeileis, A. (2008). Conditional

Variable Importance for Random Forests. BMC Bioinformatics, 9(307).

http://www.biomedcentral.com/1471-2105/9/307

Strobl, C., Boulesteix, A.L., Zeileis, A. & Hothorn, T. (2007). Bias in Random Forest

Variable Importance Measures: Illustrations, Sources and a Solution. BMC Bioinformatics, 8,

25. http://www.biomedcentral.com/1471-2105/8/25

Swenson N. G., Anglada-Cordero P. & Barone J. A. (2011). Deterministic tropical tree

community turnover: evidence from patterns of functional beta diversity along an elevational

gradient. Proceedings of the Royal Society B, 278, 877–884.

Swenson, N. G. (2011). Phylogenetic Beta Diversity Metrics, Trait Evolution and Inferring

the Functional Beta Diversity of Communities. PLoS ONE 6(6), e21264.

https://doi.org/10.1371/journal.pone.0021264

Tamás, J., Podani, J. & Csontos, P. (2001). An extension of presence/absence coefficients to

abundance data: a new look at absence. Journal of Vegetation Science, 12, 401-410.

doi:10.2307/3236854

Tuomisto, H. (2010a). A diversity of beta diversities: straightening up a concept gone awry.

Part 1. Defining beta diversity as a function of alpha and gamma diversity. Ecography, 33, 2-

22. doi:10.1111/j.1600-0587.2009.05880.x

Tuomisto, H. (2010b). A diversity of beta diversities: straightening up a concept gone awry.

Part 2. Quantifying beta diversity and related phenomena. Ecography, 33, 23-45.

doi:10.1111/j.1600-0587.2009.06148.x

Villéger, S., Mason, N.W.H. & Mouillot, D. (2008). New multidimensional functional

diversity indices for a multifaceted framework in functional ecology. Ecology, 89, 2290-2301.

doi:10.1890/07-1206.1

Violle, C., Navas, M.␣L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I. & Garnier, E.

(2007). Let the concept of trait be functional! Oikos, 116, 882-892. doi:10.1111/j.0030-

1299.2007.15559.x

Whittaker, R. H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California.

Ecological Monographs, 30, 280–338.

Whittaker, R. H. (1972). Evolution and measurement of species diversity. Taxon, 21, 213-

251.doi:10.2307/1218190

Zelený, D. (2018). Which results of the standard test for community weighted mean approach

are too optimistic? Journal of Vegetation Science 29, 953-966.

Tables and Figures

Table 1. Similarity and dissimilarity forms of resemblance indices for presence-absence data

Name of the index	Similarity version	Dissimilarity version
Sørensen	$s_S = \dfrac{2a}{2a + b + c} = \dfrac{a}{(S_j + S_k)/2}$	$d_S = \dfrac{b + c}{2a + b + c} = \dfrac{b + c}{S_j + S_k}$
Ochiai	$s_O = \dfrac{a}{\sqrt{(a + b)(a + c)}} = \dfrac{a}{\sqrt{S_j S_k}}$	$d_O = \dfrac{b + c}{\sqrt{S_j S_k}}$
Kulczynski	$s_K = \dfrac{1}{2}\left(\dfrac{a}{a + b} + \dfrac{a}{a + c}\right) = \dfrac{a}{2/(1/S_j + 1/S_k)}$	$d_K = \dfrac{1}{2}\left(\dfrac{b}{a + b} + \dfrac{c}{a + c}\right) = \dfrac{1}{2}\left(\dfrac{b}{S_j} + \dfrac{c}{S_k}\right)$
Simpson	$s_{Si} = \dfrac{a}{a + \min(b, c)} = \dfrac{a}{\min(S_j, S_k)}$	$d_{Si} = \dfrac{b + c}{\min(S_j, S_k)}$
Jaccard	$s_J = \dfrac{a}{a + b + c} = \dfrac{a}{S_{jk}}$	$d_J = \dfrac{b + c}{a + b + c} = \dfrac{b + c}{S_{jk}}$
Sokal & Sneath	$s_{SS} = \dfrac{a}{a + 2(b + c)}$	$d_{SS} = \dfrac{2(b + c)}{a + 2(b + c)}$

975   Table 2. Classification of trait-based dissimilarity indices. In columns of input data type X-es indicate, if abundance (A), relative abundance (R),

976   and presence-absence data can be used as input.

Class	Approach	Family	References	Input Data tpye			R function
				A	R	P/A	
Summary-based	*Typical value*	*CWM-based*	Ricotta et al. (2015)	X	X	X	FD:::functcomp
	Distribution-based	*CDF-based*	Appendix S3	X	X	X	our new functions, see Data availability
Direct dissimilarity	*Probabilistic*	*DISC/D_Q*	Rao 1982, Pavoine & Ricotta (2014)	X	X	X	adiv::SQ
		dsimcom	Pavoine & Ricotta (2014)	X	X	X	adiv:::dsimcom
	Ordinariness-based	*dissABC*	Pavoine & Ricotta (2015)	X	X	X	adiv:::dissABC
		generalized_Tradidiss	Pavoine & Ricotta (2019)	X	X		adiv:::generalized_Tradidiss
	Diversity	*multiplicative beta*	Chao et al. (2012)		X		our new functions, see Data

partitioning						availability
Nearest neighbour	D_{CW}, $D_{CW}(Q)$	Clarke & Warwick (1998), Ricotta & Bacaro (2010)		X	X	our new functions, see Data availability
	D_{IP}	Izsák & Prince (2001), Ricotta & Bacaro (2010)		X	X	our new functions, see Data availability
	PADDis	Ricotta et al. (2016)			X	adiv:::PADDis
Classification-based	not discussed	not discussed	Hérault & Honnay (2007), Nipperess et al. (2010), Cardoso et al. (2014), Pavoine (2016)			

978   Table 3. Kendall tau correlations between environmental distance and the functional

979   dissimilarity measures at different values of sigma and with abundance data type

	Sigma=0.01	Sigma=0.1	Sigma=0.25	Sigma=0.5	Sigma=1	Sigma=5
CWMdis	0.974	0.905	0.846	0.828	0.649	0.251
CDFdis	0.974	0.904	0.845	0.83	0.646	0.255
D(Q)	0.974	0.912	0.832	0.828	0.637	0.243
dsimcom.SS	0.974	0.911	0.832	0.829	0.638	0.243
dsimcom.Jac	0.974	0.911	0.832	0.829	0.638	0.243
dsimcom.Sor	0.974	0.911	0.832	0.829	0.638	0.243
dsimcom.Och	0.974	0.911	0.832	0.829	0.639	0.243
dsimcom.Beta	0.974	0.911	0.832	0.829	0.638	0.243
dissABC.Jac	0.967	0.899	0.82	0.813	0.617	0.243
dissABC.Sor	0.967	0.899	0.82	0.813	0.617	0.243
dissABC.SS	0.967	0.899	0.82	0.813	0.617	0.243
dissABC.Och	0.968	0.899	0.819	0.814	0.618	0.243
dissABC.Kul	0.968	0.898	0.819	0.814	0.619	0.243
dissABC.Si	0.954	0.867	0.789	0.791	0.616	0.243
Tradidiss.GC.even	0.974	0.908	0.816	0.829	0.626	0.245
Tradidiss.MS.even	0.974	0.908	0.814	0.828	0.623	0.243
Tradidiss.PE.even	0.974	0.907	0.828	0.831	0.639	0.25
Tradidiss.GC.uneven	0.967	0.901	0.827	0.815	0.622	0.244
Tradidiss.MS.uneven	0.966	0.899	0.823	0.813	0.618	0.243
Tradidiss.PE.uneven	0.969	0.905	0.837	0.821	0.637	0.249
$\beta_{turnover}$	0.974	0.911	0.837	0.829	0.641	0.251
$\beta_{heterogeneity}$	0.974	0.911	0.837	0.829	0.641	0.251
$\beta_{segregation}$	0.974	0.911	0.837	0.829	0.641	0.251

D_{IP}	0.923	0.778	0.68	0.565	0.338	0.034
D_{CW}	0.923	0.778	0.68	0.565	0.338	0.034
Bray-Curtis (species-based)	0.711	0.832	0.778	0.678	0.455	0.086

982   Table 4. Kendall tau correlations between environmental distance and the functional

983   dissimilarity measures at different values of sigma and with presence/absence data type

	Sigma=0.01	Sigma=0.1	Sigma=0.25	Sigma=0.5	Sigma=1	Sigma=5
CWMdis	0.944	0.818	0.691	0.568	0.353	0.014
CDFdis	0.943	0.820	0.700	0.594	0.306	-0.001
D(Q)	0.941	0.818	0.701	0.59	0.313	-0.003
dsimcom.SS	0.944	0.821	0.705	0.593	0.316	-0.003
dsimcom.Jac	0.944	0.821	0.705	0.593	0.316	-0.003
dsimcom.Sor	0.944	0.821	0.705	0.593	0.316	-0.003
dsimcom.Och	0.944	0.822	0.707	0.592	0.323	0.000
dsimcom.Beta	0.944	0.821	0.705	0.593	0.316	-0.003
dissABC.Jac	0.946	0.819	0.704	0.592	0.292	-0.006
dissABC.Sor	0.946	0.819	0.704	0.592	0.292	-0.006
dissABC.SS	0.946	0.819	0.704	0.592	0.292	-0.006
dissABC.Och	0.946	0.819	0.704	0.591	0.293	-0.006
dissABC.Kul	0.946	0.819	0.704	0.591	0.294	-0.006
dissABC.Si	0.939	0.835	0.701	0.556	0.324	0.019
Tradidiss.GC.even	0.945	0.820	0.707	0.593	0.305	-0.005
Tradidiss.MS.even	0.945	0.819	0.707	0.592	0.304	-0.005
Tradidiss.PE.even	0.947	0.821	0.704	0.595	0.323	-0.002
Tradidiss.GC.uneven	0.946	0.819	0.702	0.592	0.308	-0.006
Tradidiss.MS.uneven	0.946	0.819	0.703	0.592	0.307	-0.006
Tradidiss.PE.uneven	0.947	0.820	0.698	0.593	0.326	-0.003
$\beta_{turnover}$	0.943	0.817	0.699	0.585	0.331	-0.003
$\beta_{heterogeneity}$	0.943	0.817	0.699	0.585	0.331	-0.003
$\beta_{segregation}$	0.943	0.817	0.699	0.585	0.331	-0.003

D_{IP}	0.905	0.696	0.597	0.435	0.158	-0.017
D_{CW}	0.904	0.694	0.593	0.431	0.160	-0.017
PADDis.Jac	0.904	0.679	0.575	0.418	0.154	-0.025
PADDis.Sor	0.905	0.696	0.597	0.435	0.158	-0.017
PADDis.SS	0.902	0.662	0.546	0.396	0.144	-0.034
PADDis.Och	0.904	0.697	0.596	0.436	0.159	-0.017
PADDis.Simp	0.881	0.620	0.474	0.343	0.158	0.030
PADDis.Kul	0.904	0.694	0.593	0.431	0.160	-0.017
Sørensen (species-based)	0.698	0.724	0.606	0.415	0.127	0.048

Figure 1. Heat maps showing the interactive effects of niche width (sigma), transformation of

between-species dissimilarities (lin = linear, exp = exponential), data type (ABUND =

abundance, P/A = presence/absence), and dissimilarity index (1 – *CWMdis*, 2 – *CDFdis*, 3 –

D_Q, 4 – *dsimcom*/Sørensen, 5 – *dissABC*/Sørensen, 6 – *generalized_Tradidiss*/generalized

Canberra, uneven weighting, 7 – $\beta_{turnover}$, 8 – D_{CW}) on the correlation with environmental

distance