Estimating the rate of plasmid transfer with an adapted Luria–Delbrück fluctuation analysis and a case study on the evolution of plasmid transfer rate

Olivia Kosterlitz[1,2*], Clint Elg[2,3], Adamaris Muñiz Tirado[1], Claire Wate[1], Ivana Bozic[4], Eva M. Top[2,3], Benjamin Kerr[1,2*]

[1] Biology Department, University of Washington, Seattle, WA, USA.

[2] BEACON Center for the Study of Evolution in Action, East Lansing, MI, USA.

[3] Department of Biological Sciences and Institute for Bioinformatics and Evolutionary Studies, University of Idaho, Moscow, ID, USA.

[4] Department of Applied Mathematics, University of Washington, Seattle, WA, USA.

* e-mail: livkost@uw.edu; kerrb@uw.edu

Author contributions

O.K. and B.K. conceived of the presented ideas. B.K. and I.B. developed the theory. O.K. developed and performed the simulations. C.E. facilitated part of the simulations. O.K. developed the experimental protocol. O.K., A.M.T., and C.W. performed the experiments. O.K. and B.K. wrote the manuscript. E.M.T. and B.K. supervised the project. All authors discussed the results and contributed to the final manuscript.

Abstract

To increase our basic understanding of the ecology and evolution of conjugative plasmids, we need a reliable estimate of their rate of transfer between bacterial cells. Accurate estimates of plasmid transfer have remained elusive given biological and experimental complexity. Current methods to measure transfer rate can be confounded by many factors, such as differences in growth rates between plasmid-containing and plasmid-free cells. However, one of the most problematic factors involves situations where the transfer occurs between different strains or species and the rate that one type of cell donates the plasmid is not equal to the rate at which the other cell type donates. Asymmetry in these rates has the potential to bias transfer estimates, thereby limiting our capabilities for measuring transfer within diverse microbial communities. We develop a novel low-density method ("LDM") for measuring transfer rate, inspired by the classic fluctuation analysis of Luria and Delbrück. Our new approach embraces the stochasticity of conjugation, which departs in important ways from the current deterministic population dynamic methods. In addition, the LDM overcomes obstacles of traditional methods by allowing different growth and transfer rates for each population within the assay. Using stochastic simulations, we show that the LDM has high accuracy and precision for estimation of transfer rates compared to other commonly used methods. Lastly, we implement the LDM to estimate transfer on an ancestral and evolved plasmid-host pair, in which plasmid-host co-evolution increased the persistence of an IncP-1β conjugative plasmid in its *Escherichia coli* host. Our method revealed the increased persistence can be at least partially explained by an increase in transfer rate after plasmid-host co-evolution.

Introduction

A fundamental rule of life and the basis of heredity involves the passage of genes from parents to their offspring. Plasmids in bacteria can violate this rule of strict vertical inheritance by shuttling DNA among neighboring, non-related bacteria through horizontal gene transfer (HGT). Plasmids are independently replicating extrachromosomal DNA molecules, and some encode for their own horizontal transfer, a process termed conjugation. Bacteria containing such conjugative plasmids are governed by two modes of inheritance: vertical and horizontal. In an abstract sense, conjugation in prokaryotes is

similar to the genetic recombination that occurs in sexually reproducing eukaryotes; specifically, conjugation is a fundamental mechanism for producing new combinations of genetic elements. Since conjugation facilitates gene flow across vast phylogenetic distances, the expansive gene repertoire in the "accessory" genome is shared among many microbial species[1]. While this complicates phylogenetic inference, it reinforces the notion that conjugation has a central role in shaping the ecology and evolution of microbial communities[2,3]. Notably, conjugative plasmids are commonly the vehicle for the spread of antimicrobial resistance (AMR) genes and the emergence of multi-drug resistance (MDR) in clinical pathogens[4]. While such plasmids may impose growth costs on their bacterial hosts in the absence of antibiotics[5], theoretical and experimental studies have demonstrated how conjugation itself may be a critical process for the invasion and maintenance of costly plasmids[6,7,8]. To model the spread of antibiotic resistance, improve phylogenetic inference, and increase our understanding of the fundamental process of conjugation and its impacts on the basic biology of bacteria, accurate measures of conjugation are of the utmost importance.

The basic approach to measure conjugation involves mixing plasmid-containing bacteria, called "donors", with plasmid-free bacteria, called "recipients". After donors and recipients are incubated together for some period of time, which is hereafter denoted as $\tilde{t}$, the population is assessed for recipients that have received the plasmid from the donor; these transformed recipients are called "transconjugants". To describe conjugation the earliest and some of the most widely used approaches measure the densities of donors, recipients, and transconjugants after the incubation time $\tilde{t}$ ($D_{\tilde{t}}$, $R_{\tilde{t}}$, and $T_{\tilde{t}}$, respectively) to calculate various ratios (i.e., conjugation frequency or efficiency) including $T_{\tilde{t}}/R_{\tilde{t}}$, $T_{\tilde{t}}/D_{\tilde{t}}$, $T_{\tilde{t}}/(D_{\tilde{t}} + R_{\tilde{t}} + T_{\tilde{t}})$[9], see Zhong *et. al.*[10] and Huisman *et. al.*[11] for a comprehensive overview. These conjugation ratios are descriptive statistics and have been shown to vary with initial densities (D_0 and R_0) and incubation time ($\tilde{t}$)[10,11]. While these statistics may be useful for within study comparisons of the combined processes of conjugation and transconjugant growth, they are not estimating the rate of conjugation events. To see this, consider two plasmids, A and B, and assume both have the same conjugation rate, but plasmid A results in a higher transconjugant growth rate. Assuming all other parameters (e.g., donor growth, recipient growth, etc.) are the same and growth occurs during the assay, all of the above ratios would be higher for A, even though its conjugation rate

doesn't differ from plasmid B. Clearly, an estimate of the rate of conjugation itself is
desirable.

The conjugation rate (i.e., plasmid transfer rate) is the "gold-standard"
measurement to describe conjugation akin to measuring other fundamental rate
parameters, such as growth rate or mutation rate. Such metrics are useful for cross-
literature comparison to understand basic microbial ecology and evolution[12]. There are
only a few widely used methods to measure conjugation rate, which all are derived from
the Levin *et. al.*[13] mathematical model describing the change in density of donors,
recipients and transconjugants over time (given by dynamic variables D_t, R_t, and T_t,
respectively). In this model, each population type grows exponentially at the same growth
rate ψ. In addition, the transconjugant density increases as a result of conjugation events
both from donors to recipients and from existing transconjugants to recipients at the same
conjugation rate γ. The recipient density decreases due to these conjugation events. The
densities of these dynamic populations are described by the following differential
equations (where the t subscript is dropped from the dynamic variables for notational
convenience):

$$\frac{dD}{dt} = \psi D, \qquad [1]$$

$$\frac{dR}{dt} = \psi R - \gamma R(D + T), \qquad [2]$$

$$\frac{dT}{dt} = \psi T + \gamma R(D + T). \qquad [3]$$

Each current rate estimation method solved the set of ordinary differential
equations from the Levin *et. al.* model (or a slight variation) to find an estimate for the
conjugation rate γ. These estimates differ by the assumptions used to find the analytical
solution. Two estimates for conjugation rate standout as widely used in the literature[7,14,15].
Levin *et. al.*[13] derived the first of these estimates for conjugation rate γ in terms of the
density of donors, recipients, and transconjugants ($D_{\tilde{t}}$, $R_{\tilde{t}}$, and $T_{\tilde{t}}$, respectively) after the
incubation time $\tilde{t}$:

$$\gamma = \frac{T_{\tilde{t}}}{D_{\tilde{t}} R_{\tilde{t}} \tilde{t}}. \qquad [4]$$

88 We label the expression in equation [4] as the "TDR" estimate for the conjugation rate,
89 where TDR stands for the dynamic variables used in the estimate. Besides the model's
90 homogeneous growth rates and conjugation rates, the most notable assumption used in

the TDR derivation is that there is little to no change in the population densities due to growth. Thus, laboratory implementation that respects this assumption can be difficult; albeit chemostats and other implementation methods have been developed to circumvent this constraint[7,13].

Simonsen *et. al.*[16] derived the second of these widely used estimates for conjugation rate γ, which importantly expands application beyond the TDR method by allowing for population growth. In addition to $D_{\tilde{t}}$, $R_{\tilde{t}}$, and $T_{\tilde{t}}$, this estimate depends on the initial and final density of all bacteria (N_0 and $N_{\tilde{t}}$, respectively) as well as the population growth rate (ψ).

$$\gamma = \psi \ln\left(1 + \frac{T_{\tilde{t}}}{R_{\tilde{t}}}\frac{N_{\tilde{t}}}{D_{\tilde{t}}}\right)\frac{1}{(N_{\tilde{t}} - N_0)} \qquad [5]$$

We term equation [5] as the "SIM" estimate for the conjugation rate, where SIM stands for "Simonsen *et. al.* Identicality Method" since the underlying model still assumes all strains are *identical* with regards to growth rates and conjugation rates. The laboratory implementation for the SIM is straightforward, requiring measurements of cell densities over some period of time. Because several of these densities are acquired at the end of an assay, the SIM is often referred to as "the end-point method". However, we forego this term since other methods we discuss are also end-point methods.

Huisman *et. al.*[11] recently updated the SIM estimate, further extending its application by relaxing the assumption of identical growth and transfer rates for all strains. Specifically, their model extension allowed each strain to have its own growth rate (ψ_D, ψ_R, and ψ_T denote the growth rates for donors, recipients, and transconjugants, respectively) and each plasmid-bearing strain to have a distinct transfer rate (γ_D and γ_T denote the rate of conjugation to recipients from donors and transconjugants, respectively). They derived an estimate of γ_D by assuming that the rate of production of transconjugants through cell division and conjugation from donors to recipients was much greater than conjugation from transconjugants to recipients (e.g., $\gamma_T \leq \gamma_D$ is sufficient to ensure this assumption under many standard assay conditions):

$$\gamma_D = (\psi_D + \psi_R - \psi_T)\frac{T_{\tilde{t}}}{\left(D_{\tilde{t}}R_{\tilde{t}} - D_0 R_0 e^{\psi_T \tilde{t}}\right)}. \qquad [6]$$

We term equation [6] as the ASM estimate for donor-recipient conjugation rate, where ASM stands for "Approximate Simonsen *et. al.* Method". The descriptions provided for the aforementioned conjugation rate methods are cursory but see the Supplemental

Information (SI section 1-6) for a more detailed historical overview, mathematical derivations, laboratory implementations, and a fuller discussion of how methods compare.

Deterministic numerical simulations have been used to test the accuracy of the aforementioned methods (TDR, SIM, and ASM), revealing important limitations of each approach[11,16,17]. For instance, simulations have demonstrated that all the estimates can become inaccurate when the donor-recipient conjugation rate (γ_D, hereafter "donor conjugation rate" for simplicity) differs from the transconjugant-recipient conjugation rate (γ_T, hereafter "transconjugant conjugation rate" for simplicity). Such a situation is likely when the donor and recipient populations are different strains or species. While deterministic simulations have revealed estimate limitations, the deterministic framework itself encompasses an additional limitation. Conjugation is an inherently stochastic process (i.e., random process), and the effects of such stochasticity on the accuracy and precision of rate estimates are currently unknown.

Here we derive a novel estimate for conjugation rate, inspired by the Luria–Delbrück fluctuation experiment, by explicitly tracking transconjugant dynamics as a stochastic process (i.e., a continuous time branching process). In addition, our method allows for unrestricted heterogeneity in growth rates and conjugation rates. Thus, our method fills a gap in the methodological toolkit by allowing accurate estimation of conjugation rates in a wide variety of strains and species. For example, over the past few years several studies have called for and recognized the importance of evaluating the biology of plasmids in microbial communities[4,18,19]. Our method is available to support this direction of research by clearly disentangling cross-species and within-species conjugation rates. Thus, our approach can provide accurate conjugation rate parameters for mechanistic models describing plasmid dynamics in microbial communities, where key assumptions of previous estimates can be violated. We used stochastic simulations to validate our estimate and compare its accuracy and precision to the aforementioned estimates. We also developed a high throughput protocol for the laboratory by using microtiter plates to rapidly screen many mating cultures (i.e., donor-recipient co-cultures) for the existence of transconjugants. Our protocol is unique compared to the other estimates and circumvents problems involving spurious laboratory results that can bias other approaches. Finally, we implement our approach to investigate the evolution of conjugation rate in an *Escherichia coli* host containing an IncP-1β conjugative plasmid.

Theoretical Framework

Conjugation is a stochastic process that transforms the genetic state of a cell. This is analogous to mutation, which is also a process that transforms the genetic state of a cell. While mutation transforms a wild-type cell to a mutant, conjugation transforms a recipient cell to a transconjugant.

This analogy inspired us to revisit the way Luria and Delbrück handled the mutational process in their classic paper on the nature of bacterial mutation[20], outlined in Figure 1a-d. For this process, assume that the number of wild-type cells, N_t, is expanding exponentially. Let the rate of mutant formation be given by μ. In Figure 1a, we see that the number of mutants in a growing population increases due to mutation events (highlighted purple cells) and due to faithful reproduction by mutants (non-highlighted purple cells). The rate at which mutants are generated (highlighted purple cells) is μN_t, which grows as the number of wild-type cells increases (Figure 1b). However, the rate of transformation per wild-type cell is the mutation rate μ, which is constant (Figure 1c). Since mutations are random, parallel cultures will vary in the number of mutants depending if and when mutation events occur. As seen in Figure 1d, for sufficiently small wild-type populations growing over sufficiently small periods, some replicate populations will not contain any mutant cell (gray shading) while other populations exhibit mutants (yellow shading). Indeed, the cross-replicate fluctuation in the number of mutants was a critical component of the Luria-Delbrück experiment.

To apply this strategy to estimate the conjugation rate, we can similarly think about an exponentially growing population of recipients (Figure 1e). But now there is another important cell population present (the donors). The transformation of a recipient is simply the generation of a transconjugant (highlighted purple cells) via conjugation with a donor. If we ignore conjugation from transconjugants, the rate at which transconjugants are generated is $\gamma_D D_t R_t$ (Figure 1f). However, when we consider the rate of transformation per recipient cell, we do not have a constant rate. Rather, this transformation rate per recipient is $\gamma_D D_t$, which grows with the donor population (Figure 1g). It is as if we are tracking a mutation process where the mutation rate is exponentially increasing. Yet the rate of transformation per recipient *and* donor is constant, which is the donor conjugation rate γ_D. As with mutation, conjugation is random which results in a distribution in the

184 number of transconjugants among parallel cultures depending on the time points at which
185 transconjugants arise. As seen in Figure 1h, under certain conditions, some replicate
186 populations will not contain any transconjugant cell (gray shading) while other populations
187 will exhibit transconjugants (yellow shading).

Figure 1: Schematic comparing the process of mutation (a-d) to the process of conjugation (e-h). (a) In a growing population of wild-type cells, mutants arise (highlighted purple cells) and reproduce (non-highlighted purple cells). (b) The rate at which mutants are generated grows as the number of wild-type cells increases at rate μN_t. (c) The rate of transformation per wild-type cell is the mutation rate μ. (d) Wild-type cells growing in 9 separate populations where mutants arise in a proportion of the populations (yellow background) at different cell divisions. Here, for each tree, time

progresses from top to bottom (rather than left to right as in part a). (e) In a growing population of donors and recipients, transconjugants arise (highlighted purple cells) and reproduce (non-highlighted purple cells). (f) The rate at which transconjugants are generated grows as the numbers of donors and recipients increase at rate $\gamma_D D_t R_t$. (g) The rate of transformation per recipient cell grows as the number of donors increases at a rate $\gamma_D D_t$, where γ_D is the constant conjugation rate. (h) Donor and recipient cells growing in 9 separate populations where transconjugants arise in a proportion of the populations (yellow background) at different points in time (which again, here, progresses top to bottom for each tree).

Using this analogy, here we describe a new approach for estimating conjugation
rate which embraces conjugation as a stochastic process[21]. Let the density of donors,
recipients and transconjugants in a well-mixed culture at time t be given by the variables
D_t, R_t, and T_t. In all that follows, we will assume that the culture is inoculated with donors
and recipients, while transconjugants are initially absent (i.e., $D_0 > 0$, $R_0 > 0$, and $T_0 =$
0). The donor and recipient populations grow according to the following standard
exponential growth equations

$$D_t = D_0 e^{\psi_D t}, \tag{7}$$

$$R_t = R_0 e^{\psi_R t}, \tag{8}$$

where ψ_D and ψ_R are the growth rates for donor and recipient cells, respectively. With
equations [7] and [8], we are making a few assumptions, which also occur in some of the
previous methods (see SI Table 3 for a comparison). First, we assume that donors and
recipients grow exponentially. Given that we focus on low-density culture conditions
where nutrients are plentiful, this assumption will be reasonable. Second, we assume that
population growth for these two strains is deterministic (i.e., D_t and R_t are not random
variables). As long as the numbers of donors and recipients are not too small (i.e., $D_0 \gg$
0 and $R_0 \gg 0$), this assumption is acceptable. Lastly, we assume the loss of recipient cells
to transformation into transconjugants can be ignored. For what follows, the rate of
generation of transconjugants per recipient cell (as in Figure 1g) is very small relative to
the per capita recipient growth rate, making this last assumption justifiable.
The population growth of transconjugants is modeled using a continuous-time
stochastic process. The number of transconjugants, T_t, is a random variable taking on
non-negative integer values. In this section, we will assume the culture volume is 1 ml
and thus the number of transconjugants (i.e., cfu) is equivalent to the density of
transconjugants (i.e., cfu per ml). For a very small interval of time, Δt, the current number
of transconjugants will either increase by one or remain constant. The probabilities of
each possibility are given as follows:

$$\Pr\{T_{t+\Delta t} = T_t + 1\} = \gamma_D D_t R_t \Delta t + \gamma_T T_t R_t \Delta t + \psi_T T_t \Delta t, \quad\quad [9]$$

$$\Pr\{T_{t+\Delta t} = T_t\} = 1 - (\gamma_D D_t R_t + \gamma_T T_t R_t + \psi_T T_t)\Delta t. \quad\quad [10]$$

The three terms on the right-hand side of equation [9] illustrate the processes enabling
the transconjugant population to increase. The first term gives the probability that a donor
transforms a recipient into a transconjugant via conjugation (γ_D). The second term gives
the probability that a transconjugant transforms a recipient via conjugation (γ_T). The third
term measures the probability that a transconjugant cell divides (ψ_T is the transconjugant
growth rate). Equation [10] is simply the probability that none of these three processes
occur.

We focus on a situation where there are no transconjugants. Therefore, the only
process that can change the number of transconjugants is conjugation of the plasmid
from a donor to a recipient. Using equation [10] with $T_t = 0$, we have

$$\Pr\{T_{t+\Delta t} = 0 \mid T_t = 0\} = 1 - \gamma_D D_t R_t \Delta t. \quad\quad [11]$$

223   We let the probability that we have zero transconjugants at time t be denoted by $p_0(t)$
224   (i.e., $p_0(t) = \Pr\{T_t = 0\}$). In SI section 7, we derive the following expression for $p_0(t)$ at
225   time $t = \tilde{t}$:

$$p_0(\tilde{t}) = \exp\left\{\frac{-\gamma_D D_0 R_0}{\psi_D + \psi_R}\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right)\right\}. \quad\quad [12]$$

226   Solving equation [12] for γ_D yields a new measure for the donor conjugation rate:

$$\gamma_D = -\ln p_0(\tilde{t})\left(\frac{\psi_D + \psi_R}{D_0 R_0\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right)}\right). \quad\quad [13]$$

227   This expression is similar in form to the mutation rate derived by Luria and Delbrück in
228   their classic paper on the nature of bacterial mutation, which is not a coincidence.

229   In SI section 8, we rederive the Luria-Delbrück result, which can be expressed as

$$\mu = -\ln p_0(\tilde{t})\left(\frac{\psi_N}{N_0\left(e^{\psi_N \tilde{t}} - 1\right)}\right). \quad\quad [14]$$

In the mutational process modeled by Luria and Delbrück, N_0 is the initial wild-type
population size, which grows exponentially at rate ψ_N. For Luria and Delbrück, $p_0(\tilde{t})$ refers
to the probability of zero mutants at time $\tilde{t}$ (as in a gray-shaded tree in Figure 1d), whereas
$p_0(\tilde{t})$ in the conjugation estimate refers to the probability of zero transconjugants (as in a
gray-shaded tree in Figure 1h). Comparing equation [14] to equation [13], conjugation

can be thought of as a mutation process with initial wild-type population size $D_0 R_0$ that

grows at rate $\psi_D + \psi_R$. We explore further the connection between mutation and

conjugation in SI section 13.

We label the expression in equation [13] as the LDM estimate for donor

conjugation rate, where LDM stands for "Low Density Method" as the assumption of

exponential growth is well grounded when the cell populations are growing at low density.

Indeed, this is how we conduct the assay to compute this estimate. The acronym can

alternatively stand for "Luria-Delbrück Method" given the connection to their approach.

Simulation Methods

To explore the accuracy and precision of the LDM estimate (as well as other

estimates), we developed a stochastic simulation framework using the Gillespie

algorithm. We extended the base model (equations [1] - [3]) to include plasmid loss due

to segregation at rate τ. Specifically, transconjugants are transformed into plasmid-free

recipients due to plasmid loss via segregation at rate τ_T. Similarly, the donors become

plasmid-free cells through segregative loss of the plasmid at rate τ_D; therefore, the

extended model tracks the change in density of a new population type, plasmid-free

former donors (F). In total, the extended model describes the density changes in four

populations (D, R, T, and F) influenced by the values of various biological parameters:

growth rates (ψ_D, ψ_R, ψ_T, and ψ_F), conjugation rates (γ_D and γ_T) and segregation rates

(τ_D and τ_T). For each simulation, we specify starting cell densities ($D_0 > 0$ and $R_0 > 0$,

but always $T_0 = F_0 = 0$) and allow the four populations to grow and transform

stochastically for a specified incubation time $\tilde{t}$ in a 1 ml culture volume. The simulations

occur in sets, where each set involves a sweep across values of some key parameter.

Each sweep varies a biological process (i.e., growth, conjugation, or segregation) in

isolation to determine the effect on accuracy and precision of the donor conjugation rate

estimates. For instance, the transconjugant conjugation rate (γ_T) could be varied while all

other parameters are held constant. For each parameter setting within a sweep, ten

thousand parallel populations were simulated, and then conjugation rate was calculated

to yield replicate estimates for each method of interest (TDR, SIM, ASM, and LDM).

Notably, the LDM requires an estimate of the probability that a population has no

transconjugants at the end of the assay, which we denote $\hat{p}_0(\tilde{t})$. The maximum likelihood

estimate of this probability is the fraction of populations (i.e., parallel simulations) that

have no transconjugants at the specified incubation time $\tilde{t}$ (see SI section 9 for details).

Thus, for LDM, we calculated $\hat{p}_0(\tilde{t})$ at the specified incubation time $\tilde{t}$ by reserving multiple

parallel populations for each conjugation rate estimate. For more details on the

simulations such as the extended model equations, selection of the specified incubation

time $\tilde{t}$, and calculation of each estimate see SI section 11.

Simulation Results

Exploring accuracy and sensitivity of the LDM and other conjugation methods

The underlying assumptions used to derive the various estimates for conjugation

rate (i.e., TDR, SIM, ASM, and LDM) entail simplifications of biological systems. To

explore the effects of violating these assumptions on the LDM and other conjugation rate

estimates, we used a stochastic simulation framework to investigate their accuracy across

a range of theoretical conditions. All of our simulated populations *increase in size* over

the incubation time, so a fundamental assumption of the TDR approach (i.e., no change

in the density due to growth) is broken for all of the runs. Here we divide our simulations

into three scenarios that break additional assumptions of at least one conjugation model

considered to this point: unequal growth rates, unequal conjugation rates, and a non-zero

segregation rate. When growth rates are different between populations, a modeling

assumption is violated for the TDR and SIM estimates. When donor and transconjugant

conjugation rates are unequal, a modeling assumption can be violated for the TDR, SIM,

and ASM estimates. When the segregation rate is non-zero, a modeling assumption is

violated for all of the estimates considered. We note that some of our parameter values

fall outside of reported values; these choices were made either to improve computational

speed or to better illustrate the effects of differences (e.g., in growth or transfer rates).

However, many of the parameter values used to explore the effects of incubation time

were set to reported parameter values.

The effect of unequal growth rates

Plasmid-containing hosts often incur a cost or gain a benefit in terms of growth rate

for plasmid carriage relative to plasmid-free hosts. We simulated a range of growth-rate

effects on plasmid-containing hosts from large plasmid costs ($\psi_D = \psi_T \ll \psi_R$) to large plasmid benefits ($\psi_D = \psi_T \gg \psi_R$). The LDM estimate had high accuracy and precision across all parameter settings (Figure 2a). The ASM estimate performed similarly to the LDM estimate with regards to the mean value for each parameter setting (Figure 2a, black horizontal lines). Given that the ASM was developed to allow for unequal growth rates, its mean accuracy conforms to expectations. However, it is also worth noting that the distribution of the ASM estimate was skewed for some of the parameter settings. In other words, the mean (Figure 2a, black horizontal lines) is separated from the median (Figure 2a, green horizontal line); therefore, the average of a small number of replicates could have led to a deviant mean estimate. The TDR and SIM estimates were inaccurate for some to all of the parameter settings and also had skewed distributions for some parameter settings. With regards to precision, the LDM estimate outperformed the other estimates due to a smaller variance at each parameter setting.

Other than unequal growth rates due to plasmid carriage, there are other biological scenarios where growth rates can differ between donors, recipients and transconjugants. In fact, often it is the case that donors and recipients are different strains or species. We simulated donors growing faster ($\psi_D > \psi_T = \psi_R$) or slower ($\psi_D < \psi_T = \psi_R$) than recipients and transconjugants. Similar to the plasmid carriage scenario, the LDM exhibited high accuracy and precision relative to other metrics (SI Figure 1).

The effect of unequal conjugation rates

The conjugation rate can vary between different bacterial strains and species. We explored different parameter settings again, ranging from a relatively low transconjugant conjugation rate ($\gamma_T \ll \gamma_D$) to a relatively high transconjugant conjugation rate ($\gamma_T \gg \gamma_D$) compared to the donor conjugation rate. The LDM estimate had high accuracy and precision across all parameter settings and outperformed the other estimates, especially when transconjugant conjugation rate greatly exceeded donor conjugation rate (Figure 2b). This result aligns with our expectation given that a relatively high transfer rate from transconjugants to recipients can violate a modeling assumption for all the other estimates.

Figure 2 : The effect of violating various modeling assumptions on estimating conjugation rate. The Gillespie algorithm was used to simulate population dynamics. The parameter values were $\psi_D = \psi_R = \psi_T = 1$, $\gamma_D = \gamma_T = 1 \times 10^{-6}$, and $\tau_D = \tau_T = 0$ unless otherwise indicated. The dynamic variables were initialized with $D_0 = R_0 = 1 \times 10^2$ and $T_0 = F_0 = 0$. All incubation times are short but are specific to each parameter setting (see SI section 11 for details). Conjugation rate was estimated using 10,000 populations and summarized using boxplots. The gray dashed line indicates the "true" value for the donor conjugation rate. The box contains the 25th to 75th percentile. The vertical lines connected to the box contain 1.5 times the interquartile range above the 75th percentile and below the 25th percentile with the caveat that the whiskers were always constrained to the range of the data. The colored line in the box indicates the median. The solid black line indicates the mean. The boxplots in gray indicate the baseline parameter setting, and all colored plots represent deviation of one or two parameters from baseline. (a) Unequal growth rates were explored over a range of growth rates for the plasmid-bearing strains, namely $\psi_D = \psi_T \in \{0.0625, 0.125, 0.25, 0.5, 1, 2, 4, 8\}$. (b) Unequal conjugation rates were probed over a range of transconjugant conjugation rates, namely $\gamma_T \in \{10^{-9}, 10^{-8}, 10^{-7}, 10^{-6}, 10^{-5}, 10^{-4}, 10^{-3}, 10^{-2}\}$. (c) We also explored the process of segregation by considering a range of segregative loss rates $\tau_D = \tau_T \in \{0, 0.0001, 0.001, 0.001, 0.01\}$.

The effect of a non-zero segregation rate

Plasmid loss due to segregation is a common occurrence in plasmid populations and violates a model assumption underlying all the conjugation rate estimates. We simulated a range of segregative loss rates, ranging from low ($\tau_D = \tau_T = 0.0001$) to high ($\tau_D = \tau_T = 0.1$). The LDM had high accuracy and precision across all parameter settings (Figure 2c). The effect of segregation was undetectable even for an extremely high segregation rate ($\tau_D = \tau_T = 0.1$). Similarly, the effect of segregative loss was undetectable on the other conjugation estimates compared to their performance without any segregative loss (Figure 2c, colored vs. grey box plots). Thus, we find that all estimates appear robust with regards to an introduction of the process of segregation.

Case studies to explore the effects of incubation time

The incubation time $\tilde{t}$ is an important consideration for executing all conjugation rate assays in the laboratory. Often the incubation time is variable for estimates of conjugation rate reported in the literature. We explored the effects of altering incubation time by estimating conjugation rate at 30-minute intervals. Since these estimates rely on measuring different quantities, the range for the 30-minute sampling intervals differ between estimates. For the TDR, SIM, and ASM estimates, the range for these intervals was determined by the condition that at least 90 percent of the simulated populations contained transconjugants at the incubation time $\tilde{t}$. Specifically, we note that equations [4], [5], and [6] all require $T_{\tilde{t}} > 0$ for a non-zero conjugation rate estimate. This contrasts with the LDM since this estimate needs some parallel populations to be absent of transconjugants. Specifically, equation [13] requires $0 < p_0(\tilde{t}) < 1$ for a non-zero finite conjugation rate estimate. Thus, for the LDM estimate, the range for the sampling intervals was determined by the condition that at least one parallel population had zero transconjugants. Therefore, estimates for the LDM are shown at earlier incubation times than those for the other methods (Figure 3).

Figure 3 : The effect of incubation time ($\tilde{t}$) on estimating conjugation rate. The Gillespie algorithm was used to simulate population dynamics with $\psi_D = \psi_R = \psi_T = 1$, $\gamma_D = \gamma_T = 1 \times 10^{-14}$, and $\tau_D = \tau_T = 0$ unless otherwise indicated. The dynamic variables were initialized with $D_0 = R_0 = 1 \times 10^5$ and $T_0 = F_0 = 0$. Conjugation rate was estimated for 100 populations and summarized using boxplots. The gray dashed line indicates the "true" value for the donor conjugation rate. Each box contains the 25th to 75th percentile. The vertical lines connected to the box contain 1.5 times the interquartile range above the 75th percentile and below the 25th percentile with the caveat that the whiskers were always constrained to the range of the data. The colored line in the box indicates the median. The solid black line indicates the mean. (a) An unequal growth rate was simulated with $\psi_D = \psi_T = 0.07$. (b) An unequal conjugation rate was simulated with $\gamma_T = 1 \times 10^{-9}$. (c) We simulated a positive segregation rate with $\tau_D = \tau_T = 0.01$.

We ran simulations with parameter values for growth and conjugation rates which are reported in the literature ($\psi_D = \psi_R = \psi_T = \psi_F = 1$, $\gamma_D = \gamma_T = 1 \times 10^{-14}$) and reasonable initial densities ($D_0 = R_0 = 1 \times 10^5$). As before, each biological scenario examined a perturbation to a single biological parameter (i.e., growth, conjugation, or segregation) to determine the effect on accuracy and precision. For the scenario exploring growth differences, we added a plasmid cost (Figure 3a, $\psi_D = \psi_T = 0.7$). For the scenario highlighting unequal conjugation rates, we increased the conjugation rate of transconjugants (Figure 3b, $\gamma_T = 1 \times 10^{-9}$). For the scenario incorporating segregation, we simply chose a rather high segregation rate (Figure 3c, $\tau_D = \tau_T = 0.01$).

The LDM estimate had high accuracy over all time points for all scenarios with precision increasing through time. The other estimates also became more precise over time. However, their greater precision over time was sometimes accompanied by increasing inaccuracy (Figure 3b). Again, the LDM estimate performed as well or better than other estimates across incubation times.

The LDM Implementation

In this section we describe the general experimental procedure for estimating donor conjugation rate (γ_D) using the LDM estimate in the laboratory. The assay can accommodate a wide variety of microbial species and conjugative plasmids since our estimate allows for distinct growth and conjugation rates. In SI section 6, we rearrange equation [13] to this alternative form for implementing the LDM in the laboratory:

$$\gamma_D = f\left\{\frac{1}{\tilde{t}}[-\ln\hat{p}_0(\tilde{t})]\frac{\ln D_{\tilde{t}}R_{\tilde{t}} - \ln D_0 R_0}{D_{\tilde{t}}R_{\tilde{t}} - D_0 R_0}\right\}. \qquad [15]$$

Similar to previous conjugation estimates, the LDM requires measurement of initial and final densities of donors and recipients ($D_0, R_0, D_{\tilde{t}}$, and $R_{\tilde{t}}$). In addition, the LDM requires an estimate of the probability that a population has no transconjugants at the end of the assay, which we denote $\hat{p}_0(\tilde{t})$. The maximum likelihood estimate of this probability is the fraction of populations (i.e., parallel mating cultures) that have no transconjugants at the specified incubation time $\tilde{t}$ (see SI section 9 for details). Cell densities are generally measured in $\frac{\text{cfu}}{\text{ml}}$ units and the conjugation rate in $\frac{\text{ml}}{\text{h} \cdot \text{cfu}}$ units, thus, the standard unit of volume in these assays is a milliliter. If exactly 1 ml is used as the volume for each mating culture (as assumed in previous sections), then the quantity in braces in equation [15] is

the estimate for the conjugation rate. However, smaller mating volumes can be advantageous (e.g., 100 μl in a well inside a microtiter plate). When the mating volume deviates from 1 ml, we must add a correction factor f which is the number of experimental volume units (evu) per ml (e.g., for a mating volume of 100 μl, we would have $f = 10\frac{\text{evu}}{\text{ml}}$, see SI section 10 for details).

For a given incubation time ($\tilde{t}$) and initial densities of donors (D_0) and recipients (R_0), there will be some probability that transconjugants form $(1 - p_0(\tilde{t}))$. The combinations of times and densities where this probability is not close to zero or one is hereafter referred to as "the conjugation window". That is, for these time-density combinations, there is a good chance that some mating cultures will result in transconjugants, while others do not. For fixed initial donor and recipient densities, the conjugation window refers to the time period where transconjugants are first expected to form. Determining the conjugation window provides the researcher with *target* initial densities of donors (D_0') and recipients (R_0') and *target* incubation times ($\tilde{t}'$). Note we add primes to the *target* time-density combination to set them apart from D_0, R_0, and $\tilde{t}$ in equation [15] which will be gathered in the conjugation assay itself.

To determine the conjugation window and find a time-density combination, we mix exponentially growing populations of donors and recipients in a large array of parallel mating cultures for a full factorial treatment of initial densities and incubation times (Figure 4a and 4b). For ease of presentation, we will assume that donors and recipients start at equal proportion and we will refer to the "initial density" as the total cell density of these two strains. In our schematic, columns of a 96 deep well microtiter plate are grouped by the value of the initial density (3 columns per value; Figure 4a) and rows are grouped by the value of the incubation time (2 rows per value; Figure 4b). At each time in the set of incubation times being explored, growth medium selecting for transconjugants (see SI section 12 for explanation) is added. This dilutes the mating cultures by ten-fold, which hinders further conjugation and simultaneously permits the growth of any transconjugant cells that previously formed (Figure 4b). After a longer incubation, we assess the mating cultures (3 columns x 2 rows = 6 wells; Figure 4c) within each time-density treatment for presence or absence of transconjugants (i.e., turbid culture or non-turbid culture). There are three possible outcomes for each treatment: all matings have transconjugants, none of the matings have transconjugants, or there is both transconjugant-containing and

transconjugant-free matings. We are interested in the last outcome (Figure 4c, gray dots). These treatments meet the $\hat{p}_0(\tilde{t})$ condition (i.e., $0 < \hat{p}_0(\tilde{t}) < 1$) thus identifying a valid time-density combination for the LDM conjugation assay. As a general expectation, mating cultures with high donor conjugation rates (γ_D) will require shorter incubation times than mating cultures with low donor conjugation rates for a given initial density. We note that there will generally be multiple time-density treatments in the conjugation window and the final treatment choice may be determined by experimental tractability. Thus, this choice is not constrained and can be made based on an incubation time that is practically feasible. In addition, the LDM does not require the initial density of donors and recipients to be identical (as was assumed for this presentation). Indeed, in certain circumstances it may be easier to have donor and recipient initial densities be unequal.

The chosen treatment yields the target time-density combination (D_0', R_0', $\tilde{t}'$), and we proceed with executing the LDM conjugation assay to gather all components to estimate the conjugation rate. We mix exponentially growing populations of donors and recipients, inoculate many mating cultures at the target initial density ($D_0'+R_0'$) in a 96 deep well plate, and incubate in non-selective growth medium for the target incubation time ($\tilde{t}'$) (Figure 4d). We note that the incubation time $\tilde{t}$ may deviate slightly from the target incubation time $\tilde{t}'$ due to timing constraints in the laboratory, but small deviations are generally permissible. Thus, the executed incubation time $\tilde{t}$ should be used to calculate the conjugation rate. To estimate the initial densities (D_0 and R_0), three mating cultures at the start of the assay are diluted and plated on donor-selecting and recipient-selecting agar plates. We note that the initial densities (D_0 and R_0) may deviate from the target initial densities (D_0' and R_0'), but small deviations are permissible. After the incubation time $\tilde{t}$, final densities ($D_{\tilde{t}}$ and $R_{\tilde{t}}$) are obtained by dilution-plating from the same mating cultures. Growth medium selecting for transconjugants is subsequently added to the remaining mating cultures. To obtain $\hat{p}_0(\tilde{t})$, we assess the proportion of mating cultures that are non-turbid after incubation in transconjugant-selecting medium, as these cultures have no transconjugants (Figure 4d). This differs from the traditional Luria–Delbrück method since no plating is required to obtain $\hat{p}_0(\tilde{t})$. The binary output (turbidity vs. no turbidity) is assessed in a high throughput manner using a standard microtiter spectrophotometer. Using the obtained densities (D_0, R_0, $D_{\tilde{t}}$, and $R_{\tilde{t}}$), the incubation time ($\tilde{t}$), the proportion of transconjugant-free populations ($\hat{p}_0(\tilde{t})$), and correcting for the

450 experimental culture volume (f), the LDM estimate for donor conjugation rate (γ_D) can be

451 calculated via equation [15].

Finding the conjugation window

Executing the LDM conjugation assay

Figure 4 : Overview for finding the conjugation window (a-c) and executing the LDM conjugation assay (d). (a) Microtiter plate map designating placement of donors (red), recipients (blue), transconjugants (purple), and mating cultures over 10-fold dilutions (different shades of gray). For simplicity, donors and recipients are at the same proportion in each mating culture. The control wells are the last three columns. (b) Using the microtiter plate from part a, transconjugant-selecting medium (green well fill) is added at each time period designated by two rows in the microtiter plate. Two example wells from different time-density treatments are highlighted on the left where the gray background shading refers to the initial density treatment labeled in part a. In the top example well, transconjugant-selecting medium is added immediately, inhibiting growth of donor and recipient cells (grey dashed cells), and resulting in a non-turbid well as no transconjugants formed. In the bottom example well, the donor and recipient populations in the mating culture grow until transconjugant-selecting medium is added at 9-hours, inhibiting growth of donors and recipients, and permitting growth of the formed transconjugants. (c) After a lengthy incubation of the microtiter plate from part b, there are two well-types in the microtiter plate (bottom-left): transconjugant-containing (green well with purple cross) and

transconjugant-free (green well). For each treatment (time-density pair), the 6 mating wells are considered as a group, resulting in one of three outcomes (top): all transconjugant-free wells (white dot), all transconjugant-containing wells (black dot), a proportion of both well-types (gray dot). Any treatment with a gray dot represents a viable combination of initial densities ($D_0' + R_0'$) and incubation time ($\tilde{t}'$). (d) The microtiter plate is set up with many matings (gray) at the chosen target density ($D_0' + R_0'$) in addition to control wells. Three matings are sampled to determine the actual initial densities (D_0 and R_0) and final densities ($D_{\tilde{t}}$ and $R_{\tilde{t}}$). After the incubation time ($\tilde{t}$), which is as close to the target incubation time ($\tilde{t}'$) as possible, transconjugant-selecting medium is added to the microtiter plate (excluding the matings used for density-plating). After a lengthy incubation, the proportion of transconjugant-free (i.e., non-turbid) wells is calculated, yielding $\hat{p}_0(\tilde{t})$. Using the actual incubation time ($\tilde{t}$), initial densities (D_0, R_0), final densities ($D_{\tilde{t}}$, $R_{\tilde{t}}$), and correcting for the experimental culture volume (f), the LDM estimate is used to calculate donor conjugation rate (γ_D). Part d can be repeated to obtain replicate LDM estimates.

Application of the LDM

To explore the logistics of implementing the LDM, we measured the conjugation rate of a conjugative, IncP-1β plasmid (pALTS29) encoding chloramphenicol resistance in an *Escherichia coli* host[4]. Jordt *et. al.*[4] used this plasmid-host pair as an ancestral strain which was propagated in chloramphenicol antibiotic for 400 generations to explore mechanisms of plasmid-host co-evolution. The evolved lineage acquired genetic modifications in the host chromosome and the plasmid. The ancestral plasmid was gradually lost from the ancestral host population when these plasmid-bearing cells were propagated under conditions that did not select for the plasmid (Figure 5a, dashed line). In contrast, the evolved plasmid was considerably more persistent in the absence of selection in its co-evolved host (Figure 5a, solid line).

This pattern of increased plasmid persistence after plasmid-host co-evolution is consistent with previous studies[22,23]. In most cases, the cost of plasmid carriage is assumed to decrease with plasmid-host co-evolution; however, changes in other factors, such as an increased rate of conjugation or a decreased rate of segregational loss, may also result in increased plasmid persistence. Using the LDM method, we showed that the observed increase in plasmid persistence in the Jordt. *et. al.*[4] study is at least in part due to a significant increase in the conjugation rate in the evolved lineage (Figure 5b, Mann-Whitney U test, p-value = 0.029).

Figure 5 : LDM conjugation rate estimates of an ancestral and evolved lineage of *E. coli* containing a conjugative IncP-1β plasmid. (a) Plasmid persistence in the absence of chloramphenicol antibiotic was higher in the evolved plasmid-host pair (solid line) than the ancestral plasmid-host pair (dashed line). Data and figure adapted from Jordt *et. al.*[4] (b) Conjugation rate significantly increased in the evolved (Evo) plasmid-host pair compared to the ancestral (Anc) plasmid-host pair. The points represent a conjugation rate estimate from an LDM conjugation assay. The black line is the mean of four replicates. The colored line is the median of the four replicates. The asterisk indicates a significant difference from a Mann-Whitney U test.

Discussion

Here we have presented a new method for estimating the rate of plasmid conjugative transfer from a donor cell to a recipient cell. We derived our LDM estimate using a mathematical approach that captures the stochastic process of conjugation, which was inspired by the method Luria and Delbrück applied to the process of mutation. This departs from the mathematical approach for other conjugation rate estimates which assume underlying deterministic frameworks guiding the dynamics of transconjugants[11,13,16]. One noteworthy conjugation assay from Johnson & Kroer also embraced the stochastic nature of conjugation[21]. Similar to one aspect of our approach, they survey a series of mating cultures for the presence of transconjugants and use the fraction of transconjugant-free cultures to derive a conjugation metric. However, their metric yields the number of conjugation events and does not provide an estimate for the conjugation rate. Thus, the population dynamics of donors and recipients, as well as the incubation time, are not considered. The LDM captures positive features of both the Johnson & Kroer approach and the deterministic-based approaches: specifically, it both recognizes the stochastic nature of conjugation and provides an analytic estimate for the conjugation rate.

Beyond the incorporation of stochasticity, the model and derivation behind the LDM estimate relaxes assumptions that constrained former approaches. This makes calculating conjugation rates accessible to a wide range of experimenters that use different plasmid-donor-recipient combinations. The LDM estimate makes no restrictive assumptions about growth rates or conjugation rates of different bacterial strains. Other conjugation rate estimates assume little to no change in the density due to growth (TDR), growth rates are identical across all strains (TDR and SIM), conjugation rates are identical for donors and transconjugants (SIM), or conjugation events from transconjugants are few (TDR and ASM)[11,13,16]. By relaxing these assumptions, we have increased the applicability of this new estimate, and our simulation results confirmed the high accuracy of the LDM under such scenarios. In contrast, the other conjugation methods (TDR, SIM,

and ASM) sometimes yielded inaccurate estimates when their underlying assumptions were violated. Because violations of these assumptions are likely in natural plasmid systems, these performance differences could be important. For instance, unequal growth rates can occur among isogenic strains if the plasmid increases or decreases the growth rate of a host[5,24]. Also, growth rates may be significantly different when the donor and recipient are different strains or species. For example, some strains of *Escherichia coli* can divide every 20 minutes while some strains of *Shewanella oneidensis* only divide every 45 minutes[25,26]. Consequently, the growth rates of all the cell types will differ due to these combined effects in many plasmid transfer experiments. Such compounding effects could increase the error in measuring conjugation rate with the TDR and SIM estimates. Moreover, unequal growth rates are not likely to occur in isolation, but in tandem with other modeling violations such as unequal plasmid transfer rates. The consequences of such scenarios remain to be explored.

Perhaps the most significant advantage of our LDM method is the capability to accurately estimate the donor conjugation rate when it deviates from the transconjugant conjugation rate. For the previous methods, the error in estimating the donor conjugation rate increases as the transconjugant conjugation rate gets larger than the donor conjugation rate. These differences are likely to occur in plasmid transfer experiments. For instance, Dimitriu *et. al.*[27] used the TDR approach to measure donor conjugation rate where the recipient was a clone-mate or a different species. The authors observed cross-species conjugation rates that were up to 4 orders of magnitude lower than the within-species conjugation rates. If this pattern held generally, then a cross-species mating assay would violate a key assumption of previous methods, in that the transconjugant conjugation rate would be much higher than the donor conjugation rate. Indeed, the "unequal conjugation rates" simulations resulted in inflated conjugation estimates for the other methods (Figure 2b). For these reasons, reported cross-species conjugation rates may also be overestimates. In any case, unequal conjugation rates are likely, especially in microbial communities with diverse species. In addition, unequal conjugation rates can arise due to a molecular mechanism encoded on the conjugative plasmid, such as transitory de-repression[13,28]. In this case, a conjugation event between donor and recipient will temporarily elevate the conjugation rate of the newly formed transconjugant. This phenomenon would lead to similar outcomes as described above for different

species. The LDM method can accurately estimate donor conjugation rates in systems with unequal conjugation rates, whether the differences are taxonomic or molecular in origin.

The LDM estimate also has advantages in terms of precision. Since conjugation is a stochastic process, the number of transconjugants at any given time is a random variable with a certain distribution. Therefore, estimates that rely on the number of transconjugants (TDR, SIM and ASM) or the probability of their absence (LDM) will also fall into distributions (Figure 2). Even in cases where the mean (first moment of the distribution) is close to the actual conjugation rate, the variance (second central moment) may differ among estimates. In fact, the LDM estimate had smaller variance compared to other estimates, even under parameter settings where different estimates shared similar accuracy (Figure 2). This greater precision likely originates from the difference in the distribution of the number of transconjugants ($T_{\tilde{t}}$) and the distribution of the probability of transconjugant absence ($p_0(\tilde{t})$), something we explore analytically in SI section 14. Beyond the mean and variance, other features of these distributions (i.e., moments) may also be important. For certain parameter settings, the estimates relying on transconjugant numbers were asymmetric (the third moment was non-zero). In such cases, a small number of replicate estimates could lead to bias. For example, when there is a strong growth benefit to carrying the plasmid, the distribution for the ASM estimate is skewed even though the mean of this distribution reflects the actual conjugation rate (see Figure 2a). Typically a small number of conjugation assays is often the standard in studies; thus, the general position and shape of these estimate distributions may matter. Over the portion of parameter space we explored, the LDM distribution facilitated accurate and precise estimates through its position (a mean reflecting the true value) and its shape (a small variance and a low skew).

For all the methods considered, the distribution of conjugation estimates changed with time (Figure 3). Therefore, the incubation time becomes an important parameter. In our simulations, the estimate distribution generally narrowed with longer incubation times (Figure 3). This would suggest that one advantage of longer incubation is greater precision. However, when underlying assumptions are violated, there can be an interesting interaction between accuracy and precision over time. For instance, consider the case where transconjugant conjugation rate exceeds donor conjugation rate (Figure

3b). For the SIM and ASM metrics, as incubation time rises, there is a shift from an accurate imprecise estimate to an inaccurate precise estimate. For an assay with a limited number of replicates in such a case, the optimal incubation time may be intermediate (a topic worthy of formal investigation). Over a range of incubation times where $p_0(\tilde{t})$ is not close to 0 or 1, the LDM distribution is generally immune to these problems, exhibiting constant high accuracy and precision over time (Figure 3).

To execute a conjugation assay using any of the methods described, the experimenter needs to make several decisions regarding various protocol features such as incubation time and density of the mating culture. All of these choices need to conform closely to the modeling assumptions of each method and ideally maximize the accuracy of the conjugation rate estimate. For instance, the TDR assay seems simpler than the other methods, since it only requires measurement of the final densities of cells. However, implementing the TDR assay can be difficult because it assumes little to no change in density due to growth. For a growth-based TDR implementation, a very short incubation time could limit cell growth, but the density of the culture may need to be very high so transconjugants can form during the assay. Previous studies have shown that for many plasmids the conjugation rate is maximal during exponential phase[13,17,29]; therefore, using high density culture may underestimate the conjugation rate. A chemostat implementation could circumvent some of these issues by allowing lower density culture, exponential growth, and longer incubation times. However, unequal growth rates in a chemostat would lead to changes in the density of the donors and recipients. For the SIM approach, a standard growth-based 24-hour conjugation assay is often used[5,30,31,32,33]. This serves as an easy default choice and has many laboratory benefits for efficiency. Even so, the 24-hour incubation involves more than initial and final sampling, as two additional samplings are required to measure the maximum population growth rate during exponential phase (SI section 2). If these additional samplings are forgotten for the SIM approach, highly inaccurate estimates of the conjugation rate can result. In addition, the underlying SIM assumptions of equal growth rates and conjugation rates are easy to violate, and inaccuracies due to such violations can grow with incubation time (Figure 3b). For the ASM, a standard incubation time is not recommended; rather, this time is left to the experimenter to determine. However, the authors suggest short incubation times to prevent biased estimates if modeling assumptions are violated (e.g., relatively high

transconjugant transfer rates). The authors mention that a follow-up assay to measure the transconjugant transfer rate may be needed to determine if the incubation time was short enough to prevent deviations. In contrast, the LDM protocol is designed to lead the experimenter to valid incubation times by explicitly identifying the conjugation window where $p_0(\tilde{t})$ can be estimated. Therefore, the experimenter can be sure the protocol occurred during a usable incubation time if the assay results in a $\hat{p}_0(\tilde{t})$ between 0 and 1. We note that identifying the conjugation window does add an extra step in the LDM approach; however, a similar assay could help an experimenter choose an optimal incubation time for the other assays. Even so, the effects of these implementation choices on the accuracy and precision of conjugation rate estimates are currently understudied.

The LDM protocol offers an additional advantage by eliminating plating of the transconjugants. For the other methods, this is typically a source of error since the transconjugant density is often determined by plating the mating culture on transconjugant selecting medium. With short culture incubation times before plating, which will increase the accuracy of the other methods, the transconjugant density is extremely low compared to the donor and recipient densities. Therefore, plating high density cultures is needed to be able to detect the transconjugants. This can lead to conjugation continuing on the agar plates, which produces extra transconjugant colonies that did not result from conjugation events before plating[,15,34]. Such 'matings on transconjugant-selecting agar plates' can occur before the donors and recipients are completely inhibited by the transconjugant-selecting agent (typically an antibiotic). Such events will lead to an inflation of the transconjugant density and thus inaccuracy in the conjugation rate estimate. We note that the LDM avoids this problem since it relies only on recording transconjugant presence/absence in the liquid mating cultures.

We applied our high-throughput LDM protocol to our previous evolution experiment to estimate the conjugation rate for an IncP-1β plasmid in an *E. coli* host. Interestingly, we observed the evolution of conjugation rate. This comes as a bit of a surprise given that the environment in which the *E. coli* evolved contained an antibiotic selecting for plasmid carriage. Thus, plasmid-free cells would have been rare leading to few opportunities for plasmid conjugation into such cells. Bacterial cells containing a conjugative plasmid in a community with an abundance of plasmid-free cells is the scenario in which conjugation rates are expected to and have been observed to evolve[35].

Therefore, it is not immediately obvious what mechanism led to the increase in the
conjugation rate in our system. It is possible that increased conjugation rate could have
been a pleiotropic effect from a different trait under direct selection. Alternatively, higher
conjugation rate could evolve if (1) a mutation on the plasmid led to a higher transfer rate
and (2) if donor cells can conjugate with other donors in our systems[36]. In such a case,
the mutated plasmid would transfer quickly even under plasmid-selecting conditions and
it could spread through the population as long as any associated costs (e.g., lower growth
of its host or lower replication efficiency within mixed-plasmid hosts) were sufficiently low.
This hypothetical sequence of events could lead to an increased conjugation rate.
Regardless of the mechanism, it is interesting that bacterial populations in environments
containing an antibiotic could lead to the evolution of higher conjugation rates of plasmids
encoding resistance to that antibiotic. Importantly, such evolution could lead to higher
frequency of antibiotic resistance in microbial communities, even in the absence of the
antibiotic, due to more plasmid conjugation events.

In conclusion, the LDM offers new possibilities for measuring the conjugation rate
for many types of plasmids and species. We have presented evidence that supports our
method being more accurate and precise than other widely used approaches. We note
that we did not fully review all the conjugation assays currently in use in the literature[15,37],
focusing instead on the least laborious and most widely used methods. Importantly, the
LDM eliminates the possible bias caused by high transconjugant conjugation rates. Due
to the sensitivity of the LDM conjugation assay and the ease of implementation, we were
able to show the evolution of conjugation rate for an IncP-1β conjugative plasmid in an
Escherichia coli host. However, the LDM method can be applied even more broadly given
the flexibility of the underlying theoretical framework and the relative ease of assay
implementation.

Acknowledgements
This work is supported by the National Institute of Allergy and Infectious Diseases
Extramural Activities grant no. R01 AI084918 of the National Institutes of Health. O.K. is
supported by the NSF Graduate Research Fellowship grant no. DGE-1762114. C.E. is
supported by the NSF Graduate Research Fellowship. We thank Hannah Jordt, Simon

Snoeck, and members of the Kerr and Top laboratories for useful suggestions on the
manuscript.

Supplemental Information

SI section 1 : Historical theoretical perspective

In this section, we highlight three key methods for estimating conjugation rate.
While outlining the theoretical framework, we highlight the key distinctions and theoretical
assumptions of each approach. Levin *et. al.*[13] introduced a simple mathematical model
for plasmid population dynamics to isolate the conjugation rate parameter (γ)[13]. In their
model, the donor population grows exponentially at a rate ψ (equation [1]). The number
of transconjugants increases due to conjugation events (2nd term equation [3]) by donors
and by existing transconjugants as well as exponential growth of the existing
transconjugant pool (1st term equation [3]). The recipient population grows exponentially
at a rate ψ and is depleted at the rate that new transconjugants form due to conjugation
events (2nd term equation [2]). For ease of reference, model variables and parameters
are in SI Table 1 (the LDM is included for a complete comparison).

SI Table 1: Variables and parameters used in plasmid dynamic models

Variable/Parameter	Description	Relevant Estimate	Units
D	Donor density	TDR, SIM, ASM, LDM	$\dfrac{\text{cfu}}{\text{ml}}$
R	Recipient density	TDR, SIM, ASM, LDM	
T	Transconjugant density	TDR, SIM, ASM, LDM	
ψ	Growth rate (not population specific)	TDR, SIM	$\dfrac{1}{\text{h}}$
ψ_D	Donor growth rate	ASM, LDM	
ψ_R	Recipient growth rate	ASM, LDM	
ψ_T	Transconjugant growth rate	ASM, LDM	
γ	Conjugation rate (not population specific)	TDR, SIM	$\dfrac{\text{ml}}{\text{cfu} \cdot \text{h}}$
γ_D	Donor-recipient conjugation rate	ASM, LDM	
γ_T	Transconjugant-recipient conjugation rate	ASM, LDM	

Equations [1] - [3] contains four notable assumptions. First, conjugation is

described by mass-action kinetics, where conjugation events are proportional to the

product of donor and recipient cell densities, which is a reasonable assumption in well-

mixed liquid cultures[13]. Second, the model assumes a negligible rate of segregation, a

process whereby a dividing plasmid-containing cell produces one plasmid-containing

daughter cell and one plasmid-free daughter. These first two assumptions also exist in all

of the other conjugation estimate methods we discuss. Third, the growth rate is the same

for all cell types (donors, recipients, and transconjugants). Fourth, the plasmid

conjugation rate is the same from donors to recipients as from transconjugants to

recipients.

Levin *et. al.*[13] used their plasmid dynamic model to derive an estimate for plasmid

conjugation rate (γ) by making three additional simplifying assumptions. First, the change

in cell density of donors due to growth is assumed to be negligible. Likewise, the change

in cell density of recipients due to growth and to conjugation (i.e., transformation into

transconjugants) is assumed to be negligible. Finally, transconjugants are assumed to be

rare in the population resulting in negligible growth of transconjugants and conjugation

from transconjugants to recipients. Thus, the increase in transconjugants cell density is

through plasmid conjugation from donors to recipients (i.e., $dT/dt = \gamma DR$). Using these

simplifying assumptions, Levin *et. al.*[13] solved for an expression of the conjugation rate γ

(equation [4]) in terms of the density of donors, recipients, and transconjugants ($D_{\tilde{t}}$, $R_{\tilde{t}}$,

and $T_{\tilde{t}}$, respectively) after a period of incubation $\tilde{t}$ (see SI section 3 for a few different

approaches to this derivation). We label the expression in equation [4] as the "TDR"

estimate for the conjugation rate, where the letters in this acronym come from the dynamic

variables used in the estimate. TDR is a commonly used estimate[14,13,15,17]. However, the

simplifying assumptions used to derive TDR, such as no change in density due to growth,

can make implementation in the laboratory difficult. For ease of reference, all variables

used in the conjugation estimates are in SI Table 2. All assumptions underlying each

estimate are in SI Table 3. Both tables include LDM for a full comparison.

SI Table 2: Variables and parameters used to estimate* conjugation rate

Variable/Parameter	Description	Relevant Estimate	Units
$\tilde{t}$	Incubation time (final sampling time)	TDR, SIM**, ASM, LDM	h
D_0, R_0	Initial donor and recipient densities	ASM, LDM	$\dfrac{\text{cfu}}{\text{ml}}$
$D_{\tilde{t}}, R_{\tilde{t}}$	Final donor and recipient densities	TDR, SIM, ASM, LDM	
$T_{\tilde{t}}$	Final transconjugant density	TDR, SIM, ASM	
$N_0, N_{\tilde{t}}$	Initial and final total population density	SIM	
ψ_T	Transconjugant growth rate	ASM	h^{-1}
$p_0(\tilde{t})$	Probability a population has no transconjugants	LDM	$\exp\left(\dfrac{\text{cfu}}{\text{ml}}\right)$

* The laboratory estimates are used here (see SI section 6)
** If the SIM assay is conducted on exponentially growing cultures (see SI section 2), $\tilde{t}$ along with N_0 and $N_{\tilde{t}}$ can be used to estimate ψ (otherwise, an independent estimate of ψ is needed).

SI Table 3: Summary of modeling assumptions

Assumption	TDR	SIM	ASM	LDM
Conjugation events follow mass-action kinetics	X	X	X	X
There is no segregative loss of the plasmid	X	X	X	X
The cell populations do not change in size due to growth	X			
Processes of conjugation and growth are not resource dependent*	X		X	X
The cell populations grow exponentially			X	X
The growth rate is identical for all cell types	X	X		
The transconjugant conjugation rate is not high relative to the donor conjugation rate	X	X	X	

* The SIM model is able to incorporate resource-dependent growth and conjugation because (1) growth and transfer rates are homogeneous and (2) the functional form for resource dependence is the same for growth and transfer.

In contrast to the assumption behind the TDR estimate, the model explored by Simonsen *et. al.*[16] allows for population growth. Indeed, they allowed for the rate of population growth to change with the level of a resource in the environment, adding a dynamic variable for the resource concentration. In addition, the conjugation rate can also change with the resource concentration. The authors focus on a case where both growth and conjugation rates vary with resource concentration according to the Monod function. This choice was informed by experimental results showing that cells enter stationary phase and conjugation ramps down to a negligible level as resources are depleted[13]. This pattern occurs for various plasmid incompatibility groups, but not all[29]. Simonsen *et. al.*[16] used this updated model to derive an estimate for plasmid conjugation rate (equation [5], see SI section 4 for the derivation). We term equation [5] as the "SIM" estimate for conjugation rate, where SIM stands for "Simonsen *et. al.* Identicality Method" since the underlying model assumes that all strains are *identical* with regards to growth, and donors and transconjugants are *identical* with regards to conjugation rate (as in equations [1]-[3]). The SIM estimate involves measuring the density of the initial population (N_0), and the final density of donors ($D_{\tilde{t}}$), recipients ($R_{\tilde{t}}$), transconjugants ($T_{\tilde{t}}$), and the total population ($N_{\tilde{t}}$) after the incubation time $\tilde{t}$ (e.g., the end of an assay). The SIM estimate is a commonly used since it allows for the use of batch culture in the laboratory. Thus, it circumvents the constraints of the laboratory implementation of TDR; however, the underlying model (an enriched version of equations [1]-[3]) holds the same assumptions as before: homogeneous growth rates and conjugation rates.

Recently, Huisman *et. al.*[11] updated the SIM model by introducing population specific growth rates for donors, recipients, and transconjugants (ψ_D, ψ_R, and ψ_T, respectively) and population specific conjugation rates for donors and transconjugants (γ_D and γ_T). Importantly, the updated model is an *approximation* of the SIM due to three additional assumptions. First, conjugation and growth rates are assumed to be constant until resources are depleted, eliminating the additional resource concentration equation. Second, the increase in recipients due to growth greatly outpaces the decrease in recipients due to conjugation (i.e., $\psi_R R \gg \gamma_D DR + \gamma_T TR$). Third, the increase in transconjugants due to growth or plasmid conjugation from donors to recipients greatly outpaces the increase in transconjugants due to plasmid conjugation from transconjugants to recipients (i.e., $\psi_T T + \gamma_D DR \gg \gamma_T TR$). These model conditions are

reasonable if the system starts with donors and recipients present but transconjugants are absent, the system is tracked for a short period of time $\tilde{t}$, conjugation rates are low relative to growth rates, and the transconjugant conjugation rate (γ_T) is not much higher than the donor conjugation rate (γ_D). With these added assumptions, equations [1]-[3] can be reformulated as the following *approximate* system of equations:

$$\frac{dD}{dt} = \psi_D D, \qquad\qquad [1.1]$$

$$\frac{dR}{dt} = \psi_R R, \qquad\qquad [1.2]$$

$$\frac{dT}{dt} = \psi_T T + \gamma_D DR, \qquad\qquad [1.3]$$

Huisman *et. al.* use this model to derive an estimate for conjugation rate (equation [6]) where different cell types now can have different growth rates (see SI section 5 for the derivation). We term equation [6] as the ASM estimate for donor conjugation rate, where ASM stands for "Approximate Simonsen *et. al.* Method".

SI section 2 : Comparison of laboratory implementations

In this section, we compare the laboratory implementations for various end-point estimates: TDR, SIM, and ASM. Each method is explained either as recommended by its authors or the most simplified protocol to acquire the information for the estimate. For each, we describe proper laboratory implementation for the approaches based on the model and derivation assumptions used to acquire the estimate. Note in this section, we do not explore the assumptions that are violated due to the biological samples being tested (i.e., specific species or plasmids) which can result in violations such as unequal conjugation rates or growth rates. These are explored in the main text via stochastic simulations. Thus, we focus solely on the parameters under the experimenter's control. For ease of reference, key implementation differences are highlighted in SI Table 4 (LDM is included for a full comparison).

SI Table 4: Comparison of implementations

Summary	TDR	SIM	ASM	LDM
Assay conditions minimizing the change in density due to growth	X			
Minimize incubation time necessary for producing transconjugants			X	

An incubation time results in a subset of parallel populations having no transconjugants				X
Assay occurs over a period of exponential cell growth		X*	X	X
Assay requires multiple parallel mating cultures per replicate				X
Assay requires a measurement of transconjugant density	X	X	X	
Assay requires a measurement of population growth rate		X*		
Assay requires a measurement of transconjugant growth rate			X	
* For the SIM assay, either the entire assay is conducted over exponentially growing cultures or an independent estimate for (maximum) population growth rate is needed.				

The TDR estimate has a simple form (equation [4]). Donors and recipients are

mixed in non-selective growth medium and incubated for a specified time $\tilde{t}$. Typically,

densities after the incubation time are determined using selective plating. The derivation

assumes the density in donors and recipients does not change which sets specific

constraints on incubation. In the original study, Levin *et. al.*[9], used a chemostat to keep

the population constant. Other studies shorten the incubation time $\tilde{t}$ such that population

growth is negligible and use various laboratory tools to detect the small number of

transconjugants[15,17].

SIM circumvents the incubation limitations. Under conditions where the population

is growing exponentially over the entire assay, we can rework the SIM estimate (equation

[5]) into an alternative form for the laboratory (see SI section 6 for details).

$$\gamma = \frac{1}{\tilde{t}}\left[\ln\left(1 + \frac{T_{\tilde{t}}}{R_{\tilde{t}}}\frac{N_{\tilde{t}}}{D_{\tilde{t}}}\right)\right]\frac{\ln N_{\tilde{t}} - \ln N_0}{N_{\tilde{t}} - N_0} \qquad [2.1]$$

Donor and recipient populations in exponential phase are mixed in non-selective growth

medium. The initial population density (N_0) is determined by dilution plating on non-

selective medium. After the mating mixture is incubated (for a period of $\tilde{t}$), the final

densities ($D_{\tilde{t}}$, $R_{\tilde{t}}$, $T_{\tilde{t}}$, and $N_{\tilde{t}}$) are determined by dilution plating on selective and non-

selective media. To implement SIM as written in equation [2.1], the specified incubation

time $\tilde{t}$ must occur well before stationary phase is reached to collect proper data for

estimating the population growth rate ($\psi = (\ln N_{\tilde{t}} - \ln N_0)/\tilde{t}$). There is an alternative

option for implementing the SIM using equation [5]. The donor and recipient populations

are mixed and incubated under batch culture conditions (including lag, exponential, and

stationary phase). However, the (maximum) population growth rate (ψ) needs to be

determined with two additional samplings from the mixed population at times t_a and t_b,

both occurring *within exponential phase:*

$$\psi = \frac{\ln{(N_{t_b}/N_{t_a})}}{t_b - t_a} \tag{2.2}$$

The population densities N_{t_a} and N_{t_b} can be estimated either through colony counts from

plating or optical density from a spectrophotometer. Either way, the timing of exponential

phase is important for this approach and at least some analysis during this phase is

required regardless of the implementation strategy.

For the ASM estimate, we can rework equation [6] into an alternate form for the

laboratory (see SI section 6 for details).

$$\gamma_D = \left\{\frac{1}{\tilde{t}}\left(\ln D_{\tilde{t}} R_{\tilde{t}} - \ln D_0 R_0\right) - \psi_T\right\}\frac{T_{\tilde{t}}}{\left(D_{\tilde{t}} R_{\tilde{t}} - D_0 R_0 e^{\psi_T \tilde{t}}\right)} \tag{2.3}$$

Donor and recipient populations in exponential phase are mixed in non-selective medium.

Initial densities (D_0 and R_0) are determined by plating dilutions on the appropriate

selective media. After donor and recipient incubate ($\tilde{t}$), final densities ($D_{\tilde{t}}$, $R_{\tilde{t}}$, and $T_{\tilde{t}}$) are

determined by plating dilutions on the appropriate selective media. From the

transconjugant-selecting agar plates, a transconjugant clone is incubated in monoculture

then sampled twice (at times t_a and t_b) in exponential phase to measure the

transconjugant growth rate ($\psi_T = \ln\left(T_{t_b}/T_{t_a}\right)/(t_b - t_a)$). The authors point out a critical

consideration for proper implementation of the ASM is the incubation time $\tilde{t}$. Not only is

sampling in exponential phase important, but if the incubation time $\tilde{t}$ is too long and

passes a critical time (t_{crit}) the approximations used to derive the ASM break down. To

avoid overshooting t_{crit}, the authors recommend sampling as soon as measurable

transconjugants arise. To determine that the incubation time used was below the critical

time (t_{crit}), a second assay is recommended by the authors to measure the

transconjugant conjugation rate γ_T, which will determine if the original incubation time $\tilde{t}$

was below t_{crit} for measuring the donor conjugation rate. This second assay would have

the transconjugant clone become the donor in the mixture, while a newly marked recipient

must be used so that donors and recipients can be distinguished using selective plating.

Each method has aspects of implementation in common. Each one shares the

basic approach of mixing donors and recipients over some incubation time $\tilde{t}$. Each

estimate requires reliable selectable markers to differentiate donors, recipients, and

transconjugants. However, all estimates have some constraints on initial densities and

time of measurement. This can occur because the experimenter needs to capture

conjugation events (all estimates require this), avoid population growth (TDR), or keep

growth exponential (ASM, and at least parts of SIM). Even so, each method has clear

distinctions. The most notable is the incubation time $\tilde{t}$ (i.e., the end of the assay). The

TDR method is constrained to conditions where no change in population size due to

growth can occur. For SIM, initial and final sampling are not constrained to a particular

phase of growth; however, measurement of the growth rate must occur during the

exponential growth phase. For ASM, initial sampling is in early exponential phase, and

the final sampling needs to occur during a specific time window. In other words, the assay

needs to be long enough that measurable transconjugants appear, but short enough so

that assumptions aren't violated (which can occur if transconjugant density becomes too

large).

SI section 3 : TDR derivation

We present a few derivations for the TDR method. The derivations differ by the

assumptions used to solve for TDR. We use this section to show all of the conditions

where the TDR method can be used. In addition, we use this section to walk through all

of the mathematical steps and provide an easy reference for comparing the derivations

of all four methods (TDR, SIM, ASM, and LDM). The original condition presented by Levin

et. al.[13] is the last derivation described in this section.

Here we assume that we are considering conditions where population growth is

not occurring. These conditions could involve an environment hindering cell division.

Alternatively, the environment could promote growth, but we restrict the time period

sufficiently so that population change is negligible. Or finally, if all strains possess the

same growth rate, the population could be growing over longer periods of time but doing

so in a continuous flow-through device (e.g., a chemostat) such that population density

remains constant (see Levin *et. al.*[13]). Because we assume no segregative loss of the

plasmid, the donor population must remain constant

$$D_t = D_0, \qquad\qquad [3.1]$$

for any time t under consideration. Importantly, even though growth is not occurring,

conjugation can proceed. We assume $T_0 = 0$, but the population of transconjugants can

increase over time. Because every transconjugant was formerly a recipient cell, it must

be the case that

$$R_t + T_t = R_0,$$

for any time t under consideration. Therefore,

$$R_t = R_0 - T_t. \qquad [3.2]$$

839   The dynamics of transconjugants is given by

$$\frac{dT_t}{dt} = \gamma_D D_t R_t + \gamma_T T_t R_t \qquad [3.3]$$

840   By substituting terms from equations [3.1] and [3.2]

841
$$\frac{dT_t}{dt} = \gamma_D D_0 (R_0 - T_t) + \gamma_T T_t (R_0 - T_t),$$

842
$$\frac{dT_t}{dt} = (\gamma_D D_0 + \gamma_T T_t)(R_0 - T_t),$$

843
$$\frac{dT_t}{dt} = -\gamma_T \left(T_t + \frac{\gamma_D D_0}{\gamma_T} \right)(T_t - R_0),$$

844   We can solve this differential equation by a separation of variables:

845
$$\int_0^{\tilde{t}} \frac{dT_t}{-\gamma_T \left(T_t + \frac{\gamma_D D_0}{\gamma_T} \right)(T_t - R_0)} = \int_0^{\tilde{t}} dt.$$

846   The following identity is relevant here:

847
$$\frac{d\left\{ \frac{1}{a(b-c)} \ln \frac{x-b}{x-c} \right\}}{dx} = \frac{1}{a(x-b)(x-c)}.$$

848   Letting $x = T_t$, $a = -\gamma_T$, $b = -\frac{\gamma_D D_0}{\gamma_T}$, and $c = R_0$, we can proceed as follows:

849
$$\left\{ \frac{1}{\gamma_T \left(\frac{\gamma_D D_0}{\gamma_T} + R_0 \right)} \ln \frac{T_t + \frac{\gamma_D D_0}{\gamma_T}}{T_t - R_0} \right\} \Bigg|_0^{\tilde{t}} = (t)|_0^{\tilde{t}},$$

850
$$\frac{1}{\gamma_T \left(\frac{\gamma_D D_0}{\gamma_T} + R_0 \right)} \left(\ln \frac{T_{\tilde{t}} + \frac{\gamma_D D_0}{\gamma_T}}{T_{\tilde{t}} - R_0} - \ln \frac{T_0 + \frac{\gamma_D D_0}{\gamma_T}}{T_0 - R_0} \right) = \tilde{t}.$$

851   Because $T_0 = 0$,

852
$$\frac{1}{\gamma_T \left(\frac{\gamma_D D_0}{\gamma_T} + R_0 \right)} \left(\ln \frac{T_{\tilde{t}} + \frac{\gamma_D D_0}{\gamma_T}}{T_{\tilde{t}} - R_0} - \ln \frac{\frac{\gamma_D D_0}{\gamma_T}}{-R_0} \right) = \tilde{t},$$

853
$$\frac{1}{\gamma_T \left(\frac{\gamma_D D_0}{\gamma_T} + R_0 \right)} \left(\ln \frac{-R_0 \left(T_{\tilde{t}} + \frac{\gamma_D D_0}{\gamma_T} \right)}{\frac{\gamma_D D_0}{\gamma_T}(T_{\tilde{t}} - R_0)} \right) = \tilde{t},$$

$$\ln\frac{1+\dfrac{\gamma_T T_{\tilde{t}}}{\gamma_D D_0}}{1-\dfrac{T_{\tilde{t}}}{R_0}} = (\gamma_D D_0 + \gamma_T R_0)\tilde{t},\qquad[3.4]$$

854
$$\frac{1+\dfrac{\gamma_T T_{\tilde{t}}}{\gamma_D D_0}}{1-\dfrac{T_{\tilde{t}}}{R_0}} = \exp\{(\gamma_D D_0 + \gamma_T R_0)\tilde{t}\},$$

855
$$1+\frac{\gamma_T T_{\tilde{t}}}{\gamma_D D_0} = \left(1-\frac{T_{\tilde{t}}}{R_0}\right)\exp\{(\gamma_D D_0 + \gamma_T R_0)\tilde{t}\},$$

856
$$\frac{\gamma_T T_{\tilde{t}}}{\gamma_D D_0} + \frac{T_{\tilde{t}}}{R_0}\exp\{(\gamma_D D_0 + \gamma_T R_0)\tilde{t}\} = \exp\{(\gamma_D D_0 + \gamma_T R_0)\tilde{t}\} - 1,$$

857
$$T_{\tilde{t}}\left[\frac{\gamma_T}{\gamma_D D_0} + \frac{1}{R_0}\exp\{(\gamma_D D_0 + \gamma_T R_0)\tilde{t}\}\right] = \exp\{(\gamma_D D_0 + \gamma_T R_0)\tilde{t}\} - 1,$$

$$T_{\tilde{t}} = \frac{R_0(\exp\{(\gamma_D D_0 + \gamma_T R_0)\tilde{t}\} - 1)}{\dfrac{\gamma_T R_0}{\gamma_D D_0} + \exp\{(\gamma_D D_0 + \gamma_T R_0)\tilde{t}\}}.\qquad[3.5]$$

858 Thus, equation [3.5] is a general solution for the number of transconjugants at any time.

859 If we assume that $\gamma_T = 0$, then we can rewrite [3.4] as

860
$$\ln\frac{1}{1-\dfrac{T_{\tilde{t}}}{R_0}} = \gamma_D D_0 \tilde{t},$$

861
$$-\ln\left(1-\frac{T_{\tilde{t}}}{R_0}\right) = \gamma_D D_0 \tilde{t},$$

$$\gamma_D = \frac{-\ln\left(1-\dfrac{T_{\tilde{t}}}{R_0}\right)}{D_0 \tilde{t}}.\qquad[3.6]$$

862 When $R_0 \gg T_{\tilde{t}}$, a first-order Taylor approximation ensures $-\ln\left(1-\frac{T_{\tilde{t}}}{R_0}\right) \approx \frac{T_{\tilde{t}}}{R_0}$, and

863 therefore

864
$$\gamma_D \approx \frac{\dfrac{T_{\tilde{t}}}{R_0}}{D_0 \tilde{t}},$$

865
$$\gamma_D \approx \frac{T_{\tilde{t}}}{D_0 R_0 \tilde{t}}.$$

866 Because $D_{\tilde{t}} = D_0$ (see [3.1]) and $R_{\tilde{t}} \approx R_0$ (when $R_0 \gg T_{\tilde{t}}$ by [3.2]), we have

867
$$\gamma_D \approx \frac{T_{\tilde{t}}}{D_{\tilde{t}} R_{\tilde{t}} \tilde{t}}.$$

868 On the other hand, if we assume $\gamma_D = \gamma_T = \gamma$, then we can rewrite [3.4] as

869
$$\ln \frac{1 + \frac{T_{\tilde{t}}}{D_0}}{1 - \frac{T_{\tilde{t}}}{R_0}} = \gamma(D_0 + R_0)\tilde{t},$$

870
$$\gamma = \frac{1}{\tilde{t}(D_0 + R_0)} \ln \frac{1 + \frac{T_{\tilde{t}}}{D_0}}{1 - \frac{T_{\tilde{t}}}{R_0}},$$

$$\gamma = \frac{1}{\tilde{t}(D_0 + R_0)} \left\{ \ln\left(1 + \frac{T_{\tilde{t}}}{D_0}\right) - \ln\left(1 - \frac{T_{\tilde{t}}}{R_0}\right) \right\}. \qquad [3.7]$$

871   When $D_0 \gg T_{\tilde{t}}$ and $R_0 \gg T_{\tilde{t}}$, first-order Taylor approximations ensure

872
$$\gamma \approx \frac{1}{\tilde{t}(D_0 + R_0)} \left(\frac{T_{\tilde{t}}}{D_0} + \frac{T_{\tilde{t}}}{R_0}\right),$$

873
$$\gamma \approx \frac{T_{\tilde{t}}}{\tilde{t}(D_0 + R_0)} \left(\frac{1}{D_0} + \frac{1}{R_0}\right),$$

874
$$\gamma \approx \frac{T_{\tilde{t}}}{\tilde{t}(D_0 + R_0)} \left(\frac{R_0 + D_0}{D_0 R_0}\right),$$

875
$$\gamma \approx \frac{T_{\tilde{t}}}{D_0 R_0 \tilde{t}}.$$

876   Because $D_{\tilde{t}} = D_0$ and $R_{\tilde{t}} \approx R_0$ (when $R_0 \gg T_{\tilde{t}}$), we have

877
$$\gamma \approx \frac{T_{\tilde{t}}}{D_{\tilde{t}} R_{\tilde{t}} \tilde{t}}.$$

878   So we have general expressions for donor conjugation rate when $\gamma_T = 0$ (equation [3.6])

879   or when $\gamma_D = \gamma_T$ (equation [3.7]). However, when $D_0 \gg T_t$ and $R_0 \gg T_t$, for all t under

880   consideration, both of these measures are well approximated by equation [4]. Here we

881   extend the application of equation [4] even further. When $R_0 \gg T_t$, then $R_t \approx R_0$. Let us

882   assume $R_t = R_0$. We will also assume

$$\gamma_D D_0 R_0 \gg \gamma_T T_t R_0, \qquad [3.8]$$

namely, the rate of formation of transconjugants by donors is much greater than the

formation by transconjugants. Of course, if $\gamma_T = 0$ or $\gamma_D = \gamma_T$, then $D_0 \gg T_t$ ensures

assumption [3.8]. However, assumption [3.8] does not require $\gamma_T = 0$ or $\gamma_D = \gamma_T$ (indeed,

assumption [3.8] is satisfied when $\gamma_D \geq \gamma_T$ and $D_0 \gg T_t$, and $R_0 > 0$). In general, this

inequality is satisfied when $T_0 = 0$, D_0 and R_0 are is large, γ_T is not dramatically higher

than γ_D, and the period is small. Under assumption [3.8], the dynamics can be well

approximated by a simplified version of the differential equation [3.3] (where the

transconjugant conjugation term is gone). This is the differential equation originally solved
by Levin *et. al.*,

$$\frac{dT_t}{dt} = \gamma_D D_0 R_0.$$

893   Because everything on the right-hand-side of the equation is a constant, the solution is
894   straightforward:

$$\int_0^{\tilde{t}} dT_t = \int_0^{\tilde{t}} \gamma_D D_0 R_0 dt,$$

$$(T_{\tilde{t}})|_0^{\tilde{t}} = (\gamma_D D_0 R_0 t)|_0^{\tilde{t}},$$

$$T_{\tilde{t}} - T_0 = \gamma_D D_0 R_0 \tilde{t}.$$

898   Because $T_0 = 0$,

$$T_{\tilde{t}} = \gamma_D D_0 R_0 \tilde{t},$$

$$\gamma_D = \frac{T_{\tilde{t}}}{D_0 R_0 \tilde{t}}.$$

Because $D_{\tilde{t}} = D_0$ and $R_{\tilde{t}} = R_0$ (by assumption), we have recovered equation [4].

SI section 4 : SIM derivation

In this section, we walk through all of the mathematical steps in the Simonsen *et.*
al. derivation. We include this derivation as a quick reference to help the reader compare
the derivations of all four methods (TDR, SIM, ASM, and LDM). Simonsen *et. al.* modified
the model (equations [1]-[3]) by adding a dynamic variable for resource concentration (C).
The dynamics of this resource incorporate an additional parameter for the amount of
resource needed to produce a new cell (e). Both the growth rate and conjugation rate are
taken to be functions of the resource concentration.

$$\frac{dD}{dt} = \psi(C)D, \qquad\qquad [4.1]$$

$$\frac{dR}{dt} = \psi(C)R - \gamma(C)R(D + T), \qquad\qquad [4.2]$$

$$\frac{dT}{dt} = \psi(C)T + \gamma(C)R(D + T), \qquad\qquad [4.3]$$

$$\frac{dC}{dt} = -\psi(C)(R + D + T)e. \qquad\qquad [4.4]$$

The other variables are consistent with their use in equations [1]-[3]. In Simonsen *et. al.*[16],
when resources are depleted, growth and conjugation stop. A Monod function introduces
batch culture dynamics (i.e., exponential and stationary phase) making growth and

conjugation both increase in a saturated manner with resource concentration, where the

resource concentration yielding ½ the maximal rate is given by the parameter Q.

Importantly, conjugation and growth are assumed to have the same functional form:

917
$$\psi(C) = \frac{\psi_{max}C}{Q + C},$$

918
$$\gamma(C) = \frac{\gamma_{max}C}{Q + C},$$

919   where Q is the half saturation constant.

920   Here we derive the estimation of conjugation rate from Simonsen *et. al.*[16] focusing

921   on the exponential phase of growth where resources are common (C is large) thus the

922   following are reasonable assumptions:

923
$$\psi(C) = \psi_{max}$$

924
$$\gamma(C) = \gamma_{max}$$

Under these assumptions, conjugation rate (γ) and growth rate (ψ) are constant. As

resources are assumed to remain abundant such that there are no changes to the rates

of growth and conjugation, equation [4.4] can be dropped. Simonsen *et. al.*[16] assume that

all strains have the same growth rate ($\psi_D = \psi_R = \psi_T = \psi_{max}$) and conjugation rates from

donors and transconjugants are the same ($\gamma_D = \gamma_T = \gamma_{max}$). Additionally, Simonsen *et.*

al.[16] assume no segregative loss. Thus, with our assumption of exponential growth, we

recover equations [1]-[3]. We define $N = D + R + T$. Therefore, using equations [1]-[3]

932
$$\frac{dN}{dt} = \frac{dD}{dt} + \frac{dR}{dt} + \frac{dT}{dt},$$

933
$$\frac{dN}{dt} = \psi_{max}D + \psi_{max}R - \gamma_{max}R(D + T) + \psi_{max}T + \gamma_{max}R(D + T),$$

934
$$\frac{dN}{dt} = \psi_{max}(D + R + T),$$

$$\frac{dN}{dt} = \psi_{max}N. \hspace{2cm} [4.5]$$

935   The general solutions for equation [1] and [4.5] are found by (1) separating variables, (2)

936   integrating, and (3) solving for D_t or N_t, respectively.

937
$$D_t = D_0 e^{\psi_{max}t},$$

938
$$N_t = N_0 e^{\psi_{max}t}.$$

939   Define the fraction of donors in the population to be $X_t = D_t/N_t$. We note

940
$$X_t = \frac{D_t}{N_t} = \frac{D_0 e^{\psi_{max}t}}{N_0 e^{\psi_{max}t}} = \frac{D_0}{N_0} = X_0,$$

941   Thus, this fraction does not change over time (i.e., X_t is a constant). Lastly, we define a

942   fraction $Y_t = T_t/R_t$. Using equations [2] and [3],

943
$$\frac{dY}{dt} = \frac{d(T/R)}{dt} = \frac{\frac{dT}{dt}R - \frac{dR}{dt}T}{R^2},$$

944
$$\frac{dY}{dt} = \frac{\{\psi_{max}T + \gamma_{max}R(D+T)\}R - \{\psi_{max}R - \gamma_{max}R(D+T)\}T}{R^2},$$

945
$$\frac{dY}{dt} = \frac{\psi_{max}TR + \gamma_{max}(D+T)R^2 - \psi_{max}TR + \gamma_{max}(D+T)TR}{R^2},$$

946
$$\frac{dY}{dt} = \frac{\gamma_{max}(D+T)R^2 + \gamma_{max}(D+T)TR}{R^2},$$

947
$$\frac{dY}{dt} = \frac{\gamma_{max}(D+T)R + \gamma_{max}(D+T)T}{R},$$

948
$$\frac{dY}{dt} = \frac{\gamma_{max}(DR + TR + DT + T^2)}{R},$$

949
$$\frac{dY}{dt} = \gamma_{max}\frac{DT + RT + T^2 + DR}{R},$$

950
$$\frac{dY}{dt} = \gamma_{max}\left(\frac{T(D+R+T)}{R} + D\right).$$

951   Because $N = D + R + T$,

952
$$\frac{dY}{dt} = \gamma_{max}\left(\frac{TN}{R} + D\right),$$

953
$$\frac{dY}{dt} = \gamma_{max}N\left(\frac{T}{R} + \frac{D}{N}\right).$$

954   Because $X = D/N$ and $Y = T/R$,

955
$$\frac{dY}{dt} = \gamma_{max}N(Y + X),$$

956
$$\left(\frac{1}{Y+X}\right)\frac{dY}{dt} = \gamma_{max}N.$$

957   Equation [4.5] ensures $N = \left(\frac{1}{\psi_{max}}\right)\frac{dN}{dt}$, which yields

958
$$\left(\frac{1}{Y+X}\right)\frac{dY}{dt} = \gamma_{max}\left(\frac{1}{\psi_{max}}\right)\frac{dN}{dt},$$

959
$$\psi_{max}\left(\frac{1}{Y+X}\right)\frac{dY}{dt} = \gamma_{max}\frac{dN}{dt},$$

960
$$\gamma_{max}(dN) = \psi_{max}\left(\frac{dY}{Y+X}\right).$$

961   To emphasize that X is a constant, we write this as X_0 and then integrate both sides,

962
$$\gamma_{max} \int_0^{\bar{t}} dN = \psi_{max} \int_0^{\bar{t}} \left(\frac{dY}{Y + X_0}\right),$$

963 Making the time dependence of the variables explicit via subscripts

964
$$\gamma_{max} N_t |_0^{\bar{t}} = \psi_{max} \ln(Y_t + X_0)|_0^{\bar{t}},$$

965
$$\gamma_{max}(N_{\bar{t}} - N_0) = \psi_{max}\{\ln(Y_{\bar{t}} + X_0) - \ln(Y_0 + X_0)\}.$$

966 Because $X_0 = X_{\bar{t}}$

967
$$\gamma_{max}(N_{\bar{t}} - N_0) = \psi_{max}\{\ln(Y_{\bar{t}} + X_{\bar{t}}) - \ln(Y_0 + X_{\bar{t}})\},$$

968
$$\gamma_{max}(N_{\bar{t}} - N_0) = \psi_{max} \ln\left(\frac{Y_{\bar{t}} + X_{\bar{t}}}{Y_0 + X_{\bar{t}}}\right).$$

969 Because $X_{\bar{t}} = D_{\bar{t}}/N_{\bar{t}}$ and $Y_{\bar{t}} = T_{\bar{t}}/R_{\bar{t}}$,

970
$$\gamma_{max}(N_{\bar{t}} - N_0) = \psi_{max} \ln\left(\frac{\frac{T_{\bar{t}}}{R_{\bar{t}}} + \frac{D_{\bar{t}}}{N_{\bar{t}}}}{\frac{T_0}{R_0} + \frac{D_{\bar{t}}}{N_{\bar{t}}}}\right).$$

971 Because $T_0 = 0$,

972
$$\gamma_{max}(N_{\bar{t}} - N_0) = \psi_{max} \ln\left(\frac{\frac{T_{\bar{t}}}{R_{\bar{t}}} + \frac{D_{\bar{t}}}{N_{\bar{t}}}}{\frac{D_{\bar{t}}}{N_{\bar{t}}}}\right),$$

973
$$\gamma_{max}(N_{\bar{t}} - N_0) = \psi_{max} \ln\left(\frac{T_{\bar{t}}}{R_{\bar{t}}}\frac{N_{\bar{t}}}{D_{\bar{t}}} + 1\right),$$

974
$$\gamma_{max} = \psi_{max} \ln\left(1 + \frac{T_{\bar{t}}}{R_{\bar{t}}}\frac{N_{\bar{t}}}{D_{\bar{t}}}\right)\frac{1}{N_{\bar{t}} - N_0}.$$

Dropping the "max" subscripts, we have recovered equation [5].

SI section 5 : ASM derivation

In this section, we walk through all of the mathematical steps in the Huisman *et.*

al.[11] derivation. We include this derivation as a quick reference to help the reader compare

the derivations of all four methods (TDR, SIM, ASM, and LDM). We start with a system of

ordinary differential equations described by Huisman *et. al.*[11], the Extended Simonsen

Model (ESM):

$$\frac{dD}{dt} = \psi_D(C)D, \qquad\qquad\qquad [5.1]$$

$$\frac{dR}{dt} = \psi_R(C)R - (\gamma_D(C)D + \gamma_T(C)T)R, \qquad\qquad\qquad [5.2]$$

$$\frac{dT}{dt} = \psi_T(C)T + (\gamma_D(C)D + \gamma_T(C)T)R, \qquad [5.3]$$

$$\frac{dC}{dt} = -(\psi_D(C)D + \psi_R(C)R + \psi_T(C)T)e, \qquad [5.4]$$

983 Where ψ subscripts specify population specific growth rates, γ subscripts specify donor

984 and transconjugant specific conjugation rates, and other variables are consistent with

985 equations [1]-[3] or equations [4.1]-[4.4]. As with the SIM, growth and conjugation rate are

986 both dependent on resource concentration. When resources are depleted, growth and

987 conjugation stop.

988
$$\psi_A(C) = \frac{\psi_{A,max}C}{Q + C},$$

989
$$\gamma_B(C) = \frac{\gamma_{B,max}C}{Q + C},$$

where $A \in \{D, R, T\}$ and $B \in \{D, T\}$.

Under the assumption that growth rate and conjugation rate are constant (where

$\psi_A(C) = \psi_{A,max}$ and $\gamma_B(C) = \gamma_{B,max}$), equation [5.4] can be dropped. Although the

maximal rates of growth and conjugation are assumed, in what follows, we drop the "max"

subscript for notational convenience. This new set of simplified ordinary differential

equations is termed the Approximate Simonsen *et. al.* Method ('ASM').

$$\frac{dD}{dt} = \psi_D D, \qquad [5.5]$$

$$\frac{dR}{dt} = \psi_R R - (\gamma_D D + \gamma_T T)R, \qquad [5.6]$$

$$\frac{dT}{dt} = \psi_T T + (\gamma_D D + \gamma_T T)R, \qquad [5.7]$$

996 Under initial culture conditions, the final simple set of equations [1.1] - [1.3] arise

997 by assuming the recipient population is dominated by growth $\psi_R R \gg (\gamma_D D + \gamma_T T)R$ and

998 the transconjugant population is dominated by growth and conjugation from donors $\psi_T T +$

999 $\gamma_D DR \gg \gamma_T TR$. The solutions to differential equations [1.1] and [1.2] are:

$$D_t = D_0 e^{\psi_D t} \qquad [5.8]$$

$$R_t = R_0 e^{\psi_R t} \qquad [5.9]$$

1000 Here we will derive the solution for the differential equation [1.3] for transconjugant density

1001 T. We know that $D_t = D_0 e^{\psi_D t}$ and $R_t = R_0 e^{\psi_R t}$, therefore

$$\frac{dT}{dt} = \psi_T T + \gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t} \qquad [5.10]$$

1002 We propose that the transconjugant density can be written as a product of time dependent

1003 functions:

$$T_t = u_t v_t \qquad [5.11]$$

1004 We'll drop the t subscripts for notational ease. By the product rule,

$$\frac{dT}{dt} = u\frac{dv}{dt} + v\frac{du}{dt} \qquad [5.12]$$

1005 We can rewrite equation [5.10] by plugging in equations [5.11] and [5.12] as follows:

$$u\frac{dv}{dt} + v\frac{du}{dt} = \psi_T uv + \gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t}$$

$$u\frac{dv}{dt} + v\left(\frac{du}{dt} - \psi_T u\right) = \gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t} \qquad [5.13]$$

1007 We have some freedom to pick u_t as we please. So, let's choose a function such that the

1008 second term of the left-hand side of equation [5.13] is zero:

$$\frac{du}{dt} - \psi_T u = 0$$

$$\frac{du}{dt} = \psi_T u$$

1011 The solution to this differential equation is:

$$u_t = u_0 e^{\psi_T t} \qquad [5.14]$$

1012 where u_0 is a constant.

1013 Thus, we can rewrite equation [5.13] as:

$$u_0 e^{\psi_T t}\frac{dv}{dt} = \gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t}$$

$$u_0 \frac{dv}{dt} = \frac{\gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t}}{e^{\psi_T t}}$$

$$u_0 \frac{dv}{dt} = \gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t} e^{-\psi_T t}$$

$$u_0 \frac{dv}{dt} = \gamma_D D_0 R_0 e^{(\psi_D + \psi_R - \psi_T)t}$$

1018 To solve this equation, we integrate

$$u_0 \int dv = \gamma_D D_0 R_0 \int e^{(\psi_D + \psi_R - \psi_T)t} dt$$

$$u_0 v_t = \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} e^{(\psi_D + \psi_R - \psi_T)t} + c \qquad [5.15]$$

1020 where c is a constant of the integration. To solve for c, plug in $t = 0$,

1021
$$u_0 v_0 = \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} e^{(\psi_D + \psi_R - \psi_T)0} + c$$

1022
$$u_0 v_0 = \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} + c$$

1023
$$c = u_0 v_0 - \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T}$$

1024 Because $T_0 = u_0 v_0$,

1025
$$c = T_0 - \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T}$$

1026 So, we now can find the solution for $v_{\tilde{t}}$ by plugging in our solution for c into equation

1027 [5.15]:

1028
$$u_0 v_{\tilde{t}} = \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} e^{(\psi_D + \psi_R - \psi_T)\tilde{t}} + T_0 - \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T}$$

1029
$$u_0 v_{\tilde{t}} = T_0 + \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} \left\{ e^{(\psi_D + \psi_R - \psi_T)\tilde{t}} - 1 \right\}$$

$$v_{\tilde{t}} = \frac{1}{u_0} \left(T_0 + \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} \left\{ e^{(\psi_D + \psi_R - \psi_T)\tilde{t}} - 1 \right\} \right) \qquad [5.16]$$

1030 Because $T_{\tilde{t}} = u_{\tilde{t}} v_{\tilde{t}}$ through substitution of equations [5.14] and [5.16], we have

1031
$$T_{\tilde{t}} = \left[u_0 e^{\psi_T \tilde{t}} \right] \left[\frac{1}{u_0} \left(T_0 + \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} \left\{ e^{(\psi_D + \psi_R - \psi_T)\tilde{t}} - 1 \right\} \right) \right]$$

1032
$$T_{\tilde{t}} = e^{\psi_T \tilde{t}} \left(T_0 + \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} \left\{ e^{(\psi_D + \psi_R - \psi_T)\tilde{t}} - 1 \right\} \right)$$

1033
$$T_{\tilde{t}} = T_0 e^{\psi_T \tilde{t}} + \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} \left\{ e^{(\psi_D + \psi_R)\tilde{t}} - e^{\psi_T \tilde{t}} \right\}$$

1034 Because $T_0 = 0$, we arrive at the result in Huisman *et al.*[11]

1035
$$T_{\tilde{t}} = \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R - \psi_T} \left\{ e^{(\psi_D + \psi_R)\tilde{t}} - e^{\psi_T \tilde{t}} \right\}$$

1036 Given that $D_{\tilde{t}} = D_0 e^{\psi_D \tilde{t}}$ and $R_{\tilde{t}} = R_0 e^{\psi_R \tilde{t}}$, this can be rewritten as

1037
$$T_{\tilde{t}} = \frac{\gamma_D}{\psi_D + \psi_R - \psi_T} \left\{ D_{\tilde{t}} R_{\tilde{t}} - D_0 R_0 e^{\psi_T \tilde{t}} \right\}$$

1038 Solving for γ_D gives equation [6].

1039

SI section 6 : Alternative laboratory forms for conjugation estimates

To rearrange the SIM estimate, we start with equation [5]. If the entire period from $t = 0$ to $t = \tilde{t}$ involves exponential growth, then $N_{\tilde{t}} = N_0 e^{\psi \tilde{t}}$. In such a case, $\psi = (1/\tilde{t}) \ln(N_{\tilde{t}}/N_0)$. We arrive at the alternative laboratory form for SIM which is equation [2.1].

$$\gamma = \frac{1}{\tilde{t}} \left[\ln \left(1 + \frac{T_{\tilde{t}}}{R_{\tilde{t}}} \frac{N_{\tilde{t}}}{D_{\tilde{t}}} \right) \right] \frac{\ln N_{\tilde{t}} - \ln N_0}{N_{\tilde{t}} - N_0}.$$

To rearrange the ASM estimate, we start with equation [6]. While the equations $\psi_D = (1/\tilde{t}) \ln(D_{\tilde{t}}/D_0)$ and $\psi_R = (1/\tilde{t}) \ln(R_{\tilde{t}}/R_0)$ again provide estimates on donor and recipient growth rates, we cannot express the transconjugant growth rate (ψ_T) as a simple expression of time and initial/final densities of members of the mating culture. However, data from a transconjugant monoculture supply an estimate on this parameter. Thus, we arrive at the laboratory form for ASM which is equation [2.3].

$$\gamma_D = \left\{ \frac{1}{\tilde{t}} (\ln D_{\tilde{t}} R_{\tilde{t}} - \ln D_0 R_0) - \psi_T \right\} \frac{T_{\tilde{t}}}{\left(D_{\tilde{t}} R_{\tilde{t}} - D_0 R_0 e^{\psi_T \tilde{t}} \right)}.$$

To rearrange the LDM estimate, we start with equation [13]. Since $D_{\tilde{t}} = D_0 e^{\psi_D \tilde{t}}$ and $R_{\tilde{t}} = R_0 e^{\psi_R \tilde{t}}$, it is the case that $\psi_D = (1/\tilde{t}) \ln(D_{\tilde{t}}/D_0)$ and $\psi_R = (1/\tilde{t}) \ln(R_{\tilde{t}}/R_0)$. So, we have

$$\gamma_D = \frac{1}{\tilde{t}} \ln p_0(\tilde{t}) \frac{\ln \left(\frac{D_{\tilde{t}}}{D_0} \right) + \ln \left(\frac{R_{\tilde{t}}}{R_0} \right)}{D_0 R_0 - D_{\tilde{t}} R_{\tilde{t}}}$$

After rearrangement, we have

$$\gamma_D = \frac{1}{\tilde{t}} \{ -\ln p_0(\tilde{t}) \} \frac{\ln(D_{\tilde{t}} R_{\tilde{t}}) - \ln(D_0 R_0)}{D_{\tilde{t}} R_{\tilde{t}} - D_0 R_0}.$$

In the laboratory, we measure an estimate ($\hat{p}_0(\tilde{t})$) of the probability that a population has no transconjugants ($p_0(\tilde{t})$) which is simply the fraction of the populations (i.e., parallel cultures) that have no transconjugants at the incubation time $\tilde{t}$ (see SI section 9 for details). In addition, if a 1 ml volume is not used for each mating culture (assuming that all cell densities are measured in $\frac{\text{cfu}}{\text{ml}}$ units), then we must add a correction factor f which is the number of experimental volume units (evu) per ml (see SI section 10 for details and an example). Thus, we arrive at the laboratory form for the LDM which is equation [15].

$$\gamma_D = f \left\{ \frac{1}{\tilde{t}} [-\ln \hat{p}_0(\tilde{t})] \frac{\ln D_{\tilde{t}} R_{\tilde{t}} - \ln D_0 R_0}{D_{\tilde{t}} R_{\tilde{t}} - D_0 R_0} \right\}$$

SI section 7 : Derivation of $p_0(\tilde{t})$ for the LDM estimate

We define $p_n(t)$ to be the probability that there are n transconjugants at time t, where n is a non-negative integer (i.e., $p_n(t) = \Pr\{T_t = n\}$). We focus here on the probability that transconjugants are absent (namely, where $n = 0$) and derive an expression for $p_0(t)$. By definition $p_0(t + \Delta t) = \Pr\{T_{t+\Delta t} = 0\}$. However, $T_{t+\Delta t} = 0$ implies $T_t = 0$, so we can write

$$p_0(t + \Delta t) = \Pr\{(T_{t+\Delta t} = 0) \cap (T_t = 0)\} = \Pr\{T_{t+\Delta t} = 0 \mid T_t = 0\} \Pr\{T_t = 0\},$$

Given that $p_0(t) = \Pr\{T_t = 0\}$, we can use equation [11] to write the following time-increment recursion for $p_0(t)$:

$$p_0(t + \Delta t) = (1 - \gamma_D D_t R_t \Delta t) p_0(t).$$

This can be simplified as follows

$$\frac{p_0(t + \Delta t) - p_0(t)}{\Delta t} = -\gamma_D D_t R_t p_0(t).$$

Taking the limit as $\Delta t \to 0$ gives

$$\lim_{\Delta t \to 0} \frac{p_0(t + \Delta t) - p_0(t)}{\Delta t} = \frac{dp_0(t)}{dt}.$$

Therefore, we have the following differential equation:

$$\frac{dp_0(t)}{dt} = -\gamma_D D_t R_t p_0(t).$$

We are assuming $D_t = D_0 e^{\psi_D t}$ and $R_t = R_0 e^{\psi_R t}$. We note that these assumptions are reasonable if the densities of donors and recipients are reasonably large and the rate of transconjugant generation per recipient cell ($\gamma_D D_t + \gamma_T T_t$, or if $T_t = 0$, simply $\gamma_D D_t$) remains very small relative to per capita recipient growth rate. Under these assumptions, our differential equation becomes:

$$\frac{dp_0(t)}{dt} = -\gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t} p_0(t).$$

We solve this differential equation via separation of variables, integrating from 0 to our incubation time of interest $\tilde{t}$:

$$\int_0^{\tilde{t}} \frac{dp_0(t)}{p_0(t)} = \int_0^{\tilde{t}} -\gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t} dt,$$

$$\ln p_0(t)\big|_0^{\tilde{t}} = \frac{-\gamma_D D_0 R_0}{\psi_D + \psi_R} e^{(\psi_D + \psi_R)t} \bigg|_0^{\tilde{t}},$$

$$\ln p_0(\tilde{t}) - \ln p_0(0) = \frac{-\gamma_D D_0 R_0}{\psi_D + \psi_R} e^{(\psi_D + \psi_R)\tilde{t}} - \frac{-\gamma_D D_0 R_0}{\psi_D + \psi_R}.$$

1095   Given that $p_0(0) = 1$,

1096
$$\ln p_0(\tilde{t}) = \frac{-\gamma_D D_0 R_0}{\psi_D + \psi_R}\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right),$$

1097
$$p_0(\tilde{t}) = \exp\left\{\frac{-\gamma_D D_0 R_0}{\psi_D + \psi_R}\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right)\right\},$$

1098   which is equation [12].

1099

1100   **SI section 8 : Derivation of mutation rate from the Luria-Delbrück experiment**

1101   Here we derive the classic estimate of mutation rate from Luria and Delbrück. We

1102   assume that there is a population of wild-type cells that grow according to the following

1103   equation:

$$N_t = N_0 e^{\psi_N t}, \qquad\qquad [8.1]$$

where N_t is the number of wild type cells at time t and ψ_N is the per capita growth rate.

The wild-type population dynamics are assumed to be deterministic (a reasonable

assumption if the initial population size is reasonably large, i.e., $N_0 \gg 0$). We are also

ignoring the loss of wild-type cells to mutational transformation, but this omission is

reasonable if the mutation rate is very small relative to per capita wild-type growth rate.

Let the number of mutants be given by a random variable M_t. This variable takes

on non-negative integer values. For a very small interval of time, Δt, the current number

of mutants will either increase by one or remain constant. The probabilities of each

possibility are given as follows:

$$\Pr\{M_{t+\Delta t} = M_t + 1\} = \mu N_t \Delta t + \psi_M M_t \Delta t, \qquad\qquad [8.2]$$

$$\Pr\{M_{t+\Delta t} = M_t\} = 1 - (\mu N_t + \psi_M M_t)\Delta t. \qquad\qquad [8.3]$$

The two terms on the right-hand side of equation [8.2] give the ways that a mutant can be

generated. The first term measures the probability that a wild-type cell undergoes a

mutation (μ is the mutation rate). The second term gives the probability that a mutant cell

divides and produces two mutant cells (ψ_M is the mutant growth rate). Equation [8.3] is

the probability that neither of these processes occur.

Analogous to the procedure in SI section 7 (with $p_0(t) = \Pr\{M_t = 0\}$):

$$p_0(t + \Delta t) = (1 - \mu N_t \Delta t)p_0(t).$$

1120   By rearranging, taking the limit as $\Delta t \to 0$, and utilizing equation [8.1], we have

1121
$$\frac{dp_0(t)}{dt} = -\mu N_0 e^{\psi_N t} p_0(t).$$

1122　This differential equation can be solved in an analogous way as well

1123
$$\int_0^{\tilde{t}} \frac{dp_0(t)}{p_0(t)} = \int_0^{\tilde{t}} -\mu N_0 e^{\psi_N t} dt,$$

1124
$$\ln p_0(t)\big|_0^{\tilde{t}} = \frac{-\mu N_0}{\psi_N} e^{\psi_N t} \bigg|_0^{\tilde{t}},$$

1125
$$\ln p_0(\tilde{t}) - \ln p_0(0) = \frac{-\mu N_0}{\psi_N} e^{\psi_N \tilde{t}} - \frac{-\mu N_0}{\psi_N}.$$

1126　Because we assume $M_0 = 0$, we must have $p_0(0) = 1$, and

1127
$$\ln p_0(\tilde{t}) = \frac{-\mu N_0}{\psi_N} \left(e^{\psi_N \tilde{t}} - 1 \right).$$

1128　Solving for the mutation rate μ

1129
$$\mu = -\ln p_0(\tilde{t}) \frac{\psi_N}{N_0 \left(e^{\psi_N \tilde{t}} - 1 \right)},$$

1130　which is equation [14]. This equation can also be expressed as:

1131
$$\mu = -\ln p_0(\tilde{t}) \frac{\psi_N}{N_{\tilde{t}} - N_0}.$$

1132　　　In Luria and Delbrück's original formulation, they measured time in units

1133　proportional to the bacterial division cycles. Specifically, a new time variable z is defined:

1134
$$z = \frac{t}{t_d / \ln 2},$$

1135　where t_d is the period required for population doubling during exponential growth.

1136　Because

1137
$$N_0 e^{\psi_N t} = N_0 2^{t/t_d},$$

1138　it is the case that

1139
$$\psi_N = \frac{1}{t_d / \ln 2}.$$

1140　Therefore, $z = \psi_N t$ and the equation $N_t = N_0 e^{\psi_N t}$ can be expressed as

1141
$$N_z = N_0 e^z$$

1142　Performing the same analysis on this nondimensionalized equation gives the original

1143　formulation (their equations [4] and [5], where our μ is given by their "a" and our $\tilde{z}$ is given

1144　by their "t"):

1145
$$\mu = \frac{-\ln p_0(\tilde{z})}{N_{\tilde{z}} - N_0}.$$

1146

SI section 9 : The LDM MLE derivation

We start by focusing on $p_0(\tilde{t})$, the probability that a population will have no

transconjugants at time $\tilde{t}$ (for notational convenience we'll drop the time variable). What

we actually measure is the number of independent populations (or wells) that have no

transconjugants (call this w) out of the total number of populations (or wells) tracked (call

this W). What is our best estimate of p_0, given our data w? That is, what value p_0

maximizes the likelihood function:

$$\mathcal{L}(p_0) = \Pr\{w|p_0\} = \binom{W}{w}(p_0)^w(1-p_0)^{W-w}$$
1154

1155   Because $-\ln(x)$ is a monotonic decreasing function, the value p_0 that maximizes $\mathcal{L}(p_0)$

1156   will be the same value that minimizes:

$$L(p_0) = -\ln\{\mathcal{L}(p_0)\}$$
1157

1158   This can be rewritten as follows:

$$L(p_0) = -\ln\left\{\binom{W}{w}(p_0)^w(1-p_0)^{W-w}\right\}$$
1159

$$L(p_0) = -\ln\binom{W}{w} - w\ln p_0 - (W-w)\ln(1-p_0)$$
1160

1161   To find the maximum likelihood estimate, we find the critical points of L by setting its

1162   derivative to zero, or:

$$\frac{dL}{dp_0} = -\frac{w}{p_0} + \frac{W-w}{1-p_0} = 0$$
1163

1164   The maximum likelihood estimate for p_0 (which we'll denote $\hat{p}_0$) solves the above

1165   equation, or

$$\frac{W-w}{1-\hat{p}_0} = \frac{w}{\hat{p}_0}$$
1166

$$(W-w)\hat{p}_0 = w(1-\hat{p}_0)$$
1167

$$W\hat{p}_0 - w\hat{p}_0 = w - w\hat{p}_0$$
1168

$$W\hat{p}_0 = w$$
1169

$$\hat{p}_0 = \frac{w}{W}$$
1170

1171   This answer makes intuitive sense: the most likely estimate for p_0 (the probability that a

1172   population has no transconjugants) is simply the fraction of the populations that have no

1173   transconjugants.

To double check that this estimate actually corresponds to a *minimum* of $L(p_0)$, consider the second derivative:

$$\frac{d^2L}{dp_0{}^2} = \frac{w}{p_0{}^2} + \frac{W-w}{(1-p_0)^2}$$

Evaluating this derivative at the critical point yields:

$$\left.\frac{d^2L}{dp_0{}^2}\right|_{p_0=\hat{p}_0} = \frac{w}{\left(\frac{w}{W}\right)^2} + \frac{W-w}{\left(\frac{W-w}{W}\right)^2}$$

$$\left.\frac{d^2L}{dp_0{}^2}\right|_{p_0=\hat{p}_0} = W^2\left(\frac{1}{w} + \frac{1}{W-w}\right)$$

As long as $0 < w < W$,

$$\left.\frac{d^2L}{dp_0{}^2}\right|_{p_0=\hat{p}_0} > 0,$$

which means that L is convex at $\hat{p}_0$ and therefore, this value is a local minimum. In turn, this means that $\hat{p}_0$ is a local maximum for $\mathcal{L}$.

SI section 10 : Experimental volume unit conversion using f

In this section, we walk through the addition of f to the LDM estimate. This is important to maintain the typical units $\frac{\text{ml}}{\text{h}\cdot\text{cfu}}$ used for reporting the conjugation rates. In the original differential equations [1]-[3], the units of the dynamic variables were $\frac{\text{cfu}}{\text{ml}}$. If we want to deal with numbers instead of density, the let us define a new volume unit termed the "evu" standing for "experimental volume unit" where we will assume there are f evu's per ml. Focusing on the number of donors in the experiment (which we label $\breve{D}$), we have the following conversion:

$$\breve{D}\,\frac{\text{cfu}}{\text{evu}} = \frac{\left(D\,\frac{\text{cfu}}{\text{ml}}\right)}{\left(f\,\frac{\text{evu}}{\text{ml}}\right)},$$

Equivalently,

$$\breve{R} = \frac{R}{f}$$

$$\breve{T} = \frac{T}{f}$$

In our original differential equations, let us multiply both sides of all the differential equations by $1/f$, yielding:

1199
$$\frac{1}{f}\frac{dD}{dt} = \psi \frac{1}{f}D,$$

1200
$$\frac{1}{f}\frac{dR}{dt} = \psi \frac{1}{f}R - \gamma \frac{1}{f}R(D+T),$$

1201
$$\frac{1}{f}\frac{dT}{dt} = \psi \frac{1}{f}T + \gamma \frac{1}{f}R(D+T).$$

1202 This can be reworked as

1203
$$\frac{d\breve{D}}{dt} = \psi\breve{D},$$

1204
$$\frac{d\breve{R}}{dt} = \psi\breve{R} - \gamma\breve{R}(D+T),$$

1205
$$\frac{d\breve{T}}{dt} = \psi\breve{T} + \gamma\breve{R}(D+T).$$

1206 We have a mix of densities $\left(\frac{\text{cfu}}{\text{ml}}\right)$ and numbers $\left(\frac{\text{cfu}}{\text{evu}}\right)$ here, so we do the following:

1207
$$\frac{d\breve{D}}{dt} = \psi\breve{D},$$

1208
$$\frac{d\breve{R}}{dt} = \psi\breve{R} - f\gamma\breve{R}\left(\frac{D+T}{f}\right),$$

1209
$$\frac{d\breve{T}}{dt} = \psi\breve{T} + f\gamma\breve{R}\left(\frac{D+T}{f}\right).$$

1210 Again, making substitutions:

1211
$$\frac{d\breve{D}}{dt} = \psi\breve{D},$$

1212
$$\frac{d\breve{R}}{dt} = \psi\breve{R} - \{f\gamma\}\breve{R}(\breve{D}+\breve{T}),$$

1213
$$\frac{d\breve{T}}{dt} = \psi\breve{T} + \{f\gamma\}\breve{R}(\breve{D}+\breve{T}).$$

1214 If we let

1215
$$\breve{\gamma} = f\gamma$$

1216 we note the units on $\breve{\gamma}$ are $\frac{\text{evu}}{\text{h}\cdot\text{cfu}}$. Incorporating this relation, we have

1217
$$\frac{d\breve{D}}{dt} = \psi\breve{D},$$

1218
$$\frac{d\breve{R}}{dt} = \psi\breve{R} - \breve{\gamma}\breve{R}(\breve{D}+\breve{T}),$$

1219
$$\frac{d\breve{T}}{dt} = \psi\breve{T} + \breve{\gamma}\breve{R}(\breve{D}+\breve{T}).$$

1220 This set of equations tracks the number of cells (per evu). Thus, if the above equations

1221 were used, then the derivations of the LDM estimate could flow exactly like we show in

1222 the "Theoretical Framework" section and SI section 7. That is, the following will be correct:

1223
$$\check{\gamma} = -\ln p_0(\tilde{t}) \left(\frac{\psi_D + \psi_R}{\breve{D}_0 \breve{R}_0 \left(e^{(\psi_D + \psi_R)\tilde{t}} - 1 \right)} \right)$$

1224 Note, no correction is needed on $p_0(\tilde{t})$ as everything is in terms of numbers, which was

1225 how this quantity was derived. Because $\breve{D} = \frac{D}{f}$ and $\breve{R} = \frac{R}{f}$, we can rewrite the above as

1226
$$\check{\gamma} = -\ln p_0(\tilde{t}) \left(\frac{\psi_D + \psi_R}{\dfrac{D_0}{f} \dfrac{R_0}{f} \left(e^{(\psi_D + \psi_R)\tilde{t}} - 1 \right)} \right)$$

1227 Or:

1228
$$\frac{\check{\gamma}}{f} = f \left\{ -\ln p_0(\tilde{t}) \left(\frac{\psi_D + \psi_R}{D_0 R_0 \left(e^{(\psi_D + \psi_R)\tilde{t}} - 1 \right)} \right) \right\}$$

1229 Because $\gamma = \frac{\check{\gamma}}{f}$, we have

1230
$$\gamma = f \left\{ -\ln p_0(\tilde{t}) \left(\frac{\psi_D + \psi_R}{D_0 R_0 \left(e^{(\psi_D + \psi_R)\tilde{t}} - 1 \right)} \right) \right\}$$

1231 Note that if our evu was 1 ml, then $f = 1$ and we could use our estimate exactly as written.

1232 Generally, we have to correct our original metric by multiplying by f.

1233   Now just as a check, let's see what happens when we flip the unit of volume for a

1234 different metric, say SIM. Suppose that we use the $\frac{\text{cfu}}{\text{evu}}$ units for cell densities, which

1235 means we are calculating

1236
$$\check{\gamma} = \psi \ln \left(1 + \frac{\check{T}_{\tilde{t}}}{\check{R}_{\tilde{t}}} \frac{\check{N}_{\tilde{t}}}{\check{D}_{\tilde{t}}} \right) \frac{1}{(\check{N}_{\tilde{t}} - \check{N}_0)}$$

1237 where $\check{\gamma}$ has units $\frac{\text{evu}}{\text{h} \cdot \text{cfu}}$. Using our conversion, we can rewrite the right-hand side as

1238 follows:

1239
$$\check{\gamma} = \psi \ln \left(1 + \frac{T_{\tilde{t}}/f}{R_{\tilde{t}}/f} \frac{N_{\tilde{t}}/f}{D_{\tilde{t}}/f} \right) \frac{1}{(N_{\tilde{t}} - N_0)/f}$$

1240 Canceling and rearranging yields

1241

1242
$$\frac{\check{\gamma}}{f} = \psi \ln \left(1 + \frac{T_{\tilde{t}}}{R_{\tilde{t}}} \frac{N_{\tilde{t}}}{D_{\tilde{t}}} \right) \frac{1}{(N_{\tilde{t}} - N_0)}$$

1243 However, $\gamma = \frac{\check{\gamma}}{f}$, so we have

1244

$$\gamma = \psi \ln \left(1 + \frac{T_{\tilde{t}}}{R_{\tilde{t}}} \frac{N_{\tilde{t}}}{D_{\tilde{t}}} \right) \frac{1}{(N_{\tilde{t}} - N_0)}$$

1245 Therefore, if we flip to cell densities in the new unit $\frac{\text{cfu}}{\text{ml}}$, we will automatically get the

1246 conjugation rate in the new unit $\frac{\text{ml}}{\text{h} \cdot \text{cfu}}$. The same applies to the TDR and ASM estimates.

1247 Therefore, there are no correction factors needed for these metrics as volume units

1248 change. Note, in these cases, the numerical value of the transfer rate in the new units is

1249 *not* generally the same as the rate in the old units ($\gamma \neq \check{\gamma}$ when $f \neq 1$).

1250

SI section 11 : Stochastic simulation framework

We developed a stochastic simulation framework using the Gillespie algorithm. We

extended the base model (equations [1] - [3]) to include plasmid loss due to segregation

at rate τ. Thus, transconjugants are transformed into plasmid-free recipients due to

plasmid loss via segregation at rate τ_T. The donors are transformed into plasmid-free

cells due to plasmid loss via segregation at rate τ_D. Therefore, the extended model

(equations [11.1] - [11.4]) tracks the change in density of a new population type, plasmid-

free *former* donors (F). In total, the extended model describes the change in density of

four populations (D, R, T, and F) due to various biological parameters: growth rates (ψ_D,

ψ_R, ψ_T, and ψ_F), conjugation rates (γ_D and γ_T) and segregation rates (τ_D and τ_T). For

simplicity, here we assume donor-to-recipient conjugation rate is equal to donor-to-

plasmid-free-former-donor conjugation rate and represent both with γ_D. In addition, we

assume transconjugant-to-recipient conjugation rate is equal to transconjugant-to-

plasmid-free-donor conjugation rate and represent both with γ_T.

$$\frac{dD}{dt} = \psi_D D + (\gamma_D D + \gamma_T T)F - \tau_D D, \qquad [11.1]$$

$$\frac{dR}{dt} = \psi_R R - (\gamma_D D + \gamma_T T)R + \tau_T T, \qquad [11.2]$$

$$\frac{dT}{dt} = \psi_T T + (\gamma_D D + \gamma_T T)R - \tau_T T. \qquad [11.3]$$

$$\frac{dF}{dt} = \psi_F F - (\gamma_D D + \gamma_T T)F + \tau_D D. \qquad [11.4]$$

1265 Each simulation examined a biological process (i.e., growth, conjugation, or

1266 segregation) in isolation by manipulating one or two of the relevant parameters in order

1267 to determine the effect on accuracy and precision of various estimates. A simulation was

initiated with a positive density of donors (D_0), a positive density of recipients (R_0), biological parameters for each population type, and then run for a specified incubation time $\tilde{t}$. For each initial parameter setting, we simulated 10,000 parallel populations and calculated the conjugation rate using each method (TDR, SIM, ASM, and LDM). Each estimate has different requirements for calculating the conjugation rate. The LDM estimate needs multiple populations to calculate $\hat{p}_0(\tilde{t})$. Thus, for a single estimate via LDM, we reserved 100 parallel populations, where 99 were used to calculate $\hat{p}_0(\tilde{t})$ and the remaining population was used to calculate initial and final cell densities. In other words, the 10,000 parallel populations yielded 100 replicate LDM estimates for each parameter setting. For the other methods, we calculated a conjugation rate for each simulated population. In other words, the 10,000 simulated populations yielded 10,000 replicate estimates for each initial parameter setting.

The window of time for population growth (i.e., incubation time $\tilde{t}$) is different for the LDM vs. the other methods. A non-zero conjugation rate can be calculated for TDR, SIM, and ASM whenever there is a non-zero number of transconjugants. This contrasts with the LDM since conjugation rate is calculated when a proportion of populations have zero transconjugants. Given different incubation time requirements for each method, we use a deterministic simulation for each parameter setting to find the incubation time ($\tilde{t}$) where we calculated conjugation rate. For the LDM, the incubation time was chosen to occur where *on average* a single transconjugant was present in the population. For the other methods, the incubation time occurred where *on average* ten transconjugant cells were present in the population. For Figure 2, we used a "baseline" set of parameters ($\psi_D = \psi_R = \psi_T = \psi_F = 1, \gamma_D = \gamma_T = 1 \times 10^{-6}, \tau_D = \tau_T = 0$) and initial densities ($D_0 = R_0 = 1 \times 10^2$) unless otherwise indicated. Given that the Gillespie algorithm is computationally expensive, we chose low initial densities and an unusually high conjugation rate. See below and Figure 3 for more realistic rates.

For Figure 3, the conjugation rate was estimated for each method at 30-minute intervals. The 30-minute time intervals occur over an earlier time interval for the LDM vs. the other methods for reasons described above. For the 30-minute time intervals, we applied estimate-specific filters. For the TDR, SIM, and ASM estimates, the range for these intervals was determined by the condition that at least 90 percent of the simulated populations contained transconjugants at the incubation time $\tilde{t}$. For the LDM estimate,

 the range for these intervals was determined by the condition that at least one parallel

 population had zero transconjugants. Therefore, estimates for the LDM are shown at

 earlier incubation times than those for the other methods (Figure 3). Given that Figure 3

 explores only three parameter combinations (i.e., one for each panel), the computational

 expense of the Gillespie algorithm was of less concern than in Figure 2 where there are

 many more parameter settings per figure panel. Thus, for Figure 3, we set a conjugation

 rate that is often reported in the literature ($\gamma_D = \gamma_T = 1 \times 10^{-14}$) and more reasonable

 initial densities ($D_0 = R_0 = 1 \times 10^5$) unless otherwise indicated. We note that the

 Gillespie algorithm is computationally expensive when the densities are very large.

 Therefore, due to the longer incubation times needed for TDR, SIM, and ASM, only 100

 populations of the 10,000 were incubated through the later time intervals (i.e., in Figure

 3, each time point has 100 conjugation rate estimates).

SI Figure 1: The effect of different host growth rates on conjugation rate estimation. The parameter values were $\psi_D = \psi_R = \psi_T = 1$, $\gamma_D = \gamma_T = 1 \times 10^{-6}$, and $\tau_D = \tau_T = 0$ unless otherwise indicated. The dynamic variables were initialized with $D_0 = R_0 = 1 \times 10^2$ and $T_0 = F_0 = 0$. All incubation times are short but are specific to each parameter setting. Conjugation rate was estimated using 10,000 populations and summarized using boxplots. The gray dashed line indicates the "true" value for the donor conjugation rate. The box contains the 25th to 75th percentile. The vertical lines connected to the box contain 1.5 times the interquartile range above the 75th percentile and below the 25th percentile with the caveat that the whiskers were always constrained to the range of the data. The colored line in the box indicates the median. The solid black line indicates the mean. The boxplots in gray indicate the baseline parameter setting, and all colored plots represent deviation of one or two parameters from baseline. In this plot, we explored unequal host growth rates by tracking simulated populations over a range $\psi_R = \psi_T \in \{0.0625, 0.125, 0.24, 0.5, 1, 2, 4, 8\}$. This captures the situation in which the recipient (and therefore transconjugant) is a different strain or species from the donor and differs in growth rate.

SI section 12 : The LDM protocol

 Here we provide a general protocol for implementing the LDM in the laboratory.

 Some aspects need to be tailored to the specific experimental system, such as selection

 markers and incubation times. At the end of the protocol, we summarize some protocol

additions that may be useful if the experimental system allows. For the main body of this section, we left out these "additions" and only included the minimum supplies, strains, and selection conditions needed to estimate conjugation rate using the LDM. For a more general overview of the LDM see the "The LDM Implementation" section.

To start, identify the donor and recipient strains to be used (SI Table 5). The recipients must contain a selectable marker unique to the recipient host strain (S_{RH}). The donors must also contain a unique selectable marker, but it will be encoded on the donor's conjugative plasmid (S_P). The combination of the unique selectable markers allows for selection of the transconjugant cell type, a recipient strain containing the conjugative plasmid.

SI Table 5: Overview of cell types in the conjugation assay.

Cell type	Cell type description	Selectable marker (S)
R	Plasmid-free recipients	Recipient host marker (S_{RH})
D	Plasmid donors	Donor plasmid marker (S_P)
T	Transconjugants	Recipient host marker (S_{RH}) Donor plasmid marker (S_P)

Part I. <u>Characterize cell types (estimated time = 4 days)</u>

Four key pieces of information necessary to proceed with Part II are determined here. (1) The approximate cell density for donors and recipients after overnight growth. (2) The minimum inhibitory concentrations (MIC) for the donors, recipients, and transconjugants. If the selectable markers involve antibiotic resistance, this will identify the antibiotic concentration (selection for $S_{RH} + S_P$) to select for transconjugants and inhibit donors and recipients. (3) Verifying the conjugative plasmid will transfer horizontally between cells in liquid culture. Note that some plasmids transfer inefficiently in liquid culture. (4) The length of lag phase for donor and recipient cells. This information is needed because the donors and recipients are mixed in the exponential phase when determining "the conjugation window" (Part II) and executing the LDM conjugation assay (Part III).

Day I.1

Step I.1.1 Start three replicate overnight cultures from freezer stocks of the donors and recipients (SI Figure 2a). Include selection for S_P in the donor cultures to ensure plasmid maintenance.

Step I.1.2 Incubate cultures.

Day I.2

Step I.2.1 Transfer the donor and recipient cultures created in Step I.1.1 using a 1:100 dilution (SI Figure 2b). Include selection for S_P in the donor cultures to ensure plasmid maintenance.

Step I.2.2 To start three replicate mating cultures, dilute the donor and recipient cultures created in Step I.1.1 using a 1:100 dilution. Mix the diluted donors and recipients at a 1:1 ratio (SI Figure 2b).

Note: These mating cultures allow for plasmid conjugation between donors and recipients to create transconjugants. This will (1) verify that transconjugants can form in the liquid condition, (2) generate the transconjugant strain that will verify the transconjugant-selecting conditions during the MIC assay.

Note: If antibiotic selection is used to maintain the donor plasmid, wash donor cells (through three repeated rounds of centrifugation, removal of the supernatant, addition of an equal volume of saline, and vortexing) before this step.

Step I.2.3 Incubate cultures.

Day I.3

Step I.3.1 Determine the overnight culture density for the donor and recipient cultures created in Step I.2.1, for example by creating a dilution series and separately plating each culture on non-selective medium.

Note: Gathering density information for donors and recipients can help the experimenter tailor Part II most effectively (see Figure 4a)

Step I.3.2 Determine the minimum inhibitory concentration (MIC) for all cell types (donors, recipients and transconjugants) using transconjugant selection (selection for $S_{RH} + S_P$).

Note: The MIC needs to be in the growth format planned for the remaining phases (container, temperature, growth medium type, etc.). Since drug

interaction effects can occur, it is not appropriate to assume that the MIC values established for two strains with a single selectable marker can be combined to yield conditions that will inhibit a strain with both selectable markers (i.e., the MIC for the donor and the recipient are not necessarily concentrations that will inhibit the transconjugant).

I.3.2.1 Create the MIC microtiter plate with a twofold gradient of transconjugant-selecting medium across the rows. See SI Figure 2c, for example.

I.3.2.2 Dilute the donor, recipient and mating cultures created on Day I.2 using a 1:100 dilution. Inoculate the MIC plate with the diluted donors, recipients, and mating cultures in the appropriate columns.

Note: If transconjugants are at too of low of a percentage in the mating cultures, a transconjugant may not get into the MIC microtiter plate. Thus, the MIC for the "transconjugant" may appear to be equal to the maximum MIC of the donors and recipients. In this case, a mating culture should be plated on transconjugant-selecting agar plates to isolate a transconjugant and then determine the MIC.

I.3.2.3 Incubate the MIC microtiter plate.

Step I.3.3 Characterize strain's growth using a spectrophotometer.

I.3.3.1 Dilute the donor and recipient cultures created on Day I.2. using a 1:100 dilution and inoculate three replicate wells for both the diluted donors and recipients.

I.3.3.2 Incubate the microtiter plate in a spectrophotometer using the discontinuous kinetic reading setting which will measure the optical density (O.D.) of each growing culture at frequent intervals.

Note: The spectrophotometer's setting for the kinetic read should include shaking and temperature control which matches the growth conditions used for culturing the strains.

Day I.4

Step I.4.1 Calculate the overnight culture density by counting the cfu on the agar plates from Step I.3.1.

Step I.4.2 Determine the transconjugant selection MIC for each cell type by analyzing the MIC microtiter plate created in Step I.3.2.

Note: The transconjugant-selecting concentration to be chosen for Part II should be below the MIC for transconjugants and above the MIC for donors and recipients. All the tested concentrations in SI Figure 2d are usable for Part II.

Step I.4.3 Analyze the growth kinetic readings from Step I.3.3 to determine the length of lag phase for the donor and recipient strains.

Note: In Part II and Part III, the donors and recipient strains need to grow through lag phase and be mixed while in exponential phase. The length of lag phase may differ between donors and recipients.

Part II. Determine the conjugation window (estimated time = 4 days)

Day II.1

Step II.1.1 Start overnight cultures by repeating Day I.1 and adding in three replicate wells for the transconjugant strain isolated in Part I.

Day II.2

Step II.2.1 Transfer cultures by repeating Step I.2.1 and Step I.2.3 from Day I.2 including the transconjugant cultures.

Day II.3

Step II.3.1 Transfer the donor and recipient cultures created in Step II.2.1 using a 1:100 dilution. Add selection for S_P to the donor cultures to ensure plasmid maintenance.

Step II.3.2 Incubate the donor and recipient cultures created in Step II.3.1 until both are in exponential phase. The incubation time for each strain was determined in Step I.4.3.

Step II.3.3 Make a 10-fold dilution series of the donor and recipient cultures after the incubation in Step II.3.2 and plate appropriate dilutions to determine densities. Also, dilute the transconjugant cultures created in Step II.2.1.

Note: Reserve a portion of these dilutions to create control wells in a future step.

1433   Step II.3.4 Mix a portion of the corresponding donor and recipient dilutions created in
1434   Step II.3.3.
1435   *Note: If antibiotic selection is used to maintain the donor plasmid, wash*
1436   *donor cells as described previously.*
1437   Step II.3.5 Add the diluted mixed cultures created in Step II.3.4 into the corresponding
1438   columns in a deep-well microtiter plate (Figure 4a gray columns). Create
1439   multiple replicate wells for each density-time treatment (see Figure 4a and
1440   4b for an example microtiter plate map).
1441   *Note: The culture will be diluted 1:10 with transconjugant-selecting medium*
1442   *in a future step. Thus, add a culture volume which allows for a 1:10 dilution*
1443   *in the deep-well. This will vary depending on the volume capacity of the*
1444   *deep-well plate used.*
1445   Step II.3.6 Add the diluted donor, recipient, and transconjugant cultures created in
1446   Step II.3.3 into the separate corresponding columns (Figure 4a red, blue,
1447   and purple column).
1448   *Note: Multiple dilutions were created for the donors, recipients, and*
1449   *transconjugants in Step II.3.3. All of the dilutions can be used for controls*
1450   *by using an additional deep-well plate. Figure 4a shows a simple scenario*
1451   *where the densest dilution is used for the control column.*
1452   Step II.3.7 Add the diluted cultures for the donors and recipients created in Step II.3.3
1453   into additional separate columns (not shown in Figure 4) to be used as
1454   mating controls at each time point.
1455   *Note: The mating controls are created at each time point by mixing donors*
1456   *and recipients and immediately adding transconjugant-selecting media. The*
1457   *mating controls confirm that plasmid conjugation is not occurring after the*
1458   *transconjugant-selecting medium is added.*
1459   Step II.3.8 At each time point,
1460   II.3.8.1   Mix the donor and recipient cultures created for the mating controls in
1461   Step II.3.7 into an empty well in the corresponding row (i.e., the mating
1462   control column).
1463   II.3.8.2   Add transconjugant-selecting medium to all wells in the designated
1464   row(s) which dilutes the cultures ten-fold (see Figure 4b).

Step II.3.9 Incubate the microtiter plate until turbidity is stable.

Note: Since transconjugants are rare, it will take longer for cultures to reach

stationary phase. Thus, even for fast-growing bacteria this incubation may

be longer than one day.

Note: The donor and recipient control wells should stay clear throughout

incubation. If turbidity starts to occur, antibiotic may be degrading.

Day II.4

Step II.4.1 Calculate the culture density from Step II.3.1 by counting the cfu on the agar

plates created in Step II.3.3.

Step II.4.2 Record the turbidity results from the deep-well plate(s) (Figure 4c).

Step II.4.3 Analyze the data from Step II.4.1 and Step II.4.2 then select a target

incubation time $(\tilde{t}')$ and target initial density for the donors (D_0') and

recipients (R_0') to use in Part III (see Figure 4c for an example).

1478

Part III. <u>Executing the actual LDM conjugation assay (estimated time = 4 days)</u>

Day III.1

Step III.1.1 Start overnight cultures by repeating Step II.1.1.

Day III.2

Step III.2.1 Transfer cultures by repeating Step II.2.1.

Day III.3

Step III.3.1 Create donor and recipient cultures in exponential phase by repeating Step

II.3.1 and Step II.3.2.

Step III.3.2 Dilute the donor and recipient cultures from Step III.3.1 to the target initial

densities (D_0', R_0'). Dilute the transconjugant cultures created in Step III.2.1.

Note: Reserve a portion of the dilutions to create control wells.

Step III.3.3 Mix a portion of the diluted donor and recipient cultures created in Step

III.3.2.

Step III.3.4 Add the diluted mixed culture created in Step III.3.3 into the corresponding

wells in a deep-well microtiter plate (Figure 4d gray wells).

Note: Add additional culture volume to the mating wells to be used to

determine the initial and final densities. Thus, after the additional volume is

removed for the initial density plating (D_0, R_0) the remaining culture volume will match the other mating wells during the incubation time.

Step III.3.5 Add the diluted donor, recipient, and transconjugant cultures created in Step III.3.2 into the corresponding separate wells (Figure 4d red, blue, and purple wells).

Step III.3.6 Add the diluted donor and recipient cultures created in Step III.3.2 into additional wells (not shown in Figure 4) to be used as mating controls.

Note: The mating controls are created after the incubation time $(\tilde{t})$ by mixing donor and recipient cultures and immediately adding transconjugant-selecting medium. The mating controls confirm that the donors and recipients cannot grow in the transconjugant-selecting medium and that plasmid conjugation is not occurring after transconjugant-selecting medium is added. One deep-well plate can be used for Part III by rearranging the microtiter plate map shown in Figure 4d to include the extra wells needed for the mating controls).

Step III.3.7 Determine initial densities of donor and recipients (D_0, R_0) by dilution-plating from three designated mating wells.

Step III.3.8 Incubate the deep well plate for the chosen time $(\tilde{t})$.

Step III.3.9 Determine final densities of donor and recipients $(D_{\tilde{t}}, R_{\tilde{t}})$ by dilution-plating from the three designated mating wells used in Step III.3.7.

Step III.3.10 Add transconjugant-selecting medium to all wells using a 1:10 dilution (see Figure 4d).

Step III.3.11 Incubate the microtiter plate.

Note: Since transconjugants are rare, it will take longer for cultures to reach stationary phase. Thus, even for fast-growing bacteria this incubation may be longer than one day.

Day III.4

Step III.4.1 Calculate the initial (D_0, R_0) and final densities $(D_{\tilde{t}}, R_{\tilde{t}})$ by counting the cfu on the agar plates from Step III.3.7 and Step III.3.9.

Step III.4.2 Record the turbidity results from the deep-well plate (Figure 4d).

Step III.4.3 Analyze the data from Step III.4.1 and Step III.4.2 to calculate the donor conjugation rate using the LDM estimate (equation [15]).

Protocol "additions" to consider:

- Add an additional selectable marker to either the donor host strain (S_{DH}) or the recipient host strain (S_{RH2}). Since the selectable marker in the recipient host strain (S_{RH}) is often chromosomally encoded through a single mutation, the donor may easily become resistant to the selecting agent (e.g., corresponding antibiotic) through a similar or different mutation in the donor chromosome. Thus, it becomes valuable to be able to distinguish transconjugant-containing populations from mutant-donor-containing populations through an additional marker. This type of mutant arose in our experiments. However, our donor strain had an additional selectable marker (S_{DH}), thus the final microtiter plate in the LDM assay was pin replicated into medium selecting for such donors (selection for $S_{DH} + S_P + S_{RH}$). All mutant-donor-containing populations were identified and eliminated from the LDM calculation. If an additional selectable marker was available for the recipient host strain (S_{RH2}), the transconjugant-selecting medium could have all three selective agents (selection for $S_P + S_{RH} + S_{RH2}$); here, a spurious result would require two mutations in the same donor lineage, which occurs with a very low probability.

- Adding additional densities and time intervals. In Figure 4, we have shown a version of Part II. This portion of the assay is highly customizable depending on the strains that are used. If the experimenter has an idea of the conjugation window, then Part II could be simplified. If the experimenter is working with strains that have large unknowns, then adding additional densities or time intervals is advisable.

- For a more detailed protocol, see https://github.com/livkosterlitz/LDMprotocol/wiki.

SI Figure 2: Protocol overview for determining the minimum inhibitory concentration in Part I. (a) On day 1, the donors (red) and recipients (blue) are inoculated into three replicate wells in the standard 96-well microtiter plate. (b) On day 2 after incubation of the microtiter plate in a, the grown donor (red crosses) and recipient (blue crosses) replicate cultures are transferred (and diluted) into new wells. In addition, the grown donor a nd recipient are mixed to inoculate

three replicate mating wells (gray) to allow for plasmid transfer between donors and recipients to create transconjugants. (c) On day 3 after incubation, the drug microtiter plate is created using transconjugant-selecting medium (selection for $S_{RH} + S_P$) over a two-fold gradient. The bottom row of the microtiter plate contains non-selective medium thus serves as a positive control for bacterial growth. The corresponding grown culture from part (b) is diluted and inoculated into the designated columns in the drug microtiter plate. (d) After incubation of the microtiter plate in (c), MIC is determined for the donors, recipients, and for transconjugants (purple) that arose in the mating cultures. For recipients (blue cross) and donors (red cross), the MIC for transconjugant selecting medium is below the 1X drug concentration. For transconjugants (purple cross), the MIC for transconjugant-selecting medium is above the 64X drug concentration. In the bottom row with non-selective medium, all cell types can be present (mixed cross with blue, red and purple).

SI section 13 : Probability generating function and low-order moments

Keller and Antal (2015)[38] studied a generalization of the process explored by Luria and Delbrück (1943)[20]. To start, Keller and Antal consider a wildtype population expanding from a single cell as follows:

$$N_t = f(t) = e^{\delta t}.$$

Each wildtype cell generates a mutant cell at a rate v', which grows as a stochastic birth process with rate α (Keller and Antal studied a supercritical birth-death process, but we will focus on the special case of a pure birth process). In this case, mutants form at a rate $v'f(t)$, such that the times of mutant arrival conform to a non-homogeneous Poisson process. We note that if we start with N_0 cells, then mutants form at a rate $N_0 v' f(t)$. Alternatively, we can set $v = N_0 v'$, such that mutants form at a rate $v f(t)$, which is the case explored by Keller and Antal.

Keller and Antal derive the probability generating function for the total number of mutants at an arbitrary time:

$$G(z,t) = \exp\left\{\frac{v}{\delta}\left(F\left(1,\kappa;1+\kappa;\frac{z}{z-1}e^{-\alpha t}\right) - e^{\delta t}F\left(1,\kappa;1+\kappa;\frac{z}{z-1}\right)\right)\right\},$$

where F is the Gaussian hypergeometric function and $\kappa = \frac{\delta}{\alpha}$.

Our process of interest (the formation and growth of transconjugants) can be seen as an instance of their formulation by making the following substitutions:

$$\delta = \psi_D + \psi_R,$$

$$v = \gamma_D D_0 R_0,$$

$$\alpha = \psi_T.$$

With these substitutions, the generating function becomes:

1579
$$G(z,t) = \exp\left\{ \frac{\gamma_D D_0 R_0}{\psi_D + \psi_R} \left(F\left(1, \frac{\psi_D + \psi_R}{\psi_T}; 1 + \frac{\psi_D + \psi_R}{\psi_T}; \frac{z}{z-1} e^{-\psi_T t}\right) \right. \right.$$

1580
$$\left. \left. - e^{(\psi_D + \psi_R)t} F\left(1, \frac{\psi_D + \psi_R}{\psi_T}; 1 + \frac{\psi_D + \psi_R}{\psi_T}; \frac{z}{z-1}\right) \right) \right\}.$$

1581   Because

1582
$$G(z,t) = \sum_{n=0}^{\infty} p_n(t) z^n,$$

1583   the probability of zero transconjugants now becomes straightforward (given

1584 $F\left(1, \frac{\psi_D + \psi_R}{\psi_T}; 1 + \frac{\psi_D + \psi_R}{\psi_T}; 0\right) = 1$):

1585
$$p_0(t) = G(0,t) = \exp\left\{ \frac{-\gamma_D D_0 R_0}{\psi_D + \psi_R} \left(e^{(\psi_D + \psi_R)t} - 1 \right) \right\},$$

1586   which agrees with the result from SI section 7.

1587

1588   Making the appropriate substitutions, we can also write the mean and variance (eqs. 8

1589   and 9 from Keller and Antal) for the transconjugants:

1590

1591
$$E[T_t] = \begin{cases} \gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t} t & \text{if } \psi_D + \psi_R = \psi_T \\ \dfrac{\gamma_D D_0 R_0 \left(e^{(\psi_D + \psi_R)t} - e^{\psi_T t} \right)}{\psi_D + \psi_R - \psi_T} & \text{if } \psi_D + \psi_R \neq \psi_T \end{cases}$$

1592
$$\text{Var}[T_t] = \begin{cases} \dfrac{2\gamma_D D_0 R_0 \left(e^{2(\psi_D + \psi_R)t} - e^{(\psi_D + \psi_R)t} \right)}{\psi_D + \psi_R} - \gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t} t & \text{if } \psi_D + \psi_R = \psi_T \\ \dfrac{2\gamma_D D_0 R_0 \left(e^{(\psi_D + \psi_R)t/2} - e^{(\psi_D + \psi_R)t} \right)}{\psi_D + \psi_R} + 2\gamma_D D_0 R_0 e^{(\psi_D + \psi_R)t} t & \text{if } \psi_D + \psi_R = 2\psi_T \\ \gamma_D D_0 R_0 \left\{ \dfrac{2e^{2\psi_T t}(\psi_T - (\psi_D + \psi_R)) - e^{\psi_T t}(2\psi_T - (\psi_D + \psi_R)) + (\psi_D + \psi_R)e^{(\psi_D + \psi_R)t}}{(2\psi_T - (\psi_D + \psi_R))(\psi_T - (\psi_D + \psi_R))} \right\} & \text{otherwise} \end{cases}$$

1593   We provide derivations for these expressions in Box 1. In all cases, the variance grows

1594   relative to the mean over time (see Box 2 below for the derivations).

1595

Box 1 : Derivations of the first and second central moments

In this Box, we derive the equations for the moments using terminology close to what

Keller and Antal used (focusing on the case where $\kappa \notin \{1,2\}$). From above, $N = e^{\delta t}$ and

$N^{-1/\kappa} = e^{\alpha t}$. Keller and Antal denote the number of mutants (analogous to our

transconjugants) as B. Also, letting $\xi = \frac{z}{z-1}$ and $\mu = \frac{\nu}{\alpha}$ and dropping the time argument for

notational convenience, the probability generating function (PGF) can be written as:

1602

$$G_B(z) = \exp\left\{\frac{N\mu}{\kappa}\left[\frac{1}{N}F\left(1,\kappa;1+\kappa;\xi N^{-\frac{1}{\kappa}}\right) - F(1,\kappa;1+\kappa;\xi)\right]\right\}.$$

The expected value for the number of mutants can be obtained from the PGF in the usual way:

$$E[B] = G_B'(1-) = \lim_{z\to 1-} G_B(z)\frac{N\mu}{1+\kappa}\frac{1}{(z-1)^2}\Sigma,$$

where

$$\Sigma = -N^{-1-\frac{1}{\kappa}}F\left(2,1+\kappa;2+\kappa;\xi N^{-\frac{1}{\kappa}}\right) + F(2,1+\kappa;2+\kappa;\xi).$$

We use the following identity

$$F(a,b;c;z) = (1-z)^{c-a-b}F(c-a,c-b;c;z),$$

to obtain

$$F(2,1+\kappa;2+\kappa;\xi) = (1-\xi)^{-1}F(\kappa,1,2+\kappa,\xi)$$

$$= (1-z)F\left(\kappa,1,2+\kappa,\frac{z}{z-1}\right)$$

Using a related identity

$$F(a,b;c;z) = (1-z)^{-b}F\left(c-a,b;c;\frac{z}{z-1}\right),$$

we get:

$$F(2,1+\kappa;2+\kappa;\xi) = (1-z)F\left(\kappa,1,2+\kappa,\frac{z}{z-1}\right)$$

$$= (1-z)^2 F(2,1,2+\kappa,z)$$

We will use a similar procedure for the first term in Σ, obtaining:

$$F(2,1+\kappa;2+\kappa;\xi N^{-1/\kappa}) = (1-\xi N^{-1/\kappa})^{-1}F(\kappa,1,2+\kappa;\xi N^{-1/\kappa})$$

$$= (1-\xi N^{-1/\kappa})^{-1}F\left(\kappa,1,2+\kappa;\frac{x}{x-1}\right)$$

$$= (1-\xi N^{-1/\kappa})^{-1}(1-x)F(2,1,2+\kappa;x)$$

$$= \frac{(z-1)^2}{(zN^{-1/\kappa}-z+1)^2}F(2,1;2+\kappa;x)$$

where $x = \frac{zN^{-1/\kappa}}{zN^{-1/\kappa}-z+1}$.

In sum, we have shown that the first derivative of the PGF can be expressed in the following way:

$$G_B'(z) = G_B(z)\frac{N\mu}{1+\kappa}\left(-\frac{N^{-1-\frac{1}{\kappa}}}{(zN^{-\frac{1}{\kappa}}-z+1)^2}F(2,1;2+\kappa;x) + F(2,1,2+\kappa,z)\right). \quad \text{[B1.1]}$$

Using the fact that $\lim_{z \to 1-} G_B(z) = 1$ and the Gauss identity $F(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}$ (which holds if $c > a + b$) we obtain

$$E[B] = G_B'(1-) = \frac{N\mu}{\kappa - 1}(-N^{-1+1/\kappa} + 1).$$

To calculate the variance for the number of mutants, we note that

$$G_B''(1-) = E[B^2] - E[B],$$

and thus

$$\mathrm{Var}[B] = G_B''(1-) + E[B] - (E[B])^2$$

$$= G_B''(1-) + G_B'(1-) - (G_B'(1-))^2$$

From equation [B1.1] we can express the first derivative of the PGF as:

$$G_B'(z) = G_B(z)\frac{N\mu}{1 + \kappa}\Sigma_1.$$

It follows that

$$\mathrm{Var}[B] = G_B''(1-) + G_B'(1-) - (G_B'(1-))^2$$

$$= \frac{N\mu}{1 + \kappa} \lim_{z \to 1-} (\Sigma_1 + \Sigma_1')$$

$$= E[B] + \frac{N\mu}{1 + \kappa} \lim_{z \to 1-} \Sigma_1'$$

To calculate the derivative of Σ_1, we note that

$$\Sigma_1 = -\frac{N^{-1-\frac{1}{\kappa}}}{(zN^{-\frac{1}{\kappa}} - z + 1)^2} F(2,1; 2 + \kappa; x) + F(2,1,2 + \kappa, z).$$

We can calculate the derivative of the second term:

$$\frac{\partial F(2,1,2 + \kappa, z)}{\partial z} = \frac{2}{2 + \kappa} F(3,2; 3 + \kappa; z),$$

and of the first term:

$$\frac{\partial[-\frac{N^{-1-1/\kappa}}{(zN^{-1/\kappa} - z + 1)^2} F(2,1; 2 + \kappa; x)]}{\partial z}$$

$$= -N^{-1-1/\kappa}\left(-\frac{2(N^{-1/\kappa} - 1)}{(N^{-1/\kappa}z - z + 1)^3} F(2,1; 2 + \kappa; x)\right.$$

$$\left. + \frac{2N^{-1/\kappa}}{(N^{-1/\kappa}z - z + 1)^4}\frac{1}{2 + \kappa} F(3,2; 3 + \kappa; x)\right)$$

Taken together, and using Gauss' equality,

$$\lim_{z \to 1-} \Sigma_1' = 2(N^{-1+1/\kappa} - N^{-1+2/\kappa})\frac{\kappa + 1}{\kappa - 1} + \frac{2}{2 + \kappa}\frac{(\kappa + 2)(\kappa + 1)}{(\kappa - 1)(\kappa - 2)}(1 - N^{-1+2/\kappa})$$

1652

$$= 2N^{-1+2/\kappa}\frac{\kappa+1}{\kappa-1}\left(-1-\frac{1}{\kappa-2}\right) + 2N^{-1+1/\kappa}\frac{\kappa+1}{\kappa-1} + \frac{2(\kappa+1)}{(\kappa-1)(\kappa-2)}$$

1653

$$= 2N^{-1+2/\kappa}\frac{\kappa+1}{2-\kappa} + 2N^{-1+1/\kappa}\frac{\kappa+1}{\kappa-1} + \frac{2(\kappa+1)}{(\kappa-1)(\kappa-2)}$$

1654 Coming back to

1655

$$\mathrm{Var}[B] = E[B] + \frac{N\mu}{1+\kappa}\lim_{z\to 1-}\Sigma_1',$$

1656 we obtain

1657

$$\mathrm{Var}[B] = N\mu\left[\frac{2}{2-\kappa}N^{-1+\frac{2}{\kappa}} + \frac{1}{\kappa-1}N^{-1+\frac{1}{\kappa}} + \frac{\kappa}{(\kappa-1)(\kappa-2)}\right].$$

1658 Making the appropriate substitutions yields our expressions for the mean and variance

1659 for our transconjugants.

1660

1661 **Box 2 : Behavior of the variance relative to the mean over time**

1662 In this Box, we will show that the variance in transconjugant numbers grows relative to

1663 the mean over time. To do so, we will need to consider several cases.

1664

1665 Let's consider the ratio of the variance to the mean, which we denote $\rho_t = \frac{\mathrm{Var}[T_t]}{E[T_t]}$. We will

1666 start by focusing on the case where $\psi_D + \psi_R \neq \psi_T$ and $\psi_D + \psi_R \neq 2\psi_T$:

1667

$$\rho_t = \frac{\gamma_D D_0 R_0 \left\{\frac{2e^{2\psi_T t}(\psi_T - (\psi_D + \psi_R)) - e^{\psi_T t}(2\psi_T - (\psi_D + \psi_R)) + (\psi_D + \psi_R)e^{(\psi_D+\psi_R)t}}{(2\psi_T - (\psi_D + \psi_R))(\psi_T - (\psi_D + \psi_R))}\right\}}{\frac{\gamma_D D_0 R_0 (e^{(\psi_D+\psi_R)t} - e^{\psi_T t})}{\psi_D + \psi_R - \psi_T}}.$$

1668 This can be simplified as follows:

1669

$$\rho_t = \frac{2e^{2\psi_T t}(\psi_T - (\psi_D + \psi_R)) - e^{\psi_T t}(2\psi_T - (\psi_D + \psi_R)) + (\psi_D + \psi_R)e^{(\psi_D+\psi_R)t}}{(\psi_D + \psi_R - 2\psi_T)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})}.$$

1670 Letting $\kappa = \frac{\psi_D+\psi_R}{\psi_T}$,

1671

$$\rho_t = \frac{2e^{2\psi_T t}(1-\kappa) - e^{\psi_T t}(2-\kappa) + \kappa e^{(\psi_D+\psi_R)t}}{(\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})},$$

1672

$$\rho_t = \frac{\kappa e^{(\psi_D+\psi_R)t} - 2(\kappa-1)e^{2\psi_T t} + (\kappa-2)e^{\psi_T t}}{(\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})}.$$

1673 To determine the behavior of this ratio over time we take the derivative with respect to

1674 time:

$$\frac{d\rho_t}{dt}$$

$$= \frac{\left\{\begin{array}{l}\left(\kappa(\psi_D + \psi_R)e^{(\psi_D+\psi_R)t} - 4\psi_T(\kappa-1)e^{2\psi_T t} + (\kappa-2)\psi_T e^{\psi_T t}\right)(\kappa-2)\left(e^{(\psi_D+\psi_R)t} - e^{\psi_T t}\right) \\ -\left(\kappa e^{(\psi_D+\psi_R)t} - 2(\kappa-1)e^{2\psi_T t} + (\kappa-2)e^{\psi_T t}\right)(\kappa-2)\left((\psi_D+\psi_R)e^{(\psi_D+\psi_R)t} - \psi_T e^{\psi_T t}\right)\end{array}\right\}}{\left((\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})\right)^2}$$

$$\frac{d\rho_t}{dt} = \frac{\left\{\begin{array}{l}\left(\kappa(\psi_D+\psi_R)e^{(\psi_D+\psi_R)t} - 4\psi_T(\kappa-1)e^{2\psi_T t} + (\kappa-2)\psi_T e^{\psi_T t}\right)(\kappa-2)\left(e^{(\psi_D+\psi_R)t}\right) \\ -\left(\kappa(\psi_D+\psi_R)e^{(\psi_D+\psi_R)t} - 4\psi_T(\kappa-1)e^{2\psi_T t} + (\kappa-2)\psi_T e^{\psi_T t}\right)(\kappa-2)\left(e^{\psi_T t}\right) \\ -\left(\kappa e^{(\psi_D+\psi_R)t} - 2(\kappa-1)e^{2\psi_T t} + (\kappa-2)e^{\psi_T t}\right)(\kappa-2)\left((\psi_D+\psi_R)e^{(\psi_D+\psi_R)t}\right) \\ +\left(\kappa e^{(\psi_D+\psi_R)t} - 2(\kappa-1)e^{2\psi_T t} + (\kappa-2)e^{\psi_T t}\right)(\kappa-2)\left(\psi_T e^{\psi_T t}\right)\end{array}\right\}}{\left((\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})\right)^2},$$

$$\frac{d\rho_t}{dt} = \frac{\left\{\begin{array}{l}\left(-4\psi_T(\kappa-1)e^{2\psi_T t} + (\kappa-2)\psi_T e^{\psi_T t}\right)(\kappa-2)\left(e^{(\psi_D+\psi_R)t}\right) \\ -\left(\kappa(\psi_D+\psi_R)e^{(\psi_D+\psi_R)t} - 4\psi_T(\kappa-1)e^{2\psi_T t}\right)(\kappa-2)\left(e^{\psi_T t}\right) \\ -\left(-2(\kappa-1)e^{2\psi_T t} + (\kappa-2)e^{\psi_T t}\right)(\kappa-2)\left((\psi_D+\psi_R)e^{(\psi_D+\psi_R)t}\right) \\ +\left(\kappa e^{(\psi_D+\psi_R)t} - 2(\kappa-1)e^{2\psi_T t}\right)(\kappa-2)\left(\psi_T e^{\psi_T t}\right)\end{array}\right\}}{\left((\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})\right)^2},$$

$$\frac{d\rho_t}{dt} = \frac{\left\{\begin{array}{l}(\psi_D+\psi_R-2\psi_T)2(\kappa-1)(\kappa-2)e^{2\psi_T t}\left(e^{(\psi_D+\psi_R)t}\right) \\ -(\psi_D+\psi_R-\psi_T)(\kappa-2)^2 e^{\psi_T t}\left(e^{(\psi_D+\psi_R)t}\right) \\ -(\psi_D+\psi_R-\psi_T)\kappa(\kappa-2)e^{\psi_T t}\left(e^{(\psi_D+\psi_R)t}\right) \\ +2\psi_T(\kappa-1)(\kappa-2)e^{3\psi_T t}\end{array}\right\}}{\left((\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})\right)^2},$$

$$\frac{d\rho_t}{dt} = \frac{\left\{\begin{array}{l}(\psi_D+\psi_R-2\psi_T)2(\kappa-1)(\kappa-2)e^{2\psi_T t}\left(e^{(\psi_D+\psi_R)t}\right) \\ -(\psi_D+\psi_R-\psi_T)2(\kappa-1)(\kappa-2)e^{\psi_T t}\left(e^{(\psi_D+\psi_R)t}\right) \\ +2\psi_T(\kappa-1)(\kappa-2)e^{3\psi_T t}\end{array}\right\}}{\left((\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})\right)^2},$$

$$\frac{d\rho_t}{dt} = \frac{(\kappa-1)(\kappa-2)\{(\kappa-2)e^{(\psi_D+\psi_R+2\psi_T)t} - (\kappa-1)e^{(\psi_D+\psi_R+\psi_T)t} + e^{3\psi_T t}\}}{\left(\frac{1}{2\psi_T}\right)\left((\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})\right)^2},$$

$$\frac{d\rho_t}{dt} = \frac{(\kappa-1)(\kappa-2)e^{\psi_T t}\{(\kappa-2)e^{(\psi_D+\psi_R+\psi_T)t} - (\kappa-1)e^{(\psi_D+\psi_R)t} + e^{2\psi_T t}\}}{\left(\frac{1}{2\psi_T}\right)\left((\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})\right)^2},$$

$$\frac{d\rho_t}{dt} = \frac{(\kappa-1)(\kappa-2)\{(\kappa-2)e^{(\kappa+1)\psi_T t} - (\kappa-1)e^{\kappa\psi_T t} + e^{2\psi_T t}\}}{\left(\frac{e^{-\psi_T t}}{2\psi_T}\right)\left((\kappa-2)(e^{(\psi_D+\psi_R)t} - e^{\psi_T t})\right)^2}.$$

The denominator is always positive, so the sign of this derivative is governed by the numerator

$$\text{sign}\left[\frac{d\rho_t}{dt}\right] = \text{sign}\left[(\kappa-1)(\kappa-2)\{(\kappa-2)e^{(\kappa+1)\psi_T t} - (\kappa-1)e^{\kappa\psi_T t} + e^{2\psi_T t}\}\right].$$

For $\kappa > 2$, which is when the growth rate of the transconjugant is less than the average

of the donor and recipient growth rates, we have

$$\text{sign}\left[\frac{d\rho_t}{dt}\right] = \text{sign}\left[(\kappa-2)e^{(\kappa+1)\psi_T t} - (\kappa-1)e^{\kappa\psi_T t} + e^{2\psi_T t}\right].$$

For any $t > 0$, for $\kappa = 2$

$$\text{sign}\left[\frac{d\rho_t}{dt}\right] = \text{sign}\left[0 - e^{2\psi_T t} + e^{2\psi_T t}\right] = 0.$$

Now letting $g(\kappa) = (\kappa-2)e^{(\kappa+1)\psi_T t} - (\kappa-1)e^{\kappa\psi_T t} + e^{2\psi_T t}$, we have

$$g'(\kappa) = e^{(\kappa+1)\psi_T t} + (\kappa-2)\psi_T t e^{(\kappa+1)\psi_T t} - e^{\kappa\psi_T t} - (\kappa-1)\psi_T t e^{\kappa\psi_T t},$$

$$g'(\kappa) = \{1 + (\kappa-2)\psi_T t\}e^{(\kappa+1)\psi_T t} - \{1 + (\kappa-1)\psi_T t\}e^{\kappa\psi_T t}.$$

Letting $h(x) = \{1 + (\kappa-1-x)\psi_T t\}e^{(\kappa+x)\psi_T t}$, we see

$$g'(\kappa) = h(1) - h(0).$$

But we see that

$$h'(x) = -\psi_T t e^{(\kappa+x)\psi_T t} + \{1 + (\kappa-1-x)\psi_T t\}\psi_T t e^{(\kappa+x)\psi_T t},$$

$$h'(x) = \psi_T t e^{(\kappa+x)\psi_T t}(-1 + \{1 + (\kappa-1-x)\psi_T t\}),$$

$$h'(x) = \psi_T t e^{(\kappa+x)\psi_T t}(\kappa-1-x)\psi_T t.$$

If $\kappa > 2$ and $t > 0$, then $h'(x) > 0$ for $0 \le x \le 1$. Since $h(x)$ is a continuous function, we

must have

$$h(1) > h(0).$$

This implies that for $\kappa > 2$ and $t > 0$,

$$g'(\kappa) > 0.$$

Since $g(\kappa)$ is also continuous, and since $g(2) = 0$, we now know that for $\kappa > 2$ and $t >$

0,

$$g(\kappa) > 0.$$

Therefore $\text{sign}\left[\frac{d\rho_t}{dt}\right] = 1$, or the derivative is positive for $t > 0$ and $\kappa > 2$, which means

the variance grows relative to the mean over time.

For $1 < \kappa < 2$,

$$\text{sign}\left[\frac{d\rho_t}{dt}\right] = \text{sign}\left[-(\kappa-2)e^{(\kappa+1)\psi_T t} + (\kappa-1)e^{\kappa\psi_T t} - e^{2\psi_T t}\right].$$

Letting $\varphi = -(\kappa-2)$, we know that $0 < \varphi < 1$, and we can rewrite the above

$$\text{sign}\left[\frac{d\rho_t}{dt}\right] = \text{sign}\left[\varphi e^{(3-\varphi)\psi_T t} + (1-\varphi)e^{(3-(1+\varphi))\psi_T t} - e^{2\psi_T t}\right].$$

Because $e^{(3-x)\psi_T t}$ is convex $\left(t > 0 \implies \frac{d^2 e^{(3-x)\psi_T t}}{dx^2} > 0\right)$, Jensen's inequality ensures

$$\varphi e^{(3-\varphi)\psi_T t} + (1-\varphi)e^{(3-(1+\varphi))\psi_T t} - e^{2\psi_T t} > e^{\left(3-\left(\varphi^2+(1-\varphi)(1+\varphi)\right)\right)\psi_T t} - e^{2\psi_T t},$$

$$\varphi e^{(3-\varphi)\psi_T t} + (1-\varphi)e^{(3-(1+\varphi))\psi_T t} - e^{2\psi_T t} > e^{\left(3-\left(\varphi^2+1-\varphi^2\right)\right)\psi_T t} - e^{2\psi_T t},$$

$$\varphi e^{(3-\varphi)\psi_T t} + (1-\varphi)e^{(3-(1+\varphi))\psi_T t} - e^{2\psi_T t} > e^{2\psi_T t} - e^{2\psi_T t},$$

$$\varphi e^{(3-\varphi)\psi_T t} + (1-\varphi)e^{(3-(1+\varphi))\psi_T t} - e^{2\psi_T t} > 0.$$

Therefore $\text{sign}\left[\frac{d\rho_t}{dt}\right] = 1$, or the derivative is positive for $t > 0$ and $1 < \kappa < 2$, which means the variance grows relative to the mean over time.

Next, we turn to $0 < \kappa < 1$, where

$$\text{sign}\left[\frac{d\rho_t}{dt}\right] = \text{sign}\left[(\kappa-2)e^{(\kappa+1)\psi_T t} - (\kappa-1)e^{\kappa\psi_T t} + e^{2\psi_T t}\right].$$

For any $t > 0$, for $\kappa = 1$

$$\text{sign}\left[\frac{d\rho_t}{dt}\right] = \text{sign}\left[-e^{2\psi_T t} - 0 + e^{2\psi_T t}\right] = 0.$$

Now letting $g(\kappa) = (\kappa-2)e^{(\kappa+1)\psi_T t} - (\kappa-1)e^{\kappa\psi_T t} + e^{2\psi_T t}$, we have from above

$$g'(\kappa) = \{1 + (\kappa-2)\psi_T t\}e^{(\kappa+1)\psi_T t} - \{1 + (\kappa-1)\psi_T t\}e^{\kappa\psi_T t}.$$

Letting $h(x) = \{1 + (\kappa-1-x)\psi_T t\}e^{(\kappa+x)\psi_T t}$, we see

$$g'(\kappa) = h(1) - h(0).$$

But we see from above that

$$h'(x) = \psi_T t e^{(\kappa+x)\psi_T t}(\kappa-1-x)\psi_T t.$$

If $0 < \kappa < 1$ and $t > 0$, then $h'(x) < 0$ for $0 \le x \le 1$. Since $h(x)$ is a continuous function, we must have

$$h(1) < h(0).$$

This implies that for $0 < \kappa < 1$ and $t > 0$,

$$g'(\kappa) < 0.$$

Since $g(\kappa)$ is continuous, and since $g(1) = 0$, we now know that for $0 < \kappa < 1$ and $t > 0$, $g(\kappa) > 0$. Therefore $\text{sign}\left[\frac{d\rho_t}{dt}\right] = 1$, or the derivative is positive for $t > 0$ and $0 < \kappa < 1$, which means the variance grows relative to the mean over time.

Now consider the special case of $\kappa = 2$

$$\rho_t = \frac{\dfrac{2\gamma_D D_0 R_0\left(e^{\psi_T t} - e^{2\psi_T t}\right)}{2\psi_T} + 2\gamma_D D_0 R_0 e^{2\psi_T t} t}{\dfrac{\gamma_D D_0 R_0(e^{2\psi_T t} - e^{\psi_T t})}{2\psi_T - \psi_T}},$$

$$\rho_t = \frac{\left(e^{\psi_T t} - e^{2\psi_T t}\right) + 2\psi_T t e^{2\psi_T t}}{(e^{2\psi_T t} - e^{\psi_T t})},$$

$$\rho_t = \frac{(2\psi_T t - 1)e^{2\psi_T t} + e^{\psi_T t}}{(e^{2\psi_T t} - e^{\psi_T t})}.$$

Taking the derivative with respect to time yields

$$\frac{d\rho_t}{dt} = \frac{\left\{\begin{array}{c}\left(2\psi_T e^{2\psi_T t} + 2\psi_T(2\psi_T t - 1)e^{2\psi_T t} + \psi_T e^{\psi_T t}\right)\left(e^{2\psi_T t} - e^{\psi_T t}\right) \\ -\left((2\psi_T t - 1)e^{2\psi_T t} + e^{\psi_T t}\right)\left(2\psi_T e^{2\psi_T t} - \psi_T e^{\psi_T t}\right)\end{array}\right\}}{(e^{2\psi_T t} - e^{\psi_T t})^2},$$

$$\frac{d\rho_t}{dt} = \psi_T \frac{\left(4\psi_T t e^{2\psi_T t} + e^{\psi_T t}\right)\left(e^{2\psi_T t} - e^{\psi_T t}\right) - \left((2\psi_T t - 1)e^{2\psi_T t} + e^{\psi_T t}\right)\left(2e^{2\psi_T t} - e^{\psi_T t}\right)}{(e^{2\psi_T t} - e^{\psi_T t})^2},$$

$$\frac{d\rho_t}{dt} = \psi_T \frac{\left\{\begin{array}{c}4\psi_T t e^{4\psi_T t} + e^{3\psi_T t} - 4\psi_T t e^{3\psi_T t} - e^{2\psi_T t} \\ -2(2\psi_T t - 1)e^{4\psi_T t} - 2e^{3\psi_T t} + (2\psi_T t - 1)e^{3\psi_T t} + e^{2\psi_T t}\end{array}\right\}}{(e^{2\psi_T t} - e^{\psi_T t})^2},$$

$$\frac{d\rho_t}{dt} = \psi_T \frac{\{4\psi_T t - 2(2\psi_T t - 1)\}e^{4\psi_T t} - \{4\psi_T t + 1 - (2\psi_T t - 1)\}e^{3\psi_T t}}{(e^{2\psi_T t} - e^{\psi_T t})^2},$$

$$\frac{d\rho_t}{dt} = \psi_T \frac{2e^{4\psi_T t} - \{2\psi_T t + 2\}e^{3\psi_T t}}{(e^{2\psi_T t} - e^{\psi_T t})^2},$$

$$\frac{d\rho_t}{dt} = 2\psi_T e^{3\psi_T t} \frac{e^{\psi_T t} - \psi_T t - 1}{(e^{2\psi_T t} - e^{\psi_T t})^2}.$$

Using the Taylor expansion

$$\frac{d\rho_t}{dt} = 2\psi_T e^{3\psi_T t} \frac{\left(1 + \psi_T t + \dfrac{(\psi_T t)^2}{2!} + \dfrac{(\psi_T t)^3}{3!} + \dfrac{(\psi_T t)^4}{4!} \cdots\right) - \psi_T t - 1}{(e^{2\psi_T t} - e^{\psi_T t})^2},$$

$$\frac{d\rho_t}{dt} = 2\psi_T e^{3\psi_T t} \frac{\left(\dfrac{(\psi_T t)^2}{2!} + \dfrac{(\psi_T t)^3}{3!} + \dfrac{(\psi_T t)^4}{4!} \cdots\right)}{(e^{2\psi_T t} - e^{\psi_T t})^2}.$$

Therefore, $\frac{d\rho_t}{dt} > 0$ for $t > 0$ and $\kappa = 2$, which means the variance grows relative to the mean over time.

Finally, consider the special case of $\kappa = 1$

$$\rho_t = \frac{\dfrac{2\gamma_D D_0 R_0\left(e^{2\psi_T t} - e^{\psi_T t}\right)}{\psi_T} - \gamma_D D_0 R_0 e^{\psi_T t} t}{\gamma_D D_0 R_0 e^{\psi_T t} t},$$

$$\rho_t = \frac{2e^{2\psi_T t} - (2 + \psi_T t)e^{\psi_T t}}{\psi_T t e^{\psi_T t}},$$

$$\rho_t = \frac{2e^{\psi_T t} - (2 + \psi_T t)}{\psi_T t}.$$

Using the Taylor expansion

$$\rho_t = \frac{2\left(1 + \psi_T t + \frac{(\psi_T t)^2}{2!} + \frac{(\psi_T t)^3}{3!} + \frac{(\psi_T t)^4}{4!} \ldots\right) - (2 + \psi_T t)}{\psi_T t},$$

$$\rho_t = \frac{\psi_T t + 2\left(\frac{(\psi_T t)^2}{2!} + \frac{(\psi_T t)^3}{3!} + \frac{(\psi_T t)^4}{4!} \ldots\right)}{\psi_T t},$$

$$\rho_t = 1 + 2\left(\frac{\psi_T t}{2!} + \frac{(\psi_T t)^2}{3!} + \frac{(\psi_T t)^3}{4!} \ldots\right).$$

Once again, the variance grows relative to the mean over time.

Overall, we have shown that for all $\kappa > 0$ and $t > 0$, the ratio of the transconjugant variance to the mean number of transconjugants is amplified over time.

SI section 14 : Variance in Estimates

Here we will focus on two estimates, ASM and LDM, and ask about their variance (enabling us to compare precision). We will focus exclusively on the contributions to this variance coming from the stochasticity in the transconjugant numbers (i.e., ignoring contributions coming from assessment of initial and final donor and recipient populations).

We start with the ASM estimate (here we express the estimate in terms of growth rate parameters):

$$\gamma_D = \frac{\psi_D + \psi_R - \psi_T}{D_0 R_0 \left(e^{(\psi_D + \psi_R)\tilde{t}} - e^{\psi_T \tilde{t}}\right)} T_{\tilde{t}}.$$

Because we are only focusing on the contribution of the transconjugant variation, all parameters (initial densities and growth rates will be taken to be fixed). Thus, we can think about the ASM estimate as a random variable Γ_{ASM}, where

$$\Gamma_{ASM} = c_1 T_{\tilde{t}},$$

where the constant c_1 is

$$c_1 = \frac{\psi_D + \psi_R - \psi_T}{D_0 R_0 \left(e^{(\psi_D + \psi_R)\tilde{t}} - e^{\psi_T \tilde{t}}\right)}.$$

1787   The variance of the ASM estimate is then

1788
$$\mathrm{var}(\Gamma_{ASM}) = c_1{}^2\{\mathrm{var}(T_{\tilde{t}})\}.$$

1789   But we have a closed form expression for $\mathrm{var}(T_{\tilde{t}})$. If $\psi_T \notin \{\psi_D + \psi_R, (\psi_D + \psi_R)/2\}$, we

1790   have:

1791
$$\mathrm{var}(\Gamma_{ASM}) = \left(\frac{\psi_D+\psi_R-\psi_T}{D_0 R_0\left(e^{(\psi_D+\psi_R)\tilde{t}}-e^{\psi_T\tilde{t}}\right)}\right)^2 \gamma_D D_0 R_0 \left\{\frac{(\psi_D+\psi_R)e^{(\psi_D+\psi_R)\tilde{t}}}{(\psi_D+\psi_R-\psi_T)(\psi_D+\psi_R-2\psi_T)} + \frac{e^{\psi_T\tilde{t}}}{\psi_D+\psi_R-\psi_T} - \frac{2e^{2\psi_T\tilde{t}}}{\psi_D+\psi_R-2\psi_T}\right\},$$

1792
$$\mathrm{var}(\Gamma_{ASM}) = \frac{\gamma_D}{D_0 R_0}\left(\frac{\psi_D+\psi_R-\psi_T}{e^{(\psi_D+\psi_R)\tilde{t}}-e^{\psi_T\tilde{t}}}\right)^2 \left\{\frac{(\psi_D+\psi_R)e^{(\psi_D+\psi_R)\tilde{t}}+(\psi_D+\psi_R-2\psi_T)e^{\psi_T\tilde{t}}-(\psi_D+\psi_R-\psi_T)2e^{2\psi_T\tilde{t}}}{(\psi_D+\psi_R-\psi_T)(\psi_D+\psi_R-2\psi_T)}\right\},$$

1793
$$\mathrm{var}(\Gamma_{ASM}) = \frac{\gamma_D(\psi_D+\psi_R-\psi_T)}{D_0 R_0}\left\{\frac{(\psi_D+\psi_R)e^{(\psi_D+\psi_R)\tilde{t}}+(\psi_D+\psi_R-2\psi_T)e^{\psi_T\tilde{t}}-(\psi_D+\psi_R-\psi_T)2e^{2\psi_T\tilde{t}}}{(\psi_D+\psi_R-2\psi_T)\left(e^{(\psi_D+\psi_R)\tilde{t}}-e^{\psi_T\tilde{t}}\right)^2}\right\}.$$

1794   The formulas for $\psi_T = \psi_D + \psi_R$ and $2\psi_T = \psi_D + \psi_R$ could also be derived via simple

1795   substitution (note, $\lim_{\psi_T \to \psi_D+\psi_R} c_1 = 1/(D_0 R_0 t e^{(\psi_D+\psi_R)\tilde{t}})$. These formulas allow us to project

1796   variance in the ASM estimate over time due to transconjugant variation if all parameters

1797   are known.

1798

1799   We now turn to the LDM estimate:

1800
$$\gamma_D = -\ln \hat{p}_0(\tilde{t})\left(\frac{\psi_D + \psi_R}{D_0 R_0\left(e^{(\psi_D+\psi_R)\tilde{t}} - 1\right)}\right).$$

1801   What we actually measure is the number of populations (or wells) that have no

1802   transconjugants (call this w) out of the total number of populations (or wells) tracked (call

1803   this W). As we show in SI section 9, the maximum likelihood estimate of $p_0(\tilde{t})$ is

1804
$$\hat{p}_0(\tilde{t}) = \frac{w}{W}.$$

1805   Of course, from experiment to experiment, there will be variance in the number of

1806   populations with no transconjugants. Let us consider a random variable F, which

1807   represents the fraction of total populations that have no transconjugants. The expectation

1808   of F is (we drop the time argument for notational convenience):

1809
$$E[F] = \sum_{w=0}^{W} \binom{W}{w}(p_0)^w(1-p_0)^{W-w}\left(\frac{w}{W}\right),$$

1810
$$E[F] = \frac{1}{W}\sum_{w=1}^{W}\frac{W!}{w!\,(W-w)!}(p_0)^w(1-p_0)^{W-w}\,w,$$

1811
$$E[F] = \frac{1}{W}\sum_{w=1}^{W}\frac{W!}{(w-1)!\,(W-w)!}(p_0)^w(1-p_0)^{W-w},$$

$$1812 \qquad E[F] = p_0 \sum_{w=1}^{W} \frac{(W-1)!}{(w-1)! \left((W-1)-(w-1)\right)!} (p_0)^{w-1} (1-p_0)^{(W-1)-(w-1)}.$$

1813 Letting $i = w - 1$, we have

$$1814 \qquad E[F] = p_0 \sum_{i=0}^{W-1} \frac{(W-1)!}{i! \left((W-1)-i\right)!} (p_0)^i (1-p_0)^{(W-1)-i}.$$

1815 However, because $\sum_{i=0}^{W-1} \binom{W-1}{i} (p_0)^i (1-p_0)^{(W-1)-i} = 1$, we have

$$1816 \qquad E[F] = p_0,$$

1817 which makes sense. The second central moment of F is

$$1818 \qquad \text{var}[F] = \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w} \left(\frac{w}{W} - p_0\right)^2,$$

$$1819 \qquad \text{var}[F] = \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w} \left(\left(\frac{w}{W}\right)^2 - 2p_0 \left(\frac{w}{W}\right) + (p_0)^2\right),$$

$$1820 \qquad \text{var}[F] = \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w} \left(\frac{w}{W}\right)^2 - 2p_0 \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w} \left(\frac{w}{W}\right)$$

$$1821 \qquad + (p_0)^2 \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w},$$

$$1822 \qquad \text{var}[F] = \left\{\left(\frac{1}{W^2}\right) \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w} w^2\right\} - 2p_0 E[F] + (p_0)^2,$$

$$1823 \qquad \text{var}[F] = \left\{\left(\frac{1}{W^2}\right) \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w} w(w-1+1)\right\} - 2(p_0)^2 + (p_0)^2,$$

$$1824 \qquad \text{var}[F] = \left\{\left(\frac{1}{W^2}\right) \sum_{w=2}^{W} \frac{W!}{w! \, (W-w)!} (p_0)^w (1-p_0)^{W-w} w(w-1)\right.$$

$$1825 \qquad \left. + \left(\frac{1}{W^2}\right) \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w} w\right\} - (p_0)^2,$$

$$1826 \qquad \text{var}[F] = \left\{\left(\frac{1}{W^2}\right) \sum_{w=2}^{W} \frac{W!}{(w-2)! \, (W-w)!} (p_0)^w (1-p_0)^{W-w}\right.$$

$$1827 \qquad \left. + \left(\frac{1}{W}\right) \sum_{w=0}^{W} \binom{W}{w} (p_0)^w (1-p_0)^{W-w} \left(\frac{w}{W}\right)\right\} - (p_0)^2,$$

1828
$$\text{var}[F] = \left(\frac{W(W-1)(p_0)^2}{W^2}\right)\left\{\sum_{w=2}^{W}\frac{(W-2)!}{(w-2)!\,((W-2)-(w-2))!}(p_0)^{w-2}(1\right.$$

1829
$$\left.-p_0)^{(W-2)-(w-2)}\right\} + \frac{p_0}{W} - (p_0)^2.$$

1830   Letting $j = w - 2$, we have

1831
$$\text{var}[F] = \left(\frac{(W-1)(p_0)^2}{W}\right)\left\{\sum_{j=0}^{W-2}\binom{W-2}{j}(p_0)^j(1-p_0)^{(W-2)-j}\right\} + \frac{p_0}{W} - (p_0)^2,$$

1832
$$\text{var}[F] = \left(\frac{(W-1)(p_0)^2}{W}\right) + \frac{p_0}{W} - (p_0)^2,$$

1833
$$\text{var}[F] = \frac{(W-1)(p_0)^2 + p_0 - W(p_0)^2}{W},$$

1834
$$\text{var}[F] = \frac{-(p_0)^2 + p_0}{W},$$

1835
$$\text{var}[F] = \frac{p_0(1-p_0)}{W}.$$

1836

1837   Because we are only focusing on the contribution of the transconjugant variation, all

1838   parameters (initial densities and growth rates will be taken to be fixed). Thus, we can think

1839   about the LDM estimate as a random variable Γ_{LDM},

1840
$$\Gamma_{\text{LDM}} = c_2 \ln F,$$

1841   where the constant c_2 is

1842
$$c_2 = -\left(\frac{\psi_D + \psi_R}{D_0 R_0\left(e^{(\psi_D+\psi_R)\tilde{t}} - 1\right)}\right).$$

1843

1844   The variance of the LDM estimate is then

1845
$$\text{var}(\Gamma_{\text{LDM}}) = c_2{}^2\{\text{var}(\ln F)\}.$$

1846   Here we use a first-order Taylor series approximation for $\ln F$ centered at $\text{E}[F]$ in the

1847   following way:

1848
$$\ln F \approx \ln(\text{E}[F]) + \frac{1}{\text{E}[F]}(F - \text{E}[F]),$$

1849
$$\ln F \approx \frac{F}{\text{E}[F]} + \ln(\text{E}[F]) - 1.$$

1850

1851

1852 So, because $\ln(E[F]) - 1$ is constant, we have

$$\text{var}[\ln F] \approx \text{var}\left\{\frac{F}{E[F]}\right\},$$

1853

$$\text{var}[\ln F] \approx \left(\frac{1}{E[F]}\right)^2 \text{var}[F],$$

1854

$$\text{var}[\ln F] \approx \left(\frac{1}{p_0}\right)^2 \left(\frac{p_0(1 - p_0)}{W}\right),$$

1855

$$\text{var}[\ln F] \approx \frac{1}{W}\left(\frac{1}{p_0} - 1\right).$$

1856

1857 This approximation will be accurate when the deviation between F and $E[F]$ is very small

1858 (i.e., $\frac{|F - E[F]|}{E[F]} \ll 1$). As W (the number of replicate populations in the experiment) gets large,

1859 the distribution of F will tighten around $E[F]$, making the approximation more reasonable.

1860

1861 Now, we have the following expression for p_0 (reintroducing the time argument):

$$p_0(\tilde{t}) = \exp\left\{\frac{-\gamma_D D_0 R_0}{\psi_D + \psi_R}\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right)\right\}.$$

1862

1863 Therefore,

$$\text{var}[\ln F_{\tilde{t}}] \approx \frac{1}{W}\left(\exp\left\{\frac{\gamma_D D_0 R_0}{\psi_D + \psi_R}\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right)\right\} - 1\right),$$

1864

1865 where we make the time dependence of F clear. Returning to the variance for the LDM

1866 estimate,

$$\text{var}(\Gamma_{\text{LDM}}) \approx \frac{1}{W}\left(\frac{\psi_D + \psi_R}{D_0 R_0\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right)}\right)^2 \left(\exp\left\{\frac{\gamma_D D_0 R_0}{\psi_D + \psi_R}\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right)\right\} - 1\right).$$

1867

1868

1869 If we define

$$\xi_{\tilde{t}} = \frac{\psi_D + \psi_R}{D_0 R_0\left(e^{(\psi_D + \psi_R)\tilde{t}} - 1\right)},$$

1870

1871 then we have

$$\text{var}(\Gamma_{\text{LDM}}) \approx \frac{\xi_{\tilde{t}}^2}{W}\left(e^{\left(\frac{\gamma_D}{\xi_{\tilde{t}}}\right)} - 1\right).$$

1872

1873

In SI Figure 3, we explore the variances (approximate in the case of LDM) as a function

of time. The LDM estimates (for two different numbers of populations) are more precise

(lower variance) for much of the time range. However, if the time gets too high ($\tilde{t} \approx 5$ for

the parameter set shown in SI Figure 3), then the LDM variance blows up (while the ASM

variance remains very low). In a case like this, the LDM is predicted to be more precise

when the time of the assay is sufficiently low.

SI Figure 3 : The variance of the ASM (green) and LDM (pink) estimates. Different numbers of populations (W) are used for the LDM estimates, as indicated. The parameters used here are $\gamma_D = 10^{-12}$, $D_0 = R_0 = 10^4$, $\psi_D = 1$, and $\psi_R = \psi_T = 1.5$.

1880   To show mathematically that the LDM estimate is more precise for short times, we will

1881   approximate the expressions for the variance when $\tilde{t}$ is very small. This enables us to

1882   use a first-order Maclaurin approximation $e^{c\tilde{t}} \approx 1 + c\tilde{t}$. This allows the following

1883   approximation of the variance for the ASM estimate:

1884 $\mathrm{var}(\Gamma_{\mathrm{ASM}})$

1885
$$\approx \frac{\gamma_D(\psi_D + \psi_R - \psi_T)}{D_0 R_0}\left\{\frac{(\psi_D + \psi_R)(1 + (\psi_D + \psi_R)\tilde{t}) + (\psi_D + \psi_R - 2\psi_T)(1 + \psi_T\tilde{t}) - 2(\psi_D + \psi_R - \psi_T)(1 + 2\psi_T\tilde{t})}{(\psi_D + \psi_R - 2\psi_T)((\psi_D + \psi_R)\tilde{t} - \psi_T\tilde{t})^2}\right\}.$$

1886   We can then simplify this expression:

1887
$$\mathrm{var}(\Gamma_{\mathrm{ASM}}) \approx \frac{\gamma_D(\psi_D + \psi_R - \psi_T)}{D_0 R_0}\left\{\frac{(\psi_D + \psi_R)(\psi_D + \psi_R)\tilde{t} + (\psi_D + \psi_R - 2\psi_T)\psi_T\tilde{t} - 2(\psi_D + \psi_R - \psi_T)2\psi_T\tilde{t}}{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R - \psi_T)^2\tilde{t}^2}\right\},$$

1888
$$\mathrm{var}(\Gamma_{\mathrm{ASM}}) \approx \frac{\gamma_D}{D_0 R_0 \tilde{t}}\left\{\frac{(\psi_D + \psi_R)(\psi_D + \psi_R) + (\psi_D + \psi_R - 2\psi_T)\psi_T - 2(\psi_D + \psi_R - \psi_T)2\psi_T}{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R - \psi_T)}\right\},$$

1889 $\mathrm{var}(\Gamma_{\mathrm{ASM}})$

1890
$$\approx \frac{\gamma_D}{D_0 R_0 \tilde{t}}\left\{\frac{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R) + 2\psi_T(\psi_D + \psi_R) + (\psi_D + \psi_R - 2\psi_T)\psi_T - 2(\psi_D + \psi_R - \psi_T)2\psi_T}{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R - \psi_T)}\right\},$$

1891 $\mathrm{var}(\Gamma_{\mathrm{ASM}})$

1892
$$\approx \frac{\gamma_D}{D_0 R_0 \tilde{t}}\left\{\frac{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R) + 2\psi_T\psi_D + 2\psi_T\psi_R + \psi_T\psi_D + \psi_T\psi_R - 2\psi_T^2 - 4\psi_T\psi_D - 4\psi_T\psi_R + 4\psi_T^2}{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R - \psi_T)}\right\},$$

1893
$$\mathrm{var}(\Gamma_{\mathrm{ASM}}) \approx \frac{\gamma_D}{D_0 R_0 \tilde{t}}\left\{\frac{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R) - \psi_T\psi_D - \psi_T\psi_R + 2\psi_T^2}{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R - \psi_T)}\right\},$$

1894

$$\text{var}(\Gamma_{\text{ASM}}) \approx \frac{\gamma_D}{D_0 R_0 \tilde{t}} \left\{ \frac{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R) - (\psi_D + \psi_R - 2\psi_T)\psi_T}{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R - \psi_T)} \right\},$$

$$\text{var}(\Gamma_{\text{ASM}}) \approx \frac{\gamma_D}{D_0 R_0 \tilde{t}} \left\{ \frac{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R - \psi_T)}{(\psi_D + \psi_R - 2\psi_T)(\psi_D + \psi_R - \psi_T)} \right\},$$

$$\text{var}(\Gamma_{\text{ASM}}) \approx \frac{\gamma_D}{D_0 R_0 \tilde{t}}.$$

We now turn to the variance for the LDM (which we note is already an approximation). For very small $\tilde{t}$,

$$\xi_{\tilde{t}} \approx \frac{\psi_D + \psi_R}{D_0 R_0 (1 + (\psi_D + \psi_R)\tilde{t} - 1)},$$

$$\xi_{\tilde{t}} \approx \frac{1}{D_0 R_0 \tilde{t}}.$$

Thus, we have

$$\text{var}(\Gamma_{\text{LDM}}) \approx \frac{\left(\frac{1}{D_0 R_0 \tilde{t}}\right)^2}{W} \left(e^{(\gamma_D D_0 R_0 \tilde{t})} - 1 \right).$$

Using the Maclaurin approximation again yields

$$\text{var}(\Gamma_{\text{LDM}}) \approx \frac{\left(\frac{1}{D_0 R_0 \tilde{t}}\right)^2}{W} (1 + \gamma_D D_0 R_0 \tilde{t} - 1),$$

$$\text{var}(\Gamma_{\text{LDM}}) \approx \frac{\gamma_D}{W D_0 R_0 \tilde{t}}.$$

Our LDM assay requires $W > 1$. Therefore when $\tilde{t}$ is very small

$$\text{var}(\Gamma_{\text{LDM}}) < \text{var}(\Gamma_{\text{ASM}}).$$

Again, we note the caveat that our estimate for the variance of the LDM estimate was already an approximation.

References

1. Redondo-Salvo, S. *et al.* Pathways for horizontal gene transfer in bacteria revealed by a global map of their plasmids. *Nat. Commun.* **11**, 3602 (2020).

2. Loftie-Eaton, W. *et al.* Compensatory mutations improve general permissiveness to antibiotic resistance plasmids. *Nat. Ecol. Evol.* **1**, 1354–1363 (2017).

3. Norman, A., Hansen, L. H. & Sørensen, S. J. Conjugative plasmids: vessels of the communal gene pool. *Philos. Trans. R. Soc. B Biol. Sci.* **364**, 2275–2289 (2009).

4. Jordt, H. *et al.* Coevolution of host–plasmid pairs facilitates the emergence of novel

multidrug resistance. *Nat. Ecol. Evol.* 1–7 (2020) doi:10.1038/s41559-020-1170-1.

5. Dahlberg, C. & Chao, L. Amelioration of the Cost of Conjugative Plasmid Carriage in

Eschericha coli K12. *Genetics* **165**, 1641–1649 (2003).

6. Stewart, F. M. & Levin, B. R. The Population Biology of Bacterial Plasmids: A Priori

Conditions for the Existence of Conjugationally Transmitted Factors. *Genetics* **87**,

209–228 (1977).

7. Lopatkin, A. J. *et al.* Persistence and reversal of plasmid-mediated antibiotic

resistance. *Nat. Commun.* **8**, 1–10 (2017).

8. Fox, R. E., Zhong, X., Krone, S. M. & Top, E. M. Spatial structure and nutrients

promote invasion of IncP-1 plasmids in bacterial populations. *ISME J.* **2**, 1024–1039

(2008).

9. Alderliesten, J. B. *et al.* Effect of donor-recipient relatedness on the plasmid

conjugation frequency: a meta-analysis. *BMC Microbiol.* **20**, 135 (2020).

10. Zhong, X., Droesch, J., Fox, R., Top, E. M. & Krone, S. M. On the meaning and

estimation of plasmid transfer rates for surface-associated and well-mixed bacterial

populations. *J. Theor. Biol.* **294**, 144–152 (2012).

11. Huisman, J. S. *et al.* Estimating the rate of plasmid transfer in liquid mating cultures.

bioRxiv 2020.03.09.980862 (2020) doi:10.1101/2020.03.09.980862.

12. Simonsen, L. The Existence Conditions for Bacterial Plasmids: Theory and Reality.

Microb. Ecol. **22**, 187–205 (1991).

13. Levin, B. R., Stewart, F. M. & Rice, V. A. The kinetics of conjugative plasmid

transmission: Fit of a simple mass action model. *Plasmid* **2**, 247–260 (1979).

14. Zhong, X., Droesch, J., Fox, R., Top, E. M. & Krone, S. M. On the meaning and

estimation of plasmid transfer rates for surface-associated and well-mixed bacterial

populations. *J. Theor. Biol.* **294**, 144–152 (2012).

15. Bethke, J. H. *et al.* Environmental and genetic determinants of plasmid mobility in

pathogenic Escherichia coli. *Sci. Adv.* **6**, eaax3173 (2020).

16. Simonsen, L., Gordon, D. M., Stewart, F. M. & Levin, B. R. Estimating the rate of

plasmid transfer: an end-point method. *Microbiology,* **136**, 2319–2325 (1990).

17. Lopatkin, A. J. *et al.* Antibiotics as a selective driver for conjugation dynamics. *Nat.*

Microbiol. **1**, 1–8 (2016).

18. San Millan, A. Evolution of Plasmid-Mediated Antibiotic Resistance in the Clinical

Context. *Trends Microbiol.* **26**, 978–985 (2018).

19. Li, L. *et al.* Plasmids persist in a microbial community by providing fitness benefit to

multiple phylotypes. *ISME J.* 1–12 (2020) doi:10.1038/s41396-020-0596-4.

20. Luria, S. E. & Delbrück, M. Mutations of Bacteria from Virus Sensitivity to Virus

Resistance. *Genetics* **28**, 491–511 (1943).

21. Johnsen, A. R. & Kroer, N. Effects of stress and other environmental factors on

horizontal plasmid transfer assessed by direct quantification of discrete transfer

events. *FEMS Microbiol. Ecol.* **59**, 718–728 (2007).

22. Loftie-Eaton, W. *et al.* Evolutionary Paths That Expand Plasmid Host-Range:

Implications for Spread of Antibiotic Resistance. *Mol. Biol. Evol.* **33**, 885–897 (2016).

23. Yano, H. *et al.* Evolved plasmid-host interactions reduce plasmid interference cost.

Mol. Microbiol. **101**, 743–756 (2016).

24. Valle, A. A. *et al.* The distribution of plasmid fitness effects explains plasmid

persistence in bacterial communities. *bioRxiv* 2020.08.01.230672 (2020)

doi:10.1101/2020.08.01.230672.

25. Abboud, R. *et al.* Low-Temperature Growth of Shewanella oneidensis MR-1. *Appl.*

Environ. Microbiol. **71**, 811–816 (2005).

26. Gibson, B., Wilson, D. J., Feil, E. & Eyre-Walker, A. The distribution of bacterial

doubling times in the wild. *Proc. R. Soc. B Biol. Sci.* **285**, (2018).

27. Dimitriu, T., Marchant, L., Buckling, A. & Raymond, B. Bacteria from natural

populations transfer plasmids mostly towards their kin. *Proc. R. Soc. B Biol. Sci.*

286, 20191110 (2019).

28. Lundquist, P. D. & Levin, B. R. Transitory Derepression and the Maintenance of

Conjugative Plasmids. *Genetics* **113**, 483–497 (1986).

29. Sysoeva, T. A., Kim, Y., Rodriguez, J., Lopatkin, A. J. & You, L. Growth-stage-

dependent regulation of conjugation. *AIChE J.* **66**, e16848 (2020).

30. Turner, P. E. *et al.* Antibiotic resistance correlates with transmission in plasmid

evolution. *Evolution* **68**, 3368–3380 (2014).

31. Turner, P. E., Cooper, V. S. & Lenski, R. E. Tradeoff Between Horizontal and

Vertical Modes of Transmission in Bacterial Plasmids. *Evolution* **52**, 315–329

(1998).

32. Kottara, A., Hall, J. P. J., Harrison, E. & Brockhurst, M. A. Multi-host environments

select for host-generalist conjugative plasmids. *BMC Evol. Biol.* **16**, 70 (2016).

33. Gordon, D. M. Rate of plasmid transfer among Escherichia coli strains isolated from

natural populations. *Microbiology,* **138**, 17–21 (1992).

34. Philipsen, K. R., Christiansen, L. E., Hasman, H. & Madsen, H. Modelling conjugation with stochastic differential equations. *J. Theor. Biol.* **263**, 134–142 (2010).

35. Dimitriu, T., Matthews, A. & Buckling, A. *Increased copy number couples the evolution of plasmid horizontal transmission and antibiotic resistance*. http://biorxiv.org/lookup/doi/10.1101/2020.08.12.248336 (2020) doi:10.1101/2020.08.12.248336.

36. Szpirer, C., Top, E., Couturier, M. & Mergeay, M. Retrotransfer or gene capture: a feature of conjugative plasmids, with ecological and evolutionary significance. *Microbiology,* **145**, 3321–3329 (1999).

37. Kneis, D., Hiltunen, T. & Heß, S. A high-throughput approach to the culture-based estimation of plasmid transfer rates. *Plasmid* **101**, 28–34 (2019).

38. Keller, P. & Antal, T. Mutant number distribution in an exponentially growing population. *J. Stat. Mech. Theory Exp.* **2015**, P01011 (2015).