The female pheromone (*Z*)-4-undecenal mediates flight attraction and courtship in *Drosophila melanogaster*

Felipe Borrero-Echeverry 1,2, Marit Solum 1, Federica Trona 1, Erika A. Wallin 3, Marie Bengtsson 1, Peter Witzgall 1, Sebastien Lebreton 1,4

1 Chemical Ecology Unit, Department of Plant Protection Biology, Swedish University of Agricultural Sciences, 230 53 Alnarp, Sweden

2 Corporación Colombiana de Investgación Agropecuaria, Agrosavia, Mosquera, Colombia

3 Department of Chemical Engineering, Mid Sweden University, Holmgatan 10, 85170 Sundsvall, Sweden

4 IRSEA, Research Institute for Semiochemistry and Applied Ethology, Quartier Salignan, 84400 Apt, France

Felipe Borrero-Echeverry - fborrero@agrosavia.co
Marit Solum - marit.solum@slu.se
Federica Trona - federica.trona@scuole.provincia.tn.it
Erika A. Wallin - erika.wallin@miun.se
Marie Bengtsson - marie.bengtsson@slu.se
Peter Witzgall - peter.witzgall@slu.se
Sebastien Lebreton - s.lebreton@group-irsea.com

Author for correspondence: peter.witzgall@slu.se

Summary

[27 words]

The female-produced, species-specific volatile pheromone of *D. melanogaster* attracts both sexes from a distance, alone and in concert with food odorants, and elicits courtship in experienced males.

Abstract

[249 words]

Specific mate communication and recognition underlies reproduction and hence speciation. Mate communication evolves during adaptation to ecological niches and makes use of social signals and habitat cues.

Our study provides new insights in *Drosophila melanogaster* premating olfactory communication, showing that female pheromone (*Z*)-4-undecenal (*Z*4-11Al) and male pheromone cVA interact with food odour in a sex-specific manner. Furthermore, *Z*4-11Al, which mediates upwind flight attraction in both sexes, also elicits courtship in experienced males.

Twin variants of the olfactory receptor Or69a are co-expressed in the same olfactory sensory neurons, and feed into the same glomerulus in the antennal lobe. *Z*4-11Al is perceived via Or69aB, while the food odorant (*R*)-linalool is a main ligand for the other variant, Or69aA. That *Z*4-11Al mediates courtship in experienced males, not (*R*)-linalool, is probably due to courtship learning. Behavioural discrimination is reflected by calcium imaging of the antennal lobe, showing distinct glomerular activation patterns by these two compounds.

Male sex pheromone cVA is known to affect male and female courtship at close range, but does not elicit upwind flight attraction as a single compound, in to contrast to *Z*4-11Al. A blend of cVA and the food odour vinegar attracted females, while a blend of female pheromone *Z*4-11Al and vinegar attracted males instead.

Sex-specific upwind flight attraction to blends of food volatiles and male and female pheromone, respectively, adds a new element to *Drosophila* olfactory premating communication and is an unambiguous paradigm for identifying the behaviourally active components, towards a more complete concept of food-pheromone odour objects.

Keywords

Specific Mate Recognition • Olfaction • Neuroethology • (*Z*)-11-octadecenyl acetate • cis-vaccenyl acetate • (*Z*)-4-undecenal

Introduction

Structure and function of the olfactory system have been studied in *Drosophila* like in no other insect, from peripheral odorant perception to central pathways generating

59	behavioural output. And, a particular emphasis of this work has been placed on
60	courtship, inspired by a characteristic behavioural response (Depetris-Chauvin et al.
61	2015, Kohl et al. 2015, Auer and Benton 2016, Grabe et al. 2016, Bates et al. 2020).
62	The combined molecular and physiological know-how enables even investigations of the
63	evolutionary development of olfactory channels in response to pheromones (Khallaf et
64	al. 2020 a,b) and host odorants (Dekker et al. 2006, Markow 2019, Auer et al. 2020),
65	across *Drosophila* phylogenies.

66	What is still unclear is whether olfactory tuning and response to pheromones and host
67	or food odours evolve independently. And, although odorants enable evaluation and
68	recognition of the emitter or source from a distance, little attention has been paid to
69	the role of species-specific sex pheromones in distant mate recognition and flight
70	attraction, prior to courtship enactment.

71	Drosophila cuticular hydrocarbon (CHC) profiles are sexually dimorphic, they vary and
72	rapidly diverge between species and therefore contribute to sexual, reproductive
73	isolation (Howard et al. 2003, Legendre et al. 2008, Alves et al. 2010, De Oliveira et al.
74	2011, Davis et al. 2020). Naturally, this congruently applies to their volatile oxidation
75	products, particularly monounsaturated aldehydes, which derive from diunsaturated
76	CHCs.

77	A CHC produced by female flies, (Z,Z)-7,11-heptacosadiene (7,11-HD), has been shown
78	to afford reproductive isolation between *D. melanogaster* and its sibling species *D.*
79	*simulans*. It is perceived by gustatory receptors owing to its low volatility (Billeter et al.
80	2009, Thistle et al. 2012, Toda et al. 2012, Billeter and Wolfner 2018, Seeholzer et al.
81	2018, Sato and Yamamoto 2020). Oxidation of 7,11-HD gives rise to the volatile
82	pheromone (Z)-4-undecenal (Z4-11Al) which is perceived by one of the two variants of
83	the olfactory receptor DmelOr69a (Or69a) and elicits flight attraction in males and
84	females (Lebreton et al. 2017). The Or69a gene encodes two or more proteins, in a
85	range of *Drosophila* species (Robertson et al. 2003, Conceicao and Aguade 2008). In *D.*
86	*melanogaster*, the two isoforms of Or69a, Or69aA and Or69aB, are tuned to food
87	odorants, including (R)-linalool and to the female pheromone Z4-11Al , respectively.
88	They are expressed in the same olfactory sensory neurons (OSNs), and afford
89	combined input of sex and food stimuli (Lebreton et al. 2017).

90	In comparison, *cis*-vaccenyl acetate (cVA) is a pheromone produced by males that
91	increases female receptivity and reduces male attraction to recently mated females
92	(Bartelt et al. 1985, Ejima et al. 2007, Kurtovic et al. 2007, Keleman et al. 2012,
93	Lebreton et al. 2014). However, cVA cannot account for specific mate communication
94	since it is shared by many other *Drosophila* species (El-Sayed 2020). The processing of
95	cVA stimuli in sexually dimorphic neural pathways has been mapped from antennal
96	input to third-order neurons in the lateral horn, where gender-specific courtship
97	behaviour is generated (Kohl et al. 2013, Clowney et al. 2015).

cVA activates the sexually dimorphic fruitless (*Fru*) circuitry that controls male

courtship (Manoli et al. 2005, Stockinger et al. 2005, Billeter et al. 2006, Cachero et al.

2010), whereas *Z*4-11Al is not part of this circuit. Since *Z*4-11Al is produced by

females, we investigated its effect on flight attraction and courtship, which are

successive steps in male reproductive behaviour. An upwind flight assay confirms that

*Z*4-11Al elicits attraction in naive males (Lebreton et al. 2017), while it elicits courtship

only in experienced, previously mated males.

Remarkably, males distinguished between *Z*4-11Al and the food odorant (*R*)-linalool,

which did not elicit courtship. Both compounds are perceived via the two isoforms of

Or69a, that are co-expressed in the same OSNs, while functional imaging of the

antennal lobe (AL) showed nonetheless different activation patterns for *Z*4-11Al and

(*R*)-linalool.

Food odours act as aphrodisiacs and promote courtship in *Drosophila* (Grosjean et al.

2011, Gorter et al. 2016, Ando et al. 2020). Vinegar has traditionally been used as a

source of food odorants in fruit fly research, and has a prominent synergistic effect on

cVA perception and attraction (Lebreton et al. 2015, Das et al. 2017, Cazale-Debat et

al. 2019). Vinegar headspace contains several active compounds (Becher et al. 2010)

but greatly varies in composition between makes and types (Callejón et al. 2009,

Chinnici et al. 2009). Over-ripe fermenting fruit, on the other hand, is a feeding,

mating, and oviposition site for fruit flies. Yeasts growing on fruit release a very rich

volatome, that accounts for strong fly attraction (Becher et al. 2012, Buser et al. 2014,

Christiaens et al. 2014, Ljunggren et al. 2019).

We compared the effect of vinegar and yeast headspace on sex attraction to

pheromone. Blends of vinegar odour and male or female pheromone produced sex-

specific flight attraction of females and males, respectively. In comparison, both sexes

respond to blends of yeast and cVA or *Z*4-11Al.

Materials and methods

Insects and chemicals

Flies were reared on a standard sugar-yeast-cornmeal diet at room temperature (19 to

22°C) under a 16:8-h L:D photoperiod. Newly eclosing flies were anesthetized with CO_2

and sexed under a microscope. Virgin flies were identified by the presence of

meconium, and were kept together with flies of the same sex. Mated males were kept

together with females, and were separated 2 d before experiments. Flies were kept in

30-mL Plexiglas vials with fresh food, and were six d old when used for experiments.

Isomeric purity of (*Z*)-4-undecenal (Z4-11Al) and (*E*)-4-undecenal (E4-11Al) was

98.6% and 97.8%, repectively, according to gas chromatography coupled to mass

spectrometry (6890 GC and 5975 MS, Agilent Technologies, Santa Clara, CA, USA).

Chemical purity of the synthetic aldehydes was >99.9%, and of the other test

chemicals >97%. Heptane (redistilled; Merck, Darmstadt, Germany) was used as

solvent.

Functional imaging

The head capsule was opened by incising the cuticle between the antennae and the

eyes. With the brain immersed in Ringer's saline, the ALs were exposed by removing

muscle tissue, glands and trachea. In vivo recordings of illuminated preparations were

processed using custom software (Strutz et al. 2014).

We used a Till Photonic imaging system with an upright Olympus microscope (BX51WI)

and a 20x Olympus objective (XLUM Plan FL 20x/0.95W). A Polychrome V provided light

excitation (475 nm), which was then filtered (excitation: SP500, dicroic: DCLP490,

emission LP515) and captured by a CCD camera (Sensicam QE, PCO AG) with

symmetrical binning. For each measurement, a series of 300 frames was taken (1 Hz).

Data were analyzed using IDL (Research Systems Inc., Boulder, CO, USA).

A 3-D map of the fruit fly AL (Grabe et al. 2015) served to link the active area to

individual glomeruli. All experimental flies contained the calcium dependent fluorescent

sensor G-CaMP 3.0 (Nakai et al. 2001) together with a promoter Gal4 insertion to direct

expression of the calcium sensor to specific neuron populations. Stimulus-evoked

fluorescence in these flies arises from the population of labeled neurons that are

sensitive to the specific odour.

We tested the physiological responses in input neurons, i.e. the axonal terminals of

OSNs in the AL. Mass labelling of olfactory sensory neurons (OSNs) was achieved using

the transgenic line Orco-GAL4 that drives expression in at least 60% of all OSNs

(Larsson et al. 2004). Additionally, for mass labelling of PNs, the transgenic line

GH146-GAL4, expressing G-CaMP in 83 out of the 150 PNs in each AL (Vosshall and

Stocker 2007) has been used. Fly lines were obtained from the Bloomington *Drosophila*

Stock Center (Indiana University, Bloomington, IN, USA). Transgenic flies have been

generated using standard procedures.

Courtship assay

Wild-type flies for courtship experiments were the *D. melanogaster* strains Dalby-HL

(Ruebenbauer et al. 2008) and Canton-S, and and the sister species *D. simulans*.

Orco-Gal4/uas-Or69aRNAi flies, and the Orco-Gal4 and uas-Or69aRNAi (VDCR, Vienna)

parental lines were used to confirm the effect of *Z*4-11Al on courtship behavior.

Canton-S served as comparison for knockouts of the same background. We resorted to

the Orco-Gal4 line, since males of the Or69a-Gal4 line did not court. Because of the

broad expression of the Or69aRNAi, we cannot exclude that other Ors involved in
courtship in experienced males are also affected.

Courtship experiments were done on a light box (37x28x2.5 cm; color temperature
5.000 ± 5% Kelvin; Kaiser Slimlite 2422, Kaiser Fototechnik GmbH & Co. KG, Buchen,
Germany). The courtship arena consisted of three glass plates (17x13x0.5 cm) placed
on top of each other. Twelve circular holes (ø 3 cm) were cut in the middle plate to
form 12 circular single-pair mating chambers. All glassware was heated to 350°C for 8
h before use. Tests were done between 2 and 5 h after onset of scotophase (16:8 L:D
photoperiod). Target flies, unmated males or females, were anesthetized on ice and
decapitated before experiments. Test flies were either mated or unmated males. A
single target fly was added to each mating chamber, and 1 µl heptane (control) or 1 µl
heptane containing 1 ng Z4-11Al was applied onto its abdomen. After solvent
evaporation (2 min), a single live test fly was introduced into each mating chamber and
observed during 20 min. Males were scored when showing vigorous courtship
behaviour, including wing vibration, licking, and attempted copulation. Treatments (n =
80) and controls (n = 80) were conducted simultaneously, and each fly was tested
once.

Food odour collection
Brewer's yeast, *Saccharomyces cerevisiae* (strain S288C), was grown in minimal media
(Merico et al. 2007) during 20 h in a shaking incubator at 25°C and 260 RPM. 50 mL
yeast broth or white vine vinegar (7.1%, Zeta, Sweden) were filled into a wash bottle
and charcoal filtered air was bubbled through the bottle at a rate of 200 mL/min.
Headspace was trapped in 4x40-mm glass tubes holding 300 mg of Porapak Q 50/30
(Waters Corporation, Milford, Ma, USA) during 2 h (Becher et al. 2012). These air filters
were eluted with 1.2 mL redistilled ethanol (Labscan), the eluate was sprayed during 2
h of wind tunnel experiments.

Wind tunnel assay
Upwind flight attraction was observed in a glass wind tunnel (30x30x100 cm). The wind
tunnel was lit diffusely from above, at 13 lux, temperature ranged from 20°C to 24°C,
relative humidity from 38% to 48% and charcoal filtered air (Camfil AB, Malmö,
Sweden), at a velocity of 0.25 m/s, was produced by a fan (Fischbach GmbH,
Neunkirchen, Germany). Yeast headspace and odour blends were delivered from the
centre of the upwind end of the wind tunnel via a piezo-electric microsprayer (Becher et
al. 2010).

Forty flies were flown individually to each treatment. Flies were scored when flying
upwind over 80 cm in the wind tunnel centre, from the release cage towards the odour
source. Unmated, fed, 3-d-old males and females were flown towards yeast and vinegar
headspace alone and in blends with Z4-11Al sprayed at 10 ng/min (Lebreton et al.

2017) and 300 ng/min *cis*-vaccenyl acetate (cVA) (Lebreton et al. 2015). Isomeric

purity of the synthetic compounds, according to gas chromatography, was 98.6%. and

97.3%, respectively.

Statistical analysis

Generalized linear models (GLM) with a Bernoulli binomial distribution were used to

analyse wind tunnel data. Landing at source and sex were used as the target effects.

Post-hoc Wald pairwise comparison tests were used to identify differences between

treatments. Courtship assays were analysed using chi-squared tests to compare

between treatments and their respective control. Statistical analysis was calculated with

R (R Core Team 2013) and SPSS Version 22 (IBM Corp.).

Results

Flight attraction to blends of pheromone and food odour

The male-produced pheromone cVA (Bartelt et al. 1985) is a core paradigm for

behavioural and neurophysiological studies in *Drosophila*. The volatile, female-produced

and species-specific pheromone *Z*4-11Al has been discovered more recently (Lebreton

et al. 2017). We therefore compared flight attraction to male pheromone cVA and

female pheromone *Z*4-11Al, in *D*. *melanogaster* males and females. Pheromones were

tested alone and in blends with odorants from vinegar and yeast, signalling food

sources (Figure 1).

Male pheromone cVA alone attracted only few flies, while a blend of cVA and vinegar

was highly attractive to females, not to males (Figure 1b). In comparison, both sexes

responded to a blend of cVA and yeast headspace.

Female pheromone *Z*4-11Al alone attracted males and females. Blending vinegar with

*Z*4-11Al reduced attraction in both sexes, but significantly more males than females

responded. And, a blend of yeast headspace and *Z*4-11Al attracted as many flies as *Z*4-

11Al alone (Figure 1c).

Yeast headspace by itself attracted more males and females than vinegar headspace

(Figure 1a). Females attraction to yeast headspace (Figure 1a) and to the blends of

yeast headspace and cVA and *Z*4-11Al, respectively, was not significantly different

(Figures 1b,c).

Taken together, only few flies responded to the male pheromone cVA, while the female

pheromone *Z*4-11Al strongly attracted both sexes. Blending vinegar with cVA and *Z*4-

11Al, respectively, produced an inverse response pattern in male and female flies. Our

results show further that vinegar, a frequently used food odorant source for *Drosophila*,
is an inferior attractant in comparison with yeast aroma (Figure 1).

Male courtship in response to *Z*4-11Al

Male-produced cVA mediates aggregation, suppresses male-to-male courtship, and
males learn to avoid mated females tainted with cVA (Ejima et al. 2007, Kurtovic et al.
2007, Keleman et al. 2012). Since female-produced *Z*4-11Al elicits male flight
attraction (Figure 1c), we asked whether *Z*4-11Al also has an effect on courtship,
following landing at females. Males were tested with decapitated target flies, laced with
blank solvent or synthetic test chemicals.

Compared with solvent control, *Z*4-11Al elicited a significant response in experienced
males that had mated earlier, and not in naive, unmated males. Males scored showed
vigorous courtship including attempted copulation (Figure 2a). More experienced males
courted decapitated females than males, plausibly because cVA, present on the cuticula
of male flies, has an antagonistic effect on experienced males. That a few males
nonetheless courted decapitated males further confirms a role of *Z*4-11Al in male
courtship (Figure 2a). Tests with Canton-S test males corroborated results obtained
with males of the Dalby strain (Figure 2b).

Importantly, males also courted decapitated *D*. *simulans* target females, laced with
synthetic *Z*4-11Al (Figure 2b). This experiment rules out a contributing role of the
cuticular hydrocarbon pheromone 7,11-HD, which is specific for *D*. *melanogaster* and
not found in *D*. *simulans* females (Billeter et al. 2009), and underlines that *Z*4-11Al
elicits male courtship, in the absence of 7,11-HD.

On the other hand, *D*. *simulans* males did not respond in a significant manner to Dalby
females painted with *Z*4-11Al (Figure 2b). This was expected, since *D*. *simulans* males
are not attracted to *Z*4-11Al (Lebreton et al. 2017) and since *D*. *simulans* females do
not produce 7,11HD, which is the precursor for *Z*4-11Al (Billeter et al. 2009, Lebreton
et al. 2017).

Moreover, we used a RNAi fly line to corroborate that Or69a encodes *Z*4-11Al-mediated
courtship. Significantly fewer males responded when olfactory sensory neurons (OSNs)
expressing Or69a were disrupted (Figure 2c).

Finally, we tested three non-pheromonal chemicals: *E*4-11Al, the geometric isomer of
*Z*4-11Al, which is perceived via the Or69aB variant; (*R*)-linalool, a food compound,
which is the main ligand of the other variant, Or69aA; and ethyl butyrate, which is a
ligand for Or22a (DM2 glomerulus in the AL), for Or43b (VM2 glomerulus) and also for
Or85a (DM5 glomerulus; see also Figure 3). None of these compounds elicited a
significant courtship response in experienced males (Figure 2d).

Activation of antennal lobe glomeruli in response to *Z*4-11Al

The D glomerulus in the antennal lobe (AL) collects input from ab9A olfactory sensory neurons (OSNs) that co-express Or69aA and Or69aB, the two isoforms of Or69a (Couto et al. 2005). Single sensilllum recordings (SSR) have further shown that the food odour (*R*)-linalool and female pheromone *Z*4-11Al are the respective key ligands for Or69aA and Or69aB (Lebreton et al. 2017).

We used calcium imaging of the AL to first confirm the role of the Or69a channel including the D glomerulus in the perception of *Z*4-11Al. Moreover, we compared the AL response to the key ligands (*R*)-linalool and *Z*4-11Al. Both compounds elicit upwind flight (Lebreton et al. 2017), but only *Z*4-11Al participates in male courtship, which goes to show that the flies discriminate between these two compounds (Figure 2).

In males and females, *Z*4-11Al activated the D glomerulus, as expected, but also DM2 and VM2 (Figure 3a). These broadly-tuned glomeruli have also been shown to respond to a mix of cVA and vinegar (Lebreton et al. 2015). However, neither ab3A OSNs (expressing Or22a, connecting to DM2), nor ab8A (expressing Or43b, connecting to VM2; Grabe et al. 2015, Münch and Galizia 2016) responded to *Z*4-11Al, according to SSR (Lebreton et al. 2017). Simultaneous activation of DM2 and VM2 glomeruli by *Z*4-11Al might accordingly reflect interglomerular communication in the AL, via local neurons (Wilson 2013).

(*R*)-linalool, on the other hand, is also a ligand for ab7A OSNs expressing Or98a, which accounts for additional activation of the VM5v glomerulus (Couto et al. 2005, Münch and Galizia 2016). In comparison, activation of DM2 and VM2 glomeruli by (*R*)-linalool was much lower compared to *Z*4-11Al (Figure 3a).

Activation of the DM2 and VM2 glomeruli by *Z*4-11Al was substantiated by imaging Or22a-Gal4 males and Or43b-Gal4 males, respectively. Ethyl butyrate is a diagnostic stimulus for Or22a, Or43b and Or85a (DM2, VM2 and DM5 glomeruli) (Figure 3b; Grabe et al. 2015, Münch and Galizia 2016).

Imaging of the glomerular activation pattern of projection neurons (PNs), connecting the AL to the lateral horn (LH), where behavioural responses are generated, corroborated activity of *Z*4-11Al in the D and DM2 glomeruli (Figure 3c). A possible response of VM5v to linalool does not show, since GH146-Gal4 does not label this glomerulus (Grabe et al. 2015).

Discussion

Volatile chemicals that are cognate ligands for odorant receptors (ORs) are selectively perceived against noisy backgrounds and they capacitate behavioural decisions at a

distance, since they deliver reliable information about the source, including identity and
physiological state of the emitter, in the case of pheromones.

The female pheromone *Z*4-11Al attracts naive males and females by flight (Figure 1).
At close range, *Z*4-11Al also elicits courtship, but only in experienced, previously mated
males (Figure 2). A response in naive males is not expected, since neurons mediating
perception of *Z*4-11Al are not part of the sexually dimorphic fruitless circuitry that
elicits innate male courtship (Manoli et al. 2005, Stockinger et al. 2005).

Importantly, males respond also to decapitated *D. simulans* females laced with *Z*4-
11Al. This shows that a response to *Z*4-11Al does not require presence of the close-
range CHC pheromone 7,11-HD, since 7,11-HD is not produced by *D. simulans* (Billeter
et al. 2009, Lebreton et al. 2017, Billeter and Wolfner 2018, Sato and Yamamoto
2020).

Courtship discrimination between Or69a ligands

Female pheromone Z4-11Al and the food odorant (*R*)-linalool are main ligands for
Or69aB and Or69aA, which are co-expressed in ab9A OSNs, branching to the D
glomerulus in the AL (Couto et al. 2005, Lebreton et al. 2017).

Whereas both compounds elicit innate flight attraction (Lebreton et al. 2017), males
distinguish between Z4-11Al and (*R*)-linalool during courtship (Figure 2a,d), probably
because males learn to associate Z4-11Al with females during mating. Courtship
learning confirms production of Z4-11Al by females. The question arises whether males
would even learn to associate (*R*)-linalool or other food-related compounds with
females.

A differential response to Z4-11Al and (*R*)-linalool is reflected by their respective
activation patterns in the AL (Figure 3a). (*R*)-linalool is also a ligand for Or98a, which
accounts for stimulation of the VM5v glomerulus (Couto et al. 2005, Münch and Galizia
2016)

In contrast, direct activation of DM2 and VM2 by Z4-11Al is not supported by SSR data
(Lebreton et al. 2017) and lateral activation by local AL neurons is a possible explanation,
instead (Wilson 2013). DM2 and VM2 are respond broadly, also to blends of vinegar and cVA
(Lebreton et al. 2015).

Finally, we do not know whether the olfactory input of Or69aA and Or69aB through ab9A is
entirely equivalent. OSN spiking trains generated by Or69aA and Or69aB may deliver non-
congruent messages that may be distinguishable at the AL level. This is reminiscent of Or85e
and Or33c which are coexpressed in ab3A OSNs, where it is yet unclear whether different
response profiles and spiking patterns enable behavioural discrimination between the
respective Or ligands (Goldman et al. 2005).

Roles of cVA and Z4-11Al in upwind flight and courtship

Male courtship in *D. melanogaster* has been anatomically and physiologically dissected at a neuronal circuit level, with a particular emphasis on the sex-dimorphic behavioural effect of cVA. A central node is the sexually dimorphic P1 interneuron cluster that integrates input from all sensory modalities to activate male courtship, and that is regulated by social experience. Male-produced pheromone cVA provides antagonistic olfactory input to the P1 node, preventing courtship of recently mated females perfumed with male cVA (Manoli et al. 2005, Stockinger et al. 2005, Kimura et al. 2008, Ruta et al. 2010, Kohl et al. 2013, Yamamoto and Koganezawa 2013, Clowney et al. 2015, Kohl et al. 2015, Auer & Benton 2016).

cVA does not elicit long-range flight by itself, only in combination with fly food (Figure 1; Bartelt et al. 1985, Lebreton et al. 2015) and is shared by many other *Drosophila*, including the sibling species *D. simulans* (Schaner et al. 1987, El-Sayed 2020). Females, on the other hand, produce the species-specific CHC 7,11-HD that feeds gustatory input into P1 to activate male courtship and to achieve reproductive isolation towards the sibling species *D. simulans* (Billeter et al. 2009, Thistle et al. 2012, Toda et al. 2012, Seeholzer et al. 2018). What is yet unclear is whether P1 also receives excitatory olfactory input.

Does Z4-11Al feed into P1? Z4-11Al encodes species-specificity, mediates long-range attraction (Figure 1; Lebreton et al. 2017) and contributes to courtship, independently of 7,11-HD (Figure 2a,b). Z4-11Al is perceived prior to and during upwind flight, and since females release Z4-11Al, it follows that stimulation is sustained at close-range. Courtship and upwind flight are not isolated episodes, they are inextricably interconnected parts of a behavioural sequence that culminates in copulation.

A behavioural paradigm for bioactive odorant identification

An unambigious, stereotypical behavioural response is substrate for investigations of the genetic basis, neural circuitry and physiology of mate communication in *D. melanogaster*. We here show that upwind flight attraction is a behavioural paradigm that is suitable to extend investigations of reproductive behaviour to include long-range communication, and even decision-making upon odorant stimulation.

Comparison of pheromone blend attraction with vinegar and yeast shows that yeast overrides differences between pheromones (Figure 1). Vinegar, derived from acetic acid bacteria fermentation (Lynch et al. 2019), is a widely used standard food attractant for *D. melanogaster*, although live yeast aroma is much richer in composition (Callejón et al. 2009, Chinnici et al. 2009, Becher et al. 2010, Ljunggren et al. 2019). Yeast growing on ripe fruit is also a biologically more relevant attractant, since it attracts flies for oviposition and since yeast is a sufficient substrate for larval development (Becher et al. 2012, Grangeteau et al. 2018, Quan and Eisen 2018, Murgier et al. 2019).

Reproductive behaviour in *D*. *melanogaster* is mediated by an ensemble of pheromones
and food odorants, emanating from food substrates and from aggregating or mating
flies. Our knowledge of *D*. *melanogaster* pheromone chemistry is fairly comprehensive,
compared to behaviourally salient food odorants, owing to the extraordinary complexity
of microbial headspace.

Flies recognize food sources despite very considerable, inherent variation in the
bouquets emanating from fermenting fruit, according to fruit substrates, microbial
community composition and fermentation state. Towards an understanding of how
neural representations of such complex and variable odours are generated (Endo et al.
2020) and how habitat cues are integrated with pheromonal signals we next need to
identify the key odorants in microbial food headspace. An upwind flight bioassay is
suitable to describe the key bioactive chemical constitutents that encode reliable
recognition and evaluation of food odour objects and trigger the decision to engage in
orientation flights.

Acknowledgements

401 We thank two anonymous reviewers for constructive criticism.

Competing interests

403 No competing interests declared.

Funding

405 This study was supported by the Colombian Corporation for Agricultural Research
406 (Corpoica), the Colombian Administrative Department of Science, Technology, and
407 Innovation (Colciencias), the Linnaeus environment "Insect Chemical Ecology, Ethology
408 and Evolution" IC-E3 (Formas, SLU, Sweden), and the Faculty of Landscape
409 Architecture, Horticulture, and Crop Production Science (SLU, Alnarp, Sweden).

Data availability

411 Data will be made avaliable at Dryad.

References

[70]

Alves, H., Rouault, J. D., Kondoh, Y., Nakano, Y., Yamamoto, D., Kim, Y. K., & Jallon, J. M. (2010). Evolution of cuticular hydrocarbons of Hawaiian Drosophilidae. Behavior Genetics, 40(5), 694-705. (doi:10.1007/s10519-010-9364-y)

Ando, Y., Yoshimizu, T., & Matsuo, T. (2020). Food availability reverses the effect of hunger state on copulation rate in Drosophila prolongata females. Animal Behaviour, 166, 51-59. (doi:10.1016/j.anbehav.2020.06.003)

Auer, T. O., & Benton, R. (2016). Sexual circuitry in Drosophila. Current Opinion in Neurobiology 38:18-26. (doi:10.1016/j.conb.2016.01.004)

Auer TO, Khallaf MA, Silbering AF, Zappia G, Ellis K, Alvarez-Ocana R, Arguello JR, Hansson BS, Jefferis GS, Caron SJ, Knaden M, Benton R. 2020. Olfactory receptor and circuit evolution promote host specialization. Nature 579:402-408. (doi:10.1038/s41586-020-2073-7)

Bartelt, R.J., Schaner, A.M. & Jackson, L.L. 1985 cis-Vaccenyl acetate as an aggregation pheromone in *Drosophila melanogaster*. *J. Chem. Ecol.* 11, 1747-1756. (doi:10.1007/BF01012124)

Bates AS, Schlegel P, Roberts RJV, Drummond N, Tamimi IFM, Turnbull R, Zhao X, Marin EC, Popovici PD, Dhawan S, Jamasb A, Javier A, Capdevila LS, Li F, Rubin GM, Waddell S, Bock DD, Costa M, Jefferis GSXE. 2020. Complete connectomic reconstruction of olfactory projection neurons in the fly brain. Current Biology 30, 3183-3199. (doi.org/10.1016/j.cub.2020.06.042)

Becher PG, Bengtsson M, Hansson BS, Witzgall P. 2010. Flying the fly: long-range flight behavior of *Drosophila melanogaster* to attractive odors. J chem Ecol 36:599-607 (doi: 10.1007/s10886-010-9794-2)

Becher, P.G., Flick, G., Rozpędowska, E., Schmidt, A., Hagman, A., Lebreton, S., Larsson, M.C., Hansson, B.S., Piškur, J., Witzgall, P., et al. 2012 Yeast, not fruit volatiles mediate Drosophila melanogaster attraction, oviposition and development. Funct. Ecol. 26, 822-828. (doi:10.1111/j.1365-2435.2012.02006.x).

Billeter, J. C., Rideout, E. J., Dornan, A. J, Goodwin SF. Control of male sexual behavior in Drosophila by the sex determination pathway. Curr. Biol. 16, R766–R776 (2006).

Billeter JC, Atallah J, Krupp JJ, Millar JG, Levine JD. 2009. Specialized cells tag sexual and species identity in Drosophila melanogaster. Nature 461:987-U250 (doi:10.1038/nature08495)

Billeter, J. C., & Wolfner, M. F. (2018). Chemical cues that guide female reproduction in Drosophila melanogaster. Journal of chemical Ecology 44:750-769. (doi:10.1007/s10886-018-0947-z)

Buser, C.C., Newcomb, R.D., Gaskett, A.C. & Goddard, M.R. 2014 Niche construction initiates the evolution of mutualistic interactions. Ecol. Lett. 17, 1257-1264. (doi:10.1111/ele.12331).

Cachero, S., Ostrovsky, A. D., Yu, J. Y., Dickson, B. J. & Jefferis, G. S. X. E. Sexual dimorphism in the fly brain. Curr. Biol. 20, 1589–1601 (2010).

Callejón, R.M., Tesfaye, W., Torija, M.J., Mas, A., Troncoso, A.M. and Morales, M.L.,
2009. Volatile compounds in red wine vinegars obtained by submerged and surface
acetification in different woods. Food Chemistry, 113(4), 1252-1259.
(doi:10.1016/j.foodchem.2008.08.027)

Cazale-Debat, L., Houot, B., Farine, J. P., Everaerts, C., & Ferveur, J. F. (2019). Flying
Drosophila show sex-specific attraction to fly-labelled food. Scientific reports, 9(1),
1-13. (doi:10.1038/s41598-019-51351-1)

Chinnici, F., Duran Guerrero, E., Sonni, F., Natali, N., Natera Marin, R. and Riponi, C.,
2009. Gas Chromatography− Mass Spectrometry (GC− MS) characterization of
volatile compounds in quality vinegars with protected European geographical
indication. Journal of agricultural and food chemistry, 57(11), 4784-4792.
(doi:10.1021/jf804005w)

Christiaens, J.F., Franco, L.M., Cools, T.L., De Meester, L., Michiels, J., Wenseleers, T.,
Hassan, B.A., Yaksi, E. & Verstrepen, K.J. 2014 The fungal aroma gene ATF1
promotes dispersal of yeast cells through insect vectors. Cell Reports 9, 425-432.
(doi:10.1016/j.celrep.2014.09.009).

Clowney, E.J., Iguchi, S., Bussell, J.J., Scheer, E. & Ruta, V. 2015 Multimodal
Chemosensory Circuits Controlling Male Courtship in Drosophila. *Neuron* 87, 1036-
1049. (doi:10.1016/j.neuron.2015.07.025).

Conceicao IC, Aguade M. 2008. High incidence of interchromosomal transpositions in
the evolutionary history of a subset of Or genes in Drosophila. J Molec Evol.
66:325-32. (doi:10.1007/s00239-008-9071-y)

Couto A, Alenius M, Dickson BJ. 2005. Molecular, anatomical, and functional
organization of the *Drosophila* olfactory system. Current Biology 15:1535-1547
(doi:10.1016/j.cub.2005.07.034)

Das S, Trona F, Khallaf MA, Schuh E, Knaden M, Hansson BS, Sachse S. 2017. Electrical
synapses mediate synergism between pheromone and food odors in Drosophila
melanogaster. Proc Ntl Acad Sci USA 114:E9962-E9971.
(doi:10.1073/pnas.1712706114)

Davis, J. S., Pearcy, M. A., Yew, J. Y., & Moyle, L. C. (2020). A shift to shorter cuticular
hydrocarbons causes sexual isolation in the Drosophila americana species group.
bioRxiv (doi:10.1101/2020.09.07.286294)

De Oliveira, C. C., Manfrin, M. H., de M Sene, F., Jackson, L. L., & Etges, W. J. 2011.
Variations on a theme: diversification of cuticular hydrocarbons in a clade of
cactophilic Drosophila. BMC Evolutionary Biology 11:179. (doi:10.1186/1471-2148-
11-179)

Dekker, T., Ibba, I., Siju, K. P., Stensmyr, M. C., & Hansson, B. S. (2006). Olfactory
shifts parallel superspecialism for toxic fruit in Drosophila melanogaster sibling, D.
sechellia. Current Biology 16:101-109. (doi:10.1016/j.cub.2005.11.075)

Depetris-Chauvin, A., Galagovsky, D. & Grosjean, Y. 2015 Chemicals and
chemoreceptors: ecologically relevant signals driving behavior in Drosophila. *Front
Ecol Evol* **3**. (doi:10.3389/fevo.2015.00041).

Ejima A, Smith BPC, Lucas C, Van Naters WV, Miller CJ, Carlson JR, Levine JD, Griffith
LC. 2007. Generalization of courtship learning in Drosophila is mediated by cis-
vaccenyl acetate. Curr Biol 17:599-605. (doi:10.1016/j.cub.2007.01.053)

El-Sayed, A. M. (2020). The pherobase: database of pheromones and semiochemicals.
www.phero base.com

Endo K, Tsuchimoto Y, Kazama H. 2020. Synthesis of conserved odor object
representations in a random, divergent-convergent network. Neuron 108, 367–
381. (doi:10.1016/j.neuron.2020.07.029)

Goldman, A. L., van Naters, W. V. D. G., Lessing, D., Warr, C. G., & Carlson, J. R.
(2005). Coexpression of two functional odor receptors in one neuron. Neuron
45:661-666. (doi:10.1016/j.neuron.2005.01.025)

Gorter, J.A., Jagadeesh, S., Gahr, C., Boonekamp, J.J., Levine, J.D. and Billeter, J.C.,
2016. The nutritional and hedonic value of food modulate sexual receptivity in
Drosophila melanogaster females. Scientific reports, 6, 19441
(doi:10.1038/srep19441)

Grabe, V., Strutz, A., Baschwitz, A., Hansson, B.S. and Sachse, S., 2015. Digital in vivo
3D atlas of the antennal lobe of *Drosophila melanogaster*. Journal of Comparative
Neurology, 523(3), pp.530-544 (doi:10.1002/cne.23697).

Grabe, V., Baschwitz, A., Dweck, H.K., Lavista-Llanos, S., Hansson, B.S. and Sachse,
S., 2016. Elucidating the neuronal architecture of olfactory glomeruli in the
Drosophila antennal lobe. Cell Reports, 16(12), pp.3401-3413.
(doi:10.1016/j.celrep.2016.08.063)

Grangeteau, C., Yahou, F., Everaerts, C., Dupont, S., Farine, J.P., Beney, L. and
Ferveur, J.F. 2018. Yeast quality in juvenile diet affects Drosophila melanogaster
adult life traits. Scientific Reports 8, 1-9. (doi:10.1038/s41598-018-31561-9)

Grosjean Y, Rytz R, Farine JP, Abuin L, Cortot J, Jefferis GSXE, Benton R. 2011. An
olfactory receptor for food-derived odours promotes male courtship in Drosophila.
Nature 478:236-U123. (doi: 10.1038/nature10428)

Howard, R.W., Jackson, L.L., Banse, H. and Blows, M.W., 2003. Cuticular hydrocarbons
of Drosophila birchii and D. serrata: identification and role in mate choice in D.
serrata. Journal of Chemical Ecology 29(4):961-976.
(doi:10.1023/A:1022992002239)

Keleman, K., Vrontou, E., Krüttner, S., Jai, Y. Y., Kurtovic-Kozaric, A., & Dickson, B. J.
(2012). Dopamine neurons modulate pheromone responses in Drosophila courtship
learning. Nature 489:145-149. (doi:10.1038/nature11345)

Khallaf MA, Auer TO, Grabe V, Depetris-Chauvin A, Ammagarahalli B, Zhang D-D,
Lavista-Llanos S, Kaftan F, Weißflog J, Matzkin LM, Rollmann SM, Löfstedt C,
Svatos A, Dweck HKM, Sachse S, Benton R, Hansson BS, Knaden M. 2020a. Mate
discrimination among subspecies through a conserved olfactory pathway. Sci. Adv.
6: eaba5279. (doi:10.1126/sciadv.aba5279)

Khallaf, M.A., Cui, R., Weißflog, J., Svatoš, A., Dweck, H., Valenzano, D.R., Hansson,
B.S. and Knaden, M., 2020b. Large-scale characterization of sex pheromone
communication systems in Drosophila. bioRxiv, September 22
(doi:10.1101/2020.09.21.305854)

Kimura, K., Hachiya, T., Koganezawa, M., Tazawa, T., Yamamoto, D. 2008. Fruitless
and Doublesex coordinate to generate male-specific neurons that can courtship.
Neuron 59, 759-769. (doi:10.1038/nrn3567)

Kohl, J., Ostrovsky, Aaron D., Frechter, S. & Jefferis, Gregory S. 2013 A Bidirectional
Circuit Switch Reroutes Pheromone Signals in Male and Female Brains. *Cell* 155,
1610-1623. (doi:10.1016/j.cell.2013.11.025).

Kohl J, Huoviala P, Jefferis GSXE. 2015. Pheromone processing in Drosophila. Curr Op
Neurobiol 34:149-157. (doi:10.1016/j.conb.2015.06.009)

Kurtovic A, Widmer A, Dickson BJ. 2007. A single class of olfactory neurons mediates
behavioural responses to a Drosophila sex pheromone. Nature 446:542-546.
(doi:10.1038/nature05672).

Larsson, M. C., Domingos, A. I., Jones, W. D., Chiappe, M. E., Amrein, H., & Vosshall,
L. B. (2004). Or83b encodes a broadly expressed odorant receptor essential for
Drosophila olfaction. Neuron 43:703-714.(doi:10.1016/j.neuron.2004.08.019)

Lebreton S, Grabe V, Omondi AB, Ignell R, Becher PG, Hansson BS, Sachse S, Witzgall
P. 2014. Love makes smell blind: mating suppresses pheromone attraction in
Drosophila females via OR65a olfactory neurons. Sci Rep 4:7119
(doi:10.1038/srep07119)

Lebreton S, Trona S, Borrero-Echeverry F, Bilz F, Grabe V, Becher PG, Carlsson MA,
Nässel DR, Hansson BS, Sachse S, Witzgall P. 2015. Feeding regulates sex
pheromone attraction and courtship in *Drosophila* females. Sci Rep 5:13132
(doi:10.1038/srep13132)

Lebreton, S., Borrero-Echeverry, F., Gonzalez, F., Solum, M., Wallin, E.A., Hedenström,
E., Hansson, B.S., Gustavsson, A.-L., Bengtsson, M., Birgersson, G., et al. 2017 A
Drosophila female pheromone elicits species-specific long-range attraction via an
olfactory channel with dual specificity for sex and food. BMC Biol. 15, 88.
(doi:10.1186/s12915-017-0427-x).

Legendre, A., Miao, X. X., Da Lage, J. L., & Wicker-Thomas, C. (2008). Evolution of a
desaturase involved in female pheromonal cuticular hydrocarbon biosynthesis and
courtship behavior in *Drosophila*. Insect biochemistry and molecular biology, 38(2),
244-255. (doi:10.1016/j.ibmb.2007.11.005)

Ljunggren J, Borrero-Echeverry F, Chakraborty A, Lindblom TU, Hedenström E, Karlsson
M, Witzgall P, Bengtsson M. 2019. Yeast volatomes differentially effect larval
feeding in an insect herbivore. Appl Environm Microbiol 85: e01761-19
(doi:10.1128/AEM.01761-19)

Lynch, K.M., Zannini, E., Wilkinson, S., Daenen, L. and Arendt, E.K., 2019. Physiology
of acetic acid bacteria and their role in vinegar and fermented beverages.
Comprehensive Reviews in Food Science and Food Safety, 18, 587-625.
(doi:10.1111/1541-4337.12440)

Manoli, D. S., Foss, M., Villella, A., Taylor, B. J., Hall, J. C., & Baker, B. S. 2005. Male-
specific fruitless specifies the neural substrates of Drosophila courtship behaviour.
Nature 436:395-400. (doi:10.1038/nature03859)

Markow, T. A. (2019). Host use and host shifts in *Drosophila*. Current Opinion in Insect
Science 31:139-145.(doi:10.1016/j.cois.2019.01.006)

Merico, A., Sulo, P., Piskur, J. & Compagno, C. (2007) Fermentative lifestyle in yeasts
belonging to the Saccharomyces complex. FEBS Journal, 274, 976–989.
(doi:10.1111/j.1742-4658.2007.05645.x)

Münch D, Galizia CG. 2016. DoOR 2.0 - Comprehensive mapping of *Drosophila*
melanogaster odorant responses. Scientific Reports 6:21841
(doi:10.1038/srep21841)

Murgier, J., Everaerts, C., Farine, J.P. and Ferveur, J.F., 2019. Live yeast in juvenile
diet induces species-specific effects on Drosophila adult behaviour and fitness.
Scientific Reports 9,1-12. (doi:10.1038/s41598-019-45140-z)

Nakai, J., Ohkura, M., & Imoto, K. (2001). A high signal-to-noise Ca2+ probe composed
of a single green fluorescent protein. Nature Biotechnology 19:137-
141.(doi:10.1038/84397)

Quan AS, Eisen MB (2018) The ecology of the *Drosophila*-yeast mutualism in wineries.
PLoS One 13(5): e0196440. (doi:10.1371/journal.pone.0196440)

R Core Team. 2013 R: A Language and Environment for Statistical Computing. Vienna,
Austria, R Foundation for Statistical Computing.

Robertson HM, Warr CG, Carlson JR. 2003. Molecular evolution of the insect
chemoreceptor gene superfamily in Drosophila melanogaster. Proc Natl Acad Sci
USA 100:14537-42. (doi:10.1073/pnas.233584710)

Ruebenbauer, A., Schlyter, F., Hansson, B.S., Löfstedt, C., Larsson, M.C. 2008. Genetic
variability and robustness of host odor preference in *Drosophila melanogaster*.
Curr. Biol. 18: 1438-1443. (doi:10.1016/j.cub.2008.08.062)

Ruta V, Datta SR, Vasconcelos ML, Freeland J, Looger LL, Axel R. 2010. A dimorphic
pheromone circuit in Drosophila from sensory input to descending output. Nature
2010, 468:686-690. (doi:10.1038/nature09554)

Sato K and Yamamoto D (2020) Contact-chemosensory evolution underlying
reproductive isolation in Drosophila species. Front. Behav. Neurosci. 14:597428.
(10.3389/fnbeh.2020.597428)

Schaner, A.M., Bartelt, R.J. & Jackson, L.L. 1987. (Z)-11-octadecenyl acetate, an
aggregation pheromone inDrosophila simulans . J Chem Ecol 13, 1777–1786
(doi:10.1007/BF00980218)

Seeholzer, L. F., Seppo, M., Stern, D. L., & Ruta, V. (2018). Evolution of a central
neural circuit underlies Drosophila mate preferences. Nature, 559(7715), 564-569.
(doi:10.1038/s41586-018-0322-9)

Stockinger P, Kvitsiani D, Rotkopf S, Tirian L, Dickson BJ. 2005. Neural circuitry that
governs Drosophila male courtship behavior. Cell 121:795-807
(doi:10.1016/j.cell.2005.04.026).

Strutz, A., Soelter, J., Baschwitz, A., Farhan, A., Grabe, V., Rybak, J., Knaden M,
Schmuker M, Hansson BS, Sachse, S. (2014). Decoding odor quality and intensity
in the Drosophila brain. Elife, 3, e04147. (doi:10.7554/eLife.04147)

Thistle, R., Cameron, P., Ghorayshi, A., Dennison, L., & Scott, K. (2012). Contact
chemoreceptors mediate male-male repulsion and male-female attraction during
Drosophila courtship. Cell, 149(5), 1140-1151. (doi:10.1016/j.cell.2012.03.045)

Toda, H., Zhao, X., & Dickson, B. J. (2012). The Drosophila female aphrodisiac
pheromone activates ppk23+ sensory neurons to elicit male courtship behavior.
Cell Reports 1:599-607. (doi:10.1016/j.celrep.2012.05.007)

Vosshall, L. B., & Stocker, R. F. (2007). Molecular architecture of smell and taste in Drosophila. Annu. Rev. Neurosci. 30:505-533. (doi:10.1146/annurev.neuro.30.051606.094306)

Wilson RI. 2013. Early olfactory processing in *Drosophila*: mechanisms and principles. Annu Rev Neurosci 36:217-241 (doi:10.1146/annurev-neuro-062111-150533)

Yamamoto, D., & Koganezawa, M. (2013). Genes and circuits of courtship behaviour in Drosophila males. Nature Reviews Neuroscience 14:681-692. (doi:10.1038/nrn3567)

Figure legends

Figure 1

Odour-mediated upwind flight attraction of fruit fly *Drosophila melanogaster* males (blue bars) and females (ochre bars) to (a) vinegar and yeast headspace, (b) 10 ng/min male sex pheromone cVA, (c) 10 ng/min female sex pheromone *Z*4-11Al, alone and in blends with vinegar or yeast headspace, respectively. Letters of the corresponding colours show differences between treatments using Wald pairwise comparisons. Asterisks show treatments with significant differences between sexes (n=40).

Figure 2

Effect of *Z*4-11Al on courtship in *D. melanogaster* males. Decapitated target flies were painted with 1 ng test compound or heptane (solvent control). The number of courting and non-courting males in each test was compared using a chi-square test, asterisks show significant differences.

(a) Proportion of unmated or mated *D. melanogaster* (Dalby) test males courting unmated decapitated female or male target flies treated with 1 ng *Z*4-11Al or heptane (solvent control) (n=80, P=0.006 and P=0.007).

(b) Proportion of mated Canton-S and Dalby strain *D. melanogaster* males and *D. simulans* males, courting decapitated Dalby or *D. simulans* females, treated with 1 ng *Z*4-11Al or heptane (n=80, P=0.009 and P=0.039).

(c) Effect of RNA interference, in Orco-Gal4/uas-Or69aRNAi mated males, courting Dalby target females, treated with 1 ng *Z*4-11Al or heptane. Parental lines, Orco-Gal4 and uas-Or69aRNAi, show significant courtship behaviour (n=80, P=0.036 and P=0.032).

(d) Proportion of mated Dalby males courting decapitated Dalby females, treated with *E*4-11Al, (*R*)-linalool or ethyl butyrate (n=80).

 Figure 3

Calcium imaging responses in the antennal lobe (AL). Colours show the median
normalized calcium activity (Δ F/F [%]) in response to controls and odor applications,
according to the colour bar on the left (*n*=10; mean±SD).

(a) Calcium activity in Orco-Gal4 males (top) and females (below) in response to
control (air, mineral oil), to 60 and 600 ng *Z*4-11Al, and to 600 ng (*R*)-linalool. *Z*4-11Al
and (*R*)-linalool are key ligands for the D glomerulus (Lebreton et al. 2017).

(b) Responses of Or22a-Gal4 males to *Z*4-11Al in the DM2 glomerulus and of Or43b-
Gal4 males in the DM5 and VM2 glomeruli; ethyl butyrate is a diagnostic stimulus for
these glomeruli.

(c) Projection neuron (PN) responses in GH146-Gal4 males to *Z*4-11Al and (*R*)-linalool
in D and DM2 glomeruli.

Figure 1. Odour-mediated upwind flight attraction of fruit fly Drosophila melanogaster males (blue bars) and females (ochre bars) to
(a) vinegar and yeast headspace,
(b) 10 ng/min male sex pheromone cVA,
(c) 10 ng/min female sex pheromone Z4-11Al, alone and in blends with vinegar or yeast headspace, respectively.
Letters of the corresponding colours show differences between treatments using Wald pairwise comparisons. Asterisks show treatments with significant differences between sexes (n=40).

Figure 2. Effect of Z4-11Al on courtship in D. melanogaster males. Decapitated target flies were painted with 1 ng test compound or heptane (solvent control). The number of courting and non-courting males in each test was compared using a chi-square test, asterisks show significant differences.

(a) Proportion of unmated or mated D. melanogaster (Dalby) test males courting unmated decapitated female or male target flies treated with 1 ng Z4-11Al or heptane (solvent control) (n=80, P=0.006 and P=0.007).

(b) Proportion of mated Canton-S and Dalby strain D. melanogaster males and D. simulans males, courting decapitated Dalby or D. simulans females, treated with 1 ng Z4-11Al or heptane (n=80, P=0.009 and P=0.039).

(c) Effect of RNA interference, in Orco-Gal4/uas-Or69aRNAi mated males, courting Dalby target females, treated with 1 ng Z4-11Al or heptane. Parental lines, Orco-Gal4 and uas-Or69aRNAi, show significant courtship behaviour (n=80, P=0.036 and P=0.032).

(d) Proportion of mated Dalby males courting decapitated Dalby females, treated with E4-11Al, (R)-linalool or ethyl butyrate (n=80).

Figure 3. Calcium imaging responses in the antennal lobe (AL). Colours show the median normalized calcium activity (Δ F/F [%]) in response to controls and odor applications, according to the colour bar on the left (n=10; mean±SD).
(a) Calcium activity in Orco-Gal4 males (top) and females (below) in response to control (air, mineral oil), to 60 and 600 ng Z4-11Al, and to 600 ng (R)-linalool. Z4-11Al and (R)-linalool are key ligands for the D glomerulus (Lebreton et al. 2017).
(b) Responses of Or22a-Gal4 males to Z4-11Al in the DM2 glomerulus and of Or43b-Gal4 males in the DM5 and VM2 glomeruli; ethyl butyrate is a diagnostic stimulus for these glomeruli.
(c) Projection neuron (PN) responses in GH146-Gal4 males to Z4-11Al and (R)-linalool in D and DM2 glomeruli.