Compensating complete loss of signal recognition particle during co-translational

protein targeting by the translation speed and accuracy

Liuqun Zhao[a,b,c], Gang Fu[a,c], Yanyan Cui[a,c], Zixiang Xu[a,d], Dawei Zhang[a,b,c,d*]

[a]Tianjin Institute of Industrial Biotechnology, Chinese Academy of Sciences, Tianjin,

China

[b]University of Chinese Academy of Sciences, Beijing, China

[c]Key Laboratory of Systems Microbial Biotechnology, Chinese Academy of Sciences,

Tianjin, China.

[d]National Engineering Laboratory for Industrial Enzymes, Chinese Academy of

Sciences, Tianjin, China.

Running Head: Translational control regulates protein targeting

[*]For correspondence. E-mail: zhang_dw@tib.cas.cn

ABSTRACT

Signal recognition particle (SRP) is critical for delivering co-translational proteins to the bacterial inner membrane. Previously, we identified SRP suppressors in *Escherichia coli* that inhibit translation initiation and elongation, which provides an insight into the mechanism of bypassing the requirement of SRP. Suppressor mutations tend to be located in regions that govern protein translation under evolutionary pressure. To verify this hypothesis, we re-executed the suppressor screening of SRP. Here we isolated a novel SRP suppressor mutation located in the Shine-Dalgarno sequence of S10 operon, which partially offset the targeting defects of SRP-dependent proteins. We found that the suppressor mutation slowed the translation rate of proteins, especially of inner membrane proteins, which could compensate for the targeting defects of inner membrane proteins via extending the time window of protein targeting. Furthermore, the fidelity of translation was decreased in suppressor cells, suggesting that the quality control of translation was inactivated to provide a survival advantage under the loss of SRP stress. Our results demonstrate that the inefficient protein targeting due to SRP deletion can be rescued through modulating translational speed and accuracy.

SUMMARY

The signal recognition particle (SRP)-dependent delivery pathway is essential for membrane protein biogenesis. Previously, we reported that SRP is nonessential in *Escherichia coli* and slowing translation speed plays a critical role in membrane protein targeting. Here we identified a novel SRP suppressor that is also involved in translation. We found that translation speed and accuracy regulate membrane protein targeting. A slowdown of translation speed extends the time window for protein targeting. Meanwhile, a moderate decrease in translation fidelity ensures a suitable translation speed for better cell growth. These results provide that translation control could be a practical way to compensate for the loss of SRP.

KEYWORDS: signal recognition particle, suppressor, translational control, co-translational protein targeting

INTRODUCTION

The signal recognition particle (SRP) is a highly conserved ribonucleoprotein complex involved in co-translational targeting of the ribosome-nascent chain complex to the endoplasmic reticulum of eukaryotes or the inner membrane of prokaryotes (1, 2). Although the size and composition of SRP are variable in different species, the key subunits of SRP are evolutionarily conserved (3). The *Escherichia coli* SRP is much simpler than that in eukaryotes, as it contains an essential and highly conserved subunit called the fifty-four homologue (Ffh), which is homologous to mammalian SRP54, and a small stable 4.5S RNA, which is homologous to domain IV in the mammalian 7S SRP RNA (4-6). The *E. coli* SRP components can replace their mammalian homologues to mediate efficient co-translational protein targeting of mammalian proteins (4, 5). This suggests that the subunit SRP54 and domain IV of the 7S SRP RNA form the core conserved elements of SRP, and SRP is remarkably conserved from bacteria to mammals. SRP is primarily responsible for delivering integral membrane proteins (7, 8). It recognizes hydrophobic transmembrane domains or signal sequences when they emerge from the ribosome exit tunnel (9). Further, the nascent polypeptides must be targeted by SRP in a limited time window before they lose their competence due to aberrant aggregation (10, 11). Given the crowded cellular environment, it is challenging to correctly translocate newly synthesized proteins from the cytosol to the membrane.

SRP is essential in all organisms (12), except for the eukaryote *Saccharomyces cerevisiae* (13) and prokaryotes *Streptococcus mutans* (14) and *E. coli* (15). In these cases, the global repression of protein synthesis is associated with the loss of SRP (13,

15, 16). However, the precise mechanism of this tolerance remains poorly understood

(13-16). One of the best-understood mechanisms is slowing translation elongation rate

that extends the time window for targeting translating ribosomes. Previous studies have

demonstrated that slowing translation speed contributes to enhancing protein folding

and targeting (17-19). In co-translational translocation, protein folding and targeting are

inherently coupled to its elongation (20). The kinetic competition between protein

translation elongation and targeting modulates the efficiency of the co-translational

targeting pathway (17, 21). In eukaryotes, the Alu domain of SRP arrests nascent chain

translation elongation during its targeting, which is thought to pause translation

elongation until the targeting is completed (22, 23). Further, the Shine-Dalgarno-like

sequence pauses translation before the second transmembrane domain exposed in *E.*

coli, which facilitates the proper folding and targeting of membrane proteins (24). Thus,

the translation elongation is at the center of protein folding and targeting.

In an earlier study, we identified SRP suppressors: two translation initiation factors

IF2 and IF3, and a ribosomal protein RS3. The suppressor mutations decrease the

translation initiation and elongation rate (15). There are two possible explanations for

the slowing translation rate. First, it is possible that suppressor mutations directly inhibit

the translation initiation and then decrease the translation elongation rate because the

translation initiation rate is closely correlated with the elongation rate (25).

Alternatively, the ribosome is a hub in protein homeostasis, which modulates protein

translation and folding (19, 26). Here, we isolated an SRP suppressor located at the

Shine-Dalgarno (SD) sequence of ribosome S10 operon that could affect ribosomes

biogenesis. We addressed how the suppressor regulates the translation process and alleviates the fitness loss. Our results suggested that ribosome mutation suppressed the loss of SRP, although the cell growth was severely inhibited and the targeting of SRP-dependent proteins was not completely compensated. This SRP suppressor reduced translation rate, especially of inner membrane proteins. Moreover, in response to SRP deletion stress, the degree of mistranslation was increased to improve cell viability. Overall, our results showed that mutations in the ribosome may contribute to protecting cells from lethal damages caused by the loss of SRP.

RESULTS

Identification suppressors of SRP. Previous study in our laboratory demonstrated that SRP suppressors were all associated with protein translation (15). To determine whether suppressor mutations of SRP deletion were all mapped to chromosomal loci that influence protein translation, we used the same suppressor approach as previously described to screen SRP suppressors (15). We obtained another suppressor strain MY1901 that could survive when SRP was deleted (Fig. 1A). The growth rate of MY1901 was significantly reduced compared with that of wild-type strain MG1655, suggesting that the MY1901 strain had a severe growth defect. We also found that a longer lag time during the growth course of MY1901 than that of MG1655 (Fig. 1A), indicating that the lag time before regrowth may provide ample time for the cell to adapt to the absence of SRP. Whole-genome sequencing of the suppressor strain MY1901 and the original strain MG1655 allowed us to identify the suppressor mutation site in the SD sequence of ribosome S10 operon (Fig. 1B; Table S2). The S10 operon encodes 11

different ribosomal proteins and is responsible for protein synthesis (27). To determine

whether restoration of the wild-type alleles could revert to the wild-type strain, the Ffh

expression and reverting suppressor mutation to the wild-type allele in MY1901 strain

were carried out (Fig. 1C). The expression of Ffh markedly shortened the lag time and

increased the growth rate (Fig. 1D). Further, reverting the suppressor mutation to the

wild-type allele makes the growth rate of strain MY1901FS close to that of wild-type

strain MG1655 (Fig. 1D). Thus, the deletion of Ffh and suppressor mutation indeed

reduced cell growth. These results suggest that the mutation in the SD sequence of S10

operon contributes to the cell survival without SRP and this mutation is a novel

suppressor of SRP.

We hypothesized that the evolutionarily selective forces shaped the translation

process during screening suppressors in SRP-deletion cells. Thus, we tried to improve

cell growth through laboratory evolution. The strain MY1901 were analyzed by

continual subculture without any pressure over eighty rounds. The evolved strain

showed insignificant change in the growth rate but shorter lag time before regrowth and

an increased biomass compared with the initial strains (Fig. S1). The laboratory evolved

strains were sequenced and new changes in genes were identified (Table S2). In the

evolved strain MY1901, we found a mutation in the ribosomal protein S10 (rpsJ)

belonging to the S10 operon, confirming that the S10 operon plays an important role in

cell survival. Additionally, we obtained mutations in the DNA replication initiator

protein DnaA (28) and RNA polymerase rpoBC operon (29) (Table S2), which regulate

gene expression. Taken together, protein synthesis plays a critical role in suppressing

the loss of SRP.

Effects of SRP suppressor on protein synthesis. To examine the differences between

wild-type and mutant SD sequence, the mfold web server (30) was used to measure

mRNA structures of the S10 leader sequence containing the SD sequence. S10 leader

is required for the regulation of S10 operon (31). We found a 2.4 kcal mol^{-1} difference

in the minimum free energy of the thermodynamic ensemble of the predicted structure

of mutant SD compared to the wild-type SD (Fig. S2), suggesting that the S10 leader

with mutant SD was less stable. Because translation initiation is partially modulated by

the SD sequence (32), the effects of the mutant SD sequence on protein translation were

determined. The expression level of the green fluorescent protein (GFP) was used to

characterize protein translation levels in cells. Regardless of whether under the original

promoter P_{S10} or the arabinose inducible promoter P_{araBAD}, the expression level of GFP

with the mutated SD (SD*) was decreased (Fig. 2A), confirming that the suppressor

mutation weakened the binding of mRNA SD sequence to rRNA in ribosomes. This

result suggested that the translation initiation rate of S10 operon was reduced. To

examine whether the suppressor mutation decreases the expression levels of ribosomal

proteins, the whole-cell lysates proteome of strain MY1901 was analyzed (Data set

S1A). The expression level of each gene was normalized by that in wild-type strain

MG1655 (15). Unexpectedly, the levels of S10 operon and even the overall ribosomal

proteins were upregulated (Fig. 2B; Data set S1B). Given that the protein synthesis is

closely dependent on the ribosome content (33), there is a possibility that the protein

synthesis is enhanced in suppressor cells. However, the suppressor strain MY1901

showed a lower protein synthesis level because of a severe cell growth (Fig. 1A). Thus,

the increased pool of ribosomal proteins may be caused by the stalled ribosomes at the

translation initiation site, which are consistent with our previous study (15). Taken

together, although ribosomal protein biogenesis was increased, the protein synthesis

was reduced because of the inefficient translation.

Translation speed and accuracy in suppressor cells. Why did some ribosome

proteins accumulate in suppressor strain MY1901? One possibility is that ribosome

slowly translates mRNAs in suppressor cells. To test this possibility, we carried out

polysome profiling, which could detect various defects of translation. The polysome

profile of suppressor cells was changed relative to wild-type cells (Fig. 3A). The 30S

to 50S (30S/50S) ratio was not significantly altered in suppressor and wild-type strains,

but the 30S and 50S peaks of suppressor cells were higher than that of wild-type cells

(Fig. 3A), indicating an increase in the proportion of free subunits. The 70S peak of the

wild-type cells was almost indistinguishable from that of suppressor cells, suggesting

that some 70S ribosomes may pause at the translation initiation. We observed that the

polysome to 70S monosome (P/M) ratio was decreased in suppressor cells (Fig. 3A),

suggesting that translation elongation was inhibited in suppressor cells. Thus, the

suppressor mutation plays a role in protein translation. Translation can be divided into

four phases: initiation, elongation, termination, and ribosome recycling (34). We next

compared the protein abundances of translation associated factors via whole-cell lysates

proteome analysis. Ribosome recycling factor (RRF) and elongation factor-G (EF-G)

are involved in ribosome recycling (35). The expression of RRF and EF-G was

downregulated in suppressor strain (Fig. 3B; Data set S1B), indicating that the ribosome

recycling was reduced, which may be caused by slowing translation in suppressor cells.

Although the expression of release factors mediating translation termination was

upregulated, the efficiency of translation termination at three stop codons was only

slightly increased (Fig. 3B and C). This data showed that the translation termination

rate was not significantly impacted in suppressor cells. We observed that the expression

level of elongation factors in suppressor cells was inconsistent and changed little

compared to that in wild-type cells (Fig. 3B; Data set S1B). To exactly examine the

efficiency of translation elongation, we measured its fidelity and speed through

experiments. In strain MY1901, we observed a marked increase in translational

frameshifting readthrough relative to that in wild-type strain (Fig. 3D), thus the fidelity

of translation elongation was decreased. Additionally, a LacZ induction assay was used

to measure the translation elongation speed (36). We used a cytoplasmic protein FusA

and an inner membrane protein MsbA to define the level of translation elongation rates

(Fig. S3). When cells were grown at the same rich growth media (Glucose + cAA), the

growth rates of wild-type and suppressor cells were approximately 1.0 h^{-1}(15) and 0.6

h^{-1}, respectively (Fig. S3A; Table S3). As expected, the translation elongation rate of

suppressor cells was markedly decreased compared with that of wild-type cells (Fig.

3E; Fig. S3B to F). As the translation elongation closely depends on growth rate (37),

we further reduced the growth rate of wild-type cells to 0.6 h^{-1} that was similar to that

of suppressor cells (Fig. S3A). The elongation rate of the MsbA in suppressor cell was

reduced by approximate 2.0 aa s^{-1} (amino acids per second) compared with that in wild-

type cells, but the elongation rate of the FusA was similar in both wild-type and

suppressor cells (Fig. 3E; Fig. S3B toD). We also observed that the elongation rate of

MsbA was slower than that of FusA (Fig. 3E; Fig. S3B to F), which is consistent with

the issue that the translation speed of inner membrane proteins is slowed down during

targeting, but not that of cytoplasmic proteins (24). These data suggested that the effect

of selectively reducing translation elongation rate of inner membrane proteins was

enhanced in suppressor cells. For the expression of translation initiation factors in

suppressor cells, IF1 and IF3 were downregulated but IF2 was upregulated (Fig. 3B).

Given that IF2 accelerates ribosomal subunit joining, whereas IF1 and IF3 slowed down

subunit association (38, 39), the capacity of formation of 70S initiation complex may

not be significantly impaired although cells showed severe defects in growth. We next

examined the translation initiation rate by a computational model called the

homogeneous ribosome flow model (HRMF), in which the translation elongation rate

was assumed to be constant (40). The translation initiation rate can be estimated by the

measurable translation rate and elongation rate (Table S4). The translation initiation

rate in suppressor cells showed similar trend to the translation elongation rate. The

initiation rate of FusA was not markedly changed in the suppressor and wild-type cells

when grown at the same growth rate (Fig. 3F). However, the inner membrane protein

MsbA had a slower translation initiation rate in suppressor cells than that in wild-type

cells when grown at the same growth rate (Fig. 3F). These results suggested that the

translation initiation rate was selectively decreased in suppressor cells. As translation

initiation factors play a vital role in translation initiation fidelity (39), we addressed

whether the suppressor was detrimental to the fidelity of start codon selection and
initiator tRNA binding. We changed the start codon of GFP from AUG to GUG, UUG,
AUA and AUC and the initiator tRNA codon of GFP from AUG: CAU to UAG: CUA,
CAC: GUG and UAC: GUA. We measured the GFP fluorescence of these GFP variants
in the wild-type and suppressor cells. We observed that the fidelity of translation
initiation in suppressor cells was decreased at different levels compared with that in
wild-type cells (Fig. 3G), indicating that suppressor mutation may impair the fidelity
of translation initiation. Taken together, the suppressor cell trades translation speed and
accuracy for cell survival in the absence of SRP.

SRP-dependent protein targeting in suppressor cells. In *E. coli*, most inner
membrane proteins are delivered by the co-translational SRP pathway (41). In principle,
SRP-dependent proteins are not properly targeted to the bacterial cytoplasmic
membrane after Ffh depletion (42, 43). To test whether the suppressor mutation could
suppress protein targeting defects, we first examined cell morphological changes by
scanning electron microscopy (SEM) and transmission electron microscopy (TEM).
SEM images showed that suppressor strain MY1901 still had typical rod morphology
but had rougher surface relative to the wild-type strain MG1655 (Fig. 4A). TEM images
showed that suppressor strain MY1901 retained cell wall integrity but had damaged
inner membrane structure (Fig. 4A). MY1901 displayed a significant detachment of the
inner membrane from the outer membrane (Fig. 4A). Thus, the suppressor mutation
partially offsets the effects of the loss of SRP pathway on inner membrane protein
translocation. To gain insight into the localization of inner membrane proteins, we

performed proteomic analysis of inner membrane proteins in strains wild-type

MG1655(15) and suppressor MY1901 (Data set S1C). The *E. coli* strain HDB51 was

used as a control, in which the expression of Ffh was induced by arabinose (44).

Depleted Ffh can be obtained after several hours of incubation in the presence of

glucose, thus yielding the SRP⁻strain (45). The inner membrane proteome analysis of

SRP⁻ was also conducted (Data set S1C). According to the previous study, we identified

262 SRP-dependent inner membrane proteins (15). Since the completely untargeted

inner membrane proteins would aggregate or misfold in cytoplasm, the detected inner

membrane proteins by inner membrane mass spectrometry analysis are primary in their

folded and targeted state. We found that targeting levels of many identified SRP-

dependent inner membrane proteins in both MY1901 and SRP⁻ cells were higher than

their targeting levels in wild-type cells (Fig. 4B; Data set S1D), indicating that these

inner membrane proteins can target to the cytoplasmic membrane in the absence of SRP,

which is consistent with the previous study that the only part of the inner membrane

protein targeting was completely inhibited when SRP is inefficient (42). We also found

that more proteins were successfully targeted in MY1901strain than in SRP⁻ strain (Fig.

4B), suggesting that the suppressor mutation indeed plays a role in inner membrane

protein targeting without SRP. FtsQ is an SRP-dependent protein, which is often used

as a model protein for studying SRP-mediated protein targeting (17, 46). However, the

targeting of FtsQ was not significantly inhibited in strains SRP⁻ and MY1901 (Fig. 4B;

Data set S1D). This may be explained by the fact that FtsQ is a single-spanning

membrane protein that is easily targeted to the membrane (15, 47). To confirm this

result, we examined the targeting level of FtsQ by protein biotinylation (17, 48, 70).

The Avi-tag was fused to the periplasmic domain of the targeted proteins. The

biotinylated proteins would be the untargeted proteins in which the periplasmic

domains are exposed in the cytosol. Thus, the protein biotinylation can be used for

protein targeting assay. As expected, FtsQ can target to the inner membrane in all

different cells (Fig. 4C), which confirmed the prediction of proteomic analysis (Fig.

4B). We also found that the targeting levels of the SRP-independent protein EspP in all

strains were similar (Fig. 4C), suggesting that the targeting of SRP-independent

proteins was not affected by the loss of SRP. Thus, the suppressor enables SRP-

dependent protein targeting which did not occur at the cost of failed targeting of SRP-

independent proteins.

DISCUSSION

Co-translational protein targeting by SRP is an essential and conservative pathway that

delivers most inner membrane proteins to their correct subcellular destinations (3).

Early work revealed that SRP was not essential in *E. coli* when the translation initiation

and elongation rate were decreased (15). Isolation of suppressors is a useful strategy to

provide insight into certain molecular mechanisms by suggesting which cellular

component is involved in an inefficient process (49). The SRP suppressors involved in

protein translation initiation have been identified before, and these suppressors directly

affect the translation process (15). In this study, we obtained an SRP suppressor

associated with protein translation too. Additionally, the expression of heat shock

response related chaperones and proteases was not upregulated (Fig. S4A; Data set

S1B), suggesting that the heat shock response plays little role in compensating the loss

of SRP, which is consistent with the previous study (15). In strain MY1901, the

expression of SecA was not affected and other transport machinery components such

as SecYEG-YajC-SecDF, YidC, and FtsY showed a decreased level relative to those in

the wild-type strain (Fig. S4B; Data set S1E). Moreover, the expression level of these

components was similar in strains MY1901 and SRP⁻ (Fig. S4B; Data set S1E). These

results suggest that protein transport machinery components were not likely involved

in protein targeting without SRP. Since the translation consumes a large amount of

energy, it is important to properly regulate the translation (50). However, there is

another possibility that the suppressor mutation may indirectly interfere with the

general translation process, and any strategies inhibiting translation can offset the

missing SRP. To exclude this possibility, we added a low level of chloramphenicol that

is well-known to block translation elongation into the medium of SRP⁻ cells. We found

that chloramphenicol did not suppress cell growth defects in the absence of SRP (Fig.

S5), thus simply decreasing protein synthesis could not suppress the defects of SRP

deletion, which is inconsistent with the previous result that chloramphenicol suppresses

the SecY mutant (49). Together with the previous study (15), the SRP suppressors tend

to be identified in translation machinery.

Consistent with the previous observation (15), we observed that deletion of SRP

led to inefficient protein synthesis, suggesting a close relationship between protein

synthesis and translocation. Although the translation was repressed in strain

MY1901(Fig. 3E and F), the ribosomal protein expression was upregulated (Fig. 2B)

and the 30S and 50S ribosomal subunits were accumulated (Fig. 3A). This led to the

belief that the ribosomes may be inactive and accumulate in the cytosol. Consistent

with the earlier report (15), we observed that the assembly of 70S particles was not

significantly impaired (Fig. 3A). Thus, the 70S ribosomes initiation complex formation

process was not inhibited. Given that the protein translation initiation was impeded (Fig.

3E and F), we speculated that this impediment was due to the stalled initiation-like 70S

ribosome complex. However, we observed that the initiation time of translation was

constant in wild-type and suppressor cells under different growth rates (Table S3), thus

the initiation pausing can be negligible in suppressor cells and the suppressor mutation

did not affect the translation initiation process. Considering the translation elongation

rate is dependent on the growth rate (37), the growth defect may directly play a big role

in inhibiting translation elongation of membrane proteins. However, this possibility can

be excluded, because suppressor cells were grown under the rich nutrient condition that

can supply enough macromolecular for ribosomal synthesis and translation elongation,

and the cell growth could not automatically decrease (37, 51). Moreover, under the

same growth rate, the elongation rate of inner membrane protein in the suppressor cells

was also decreased relative to that in the wild-type cells (Fig. 3E). As the level of protein

translation and the growth rate is closely interweaved (51) and the translation

elongation plays a critical role in protein homeostasis that is important for cell growth

(52, 53), it is most likely that the cell growth rate depends on the level of elongation

rate and the slowing elongation rate are caused by the suppressor mutation. Increasing

evidence has supported the notion that the translation elongation of nascent polypeptide

regulates the targeting of SRP-dependent proteins (17, 21), thus decreasing elongation

rate contributes to the survival of SRP deletion cells. Interestingly, relative to wild-type

cells, the reduction level of translation rate of the cytoplasmic protein FusA was less

than that of the inner membrane protein MsbA in suppressor cells (Fig. 3E and F).

Hence, the suppressor mutation may be biologically favored to reduce the translation

of inner membrane proteins, which may be beneficial to cell fitness. Decreasing

translation elongation rate plays a critical role in suppressing the loss of SRP by

extending the time window for protein targeting (15). SRP binds to the ribosome-

nascent chain complex when the N-terminus of the first TMD is exposed from the

ribosome. The maximal SRP binding site is 55 amino acids from the ribosomal peptidyl

transferase center in *E. coli* (54). Assuming that ~30 amino acids can fit into the

ribosome exit tunnel (55), 25 residues would be exposed outside the tunnel. At a

translation elongation rate of ~15 aa s^{-1} in rapid growth conditions (15), the maximum

time required for protein localization is ~2 s (Fig. 5). Thus, with the help of SRP, most

translating ribosomes move to the membrane within this period in *E. coli*. Without SRP,

suppressors showed the translation elongation rate to ~11 aa s^{-1}(Fig. 3E; Table S3),

which also provides ~2 s for nascent chains of 55 amino acids to target to the inner

membrane (Fig. 5). However, it is not likely that nascent chains successfully target to

the membrane within ~2 s without SRP. To get a longer time to find the membrane, the

length of translating nascent chains is more likely longer than 55 amino acids. However,

the nascent chain cannot exceed a specific length as aggregation would prevent protein

from being targeted (10, 11), and this specific length is called the critical length (L) for

targeting. Proteins with fewer transmembrane domains (TMDs) or longer first loop length have a longer critical length (15). Furthermore, if the nascent chain exceeds a critical length of ~140 amino acids, it becomes translocation-incompetent (10, 11). In suppressor cells, the upper limit of the critical time for protein targeting would be ~10 s (Fig. 5). If the targeting time of some SRP-dependent proteins exceeds 10 s, these proteins would not be targeted to the inner membrane in suppressor cells. Thus, the suppressor extends the time window to ~2-10 s (Fig. 5). Taken together, this model shows that SRP greatly shortened the protein targeting time up to 8 s, which minimizes the cost of targeting and maintains a fast growth. Furthermore, we showed that the translation fidelity was decreased in suppressor cells (Fig. 3D and G). Earlier work revealed that mistranslation can provide a growth advantage in response to stress (56). Hence, the suppressor mutation may inactive the translation quality control mechanism to alleviate the stress response and maintain cell growth. Overall, our data suggest that in response to the deletion of SRP, suppressor cells attenuate translation elongation to give the translating ribosomes more time to find and target to the inner membrane. Additionally, suppressor mutation selectively permits SRP-dependent proteins translation and reduced the fidelity of translation, which is vital to cell survival.

As expected, the suppressor mutation can partially offset the defective targeting of inner membrane proteins (Fig. 4B), which is consistent with the previous result (15). Furthermore, we found that in the SRP-depletion strain SRP$^-$, many proteins can successfully target to the inner membrane (Fig. 4B). However, proper localization of these proteins cannot bypass the requirement of SRP (57). We speculated that the

proteins that can be correctly located in suppress strain MY1901 but not in SRP-

depletion strain SRP⁻ may be responsible for cell survival. We hypothesized that

specific membrane protein targeting defects can block the essential cellular process,

which would be responsible for the loss of cell viability. Among these localization

defective proteins, only one protein PgsA is essential for *E. coli* (Data set S1F). PgsA

catalyzes the step in the synthesis of the acidic phospholipids that are considered to be

indispensable in multiple cellular processes (58, 59). We speculated that mislocalization

of PgsA may inhibit cell growth. Furthermore, in SRP⁻ strain, some transportation

associated proteins (AsmA, Bcr, PheP, YbaL, and YidE) did not successfully target to

the inner membrane (Data set S1F), which would impair membrane traffic and decrease

energy production. Further studies are needed on the protein targeting defects that

determine whether cells devoid of SRP will survive.

MATERIALS AND METHODS

Bacterial strains, plasmids and media. All bacterial strains, plasmids and primers

used in this study are listed in Table S1. *E. coli* K-12 MG1655 derivative strains were

grown either in LB medium or on LB agar at the indicated temperature. *E. coli* HDB51

strain in which the expression of Ffh is under the control of an arabinose inducible

promoter, were grown in LB medium containing 0.2% arabinose (44). The SRP

suppressor strain MY1901 was isolated and validated as previously described. The

antibiotics kanamycin, gentamycin, ampicillin, and chloramphenicol were used at a

concentration of 50, 10, 100, and 200 μg ml^{-1}, respectively. The *lac*, *trc* and *tac*

promoters were induced with 0.02 mM isopropyl-β-D-1-thiogalactopyranoside (IPTG).

The *araBAD* promoter was induced with 0.2% arabinose.

Measurement of cell growth. The bacterial growth curves and growth rates were

estimated as previously predicted (15). For bacterial growth under oxidative and

osmotic stress, overnight cultures were diluted into liquid LB media containing 2 mM

or 8 mM H_2O_2 and 2.5% or 5% NaCl, respectively. For the thermal stress experiment,

the medium was chilled to 30°C or heated to 42°C before inoculation.

Polysome analysis, cell ultrastructure and proteomic analysis. Strains were grown

at 37°C at the early-exponential growth in LB medium and harvested by centrifugation

(15). Polysome analysis was performed as described (60, 61). The scanning electron

microscopy (SEM) and transmission electron microscopy (TEM) analyses were carried

out as described previously (15). The whole-cell lysates and inner membrane proteins

of strains MY1901 and SRP⁻ were isolated and analyzed by liquid chromatography-

tandem mass spectrometry (LC-MS/MS) as described (15, 62-64), with modifications.

For seed culture, the strain MY1901 or HDB51was inoculated in LB medium at 37°C

overnight and the strain HDB51 was grown under the addition of 0.2% arabinose. The

overnight culture of MY1901 was diluted into the fresh LB medium with an initial

OD_{600} of 0.02-0.03. Strain MY1901 was harvested in the mid-exponential phase. The

overnight culture of HDB51 was washed for three times with fresh LB medium and

then incubated in LB with addition of 0.2% glucose with an initial OD_{600} of 0.03-0.04

for several hours, yielding the SRP⁻ strain. This strain was harvested when entering the

stationary phase. Three sample replicates were prepared by performing collection of

cells from independent cultures. We used the Filter-Aided Sample Preparation (FASP)

strategy for proteome analysis (63). The inner membrane fraction was separated by

sucrose gradient (64). Then, the inner membrane was chemically treated by Na_2CO_3

and KCl to remove peripherally associated proteins (15). The MS sample of inner

membrane proteins was prepared by surface proteolysis (64). The sample was

characterized by sequential window acquisition of all theoretical spectra (SWATH)

analysis (65). The LC-MS/MS analysis was performed according to the previous study

(15). The obtained data were normalized by the median scale normalization (MedScale)

method (66).

β-galactosidase assay. The β-galactosidase activity was assayed as previously

described (67). Cells were grown in LB medium supplemented with 50 μg ml^{-1}

kanamycin at 37°C. Protein induction was performed when MG1655Δ*lacZ* and

MY1901Δ*lacZ* were grown to OD_{600} of 0.6-0.8 and 0.4-0.5, respectively. Two hours

later the cultures were harvested and assayed. β-galactosidase activity from the plasmid

encoding wild-type LacZ was used for normalization.

Measurements of translation efficiency and initiation fidelity. To determine the

effects of mutated SD sequence on protein translation efficiency, we constructed

plasmids carrying the *gfp* gene with the wild-type SD and suppressor mutation SD*

under the promoter P_{S10} or P_{araBAD}. To analyze the translation initiation fidelity, a set of

GFP variants were generated in which the start codon (AUG) or the initiator tRNA was

replaced with different non-canonical start codons and non-initiator tRNAs codons.

Several non-canonical start codons (GUG, UUG, AUA and AUC) and non-initiator

tRNAs codons (UAG: CUA, CAC: GUG and UAC: GUA) were used as potential start

codons and initiator tRNAs codons, respectively. *E. coli* cells were grown in 300 μl of

LB medium with necessary antibiotics in a 96-well deep well culture plate at 37°C

overnight followed by transferring and cellular fluorescence measurements (68). For

detecting the effects of Shine-Dalgarno (SD) sequence on translation efficiency and the

fidelity of translation initiation, the non-expression strains containing empty vectors

pJH30, pJH31 and pTrc99K lacking a reporter gene were used as controls. OD_{600} was

measured to estimate culture density, followed by fluorescence (excitation=488 nm,

emission=520 nm). Assays were carried out from six independent colonies. For the

initiation fidelity assay, the fluorescence intensity from the plasmid encoding mutant

GFP was normalized against the fluorescence intensity of wild-type or suppressor

strains containing plasmid encoding wild-type GFP.

Measurements of translation rates. Translation elongation rates were measured as

described previously (36, 37, 69). For seed culture, MG1655Δ*lacZ* and

MY1901Δ*lacZ*Δ*cat* cells were cultured in LB medium at 37°C for several hours, then

cultures were collected and washed with fresh MOPS medium. To make strains grow

well, MG1655Δ*lacZ* and MY1901Δ*lacZ*Δ*cat* it was grown in rich Glucose + cAA (0.2%

glucose + 0.2% casamino acids) MOPS medium overnight as the pre-culture. The

experiment culture was performed with an initial OD_{600} of 0.04-0.05. The subsequent

collection and measurement methods were performed as previously described (15). The

translation elongation rate was measured based on the LacZα induction assay. The

translation time of first newly synthesized LacZα, $T_α$, was estimated by measuring the

LacZα induction kinetics. The translation time of the first newly synthesized

LacZα fused protein (FusA-LacZα or MsbA- LacZα), T_{total}, was obtained by Schleif plot of the induction curve. The initiation time, T_{init}, equals $T_\alpha - \{90/[L/(T_{total}-T_\alpha)]\}$, where 90 is the 90 aa LacZα fragment and L is the length of the LacZα fusion protein (containing 10 aa linker). The translation elongation rate equals $(L+90)/(T_{total}-T_{init})$. When MY1901Δ*lacZ*Δ*cat* grew in Glucose + cAA medium, the growth rate of it was about 0.6 h^{-1}. To eliminate the cell growth effect on translation elongation rate, the wild-type strain MG1655Δ*lacZ* was grown in Glycerol + NH_4Cl (0.2% glycerol + 10 mM NH_4Cl) MOPS medium and the growth rate was similar to 0.6 h^{-1}. The measurement of the elongation rate of MG1655Δ*lacZ* with the growth rate 0.6 h^{-1} was performed similarly to that of MY1901Δ*lacZ*Δ*cat*. Translation initiation rate was estimated by a computational model homogeneous ribosome flow model (HRFM) (40, 70, 71). Based on the measured translation rate and elongation rate, the initiation rate can be calculated (15).

Protein targeting assay *in vivo*. The biotinylation of proteins has been successfully applied to SRP-dependent protein targeting *in vivo* (17, 48). *E. coli* enzyme biotin ligase (BirA) can specially ligate biotin to a 15-amino acid peptide (GLNDIFEAQKIEWHE) termed the Avi-tag (72). The BirA and Avi-tagged proteins were co-expressed for biotinylation. Cells were co-transformed with recombinant vectors p15A-birA and pJH29-EspP/FtsQ/LacZ-Avi and grown overnight at 37°C. Then overnight cultures were washed and diluted into 30 mL fresh LB medium at an initial OD_{600} of 0.02. For HDB51, the overnight culture that grown in LB medium with 0.2% arabinose was washed and diluted into 30 mL fresh LB medium at an initial OD_{600} of 0.02 with or

without arabinose to construct SRP$^+$ and SRP$^-$ cells, respectively. When cultures reached an OD$_{600}$ ~0.4-0.5 (SRP$^-$ cells were cultured for 2-3 h), protein expression was induced by 0.5 mM IPTG, and 100 μM biotin was also added at this point. After 3 h of cultivation, cells were harvested by centrifugation. For Ffh depletion, 0.2% glucose was added 2 h before harvesting of cells. Then the samples were analyzed by SDS-PAGE and immunoblotting. Biotinylated proteins were detected by streptavidin-HRP and the total amount of protein was detected by anti-FLAG antibody. Detection was performed by the DAB substrate kit (Thermo Scientific, USA).

Acknowledgments

We thank Dr. Manlu Zhu (Central China Normal University) for plasmid pKUT15-fusA-lacZα, Dr. Changhao Bi (TIB, CAS) for plasmid pRed_Cas9_ΔpoxB300, Dr. Harris D Bernstein (NIH) for strain HDB51 and plasmid pJH29 and Dr. Zhidan Zhang (TIB, CAS) for help with mass spectrometry. This work was supported by the National Key R&D Program of China (2020YFA0907800) and Tianjin Synthetic Biotechnology Innovation Capacity Improvement Project (TSBICIP-KJGG-004-03 and TSBICIP-KJGG-006).

Conflict of interest

The authors declare no conflict of interest.

Author contributions

LZ, GF and DZ designed the experiments. LZ and YC performed experiments. LZ, YC, and ZX analyzed the data. LZ and DZ wrote the manuscript.

REFERANCES

1. Walter P, Johnson AE. 1994. Signal sequence recognition and protein targeting to the endoplasmic reticulum membrane. Annu Rev Cell Biol 10:87–119.

2. Pool MR. 2005. Signal recognition particles in chloroplasts, bacteria, yeast and mammals. Mol Membr Biol 22:3–15.

3. Saraogi I, Shan SO. 2014. Co-translational protein targeting to the bacterial membrane. Biochim Biophys Acta 1843:1433–1441.

4. Bernstein HD, Zopf D, Freymann DM, Walter P. Functional substitution of the signal recognition particle 54-kDa subunit by its Escherichia coli homolog. Proc Natl Acad Sci U S A 90:5229–5233.

5. Powers T, Walter P. 1997. Co-translational protein targeting catalyzed by the Escherichia coli signal recognition particle and its receptor. EMBO J 16:4880–4886.

6. Regalia M, Rosenblad MA, Samuelsson T. 2002. Prediction of signal recognition particle RNA genes. Nucleic Acids Research 30:3368–3377.

7. Zhang X, Shan SO. 2014. Fidelity of cotranslational protein targeting by the signal recognition particle. Annu Rev Biophys 43:381–408.

8. Ulbrandt ND, Newitt JA, Bernstein HD. 1997. The E. coli signal recognition particle is required for the insertion of a subset of inner membrane proteins. Cell 88:187–196.

9. Ng DT, Brown JD, Walter P. 1996. Signal sequences specify the targeting route to the endoplasmic reticulum membrane. J Cell Biol 134:269–278.

10.   Siegel V, Walter P. 1988. The affinity of signal recognition particle for

presecretory proteins is dependent on nascent chain length. EMBO J 7:1769–

1775.

11.   Flanagan JJ, Chen JC, Miao Y, Shao Y, Lin J, Bock PE, Johnson AE. 2003.

Signal recognition particle binds to ribosome-bound signal sequences with

fluorescence-detected subnanomolar affinity that does not diminish as the

nascent chain lengthens. J Biol Chem 278:18628–18637.

12.   Akopian D, Shen K, Zhang X, Shan SO. 2013. Signal recognition particle: an

essential protein-targeting machine. Annu Rev Biochem 82:693–721.

13.   Mutka SC, Walter P. 2001. Multifaceted physiological response allows yeast to

adapt to the loss of the signal recognition particle-dependent protein-targeting

pathway. Mol Biol Cell 12:577–588.

14.   Kremer BH, van der Kraan M, Crowley PJ, Hamilton IR, Brady LJ, Bleiweis

AS. 2001. Characterization of the sat operon in Streptococcus mutans: evidence

for a role of Ffh in acid tolerance. J Bacteriol 183:2543–2552.

15.   Zhao L, Cui Y, Fu G, Xu Z, Liao X, Zhang D. 2020. Signal recognition particle

suppressor screening reveals the regulation of membrane protein targeting by

the translation rate. mBio 11: e02373-20.

16.   Hasona A, Zuobi-Hasona K, Crowley PJ, Abranches J, Ruelf MA, Bleiweis AS,

Brady LJ. 2007. Membrane composition changes and physiological adaptation

by Streptococcus mutans signal recognition particle pathway mutants. J

Bacteriol 189:1219–1230.

17.   Zhang D, Shan SO. 2012. Translation elongation regulates substrate selection

by the signal recognition particle. J Biol Chem 287:7652–7660.

18.   Siller E, DeZwaan DC, Anderson JF, Freeman BC, Barral JM. 2010. Slowing

bacterial translation speed enhances eukaryotic protein folding efficiency. J Mol

Biol 396:1310–1318.

19.   Sherman MY, Qian SB. 2013. Less is more: improving proteostasis by

translation slow down. Trends Biochem Sci 38:585–591.

20.   du Plessis DJ, Nouwen N, Driessen AJ. 2011. The Sec translocase. Biochim

Biophys Acta 1808:851–865.

21.   Chartron JW, Hunt KC, Frydman J. 2016. Cotranslational signal-independent

SRP preloading during membrane targeting. Nature 536:224–228.

22.   Walter P, Blobel G. 1981. Translocation of proteins across the endoplasmic

reticulum III. Signal recognition protein (SRP) causes signal sequence-

dependent and site-specific arrest of chain elongation that is released by

microsomal membranes. J Cell Biol 91:557–561.

23.   Mason N, Ciufo LF, Brown JD. 2000. Elongation arrest is a physiologically

important function of signal recognition particle. EMBO J 19:4164–4174.

24.   Fluman N, Navon S, Bibi E, Pilpel Y. 2014. mRNA-programmed translation

pauses in the targeting of E. coli membrane proteins. Elife 3:e03440.

25.   Riba A, Di Nanni N, Mittal N, Arhné E, Schmidt A, Zavolan M. 2019. Protein

synthesis rates and ribosome occupancies reveal determinants of translation

elongation rates. Proc Natl Acad Sci U S A 116:15023–15032.

26. Pechmann S, Willmund F, Frydman J. 2013. The ribosome as a hub for protein

quality control. Mol Cell 49:411–421.

27. Zengel JM, Lindahl L. 1994. Diverse mechanisms for regulating ribosomal

protein synthesis in Escherichia coli. Prog Nucleic Acid Res Mol Biol 47:331–

370.

28. Hughes P, Landoulsi A, Kohiyama M. 1988. A novel role for cAMP in the control

of the activity of the E. coli chromosome replication initiator protein, DnaA.

Cell 55:343–350.

29.   Ishihama A, Fukuda R. 1980. Autogenous and post-transcriptional regulation

of RNA polymerase synthesis. Mol Cell Biochem 31:177–196.

30. Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization

prediction. Nucleic Acids Res 31:3406–3415.

31. Allen T, Shen P, Samsel L, Liu R, Lindahl L, Zengel JM. 1999. Phylogenetic

analysis of L4-mediated autogenous control of the S10 ribosomal protein

operon. J Bacteriol 181:6124–6132.

32. Yang C, Hockenberry AJ, Jewett MC, Amaral LAN. 2016. Depletion of Shine-

Dalgarno Sequences Within Bacterial Coding Regions Is Expression Dependent.

G3 (Bethesda) 6:3467–3474.

33. Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T. 2010.

Interdependence of cell growth and gene expression: origins and consequences.

Science 330:1099–1102.

34. Rodnina MV. 2018. Translation in Prokaryotes. Cold Spring Harb Perspect Biol

10:a032664.

35.   Prabhakar A, Capece MC, Petrov A, Choi J, Puglisi JD. 2017. Post-termination

Ribosome Intermediate Acts as the Gateway to Ribosome Recycling. Cell Rep

20:161–172.

36.   Zhu M, Dai X, Wang YP. 2016. Real time determination of bacterial in vivo

ribosome translation elongation speed based on LacZalpha complementation

system. Nucleic Acids Res 44:e155.

37.   Dai X, Zhu M, Warren M, Balakrishnan R, Patsalo V, Okano H, Williamson J

R, Fredrick K, Wang YP, Hwa T. 2016. Reduction of translating ribosomes

enables Escherichia coli to maintain elongation rates during slow growth. Nat

Microbiol 2:16231.

38.   Naaktgeboren N, Roobol K, Voorma HO. 1977. The effect of initiation factor

IF-1 on the dissociation of 70-S ribosomes of Escherichia coli. Eur J Biochem

72:49–56.

39.   Ling C, Ermolenko DN. 2015. Initiation factor 2 stabilizes the ribosome in a

semirotated conformation. Proc Natl Acad Sci U S A 112:15874–15879.

40.   Margaliot M, Tuller T. 2012. On the steady-state distribution in the

homogeneous ribosome flow model. IEEE/ACM Trans Comput Biol Bioinform

9:1724–1736.

41.   Elvekrog MM, Walter P. 2015. Dynamics of co-translational protein targeting.

Curr Opin Chem Biol 29:79–86.

42.   Bernstein HD, Hyndman JB. 2001. Physiological basis for conservation of the

signal recognition particle targeting pathway in Escherichia coli. J Bacteriol

183:2187–2197.

43.   Wickström D, Wagner S, Baars L, Ytterberg AJ, Klepsch M, van Wijk KJ,

Luirink J, de Gier JW. 2011. Consequences of depletion of the signal recognition

particle in Escherichia coli. J Biol Chem 286:4598–4609.

44.   Lee HC, Bernstein HD. 2001. The targeting pathway of Escherichia coli

presecretory and integral membrane proteins is specified by the hydrophobicity

of the targeting signal. Proc Natl Acad Sci U S A 98:3471–3476.

45.   Zhang D, Sweredoski MJ, Graham RL, Hess S, Shan SO. 2012. Novel

proteomic tools reveal essential roles of SRP and importance of proper

membrane protein biogenesis. Mol Cell Proteomics 11:M111.011585.

46.   Scotti PA, Valent QA, Manting EH, Urbanus ML, Driessen AJ, Oudega B,

Luirink J. 1999. SecA is not required for signal recognition particle-mediated

targeting and initial membrane insertion of a nascent inner membrane protein. J

Biol Chem 274:29883–29888.

47.   Weiss DS, Chen JC, Ghigo JM, Boyd D, Beckwith J. 1999. Localization of FtsI

(PBP3) to the septal ring requires its membrane anchor, the Z ring, FtsA, FtsQ,

and FtsL. J Bacteriol 181:508–520.

48.   Jander G, Cronan J, Beckwith J. 1996. Biotinylation in vivo as a sensitive

indicator of protein secretion and membrane protein insertion. J Bacteriol

178:3049–3058.

49.   Lee CA, Beckwith J. 1986. Suppression of growth and protein secretion defects

in Escherichia coli secA mutants by decreasing protein synthesis. J Bacteriol
166:878–883.

50. Hershey JW, Sonenberg N, Mathews MB. 2012. Principles of translational
control: an overview. Cold Spring Harb Perspect Biol 4:a011528.

51. Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T. 2010.
Interdependence of cell growth and gene expression: origins and consequences.
Science 330:1099–1102.

52. Laskowska E, Kuczyńska-Wiśnik D, Lipińska B. 2019. Proteomic analysis of
protein homeostasis and aggregation. J Proteomics 198:98–112.

53. Stein KC, Frydman J. 2019. The stop-and-go traffic regulating protein
biogenesis: How translation kinetics controls proteostasis. J Biol Chem
294:2076–2084.

54. Schibich D, Gloge F, Pohner I, Björkholm P, Wade R, von Heijne G, Bukau B,
Kramer G. 2016. Global profiling of SRP interaction with nascent polypeptides.
Nature 536:219–223.

55. Bornemann T, Jöckel J, Rodnina MV, Wintermeyer W. 2008. Signal sequence-
independent membrane targeting of ribosomes containing short nascent
peptides within the exit tunnel. Nat Struct Mol Biol 15:494–499.

56. Mohler K, Ibba M. 2017. Translational fidelity and mistranslation in the cellular
response to stress. Nat Microbiol 2:17117.

57. Phillips GJ, Silhavy TJ. 1992. The E. coli ffh gene is necessary for viability and
efficient protein export. Nature 359:744–746.

58. Kikuchi S, Shibuya I, Matsumoto K. 2000. Viability of an Escherichia coli pgsA

null mutant lacking detectable phosphatidylglycerol and cardiolipin. J Bacteriol

182:371–376.

59. Gopalakrishnan AS, Chen YC, Temkin M, Dowhan W. 1986. Structure and

expression of the gene locus encoding the phosphatidylglycerophosphate

synthase of Escherichia coli. J Biol Chem 261:1329–1338.

60. Malecki M, Bárria C, Arraiano CM. 2014. Characterization of the RNase R

association with ribosomes. BMC microbiology 14:34.

61. Quan S, Zhang N, French S, Squires CL. 2005. Transcriptional polarity in rRNA

operons of Escherichia coli nusA and nusB mutant strains. J Bacteriol

187:1632–1638.

62. Wiśniewski JR, Mann M. 2012. Consecutive proteolytic digestion in an enzyme

reactor increases depth of proteomic and phosphoproteomic analysis. Anal

Chem. 84:2631–2637.

63. Wiśniewski JR, Zougman A, Nagaraj N, Mann M. 2009. Universal sample

preparation method for proteome analysis. Nature Methods 6:359–362.

64. Tsolis KC, Economou A. 2017. Quantitative Proteomics of the E. coli

Membranome. Methods Enzymol 586:15–36.

65. Jylhä A, Nättinen J, Aapola U, Mikhailova A, Nykter M, Zhou L, Beuerman R,

Uusitalo H. 2018. Comparison of iTRAQ and SWATH in a clinical study with

multiple time points. Clin Proteomics 15:24.

66. Callister SJ, Barry RC, Adkins JN, Johnson ET, Qian WJ, Webb-Robertson BJ,

Smith RD, Lipton MS. 2006. Normalization approaches for removing

systematic biases associated with mass spectrometry and label-free proteomics.

J Proteome Res 5:277–286.

67.   O'Connor M, Thomas CL, Zimmermann RA, Dahlberg AE. 1997. Decoding

fidelity at the ribosomal A and P sites: influence of mutations in three different

regions of the decoding domain in 16S rRNA. Nucleic Acids Res 25:1185–1193.

68.   Hecht A, Glasgow J, Jaschke PR, Bawazer LA, Munson MS, Cochran JR, Endy

D, Salit M. 2017. Measurements of translation initiation from all 64 codons in

E. coli. Nucleic acids research 45:3615–3626.

69.   Dai X, Zhu M, Warren M, Balakrishnan R, Okano H, Williamson JR, Fredrick

K, Hwa T. 2018. Slowdown of Translational Elongation in Escherichia coli

under Hyperosmotic Stress. mBio 9:e02375-17.

70.   Zarai Y, Margaliot M, Tuller T. 2013. Explicit expression for the steady-state

translation rate in the infinite-dimensional homogeneous ribosome flow model.

IEEE/ACM Trans Comput Biol Bioinform 10:1322–1328.

71.   Zarai Y, Margaliot M, Tuller T. 2014. Maximizing Protein Translation Rate in

the Ribosome Flow Model: The Homogeneous Case. IEEE/ACM Trans Comput

Biol Bioinform 11:1184–1195.

72.   Chen I, Howarth M, Lin W, Ting AY. 2005. Site-specific labeling of cell surface

proteins with biophysical probes using biotin ligase. Nat Methods 2:99–104.

FIGURES

FIG 1 Isolation SRP suppressor strain and its growth. (A) Growth curves and growth

rates of wild-type strain MG1655 and suppressor strain MY1901. (B) The suppressor

mutation sites in the SD sequence of S10 operon. The region and encoded ribosomal

proteins of the S10 operon are shown. The wild-type SD sequence (GGAG) mutated

into suppressor mutation (GAAG). (C) Restoration of the wild-type alleles. Plasmid

pTrc99K-Ffh was first transformed into strain MY1901, generating the strain MY1901F.

Ffh protein was expressed with the basal level in MY1901 to reduce the toxicity of

overexpression of Ffh. Then, the suppressor mutation (GAAG) was restored to wild-

type SD sequence (GGAG) by site-directed mutagenesis, generatiing the strain

MY1901FS. (D) Growth curves and growth rates of strains restored to the wild-type.

Strains MG1655 and MY1901 carrying the empty vector (EV) pTrc99K were used as

control. Solid curves are the mean of three independent measurements and error bars

represent the standard error of the mean of samples. Growth rates were figured out from

751　the exponential growth phase. All growth rates shown represent the mean growth rates

752　from three biological replicates, and error bars represent the standard error of the mean

753　value.

FIG 2 Effects of SRP suppressor on ribosomal protein synthesis. (A) Effect of mutant

SD sequence on protein translation. GFP was used as a reporter to measure gene

expression. (Top) Graphical representation of GFP expression. P_{S10} / P_{araBAD} -GFP with

wild-type SD and suppressor mutation SD* were introduced into cells. (Bottom)

Fluorescence measurement of GFP. Data represent the mean of six independent

experiments, and error bars represent the standard error of the mean value. (B) Fold

changes in the expression of ribosomal proteins in strain MY1901relative to that in

strain MG1655 (Data set S1B).

FIG 3 Effects of SRP suppressor on translation efficiency. (A) Polysome profiles of wild-type strain MG1655 and suppressor strain MY1901Δ*cat*. Data represent the mean ± SEM of three independent experiments. (B) Fold changes in the expression of translation associated factors in strain MY1901relative to that in strain MG1655 (Data set S1B). (C-D) Effect of suppressor mutation on stop-codon (C) and frameshifting (D) readthrough. β-galactosidase activities of strains harboring the mutant LacZ reporters were calculated relative to that of the same strain harboring the wild-type LacZ. (E)The translation elongation rates of FusA-LacZα and MsbA-LacZα under different growth rates (MG1655Δ*lacZ*, 1.0 h⁻¹; MG1655Δ*lacZ*, 0.6 h⁻¹; MY1901Δ*lacZ*Δ*cat*, 0.6 h⁻¹). (F) Translation initiation rates of proteins were estimated based on their corresponding translation elongation rates. (G) Initiation with various non-canonical start codons and non-initiator tRNAs codons in the wild-type and suppressor strains. Relative activities of mutant GFP were calculated relative to that of the same strain harboring the wild-type GFP. The wild-type start codon was indicated as AUG and the wild-type initiator

795 tRNA was indicated as AUG: CAU. All data are averages of six independent

796 experiments, and error bars represent the SEM value.

FIG 4 Suppressor mutation suppresses targeting defects of partial inner membrane

proteins. (A) Scanning electron microscopy (SEM) and transmission electron

microscopy (TEM) analysis of wild-type strain MG1655 and suppressor strain MY1901.

For SEM, the scale bar is 1.0 μm. For TEM, the scale bar is 500 nM. (B) Quantification

of identified SRP-dependent inner membrane proteins in strains MY1901 and SRP⁻

(Data set S1D). Protein abundance of MY1901 and SRP⁻ is relative to that of wild-

type MG1655. (C) FtsQ (left) and EspP (right) targeting assay by their biotinylation.

SRP⁺, Ffh expression in HDB51 strain; SRP⁻, Ffh depletion in HDB51 strain.

825 **FIG 5** The SRP suppressor extends the time window for protein targeting. In wild-type

826 MG1655 cells, although the critical length (L) of inner membrane proteins is variable,

827 the targeting time (T) is fast (~2 s). The suppressor cells with a slower elongation rate

828 extended the time window for protein targeting to 2-10 s. For details and references,

829 see Discussion.