Interplay between intrinsic reprogramming potential and microenvironment controls

neuroblastoma cell plasticity and identity

Authors: Cécile Thirant[1,2§], Agathe Peltier[1,2§], Simon Durand[1,2#], Amira Kramdi[1,2#], Caroline Louis-

Brennetot[1,2#], Cécile Pierre-Eugène[1,2#], Ana Costa[1,2], Amandine Grelier[1,2], Sakina Zaïdi[1,2], Nadège

Gruel[1,4], Irène Jimenez[2,3,4], Eve Lapouble[5], Gaëlle Pierron[5], Hervé J. Brisse[6], Arnaud Gauthier[7], Paul

Fréneaux[7], Sandrine Grossetête-Lalami[1,2], Laura G. Baudrin[8], Virginie Raynal[1,8], Sylvain Baulande[8],

Angela Bellini[2,3,4], Jaydutt Bhalshankar[2,3,4], Angel M. Carcaboso[9], Birgit Geoerger[10], Hermann

Rohrer[11], Didier Surdez[1,2], Valentina Boeva[12,13,14], Gudrun Schleiermacher[2,3,4], Olivier Delattre[1,2],

Isabelle Janoueix-Lerosey[1,2*]

Affiliations :

[1]Institut Curie, PSL Research University, Inserm U830, Equipe Labellisée Ligue contre le Cancer, Paris, France.

[2]SIREDO: Care, Innovation and Research for Children, Adolescents and Young Adults with Cancer, Institut Curie,

Paris, France.

[3]Institut Curie, Laboratory Recherche Translationnelle en Oncologie Pédiatrique (RTOP), Laboratoire "Gilles

Thomas", Paris, France.

[4]Institut Curie Department of Translational Research, Paris, France.

[5]Institut Curie, Unité de Génétique Somatique, Paris, France.

[6]Institut Curie, Department of Imaging, PSL Research University, Paris, France.

[7]Institut Curie, Department of Biopathology, Paris, France.

[8]Institut Curie, Genomics of Excellence (ICGex) Platform, Paris, France.

[9]Institut de Recerca Sant Joan de Déu, Barcelona, Spain.

[10]Gustave Roussy Cancer Campus, INSERM U1015, Department of Pediatric and Adolescent Oncology, Univ.

Paris-Sud, Université Paris-Saclay, Villejuif, France.

[11]Institute of Clinical Neuroanatomy, Dr. Senckenberg Anatomy, Neuroscience Center,

Goethe University, Frankfurt/M, Germany

[12]Inserm, U1016, Cochin Institute, CNRS UMR8104, Paris Descartes University, Paris, France.

[13]ETH Zürich, Department of Computer Science, Institute for Machine Learning, Zürich, Switzerland.

[14]Swiss Institute of Bioinformatics (SIB), Zürich, Switzerland.

[§,#]These authors contributed equally to this work.

* Corresponding author and lead contact: janoueix@curie.fr

Abstract

Two cell identities, noradrenergic and mesenchymal, have been characterized in neuroblastoma cell

lines according to their epigenetic landscapes relying on specific circuitries of transcription factors.

Yet, their relationship and relative contribution in patient tumors remain poorly defined. Here, we

demonstrate that the knock-out of *GATA3*, but not of *PHOX2A* or *PHOX2B*, in noradrenergic cells

induces a mesenchymal phenotype. Our results document spontaneous plasticity in several models

between both identities and show that plasticity relies on epigenetic reprogramming. We

demonstrate that an *in vivo* microenvironment provides a powerful pressure towards a

noradrenergic identity for these models. Consistently, tumor cells with a mesenchymal identity are

not detected in a series of PDX models. Further study of the intra-tumor noradrenergic

heterogeneity reveals two distinct cell populations exhibiting features of chromaffin-like or

sympathoblast-like cells. This work emphasizes that both external cues of the environment and

intrinsic factors control plasticity and cell identity in neuroblastoma.

INTRODUCTION

Neuroblastoma is a childhood cancer arising from the peripheral sympathetic nervous system,

known to be derived from multipotent neural crest cells (NCCs). Tumors mostly develop in the adrenal

gland but a subset of them originates from sympathetic ganglia along the paravertebral sympathetic

chain[1]. With respect to these localizations, neuroblastoma likely arises from the transformation of

sympathoblasts either in sympathetic ganglia or in the adrenal medulla, or from catecholamine-

secreting chromaffin cells of the adrenal medulla, or alternatively from a common sympatho-adrenal

progenitor[2,3].

The hallmark of neuroblastoma is its wide range of clinical presentations and outcomes,

ranging from spontaneous regression to fatal outcome despite multimodal therapies[1]. High-risk

neuroblastoma most often initially responds to intensive chemotherapy; however, relapses frequently

occur followed by fatal outcome. Several genes including *MYCN*[4], *ALK*[5–8] and *TERT*[9–11] have been

identified as key drivers of neuroblastoma oncogenesis.

The master transcriptional regulators controlling the gene expression program of

neuroblastoma have been recently highlighted through the characterization of the neuroblastoma

super-enhancer landscape of neuroblastoma cell lines, revealing two distinct cell identities: a

sympathetic noradrenergic identity defined by a core regulatory circuitry (CRC) module including the

PHOX2B, HAND2 and GATA3 transcription factors (TFs) and a NCC-like/mesenchymal identity, close to

that of human neural crest cells (NCCs), driven by TFs of the AP1 family among others[12,13]. Additional

TFs participating in the noradrenergic CRC have been then identified, including ISL1, TBX2 and ASCL1[14–

16]. It is likely that PHOX2A, also highly expressed in neuroblastoma cell lines and exhibiting the same

DNA binding domain as PHOX2B is involved in the noradrenergic CRC. Importantly, mesenchymal

tumor cells *in vitro* have been shown to be more resistant to standard chemotherapy[12,13] suggesting

that they may be involved in therapeutic resistance and relapses in neuroblastoma patients. Bulk RNA-

seq analyses or immunohistochemistry with few markers suggested that mesenchymal tumor cells are

present in patient tumors and that some tumors exhibit a mesenchymal identity[12,13,17]. Cellular

plasticity between the noradrenergic and mesenchymal states has been reported for a few cell

lines[13,18], still the underlying mechanisms wherein the cell phenotype switches from one to the other

state (considered as transdifferentiation) remain poorly described. In the present paper, we have

combined various approaches including genetic inactivation of specific TFs and single-cell

transcriptomic analyses to unravel TFs involved in cell identity and plasticity and better decipher the

relationship and relative contribution of cells of noradrenergic and mesenchymal identity in

neuroblastoma tumors.

RESULTS

The knock-out of *GATA3* but not of *PHOX2B* or *PHOX2A* induces a switch from a

noradrenergic to a mesenchymal identity

In order to directly address the functional role of the PHOX2A, PHOX2B and GATA3 TFs in

shaping the noradrenergic identity, we performed their individual genetic knock-out (KO) in the

noradrenergic SH-SY5Y cell line using a CRISPR-Cas9 approach. Guide RNAs were designed to induce

large deletions and create frameshift mutations in those genes, leading to truncated and nonfunctional

proteins. We obtained two *PHOX2A*$^{-/-}$ clones, a *PHOX2B*$^{-/-}$ clone and two *GATA3*$^{-/-}$ clones (**Figure S1A**).

While the *PHOX2A*$^{-/-}$ and *PHOX2B*$^{-/-}$ clones showed a morphology close to that of the parental

noradrenergic SH-SY5Y cells with neurite-like processes, *GATA3*$^{-/-}$ cells exhibited a more abundant

cytoplasm and many actin stress fibers consistent with a mesenchymal phenotype (**Figure 1A**).

PHOX2A$^{-/-}$ and *PHOX2B*$^{-/-}$ clones maintained the expression of the noradrenergic CRC TFs including

GATA3 and HAND2 (**Figures 1B and S1B**). Strikingly, both *GATA3*$^{-/-}$ clones showed an absence or highly

reduced expression of the TFs from the noradrenergic CRC, including PHOX2A, PHOX2B, and HAND2

(**Figure 1B**). *PHOX2B*$^{-/-}$ and *GATA3*$^{-/-}$ cells were characterized by a decreased proliferation compared to

the parental SH-SY5Y cell line (**Figure 1C**). Consistently with their phenotype, *GATA3*$^{-/-}$ clones displayed

mesenchymal properties such as an increased invasion ability measured in a 3D-spheroid assay (**Figure**
1D), increased migration capacity documented using a transwell assay (**Figure 1E**) and a higher
resistance to chemotherapy *in vitro* (**Figure 1F**). Bulk RNA-seq analysis showed that *PHOX2A*[-/-] and
PHOX2B[-/-] clones harbored transcriptomic profiles highly similar to that of the parental noradrenergic
SH-SY5Y cell line, confirming that *PHOX2A* or *PHOX2B* genetic invalidation did not change the cell
identity. In contrast, *GATA3*[-/-] clones, at two different passages called early and late showed a
mesenchymal transcriptomic profile, as they clustered with the mesenchymal SH-EP cell line (**Figures**
1G and S1C). Accordingly, Gene Ontology analysis on the 2,938 differentially expressed genes between
the transcriptomic profiles of the 4 *GATA3*[-/-] samples versus 5 noradrenergic neuroblastoma cell lines
(without *MYCN* amplification as the parental SH-SY5Y) retrieved categories related to neuron
differentiation, neurogenesis, sympathetic nervous development for the downregulated genes and
extracellular matrix organization, cell motility and cell migration for the upregulated genes (**Figure**
S1D).

These observations therefore demonstrated that the KO of the *PHOX2A, PHOX2B* or *GATA3*
transcription factors has different consequences on the maintenance of the noradrenergic identity and
that SH-SY5Y cells are able to transdifferentiate from a noradrenergic to a mesenchymal state upon
GATA3 KO.

Spontaneous plasticity between the noradrenergic and mesenchymal states reveals the
reprogramming potential of a subset of neuroblastoma cells

Our previous study has shown that most of the neuroblastoma cell lines (18 out of 25) exhibit
a noradrenergic identity whereas only 3 have a mesenchymal epigenetic profile. An intermediate
group of cells expressing noradrenergic TFs and mesenchymal TFs was composed of 4 samples,
including the heterogeneous SK-N-SH cell line[12]. Interestingly, plasticity properties have been reported
for this cell line[18]. Using single-cell transcriptomic sequencing with the 10X Genomics technology, we
highlight here two cell populations in the SK-N-SH cell line. Noradrenergic cells expressed *PHOX2B*
whereas *CD44*[19–21] appeared as a specific marker of the mesenchymal population confirmed by FACS
and immunofluorescence (**Figure 2A-C**). An InferCNV analysis that predict the genetic alterations at
the single-cell level[22], showed that both populations exhibited similar genetic alterations (2p, 7 and
17q gains). Only one subcluster of the mesenchymal population presented with a 1q gain (**Figure 2D**),
consistently with previous data reporting distinct genetic subclones[23]. As CD44 is a cell surface marker
it was further used to sort both populations. Bulk RNA-seq experiments confirmed that CD44[neg] and
CD44[pos] sorted cells exhibited transcriptomic profiles close to noradrenergic SH-SY5Y and

mesenchymal SH-EP cells, respectively (**Figure 2E**). Both sorted noradrenergic/CD44[neg] and

mesenchymal/CD44[pos] populations were able to give rise to a heterogeneous cell population,

demonstrating a spontaneous and bidirectional plasticity between the noradrenergic and

mesenchymal states (**Figure 2F**). As expected from previous data obtained with the SH-SY5Y and SH-

EP cell lines sub-cloned from the heterogeneous parental SK-N-SH cell line[12], the

mesenchymal/CD44[pos] population of this cell line exhibited a higher chemo-resistance compared to

the noradrenergic/CD44[neg] one (**Figure 2G**).

In order to demonstrate that plasticity between the noradrenergic and mesenchymal states is

not a property exclusively observed in the SK-N-SH sample, we generated new cell lines from a series

of 10 neuroblastoma PDX models. Cells from two models could be maintained in culture for several

months and frozen. One cell line had a pure noradrenergic phenotype. Strikingly, the other cell line

called IC-pPDXC-63 was able to grow *in vitro* as a bi-phenotypic culture, with both adherent cells and

floating neurospheres (**Figure 3A**). Bulk RNA-seq analysis confirmed that the original PDX model (IC-

pPDX-63) and its derived-cell line exhibited a transcriptomic profile highly similar to the one of the

patient tumor (NB1549) (**Figure 3B**). As shown by single-cell analysis, two main clusters of

noradrenergic cells and mesenchymal tumor cells were observed in the IC-pPDXC-63 cell line, with a

bridge of cells in-between (**Figure 3C**). Noradrenergic cells expressed *PHOX2B*, whereas the

mesenchymal population expressed the cell surface CD44 marker (**Figure 3C**). The clustering

separating noradrenergic and mesenchymal tumor cells is not biased by the cell cycle (**Figure 3C**).

Immunofluorescence confirmed that CD44 and PHOX2B were specifically expressed by adherent cells

and neurospheres, respectively (**Figure 3D**). Bulk RNA-seq analysis endorsed that CD44[pos] FACS-sorted

cells and adherent cells have a transcriptomic profile close to the mesenchymal SH-EP cells, whereas

CD44[neg] FACS-sorted cells and floating neurospheres clustered with the noradrenergic SH-SY5Y cells

(**Figure 3B**). Consistently with the observations in the SK-N-SH cell line, we documented that the

mesenchymal population of the IC-pPDXC-63 cell line exhibited a higher chemo-resistance compared

to the noradrenergic one (**Figure 3E**). Of note, inferred genomic alterations were similar in the three

populations of the IC-pPDXC-63 cell line, *i.e.*, the noradrenergic, mesenchymal and bridge cells (**Figure**

3F). We next investigated the plasticity properties of both noradrenergic and mesenchymal cells of the

IC-pPDXC-63 cell line *in vitro*. After a few days in culture, both sorted noradrenergic/CD44[neg] and

mesenchymal/CD44[pos] cells were able to reconstitute a heterogeneous cell population (**Figure 3G**), as

previously documented for the SK-N-SH cell line. These observations therefore highlight the

reprogramming potential between the noradrenergic and mesenchymal states of the IC-pPDXC-63

model and importantly, show that this ability does not rely on genetic heterogeneity.

Phenotypic plasticity relies on epigenetic reprogramming

To deeper characterize the epigenetic reprogramming contribution to cell plasticity, we next defined the super-enhancer landscape of our three different models (*GATA3* genetic inactivation, SK-N-SH and IC-pPDXC-63 cell lines) by ChIP-seq analyses for the H3K27ac mark. We added these samples in the principal component analysis (PCA) of neuroblastoma cell lines and hNCC lines based on their super-enhancer log scores[12] (**Figure 4A**). For all three models, their mesenchymal counterparts, *i.e.* *GATA3*$^{-/-}$ clones, SK-N-SH CD44^{pos} FACS-sorted and adherent IC-pPDXC-63 cells, showed an epigenetic profile close to the group II of mesenchymal identity. Consistently, their noradrenergic counterparts, SH-SY5Y, SK-N-SH CD44^{neg} FACS-sorted and floating IC-pPDXC-63 cells were part of the noradrenergic cell line group I (**Figure 4A**). For the mesenchymal cells of the three models, a decrease of the H3K27ac signal could be quantified for the super-enhancer regions of the noradrenergic CRC such as *GATA3*, *PHOX2B* and *HAND1*. On the other hand, an increase of the H3K27ac signal could be observed on some NCC-like/mesenchymal TFs such as *RUNX1*, *FOSL1*, *NR3C1,* and *TBX18* (**Figure 4B,C**). Of note, the decreased PHOX2A and HAND2 protein expressions in the *GATA3*$^{-/-}$ clones (**Figure 1B**) contrasted with high transcript levels (**Figure 1G**) and high H3K27ac scores (**Figure 4B**) suggesting a distinct transcriptional and post-transcriptional regulation for these specific genes.

Altogether, these results indicate that the transdifferentiation from a noradrenergic to a mesenchymal identity obtained after the genetic inactivation of *GATA3* or spontaneously from SK-N-SH and IC-pPDXC-63 cells is supported by an epigenetic reprogramming.

Mesenchymal neuroblastoma cells are reprogrammed to a noradrenergic phenotype *in vivo*

Since noradrenergic and mesenchymal cells of our three models presented with different properties *in vitro*, we next investigated their behavior *in vivo*. We injected the noradrenergic/CD44^{neg} and mesenchymal/CD44^{pos} sorted cell populations from the two heterogeneous SK-N-SH and IC-pPDXC-63 cell lines, as well as the *GATA3*$^{-/-}$ cells, into mice. Tumors developed in all cases, indicating that the different states displayed tumorigenic potential *in vivo*. Unexpectedly, as revealed by IHC analysis, PHOX2B expression was observed in most tumor cells from the whole set of xenografts, even those obtained after engraftment of mesenchymal populations (**Figure 5A**). Bulk RNA-seq experiments confirmed that all tumors highly expressed the TFs of the noradrenergic CRC and exhibited a noradrenergic transcriptomic profile (**Figures 5B and S2**). Xenografts of the *GATA3*$^{-/-}$ clones re-expressed the noradrenergic and neuroendocrine *DBH, NET/SLC6A2, CHGA* and *CHGB* markers compared to the clones cultured *in vitro* (**Table S1**). Consistently, PHOX2B and DBH proteins were detected in the *GATA3*$^{-/-}$ cell xenografts (**Figure 5C**). Additionally, these markers were similarly

expressed in xenografts obtained with the two CD44[pos] and CD44[neg] cell populations of the SK-N-SH and IC-pPDXC-63 cell lines (**Table S1**). Differential analyses (Fold-change >2; Bonferroni-corrected p-value <0.05) showed that xenografts obtained from CD44[pos] and CD44[neg] cells of both heterogeneous cell lines were highly similar, with less than 50 genes (data not shown) showing a differential expression, compared to more than 2,000 genes differentially expressed *in vitro*. Of note, some genes of the neurogenesis and extracellular matrix organization were still differentially expressed between SH-SY5Y and *GATA3[-/-]* cell xenografts, but of lower magnitude compared to the difference between the corresponding cells *in vitro*. Finally, to fully demonstrate the *in vivo* reprogramming of mesenchymal cells towards a noradrenergic identity, we determined the super-enhancer profiles of SH-SY5Y and *GATA3[-/-]* cell xenografts and additionally from the xenografts of SK-N-SH CD44[pos] and CD44[neg] FACS-sorted cells. The PCA clearly showed that all these tumors were part of the noradrenergic group I (**Figure 5D**). Super-enhancers marked *PHOX2B* and *HAND1* genes in xenografts of SH-SY5Y and SK-N-SH CD44[neg] cells but also in xenografts of *GATA3[-/-]* and SK-N-SH CD44[pos] cells (**Figure 5E**). In an *in vivo* environment, the mesenchymal cells therefore shifted back towards a noradrenergic identity indicating that the mouse microenvironment provided strong cues inducing a global epigenetic reprogramming.

Altogether, our analyses highlighted the plasticity properties of neuroblastoma cells using three different models (*GATA3* genetic inactivation, SK-N-SH and IC-pPDXC-63 cell lines). This plasticity was associated with a reprogramming potential from a mesenchymal state to a noradrenergic state following *in vivo* engraftment in the mouse.

Single-cell transcriptomic analyses reveal intra-tumor noradrenergic heterogeneity but no mesenchymal tumor cells in neuroblastoma PDX models

Since heterogeneity of cell identity has been observed in some neuroblastoma cell lines, we took advantage of the 10X Genomics technology to explore neuroblastoma intra-tumor heterogeneity using single-cell transcriptomic analyses on 14 PDX models, obtained at diagnosis, progression or at relapse (**Figure 6A, Table S2**). The study was designed to specifically identify human tumor cells (see methods). The integration of the human cells from all models (n=47,219 cells) with the Harmony tool[24] highlighted tumor cells of noradrenergic identity (*PHOX2B+*, *HAND2+*) (**Figures 6B-C** and **S3A**). No cluster of mesenchymal tumor cells could be identified as shown by the analysis of previously published mesenchymal signatures (**Figure 6D**). The InferCNV analyses confirmed that noradrenergic tumor cells exhibited the emblematic genetic alterations of neuroblastoma such as 17q gain[1] (**Figure S3B-C**). For several models, we could compare the InferCNV profile calculated from scRNAseq data to

copy number profiles inferred from whole-exome sequencing data and confirmed their full consistency (**Figure S3B**). We next assessed the heterogeneity of the human tumor cell populations and documented that most clusters were shared by all PDX models (**Table S4**). Several populations could be defined according to the differential expression of specific genes: cycling cells marked by the expression of *TOP2A, MKI67* and *CDK1* (clusters 2 and 6), cells driven by a *MYCN/*2p amplicon signature (clusters 0-1-2-3-13), cells expressing chromaffin markers such as *CDKN1C*, *SLC18A1*/VMAT1[25,26] and *DLK1*[27] (clusters 9-10-11) and cells with a sympathoblast-like identity such as *TFAP2B*[28] (cluster 7) (**Figures 6E, S3D-F** and **Table S4**).

Interestingly, our single-cell transcriptomic data revealed that the GR-NB7 PDX model did not express *GATA3* but expressed *PHOX2B*, *CHGA* and *DDC* (**Figure 6F).** These data confirmed that *GATA3* was dispensable to establish a noradrenergic identity *in vivo*, in agreement with the noradrenergic shift of *GATA3*[-/-] cells observed in mouse xenografts. Unsupervised hierarchical clustering using the 10% genes with the highest IQR on bulk RNA-seq data confirmed that all studied PDX models exhibited a transcriptomic profile corresponding to a noradrenergic identity (**Figure S3F**).

Altogether, the present analysis of 24 neuroblastoma samples identified no *bona fide* tumor mesenchymal cells, which is consistent with our aforementioned results demonstrating a strong pressure of the microenvironment towards a noradrenergic identity.

DISCUSSION

In the present work, we have first deciphered the distinct roles of the members of the noradrenergic CRC in shaping neuroblastoma cell identity. Whereas *Phox2b* is essential for initial sympatho-adrenal cell specification from neural crest progenitors[29–31], the KO of *PHOX2B* or *PHOX2A* did not modify the noradrenergic identity of SH-SY5Y cells. Yet, one hypothesis is functional redundancy between PHOX2A and PHOX2B. Of note, we did not succeed in obtaining double KO clones suggesting synthetic lethality for the cells. In contrast, the KO of *GATA3* in the same cells had a dramatic effect, inducing the collapse of the noradrenergic CRC. Surprisingly, whereas *GATA3* KO clones harbored mesenchymal properties *in vitro*, they reacquired a noradrenergic identity *in vivo* indicating that *GATA3* KO allows a permissive epigenetic state and the transdifferentiation towards a mesenchymal or noradrenergic state according to non-autonomous cues. Consistently with these observations, we could document a noradrenergic identity of one neuroblastoma PDX model characterized by an absence of GATA3 expression. In a previous study using siRNA and short-term transcriptomic analysis by RT-q-PCR, Durbin *et al*. reported that the knock-down of only one member of the CRC was able to induce a decrease of the expression of several members of the CRC[15]. Yet,

residual expression remained for the different TFs studied and the impact of this decrease on cell

identity has not been explored. Therefore, our results constitute the first evidence that TFs of a specific

CRC may have distinct roles in shaping cell identity in neuroblastoma.

Transdifferentiation from the noradrenergic to the mesenchymal identity has been shown

previously *in vitro* following overexpression of the PRRX1 TF[13]. The same team also recently pointed

out a role of the NOTCH signaling pathway in this process[32]. Of note, xenografts of SH-SY5Y cells

overexpressing NOTCH3 intracellular domain exhibited a mesenchymal identity. In this model, the

permanent production of intra-cellular NOTCH3 likely activates an endogenous feed-forward loop

between NOTCH receptors and ligands *in vivo.* Although we observed expression of several members

of this pathway (*NOTCH2*, *MAML2*, *HES1*) in our mesenchymal tumor cells *in vitro*, in our series of PDX

models, only expression of *MAML2* could be detected in some tumor cells; yet these cells were not

associated with a specific cluster and very spare expression was observed for *NOTCH2* and *HES1* in the

whole population of tumor cells*.*

We document here the heterogeneity and spontaneous reprogramming potential of cells of

IC-PDXC-63 and SK-N-SH models from a noradrenergic to a mesenchymal identity and conversely.

Single-cell transcriptomic analysis performed on these models first identified CD44 as a surface marker

specific of the mesenchymal identity, further allowing the use of this marker to sort each population

and analyze their respective proportions by FACS. We show that the spontaneous and bi-directional

plasticity observed *in vitro* relies on a profound epigenetic reprogramming, as revealed by the analysis

of the super-enhancer landscape of the various cell populations. Very strikingly, we demonstrate that

mesenchymal tumor cells from three different models exhibiting plasticity revert their identity towards

a noradrenergic state when engrafted in mice. These results highlight that neuroblastoma cell

phenotype is strongly influenced by the *in vivo* microenvironment which provides a powerful pressure

towards a noradrenergic state. This conclusion is reinforced by the observation that mesenchymal

tumor cells were not identified by single-cell transcriptomic analyses in our series of 14 neuroblastoma

PDX models, obtained either at diagnosis or at relapse and presenting various genetic alterations.

Altogether, our data obtained on a variety of cellular models and PDXs provide a biological explanation

for the absence of mesenchymal tumor cells *in vivo*. This observation is also in line with the very recent

study performed by Dong and colleagues that identified normal, but not tumor, mesenchymal cells in

adrenal neuroblastoma tumors by single-cell RNA sequencing[33]. Interactions between tumor cells and

several cell populations of the microenvironment may influence the tumor cell phenotype and play a

role in tumor progression, as previously demonstrated for tumor-associated inflammatory cells[34].

Future studies should decipher the cues of the microenvironment and their associated pathways that

converge to regulate cell plasticity during tumor progression.

Interestingly, some evidences suggested that treatments against neuroblastoma may impact
cell identity. Indeed, previous analyses of neuroblastoma cells selected to be resistant to cisplatin[35] or
ALK inhibitors[36] *in vitro* have reported that noradrenergic cells may acquire mesenchymal properties.
Cells exhibiting such a potential may be transiently induced upon chemotherapy treatment and revert
their state towards a noradrenergic identity when the treatment pressure decreases, the
mesenchymal cells being the drug-resistant reservoirs for noradrenergic cells. Further experiments will
allow better defining their contribution to therapeutic resistance and relapse in high-risk
neuroblastomas.

The cell of origin in neuroblastoma has been for long a matter of debate[37–39]. Some models
argue that this tumor arises from the transformation of NCCs while other models suggest that they
develop from more engaged sympatho-adrenal progenitors, able to generate both sympathetic
neurons and neuroendocrine cells of the adrenal. Recently, this hierarchical dogma of normal
differentiation has been questioned with the identification of a population of Schwann cell precursors
(SCPs) as the main reservoir of adrenal chromaffin cells in the mouse[26]. In their recent paper, Dong et
al. who analyzed human fetal adrenal gland at the single-cell level, in addition to adrenal
neuroblastoma tumors, concluded that malignant cells had a predominant chromaffin-cell-like
phenotype[33]. We nevertheless disagree with their interpretation of cell phenotypes. We strongly
believe that *CARTPT* and *INSM1,* used by Dong et al. to define sympathoblast identity, rather witness
a chromaffin-like phenotype, since they are co-expressed with *DLK1*, *CDKN1C*, *CHGA*/*CHGB*, *TH*, *DBH*
and *SLC18A1* in our data [25–27,40–44]. Integration of our single-cell RNA-seq data from 14 PDX models
revealed two distinct cell populations expressing either markers of chromaffin cells or markers of
sympathoblasts. Of note, SCP markers such as *SOX10, S100B, PLP1* and *ERBB3*[26] were not detected in
our cohort of 14 neuroblastoma cases. Nevertheless, it remains difficult to infer which cell type is the
one targeted by neoplastic transformation. Indeed, it cannot be excluded that transitions may occur
between chromaffin cells and sympathoblasts during development and/or that markers of a specific
precursor may be lost during cell transformation.

Altogether, our data obtained on several cellular models demonstrate that a subset of
neuroblastoma cells exhibits a reprogramming potential between a noradrenergic and a mesenchymal
identity and that both intrinsic properties and exogenous signals of the microenvironment dictate this
identity. A better understanding of the molecular factors that control phenotypic plasticity will
represent a key step in the design of more efficacious therapies that aim at improving the outcome of
neuroblastoma patients with high-risk disease.

ACKNOWLEDGEMENTS

This work was supported by grants from Institut Curie, Inserm, the Ligue Nationale Contre le Cancer (Equipe labellisée), the Institut National du Cancer (PLBIO18-273) and by the following associations: Association Hubert Gouin-Enfance et Cancer, Les Bagouz à Manon, les amis de Claire, Courir pour Mathieu, Dans les pas du Géant, Olivier Chape. The Mappyacts protocol is supported by the Institut National de Cancer (INCa) through the PHRC "INCa-DGOS_8519" MERRI, the Fondation ARC, the Association Imagine for Margo, the Société Française de lutte contre les Cancers et les leucémies de l'Enfant et l'adolescent (SFCE), Fédération Enfants & Santé, the associations AREMIG and Thibault BRIET. The Micchado protocol is supported by PRT-K, Association Imagine for Margo, Kickcancer, Hubert Gouin-Enfance et Cancer, Fédération Enfants & Santé, and funding support by BMS and Roche. High-throughput sequencing has been performed by the ICGex NGS platform of the Institut Curie supported by the grants ANR-10-EQPX-03 (Equipex) and ANR-10-INBS-09-08 (France Génomique Consortium) from the Agence Nationale de la Recherche ("Investissements d'Avenir" program), by ITMO Cancer Aviesan ("Equipement pour la recherche en Cancérologie" program) and by the SiRIC-Curie program -SiRIC Grant "INCa-DGOS- 4654". G.S. is supported by the Annenberg foundation. H.R. is supported by the Wilhelm-Sander-Stiftung. We are grateful to the animal facilities team, the Experimental Pathology Department, the Plateforme Génomique and the Plateforme Cytométrie of Institut Curie, U900 colleagues for help with alignment of NGS data and Julien Masliah-Planchon for help with genomic analyses. We thank Mélissa Saichi and Divya Sahu for preliminary analyses of single-cell RNA-seq and ChIP-seq data.

AUTHOR CONTRIBUTIONS

Conceptualization: C.T., V.B., O.D. and I.J.-L.; Methodology: C.T., A.P., S. Z., N.G., I. J., A.M.C., B.G. and D.S.; Formal Analysis: C.T., A.K., A.C., S.G.-L., V.B.; Investigation: C.T., A.P., S.D., C.L.-B., C.P.-E., A.C., A.G., N.G. and H.R.; Resources: E.L., G.P., H.B., A.G., P.F., L.G. B., V.R., S.B., A. B., J.B and G.S.; Writing – Original Draft: C.T., A.P. and I.J.-L.; Writing – Review and Editing: C.T., A.P., C.L.-B., H.R., O.D. and I.J.-L.; Visualization: C.T., A.P., A.K., C.L.-B., C.P.-E.; Supervision: C.T. and I.J.-L.; Project Administration: I.J.-L.; Funding Acquisition: I.J.-L, V. B. and O.D.

DECLARATION OF INTERESTS

The authors declare no potential conflicts of interest.

Figure 1

Figure 1. *GATA3*^{-/-} cells exhibit mesenchymal properties whereas *PHOX2B*^{-/-} or *PHOX2A*^{-/-} cells keep

a noradrenergic phenotype *in vitro*. (A) Phalloidin staining reveals the cell phenotype of the different

KO clones and the SH-SY5Y parental cell line (x400, scale bar = 20 μm). (**B**) Western blot analysis of

GATA3, PHOX2B, PHOX2A, HAND2 and DBH in the *PHOX2B*^{-/-} and *GATA3*^{-/-} clones, with Vinculin as a

loading control. (**C**) Proliferation curves for *PHOX2B*^-/-^ and *GATA3* ^-/-^ cells *in vitro*. 10,000 cells were

seeded in 24-well plates. Living cells were counted after 3, 7 and 10 days of culture (mean ± sd.; n = 6

replicates). (**D**) Representative images (left) and quantification (right) of spheroid invasion assays. Cells

were seeded in low adherent plates to form neurospheres for 4 days and embedded in collagen I for

72h. Scale bar = 100 µm (mean ± sd.; n = 12 replicates for SH-SY5Y and SH-EP cell lines and n = 18

replicates for *GATA3*^-/-^ clones #1 and #2). (**E**) Representative images (left) and quantification (right) of

transwell assays for the SH-SY5Y cells, the *GATA3*^-/-^ clones (#1 and #2) and mesenchymal control cells

(SH-EP). 50,000 cells were plated in a transwell insert and the living cells that have migrated into the

transwell membrane were counted 24h later. Scale bar = 20 µm (mean ± sd.; n = 6 replicates). Each

dot represents a replicate and is the mean of 6 different images of the well. (**F**) *GATA3*^-/-^ cells are more

resistant to chemotherapy than the SH-SY5Y noradrenergic parental cell line. Cell viability was

measured after 48 hours of chemotherapy treatments (Doxorubicin 50, 100, 200 nM and Etoposide

0.5, 1, 2.5 µM) (mean ± sd.; n = 6 replicates). (**G**) Heatmap showing the expression levels of the TFs of

the noradrenergic (n=7) and mesenchymal (n=15) CRC in the *PHOX2B*^-/-^ (clone#1, 3 replicates)*,

PHOX2A*^-/-^ (clones #1 and #2) and *GATA3*^-/-^ (clones #1 and #2). Both *GATA3*^-/-^ clones were analyzed at

two different time points (early (E) and late (L)). The noradrenergic SH-SY5Y and the isogenic

mesenchymal SH-EP cell lines were also included in the analysis. P-values were determined via two-

tailed unpaired Welch's t-test (**: $p<0.01$, ***: $p<0.001$).

Figure 2

Figure 2. Cell plasticity in the heterogeneous neuroblastoma SK-N-SH cell line. (**A**) Single-cell RNA-seq analysis by Seurat. The umap plot shows two main clusters, *PHOX2B* and *CD44* expression distinguish noradrenergic and mesenchymal cells, each identity including cycling cells. (**B**), (**C**) FACS and immunofluorescence analyses of the SK-N-SH cell line with the PHOX2B and CD44 markers (scale bar= 50 µm). (**D**) Genetic alterations were predicted from single-cell RNAseq data using the InferCNV tool. (**E**) Unsupervised clustering using the top 10% of genes with the highest IQR (Inter-Quartile Range) indicates that CD44^neg and CD44^pos sorted cells (two replicates) exhibit a transcriptomic profile close to

the noradrenergic SH-SY5Y and mesenchymal SH-EP cells, respectively. (**F**) CD44^{neg} (upper panel) and

CD44^{pos} (lower panel) sorted cells are able to reconstitute a heterogeneous population of both CD44^{neg}

and CD44^{pos} cells after a few weeks in culture. (**G**) Mesenchymal/CD44^{pos} sorted cells are more resistant

to doxorubicin and etoposide than noradrenergic/CD44^{neg} cells. Cell viability was measured after 72

hours of chemotherapy treatments (Doxorubicin 50, 100, 250 nM and Etoposide 0.5, 1, 2.5 μM)) (mean

± sd.; n = 6 replicates). P-values were determined via two-tailed unpaired Welch's t-test (**: p<0.01;

***: p<0.001).

Figure 3

Figure 3. The IC-pPDXC-63 model exhibits a phenotypic plasticity between noradrenergic and mesenchymal identities. (**A**) *Ex vivo* culture of the noradrenergic IC-pPDX-63 neuroblastoma model includes floating neurospheres and adherent cells. (**B**) Unsupervised clustering using the top 10% of

genes with the highest IQR (Inter-Quartile Range) shows that the transcriptomic profile of the IC-pPDX-

63 model and its derived-cell line (IC-pPDXC-63) are highly similar to that of the patient's tumor

(NB1549) from which it has been generated. (**C**) Single-cell transcriptomic analyses by Seurat highlight

both noradrenergic and mesenchymal clusters in the IC-pPDXC-63 cell line together with a bridge in-

between. Four umap plots are shown (from left to right): the cell clustering, *PHOX2B* expression

marking noradrenergic cells, *CD44* expression marking mesenchymal cells and the cell cycle. (**D**)

Immunofluorescence shows the specific expression of the PHOX2B and CD44 markers by neurospheres

and adherent cells, respectively (scale bar = 20 μm). (**E**) Mesenchymal (IC-pPDXC-63 CD44^{pos}) cells are

more resistant to chemotherapy than noradrenergic (IC-pPDXC-63 CD44^{neg}) cells. Cell viability was

measured with resazurin assay after 72 hours of chemotherapy treatments ((Doxorubicin 50, 100, 250

nM and Etoposide 0.5, 1, 2.5 μM)) (mean ± sd.; n = 6 replicates). P-values were determined via two-

tailed unpaired Welch's t-test (**: $p<0.01$; ***: $p<0.001$). (**F**) Inferred genomic profile of the IC-pPDXC-

63 cell line obtained with InferCNV on single-cell data. (**G**) Plasticity properties of the IC-pPDXC-63 cell

line. CD44^{pos} and CD44^{neg} cells were FACS sorted and put back in culture. Each sorted population

reconstituted a heterogeneous cell population after several days in culture.

Figure 4

Figure 4. Phenotypic plasticity relies on epigenetic reprogramming. (A) Principal Component Analysis

(PCA) based on neuroblastoma and hNCC super-enhancer log scores[12] that discriminates the two

neuroblastoma cell groups I (noradrenergic-NOR) and II (NCC-like/mesenchymal-MES) and in which

were added the two *GATA3*[-/-] clones at early (E) and late (L) time points as well as the floating and

adherent cells of the IC-pPDXC-63 and CD44^{neg} and CD44^{pos} sorted cells of the SK-N-SH cell lines. (**B**) Heatmap showing the H3K27ac signal on the super-enhancer regions of the TFs of the noradrenergic (NOR) and mesenchymal (MES) CRC in the two *GATA3$^{-/-}$* at both early (E) and late (L) time points, in floating and adherent cells of the IC-pPDXC-63 cell line and in the CD44^{pos} and CD44^{neg} sorted cells of the SK-N-SH cell line, and the SH-SY5Y and SH-EP control cell lines. For the TFs associated with several super-enhancers, the signal was summarized as described in the Methods section to have one value per TF. The unsupervised hierarchical clustering based on H3K27ac signals discriminated noradrenergic and mesenchymal TFs and cell identity. (**C**) Tracks of ChIP-seq profiles for H3K27ac at *PHOX2B, HAND1, RUNX1,* and *FOSL1* super-enhancers in the SH-SY5Y, SH-EP, the 4 *GATA3$^{-/-}$* samples (2 at early (E) and 2 at late (L) time-points), the floating and adherent IC-pPDXC-63 cells and the CD44^{pos} and CD44^{neg} sorted SK-N-SH cells.

Figure 5

Figure 5. All sorted mesenchymal cells from the 3 models, SK-N-SH and IC-pPDXC-63 cell lines and *GATA3⁻/⁻* cells, adopt a noradrenergic identity when engrafted in mouse. (**A**) PHOX2B immunohistochemistry of one representative mouse xenograft per group (CD44ⁿᵉᵍ or CD44ᵖᵒˢ sorted cells of SK-N-SH and IC-pPDXC-63 cell lines, SH-SY5Y control cells and *GATA3⁻/⁻* clones #1 and #2). Scale

bar = 50 µm. Similar results were obtained for all analyzed xenografts. (**B**) Heatmap showing the

expression of the TFs associated with the noradrenergic or mesenchymal identity[12] reveals that all

mouse xenografts exhibit a noradrenergic transcriptomic profile. (**C**) Western-blot analysis of GATA3,

PHOX2B, HAND2 and DBH in the SH-SY5Y (n=5) and $GATA3^{-/-}$ clone xenografts ($GATA3^{-/-}$#1: n=4; $GATA3^{-/-}$

#2: n=3) with Vinculin as a loading control. (**D**) PCA as in **Figure 4A** in which were added the xenografts

of the SH-SY5Y and $GATA3^{-/-}$ clones and the CD44^{pos} or CD44^{neg} sorted cells of the SK-N-SH cell line. (**E**)

Tracks of ChIP-seq profile for H3K27ac at *PHOX2B* and *HAND1* super-enhancers in the xenografts of

the following cells: SH-SY5Y, $GATA3^{-/-}$ clones and SK-N-SH cell populations sorted according to CD44

expression.

Figure 6

Figure 6. Single-cell transcriptomic analyses of 14 PDX models. (**A**) Schematic illustration of the overall

procedure. scRNA-seq was performed for 14 neuroblastoma PDX models obtained either at diagnosis

(light blue) or at relapse (dark blue) after tumor cell dissociation. In parallel, CNV profiles and global

transcriptomes were established from the DNA and RNA of the bulk of the tumor. (**B**) Uniform manifold

approximation and projection (umap) of the 47,219 human cells obtained after the integration by Harmony of the 14 PDX models. (**C**) *PHOX2B* and *HAND2* expression within the noradrenergic tumor cells. (**D**) Plots of published noradrenergic (NOR) and mesenchymal (MES) signatures[12,13]. (**E**) Dot plot graph illustrating cluster-specific gene expression. Four main populations of tumor cells could be defined. (**F**) The GR-NB7 PDX model does not express *GATA3* but exhibits a noradrenergic phenotype as shown by the plot of *PHOX2B*, *CHGA* and *DDC* expression in the scRNAseq data.

MATERIAL AND METHODS

RESOURCE AVAILABILITY

Lead Contact

Further information and request for resources should be directed to and will be fulfilled by the Lead

Contact, Isabelle Janoueix-Lerosey (isabelle.janoueix@curie.fr).

Material availability

Availability of the IC-pPDXC-63 model generated in this study is subjected to a Material Transfer

Agreement.

Data and code availability

ChIP-seq (RRID:SCR_001237) data of cell lines and xenografts (34 samples) are available in Gene

Expression Omnibus (GEO, RRID:SCR_005012) under the accession number GSE154907.

All single-cell RNA-seq from PDXs, all RNA-seq and the ChIP-seq data on the IC-pPDXC-63 cell line are

available in European Genome-Phenome Archive (EGA) under the accession number

EGAS00001004781 (ongoing submission).

EXPERIMENTAL MODELS

Neuroblastoma cell lines

The SK-N-SH (Cat# HTB-11, RRID:CVCL_0531) and SH-SY5Y (Cat# CRL-2266, RRID:CVCL_0019) cell lines

have been obtained from the ATCC. The SH-EP cell line has been kindly provided by M. Schwab. Cell

line authentication was done by STR profiling with PowerPlex® 16 HS System from Promega. The IC-

pPDXC-63 cell line was derived from the IC-pPDX-63 model. $GATA3^{-/-}$ clones were genetically modified

from the SH-SY5Y cell line. Cells were grown at 37°C with 5% CO_2 in a humidified atmosphere in

DMEM/HIGH glucose (Cat# SH30022.01, GE Healthcare) for SK-N-SH, SH-SY5Y and $GATA3^{-/-}$ clones and

in RPMI-1640 (Cat# SH30027.01, GE Healthcare) for IC-pPDXC-63 and SH-EP cell lines, with 10% FBS

(Cat# SV30160.03, GE Healthcare). Cells were monthly checked by qPCR (Venor® GeM qEP 11-9250,

Minerva biolabs®) for the absence of mycoplasma.

Mouse xenograft experiments

For each cell line, 5 million cells were injected subcutaneously in the flanks of 8 week-old Nude or NSG

mice with a ratio of 50/50 standard medium (DMEM: Cat# SH30022.01, GE Healthcare) and BD

Matrigel™ (Cat# 356234, BD Biosciences). Tumor volume was measured every 2 or 3 days with a

caliper. Mice were sacrificed when the tumor reached a volume of 2,000 mm^3 calculated as V = (a/2)

* b * ((a+b)/2), a and b being the largest and smallest diameters, respectively.

In vivo experiments for this study were performed in accordance with the recommendations of the

European Community (2010/63/UE) for the care and use of laboratory animals. Experimental

procedures were specifically approved by the ethics committee of the Institut Curie CEEA-IC #118

(Authorization APAFIS#11206-2017090816044613-v2 given by National Authority) in compliance with

the international guidelines.

Patient-derived Xenografts (PDX models)

Written informed consents for the establishment of PDXs were obtained for all patients from parents

or guardians. GR-NB4 (previously named MAP-GR-A99-NB-1[12]), GR-NB5 (previously named MAP-GR-

B25-NB-1[12]), GR-NB7 and GR-NB10 have been provided by Birgit Geoerger (Gustave Roussy, Villejuif,

France). IC-pPDX-63, IC-pPDX-75, IC-pPDX-109 and IC-pPDX-112 have been developed at Institut Curie.

HSJD-NB-003, HSJD-NB-004, HSJD-NB-005, HSJD-NB-007, HSJD-NB-009 and HSJD-NB-011 PDX models

have been provided by Angel Carcaboso (Institut de Recerca San Joan de Déu, Barcelona, Spain). These

models have been generated from patients under an Institutional Review Board-approved protocol or

within a clinical trial, respectively (**Table S2**).

METHOD DETAILS

CRISPR-Cas9 KO strategy

The guide RNAs used to specifically target the *PHOX2A*, *PHOX2B* and *GATA3* genes were selected from

the CRISPOR website (crispor.tefor.net) for their high predicted efficiency and specificity:

CAATTCGTACGACTCGTGCG   and   CTTGGAATCGTCGTCCTCGG   targeting   *PHOX2A*,

CCCAGCCATACAGGACTCGT   and   AAACTCTTCACGGACCACGG   targeting   *PHOX2B*,

GTACTGCGCCGCGTCCATGT and GAGCTGTACTCGGGCACGTA targeting *GATA3*. *PHOX2A, PHOX2B* and

GATA3 KO were performed by inducing a large deletion between exons 1 and 3, exons 1 and 2 or exons

2 and 3, respectively (**Figure S1A**). To screen the KO clones, 3 pairs of primers were used per gene:

surrounding   the   deleted   region   (*PHOX2A*:   5'-CCGATGGACTACTCCTACCTC-3',   5'-

GCCGGCAGCTAGAAGAGATT-3',   *PHOX2B*:   5'-GTTGGACAGCTCAGTTCCC-3'   ,   5'-

CCCTAGGTCCTTCTCACTCG-3',   *GATA3*:   5'-GCAGAATTGCAGAGTCGTCG-3',   5'-

AAGAGCTGGCTCCTACCTGT-3'), at the 1st cut site (*PHOX2A*: 5'-GGCCGATGGACTACTCCTACCT-3', 5'-

GGGGGACAGTCGCATTCAC-3',   *PHOX2B*:   5'-CAGCAATAAGACCAACCGCT-3',   5'-

GGTTCGGGTGTGACTAGGAT-3',   *GATA3*:   5'-TTGCTAAACGACCCCTCCA-3',   5'-

AAATGAACCAGGAACGGCAG-3') and at the 2nd cut site (*PHOX2A*: 5'-AGCTTTGAAAACCCGGAGCC-3', 5'-

CGGCTGCCAAGCCTTAAGTA-3',   *PHOX2B*:   5'-TCTCAAGTCCGTCACATCGC-3',   5'-

ATTTCTGATCGGCCATGGGG-3', *GATA3*: 5'-TGCGAGGTAGAGATTCCCCA-3', 5'-

GCTAGGATGGGAGGACATGC-3'). The guide RNAs, crRNA and the recombinant Cas9 protein were

purchased from Integrated DNA Technologies. Their reconstitution, assembly and delivery to the cells

were performed following manufacturer's instructions. KO efficiency was determined 2 days after cell

transfection and clones were generated by plating 200 cells in 10 cm culture dishes. KO clones were

verified by Sanger sequencing with the BigDye Terminator V1.1 Cycle Sequencing Kit (130-098-462,

Thermo Fisher).

Phalloidin staining

100,000 cells for SH-SY5Y, *PHOX2B$^{-/-}$* and *PHOX2A$^{-/-}$* clones, 50,000 cells for *GATA3$^{-/-}$* clones were plated

in a 4-well Lab-Tek chamber (Cat# 177399PK, Thermo Fisher) 24 hours before immunostaining. Cells

were fixed with 4% PFA buffer, permeabilized with a 0.2% triton solution and blocked in a 1% BSA 0.1%

triton solution. Phalloidin–Tetramethylrhodamine B isothiocyanate (Sigma Aldrich Cat# P1951,

RRID:AB_2315148) was used at 1:100 to stain actin filaments. DAPI (Cat# 62248, Thermo Fisher) was

diluted at 1: 1,000 in ProLong™ Gold (Cat# P36930, Thermo Fisher).

Immunoblotting

Proteins were extracted using a RIPA buffer (NaCl 150m M, Tris 50 mM pH=7.5, EDTA 1 mM, SDS 0.1%,

deoxycholic acid 0.25%, IGEPAL 1%, PMSF 1 mM) supplemented with protease inhibitor cocktail tablets

(Cat# 11836145001, Roche). 30 to 40 μg of proteins were used for Western blot analysis. The

antibodies targeting the PHOX2B N-terminal part (Santa Cruz Biotechnology Cat# sc-376997,

RRID:AB_2813765), PHOX2A (Santa Cruz Biotechnology Cat# sc-81978, RRID:AB_1127226) and DBH

(Santa Cruz Biotechnology Cat# sc-15318) were used at 1:500. Anti-GATA3 (Cell Signaling Technology

Cat#5852) and HAND2 (Abcam Cat# ab200040) were used at 1: 1,000 and anti-Vinculin (Abcam Cat#

ab129002, RRID:AB_11144129) was used at 1: 10,000.

Proliferation assays

To evaluate the proliferation rate of the *PHOX2B$^{-/-}$* and *GATA3$^{-/-}$* clones compared to the parental SH-

SY5Y cell line, 10,000 cells were plated in a 24-well plate in 6 replicates. Cells were then counted at day

3, 7 and 10, using a Vi-cell XR Viability Analyzer (Beckman Coulter).

Invasion and migration assays

Invasion assays were performed in 96-well low adherent plates (Corning) with 2,000 cells per well for

the SH-EP cell line (n=12), and 10,000 cells per well for the SH-SY5Y cell line (n=12) and *GATA3$^{-/-}$* clones

(n=18). After 4 days of culture, spheroids were embedded in Corning® Collagen I (cat# 354249, Corning)

and cultured in normal conditions. Invasive cells were observed 72 hours after spheroid inclusion and

the area of migration was quantified using the ImageJ software (RRID:SCR_003070).

Transwell experiments were performed using 12-well 8 µm culture inserts (Dutscher). 50,000 cells

were used per condition. After 24 hours of incubation, cells that have migrated through the insert

membrane were fixed and stained with crystal violet (Sigma Aldrich). In each replicate, migrated cells

were quantified using the mean of 6 different images, 6 replicates were performed per cell line.

Immunofluorescence

100,000 cells for SK-N-SH and IC-pPDXC-63 cell lines were plated in a 4-well Lab-Tek chamber (Cat#

177399PK, Thermo Fisher) 48 hours before immunostaining. Cells were fixed with 4% PFA buffer,

permeabilized with a 0.2% triton solution and blocked in a 1% BSA 0.1% triton solution and incubated

with anti-PHOX2B (Santa Cruz Biotechnology Cat# sc-376997, RRID:AB_2813765) and anti-CD44 (Cat#

15675-1-AP, Proteintech) at 1:100. Secondary antibodies were Cy5-Anti-Mouse (Cat# 715-175-151,

Jackson ImmunoResearch Labs / 1:100, RRID:AB_2340820) and Cy3-Anti-Rabbit (Cat# 711-165-152,

Jackson ImmunoResearch Labs / 1:100, RRID:AB_2307443) and DAPI (Cat# 62248, Thermo Fisher) was

diluted at 1: 1,000 in ProLong™ Gold (Cat# P36930, Thermo Fisher).

Chemotherapy treatments

SH-SY5Y, SH-EP or *GATA3*[−/−] clones were plated in a 24-well plate 24 hours before the chemotherapy

treatments. Seeding densities for each cell lines were optimized to reach 80% confluence in the

untreated cells. Cells were treated with 3 different doses of the conventional chemotherapy for 2 days

and cell viability was assessed using a Vi-cell XR Viability Analyzer (Beckman Coulter); 6 replicates were

performed for each cell line and each dose.

IC-pPDXC-63 and SK-N-SH cell lines were plated in 96-well plates 24 hours before the addition of

doxorubicin or etoposide. Seeding densities for each cell line were optimized to reach 80% confluence

in the untreated cells. Cells were treated with chemotherapeutic agents for 72 hours. Cell viability was

then measured using the Resazurin reagent (Sigma-Aldrich).

RNA-sequencing and analyses

RNAs were extracted from frozen tumors by mechanical crushing followed by TRIzol® reagent

(Cat#15596018, Invitrogen) and purified with the NucleoSpin RNA kit (Cat# 740955.50, Macherey-

Nagel). For the bulk scRNA-seq samples and the cell lines, extraction and purification were done

directly using this NucleoSpin RNA kit. RNA quality was assessed with a Bioanalyzer instrument and

RNAs with an RNA Integrity Number above 7 were processed for sequencing as previously described[12].

RNA sequencing libraries were prepared from 500 ng to 1 µg of total RNA using the Illumina TruSeq

Stranded mRNA Library preparation kit according to manufacturer recommendation. For the

xenografts of IC-pPDXC-63 *in vivo* sample, mRNA Library preparation was done with TruSeq RNA Exome

from Illumina. 100 bp paired-end sequencing was performed with the Illumina NovaSeq 6000

instrument (pair-ended, 100 nt).

Reads were aligned to the human reference genome hg38/GRCh38 using STAR 2.6.1a_08-27

(RRID:SCR_015899, https://github.com/alexdobin/STAR) with the following options: -

outFilterMismatchNoverLmax 0.04 --alignIntronMin 20 – lignIntronMax 1000000 –

outFilterMultimapNmax 20. Gene expression values (FPKM=fragments per kilobase per million reads)

were computed by Cufflinks v2.2.146 (RRID:SCR_014597, http://cole-trapnell-lab.github.io/cufflinks/)

and further normalization between samples was done using quantile normalization (R/Bioconductor

package LIMMA (RRID:SCR_010943)).

PHOX2B immunohistochemistry

Tumors were fixed in a 4% formol buffer (VWR) during 24 hours, embedded in paraffin and cut in 4 μm

slices. For PHOX2B immunohistochemistry, the REAL™ EnVision™ Detection System (Cat# K406511-2,

Agilent Technologies) was used and the antibody against PHOX2B (Cat# sc-376997, Santa Cruz) was

diluted at 1:200. Xenograft slices were also colored with a hematoxylin solution.

ChIP-seq and analyses

H3K27ac chromatin immunoprecipitation (ChIP) experiments were performed as previously

described[12] using the iDeal ChIP-seq kit for histones (Cat# C01010171, Diagenode) and the H3K27ac

rabbit polyclonal antibody (Abcam Cat# ab4729, RRID:AB_2118291). Illumina sequencing libraries

were prepared from the ChIP and input DNA and sequenced on the Illumina NovaSeq 6000 instrument

(single reads, 100 nt). ChIP-seq reads were mapped to the human reference genome hg19/GRCh37

using Bowtie2 v2.1.0 (http://bowtie-bio.sourceforge.net/bowtie2/index.shtml) and further analyzed

with HMCan v1.40 (RRID:SCR_010858)[45]. Super-enhancers were called with LILY software as previously

described[12]. LILY was also used to normalize HMCan density profiles between samples. The H3K27ac

signal on super-enhancers shown in the heatmap was computed as the sum of normalized H3K27ac

densities divided by the length of the super-enhancers. For TF genes associated with several super-

enhancers, the signal of the associated super-enhancers is summed and divided by their total length.

PCA was computed as described previously[12].

Flow cytometry and sorting

Flow cytometry analysis was performed with the BD™ LSRII cytometer. Cells were detached with

TrypLE™ Express Enzyme (Cat# 12604013, Gibco), suspended in PBS and permeabilized with the

IntraPrep kit (Cat# A07803, BeckmanCoulter). The cell suspension was incubated with PHOX2B [Clone

B-11] - AlexaFluor® 647 (Cat# SC-376997 AF647, Santa Cruz) and CD44-FITC (Cat# 103005, Biolegend,

RRID:AB_312956) antibodies during 40 min at 4°C.

Flow cytometry sorting was performed with the S3e™ cell sorter (Bio Rad). Cells were detached with

TrypLE™ Express Enzyme (Cat# 12604013, Gibco), suspended in PBS and incubated with CD44-FITC

antibody 30 min at 4°C in dark. Cells positive and negative for CD44 staining were sorted.

The first gating based on FSC/SSC represents 60% for IC-pPDXC-63 and 75% for SK-N-SH. Doublet cells

are eliminated by gating on SSC-W / SSC-H followed by FSC-W / FSC- H. The second gating based on

DAPI negative staining eliminates dead cells. The boundaries between positive staining and negative

staining are always more than 1 Log of fluorescence intensity. A control tube without staining is always

analyzed to determine auto-fluorescence.

Tumor dissociation into single-cell suspension

PDX tumors were cut with scalpels in small fragments. Enzymatic dissociation was realized in CO_2

independent medium (GIBCO) containing 150 µg/mL Liberase™ TL Research Grade (Cat# 5401020001,

Merk) and 150 µg/mL DNase (DN25, Sigma Aldrich), for 30 min at 37°C with 400 rpm agitation. Cell

suspension was then filtered using 70 µm cell strainer (Cat# 130-098-462, Miltenyi Biotec). The cell

suspension was washed twice with PBS. Viability was measured using Vi-cell XR Viability Analyzer

(Beckman Coulter). For some PDXs, the Mouse Cell Depletion KIT was used following the

manufacturer's instructions (Cat# 130-104-694, Miltenyi Biotec).

Single-cell RNA-sequencing experiments and preprocessing of data

Single-cell RNA-seq was performed with the 10x Genomics Chromium Single Cell 3' Kit (v3) according

to the standard protocol. Libraries were sequenced on an Illumina HiSeq2500 or NovaSeq 6000

sequencing platform.

CellRanger version 3.1.0 (10x Genomics, https://support.10xgenomics.com/) was used to demultiplex,

align and generate UMI count tables from sequencing reads. Two reference genomes were used to

align reads:

• The human reference genome (hg38/GRCh38) for *in vitro* cell lines.

• A human-mouse reference genome (GRCh38-mm10) for the 14 neuroblastoma PDX models. In this

scenario, we identified the mouse and human cells after inspection of the percentage of coverage from

GRCh38 genome. We labeled cells as either human (at least 80%), murine (less than 30%) or human-

murine doublets (between 30-80%). Only human and human-murine doublet cells were selected and

coverage plus gene information of only GRCh38 genome were retained for downstream analysis.

Summary of analyses are shown in **Table S3**. Of note, we performed two technical replicates for three

models: HSJD-NB-005, IC-pPDX-63 and IC-pPDX-75 (* in Table S3) to assess reproducibility at distinct

passages in mice (data not shown).

Doublet detection

Scrublet[46] v0.2.1   was   used   to   detect   potential   doublets   using   default   parameters

(expected_doublet_rate=0.06). Cells marked as doublets were removed from subsequent analysis

(results in **Table S3**). Doublet detection was only feasible on samples aligned to GRCh38 genome.

Quality control of single-cell data

First, all ribosomal genes (defined as *RLP/RPS* genes) were removed from the raw expression matrices.

Then, coverage thresholds were set for each sample individually; an upper threshold was set to remove

outlier cells with coverage greater than the 99th percentile, and a lower threshold was set to remove

low quality cells with coverage inferior to the 1st percentile, except for the IC-pPDXC-63 cell line for

which the limits were 1000 UMI and 500 genes detected per cell. To avoid cells with low number of

genes, the same lower threshold was applied on the number of genes thus defining a minimum

number of genes required. Finally, cells with more than 20% of reads mapping mitochondrial genes

were removed.

Normalization of single-cell data

Raw UMI counts were normalized using the "SCTransform" function of Seurat[47,48] v3.1.5

(RRID:SCR_007322). Regressed variables included cell coverage, number of features, and the

percentage of UMI from mitochondrial genes.

Dimensionality reduction and cluster identification

Normalized count data was subjected to dimensionality reduction keeping the first 30 principal

components. Uniform Manifold Approximation and Projections (umap) embeddings were calculated

using these PCs as input and cells were clustered using the "FindClusters" function of Seurat.

Cell type annotation

Marker genes that define cell clusters were identified after differential expression analysis using

Seurat's "FindAllMarkers" function. Clusters were annotated by comparing their top marker genes to

canonical cell type markers from the literature.

Generation of single-cell signature scores

To plot the expression of gene signatures in single cells, we used the "AddModuleScore" function from

Seurat R package with 100 genes in the control gene set. Expression scale was binned into 10 bins

when a gene signature is plotted and into 3 bins (1= low, 2= median, 3= high) when a single gene is

plotted.

scRNA-seq data integrations

Harmony[24] (https://github.com/immunogenomics/harmony) was used to integrated the 14 PDX

models. Downstream analysis was carried out as described in ''Dimensionality reduction and cluster

identification". Of note, when several single-cell transcriptomes are available for the same model, only

one was used for the integration to avoid over-representation.

Cell cycle analysis

We scored single cells based on expression of G2/M and S phase markers using Seurat's

"CellCycleScoring" function.

Copy number analysis in single cells

Copy number variations at the single cell level were called with R package InferCNV v1.2.1 [22]

(https://github.com/broadinstitute/inferCNV) using default parameters. Normal cells from the

microenvironment were used as reference cells. Cells with fewer than 1000 UMI were excluded and

monocytes from publicly available single cell RNA sequencing of healthy human PBMCs[49,50] (GEO:

GSE115189) were used as reference cells (n=376).

QUANTIFICATION AND STATISTICAL ANALYSES

Statistical tests were performed using GraphPad Prism 8 (RRID:SCR_002798). Significance values are

described in the figure legends. P-values were determined via two-tailed unpaired Welch's t-test

(**:p<0.01; ***:p<0.001).

SUPPLEMENTAL INFORMATION

Figure S1: Knock-out of *PHOX2A, PHOX2B* and *GATA3* genes by CRISPR-Cas9.

Figure S2: Sorted mesenchymal cells from the IC-pPDXC-63, SK-N-SH and *GATA3*$^{-/-}$ cell lines, adopt a

noradrenergic identity when engrafted in the mouse.

Figure S3: Detailed analyses of single-cell RNAseq of the 14 PDX models.

Table S1: Expression levels of noradrenergic markers, TFs of the noradrenergic and mesenchymal CRCs

in cell lines and xenografts.

Table S2: Characteristics of the 14 neuroblastoma PDX models studied by scRNA-seq.

Table S3: Filtering of human cells with high quality data for the 14 neuroblastoma PDX models.

Table S4: Lists of genes that are up-regulated in the different clusters of noradrenergic cells in a series

of 14 neuroblastoma PDX models.

Figure S1

Figure S1: Knock-out of *PHOX2A, PHOX2B* and *GATA3* genes by CRISPR-Cas9. (**A**) Two guide RNAs were chosen per gene to induce large deletions (see Methods). The PAM motifs are surrounded in blue when still presents in the modified sequence. Both *PHOX2A^-/-^* clones are homozygous and present with

a large deletion between the 1st and 3rd exons, following Cas9-induced breaks at the expected

positions. This results in a frameshift and subsequently in a truncated protein that has lost all its

functional domains, containing only the 11 first amino acids of the normal PHOX2A protein. The

PHOX2B-/- clone is heterozygous. In the first allele, the Cas9 induced one break in exon 1 in the PAM

motif instead of 3 bp upstream and one break in exon 2, 4 bp upstream of the PAM (instead of 3 bp

upstream). This results in a frameshift and subsequently in a truncated protein, in which only the first

13 amino acids of the normal PHOX2B protein are conserved. In the second allele, the Cas9 induced

only one cleave in the first exon, 3 bp upstream of the PAM, as expected and an "A" insertion was

observed. This results in a frameshift and subsequently in a truncated protein, in which only the first

15 amino acids of the normal PHOX2B protein are conserved. *GATA3*-/- clone #1 is homozygous and

presents with a large deletion between the 2nd and 3rd exons, following Cas9-induced breaks that

occurred at the expected positions. This leads to a truncated protein of 112 amino acids that has lost

all its functional domains and contains only the 35 first amino acids of the normal GATA3 protein.

GATA3-/- clone #2 is heterozygous. One allele is similar to those of *GATA3*-/- clone #1. For the second

allele, the break occurred at the expected position but the *GATA3* sequence between these two

cleavage sites was reintegrated in the reverse direction, with one "C" insertion. This results in the

translation of an aberrant protein of 36 amino acids, which contains only the first 35 amino acids of

the normal GATA3 protein (HD: homeodomain, Ala: poly-alanines track, TA: transactivation domain,

Zn: zinc finger domain) (**B**) Western blot analysis of PHOX2A, PHOX2B and GATA3 TFs in the 2 *PHOX2A*-

/- clones and in the SH-SY5Y cell line, Vinculin was used as a loading control. (**C**) Unsupervised clustering

analysis using the top 10% of genes with the highest IQR (inter-quantile range) shows that *PHOX2A*-/-

and *PHOX2B*-/- clones resemble the parental noradrenergic SH-SY5Y cell line whereas *GATA3*-/- clones

are clustered in the same branch as the SH-EP mesenchymal cell line. (**D**) Volcano plot of a differential

analysis comparing the 4 *GATA3*-/- samples with 5 noradrenergic neuroblastoma cell lines without

MYCN amplification (CLB-GA, NB-EBc1, SH-SY5Y, SJNB-1 and SK-N-FI). We used a raw p-value <0.05 and

a fold-change >2. Gene Ontology analysis using ToppGene

(https://toppgene.cchmc.org/enrichment.jsp) performed on the lists of differentially expressed genes

in the *GATA3*-/- samples compared to the other neuroblastoma cell lines.

Figure S2

Figure S2: Sorted mesenchymal cells from the IC-pPDXC-63, SK-N-SH and *GATA3*[-/-] cell lines, adopt a noradrenergic identity when engrafted in the mouse. Unsupervised hierarchical clustering based on a transcriptomic signature[13] discriminating noradrenergic and mesenchymal cells shows that all the *in vitro* mesenchymal cell populations engrafted in mice give tumors with a noradrenergic transcriptomic profile.

Figure S3

Figure S3

G

Figure S3: Detailed analyses of single-cell RNAseq of the 14 PDX models.

(**A**) The histogram highlights the contribution of the different samples to each cluster in percentage. The pie chart depicts the contribution of each sample to the total number of cells (n= 47,219). (**B**) (Top) Copy number profile inferred from WES from HSJD-NB-003 PDX model. (Bottom) InferCNV profile obtained from scRNAseq data of HSJD-NB-003. (**C**) Overview of the genomic alterations present in each

PDX model with the InferCNV profiles on the 14 PDX models, using 500 randomly-selected cells from each sample. (**D**) Individual expression of some genes representing the mains categories identified in the noradrenergic tumor cells: cell cycle, MYCN/2p-amp, sympathoblast-like and chromaffin-like. (**E**) The tumor cells are colored according to their corresponding cell cycle phase (red: G1 phase; green: G2/M phase; blue: S phase). (**F**) Plot of the signature of MYCN target genes[51] in the integration of all PDX samples. (**G**) Unsupervised hierarchical clustering using the top 10% of genes with the highest IQR (inter-quantile range) for all single cell RNAseq from PDX samples, the mesenchymal (SH-EP and hNCCs) and noradrenergic (SH-SY5Y) cell lines. This shows that all PDX models clustered in the noradrenergic (NOR) branch.

REFERENCES

1. Matthay, K. K. *et al.* Neuroblastoma. *Nat Rev Dis Primers* **2**, 16078 (2016).

2. Huber, K. The sympathoadrenal cell lineage: specification, diversification, and new

perspectives. *Dev. Biol.* **298**, 335–343 (2006).

3. Marshall, G. M. *et al.* The prenatal origins of cancer. *Nat. Rev. Cancer* **14**, 277–289

(2014).

4. Brodeur, G. M., Seeger, R. C., Schwab, M., Varmus, H. E. & Bishop, J. M. Amplification

of N-myc in untreated human neuroblastomas correlates with advanced disease stage.

Science **224**, 1121–4 (1984).

5. Chen, Y. *et al.* Oncogenic mutations of ALK kinase in neuroblastoma. *Nature* **455**, 971–4

(2008).

6. George, R. E. *et al.* Activating mutations in ALK provide a therapeutic target in

neuroblastoma. *Nature* **455**, 975–8 (2008).

7. Janoueix-Lerosey, I. *et al.* Somatic and germline activating mutations of the ALK kinase

receptor in neuroblastoma. *Nature* **455**, 967–70 (2008).

8. Mosse, Y. P. *et al.* Identification of ALK as a major familial neuroblastoma

predisposition gene. *Nature* **455**, 930–5 (2008).

9. Ackermann, S. *et al.* A mechanistic classification of clinical phenotypes in

neuroblastoma. *Science* **362**, 1165–1170 (2018).

10. Peifer, M. *et al.* Telomerase activation by genomic rearrangements in high-risk

neuroblastoma. *Nature* **526**, 700–704 (2015).

11. Valentijn, L. J. *et al.* TERT rearrangements are frequent in neuroblastoma and identify

aggressive tumors. *Nat. Genet.* **47**, 1411–1414 (2015).

12. Boeva, V. *et al.* Heterogeneity of neuroblastoma cell identity defined by transcriptional

circuitries. *Nat. Genet.* **49**, 1408–1413 (2017).

13. van Groningen, T. *et al.* Neuroblastoma is composed of two super-enhancer-associated

differentiation states. *Nat. Genet.* **49**, 1261–1266 (2017).

14. Decaesteker, B. *et al.* TBX2 is a neuroblastoma core regulatory circuitry component

enhancing MYCN/FOXM1 reactivation of DREAM targets. *Nat Commun* **9**, 4866 (2018).

15. Durbin, A. D. *et al.* Selective gene dependencies in MYCN-amplified neuroblastoma

include the core transcriptional regulatory circuitry. *Nat. Genet.* **50**, 1240–1246 (2018).

16. Wang, L. *et al.* ASCL1 is a MYCN- and LMO1-dependent member of the adrenergic

neuroblastoma core regulatory circuitry. *Nat Commun* **10**, 5622 (2019).

17. Rajbhandari, P. *et al.* Cross-Cohort Analysis Identifies a TEAD4-MYCN Positive

Feedback Loop as the Core Regulatory Element of High-Risk Neuroblastoma. *Cancer*

Discov **8**, 582–599 (2018).

18. Ross, R. A., Spengler, B. A. & Biedler, J. L. Coordinate morphological and biochemical

interconversion of human neuroblastoma cells. *J. Natl. Cancer Inst.* **71**, 741–747 (1983).

19. Fernando, J. *et al.* A mesenchymal-like phenotype and expression of CD44 predict lack of

apoptotic response to sorafenib in liver tumor cells. *International Journal of Cancer* **136**,

E161–E172 (2015).

20. Ramos, T. L. *et al.* MSC surface markers (CD44, CD73, and CD90) can identify human

MSC-derived extracellular vesicles by conventional flow cytometry. *Cell Commun.*

Signal **14**, 2 (2016).

21. Saito, S. *et al.* CD44v6 expression is related to mesenchymal phenotype and poor

prognosis in patients with colorectal cancer. *Oncology Reports* **29**, 1570–1578 (2013).

22. Tickle T, Tirosh I, Georgescu C, Brown M, & Haas B. inferCNV of the Trinity CTAT

Project. (2019).

23. Cohen, N., Betts, D. R., Rechavi, G., Amariglio, N. & Trakhtenbrot, L. Clonal expansion

and not cell interconversion is the basis for the neuroblast and nonneuronal types of the

SK-N-SH neuroblastoma cell line. *Cancer Genet. Cytogenet.* **143**, 80–84 (2003).

24. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

25. Chan, W. H. *et al.* RNA-seq of Isolated Chromaffin Cells Highlights the Role of Sex-

Linked and Imprinted Genes in Adrenal Medulla Development. *Sci Rep* **9**, 3929 (2019).

26. Furlan, A. *et al.* Multipotent peripheral glial cells generate neuroendocrine cells of the

adrenal medulla. *Science* **357**, (2017).

27. Van Limpt, V. A. E. *et al.* High delta-like 1 expression in a subset of neuroblastoma cell

lines corresponds to a differentiated chromaffin cell type. *Int J Cancer* **105**, 61–69 (2003).

28. De Preter, K. *et al.* Human fetal neuroblast and neuroblastoma transcriptome analysis

confirms neuroblast origin and highlights neuroblastoma candidate genes. *Genome Biol.*

7, R84 (2006).

29. Goridis, C. & Rohrer, H. Specification of catecholaminergic and serotonergic neurons.

Nature Reviews Neuroscience **3**, 531–541 (2002).

30. Pattyn, A., Morin, X., Cremer, H., Goridis, C. & Brunet, J. F. The homeobox gene

Phox2b is essential for the development of autonomic neural crest derivatives. *Nature*

399, 366–70 (1999).

31. Rohrer, H. Transcriptional control of differentiation and neurogenesis in autonomic

ganglia. *Eur J Neurosci* **34**, 1563–73 (2011).

32. van Groningen, T. *et al.* A NOTCH feed-forward loop drives reprogramming from

adrenergic to mesenchymal state in neuroblastoma. *Nat Commun* **10**, 1530 (2019).

33. Dong, R. *et al.* Single-Cell Characterization of Malignant Phenotypes and Developmental

Trajectories of Adrenal Neuroblastoma. *Cancer Cell* (2020)

doi:10.1016/j.ccell.2020.08.014.

34. Asgharzadeh, S. *et al.* Clinical significance of tumor-associated inflammatory cells in

metastatic neuroblastoma. *J. Clin. Oncol.* **30**, 3525–3532 (2012).

35. Piskareva, O. *et al.* The development of cisplatin resistance in neuroblastoma is

accompanied by epithelial to mesenchymal transition in vitro. *Cancer Lett.* **364**, 142–155

(2015).

36. Debruyne, D. N. *et al.* ALK inhibitor resistance in ALK(F1174L)-driven neuroblastoma

is associated with AXL activation and induction of EMT. *Oncogene* **35**, 3681–3691

(2016).

37. Lumb, R. & Schwarz, Q. Sympathoadrenal neural crest cells: the known, unknown and

forgotten? *Dev. Growth Differ.* **57**, 146–157 (2015).

38. Olsen, R. R. *et al.* MYCN induces neuroblastoma in primary neural crest cells. *Oncogene*

36, 5075–5082 (2017).

39. Tsubota, S. & Kadomatsu, K. Origin and initiation mechanisms of neuroblastoma. *Cell*

Tissue Res. **372**, 211–221 (2018).

40. Dun, S. L., Brailoiu, G. C., Yang, J., Chang, J. K. & Dun, N. J. Cocaine- and

amphetamine-regulated transcript peptide and sympatho-adrenal axis. *Peptides* **27**, 1949–

1955 (2006).

41. El Faitwri, T. & Huber, K. Expression pattern of delta-like 1 homolog in developing

sympathetic neurons and chromaffin cells. *Gene Expr Patterns* **30**, 49–54 (2018).

42. Schütz, B., Schäfer, M. K., Eiden, L. E. & Weihe, E. Vesicular amine transporter

expression and isoform selection in developing brain, peripheral nervous system and gut.

Brain Res Dev Brain Res **106**, 181–204 (1998).

43. Weihe, E., Schäfer, M. K., Erickson, J. D. & Eiden, L. E. Localization of vesicular

monoamine transporter isoforms (VMAT1 and VMAT2) to endocrine cells and neurons

in rat. *J Mol Neurosci* **5**, 149–164 (1994).

44. Wildner, H., Gierl, M. S., Strehle, M., Pla, P. & Birchmeier, C. Insm1 (IA-1) is a crucial

component of the transcriptional network that controls differentiation of the sympatho-

adrenal lineage. *Development* **135**, 473–481 (2008).

45. Ashoor, H. *et al.* HMCan: a method for detecting chromatin modifications in cancer

samples using ChIP-seq data. *Bioinformatics* **29**, 2979–2986 (2013).

46. Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of Cell

Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291.e9 (2019).

47. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nature*

Biotechnology **36**, 411–420 (2018).

48. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21

(2019).

49. Franzén, O., Gan, L.-M. & Björkegren, J. L. M. PanglaoDB: a web server for exploration

of mouse and human single-cell RNA sequencing data. *Database (Oxford)* **2019**, (2019).

50. Freytag, S., Tian, L., Lönnstedt, I., Ng, M. & Bahlo, M. Comparison of clustering tools in

R for medium-sized 10x Genomics single-cell RNA-sequencing data. *F1000Res* **7**, 1297

(2018).

51. Valentijn, L. J. *et al.* Functional MYCN signature predicts outcome of neuroblastoma

irrespective of MYCN amplification. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 19190–19195

(2012).