**Ataxin-2 is essential for neurodevelopment in *Drosophila* and is a major regulator of the cytoskeleton.**

Urko del Castillo*[1,a,b], Rosalind Norkett[1,a], Anna Serpinskaya[a], Vladimir I. Gelfand*[a]

a. Department of Cell and Developmental Biology, Feinberg School of Medicine, Northwestern University, Chicago, IL 60611.

b. Current address: Division of Hematology and Medical Oncology, Department of Medicine, Weill Cornell Medical Center, New York, NY, 10065.

*Urko del Castillo

*Vladimir I. Gelfand

1. These authors contributed equally to this work

**Email:** urd4002@med.cornell.edu, vgelfand@northwestern.edu

**Author Contributions:** UdC, RN and VG designed research, UdC, RN and AS performed research, UdC and RN analyzed data, RN, UdC and VG wrote the manuscript.

**Competing Interest Statement:** No competing interests.

**Classification:** Biological Sciences, Cell Biology

**Keywords:** Neuron, Microtubules, Actin, organelle trafficking.

**Abstract**

Ataxin-2 (Atx2) is a highly conserved RNA binding protein. Atx2 undergoes polyglutamine expansion leading to Amyotrophic Lateral Sclerosis (ALS) or Spinocerebellar Ataxia (SCA). However, the normal physiological functions of Atx2 remain unknown, likely because of functional redundancy between Atx2 and the related Atx2-Like gene in mammals. Here we use the powerful genetics of *Drosophila* to show that Atx2 is essential for normal cytoskeletal dynamics in neurons. Neuron-specific depletion of Atx2 caused multiple morphological defects in the nervous system of 3[rd] instar larvae. These include reduced brain size, impairments in optic lobe innervation and decreased dendrite outgrowth in sensory neurons. Defects in the nervous system of these larvae caused loss of the ability to crawl and were lethal at the pupal stage. Interestingly, we found severe impairments in cytoskeletal dynamics both in microtubule and actin networks. Microtubules became hyperstabilized as demonstrated by increased tubulin acetylation and resistance to microtubule depolymerizing drugs. Similarly, we found F-actin was hyperstabilized as shown by resistance to depolymerizing agents. Further, we show inhibition of microtubule-microtubule sliding, a crucial process for initial neurite outgrowth. We also demonstrated that microtubule-dependent transport of multiple cargoes in neurons, both *in vitro* and *in vivo*, was dramatically inhibited. Taken together, these data mark Atx2 as a master regulator of cytoskeletal dynamics and denote Atx2 as an essential gene in neurodevelopment, as well as a neurodegenerative factor. These data could provide insight into potential therapeutic interventions for Atx2 polyglutamine disorders.

**Significance Statement**

Neurons extend a complex array of axons and dendrites for network formation and communication. Therefore, precise neuronal development is essential for normal nervous system function and viability. Neuronal development is highly dependent upon dynamics of the actin and microtubule cytoskeleton for extension and maintenance of neurites. There are many descriptions of cytoskeleton impairment in neurodegenerative disease, however, the physiological functions of many genes mutated in neurodegenerative remain undescribed. The major finding of this work is

that the loss of Ataxin-2 at early developmental stages in *Drosophila* is lethal and causes multiple
defects in neurodevelopment via dysregulation of the cytoskeleton.

**Main Text**

**Introduction**

Ataxin-2 (Atx2) is a highly conserved RNA binding protein which is known to associate with poly
ribosomes, coordinate RNA granule assembly and regulate translation for synaptic plasticity (1–4).
Atx2 contains mRNA binding domains, intrinsically disordered domains and a polyQ domain. This
domain has been shown to undergo polyQ expansion leading to neurodegenerative diseases
Spinocerebellar Ataxia (SCA) and Amyotrophic Lateral Sclerosis (ALS) (5, 6). SCA is characterized
by decreased dendritic arbors and abnormal axons in the cerebellum (7), while ALS is
characterized by motor neuron degeneration (8). Both conditions lead to movement disorders. Due
to its causative role in neurodegenerative diseases, there has been much interest in discerning the
function of Atx2, with research largely focused on the Atx2 polyQ expansion. However, Atx2 is
expressed throughout development, indeed it is highly expressed in the developing central nervous
system, and there have also been investigations focusing on the physiological roles of wild type
Atx2. In one study, Atx2 knockout mice were generated (9). These mice displayed no obvious
abnormalities, likely due to compensation by a related protein called Atx2-like. The gene encoding
Atx2-like is absent in *Drosophila*, marking them as an ideal model organism to probe physiological
Atx2 function.

Hints as to the physiological roles of Atx2 have come from studies of its domain structure (1, 10).
Atx2 contains RNA binding domains and intrinsically disordered regions. Atx2 is known to associate
with polyribosomes (11) and can assemble into phase-separated, membraneless stress granules,
along with cognate RNAs. This action serves to concentrate or organize RNAs within the cell (12).
In the case of Atx2, incorporation into phase separated stress granules is dependent upon the
intrinsically disordered regions. Once assembled into granules, Atx2 can associate with its target
RNAs, and stabilizes these transcripts (10). In turn, this action contributes to long term habituation
or memory formation (1, 3). Via RNA organization and control of translation, Atx2 may be a critical
regulator in many cellular pathways. For example, Atx2 has been previously indicated in actin
regulation in the germline (4), microtubule organization in mitosis (13, 14), and in neurite outgrowth
and ER distribution in neurons (15).

Here we use *Drosophila* to characterize the effects of neuron specific loss of wild type Atx2 on
nervous system development. We show that depletion of Atx2 by RNAi at early developmental
stages causes severe impairments in locomotion at the larval stage and is 100% lethal at the pupae
stage of development. In affected larvae, we find impaired brain development and, at the cellular
level, decreased neurite outgrowth of axons and dendrites. We probe the reasons behind these
observations, both in *Drosophila* S2 cells as a versatile tool for protein knock down, and in
*Drosophila* neurons. We show the microtubules are hyperacetylated and much less dynamic upon
Atx2 knockdown. The actin cytoskeleton is similarly hyperstabilized. We further demonstrated that
Atx2 loss causes a dramatic reduction in organelle transport in neurons in culture and *in vivo*. This
leads to disturbed cargo distribution, with organelles becoming trapped in neurite varicosities.
Within these varicosities, the microtubule cytoskeleton is highly disorganized, showing looping
rather than linear microtubules. Together, our work describes that Atx2 is an essential gene in

neurodevelopment, required for proper cytoskeleton organization, dynamics and organelle
transport.
93
94  **Results:**
95
96  **Ataxin-2 is essential for neurodevelopment, locomotion and viability in *Drosophila*.**
97
We used the powerful genetics of *Drosophila* to probe the physiological roles of Atx2. In this way,
we were able to take advantage of the presence of a single Atx2 gene, and avoid potential
redundancy between mammalian Atx2 and Atx2-like (9). In order to study the physiological role of
Atx2 in neurons, we depleted the protein by neuron specific expression of shRNA against Atx2
using the pan-neuronal driver elav-Gal4. Crucially, this is a post-mitotic driver, so any role for Atx2
in cell division is not affected.
104
We started by investigating the effect of neuron specific Atx2 depletion on the whole animal by
looking at viability. We found that depleting Atx2 in neurons was lethal at the pupae stage and
animals expressing the Atx2 RNAi were not viable to adult stages (Fig. 1A, control; 523/528, RNAi
line 1; 0/458, RNAi line 2; 0/188 flies eclosed). We found consistent results with two different Atx2
RNAi lines; therefore, this effect is specific to Atx2 knockdown rather than an off-target effect. Atx2
knockdown was confirmed by western blot of brain lysate (Fig. S1). We also carried out locomotion
studies. We imaged 3$^{rd}$ instar larvae crawling on agar plates and calculated the velocity. Control
larvae crawled over the surface of the plate at about 0.7mm/s. However, Atx2 knockdown larvae
were largely stationary, crawling with a velocity of less than 0.1mm/s (Movie 1). This is further
shown by the color-coded tracks of each larva over time (Fig. 1B), and quantified in Fig. 1C.
115
We wanted to investigate the reasons behind these drastic effects of Atx2 depletion. We carried
out experiments to study development of the nervous system. Firstly, we dissected 3$^{rd}$ instar larvae
brains and immunostained for Fasciclin II (FasII) – an axon specific marker. We found that Atx2
RNAi brains were drastically smaller than controls. Quantification of the optic lobes (Fig. 1D, cyan)
showed a roughly 2-fold decrease in area (Fig. 1E). Further we quantified area of the retinotopic
pattern – the structure representing the photoreceptor axons (Fig. 1D, magenta). When normalized
to the area of the optic lobe, we found this structure was decreased to around 10% of control area
upon Atx2 depletion (Fig. 1F). These phenotypes demonstrate that Atx2 depletion induces
impairment of axon outgrowth and targeting.
125
We investigated the effect of the loss of Atx2 on dendritic development. To do this, we labelled
Class IV da (Dendritic arborization) neurons (sensory neurons that line the *Drosophila* body wall)
with a fluorescent reporter expressed specifically in these cells (ppk::tdTomato). Under control
conditions, these neurons extended long, branched dendritic arbors which covered the whole body
segment. However, upon loss of Atx2, we found that these dendritic arbors were drastically
shrunken (Fig. 1G). This effect was quantified in two ways, firstly the total dendrite length per cell
was decreased around 65% (Fig. 1H). These observations are consistent with our previously
published *in vitro* data where we showed depleting Atx2 decreased neurite length in cultured
neurons (15). Secondly, to further describe the effect of Atx2 depletion on the shape of the dendritic
arbor, we carried out Sholl analysis (16). This analysis describes dendritic complexity as a function
of distance from the soma. Concentric circles are drawn outwards from the soma, and number of
intersections that the dendritic arbor makes with each circle is counted. From 20μm from the soma,
the Atx2 RNAi neurons displayed many fewer intersections than control neurons and the dendritic

arbors only extended to around 100μm from the soma, whereas the dendrites of the control neurons
extended around 30% more than this (Fig. 1I). Therefore, the Atx2 knock down severely impairs
dendrite development.
142
143    Taken together, these data show that Atx2 is necessary for normal development of both axons and
144    dendrites. As a result, the loss of Atx2 severely impairs nervous system development and this
145    causes severe locomotion defects and lethality.
146
147    **Ataxin-2 depletion decreases microtubule dynamics.**
148
We and others have demonstrated that one of the major drivers of neuronal development and
neurite outgrowth is the microtubule cytoskeleton (17–21). An inability to properly regulate
microtubule dynamics leads to phenotypes similar to those we reported upon Atx2 depletion (19).
Therefore, we investigated the possibility that microtubule dynamics may be altered upon Atx2
knockdown. Initially, we chose to use *Drosophila* S2 cells to test this hypothesis as they are an
efficient system to study the cytoskeleton using dsRNA strategies (22, 23). We depleted Atx2 in
*Drosophila* S2 cells and carried out immunostaining for acetylated tubulin. Tubulin acetylation
increases with microtubule age, so is a read out for microtubule lifetime (24–26). In S2 cells, we
found a roughly 3-fold increase in acetylated tubulin signal (Fig. 2A, B). Importantly, this finding
was also apparent in *Drosophila* 3[rd] instar larvae neuronal culture *in vitro*. Upon Atx2 knockdown
with elav-Gal4, we found acetylated tubulin signal increased around 30% compared to control (Fig.
2C, D). Further, we investigated acetylated tubulin levels in brain lysate from elav>Atx2 RNAi larvae
brains. Consistent with our immunostaining data, western blotting demonstrated an increase in
acetylated tubulin in Atx2 RNAi brain lysate compared to control (Fig. 2E). Taken together, these
results suggest an increase in microtubule stability upon loss of Atx2.
164
To investigate whether the increase in acetylated tubulin was related to an alteration in microtubule
dynamics, we directly observed exchange of tubulin subunits – the building blocks of polymerized
microtubules. To do this, we depleted Atx2 by RNAi in *Drosophila* S2 cells expressing
photoconvertible EOS-tubulin. UV-light was applied to a specific region of interest and the
photoconverted signal outside the original ROI was quantified 30 minutes after photoconversion.
Incorporation of photoconverted tubulin subunits to microtubules located outside the original
photoconverted area indicates tubulin subunit exchange. Under control conditions, around 50% of
the original photoconverted signal was detected outside the ROI, whereas upon Atx2 depletion,
this value was around 25% (Fig. 2F, G, movie 2). This indicates that Atx2 is necessary for normal
subunit exchange. Additionally, we investigated the stability of microtubules after treatment with a
microtubule depolymerizing agent vinblastine by immunostaining for tubulin. Consistent with our
previous data, we found that Atx2 depletion confers resistance of microtubules to vinblastine –
intact microtubules could be detected in the Atx2 RNAi cells, whereas microtubules in the control
cells were largely depolymerized (Fig. 2H and I).
179
In order to directly assess the effect of loss of Atx2 on microtubule dynamics, we carried out live
imaging of EB1 (end binding protein 1) – a marker for polymerizing microtubule plus-ends (27).
Initially, we observed EB1-GFP comets in S2 cells (Fig. 2J and K, movie 3). We found around a
65% decrease in the number of comets per cell, as indicated by the loss in comets shown in the
temporal color code – a color coded projection of each comet over time. Furthermore, we carried
out the same assay in cultured larvae neurons expressing EB1-GFP (Fig. 2L, movie 4). Comets in
neuronal processes are demonstrated in kymographs in Fig. 2M where time is projected on the Y
axis, showing motile objects as diagonal lines. Upon Atx2 knockdown, we found a roughly 30%

decrease in the number of comets per 10μm (Fig. 2N). Notably, the total microtubule content in
these cells is unchanged, as shown by tubulin staining of extracted control and Atx2 knockdown
neurons (Fig. S2A, B). Therefore, Atx2 knockdown decreases only the dynamic microtubule
population, not the total number of microtubules. Taken together, these data show that loss of Atx2
confers increased microtubule stability and that Atx2 is essential for normal microtubule dynamics,
both in cultured S2 cells, and crucially, in neurons.
194
**Ataxin-2 depletion stabilizes the actin cytoskeleton**
In addition to the microtubule cytoskeleton, neuronal morphology is also dependent upon the actin
cytoskeleton. Therefore, we decided to study the possibility that Atx2 depletion may impact actin
dynamics. Initially, we knocked down Atx2 by RNAi in *Drosophila* S2 cells as a tractable system to
study actin dynamics. We stained for actin filaments using Rhodamine-phalloidin. We also treated
these cells with Latrunculin B (LatB), an actin depolymerizing agent, a comparable experiment to
Fig. 2H. We found that in control cells, the F-actin signal, detected by phalloidin staining, was
decreased by around 35%, indicative of actin depolymerization. However, upon Atx2 knock down,
this effect was occluded. There was no significant alteration in F-actin signal (Fig. 3A, B). To make
this observation relevant to neurodevelopment, we carried out similar experiments in cultured
neurons from *Drosophila* 3$^{rd}$ instar larvae, either wild type, or after Atx2 knock-down. We
demonstrated that after Atx2 knockdown actin filaments in neurons are more stable. As
demonstrated by Rho-phalloidin staining about 40% more F-actin survived the treatment with LatB
(Fig. 3C, D). LatB promotes depolymerization of F-actin by sequestering actin monomers. This
shows that actin polymers are less sensitive to depolymerizing agents upon loss of Atx2, and so,
that the actin network is less dynamic.
211
In order to study the potential effects of increased actin stability on neuronal morphology, we chose
to observe filopodia, thin rod like protrusions rich in F-actin, essential for generation of neurites and
the growth cone (28). We prepared primary neuronal cultures from 3$^{rd}$ instar larvae and carried out
phase contrast imaging to examine filopodia along neuronal processes. Upon loss of Atx2, the
number of filopodia projecting from neurites decreased over 50% (Fig. 3E, F). This is consistent
with a less dynamic actin network upon loss of Atx2. Without correct actin dynamics, formation of
filopodia is impaired and neurons are unable to extend normal arbors, consistent with our data
showing that dendrite and axon development is affected by Atx2 loss (Fig. 1).
220
221  Taken together, these data show that Atx2 is essential for normal actin dynamics in neurons. These
222  effects are comparable to our data demonstrating the microtubule network is stabilized upon Atx2
223  knockdown, and highlights the role of Atx2 in regulating normal cytoskeletal dynamics in neurons.
224
**Ataxin-2 is essential for organelle transport and distribution in *Drosophila* neurons**
Beyond neuronal morphology, both actin and microtubule networks are crucial for normal organelle
transport. Microtubules act as tracks for longer range transport of cargo by kinesin and dynein
motors, whereas actin is responsible for shorter range transport and cargo anchoring by myosin
motors (29–33). Therefore, we hypothesized that Atx2 depletion may alter organelle transport via
its effects on the cytoskeleton. To investigate the effect of loss of Atx2 on organelle trafficking, we
started by carrying out live imaging of mitochondria and lysosomes in *Drosophila* 3$^{rd}$ instar larvae
neurons in culture (Fig. 4A). Using the neuron specific driver elav-Gal4, we labelled mitochondria
using a genetically encoded mCherry fluorescent protein targeted to the outer mitochondrial
membrane. Time lapse microscopy showed around 45% of these organelles were motile in control
conditions. However, depletion of Atx2 caused a large decrease in the fraction of motile
mitochondria – only around 15% of mitochondria moved in the Atx2 RNAi condition, roughly 65%

less than in control neurons. This is exemplified in the kymographs in Fig. 4B and in movie 5 and
quantified in Fig. 4D. In addition to mitochondria, we studied the motility of lysosomes labeled with
lysotracker. Consistent with our data for mitochondrial transport, we found a vast decrease in
lysosome motility upon loss of Atx2 (Fig. 4C, movie 6) – in control conditions around 30% of the
organelles were moving, this dropped about two-fold in Atx2 RNAi neurons (Fig. 4E).
242
Next, we extended these findings to investigate organelle transport *in vivo* in the segmental nerves
of *Drosophila* 3[rd] instar larvae. These nerves contain the axons of motor and sensory neurons which
extend from the CNS into segments of the larvae body wall (Fig. 4F). The great advantage of this
system is that the nerves are visible in the intact *Drosophila* larva without the need for dissection
(34). We studied motility of two cargoes; mitochondria and dense core vesicles. We found that *in*
*vivo* mitochondrial transport was drastically decreased in Atx2 RNAi animals compared to control,
from around 50% of mitochondria moving to around 2% (Fig. 4 G, I, movie 7). Trafficking of dense
core vesicles was similarly impaired in Atx2 RNAi larvae – motile organelles decreased from around
70% to around 20% (Fig. 4H, J, movie 8). Taken together these data show Atx2 has an essential
role in normal organelle transport in neurons *in vitro* and *in vivo*, and that Atx2 is necessary for
normal transport of multiple cargoes.
254
To study the effect of Atx2 dependent loss of cargo transport, we investigated the distribution of
organelles throughout neurons in culture. We have previously shown that depleting Atx2 leads to
varicosities in neuronal processes and that endoplasmic reticulum accumulates in these structures
(15). Phase contrast imaging coupled with labelling of mitochondria and lysosomes demonstrated
that these organelles could be found colocalized in the axonal varicosities (Fig. 4K). Indeed, the
majority of the varicosities held both of these organelles (84%) while only around 5% each of
varicosities contained only one or neither of these organelles. This is consistent with a 'traffic jam'
leading to accumulation of organelles at these sites due to severely impaired transport.
263
Finally, we were interested in the microtubule structure at these 'traffic jam' sites. In order to
visualize microtubules at the sites, we carried out super resolution imaging on extracted Atx2 RNAi
neurons with stained for tubulin. We found that microtubules in these regions were less bundled,
and instead were disordered and looped (Fig. 4L). This disorganization of the microtubule network
is likely responsible for preventing normal organelle transport through these sites, causing their
accumulation.
**Discussion**
271
Multiple works have demonstrated the involvement of mutant Atx2 in neurodegenerative conditions
ALS and SCA (5, 6). However, an understanding of the physiological roles of the RNA-binding
protein in neurons is lacking. Here we use *Drosophila* as a model system to effectively demonstrate
the necessity of wild type Atx2 for normal nervous system development. Underlying the gross
morphological defects seen in the nervous system, we show that loss of Atx2 has large scale effects
on the microtubule and actin cytoskeleton, causing their abnormal stabilization. We also
demonstrate that Atx2 is essential for normal organelle transport and distribution. These
neurodevelopmental defects severely impair neurite outgrowth, locomotion of *Drosophila* larvae
and are lethal at later stages of development.
281
Here we show that Atx2 is essential for normal cytoskeletal dynamics, in both microtubule and actin
networks. Firstly, our data show that specific depletion of Atx2 in neurons drives an increase in
microtubule acetylation, a decrease in tubulin subunit exchange and fewer EB1 comets. Thus, we
conclude microtubules shift to a more stable, less dynamic network. A role for Atx2 in regulating

the microtubule network is consistent with previous reports of Atx2 having roles in centrosome
organization and mitotic spindle regulation in *C. elegans* (13, 14). Secondly, we also found an
increase in the stability of the actin network upon Atx2 knockdown. These data are in good
agreement with early descriptions of Atx2 function, which showed evidence of disorganized actin
bundles in retinal precursor cells, the female germline and bristles in Atx2 null *Drosophila* (4)*.*
291
Microtubule and actin dynamics are intricately linked with neurite outgrowth. Microtubules not only
serve as tracks for directed transport but they are crucial to transduce forces that are generated by
molecular motors, which control neuronal polarization and morphology (18, 35–40). Therefore, it is
highly likely that the developmental defects reported herein are dependent upon loss of microtubule
dynamics. In neurons in particular, impaired actin dynamics have important consequences on
development (41). Interrupting the actin network prevents normal growth cone formation, and so,
normal axon outgrowth (28). Incorrectly regulated actin dynamics may also impair the organization
of the microtubule cytoskeleton. Actin is necessary to guide microtubules in the axon, at filipodia
and at the growth cone (42–45). Fewer filopodia being able to form may contribute to the decrease
in dendritic branching described in DA neurons, potentially underlying the gross neurite extension
defects we describe.
303
Finally, we show that decreasing levels of Atx2 decreases organelle transport in culture and *in vivo*.
Impaired organelle transport is likely a direct consequence of the cytoskeletal defects caused by
Atx2. This is supported by our observation of microtubule looping and disorganization in the axonal
varicosities. The increase in microtubule stability and longevity would be thought to facilitate
microtubule motor- based organelle transport (46, 47). However, we see a decrease in motility of
multiple cargoes. Instead, an increase in the stability of the actin network may lead to organelles
being anchored or trapped by myosin motors (30, 31). This loss of organelle transport leads to
'traffic jams' – an accumulation of organelles - in axonal varicosities. These axonal varicosities are
also known to contain ER aggregates (15). In turn, this impairment in organelle distribution could
lead to the neuronal defects we observed with loss of Atx2 at early developmental stages (39).Other
examples of impaired organelle transport in neurodegenerative disease include mitochondria
trapped in dystrophic neurites in an Alzheimer's disease model (48, 49), and impairments in
endosome retrograde transport in ALS (50). These transport defects can be caused by mutations
in kinesins, dynein and tubulin themselves (51–53), as well as adaptors and other RNA binding
proteins such as TDP43 and Fus (46).
It will be crucial to determine the mechanisms by which Atx2 can act as a master regulator of the
neuronal cytoskeleton. It is known that Atx2 does not associate directly with actin (4), perhaps Atx2
acts indirectly via interaction with actin/microtubule binding proteins, such as profilin, another ALS
risk factor which locates to stress granules (54, 55). Another likely option is via regulation of
expression of its cognate RNAs – perhaps among these transcripts are known regulators of the
actin and microtubule cytoskeleton. Atx2 tightly regulates RNP granules and is necessary for axon
path finding (1, 10). It is essential for local translation for synaptic plasticity (2). Ongoing
experiments serving to identify Atx2 bound RNAs (TRIBE), along with RNA seq data, will prove
indispensable for identifying mechanisms of Atx2 in both neuronal development and
neurodegeneration (10).
329
Currently it is unclear whether the neurodegenerative-related polyQ expansion of Atx2 corresponds
to a gain of toxic function or simply to a loss of function of wild type Atx2. Previous reports support
both alternatives. For example, protein misfolding and accumulation of Atx2 aggregates induced
by polyQ expansion has been linked to the diseases and cell death (56, 57). These phenotypes
can be reproduced by knockin of mutant Atx2 in wild type animals, but not by Atx2 knockout mice

(9). In contrast, other works have shown that toxicity of the polyQ Atx2 is independent of its ability
to aggregate (58). Our data can't evaluate the potential for gain of toxic of function. However, the
neurodevelopmental phenotypes we observe upon Atx2 loss in flies mirror observations in
neurodegenerative disease. This may support a potential contribution of loss of functional Atx2 to
neurodegeneration. The differences in phenotypes observed in Atx2 knockdown between flies and
rodents may be explained by the presence of Atx2-like, a highly homologous gene in rodents (59).
Atx2-like might compensate for the loss of function of wild type Atx2 during rodent development. It
would be very informative to revisit Atx2 knockout and polyQ knock-in phenotypes in the absence
of Atx2-like.
344
345  **Materials and Methods**
346
347  **Fly stocks and plasmids**
348
349  Fly stocks and crosses were cultured on standard cornmeal food at room temperature based on
350  the Bloomington Stock Center's recipe. Fly stocks can be found in table S1. All experiments used
351  Atx2 shRNA line 44012 unless otherwise stated. For tubulin photoconversion experiments, S2 cells
352  were transfected with pMT-EOS-tubulin described in Barlan et al (60).
353
354  ***Drosophila* Cell Culture: Primary Neurons and S2 Cells**
355
*Drosophila* S2 cells were maintained in Insect-Xpress medium (Lonza) at 25°C. These cells came
directly from DGRC and their identity was not subsequently confirmed. Transfections were carried
out with Effectene (Qiagen) according to the manufacturer's instructions. dsRNA was added to cells
on days 1 and 3 and imaging was carried out on day 5. For EB1 experiments a stable S2 cell line
was created expressing EB1-GFP under control of the EB1 promoter in the pMT vector. Primary
neuronal cultures were prepared by dissection of brains from 3rd instar larvae and dissociation of
tissue using liberase (Roche). Cells were plated on ConA coated glass coverslips and maintained
in Schneiders medium supplemented with 20% FBS, 5 µg/ml insulin, 100 µg/ml Pen-Strep, 50 µg/ml
Gentamycin and 10 µg/ml Tetracycline.
365
366  **Survival analysis**
367
368  For line 44012 (which we designated Atx2 shRNA line 1), homozygous adults were crossed with
369  elav-gal4 flies and number of adults eclosing from pupae cases was counted. For line 36114
370  (designated Atx2 shRNA line 2), heterozygous adults were crossed with elav-gal4 flies and number
371  of 'non tubby' adults eclosing from pupae cases. Elav-gal4 flies were used as control.
372
373  **Immunostaining, Western Blotting and Antibodies**
374
375  Antibodies and dyes used in this study are detailed in table S2. Full details of immunostaining and
376  Western Blotting can be found in the supplementary information.
377
378  **Microscopy and Image Analysis**
379
380  Full details of microscopy and image analysis can be found in the supplementary information.
381
382  **Statistical Analysis**

Statistical significance between two groups was determined using the unpaired, nonparametric Mann–Whitney *U* test. Data analyses were performed with Prism v6 (GraphPad Software). Statistical significance is presented as *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Data in Figure legends are presented as mean ± standard error.

**Data availability**
All data associated with this work are included in this article and SI Appendix.

**Funding sources and Acknowledgments**

We thank members of the Gelfand lab and Dr. M. Gelfand for helpful discussion. We thank Drs. A. Lim, W. Saxton and S. Rogers for *Drosophila* lines. Research reported in this study was supported by National Institute of General Medical Sciences Grants R01GM052111 and R35GM131752 (to V.I.G.).

**References**

1.  B. Bakthavachalu, *et al.*, RNP-Granule Assembly via Ataxin-2 Disordered Domains Is Required for Long-Term Memory and Neurodegeneration. *Neuron* **98**, 754-766.e4 (2018).
2.  C. McCann, *et al.*, The ataxin-2 protein is required for microRNA function and synapse-specific long-term olfactory habituation. *Proc. Natl. Acad. Sci. U. S. A.* **108** (2011).
3.  I. P. Sudhakaran, *et al.*, FMRP and Ataxin-2 function together in long-term olfactory habituation and neuronal translational control. *Proc. Natl. Acad. Sci. U. S. A.* **111** (2014).
4.  T. F. Satterfield, S. M. Jackson, L. J. Pallanck, A drosophila homolog of the polyglutamine disease gene SCA2 is a dosage-sensitive regulator of actin filament formation. *Genetics* **162**, 1687–1702 (2002).
5.  D. Lorenzetti, S. Bohlega, H. Y. Zoghbi, The expansion of the CAG repeat in ataxin-2 is a frequent cause of autosomal dominant spinocerebellar ataxia. *Neurology* **49**, 1009–1013 (1997).
6.  A. C. Elden, *et al.*, Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. *Nature* **466**, 1069–1075 (2010).
7.  K. J. Robinson, M. Watchon, A. S. Laird, Aberrant Cerebellar Circuitry in the Spinocerebellar Ataxias. *Front. Neurosci.* **14** (2020).
8.  G. Kim, O. Gautier, E. Tassoni-Tsuchida, X. R. Ma, A. D. Gitler, ALS Genetics: Gains, Losses, and Implications for Future Therapies. *Neuron* (2020) https:/doi.org/10.1016/j.neuron.2020.08.022 (November 5, 2020).
9.  T. R. Kiehl, *et al.*, Generation and characterization of Sca2 (ataxin-2) knockout mice. *Biochem. Biophys. Res. Commun.* **339**, 17–24 (2006).
10. A. Singh, *et al.*, ANTAGONISTIC ROLES FOR ATAXIN-2 STRUCTURED AND DISORDERED DOMAINS IN RNP CONDENSATION. *bioRxiv*, 2020.07.02.184796 (2020).
11. T. F. Satterfield, L. J. Pallanck, Ataxin-2 and its Drosophila homolog, ATX2, physically assemble with polyribosomes. *Hum. Mol. Genet.* **15**, 2523–2532 (2006).
12. S. F. Banani, H. O. Lee, A. A. Hyman, M. K. Rosen, Biomolecular condensates: Organizers of cellular biochemistry. *Nat. Rev. Mol. Cell Biol.* **18**, 285–298 (2017).
13. M. D. Stubenvoll, J. C. Medley, M. Irwin, M. H. Song, ATX-2, the C. elegans Ortholog of Human Ataxin-2, Regulates Centrosome Size and Microtubule Dynamics. *PLoS Genet.* **12** (2016).
14. M. M. Gnazzo, *et al.*, The RNA-binding protein ATX-2 regulates cytokinesis through PAR-5 and ZEN-4. *Mol. Biol. Cell* **27**, 3052–3064 (2016).
15. U. del Castillo, *et al.*, Conserved role for Ataxin-2 in mediating endoplasmic reticulum dynamics. *Traffic* **20**, 436–447 (2019).
16. D. A. SHOLL, Dendritic organization in the neurons of the visual and motor cortices of the

436    cat. *J. Anat.* **87**, 387–406 (1953).

437 17. U. Del Castillo, W. Lu, M. Winding, M. Lakonishok, V. I. Gelfand, Pavarotti/MKLP1
438    regulates microtubule sliding and neurite outgrowth in Drosophila neurons. *Curr. Biol.*
439    (2015) https:/doi.org/10.1016/j.cub.2014.11.008.

440 18. M. Winding, M. T. Kelliher, W. Lu, J. Wildonger, V. I. Gelfand, Role of kinesin-1–based
441    microtubule sliding in *Drosophila* nervous system development. *Proc. Natl. Acad. Sci.* **113**,
442    E4985–E4994 (2016).

443 19. R. Norkett, U. Del Castillo, W. Lu, V. I. Gelfand, Ser/Thr kinase Trc controls neurite
444    outgrowth in drosophila by modulating microtubule-microtubule sliding. *Elife* **9** (2020).

445 20. W. Lu, P. Fox, M. Lakonishok, M. W. Davidson, V. I. Gelfand, Initial neurite outgrowth in
446    Drosophila neurons is driven by kinesin-powered microtubule sliding. *Curr. Biol.* **23**, 1018–
447    1023 (2013).

448 21. W. Yu, *et al.*, Depletion of a microtubule-associated motor protein induces the loss of
449    dendritic identity. *J. Neurosci.* **20**, 5782–91 (2000).

450 22. U. Del Castillo, H. A. J. Müller, V. I. Gelfand, Kinetochore protein Spindly controls
451    microtubule polarity in Drosophila axons. *Proc. Natl. Acad. Sci. U. S. A.* **117** (2020).

452 23. A. L. Jolly, *et al.*, Kinesin-1 heavy chain mediates microtubule sliding to drive changes in
453    cell shape. *Proc. Natl. Acad. Sci.* **107**, 12151–12156 (2010).

454 24. C. Janke, M. M. Magiera, The tubulin code and its role in controlling microtubule
455    properties and functions. *Nat. Rev. Mol. Cell Biol.* **21**, 307–326 (2020).

456 25. A. Szyk, *et al.*, Molecular basis for age-dependent microtubule acetylation by tubulin
457    acetyltransferase. *Cell* **157**, 1405–1415 (2014).

458 26. D. Portran, L. Schaedel, Z. Xu, M. Théry, M. V. Nachury, Tubulin acetylation protects long-
459    lived microtubules against mechanical ageing. *Nat. Cell Biol.* **19**, 391–398 (2017).

460 27. A. Akhmanova, M. O. Steinmetz, Control of microtubule organization and dynamics: Two
461    ends in the limelight. *Nat. Rev. Mol. Cell Biol.* **16**, 711–726 (2015).

462 28. C. H. Lin, E. M. Espreafico, M. S. Mooseker, P. Forscher, Myosin drives retrograde F-actin
463    flow in neuronal growth cones. *Neuron* **16**, 769–782 (1996).

464 29. A. E. Twelvetrees, The lifecycle of the neuronal microtubule transport machinery. *Semin.*
465    *Cell Dev. Biol.* (2020) https:/doi.org/10.1016/j.semcdb.2020.02.008.

466 30. W. Lu, *et al.*, Competition between kinesin-1 and myosin-v defines drosophila posterior
467    determination. *Elife* **9** (2020).

468 31. L. C. Kapitein, *et al.*, Myosin-V opposes microtubule-based cargo transport and drives
469    directional motility on cortical actin. *Curr. Biol.* **23**, 828–834 (2013).

470 32. K. J. De Vos, M. Hafezparast, Neurobiology of axonal transport defects in motor neuron
471    diseases: Opportunities for translational research? *Neurobiol. Dis.* **105**, 283–299 (2017).

472 33. I. Noordstra, *et al.*, Control of apico–basal epithelial polarity by the microtubule minus-end-
473    binding protein CAMSAP3 and spectraplakin ACF7. *J. Cell Sci.* **129**, 4278–4288 (2016).

474 34. R. Banerjee, J. A. White, S. Gunawardena, "In vivo visualization of moving synaptic cargo
475    complexes within Drosophila larval segmental axons" in *Methods in Molecular Biology*,
476    (Humana Press Inc., 2020), pp. 293–300.

477 35. W. Lu, P. Fox, M. Lakonishok, M. W. Davidson, V. I. Gelfand, Initial neurite outgrowth in
478    drosophila neurons is driven by kinesin-powered microtubule sliding. *Curr. Biol.* **23**, 1018–
479    1023 (2013).

480 36. R. Norkett, U. Del Castillo, W. Lu, V. I. Gelfand, Ser/Thr kinase Trc controls neurite
481    outgrowth in Drosophila by modulating microtubule-microtubule sliding. *Elife* **9** (2020).

482 37. U. Del Castillo, R. Norkett, V. I. Gelfand, Unconventional Roles of Cytoskeletal Mitotic
483    Machinery in Neurodevelopment. *Trends Cell Biol.* **29**, 901–911 (2019).

484 38. S. Lin, M. Liu, O. I. Mozgova, W. Yu, P. W. Baas, Mitotic Motors Coregulate Microtubule
485    Patterns in Axons and Dendrites. *J. Neurosci.* **32**, 14033–14049 (2012).

486 39. Y. Zheng, *et al.*, Dynein is required for polarized dendritic transport and uniform
487    microtubule orientation in axons. *Nat. Cell Biol.* **10**, 1172–1180 (2008).

488 40. D. H. Roossien, P. Lamoureux, K. E. Miller, Cytoplasmic dynein pushes the cytoskeletal
489    meshwork forward during axonal elongation. *J. Cell Sci.* **127**, 3593–3602 (2014).

490 41. M. J. Papandréou, C. Leterrier, The functional architecture of axonal actin. *Mol. Cell.*
491    *Neurosci.* **91**, 151–159 (2018).

42.    I. Hahn, *et al.*, Tau, XMAP215/Msps and Eb1 jointly regulate microtubule polymerisation
and bundle formation in axons. *bioRxiv*, 2020.08.19.257808 (2020).

43.    F. Bradke, C. C. Dotti, The role of local actin instability in axon formation. *Science (80-. ).*
**283**, 1931–1934 (1999).

44.    F. Korobova, T. Svitkina, Arp2/3 complex is important for filopodia formation, growth cone
motility, and neuritogenesis in neuronal cells. *Mol. Biol. Cell* **19**, 1561–1574 (2008).

45.    A. W. Schaefer, *et al.*, Coordination of Actin Filament and Microtubule Dynamics during
Neurite Outgrowth. *Dev. Cell* **15**, 146–162 (2008).

46.    V. K. Godena, *et al.*, Increasing microtubule acetylation rescues axonal transport and
locomotor deficits caused by LRRK2 Roc-COR domain mutations. *Nat. Commun.* **5**, 1–11
(2014).

47.    N. Kaul, V. Soppina, K. J. Verhey, Effects of α-tubulin K40 acetylation and detyrosination
on kinesin-1 motility in a purified system. *Biophys. J.* **106**, 2636–2643 (2014).

48.    J. C. Fiala, M. Feinberg, A. Peters, H. Barbas, Mitochondrial degeneration in dystrophic
neurites of senile plaques may lead to extracellular deposition of fine filaments. *Brain*
*Struct. Funct.* **212**, 195–207 (2007).

49.    S. C. Correia, G. Perry, P. I. Moreira, Mitochondrial traffic jams in Alzheimer's disease -
pinpointing the roadblocks. *Biochim. Biophys. Acta - Mol. Basis Dis.* **1862**, 1909–1917
(2016).

50.    Y. Xie, *et al.*, Endolysosomal Deficits Augment Mitochondria Pathology in Spinal Motor
Neurons of Asymptomatic fALS Mice. *Neuron* **87**, 355–370 (2015).

51.    N. Hirokawa, Y. Tanaka, Kinesin superfamily proteins (KIFs): Various functions and their
relevance for important phenomena in life and diseases. *Exp. Cell Res.* **334**, 16–25
(2015).

52.    S. L. Reck-Peterson, W. B. Redwine, R. D. Vale, A. P. Carter, The cytoplasmic dynein
transport machinery and its many cargoes. *Nat. Rev. Mol. Cell Biol.* **19**, 382–398 (2018).

53.    M. A. Tischfield, *et al.*, Human TUBB3 Mutations Perturb Microtubule Dynamics, Kinesin
Interactions, and Axon Guidance. *Cell* **140**, 74–87 (2010).

54.    M. Ralser, *et al.*, Ataxin-2 and huntingtin interact with endophilin-A complexes to function
in plastin-associated pathways. *Hum. Mol. Genet.* **14**, 2893–2909 (2005).

55.    M. D. Figley, G. Bieri, R.-M. Kolaitis, J. P. Taylor, A. D. Gitler, Neurobiology of Disease
Profilin 1 Associates with Stress Granules and ALS-Linked Mutations Alter Stress Granule
Dynamics (2014) https:/doi.org/10.1523/JNEUROSCI.0543-14.2014.

56.    S. W. Davies, *et al.*, Formation of neuronal intranuclear inclusions underlies the
neurological dysfunction in mice transgenic for the HD mutation. *Cell* **90**, 537–548 (1997).

57.    L. T. Pflieger, *et al.*, Gene co-expression network analysis for identifying modules and
functionally enriched pathways in SCA2. *Hum. Mol. Genet.* **26**, 3069–3080 (2017).

58.    D. P. Huynh, K. Figueroa, N. Hoang, S. M. Pulst, Nuclear localization or inclusion body
formation of ataxin-2 are not necessary for SCA2 pathogenesis in mouse or human. *Nat.*
*Genet.* **26**, 44–50 (2000).

59.    C. Kaehler, *et al.*, Ataxin-2-Like Is a Regulator of Stress Granules and Processing Bodies.
*PLoS One* **7** (2012).

60.    K. Barlan, W. Lu, V. I. Gelfand, The Microtubule-Binding Protein Ensconsin Is an Essential
Cofactor of Kinesin-1. *Curr. Biol.* **23**, 317–322 (2013).

536
537
538

# Figure 1

Figure 1: Ataxin-2 is essential for neurodevelopment, locomotion and viability in *Drosophila.*

A. Atx2 depletion is lethal at the pupae stage. Percentage of adults eclosing from pupae cases (Control = 99%, Atx2 RNAi line 1 = 0.0%, Atx2 RNAi line 2 = 0.0%). B. Atx2 depletion impairs locomotion. Example crawling tracks of control and elav>Atx2 RNAi 3rd instar larvae. C. Crawling velocity of control and elav>Atx2 RNAi 3rd instar larvae (Control = 0.71 mm/s ± 0.05, Atx2 RNAi = 0.03 mm/s ± 0.003). D. Atx2 depletion decreases brain size and innervation. Example images of control and elav>Atx2 RNAi brains from *Drosophila* 3rd instar larvae. Scale bar, 100μm. E. Size of optic lobes in control and elav>Atx2 RNAi brains (Control = 32931 μm$^2$ ± 1993, Atx2 RNAi = 17306 μm$^2$ ± 2028, n=18 control and 9 Atx2 RNAi brains, p<0.0001). F. Area of optic stalks normalized to optic lobes in control and Atx2 RNAi brains (Control = 0.212 ± 0.01, Atx2 RNAi = 0.026 ± 0.003, n=18 control and 9 Atx2 RNAi brains, p<0.0001). G. Atx2 depletion decreases dendrite development. Example images of DA neurons in control and elav>Atx2 RNAi neurons. Scale bar, 50μm. H. Dendrite length per cell in control and elav>Atx2 RNAi neurons (Control = 4500 μm ± 644, Atx2 RNAi = 1290 μm ± 134, n=11 animals). I. Sholl analysis of control and Atx2 RNAi neurons DA neurons showing number of intersections with distance from soma.

# Figure 2

558     Figure 2: Ataxin-2 depletion decreases microtubule dynamics.
559
A. Example images showing Atx2 depletion in *Drosophila* S2 cells increases microtubule
acetylation as quantified in B (Signal intensity Control = 2061210 ± 204596 A.U. Atx2 RNAi =
6286404 ± 608574). C. Example images showing Atx2 depletion increases microtubule
acetylation in *Drosophila* neurons in culture (scale bar, 10μm), as quantified in D (Signal Intensity
Control = 1.0 ± 0.04, n = 23 cells, Signal Intensity Atx2 RNAi = 1.3 ± 0.10, n = 27 cells, p = 0.01).
E. Western blot of Acetylated tubulin levels in Control and Atx2 RNAi brain lysate from *Drosophila*
3$^{rd}$ instar larvae. F. Example images of EOS-tubulin in S2 cells 0 and 30mins after
photoconversion (scale bar, 10μm) as quantified in G (Motile fraction Control = 0.52 ± 0.03, n =
21 cells, Atx2 RNAi = 0.26s ± 0.04, n = 22 cells, p < 0.0001). H. Example images of tubulin
staining in S2 cells showing microtubules after 25μM vinblastine treatment 1hr. Scale bar, 10μm.
I. Quantification of cells with an intact microtubule network after vinblastine treatment (Control =
26.4% ± 9.4, Atx2 RNAi = 71.9% ± 3.1). J. Temporal color code images showing motility of EB1-
GFP comets in S2 cells over 120 seconds. Scale bar, 10μm. K. Quantification of number of EB1
comets per cell. L. Temporal color code images showing motility of EB1-GFP comets in cultured
neurons over 120 seconds. Scale bar, 10μm, inset, 5μm. M. Kymographs showing motility of
comets in neuronal processes over 120 seconds. Scale bar, 10μm. N. Quantification of number of
EB1 comets per process length (Control = 0.53 ± 0.03, n=31, Atx2 RNAi = 0.37 ± 0.02, n=34, p =
0.0004).

# Figure 3

Figure 3: Ataxin-2 depletion stabilizes the actin cytoskeleton
A. Example images showing phalloidin staining for F-actin in control and Atx2 RNAi treated
*Drosophila* S2 cells treated with DMSO or 10μM LatB for 1hr. Scale bar, 10μm B. Normalized
Phalloidin Rhodamine signal (Control = 1.00 ± 0.19, Atx2 RNAi = 1.15 ± 0.11, Control + LatB =
0.65 ± 0.07, Atx2 RNAi + LatB = 1.02 ± 0.11, n = 15-24 cells, $p = 0.03$). C. Example images
showing phalloidin staining in control and Atx2 RNAi neurons treated with DMSO or 10μM LatB
for 1hr. Scale bar, 10μm. D. Quantification of Phalloidin Rhodamine intensity along length of
neuron (Signal Intensity Control LatB = 440.4 ± 53, n = 37 cells, Signal Intensity Atx2 RNAi LatB
= 786.5 ± 128, n = 27 cells, $p = 0.02$). E. Example phase contrast images showing filipodia from
neuronal processes in control and Atx2 RNAi neurons. Scale bar, 10μm. F. Average number of
filopodia per 10μm process length (Control = 2.8 ± 0.19, n = 28 cells, Atx2 RNAi = 1.2 ± 0.15, n =
28 cells, $p < 0.0001$).
599

# Figure 4

Figure 4: Ataxin-2 depletion inhibits organelle transport

A. Schematic showing region of imaging in neurons in culture. B. Example kymographs showing mitochondria motility in processes of control and Atx2 RNAi neurons. Horizontal scale bar, 10µm.

Vertical scale bar, 20s. C. Example kymographs showing lysosome motility in processes of
control and Atx2 RNAi neurons. Horizontal scale bar, 10μm. Vertical scale bar, 20s. D.
Quantification of motile mitochondria in control and Atx2 knockdown neurons (Control = 47% ±
3.7, n = 25 cells, Atx2 RNAi = 15.7% ± 2.5, n = 28 cells, p <0.0001). E. Quantification of motile
lysosomes in control and Atx2 knockdown neurons (Control = 31% ± 2.1, n = 25 cells, Atx2 RNAi
= 14.5% ± 2.4, n = 29 cells, p <0.0001). F. Schematic image of a third instar larva showing
segmental nerves. Example kymographs showing mitochondria motility (G) and dense core
vesicle motility (H) in segmental nerves *in vivo* in control and Atx2 RNAi larvae. Vertical scale
bars, 10s, horizontal scale bars,10μm for dense core vesicles, 20μm for mitochondria. I.
Quantification of motile mitochondria in control and Atx2 knockdown segmental nerves *in vivo*
(Control = 50% ± 12.5, n = 3 larvae, Atx2 RNAi = 1.5% ± 1.5, n = 3 larvae, p = 0.015). J.
Quantification of motile DCVs in control and Atx2 knockdown segmental nerves *in vivo* (Control =
66% ± 1.7, n = 9 larvae, Atx2 RNAi = 18.6% ± 7.4, n = 8 larvae, p < 0.0001). K. Example image
showing mitochondria and lysosome distribution in Atx2 RNAi cultured neuron. Scale bar, 10μm,
inset 5μm. L. Example super resolution images of microtubules in varicosities in Atx2 RNAi
neurons. Scale bars, 10μm.
621

628 Supplementary Information for

629 **Ataxin-2 is essential for neurodevelopment in *Drosophila* and is a major regulator of the**

630 **cytoskeleton**

631
632
633

635 Fig. S1: Ataxin-2 depletion in brains by western blot. Western blot demonstrating knockdown of
636 Atx2 by elav>Atx2 shRNA #44012.

638
639 Fig. S2: Microtubule content is equal control vs Atx2 RNAi neurons in culture. A. Example images
640 showing tubulin staining in extracted neurons. Scale bar, 20µm. B. Quantification of tubulin signal
641 intensity (Ctrl = 488.4 ± 67.7, n = 8 cells, Atx2 RNAi = 441.2 ± 76.1, n = 6 cells, NS p = 0.85)

642
643
644
645

646 Supplementary methods
647
648 **Immunostaining, Western Blotting and Antibodies**
649
Optic lobes, dissected from third-instar larva in 1 × PBS, were fixed in 4% (wt/wt) formaldehyde
(methanol-free) in PBT (1 × PBS, 0.1% Triton X-100) for 20 min; brains were washed five times
with PBTB (PBT + 0.2% bovine serum albumin) for 10 min and blocked in 5% (vol/vol) normal goat
serum for 1 h. Samples were incubated with primary anti-Fasciclin II antibody (1:40) at 4° C
overnight, washed and incubated with secondary anti-mouse for 2h at room temperature (RT).
Finally, samples were washed with PBT for 10min five times before mounting.
For immunostaining of S2 cells and cultured neurons, cells were fixed 5 mins in methanol (S2 cells)
or 1% glutaraldehyde and quenched in 2 mg/ml sodium borohydride (neurons). Cells were
permeabilised and blocked in wash buffer (1% BSA, 0.1% Triton X-100 in TBS) for 30mins and
incubated with primary antibody in wash buffer (Anti-Acetylated tubulin 1:100-1:250, DM1a 1:1000).
Coverslips were washed 3x in wash buffer and incubated with secondary antibodies (1:200). For
extraction, cells were incubated in extraction buffer (1% Triton X-100, 1uM taxol, 30% glycerol in
BRB-80) for 2 minutes, then in extraction buffer with 1% glutaraldehyde for 2 minutes, then 1%
glutaraldehyde in PBS for 3 minutes followed by reduction as above. For western blotting, dissected
third instar larvae brains or S2 cell lysate were prepared in 1% SDS, boiled in laemmli buffer and
analyzed by SDS-PAGE.
666
667 **Microscopy and Image Analysis**
668
To image DA neurons or S2 cells we used a Nikon Eclipse U2000 inverted microscope equipped
with a Yokogawa CSU10 spinning disk head with a perfect focus system and a 40x 1.30 or 100x
1.45 oil immersion lens, using Evolve EMCCD (Photometrics) and controlled by Nikon Elements
4.00.07 software. For EB1 imaging in S2 cells images were acquired at 1 fps. For organelle imaging
in larvae, larvae were immobilised and frames were acquisition was at 3 fps (dense core vesicles)
or 0.5 fps (mitochondria). Immunostained brains and neurons were imaged on a Nikon Ti2 inverted
microscope equipped with W1 spinning disk driven by Nikon Elements 5.20 using a 20X or 100x
1.45 oil immersion lens and CMOS prime 95B sensor (Photometrics). For microtubule organization
images, this set-up was used with a Live-SR (Gataca systems). Live imaging of lysosomes and
EB1 comets in neurons was carried out at 1fps. For photoconversion experiments we applied 405
nm light from a light-emitting diode light source (89 North Heliophor) for 5 s, using an adjustable
diaphragm to constrain light and illuminate only a specific region of interest. After photoconversion,
images were collected every ten minutes for 30 minutes. Imaging of filopodia, organelle distribution
in culture and mitochondrial trafficking was carried out on a Nikon Ti inverted microscope with a
Hamamatsu Orca Flash sensor. For locomotion assays, 3rd Instar larvae were allowed to crawl on
the surface of a petri dish containing 2% agarose supplemented with apple juice and charcoal
powder. Trajectories are presented using the Physics Temporal-Color Code plugin in Fiji.
686
Image analysis was carried out in Fiji unless otherwise stated. Total neurite length in DA neurons
was measured using the 'curve tracing' and sholl analysis plugins. Tubulin motile fraction was
calculated as 'photoconverted signal outside original photoconversion zone / original
photoconverted signal' at 30mins post photoconversion. Numbers and motility of EB1 comets,
mitochondria and lysosomes in culture or *in vivo* was carried out by generating kymographs and
analyzed using the 'kymobutler' software (1). Motile organelles were defined as those that moved
more than 2μm over the course of the movie. Comets that moved more than 1μm over the course

of the movie were counted. EB1 comets in S2 cells were counted using Nikon elements analysis
software (v5.20). Numbers of filopodia and organelles in varicosities were scored manually.
Table S1: Fly stocks and sources used in this study.
697

| Line | Identifier | Source |
|---|---|---|
| Ataxin-2 RNAi TRiP line #1 (Val20, second chromosome attP40 insertion targeting CDS 561-581) | 44012 | Bloomington Drosophila Stock Center (BDSC) |
| Ataxin-2 RNAi TRiP line #2 (Val20, third chromosome attP2 insertion targeting Atx2 CDS 474-494) | 36114 | BDSC |
| elav-gal4 | 8760 | BDSC |
| ppk-CD4-tomato | 35844 | BDSC |
| UASt-mCherry.mito.OMM | 66532 | BDSC |
| Ubi-EB1-GFP | | S. Rogers |
| D42 gal4>UAS ANF GFP | | A. Lim, W. Saxton (2) |

698
699
700 Table S2: Antibodies used in this study.
701

| Antibody | Source | Concentration |
|---|---|---|
| Anti FasII 1D4 | DSHB | 1:40 |
| Anti Acetylated alpha tubulin (K40) | protein tech 66200 | 1:100-1:250 |
| Anti alpha tubulin DM1a | Prepared in house | 1:1000 |
| Anti tubulin rb polyclonal | made in house | 1:1000 |
| Anti Hsc 70 K-19 | Santa Cruz | 1:200 |
| HRP-conjugated anti mouse | Jackson | 1:10,000 |
| HRP-conjugated anti rabbit | Jackson | 1:10,000 |
| HRP-conjugated anti goat | Jackson | 1:10,000 |
| Alexafluor 647 conjugated anti mouse IgG | Jackson | 1:250 |
| Lysotracker DND 99 | Invitrogen | 1:10,000 |

| Rhodamine conjugated Phalloidin | Jackson | 1:1000 |
| --- | --- | --- |

703
**Video Legends**
705

706  Movie 1: Crawling behavior of control (elav-gal4) and Atx2 RNAi (elav>Atx2 RNAi) *Drosophila* 3$^{rd}$
707  Instar Larvae, Related to Figure 1B.
708

709  Movie 2: Time-lapse of control (untreated cell) or Atx2 RNAi S2 cells expressing EOS-tub. Scale
710  bar, 10μm. Related to Fig. 2F.
711

712  Movie 3: Time-lapse of control (untreated cell) or Atx2 RNAi S2 cells expressing EB1-GFP. Scale
713  bar, 10μm. Related to Fig. 2J.
714

715  Movie 4: Time-lapse of primary neurons expressing EB1-GFP of two different genotypes. Control
716  (elav-gal4) and Atx2 RNAi (elav>Atx2 RNAi). Scale bar, 10μm. Related to Fig. 2L.
717

718  Movie 5: Transport of mitochondria in primary cultures of control and Atx2 RNAi expressing
719  MitomCherry. Scale bar, 10μm. Related to Fig. 4B.
720

721  Movie 6: Transport of lysosomes in primary cultures of control and Atx2 RNAi labelled with
722  lysotracker. Scale bar, 10μm. Related to Fig. 4C.
723

724  Movie 7: Transport of mitochondria in motor segmental nerves of control and Atx2 RNAi
725  expressing mitoGFP. Scale bar, 10μm. Related to Fig. 4G.
726

727  Movie 8: Transport of Dense core vesicles in motor segmental nerves of control and Atx2
728  RNAi expressing ANF-GFP. Scale bar, 10μm. Related to Fig. 4G.
729

730 Supplementary References

731 1. M. A. Jakobs, A. Dimitracopoulos, K. Franze, Kymobutler, a deep learning
732 software for automated kymograph analysis. *Elife* **8** (2019).

733 2. A. Lim, A. Rechtsteiner, W. M. Saxton, Two kinesins drive anterograde
734 neuropeptide transport. *Mol. Biol. Cell* **28**, 3542–3553 (2017).

737