A simple *Agrobacterium*-mediated stable transformation technique for the hornwort model *Anthoceros agrestis*

Eftychios Frangedakis[1], Manuel Waller[2,3], Tomoaki Nishiyama[4], Hirokazu Tsukaya[5], Xia Xu[6], Yuling Yue[2,3], Michelle Tjahjadi[6], Andika Gunadi[6], Joyce Van Eck[6,7], Fay-Wei Li[6,8], Péter Szövényi[2,3]* and Keiko Sakakibara[9]**

[1]Department of Plant Sciences, University of Cambridge, Cambridge, CB3 EA, UK

[2]Department of Systematic and Evolutionary Botany, University of Zurich, Switzerland

[3]Zurich-Basel Plant Science Center, Zurich, Switzerland

[4]Advanced Science Research Center, Kanazawa University, Ishikawa, Japan

[5]Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Tokyo, 113-0033, Japan

[6]Boyce Thompson Institute, Ithaca, New York, USA

[7]Plant Breeding and Genetics Section, Cornell University, Ithaca, New York, USA

[8]Plant Biology Section, Cornell University, Ithaca, New York, USA

[9]Department of Life Science, Rikkyo University, Tokyo, Japan

*corresponding author email address: peter.szoevenyi@uzh.ch

**corresponding author email address: bara@rikkyo.ac.jp

Funding information

Japanese Society for the Promotion of Science (JSPS) Short Term Postdoctoral Fellowship grant

no. PE14780 to E.F. MEXT and JSPS Grants-in-Aid for Scientific Research on Innovation Areas

nos. 25113001 and 19H05672 to H.S., JSPS KAKENHI 26650143 and 18K06367 to K.S.,

15H04413 to T.N., 19K22448 to T.N. and K.S. Swiss National Science Foundation grants 160004,

131726 and 184826 (to P.S.), The Deutsche Forschungsgemeinschaft (DFG – German Research

Foundation) under the Priority Programme "MAdLand – Molecular Adaptation to Land: Plant

Evolution to Change" (SPP 2237, 440370263, to P.S.), The Georges and Antoine Claraz

Foundation (to P.S., M.W and Y.Y), The University Research Priority Program 'Evolution in Action'

of the University of Zurich to P.S. UZH Forschungskredit Candoc grant no. FK-19-089 to

M.W.National Science Foundation IOS-1923011 to F.-W.L. and J.V.E.

Author Contributions

KS, EF, HT, MW and PS, conceived and designed the experiments. TN and MW identified gene

promoter regions. EF and MW performed cloning. EF generated and characterized the transgenic

lines and performed imaging. XX, MT and JVC generated the BTI lines and AG imaged the lines.

MW, PS, FW and AG analyzed the transcriptomic data. MW performed NGS of transgenic lines,

PS and MW assembled the genomes. AG and MW confirmed the insert locations. YY provided

technical assistance. KS, EF, TN, MW and PS wrote the article with contributions from all the

authors.

Abstract

We have developed a simple *Agrobacterium*-mediated method for the stable transformation of the hornwort *Anthoceros agrestis*, the fifth bryophyte species for which a genetic manipulation technique becomes available. High transformation efficiency was achieved by using thallus tissue grown under low-light conditions. We generated a total of 216 transgenic *A. agrestis* lines expressing the β-Glucuronidase (GUS), cyan, green, and yellow fluorescent proteins under the control of the CaMV 35S promoter and several endogenous promoters. Nuclear and plasma membrane localization with multiple color fluorescent proteins was also confirmed. The transformation technique described here should pave the way for detailed molecular and genetic studies of hornwort biology, providing much needed insight into the molecular mechanisms underlying symbiosis, carbon-concentrating mechanism, RNA editing, and land plant evolution in general.

Introduction

The hornworts are one of the three lineages of bryophytes that diverged from liverworts and mosses about 460 million years ago (Morris et al., 2018; One Thousand Plant Transcriptomes Initiative, 2019; Li et al., 2020). While having only around 220 extant species (Söderström et al., 2016), hornworts are key to address diverse questions about land plant evolution and terrestrialization. Hornworts display a unique combination of features (Frangedakis et al., 2020) such as a sporophyte, that is produced by an indeterminate basal meristem, and bears stomata similar to mosses and vascular plants (Renzaglia et al., 2017). In addition, it is the only extant land plant lineage (together with a few *Selaginella* species (Liu et al., 2020)), that has a single (or just a few) algal-like chloroplast(s) per cell. The chloroplasts resemble those of algae in that they may contain pyrenoids, a carbon-concentrating structure that is shared with many algal lineages (Villarreal and Renner, 2012; Li et al., 2017). Hornwort plastids are also unique by exhibiting the highest RNA editing rates amongst land plants (Yoshinaga, 1996; Yoshinaga, 1997; Kugita, 2003; Small et al., 2019). Finally, hornworts are among the very few plant lineages that can establish symbiotic relationships with both endophytic cyanobacteria (Renzaglia et al., 2009) and various glomeromycotina and mucoromycotina fungal partners (mycorrhiza) (Desirò et al., 2013).

Anthoceros agrestis has been established as an experimental model system for hornworts and
two isolates are currently available (Oxford and Bonn) (Szövényi et al., 2015). *A. agrestis,* like
other bryophytes*,* has a haploid-dominant life cycle through which the haploid gametophyte phase
alternates with the diploid sporophyte phase. The life cycle of *A. agrestis* starts with the
germination of the haploid spores which develop into an irregularly shaped thallus (Fig. 1 A-C).
Sexual reproduction occurs through fusion of the egg (produced in archegonia) and the motile
sperm (produced in antheridia). The resulting embryo develops within the gametophyte and gives
rise to the sporophyte (Fig. 1 D). *A. agrestis* can be easily grown under laboratory conditions and
its haploid-dominant life cycle makes genetic analysis straightforward. The nuclear genome of *A.*
agrestis was recently sequenced (Li et al., 2020) which is one of the smallest genomes amongst
land plants. The species is monoicous, with male and female reproductive organs produced by
the same individual. The sexual life cycle of *A. agrestis* can be completed under laboratory
conditions within approximately 2-3 months (Szövényi et al., 2015). However, transformation of
A. agrestis was not feasible until recently, posing a major obstacle to the analysis of gene function
in hornworts, and more generally to land plant evo-devo studies.

Several approaches have been used for gene delivery in bryophytes, including polyethylene
glycol (PEG)-mediated uptake of DNA by protoplasts (Schaefer et al., 1991), particle
bombardment (Cho et al., 1999; Chiyoda et al., 2008) and *Agrobacterium tumefaciens*-mediated
(revised scientific name *Rhizobium radiobacter* (Young et al., 2001)) transformation (Ishizaki et
al., 2008; Kubota et al., 2013; Althoff and Zachgo, 2020). *Agrobacterium*-mediated transformation
is a commonly used method for various plant species (Gelvin, 2003), it is relatively simple and
does not require specialised or expensive equipment. In addition, *Agrobacterium*-mediated
transformation has several advantages over other transformation methods, such as the
integration of a lower number of transgene copies into the plant genome and the ability to transfer
relatively large DNA segments with intact transgene genome integration.

In this study we report the first successful stable genetic transformation method for hornworts.
The method is based on *Agrobacterium*-mediated transformation of *A. agrestis* thallus*.* We also
report the successful expression and targeting of four different fluorescent proteins in two different
cellular compartments, the plasma membrane and the nucleus. Finally, we characterize a number
of native *A. agrestis* promoters for their potential to drive strong constitutive transgene expression
that can be useful for future hornwort genetic studies.

Results

We tested the potential of the *Agrobacterium*-mediated gene delivery method to recover stable *A.*
agrestis transgenic lines. There are several critical factors that determine the efficiency of
Agrobacterium-mediated gene delivery: (1) the selection of appropriate plant tissue for infection
and conditions of tissue preparation, (2) the type and concentration of antibiotics applied to select
for transgenic lines, (3) the choice of transformation vectors, (4) the choice of *Agrobacterium*
strains, and (5) the conditions used for co-cultivation. Each of these factors is examined in this
study.

Selection of tissue and optimal growth conditions

A. agrestis gametophyte thallus was chosen as the appropriate tissue for transformation, because
it is easily accessible, has a remarkable regenerative capacity and is haploid. *A. agrestis* thallus
tissue cultures can easily be propagated and maintained by transfer of small thallus fragments
(approximately 1 x 1 mm) onto fresh growth medium on a monthly basis (Supplemental Fig. S1).
A. agrestis is similar to several other bryophytes in that an entire plant can regenerate from a
small thallus fragment. This is in striking contrast with vascular plants, where usually the transition
to an undifferentiated tissue state (callus), after treatment with extrinsic hormones such as auxin
and cytokinin, is necessary before the regeneration of new plant tissue (Ikeuchi et al., 2013). The
use of thallus has the additional advantage that the resulting transformants have a uniform genetic
background.

We reasoned that similar to the liverwort *Marchantia polymorpha* (Kubota *et al.*, 2013),
fragmented tissue will be susceptible to *Agrobacterium* infection. In the case of *M. polymorpha,*
the apical part of the thallus is removed to induce regeneration, followed by co-cultivation with
Agrobacterium to generate transformed plants. *A. agrestis* thallus regeneration is similarly
induced by fragmentation and presumably removal of the apical parts of the thallus.
Consequently, a transformation approach similar to the one used for *M. polymorpha* was utilized
and adapted to *A. agrestis*. However, unlike in *M. polymorpha*, the *A. agrestis* notch area and
apical cells are not easily distinguishable, thus determining which part of the thallus should be
removed is not easy. Therefore, we tested whether homogenization using dispensing tools is a
suitable method for thallus tissue fragmentation. Different speed levels and duration of
homogenization were examined. We found that homogenization of 0.5 g of thallus tissue in 15

mL of sterile water for 5 seconds (see Materials and methods) is sufficient to fragment the tissue
without damaging the plants and results in rapid tissue regeneration.

In addition, we found that the light intensity used to cultivate *A. agrestis* tissue is a critical factor
for successful transformation (Fig. 1 E-G). Tissue that was grown under low light conditions, even
though smaller in size, had a more regular and flattened shape and was optimal for
transformation. (Fig. 1 F-G and Supplemental Fig. S1 A). When tissue was grown under high light
intensity (above 40 µmol m^{-2} s^{-1}), no transformants were obtained.

Selection of appropriate antibiotics

A selectable marker gene, most commonly one that confers antibiotic-resistance, is necessary for
efficient recovery of stable transgenic lines following co-cultivation with *Agrobacterium*. To identify
antibiotics and their appropriate concentration with cytotoxic effect on *A. agrestis*, untransformed
A. agrestis (Oxford and Bonn) thallus fragments were subjected to different concentrations of
hygromycin and geneticin/G418 (an analogue of neomycin and kanamycin). The tested
concentrations ranged from 0 µg/mL - 20 µg/mL for hygromycin and 0 µg/mL - 150 µg/mL for
geneticin/G418. We found that a 3-week incubation period with 10 µg/mL hygromycin was
sufficient to inhibit growth of untransformed thallus tissue (Supplemental Fig. S2 and S3), whereas
thallus tissue was not susceptible to geneticin/G418 even when supplied in high concentration
(150 µg/mL) (Supplemental Fig. S4). Thus, we selected hygromycin as an appropriate selection
agent for *A. agrestis* transformation.

Preliminary tests with the GUS reporter

Preliminary transformation experiments were performed using the Oxford isolate and the
pCAMBIA1305.2 plasmid containing the *hygromycin B phosphotransferase* gene (*hph*, conferring
hygromycin resistance) driven by the Cauliflower mosaic virus 35S (CaMV 35S, hereafter called
35S) promoter, terminated with a 35S polyadenylation signal, and a *p-35S_s::GUSPlus* (β-
Glucuronidase) transcription unit. *GUSPlus* contains a catalase intron to ensure that the observed
GUS expression is not due to the *Agrobacterium*.

Homogenized regenerating thallus tissue was co-cultivated with the *Agrobacterium* AGL1 strain

containing the pCAMBIA1305.2 plasmid, as well as *Agrobacterium* without a transformation

vector as a negative control, in liquid KNOP media supplemented with sucrose. Media were also

supplemented with 3',5'-dimethoxy-4'-hydroxyacetophenone (acetosyringone) since phenolic

compounds such as acetosyringone have been shown to be important for the virulence genes

activation (Stachel et al., 1985). Co-cultivation duration was 3 days at 22°C on a shaker without

any light supplementation (only ambient light from the room). After co-cultivation, the tissue was

spread on solid KNOP plates supplemented with cefotaxime and hygromycin. After 3-4 weeks the

tissue was transferred to fresh selective media (transformation outline in Fig. 2 and Supplemental

Fig. S5). Emergence of rhizoids on surviving tissue fragments is a reliable indicator of successful

transformation events (Fig. 3A-B). One to two months later successful transformants (plant

fragments producing rhizoids) were visible. Finally, surviving plants were subjected to a third

round of antibiotic selection to ensure false positives were eliminated. Thallus surviving selection

on hygromycin exhibited GUS expression (Fig. 3C, Supplemental Fig. S6). No mock-transformed

plants survived antibiotic selection; e.g. when transformation was carried out using

Agrobacterium lacking the transformation vector. These results indicated that *A. agrestis* Oxford

thallus tissue is susceptible to *Agrobacterium* infection and that the 35S promoter driving the *hph*

gene is sufficient for selection of transformants.

Tests with eGFP as a reporter

Tests with eGFP as a reporter

Subsequent experiments were carried out using the *A. agrestis* Oxford isolate and a construct

containing the enhanced Green Fluorescent Protein (eGFP) reporter gene (Cormack et al., 1996).

eGFP makes the identification of successful transformation events easier without the need of

laborious GUS staining. For the construction of the eGFP transformation vector, we used the

OpenPlant toolkit (Sauret-Gueto et al., 2020) which is based on the Loop assembly Type IIS

cloning system (Pollak et al., 2019). All the DNA parts described here are generated following the

common syntax (Patron et al., 2015) and are compatible with Type IIS cloning systems, such as

GoldenGate and Loop assembly, facilitating the exchange of DNA parts between different

laboratories. The transformation vector contained the *hph* gene driven by the 35S promoter and

terminated with a 35S polyadenylation signal. It also contained a *p-35S_s::eGFP-Lti6B*

transcription unit (same 35S promoter with the one driving *GUSPlus* in pCAMBIA1305.2 plasmid)

terminated by the double *nopaline synthase* (*Nos*) - 35S polyadenylation signal (Sauret-Güeto et

al., 2020) (Fig. 3E) which was fused to the Low Temperature Induced Protein 6B (Lti6B) signal

for membrane localization from *Arabidopsis thaliana* (Arabidopsis) (Cutler et al., 2000). The vector

was transformed into *A. agrestis* using the method described above and eGFP was successfully

expressed in *A. agrestis* with the expected localization in the plasma membrane (Fig. 3D, E and

I). During the course of this study, we generated 157 stable *A. agrestis* transgenic lines expressing

the *p-35S_s::eGFP-Lti6B* (Supplemental Table I). There is variability in eGFP expression patterns

between different transgenic lines (Supplemental Fig. S8) presumably due to differences in the

transgene copy number or the genome location of transgene insertion. A small fraction of

hygromycin resistant lines (four out of 157) do not show eGFP fluorescence which could be

attributed to potential silencing events or truncation of the inserted T-DNA. These plants have

been through at least five rounds of hygromycin antibiotic selection so it is unlikely they are false

positives. A total of 15 lines have been propagated vegetatively for more than two years without

abolishing transgene expression.

225   Testing additional fluorescent proteins, the Bonn isolate and transgene inheritance

Fluorescent proteins have been proven to be a powerful tool for plant cell biology studies,

permitting temporal and spatial monitoring of gene expression patterns at a cellular and

subcellular level (Berg et al., 2008). In order to expand the palette of fluorescent proteins that can

be used in *A. agrestis*, we tested the expression of the monomeric Turquoise 2 fluorescent

protein (mTurquoise2) (Kremers et al., 2006; Goedhart et al., 2012), the enhanced yellow

fluorescent (eYFP) protein (Orm et al., 1996), and the mVenus fluorescent protein (Kremers et

al., 2006). We used a construct similar to the one for the expression of eGFP protein, but with

different subcellular localization signals. mTurquoise2 and mVenus were fused to the nuclear-

localization peptide sequence of At4g19150/N7 (Cutler et al., 2000) with a linker to the amino (C)-

terminus (Cutler et al., 2000), and eYFP was fused to the membrane-targeting myristoylation

(myr) signal to the amino (N)-terminus (Resh, 1999). mTurquoise2 (Fig. 2F), eYFP (Fig. 2G) and

mVenus (Fig. 2H) were successfully expressed in *A. agrestis* and were targeted to the predicted

cellular compartments (for further information on the number of lines generated see Supplemental

Table I).

We then tested whether the protocol developed for the Oxford isolate can be used successfully

for the Bonn isolate. Four trials resulted in two successful transformants, which is considerably

less than the average number of transformants obtained for the Oxford isolate (Fig. 3J).

We finally tested whether the transgene can be stably inherited through the sexual life cycle. Two

eGFP expressing transgenic lines for both Oxford and Bonn isolates were brought to sexual

reproduction. Sporophytes were produced and young gametophytes germinating from the spores

(sporelings) were expressing eGFP indicating that the transgene and its expression was

successfully passed on to the next generation (Fig.3K-L).

Identification and selection of *A. agrestis* endogenous gene promoters

It is important to identify promoters that can be used to drive constitutive transgene expression

(i.e. high-level expression across almost all tissues and development stages). Commonly used

constitutive promoters in other bryophyte model species include the *M. polymorpha*

ELONGATION FACTOR 1 ALPHA (*EF1a*) promoter (Althoff et al., 2014), the rice *Actin1*, and the

M. polymorpha ubiquitin-conjugating enzyme E2 promoter (Sauret-Güeto et al., 2020). Using the

genomic sequence and RNA sequencing data (Li et al., 2020), (Fig. 4 A-B) we identified a series

of candidate promoter regions as constitutive *A. agrestis* promoters. In particular we selected the

promoter regions of the putative *A. agrestis* homologs of *EF1a*, *Ubiquitin*, *Actin*, and the

Arabidopsis *GAMMA TONOPLAST INTRINSIC PROTEIN* (*Tip1;1*) genes (Fig. 4 A-B,

Supplemental Table II). We amplified a 1532 bp long stretch of the 5' flanking region including the

5'UTR of the *EF1a*, a 933 bp segment for the *Ubiquitin (Ubi)*, two fragments (1729 bp and 1516

bp) for the *Actin* (that correspond to two different predicted translational start sites), and a 1368

bp putative promoter for the *Tip1;1* gene (Gene models, position of promoters and the

corresponding RNA-seq coverage tracks are shown in Supplemental Fig. S9).

The candidate promoter regions were cloned (and if necessary domesticated in order to generate

a Loop assembly cloning system compatible DNA part), fused with the eGFP or the mTurquoise2

reporter genes, and terminated with the double Nos - 35S terminator (Sauret-Güeto et al., 2020).

The *AaEF1a* (Fig. 5A) promoter region was sufficient to drive expression of eGFP throughout the

thallus. Similarly, the *AaTip1;1* promoter region was sufficient to drive expression of eGFP and

mTurquoise2 (Fig. 5B). However, only three independent lines were obtained for the *p-*

AaTip1;1::mTurquoise2-N7 construct (with one showing growth retardation probably due to the

insertion site of the T-DNA) and one for the *p-AaTip1;1::eGFP* construct (Supplemental Fig. S10).

Thus, further characterization of the *AaTip1;1* promoter is needed. The *AaUbi* (Fig. 5C) promoter

gave less uniform expression patterns throughout the thallus, and the two *AaActin* promoters

produced no detectable signal (Fig. 5D) (summary of the number of lines generated is shown in

Supplemental Table I). Our data thus indicate that out of the five candidates, *AaEF1a* is the best

promoter for driving relatively strong expression across cells of the gametophyte thallus. We

generated a total of eight p-*AaEF1a::eGFP* lines, four of which are shown in Supplemental Fig.

S10. Out of the eight hygromycin resistant lines, two do not express eGFP which could be due to

transgene silencing or truncation of the inserted T-DNA. In addition, we showed that the *AaEf1a*

promoter can drive adequate *hph* expression (Fig. 5E)*.* Finally, we were also able to successfully

express simultaneously two different transcription units, *p-AaEF1a::mTurquoise2-N7* and *p-*

35S_s::eGFP-LTI6b (Fig. 5F), which was the largest construct (approximately 7.4 kb) we

successfully introduced into the *A. agrestis* genome.

Comparison of the 35S and *AaEF1a* promoters

Expression of eGFP driven by the CaMV 35S promoter seems to be weaker in newly grown parts

of the thallus (Fig. 6A and Supplemental Fig. S8). This is similar to the expression patterns of

transgenes driven by the CaMV 35S promoter in *M. polymorpha*, which has a strong activity in all

parts of the thallus except the notch area (Althoff et al., 2014). Expression of eGFP driven by the

AaEF1a promoter seems to be stronger in the putatively younger parts of the thallus (Fig. 6B and

Supplemental Fig. S10). This is similar to the expression patterns of transgenes driven by the

EF1a promoter in *M. polymorpha*, showing a strong activity in all parts of the thallus particularly

the notch area (Althoff et al., 2014).

Transformation efficiency and optimization

In order to estimate the transformation efficiency of the protocol, we performed 10 independent

trials using approximately 2 g of tissue as starting material and the *p-35S::hph - p-35S_s::eGFP-*

LTI6b construct. The number of successful transformation events per experiments varied from 3

to 23 (Table I).

We then carried out further experiments to optimize transformation efficiency. We reasoned that

tissue susceptibility to *Agrobacterium* infection may differ during different stages of regeneration

after homogenization, thereby affecting transformation efficiency. To estimate when plant

regeneration is initiated, we set up a microscopy time course using homogenized thallus

fragments. The first cell division was observed five days after homogenization (Fig. 6C). Based

on this result, we carried out an optimization experiment starting co-cultivation at two, five and

seven days after homogenization. We found that the number of stably transformed lines

decreased when using tissue that was recovered for more than five days after homogenization

(Fig. 6D). The highest number of transformants could be obtained when using tissue two days

after homogenization. In parallel with the above mentioned experiment, we also tested another

Agrobacterium strain, the *GV3101*, for its ability to infect *A. agrestis* thallus. However, only a single

successful transformation event was obtained when the *GV3101* strain harboring the *p-35S::hph*

- *p-35S::eGFP-LTI6b* construct was used.

Verification of transgene incorporation into the *A. agrestis* genome

To confirm the genomic integration of the transgene, we sequenced and assembled the genomes

of five stable transformant lines (Fig. 7). For all five lines, we found a single integration site, with

one line showing a single full length insertion of the T-DNA. The other four lines additionally

showed one or multiple partial insertions in inverted and/or tandem directions (Supplemental

Material). To confirm the transgene integration site in the five lines, fragments overlapping the 5'-

and 3'- ends of the inserts and their adjacent genomic region were amplified by nested PCR and

Sanger-sequenced. The resulting sequences confirmed the integration sites identified by the

genome assemblies (Fig. 7G and Supplemental Material). We conclude that the transformation

method described here results in the stable integration of one or more targeted transcriptional

units into the *A. agrestis* nuclear genome.

Discussion

The protocol described in this study successfully generated stable transformants in *A. agrestis*

Oxford and Bonn isolates and may be applicable to other hornwort species. We generated a total

of 216 stable lines. We showed that transgenic lines can be propagated for more than two years

without abolishing transgene expression. Additionally, we verified that the transgene is integrated
into the genome of *A. agrestis* and can be successfully inherited.

The genome sequencing of transgenic lines showed that the integration occurs in a single locus
with one or more copies, which is similar to the reports available for other organisms based on
DNA gel blot analysis (Feldmann and David Marks, 1987; Ishizaki et al., 2008; Plackett et al.,
2014). The utilization of recent high throughput sequencing technologies combined with the
genome size of the *A. agrestis* allows precise determination of the transgene insertion site. Thus,
it should be simple to perform enhancer-trap or T-DNA based mutagenesis experiments in *A.*
agrestis.

The light conditions under which the tissue was grown significantly affected the thallus
morphology and were critical for successful transformation. It is likely that high light intensity
triggers the accumulation of secondary metabolites (such as mucilage) and/or affects the
composition of the cell wall, thereby significantly reducing transformation efficiency. Multiple
photoreceptors are present in the *A. agrestis* genome (Li et al., 2014; Li et al., 2015a; Li et al.,
2015b). Identifying which receptors determine the response to high light intensity could help to
further improve transformation efficiency. In addition, testing different methods for tissue
fragmentation, such as vortexing or use of a scalpel, or employing other types of tissues
(germinating spores or callus) might also improve transformation efficiency.

We are currently developing genome editing tools for *A. agrestis* using CRISPR/Cas9 (Jinek et
al., 2012). We are also investigating whether inducible gene expression systems such as the
glucocorticoid receptor (Schena et al., 1991) or estrogen receptor (Zuo et al., 2000) can be applied
successfully in hornworts. Finally, we are testing alternative gene delivery methods for both *A.*
agrestis Oxford and Bonn isolates, such as particle bombardment.

The development of a hornwort transformation method, in combination with the recently published
genome, will greatly facilitate more comprehensive studies of the mechanisms underpinning land
plant evolution. It can also help with engineering hornwort traits into plants with agronomic value.
For example engineering pyrenoids in crops, has the potential toimprove carbon fixation and
therefore increase crop yield (Li et al., 2017).

Materials and methods

Plant material and maintenance

The *Anthoceros agrestis* Oxford and Bonn isolates were used (Szövényi et al., 2015). *A. agrestis*

thallus tissue was propagated on KNOP medium (0.25 g/L KH_2PO_4, 0.25 g/L KCl, 0.25 g/L

$MgSO_4\cdot7H_2O$, 1 g/L $Ca(NO_3)_2\cdot4H_2O$ and 12.5 mg/L $FeSO_4\cdot7H_2O$). The medium was adjusted

to pH 5.8 with KOH and solidified using 7.5 g/L Gelzan CM (#G1910, SIGMA) in 92x16 mm petri

dishes (#82.1473.001, SARSTEDT) with 25-30 mL per plate. Plants were routinely grown in a

tissue culture room (21°C, 12 h of light and 12 h of dark, 5 µmol m^{-2} s^{-1} light intensity). In order

to subculture the thallus tissue, a small part of the thallus was cut using sterile disposable scalpels

(#0501, Swann Morton) and placed on fresh media on a monthly basis.

Co-cultivation medium

Co-cultivation medium was liquid KNOP supplemented with 2% sucrose (0.25 g/L KH_2PO_4, 0.25

g/L KCl, 0.25 g/L $MgSO_4\cdot7H_2O$, 1 g/L $Ca(NO_3)_2\cdot4H_2O$, 12.5 mg/L $FeSO_4\cdot7H_2O$ and 20 g/L

sucrose, pH 5.8 adjusted with KOH).

Tissue preparation for transformation

Approximately 2 g of thallus tissue were divided into 4 parts, and each part was homogenized in

15 mL of sterile water using a homogenizer (#727407, IKA Ultra-Turrax T25 S7 Homogenizer)

and corresponding dispensing tools (#10442743, IKA Dispersing Element), for 5 seconds, using

the lowest speed of 8000 rpm. The homogenized tissue was washed with 50 mL of sterile water

using a 100 µm cell strainer (#352360, CORNING), spread on solid KNOP medium and placed at

21°C under 12 hours light and 12 hours dark at a light intensity of 3-5 µmol m^{-2} s^{-1}. After 4 weeks

the tissue was re-homogenized in 15-20 mL of sterile water and filtered using a 100 µm cell

strainer. The re-homogenized tissue was transferred again onto 4 plates with solid KNOP medium

and was allowed to grow for 2 days at 21°C under continuous light (35 µmol m^{-2} s^{-1}, PHILIPS,

TL-D58W/835).

Agrobacterium **culture preparation**

A single *Agrobacterium* colony (AGL1 strain) was inoculated in 5 mL of LB medium supplemented
with rifampicin 10 μg/mL (#R0146, Duchefa), carbenicillin 50 μg/mL (#C0109, MELFORD) and
the plasmid-specific selection antibiotic spectinomycin 100 μg/mL (#SB0901, Bio Basic). The pre-
culture was incubated at 28°C for 2 days with shaking at 120 rpm. OD600 was ~2.7 and was
measured using an OD600 DiluPhotometer (IMPLEN).

Co-cultivation conditions

5 mL of 2 days *Agrobacterium* culture was centrifuged for 7 minutes at 2000 xg. The supernatant
was discarded and the pellet was resuspended in 5 mL liquid KNOP supplemented with 2% (w/v)
sucrose (#S/8600/60, Fisher) and 100 μM 3',5'-dimethoxy-4'-hydroxyacetophenone
(acetosyringone) (#115540050, Acros Organics, dissolved in dimethyl sulfoxide (DMSO)
(#D8418, SIGMA)). The culture was incubated with shaking (120 rpm) at 28°C for 5 hours. The
regenerating thallus tissue was transferred (1/2 tissue from one plate – 2 days after the second
homogenization) into a well of a 6-well plate with 4 mL of liquid KNOP medium supplemented with
2% (w/v) sucrose. 80 μL of *Agrobacterium* culture and acetosyringone at final concentration of
100 μM were added to the medium.

The tissue and *Agrobacterium* were co-cultivated using a 6-well plate (#140675, ThermoFisher)
for 3 days with shaking at 110 rpm at 22°C on a shaker without any additional light supplemented
(only ambient light from the room, 1-3 μmol m^{-2} s^{-1}). After 3 days, the tissue was drained using
a 100 μm cell strainer (#352360, CORNING) and moved onto solid KNOP medium plates
supplemented with 100 μg/mL cefotaxime (#BIC0111, Apollo Scientific) and 10 μg/mL
Hygromycin (#10687010, Invitrogen). After 3-4 weeks, plants were transferred to fresh solid
KNOP medium plates supplemented with 100 μg/mL cefotaxime and 10 μg/mL Hygromycin and
grown at 22°C 12 hours light and 12 hours dark at a light intensity of 35 μmol m^{-2} s^{-1} (PHILIPS,
TL-D58W/835).

GUS staining

GUS staining was performed according to Plackett et al., 2014**.**

Genomic DNA extraction

A modified CTAB protocol from (Porebski et al., 1997) was used for hornwort genomic DNA
extraction. 0.5 g of tissue was harvested and frozen in liquid nitrogen. Tissue was ground into a
fine powder using a chilled mortar and pestle and then added to 10 mL of DNA extraction buffer
(100 mM Tris-HCl pH 8, 1.4 M NaCl, 20 mM EDTA pH 8, 2% (w/v) CTAB, 0.3% (v/v) β-
mercaptoethanol and 100 mg of polyvinylpyrrolidone (PVP)/g of tissue) that had been prewarmed
at 60°C, 100 μL of RNAse A (100 mg/mL) was added and the solution was mixed well. The mix
was incubated at 60°C for 20-30 minutes and then removed from heat and allowed to cool at room
temperature for 4 minutes. 12 mL of chloroform:isoamyl alcohol (24:1) was added, mixed well by
inversion and then centrifuged at 12000 xg for 10 minutes at room temperature. The upper
aqueous phase was transferred to a new 50 mL centrifugation tube and 10 mL of
chloroform:isoamyl alcohol (24:1) was added, mixed well by inversion and then centrifuged at
6000 xg for 10 minutes at room temperature to remove any remaining PVP in the aqueous phase.
The upper aqueous phase was transferred to a new 50 mL centrifugation tube and ½ volume of
5 M NaCl was added. 2 volumes of cold (-20°C) 95% (v/v) ethanol were also added and the
contents of the tube were mixed well by inversion. The tube was spun at 20000 xg for 6 minutes.
The pellet was resuspended in 2 mL of TE buffer and the previous step was repeated. The pellet
was washed with cold 70% (v/v) ethanol. The pellet was dried and dissolved in 80 μL of TE buffer
and then stored at 4°C.

Construct generation

Constructs were generated using the OpenPlant toolkit (Sauret-Gueto et al., 2020). Full sequence
of constructs can be found in Supplemental Table III.

Promoter identification and isolation

We used RNA-seq data to find genes showing constantly high levels of expression under various
developmental stages and experimental conditions ("constitutively expressed genes"). To do so, we
estimated expression of genes under three developmental stages of the gametophyte and sporophyte
phases and in symbiosis with cyanobacteria. We retrieved raw RNA-seq data for these experiments
from (Li et al., 2020). We used trimmomatic to quality filter and trim the raw reads. Gene expression
was estimated using Salmon (Patro et al., 2017) and expressed as normalized expression counts. We
identified candidates by selecting those showing the highest average expression level and the least
gene expression variability across all conditions investigated. We then manually selected a subset of
genes taking into account their genomic location, exact expression pattern, and the length and
sequence composition of their putative promoter sequences. We also assessed the suitability of our

candidate promoters using the information available for *M. polymorpha* and *P. patens* (Supplemental
Table II).

Putative promoter sequences were amplified from genomic DNA using the KOD Hot start
polymerase (#71086-5, Merck Millipore) and cloned into pJET1.2 (#K1231, ThermoFisher) before
Sanger sequencing. Loop assembly compatible DNA parts were generated according to (Sauret-
Güeto et al., 2020). List of primers can be found in Supplemental Table IV.

Sample preparation for Imaging

A gene frame (#AB0576, ThermoFisher) was positioned on a glass slide and 30 µL of KNOP medium
with 1% (w/v) Gelzan CM (#G1910, SIGMA) was placed within the gene frame. A thallus fragment
was placed within the medium-filled gene frame together with 30 µL of milliQ water. The frame was
then sealed with a cover slip. Plants were imaged immediately using a Leica SP8X spectral
fluorescent confocal microscope.

For the regeneration test experiment, five thallus fragments were placed into a KNOP medium-
filled gene frame as described above (three slides and 15 plants in total). Images were acquired on
a daily basis, for a total duration of a week, using a Leica SP8X spectral fluorescent confocal
microscope and a 10× air objective (HC PL APO 10×/0.40 CS2).

Imaging with Confocal Microscopy.

Images were acquired on a Leica SP8X spectral confocal microscope. Imaging was conducted
using either a 10× air objective (HC PL APO 10×/0.40 CS2) or a 20× air objective (HC PL APO
20×/0.75 CS2). Excitation laser wavelength and captured emitted fluorescence wavelength
window were as follows: for mTurquoise2 (442 nm, 460−485 nm), for eGFP (488 nm, 498−516
nm), for mVenus and eYFP (515 nm, 522−540 nm), and for chlorophyll autofluorescence (488 or
515, 670−700 nm). When observing lines expressing both eGFP and mTurquoise2, sequential
scanning mode was used.

Light microscopy

Images were captured using a KEYENCE VHX-S550E microscope (VHX-J20T lens) or a Leica
M205 FA Stereomicroscope (with GFP longpass (LP) filter).

Sequencing transformant line genomes

Transformant lines were grown on solid KNOP medium containing 10 µg/mL hygromycin and 100
µg/mL cefotaxime. Genomic DNA was extracted from 600 mg fresh tissue per line using either
the DNeasy Plant Pro kit (#69204, Qiagen) (Cam-1 and Cam-2 lines) or the procedure from Li et
al (2020) (BTI1-3 lines) to reach a total yield of at least 200 ng/line. Illumina libraries were
prepared using the TruSeq DNA nano kit (#20015964, Illumina) and were sequenced on an
Illumina Novaseq 6000 platform with an expected sequencing depth of 80-150x for all five lines
in paired-end mode (read length: 151 bp).

After sequencing we quality filtered and trimmed reads using trimmomatic (command line:
ALL_TruSeq-PE.fa:2:30:10:2:keepBothReads LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20
MINLEN:36) and assembled the reads with spades3.14.1 using the --isolate --cov-cutoff auto --
only-assembler options as recommended (Nurk et al., 2013). To localize the insertion and its copy
number, we used the insert sequence as a query in a BLASTN search (Altschul et al., 1990)
against the database containing the assembly (Altschul et al., 1990) with an e-value threshold of
10^{-4}. As evidence of genomic integration, we only accepted hits covering the full-length of the
insert sequence with one or no mismatches. We manually inspected blast hits to eliminate false
positives. Finally, we used the *A. agrestis* Oxford genome sequence (Li et al., 2020) to localize
the hits on the pseudomolecules.

PCR confirmation of transformant insertion sites

Based on the genome assemblies of the transformant lines, primers were designed to amplify
regions of 0.6-1.3 kb spanning the 5'- and 3'- end of the T-DNA inserts and their respective
adjacent genomic regions. Sequences were amplified from genomic DNA with Phusion High-
Fidelity DNA Polymerase (#F-530S, ThermoFisher). For each amplified region, a second set of
nested primers were designed to amplify a shorter amplicon, using 1:100 dilution of the previous
PCR product as a template. Resulting nested PCR-products were either purified and Sanger
sequenced (Eurofins) from both ends, or cloned into pJET1.2 (#K1231, ThermoFisher) before
Sanger sequencing from both ends using pJET1.2 sequencing primers.

Acknowledgments

We would like to thank Juan Carlos Villarreal for introducing us to the world of hornworts and always generously sharing his expertise. We are also thankful to Jim Haseloff for support during the latest stage of the project. Figure 1D picture credit: John Baker, Oxford University. This research was supported by the Japanese Society for the Promotion of Science (JSPS) Short Term Postdoctoral Fellowship grant no. PE14780 to E.F., the UZH Forschungskredit Candoc grant no. FK-19-089 to M.W., the MEXT and JSPS Grants-in-Aid for Scientific Research on Innovation Areas nos. 25113001 and 19H05672 to H.S., JSPS KAKENHI 26650143 and 18K06367 to K.S., 15H04413 to T.N., 19K22448 to T.N. and K.S., the Swiss National Science Foundation (grants 160004, 131726 and 184826) to P.S., The Deutsche Forschungsgemeinschaft (DFG - German Research Foundation) under the Priority Programme "MAdLand - Molecular Adaptation to Land: Plant Evolution to Change" (SPP 2237, 440370263) to P.S., The Georges and Antoine Claraz Foundation to P.S., M.W and Y.Y, The University Research Priority Program 'Evolution in Action' of the University of Zurich to P.S., and the National Science Foundation IOS-1923011 to F.-W.L. and J.V.E.

Accession numbers

Raw sequencing reads have been submitted to the NCBI SRA under the BioProject ID PRJNA683066 (SRR13209765-SRR13209769).

FIGURES

Figure 1: Morphological features of *Anthoceros agrestis* and effect of light on growth

A) Light micrograph (LM) of a germinating spore. Scale bar: 50 µm. B) Surface view of the irregularly shaped thallus (gametophyte). Scale bar: 1 mm. C) LM showing cells of mature gametophyte tissue with single chloroplasts. Scale bar: 10 µm. D) *A. agrestis* Oxford gametophyte with sporophytes. Scale bar: 4 mm.

E) The conditions for the preparation of tissue used for transformation are critical. Plants must be propagated in axenic culture by transferring small thallus fragments (typically 1 x 1 mm) onto plates with fresh growth medium using sterile scalpels and then grown under low light conditions (3-5 µmol m^{-2} s^{-1}) for 4 weeks.

F) *A. agrestis* Oxford thallus tissue grown for 4 weeks under low light intensity (3-5 μmol m^{-2} s^{-1}).

G) *A. agrestis* Oxford thallus tissue grown for 4 weeks under high light intensity (80 μmol m^{-2} s^{-1}).

Scale bars: 1 mm.

Figure 2: Outline of *Anthoceros agrestis* transformation method

1a-c: Tissue is homogenized, transferred on growth medium, and placed under low light conditions. **2a-c**: After 4 weeks, the tissue is homogenized again and grown for two additional days. **3**: The tissue is co-cultivated with *Agrobacterium* for three days and then **4a-b**: spread on appropriate antibiotic-containing growth medium. **5**: After 3 weeks, the tissue is transferred again onto freshly prepared antibiotic-containing growth medium for a second round of selection. **6**: After approximately 4-8 weeks, putative transformants are visible.

Figure 3: Schematic representation of transformation constructs and transgenic
***Anthoceros agrestis* expressing different fluorescent proteins**

A-B) The emergence of rhizoids (shown with yellow arrows) is a reliable indicator of successfully
transformed plant fragments. In (A) the red arrowheads show false positives regenerating (green)
tissue fragments that lack rhizoids. Scale bars: 2 mm.

C) GUS activity detected as blue staining in thallus tissue fragments from a plant transformed with
the pCAMBIA1305.2 plasmid. Scale bar: 200 µm.

D) Light micrograph of thallus surface view of thallus (gametophyte), similar to the area imaged
in E-H. Scale bar: 100 µm.

E-H top: Schematic representation of constructs for the expression of two transcription units (TU):
one TU for the expression of the *hygromycin B phosphotransferase* (*hph*) gene under the control
of the cauliflower mosaic virus (CaMV) 35S promoter and one TU for the expression of *p-*
35S_s::eGFP-LTI6b (E), *p-35S_s::mTurquoise2-N7* (F), *p-35S_s::eYFP-myr* (G) and *p-*
35S:mVenus-N7 (H) TU. *hph*: *hygromycin B phosphotransferase*; 35S: CaMV 35S promoter;
eGFP: enhanced green fluorescent protein; mTurquoise2: monomeric turquoise 2 fluorescent
protein; eYFP: enhanced yellow fluorescent protein; LTI6b: Low Temperature Induced Protein
6B signal for membrane localization; N7: Arabidopsis At4g19150/N7 nuclear localization signal;
myr: myristoylation signal for membrane localization; nosT: 3′ signal of *nopaline synthase*.

E-H middle and bottom: Images of *A. agrestis* Oxford thallus tissue expressing different
combinations of CaMV 35S promoter - fluorescent protein - localization signal. E) *p-35S_s::eGFP-*
LTI6b for plasma membrane localization (white arrow). Scale bars: top: 50 µm, bottom: 20 µm, F)
p-35S_s::mTurquoise2-N7 for nuclear localization (white arrow). Scale bars: top: 20 µm, bottom:
10 µm, G) *p-35S_s::eYFP-myr* for plasma membrane localization (white arrow). Scale bars: top:
50 µm, bottom: 20 µm, H) and *p-35S::mVenus-N7* for nuclear localization (white arrow). Scale
bars: top: 50 µm, bottom: 25 µm. The bottom image is a magnification of the image in the middle.
Red, chlorophyll autofluorescence.

I) Example of transgenic *A. agrestis* plants (gametophyte thallus). Petri dish dimensions: 92 x16
mm.

J) Images of *A. agrestis* Bonn gametophyte tissue expressing the *p-35S_s::eGFP-LTI6b* TU for
eGFP plasma membrane localization. Scale bar: 50 μm. K) *A. agrestis* Bonn with mature
sporophytes indicated with white arrows. Scale bars: 2 mm L) *A. agrestis* Bonn transgenic spores
expressing the *p-35S_s::eGFP-LTI6b* TU (white arrow indicates rhizoid and yellow arrow
indicates young thallus). Top left: Light microscopy of *A. agrestis* Bonn wild type germinating
spore. Scale bar: 20 μm.

Figure 4. Identification of constitutive promoters for *Anthoceros agrestis*

A) Analysis of expression levels from RNA-seq experiments on *A. agrestis* Oxford using datasets from Li et al. 2020. To generate this dataset, gametophytes were grown under varying carbon sources in the growth medium (indicated as C depleted, C enriched and C normal), as well as two different nitrogen depleted conditions (N- with cyanobacteria symbiosis and N- without cyanobacteria symbiosis). Error bars indicate standard error based on three independent experimental replicates. (Included for comparison: "CN-Max 1, 3 and 5": highest expressing genes based on N conditions and "CN-Max 2 and 4": highest expressing genes based on C normal conditions (for gene ID, see Supplemental Table II)).

B) Analysis of expression levels from RNA-seq experiments on *A. agrestis* Bonn using data sets from Li et al. 2020. Note: Normalized expression level of *A. agrestis* BONN genes selected to represent strong and constitutive expression across various developmental stages of the gametophyte and the sporophyte phases (Included for comparison: "T-Max 1 to 5": Highest expressing genes under all conditions (for gene ID, see Supplemental Table II)). Error bars indicate standard error based on two independent experimental replicates.

Figure 5. Identification of constitutive promoters for *Anthoceros agrestis*

A-F top: Schematic representation of constructs for the expression of two/three transcription units
(TU): one TU for the expression of the *hph* gene under the control of the CaMV 35S or the *p-*
AaEF1a promoter and one TU for the expression of plasma membrane localized eGFP and/or
nucleus localized mTurquoise2 under the control of different native *A. agrestis* promoters. A) *p-*
AaEF1a, B) *p-AaTip1;1*, C) *p-AaUbi* and D) *p-AaActin_1.* E) *p-AaEF1a* driving *hph* and F) *p-*
AaEF1a driving *mTurquoise2-N7*.

A-F middle and bottom: Images of *A. agrestis* Oxford gametophyte tissue expressing different
combinations of *A. agrestis* native promoter - fluorescent protein - localization signal. A) *p-*
AaEF1a:eGFP-LTI6b for plasma membrane localization. Scale bars: top: 50 µm, bottom: 10 µm,
B) *p-AaTip1;1::mTurquoise2-N7* for nuclear localization. Scale bars: top: 20 µm, bottom: 10 µm,
C), *p-AaUbi::eGFP-LTI6b* for plasma membrane localization. Scale bars: top: 20 µm, bottom: 10
µm. The bottom images are a magnification of the images in the top. D) *p-AaActin::eGFP-LTI6b*.
Scale bars: 50 µm and E) *p-AaEF1a::hph - p-35S_s::eGFP-LTI6b.* Scale bar: 50 µm. F) Image of
A. agrestis Oxford gametophyte tissue expressing both *p-AaEF1a::mTurquoise2-N7* for nuclear
localization and *p-35S_s::eGFP-LTI6b* for plasma membrane localization. Scale bar: 50 µm. Red,
chlorophyll autofluorescence.

Figure 6: Comparison of the 35S with the *AaEF1a* promoter and factors affecting the

efficiency of *Agrobacterium*-mediated transformation of *Anthoceros agrestis*

A) Expression of eGFP driven by the CaMV 35S promoter. Younger part of the thallus indicated

with a white arrow. Scale bar: 50 µm.

B) Expression of eGFP driven by the *AaEF1a* promoter. Younger part of the thallus indicated

with white arrows. Scale bar: 100 µm.

C) Confocal microscopy images of fragmented *A. agrestis* thallus tissue taken on seven

consecutive days after homogenization. Five days after homogenization plants start to

regenerate. Yellow arrowhead indicates the fragmented thallus part. White arrows indicate new

cell divisions. Scale bars: 50 µm.

D) Number of transgenic lines obtained using *A. agrestis* thallus tissue two, five and seven days

after homogenization. Values of independent experimental replicates are shown.

Figure 7: Stable incorporation of transgene into *Anthoceros agrestis* genome

A-F Left: Light micrograph (LM) of transgenic thallus of *A. agrestis* plants. Scale bar: 500 µm.
Right: Confocal fluorescent microscopy images of thallus expressing eGFP in the plasma
membrane, driven by the CaMV 35S promoter. Scale bars: A-E 100 µm and F 50 µm. Bottom:
Location of transgene insertion in the genome (see details in Supplemental material). Black
arrows indicate directionality of T-DNA insert. G) PCR analysis of genomic DNA from transgenic
plants. L: fragment amplified from sequences spanning the 5'- end of the T-DNA inserts and their
respective adjacent genomic regions. R: fragment amplified from sequences spanning the 3'- end
of the T-DNA inserts and their respective adjacent genomic regions.

Note: A, E and F LM images were acquired using a KEYENCE VHX-S550E microscope (VHX-
J20T lens) and confocal fluorescent images with a Leica SP8X microscope, in Cambridge
University.   B-D LM and fluorescent images were acquired using a Leica M205 FA
Stereomicroscope with GFP long-pass (LP) filter, in BT Institute.

Experiment	Amount of tissue (g)	Number of transformants
1	2.7	21
2	3.1	12
3	2.9	23
4	2.5	5
5	1.7	3
6	2.7	11
7	2.6	19
8	1.9	4
9	2.23	12
10	2.1	6

692 **Table I: Number of *p-35S::eGFP-Lti6B* transgenic lines obtained from different**
693 **transformation experiments**.

References

Althoff F, Kopischke S, Zobell O, Ide K, Ishizaki K, Kohchi T, Zachgo S (2014) Comparison of the MpEF1α and CaMV35 promoters for application in Marchantia polymorpha overexpression studies. Transgenic Res **23**: 235–244

Althoff F, Zachgo S (2020) Transformation of Riccia fluitans, an Amphibious Liverwort Dynamically Responding to Environmental Changes. International Journal of Molecular Sciences **21**: 5410

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. J Mol Biol **215**: 403–410

Berg RH, Howard Berg R, Beachy RN (2008) Fluorescent Protein Applications in Plants. Fluorescent Proteins 153–177

Chiyoda S, Ishizaki K, Kataoka H, Yamato KT, Kohchi T (2008) Direct transformation of the liverwort Marchantia polymorpha L. by particle bombardment using immature thalli developing from spores. Plant Cell Rep **27**: 1467–1473

Cho SH, Chung YS, Cho SK, Rim YW, Shin JS (1999) Particle bombardment mediated transformation and GFP expression in the moss Physcomitrella patens. Mol Cells **9**: 14–19

Cormack BP, Valdivia RH, Falkow S (1996) FACS-optimized mutants of the green fluorescent protein (GFP). Gene **173**: 33–38

Cutler SR, Ehrhardt DW, Griffitts JS, Somerville CR (2000) Random GFP::cDNA fusions enable visualization of subcellular structures in cells of Arabidopsis at a high frequency. Proceedings of the National Academy of Sciences **97**: 3718–3723

Desirò A, Duckett JG, Pressel S, Villarreal JC, Bidartondo MI (2013) Fungal symbioses in hornworts: a chequered history. Proc Biol Sci **280**: 20130207

Feldmann KA, David Marks M (1987) Agrobacterium-mediated transformation of germinating seeds of Arabidopsis thaliana: A non-tissue culture approach. Molecular and General Genetics MGG **208**: 1–9

Frangedakis E, Shimamura M, Villarreal JC, Li F-W, Tomaselli M, Waller M, Sakakibara K,
Renzaglia KS, Szövényi P (2020) The Hornworts: Morphology, evolution and
development. New Phytol. doi: 10.1111/nph.16874

Gelvin SB (2003) Agrobacterium-Mediated Plant Transformation: the Biology behind the "Gene-
Jockeying" Tool. Microbiology and Molecular Biology Reviews **67**: 16–37

Goedhart J, von Stetten D, Noirclerc-Savoye M, Lelimousin M, Joosen L, Hink MA, van
Weeren L, Gadella TWJ, Royant A (2012) Structure-guided evolution of cyan fluorescent
proteins towards a quantum yield of 93%. Nature Communications. doi:
10.1038/ncomms1738

Ishizaki K, Chiyoda S, Yamato KT, Kohchi T (2008) Agrobacterium-mediated transformation
of the haploid liverwort Marchantia polymorpha L., an emerging model for plant biology.
Plant Cell Physiol **49**: 1084–1091

Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E (2012) A
Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity.
Science **337**: 816–821

Kremers G-J, Goedhart J, van Munster EB, Gadella TWJ (2006) Cyan and Yellow Super
Fluorescent Proteins with Improved Brightness, Protein Folding, and FRET Förster
Radius†,‡. Biochemistry **45**: 6570–6580

Kubota A, Ishizaki K, Hosaka M, Kohchi T (2013) Efficient Agrobacterium-mediated
transformation of the liverwort Marchantia polymorpha using regenerating thalli. Biosci
Biotechnol Biochem **77**: 167–172

Kugita M (2003) RNA editing in hornwort chloroplasts makes more than half the genes
functional. Nucleic Acids Research **31**: 2417–2423

Li F-W, Melkonian M, Rothfels CJ, Villarreal JC, Stevenson DW, Graham SW, Wong GK-S,
Pryer KM, Mathews S (2015a) Phytochrome diversity in green plants and the origin of
canonical plant phytochromes. Nature Communications. doi: 10.1038/ncomms8852

Li F-W, Nishiyama T, Waller M, Frangedakis E, Keller J, Li Z, Fernandez-Pozo N, Barker
MS, Bennett T, Blázquez MA, et al (2020) Anthoceros genomes illuminate the origin of
land plants and the unique biology of hornworts. Nat Plants **6**: 259–272

Li F-W, Rothfels CJ, Melkonian M, Villarreal JC, Stevenson DW, Graham SW, -S. Wong GK, Mathews S, Pryer KM (2015b) The origin and evolution of phototropins. Frontiers in Plant Science. doi: 10.3389/fpls.2015.00637

Li F-W, Villarreal Aguilar JC, Szövényi P (2017) Hornworts: An overlooked window into carbon-concentrating mechanisms. Trends Plant Sci **22**: 275–277

Li F-W, Villarreal JC, Kelly S, Rothfels CJ, Melkonian M, Frangedakis E, Ruhsam M, Sigel EM, Der JP, Pittermann J, et al (2014) Horizontal transfer of an adaptive chimeric photoreceptor from bryophytes to ferns. Proceedings of the National Academy of Sciences **111**: 6672–6677

Liu J-W, Li S-F, Wu C-T, Valdespino IA, Ho J-F, Wu Y-H, Chang H-M, Guu T-Y, Kao M-F, Chesson C, et al (2020) Gigantic chloroplasts, including bizonoplasts, are common in shade-adapted species of the ancient vascular plant family Selaginellaceae. Am J Bot **107**: 562–576

Morris JL, Puttick MN, Clark JW, Edwards D, Kenrick P, Pressel S, Wellman CH, Yang Z, Schneider H, Donoghue PCJ (2018) The timescale of early land plant evolution. Proc Natl Acad Sci U S A **115**: E2274–E2283

Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Prjibelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, et al (2013) Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. J Comput Biol **20**: 714–737

One Thousand Plant Transcriptomes Initiative (2019) One thousand plant transcriptomes and the phylogenomics of green plants. Nature **574**: 679–685

Orm M, Cubitt AB, Kallio K, Gross LA, Tsien RY, Remington SJ (1996) Crystal Structure of the Aequorea victoria Green Fluorescent Protein. Science **273**: 1392–1395

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C (2017) Salmon provides fast and bias-aware quantification of transcript expression. Nat Methods **14**: 417–419

Plackett ARG, Huang L, Sanders HL, Langdale JA (2014) High-efficiency stable transformation of the model fern species Ceratopteris richardii via microparticle bombardment. Plant Physiol **165**: 3–14

Pollak B, Cerda A, Delmans M, Álamos S, Moyano T, West A, Gutiérrez RA, Patron NJ,
Federici F, Haseloff J (2019) Loop assembly: a simple and open system for recursive
fabrication of DNA circuits. New Phytol **222**: 628–640

Porebski S, Grant Bailey L, Baum BR (1997) Modification of a CTAB DNA extraction protocol
for plants containing high polysaccharide and polyphenol components. Plant Molecular
Biology Reporter **15**: 8–15

Renzaglia KS, Villarreal Aguilar JC, Piatkowski BT, Lucas JR, Merced A (2017) Hornwort
stomata: architecture and fate shared with 400-Million-year-old fossil plants without leaves.
Plant Physiol **174**: 788–797

Resh MD (1999) Fatty acylation of proteins: new insights into membrane targeting of
myristoylated and palmitoylated proteins. Biochim Biophys Acta **1451**: 1–16

Sauret-Güeto S, Frangedakis E, Silvestri L, Rebmann M, Tomaselli M, Markel K, Delmans
M, West A, Patron NJ, Haseloff J (2020) Systematic Tools for Reprogramming Plant
Gene Expression in a Simple Model,. ACS Synth Biol **9**: 864–882

Schaefer D, Zryd J-P, Knight CD, Cove DJ (1991) Stable transformation of the moss
Physcomitrella patens. Molecular and General Genetics MGG **226**: 418–424

Schena M, Lloyd AM, Davis RW (1991) A steroid-inducible gene expression system for plant
cells. Proceedings of the National Academy of Sciences **88**: 10421–10425

Small ID, Schallenberg-Rüdinger M, Takenaka M, Mireau H, Ostersetzer-Biran O (2019)
Plant organellar RNA editing: what 30 years of research has revealed. Plant J. **101**: 1040–1056
808
Söderström L, Hagborg A, von Konrat M, Bartholomew-Began S, Bell D, Briscoe L, Brown
E, Cargill DC, Costa DP, Crandall-Stotler BJ, et al (2016) World checklist of hornworts
and liverworts. PhytoKeys 1–828

Stachel SE, Messens E, Van Montagu M, Zambryski P (1985) Identification of the signal
molecules produced by wounded plant cells that activate T-DNA transfer in Agrobacterium
tumefaciens. Nature **318**: 624–629

Szövényi P, Frangedakis E, Ricca M, Quandt D, Wicke S, Langdale JA (2015)
Establishment of Anthoceros agrestis as a model species for studying the biology of

hornworts. BMC Plant Biol **15**: 98

Villarreal JC, Renner SS (2012) Hornwort pyrenoids, carbon-concentrating structures, evolved
and were lost at least five times during the last 100 million years. Proc Natl Acad Sci U S A
109: 18873–18878

Yoshinaga K (1996) Extensive RNA editing of U to C in addition to C to U substitution in the
rbcL transcripts of hornwort chloroplasts and the origin of RNA editing in green plants.
Nucleic Acids Research **24**: 1008–1014

Yoshinaga K (1997) Extensive RNA editing and possible double-stranded structures
determining editing sites in the atpB transcripts of hornwort chloroplasts. Nucleic Acids
Research **25**: 4830–4834

Young JM, Kuykendall LD, Martínez-Romero E, Kerr A, Sawada H (2001) A revision of
Rhizobium Frank 1889, with an emended description of the genus, and the inclusion of all
species of Agrobacterium Conn 1942 and Allorhizobium undicola de Lajudie et al. 1998 as
new combinations: Rhizobium radiobacter, R. rhizogenes, R. rubi, R. undicola and R. vitis.
Int J Syst Evol Microbiol **51**: 89–103

Zuo J, Niu QW, Chua NH (2000) Technical advance: An estrogen receptor-based
transactivator XVE mediates highly inducible gene expression in transgenic plants. Plant J
24: 265–273