Competitive outcomes between native and invasive plants are linked to shifts in the bacterial

rhizosphere microbiome

Marina L. LaForgia[1], Hannah Kang[2], Cassandra L. Ettinger[3,4*]

[1]Department of Environmental Science and Policy, University of California, Davis, CA, USA

[2]Department of Plant Sciences, University of California, Davis, CA, USA

[3]Genome Center, University of California, Davis, CA, USA

[4]Department of Evolution and Ecology, University of California, Davis, CA, USA

* Corresponding author; email: clettinger@ucdavis.edu

Abstract

Background

Rhizosphere microbiomes have received growing attention in recent years for their role in plant health, stress tolerance, soil nutrition, and invasive species dominance. Still, relatively little is known about how these microbial communities are altered under plant competition, and even less about whether these shifts are tied to competitive outcomes between native and invasive plant species. We investigated the structure and diversity of rhizosphere bacterial and fungal microbiomes of native annual forbs and invasive annual grasses individually and in competition using high-throughput amplicon sequencing of the bacterial 16S rRNA gene and the fungal ITS region. We assessed how significant shifts in key microbial families correlate to plant competitive responses through changes in biomass (log competitive response ratios).

Results

We find that bacterial diversity and structure differ between invasive grasses and native forbs, but fungal diversity and structure do not. Further, bacterial community structures under competition are distinct from both individual forb and grass bacterial community structures. We identified four bacterial families (Burkholderiaceae, Methylophilaceae, Clostridiaceae_1, and Fibrobacteraceae) that varied in relative abundance between treatments and that were significantly correlated with plant competitive responses.

Conclusions

Invasive grass dominance may be partially due to effects on the rhizosphere community of native forbs, with changes in specific bacterial families potentially benefiting grasses at the expense of native forbs. Our study underscores the importance of considering plant-rhizosphere interactions for understanding outcomes of plant invasion on grassland ecosystems.

Keywords: rhizosphere, microbiome, bacteria, fungi, competition, invasive species, plant-

microbe interactions, annual grassland

Background

Plants and their associated soil microbial communities have coevolved complex dynamic

relationships through time. The soil directly surrounding plant roots, known as the rhizosphere, is

home to an abundance and diversity of microbes distinct from the surrounding bulk soil

microbial community [1–4]. Microbes in the rhizosphere play vital roles in plant health [1],

nutrition [5], and stress tolerance [6,7]. Though rhizosphere microbial composition is driven

predominantly by soil particle size, pH, and moisture content [8–10], these communities can also

be highly host plant-specific [11,12], with plants exhibiting strong local adaptations to their

home microbial community [13–15]. Further, there is growing evidence that microbial

community composition varies in the presence of certain neighbors [16], and can even affect

competitive outcomes between plants [17–19] with potentially broad consequences for plant

community assembly, nutrient cycling, and ecosystem resilience [20–23].

Microbes can affect plant-host competition both directly via resource availability and

indirectly through community interactions. For example, microbes can make nitrogen more

readily available to plants through the processes of decomposition, nitrification, and nitrogen

fixation [23,24], which could preferentially benefit hosts capable of quickly taking up resources

[25,26] thereby increasing these plants' competitive success relative to species with slower

uptake rates [27]. Plants may also indirectly compete for beneficial plant-growth-promoting

microbes, with these microbes boosting root growth and soil exploration of some species more

than others [4,28,29]. Conversely, competition with a novel plant, such as an invader, may bring

resident plants into contact with novel pathogens [30]. These competitive interactions can lead to

the loss of specialized microbes in the inferior competitor [31] and drive the resulting microbial

community to resemble that of the dominant competitor [18,19]. Furthermore, microbial

community changes can feed back to affect plant competitive ability [19], with the potential to

drive either coexistence or exclusion [32].

Shifts in the microbiome are a potentially important contributor to the dominance of

invasive plants and their threats to native species. Although there is strong evidence for local

adaptation between plants and their home microbiome, invasive plants are relatively novel

players, causing disruptions in important plant-microbe interactions [33] by increasing soil

microbial activity [34], reducing microbial biomass [35], decreasing microbial diversity [36],

increasing nitrification rates [25], and changing microbiome composition [37,38]. These plant-

soil feedbacks have the potential to further benefit the invader [38,39], selectively harm natives

[40], and/or benefit other invaders [41]. While it is known that invaders and natives may harbor

distinct microbial communities, and that competition between host plants can affect microbiome

composition (and vice versa), little is known of how changes in the abundance of specific

microbial taxa may be tied to the competitive dominance of invaders and/or the competitive

inferiority of natives.

Serpentine soils in California annual grassland communities are characterized by high

levels of heavy metals and magnesium and low levels of essential plant nutrients, and as a result,

are home to a unique community of native annual forbs adapted to growing in these conditions.

While some of these soils are rocky and shallow, other serpentine areas are deeper and more

finely textured and therefore hold more water and nitrogen [42], making them more favorable to

invasive annual grass establishment and growth. Invasive annual grasses are highly competitive,

spread rapidly, and produce an enormous amount of seed each year. Their invasion into finer-

textured serpentine areas is promoted through nutrient addition and increased rainfall [42], which

leads to a fungal community more like that of heavily-invaded nonserpentine soils [43]. In

native-dominated serpentine areas, microbes have been shown to increase seedling survival [44]

and facilitate their ability to thrive in heavy metal soils [45]. With the invasion of grasses,

changes in the microbiome may strengthen the advantage that these grasses have over native

forbs if they are able to recruit beneficial microbes away from natives or harbor novel microbes

harmful to native forbs. Alternatively, locally adapted microbes may be harming invasive

grasses, helping native forbs to persist in these areas.

We sought to understand how competition between native forbs and invasive grasses

shifts the bacterial and fungal rhizosphere microbiomes, and how these shifts may be shaping

competitive outcomes in this community. Our main questions were:

(Q1) Do microbiomes differ between invasive grasses and native forbs?

(Q2) Do microbiomes under grass-forb competition differ from the microbiomes of (a)

invasive grasses and (b) native forbs?

(Q3) If so, are microbiomes under grass-forb competition sourced more from invasive

grasses, native forbs, or equally from both?

(Q4) Given compositional changes in the microbiome under grass-forb competition,

which specific microbial families are driving these changes?

(Q5) Are changes in the relative abundance of these microbial families correlated with

the competitive responses of forbs and grasses to each other?

To answer these questions, we conducted a manipulative shade-house experiment in

which forbs and grasses were grown in field-collected soil both individually and in grass-forb

pairs. Individual rhizosphere microbiomes were sampled for each species grown alone, joint

rhizosphere microbiomes were sampled for each species pair, and high throughput amplicon

sequencing was used to identify both bacterial and fungal rhizosphere communities. To assess

how microbiomes differed between invasive grasses and native forbs (Q1), and between each of

these groups individually and in competition (Q2), we looked at differences in community

structure and composition as well as alpha diversity. We evaluated the contribution of individual

grass and forb microbiomes to the joint microbiomes of grass-forb pairs (Q3) by calculating the

proportion of the joint microbiome predicted to be sourced from either the forb or the grass

microbiome. We then used differential abundance analysis to identify which microbial families

were driving any observed changes (Q4). Finally, to understand whether the relative abundances

of these microbial families may be tied to the competitive responses of grasses and forbs (Q5),

we evaluated the relationship of biomass-based log competitive response ratios of forbs and

grasses to each microbial family's relative abundance. To corroborate whether these

relationships may be driven by either the forb or the grass, we also calculated the proportion of

each family in the joint microbiome predicted to originate from either the forb or the grass

microbiome. For each step, we looked at effects in both the bacterial and the fungal rhizosphere.

We focus our analysis on group-level results (i.e. native forb, invasive grass, and grass-forb

pairs) after ensuring differences in microbiomes were attributable to groups rather than

individual plant species. We also accounted for species-level differences with random effects in

mixed models.

Methods

Experimental set-up

In spring and summer 2017, we collected soil and seed from an annual grassland at the

University of California McLaughlin Natural Reserve [2] in the Inner North Coast Range (N

38°52', W 122°26'). The site has a Mediterranean climate with cool, wet winters (mean annual

precipitation 2000-2019 of 738 mm [3]) and dry, hot summers. Soil was collected from an area

of the reserve with deep, finely-textured serpentine soils and seeds were collected nearby from

six common native annual forb species (Fig. 1a; *Lasthenia californica, Clarkia purpurea,*

Agoseris heterophylla, Calycadenia pauciflora, Hemizonia congesta, and *Plantago erecta)* and

three dominant invasive annual grasses (Fig. 1b; *Avena fatua, Taeniatherum caput-medusae,* and

Bromus hordeaceus). In December 2017, seeds were sown into physan-washed pots with a

mixture of 65% field-collected soil and 35% sterile autoclaved sand to improve drainage. Each

species was grown alone (five replicates per species) to assess the microbial communities of

individual plant species. To assess the effects of competition between natives and invasives, each

forb species was grown with each grass species in a pairwise factorial design for a total of 18

different species combinations with five replicates per pair. Pots were placed in a shade-house

open to natural temperature variation in the UC Davis Orchard Park Greenhouse and soil

moisture was maintained with an automatic drip irrigation system (Fig. 1c).

Sample collection

In April 2018, we sampled rhizosphere microbial communities by gently shaking off soil

from plant roots, then submerging the roots in autoclaved nanopure water in a 50 mL conical

tube and vortexing to obtain the rhizosphere soil similar to Edwards et al. [1]. We sampled

rhizospheres prior to flowering to minimize differences in composition due to developmental

stage [16,36]. Conical tubes were then centrifuged at 4000 g for 1 minute to obtain rhizosphere

soil pellets. In plant mixtures, we sampled the joint rhizosphere microbiome as roots were

completely interwoven, suggesting strong interactions between individual rhizospheres.

To assess the competitive response of each species, we harvested, dried, and weighed

aboveground plant biomass and used it to calculate log response ratios. Roots were not included

due to the difficulty of identifying roots to species in the competition treatment. Log response

ratio was calculated as ln(weight grown together/mean weight grown alone) [46]. Values below

zero indicate competition with more negative values signifying a stronger response to

competition.

Molecular methods and sequence generation

We randomized samples using a random number generator prior to DNA extraction. We

then extracted the DNA from samples using the MoBio PowerSoil DNA Extraction kit (MO BIO

Laboratories, Inc., Carlsbad, CA, USA) with one minor modification to manufacturer's

instructions: samples were bead beat on the "homogenize" setting for 1 minute during step 5.

Samples were then sent to the Integrated Microbiome Resource (IMR) at Dalhousie University to

amplify and sequence the 16S rRNA gene using the 515FB-806RB primer set [47,48] and ITS2

region using the ITS86F-ITS4R primer set [49]. The amplification and sequencing process is

described in Comeau et al. [50]. In addition to experimental samples, we sent five subsamples of

the background field-collected soil-sand mix, three replicates each of a positive technological

control using DNA extracted from the ZymoBIOMICS Microbial Community Standard (Zymo

Research Inc., Irvine, CA, USA), and a negative technological control where no sample was

added during DNA extraction to the IMR for sequencing. The sequence reads generated for this

16S rRNA gene and ITS region amplicon project were deposited at Genbank under accession no.

PRJNA666893.

Sequence processing

Sequencing primers were removed using the Cutadapt (v. 2.3) program and the resulting

sequences were then processed using the DADA2 workflow in R . 16S rRNA gene amplicon

sequences were trimmed at 280 base pairs for forward reads and 250 base pairs for reverse reads

based on their error profiles to maintain a quality score above 10. ITS region amplicon sequences

were not trimmed to preserve the ability to merge reads due to the inherent length variation of

the ITS region. Both 16S rRNA gene and ITS region amplicon reads were then truncated at the

first quality score of 2 and reads with an expected error greater than 2 were removed. We then

merged paired end reads, created an Amplicon Sequence Variant table (ASV), and identified

chimeric reads using removeBimeraDenovo, which were subsequently removed prior to

downstream analyses (~4.1% of sequence reads for 16S rRNA gene, ~0.93% for ITS region).

For 16S rRNA gene ASVs, we assigned taxonomy using the RDP Naive Bayesian

Classifier algorithm and the SILVA high quality ribosomal RNA database v. 132 [51]. For ITS

ASVs, taxonomy was inferred with the UNITE (v. 8.0) database modified to include a

representative ITS amplicon sequence for each host plant, i.e. *H. congesta* (MF964056.1), *C.*

purpurea (MG235764.1), *P. erecta* (AY101909.1), *A. fatua* (FJ794719.1), *L. californica*

(AY043514.1), *A. heterophylla* (AY218965.1), *T. caput-medusae* (AJ608153.1), *B. hordeaceus*

(KP987344.1) [52,53]. ITS-x (v. 1.1.1) was then run on the unique ITS ASVs to pull out only

sequences that had a fungal ITS region based on kingdom-specific hidden markov models [54].

ITS-x was unable to detect a fungal ITS region in 744 of the ASVs (~13.46% of ASVs). These

ASVs were thus considered putatively non-fungal and were removed from downstream analysis.

Decontam's prevalence method was used to identify putative contaminants with a

threshold of 0.5 which will identify sequences that have a higher prevalence in negative controls

than in true samples [55]. This method identified 55 contaminants in the 16S rRNA gene data

and 32 in the ITS amplicon data which were then removed from the analysis.

For the 16S rRNA gene dataset, unique ASVs were aligned using the MAFFT Alignment

Program (v 7.402) on XSEDE with standard parameters through the CIPRES Science Gateway

V. 3.3 (https://www.phylo.org/) [56,57]. We then built a phylogenetic tree using FastreeMP on

XSEDE with standard parameters also through CIPRES [58] and rooted the tree using an

archaeal outgroup. We then created a phyloseq object from the resulting 16S ASV table, Silva

taxonomy table, phylogeny, and mapping file using the phyloseq package in R [59]. Sequences

that were taxonomically assigned to chloroplasts and mitochondria were then removed and

samples were rarefied to 9,434 reads per sample for comparison purposes.

For the ITS region data, we created a phyloseq object from the ITS ASV table, UNITE

taxonomy table, and mapping file using the phyloseq package in R. Samples were rarefied to

7,557 reads per sample for comparison purposes.

Data analysis (Q1-Q2)

To evaluate how rhizosphere communities varied in structure (beta diversity) between

groups and in competition, we used the vegan package [60] in R to conduct a Principal

Coordinates Analysis (PCoA) on weighted UniFrac of each sample for rarefied 16S rRNA gene

data and on Bray-Curtis dissimilarities for rarefied ITS region data. We tested for significant

differences in mean centroids between groups (native forbs, invasive grasses, grass-forb pairs)

using the adonis function to perform a permutational multivariate analysis of variance

(PERMANOVA) and between mean dispersions using the betadisper function in vegan. We then

conducted post-hoc analyses using the adonis.pair function with Benjamini-Hochberg p-value

corrections [61]. We also investigated whether these groups varied compared to background soil

(i.e. the community in the soil-sand mix at the start of the experiment) by performing

PERMANOVA tests to ascertain whether plant hosts were drivers of community assembly.

Finally, we investigated the degree of species-level variation within both grasses and forbs by

running PERMANOVA tests on each group separately to evaluate whether there was more

variation within or between groups.

To assess how rhizosphere communities varied in alpha diversity between grasses, forbs

and grass-forb pairs, we calculated Shannon diversity of each sample using the estimate_richness

function in the phyloseq package, then conducted a Kruskal-Wallis test to test whether alpha

diversity varied between these groups. We then conducted post-hoc analyses using a Dunn Test

with Benjamini-Hochberg p-value corrections.

Data analysis (Q3)

To evaluate whether a larger proportion of the joint microbiome community was

predicted to be sourced from forbs or grasses, we used SourceTracker, a Bayesian source

tracking classifier [62]. First we trained SourceTracker on the rarified ASV tables of forbs,

grasses, and background soil and then tested the trained Baysian model on the grass-forb pair

joint microbiomes to estimate the proportion of ASVs originating from each source. The trained

Baysian model assumes that joint microbiomes contain a combination of colonists from known

and unknown sources (i.e. anything not sampled here such as air, water, etc.) and estimates the

fraction of ASVs detected in the joint microbiome that originated from grasses, forbs,

background soil or unknown habitats. We used Wilcoxon signed-rank tests to assess significant

differences in the SourceTracker predicted proportions between different sources for the

bacterial community and fungal community, and then also tested for differences in the

proportions of each individual source between the two communities.

Data Analysis (Q4)

We used differential abundance analysis to identify microbial families whose relative

abundance varied between groups in order to investigate which families were driving observed

differences in the community structure and composition. To do this, we first summed the raw

read counts of each full data set (i.e. not rarefied) into families using the tax_glom function in

phyloseq. We used the DESeq2 R package [63] on the family counts to examine the $\log_2$fold

change (i.e. differential abundance) of families between pairwise contrasts of groups (forbs,

grasses, and grass-forb pairs). We focused on families that were identified as differentially

abundant in at least one pairwise comparison between forbs, grasses and grass-forb pairs with

Bonferroni corrected p-values < 0.01.

Data analysis (Q5)

To understand whether the relative abundances of these microbial families may be tied to

the competitive responses of grasses and forbs, we first identified which functional group was

competitively dominant by testing for differences in log competitive response ratios between

grasses and forbs in a linear mixed effect model with functional group as our predictor and a

random intercept for plant species combination to account for species-level differences.

To determine if the relative abundance of a specific microbial family was related to forb

or grass competitive response, we ran linear mixed effects models for each host group (grass and

forb) and each differentially abundant microbial family with log competitive response ratios as

the response variable and regularized log abundance of the focal family (calculated using the rld

function in the DESeq2 package; hereafter referred to as relative abundance) as the predictor

variable. Plant species combination was also included as a random intercept to account for

species-level differences.

Finally, we also calculated the proportion of each differentially abundant family in the

joint microbiome predicted to originate from either the individual forb or grass microbiomes by

grouping ASVs into taxonomic families and then summarizing the fraction of ASVs within each

family attributed to each possible source using SourceTracker.

Results

Q1: Do microbiomes differ between invasive grasses and native forbs?

Overall, the rhizosphere bacterial microbiome varied between invasive grasses and native

forbs, but the fungal microbiome did not. The structure of forb bacterial microbiomes was

significantly different from the structure of grass bacterial microbiomes ($p < 0.001$; Fig. 2).

Further, bacterial Shannon diversity (Fig. 3) was significantly higher in forb microbiomes than in

grass microbiomes ($p = 0.035$). Conversely, there were no significant differences in the structure

of fungal communities among treatments ($p = 0.072$; Fig. S1). There were also no differences in

fungal Shannon diversity among treatments ($p = 0.854$; Fig. S2). Further, microbial community

structure was significantly different between background soil and all treatment groups for both

the bacterial ($p < 0.001$, Fig. S3a) and fungal communities ($p < 0.001$, Fig. S3b).

The structure of individual microbiomes did not vary between forb species (bacterial: $p =$

0.069, Fig. S4a; fungal: $p = 0.755$, Fig. S5a), but did vary between grass species (bacterial $p =$

0.013, Fig. S4b; fungal: $p = 0.589$, Fig. S5b). The bacterial microbiomes of *T. caput-medusae*

were distinct from those of *B. hordeaceus* ($p = 0.049$) and *A. fatua* ($p = 0.047$). Statistical

differences between grass and forb bacterial microbiomes remained even after removing this

species and rerunning the analysis, suggesting larger variation between grass and forb functional

groups than within grass or forb groups. We therefore proceeded with the analysis of all species,

but focused the remainder of our results at the functional group level.

Q2: Do microbiomes under grass-forb competition differ from the microbiomes of (a) invasive

grasses and (b) native forbs?

Grass-forb pairs displayed significantly different joint bacterial microbiomes from

microbiomes of both functional groups grown alone (forb: $p < 0.001$; grass: $p < 0.001$; Fig. 2).

Grass-forb pair microbiomes also had marginally higher Shannon diversity than grass

microbiomes ($p = 0.059$), but did not significantly differ from forb microbiomes ($p = 0.292$). As

reported above, there were no significant differences found between treatments in the fungal

microbiome (either in structure or diversity), so we did not conduct post-hoc analyses to assess

pairwise contrasts between treatment groups.

Q3: Are microbiomes under grass-forb competition sourced more from invasive grasses, native

forbs, or equally from both?

For bacteria, similar fractions of the community in competition were predicted to have

originated from grasses and from forbs ($p = 0.341$, grass: $40.34 \pm 1.33\%$, forb: $42.13 \pm 1.14\%$).

Only relatively low proportions of the bacterial community came from background soil or

unknown sources (background: $8.57 \pm 0.48\%$, unknown: $8.96 \pm 0.60\%$). For fungi, the fraction

of the community predicted to originate from grasses was marginally higher than that from forbs

($p = 0.066$, grass: $38.88 \pm 2.26\%$, forb: $32.65 \pm 2.01\%$). Additionally, the fraction of the fungal

community originating from forbs was significantly lower than the fraction of the bacterial

community originating from forbs ($p < 0.001$). A small proportion of the fungal community was

predicted to derive from background soil ($5.16 \pm 0.67\%$), while a much larger proportion was

estimated to be from unknown sources ($23.31 \pm 2.11\%$). The contribution of unknown sources to

the fungal community was significantly higher than the contribution of unknown sources to the

bacterial community ($p < 0.001$).

Q4: Given compositional changes in the microbiome under grass-forb competition, which

specific microbial families are driving these changes?

Differential abundance analysis revealed a total of 19 bacterial families (Fig. S6a) that

varied between treatments, with 16 bacterial families varying between grass-forb pairs and either

grass or forb microbiomes and three additional families varying only between grass and forb

microbiomes. Of the 16 bacterial families that were differentially abundant under competition,

six families varied between both microbiomes of forbs and grasses grown alone and between

grass-forb pair microbiomes and individual microbiomes: Methylophilaceae, Fibrobacteraceae,

Clostridiaceae_1, Burkholderiaceae, Rhodocyclaceae, and Veillonellaceae (Fig. 4).

Methylophilaceae was the only family with higher relative abundance in forbs and

subsequently lower relative abundance in grass-forb pairs. Both Fibrobacteraceae and

Clostridiaceae_1 were higher in relative abundance in grass than forb microbiomes and were

subsequently higher in grass-forb pairs than in forbs alone. The relative abundance of

Burkholderiaceae was higher in grasses than in forbs, but decreased in grass-forb pairs.

Veillonellaceae displayed higher relative abundances in grasses than in forbs, and remained

higher in grass-forb pairs. Finally, the relative abundance of Rhodocyclaceae was higher in

grasses than in forbs, and lower in grass-forb pairs than in grasses. Of the remaining 13 families,

three families only displayed differential abundances between grasses and forbs and 10 families

displayed differential abundances between grass-forb pairs and forbs but not between grasses and

forbs (Fig. S7).

A total of three fungal families were differentially abundant between treatments (Fig.

S6b, Fig. S8). Of these, only Tubeufiaceae showed variation between both individual

microbiomes and grass-forb pairs, with higher relative abundances in forbs than grasses and

subsequently lower relative abundances in grass-forb pairs than in forbs. Ceratobasidiaceae

displayed lower relative abundances in grass-forb pairs than in grasses, and an unclassified

family from the order Sebacinales displayed higher abundance in grass-forb pairs than in forbs.

Q5: Are changes in the relative abundance of these microbial families correlated with the

competitive responses of forbs and grasses to each other?

Grasses were generally dominant over forbs, and this dominance was correlated with the

relative abundance of some bacterial families. Native forbs displayed stronger negative log

response ratios than invasive grasses ($p = 0.047$; Fig. 5), indicating that forbs were more

negatively affected by competition from grasses than the grasses were from forbs. Competitive

outcomes did, however, vary depending on the identity of the grass and the identity of the forb

within the species pair combination (Fig. S9), indicating a wide range of responses.

Of the six families that differentially varied between grass-forb pairs and either grasses or

forbs, the relative abundances of Methylophilaceae, Fibrobacteraceae, Clostridiaceae_1, and

Burkholderiaceae also displayed correlations with either forb or grass log response ratios (Fig.

6). Moreover, those families that displayed negative correlations with forb responses and/or

positive correlations with grass responses had higher predicted proportions originating from

grass microbiomes, while the one family that displayed the opposite correlations (positively

correlated with forb response/negative with grass response) had a higher predicted proportion

originating from forb microbiomes (Fig. 7).

The relative abundance of Methylophilaceae was significantly positively correlated with

log response ratio of forbs (est = 0.48, se = 0.14 , $p < 0.001$), and significantly negatively

correlated with log response ratio of grasses (est = -0.32, se = 0.13, $p = 0.021$), indicating that

as the relative abundance of this family decreased, forbs responded more negatively to grass

competition. Conversely, grasses were negatively correlated with higher relative abundances of

this family. This family also showed a higher proportion of ASVs originating from the forb

microbiome in grass-forb pairs.

The relative abundance of Fibrobacteraceae was significantly negatively correlated with

forb log response ratio (est = -0.51, se = 0.19, p = 0.009), but was not correlated with grass

response (est = -0.07, se = 0.18, p = 0.719; Fig 5b), indicating that as the relative abundance of

this family increased, the competitive effect of grasses on forbs intensified, while grasses were

unaffected by changes in this family. The relative abundance of Clostridiaceae_1 was

significantly positively correlated with grass log response ratio (est = 0.35, se = 0.16, p = 0.036),

but not with forbs (est = -0.26, se = 0.18, p = 0.139), indicating that as this family increased, the

competitive response of grasses to forbs weakened. A higher predicted proportion of ASVs

within these families originated from grasses, suggesting that grasses may have been driving

high relative abundance of Fibrobacteraceae and Clostridiaceae_1 in grass-forb pairs.

The relative abundance of Burkholderiaceae was negatively correlated with forb log

response ratio (est = -0.36 , se = 0.11, p = 0.002) and positively correlated with grass log

response ratio (est = 0.37, se = 0.11, p < 0.001), indicating that as the relative abundance of this

family increased, forbs were more negatively affected by competition from grasses and grasses

were less negatively affected by competition from forbs. Grasses also contributed a

disproportionate number of ASVs from this family to the joint microbiome.

In the remaining 15 bacterial families that displayed differential abundances among

treatments, we found no significant relationships to log competitive response ratios (Fig. S10).

For predicted proportions of these families originating from grasses and forbs, see Fig. S11.

Tubeufiaceae, the only fungal family that displayed differential abundance between forbs

and grasses, and between grass-forb pairs and forbs, displayed no correlation with either forb or

grass log competitive response ratios (Fig. S12). Tubeufiaceae was composed of a higher

predicted proportion of ASVs originating from forbs relative to grasses (Fig. S12).

Ceratobasidiaceae, which was significantly lower in grass-forb pairs than in grasses but did not

vary between grasses and forbs, showed positive correlations of its relative abundance to log

response ratios of grasses (est = 0.23, se = 0.04, $p < 0.001$) and negative correlations with forb

log response ratios (est = -0.11, se = 0.05, $p = 0.032$), indicating that as the relative abundance of

this family increased, forbs were more negatively affected by grasses and grasses were less

negatively affected by forbs. We found no other relationships between the remaining

differentially abundant fungal family and group log response ratios (Fig. S13).

Discussion

The rhizosphere bacterial community of native forbs is structurally distinct from those of

invasive grasses and has higher bacterial alpha diversity. Competition between grasses and forbs

in our system correlates with shifts in the bacterial rhizosphere of native forbs towards

communities dissimilar to both individual forb and grass microbiomes, with marginally higher

diversity than grasses alone. Studies in similar annual grasslands have found little change in

bacterial structure with increased invasive grass abundance [43,64,65] but invader dominance

has been linked to decreased bacterial diversity in other systems [36,66]. Contrary to Hortal et al.

[18] and Lozano et al. [19], we did not find that the new community resembled that of the

dominant competitor; instead, during competition, we observed shifts towards a novel

assemblage of bacteria distinct from both individual forb and grass microbiomes, and sourced

nearly equally from these groups. Competition is therefore associated with shifts in the

microbiomes of both invasive grasses and native forbs; however, these changes are linked with

more negative consequences for native forbs than invasive grasses.

Native annual forbs displayed a stronger negative response to competition than invasive

grasses. Plants were well-watered and harvested before shading could become problematic,

therefore virtually all competitive effects in this experiment likely resulted mainly from below-

ground competition. Given the large role played by microbes in nutrient availability and uptake,

observed competitive imbalances, and the substantial differences between forb and grass

microbiomes, it is likely these competitive outcomes are mediated by microbes. Invaded

grassland soils tend to be depleted in plant available nitrogen [35,64,65], but with increased

microbial activity and faster nitrogen cycling [25,34,65]. These microbe-driven changes may

selectively benefit fast-growing grasses over natives, especially in nutrient-poor serpentine soils

where added nitrogen has been found to preferentially aid invasives [67,68]. Invasive grasses

may further benefit from decreased microbial diversity, which has been linked to both decreased

nitrogen uptake by forbs and increased nitrogen uptake by grasses [69]. Beyond competition for

nutrients, grasses and forbs may also be indirectly competing through recruitment of microbes

harmful to natives [70] and/or helpful to invasives [39]. Other studies in serpentine grasslands

have shown that soil primed with invasive grass negatively affected the growth of native forbs

due to changes in the microbial community [71,72]. While our study was not able to disentangle

these various mechanisms, our results provide evidence that compositional shifts in the

rhizosphere microbiome may be reinforcing the competitive dominance of these grasses.

The relative abundance of 19 bacterial families showed significant variation between

grasses, forbs, and grass-forb pairs. Of these 19 families, we found links between bacterial

relative abundance and plant competitive response for four: two families were linked to

decreased competitive abilities in forbs (Burkholderiaceae, Fibrobacteraceae), one family was

linked to increased competitive ability in grasses (Clostridiaceae_1), and one family was linked

to both decreased competitive ability of grasses and increased competitive ability of forbs

(Methylophilaceae). All four of these families were differentially abundant between forb and

grass rhizospheres and between grass-forb pairs and either grasses or forbs.

We found only one bacterial family, Methylophilaceae, that was linked to increased

competitive ability in forbs. This family was more abundant in forbs than in grasses, and

declined in relative abundance in grass-forb pairs. Still, higher relative abundance of

Methylophilaceae was associated with a less negative competitive response in forbs and a more

negative response in grasses. Bacteria in this family have been found in soils with high heavy

metal content and include microbes critical for heavy metal attenuation and immobilization

[73,74]. Members of Methylophilaceae also form tight symbioses with plants [75–77] and are

even known to promote seed germination and plant growth [76]. This family may include

important taxa for serpentine-adapted species, with a decrease in relative abundance driving

negative effects through reduced heavy metal attenuation and plant growth. Forbs also

contributed a higher percentage of ASVs from this family to the joint grass-forb rhizosphere

microbiome, further supporting close associations between native forbs and this group. Invasive

grass dominance may therefore be partially due to a decrease in locally adapted microbes similar

to Cavalieri et al. [31].

Relative abundances of the bacterial families Fibrobacteraceae and Clostridiaceae_1 were

higher in grasses than in forbs and remained higher in competition treatments. Grasses were

likely driving these higher relative abundances, as they contributed a higher percentage of ASVs

from these families to the joint rhizosphere microbiome in pairs. The relative abundance of

Fibrobacteraceae was significantly negatively correlated with forb competitive response, but not

grass competitive response. Conversely, the relative abundance of Clostridiaceae_1 was

positively correlated with grass response, but did not display a relationship with forb response.

Although Fibrobacteraceae is typically lower in abundance in soils with high heavy metal

content [74], this family is also positively associated with invasive grass dominance [78].

Further, both Fibrobacteraceae and Clostridiaceae_1 are known for their cellulose-degrading

properties and are often found in rumen [79,80], and therefore may be important taxa for the

breakdown of organic matter [81]. These two families also contain known nitrogen-fixers [82–

84]. Together, these taxa may be increasing nutrient availability and thereby disproportionately

helping fast-growing invasive annual grasses [67,68].

The relative abundance of the bacterial family Burkholderiaceae was higher in grasses

than in forbs, but was found to be lower in competition. Even with this decrease, grasses may

have benefited from their presence - grass competitive response was positively correlated with

their relative abundance while forb's displayed a negative competitive response correlation.

Grasses also contributed a higher proportion of ASVs from this family to the joint rhizosphere

under competition. This family is considered a keystone member in grasslands and is often

associated with plants as endophytes or pathogens [85–88]. Burkholderiaceae contains members

known for their antimicrobial properties [89–91], nitrogen-fixing abilities [82,92], and

competitive dominance, especially in N-limited environments [93] like serpentine. Some

members of Burkholderiaceae are even known to suppress fungal pathogens including the turf

grass pathogen Rhizoctonia (family Ceratobasidiaceae) [94,95]. It is possible that the

Burkholderiacae observed in competition here are either providing positive functional benefits to

grasses, or are less harmful to grasses than forbs as pathogens.

In contrast to our bacterial rhizosphere findings, we found no significant differences in

fungal communities across treatments, either in terms of structure or diversity. This suggests

only loose fungal associations with plant hosts and a lack of plant-driven selection for specific

fungal taxa, possibly due to functional redundancy of fungi across large geographic scales [96].

Within annual grasslands, however, there is evidence of invader-driven fungal shifts [43] and

increases in fungal relative abundance [97]. While we found one fungal family that was

differentially abundant between both plants grown alone and together (Tubeufiaceae), the only

relationship between fungal family relative abundance and competition we found was in

Ceratobasidiaceae, which did not vary between grass and forb microbiomes, but was lower in

pairs relative to grasses. Ceratobasidiaceae play important ecological roles as plant pathogens,

saprotrophs or plant endophytes [98–100]. The negative correlations with forb response and

positive correlation with grass response observed here may indicate that this group is acting as a

beneficial endophyte of grasses, or may be explained by evolved pathogen resistance in grasses

and lack thereof in forbs [101,102].

Our limited fungal findings may indeed indicate that fungi are generally less important

than bacteria for competitive outcomes between plants [103]. Alternatively, fungi may be

important, but complex fungal-fungal interactions at the community level may cancel out fitness

effects on plants by beneficial and detrimental fungi [104]. In addition, experimental constraints

such as unaccounted fungal regional source pools, plant age at sampling, and methodology may

have prevented the observation of fungal relative abundance effects on host plant biomass.

Critical fungi are often locally sourced from airborne spores or local watersheds during plant

development [96] and thus may have been missing from the shade-house where our experiment

took place. This is highlighted in our findings that ~23% of the fungal community was predicted

to come from 'unknown' sources. Another limitation is plant age at time of sampling, as fungi

may form tighter associations later on in plant development or have greater effects on plant

fitness over longer time periods [16]. Finally, tightly associated fungi may be root endophytes

(e.g. Emam et al. [105]) or found in the rhizoplane (e.g. Edwards et al. [1]), rather than the

rhizosphere sampled here.

Conclusions

Our study found that native annual forbs are host to a diverse and distinct community of

microbes possibly helping them cope with serpentine's high heavy metal content, but these close

associations are disrupted by invasive grasses. The joint bacterial rhizosphere of invasive grasses

and native forbs differed from those of native forbs and invasive grasses grown alone, suggesting

that rhizosphere shifts may be related to competition. Both forbs and grasses experienced shifts

in their microbiome under competition, but invasive grasses behaved as might be expected of

generalists and were notably less affected by these changes. Moreover, the relative abundances

of four bacterial families were correlated with plant host competitive outcomes, providing

support for the role of microbe-mediated grass dominance in this system. More research is

needed to clarify whether the observed changes in plant competitive response were driven by

these candidate bacterial families and future work should focus on these families through

individual inoculations to tease apart their role in plant growth and competition. While microbial

diversity was unaffected by competition in this simplified pairwise experiment, the rhizosphere

may become homogenized as grasses invade over larger landscapes [43,66], with selective

benefits for grasses over forbs [69]. Increasing native forb abundance in this ecosystem and

inoculating soil with locally adapted microbes through restoration efforts may help to alleviate

decreases in critical bacteria and lead to better fitness outcomes for native forbs. More generally,

our study provides further affirmation of the importance of plant-microbe relationships to the

maintenance of ecosystems facing dominance by invasive species.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

The datasets and scripts supporting the conclusions of this article are available in the GitHub

repository, DOI:10.5281/zenodo.4422181. The sequence reads generated for this 16S rRNA gene

and ITS region amplicon project were deposited at Genbank under accession no. PRJNA666893.

Competing interests

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

Funding

This work was largely supported by grants from the UC Davis Center for Population Biology to

MLL and CLE and the UC Davis Center for Population Biology Hardman Foundation Research

Award to MLL. This work was also supported in part by a grant awarded to undergraduate

assistant HK from the California Native Plant Society. The funders had no role in study design,

data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

MLL, CLE, and HK conceived of, planned, and carried out the experiment. MLL and CLE

processed and analyzed the data and wrote the paper. All authors contributed to the final version

of the manuscript.

Acknowledgements

We would like to thank undergraduate intern, Vyvy Ha, with their help on this project, including

processing plant biomass and assisting with shade-house related activities. We are also grateful

to Paul Aigner for permission to use their photos. Finally, we would like to thank Susan P.

Harrison, Andrew M. Latimer and Jonathan A. Eisen for useful conversations and suggestions

related to this work, providing lab resources and space and contributing funds to this project.

References

1. Edwards J, Johnson C, Santos-Medellin C, Lurie E, Podishetty NK, Bhatnagar S, et al. Structure, variation, and assembly of the root-associated microbiomes of rice. Proceedings of the National Academy of Sciences of the United States of America. 2015;112:E911–20.

2. Berendsen RL, Pieterse CMJ, Bakker PAHM. The rhizosphere microbiome and plant health. Trends in Plant Science. 2012;17:478–86.

3. Philippot L, Raaijmakers JM, Lemanceau P, van der Putten WH. Going back to the roots: the microbial ecology of the rhizosphere. Nature Reviews Microbiology. Nature Publishing Group; 2013;11:789–99.

4. Bever JD, Platt TG, Morton ER. Microbial Population and Community Dynamics on Plant Roots and Their Feedbacks on Plant Communities. Annu Rev Microbiol. Annual Reviews; 2012;66:265–83.

5. Richardson AE, Barea J-M, McNeill AM, Prigent-Combaret C. Acquisition of phosphorus and nitrogen in the rhizosphere and plant growth promotion by microorganisms. Plant Soil. 2009;321:305–39.

6. Yang J, Kloepper JW, Ryu C-M. Rhizosphere bacteria help plants tolerate abiotic stress. Trends in Plant Science. 2009;14:1–4.

7. Doubková P, Suda J, Sudová R. The symbiosis with arbuscular mycorrhizal fungi contributes to plant tolerance to serpentine edaphic stress. Soil Biology and Biochemistry. 2012;44:56–64.

8. Xue P-P, Carrillo Y, Pino V, Minasny B, McBratney AB. Soil Properties Drive Microbial

Community Structure in a Large Scale Transect in South Eastern Australia. Scientific

Reports. Nature Publishing Group; 2018;8:11725.

9. Singh BK, Munro S, Potts JM, Millard P. Influence of grass species and soil type on

rhizosphere microbial community structure in grassland soils. Applied Soil Ecology.

2007;36:147–55.

10. de Ridder-Duine AS, Kowalchuk GA, Klein Gunnewiek PJA, Smant W, van Veen JA, de

Boer W. Rhizosphere bacterial community composition in natural stands of Carex arenaria

(sand sedge) is determined by bulk soil community composition. Soil Biology and

Biochemistry. 2005;37:349–57.

11. Aleklett K, Leff JW, Fierer N, Hart M. Wild plant species growing closely connected in a

subalpine meadow host distinct root-associated bacterial communities. PeerJ. PeerJ Inc.;

2015;3:e804.

12. Lakshmanan V. Chapter Three - Root Microbiome Assemblage is Modulated by Plant Host

Factors. In: Bais H, Sherrier J, editors. Advances in Botanical Research [Internet]. Academic

Press; 2015 [cited 2020 Dec 7]. p. 57–79. Available from:

http://www.sciencedirect.com/science/article/pii/S0065229615000658

13. Rúa MA, Antoninka A, Antunes PM, Chaudhary VB, Gehring C, Lamit LJ, et al. Home-field

advantage? evidence of local adaptation among plants, soil, and arbuscular mycorrhizal fungi

through meta-analysis. BMC Evol Biol. 2016;16:122.

14. Schultz PA, Michael Miller R, Jastrow JD, Rivetta CV, Bever JD. Evidence of a mycorrhizal

mechanism for the adaptation of Andropogon gerardii (Poaceae) to high- and low-nutrient

prairies. Am J Bot. 2001;88:1650–6.

15. Johnson NC, Wilson GWT, Bowker MA, Wilson JA, Miller RM. Resource limitation is a

driver of local adaptation in mycorrhizal symbioses. PNAS. National Academy of Sciences;

2010;107:2093–8.

16. Chaparro JM, Badri DV, Vivanco JM. Rhizosphere microbiome assemblage is affected by

plant development. The ISME Journal. Nature Publishing Group; 2014;8:790–803.

17. Siefert A, Zillig KW, Friesen ML, Strauss SY. Soil microbial communities alter conspecific

and congeneric competition consistent with patterns of field coexistence in three Trifolium

congeners. Journal of Ecology. 2018;106:1876–91.

18. Hortal S, Lozano YM, Bastida F, Armas C, Moreno JL, Garcia C, et al. Plant-plant

competition outcomes are modulated by plant effects on the soil bacterial community. Sci

Rep. 2017;7:17756.

19. Lozano YM, Hortal S, Armas C, Pugnaire FI. Soil micro-organisms and competitive ability

of a tussock grass species in a dry ecosystem. Journal of Ecology. 2019;107:1215–25.

20. Muller-Landau HC. Plant diversity rooted in pathogens. Nature. Nature Publishing Group;

2014;506:44–5.

21. Mangan SA, Schnitzer SA, Herre EA, Mack KML, Valencia MC, Sanchez EI, et al. Negative

plant–soil feedback predicts tree-species relative abundance in a tropical forest. Nature.

Nature Publishing Group; 2010;466:752–5.

22. Klironomos JN, McCune J, Hart M, Neville J. The influence of arbuscular mycorrhizae on

the relationship between plant diversity and productivity. Ecology Letters. 2000;3:137–41.

23. Heijden MGAVD, Bardgett RD, Straalen NMV. The unseen majority: soil microbes as

drivers of plant diversity and productivity in terrestrial ecosystems. Ecology Letters.

2008;11:296–310.

24. Franche C, Lindström K, Elmerich C. Nitrogen-fixing bacteria associated with leguminous

and non-leguminous plants. Plant Soil. 2009;321:35–59.

25. Hawkes CV, Wren IF, Herman DJ, Firestone MK. Plant invasion alters nitrogen cycling by

modifying the soil nitrifying community. Ecology Letters. 2005;8:976–85.

26. Lee MR, Flory SL, Phillips RP. Positive feedbacks to growth of an invasive grass through

alteration of nitrogen cycling. Oecologia. 2012;170:457–65.

27. Funk JL, Vitousek PM. Resource-use efficiency and plant invasion in low-resource systems.

Nature. 2007;446:1079–81.

28. Hodge A, Fitter AH. Microbial mediation of plant competition and community structure.

Functional Ecology. 2013;27:865–75.

29. Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MTJ. Assembly

and ecological function of the root microbiome across angiosperm plant species. PNAS.

2018;115:E1157–65.

30. Eppinga MB, Rietkerk Max, Dekker Stefan C., De Ruiter Peter C., Van der Putten Wim H.,

Van der Putten Wim H. Accumulation of local pathogens: a new hypothesis to explain exotic

plant invasions. Oikos. 2006;114:168–76.

31. Cavalieri A, Bak F, Garcia-Lemos AM, Weiner J, Nicolaisen MH, Nybroe O. Effects of

Intra- and Interspecific Plant Density on Rhizosphere Bacterial Communities. Front

Microbiol [Internet]. Frontiers; 2020 [cited 2020 Oct 26];11. Available from:

https://www.frontiersin.org/articles/10.3389/fmicb.2020.01045/full

32. Kandlikar GS, Johnson CA, Yan X, Kraft NJB, Levine JM. Winning and losing with

microbes: how microbially mediated fitness differences influence plant diversity. Ecology

Letters. 2019;22:1178–91.

33. Levine JM, Pachepsky E, Kendall BE, Yelenik SG, Lambers JHR. Plant–soil feedbacks and

invasive spread. Ecology Letters. 2006;9:1005–14.

34. Esch EH, Hernández DL, Pasari JR, Kantor RSG, Selmants PC. Response of soil microbial

activity to grazing, nitrogen deposition, and exotic cover in a serpentine grassland. Plant Soil.

2013;366:671–82.

35. Carey CJ, Blankinship JC, Eviner VT, Malmstrom CM, Hart SC. Invasive plants decrease

microbial capacity to nitrify and denitrify compared to native California grassland

communities. Biol Invasions. 2017;19:2941–57.

36. Shi S, Nuccio E, Herman DJ, Rijkers R, Estera K, Li J, et al. Successional Trajectories of

Rhizosphere Bacterial Communities over Consecutive Seasons. mBio [Internet]. American

Society for Microbiology; 2015 [cited 2020 Oct 7];6. Available from:

https://mbio.asm.org/content/6/4/e00746-15

37. Komatsu KJ, Simms EL. Invasive legume management strategies differentially impact

mutualist abundance and benefit to native and invasive hosts. Restoration Ecology.

2020;28:378–86.

38. Rodríguez-Caballero G, Caravaca F, Díaz G, Torres P, Roldán A. The invader Carpobrotus

edulis promotes a specific rhizosphere microbiome across globally distributed coastal

ecosystems. Science of The Total Environment. 2020;719:137347.

39. Rout ME, Chrzanowski TH, Westlie TK, DeLuca TH, Callaway RM, Holben WE. Bacterial

endophytes enhance competition by invasive plants. American Journal of Botany.

2013;100:1726–37.

40. Callaway RM, Cipollini D, Barto K, Thelen GC, Hallett SG, Prati D, et al. Novel Weapons:

Invasive Plant Suppresses Fungal Mutualists in America but Not in Its Native Europe.

Ecology. 2008;89:1043–55.

41. Blank RR, Mackey B, Morgan T. Do a native and two exotic grasses respond differently in

soils conditioned by native vegetation versus an exotic grass? Rhizosphere. 2016;2:38–47.

42. Eskelinen A, Harrison S. Exotic plant invasions under enhanced rainfall are constrained by

soil nutrients and competition. Ecology. 2014;95:682–92.

43. Gravuer K, Eskelinen A, Winbourne JB, Harrison SP. Vulnerability and resistance in the

spatial heterogeneity of soil microbial communities under resource additions. PNAS.

National Academy of Sciences; 2020;117:7263–70.

44. Igwe AN, Vannette RL. Bacterial communities differ between plant species and soil type,

and differentially influence seedling establishment on serpentine soils. Plant Soil.

2019;441:423–37.

45. Porter SS, Chang PL, Conow CA, Dunham JP, Friesen ML. Association mapping reveals

novel serpentine adaptation gene clusters in a population of symbiotic Mesorhizobium. The

ISME Journal. Nature Publishing Group; 2017;11:248–62.

46. Hedges LV, Gurevitch J, Curtis PS. The Meta-Analysis of Response Ratios in Experimental

Ecology. Ecology. 1999;80:1150–6.

47. Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R

gene primer greatly increases detection of SAR11 bacterioplankton. Aquatic Microbial

Ecology. 2015;75:129–37.

48. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA

primers for marine microbiomes with mock communities, time series and global field

samples. Environmental Microbiology. 2016;18:1403–14.

49. Beeck MOD, Lievens B, Busschaert P, Declerck S, Vangronsveld J, Colpaert JV.

Comparison and Validation of Some ITS Primer Pairs Useful for Fungal Metabarcoding

Studies. PLOS ONE. Public Library of Science; 2014;9:e97629.

50. Comeau AM, Douglas GM, Langille MGI. Microbiome Helper: a Custom and Streamlined

Workflow for Microbiome Research. mSystems [Internet]. American Society for

Microbiology Journals; 2017 [cited 2020 Oct 5];2. Available from:

https://msystems.asm.org/content/2/1/e00127-16

51. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal

RNA gene database project: improved data processing and web-based tools. Nucleic Acids

Res. 2013;41:D590–6.

52. Kõljalg U, Larsson K-H, Abarenkov K, Nilsson RH, Alexander IJ, Eberhardt U, et al.

UNITE: a database providing web-based methods for the molecular identification of

ectomycorrhizal fungi. New Phytologist. 2005;166:1063–8.

53. UNITE Community. UNITE general FASTA release for Fungi 2. UNITE Community

[Internet]. 2019 [cited 2020 Dec 7]; Available from:

https://plutof.ut.ee/#/doi/10.15156/BIO/786353

54. Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, et al. Improved

software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi

and other eukaryotes for analysis of environmental sequencing data. Methods in Ecology and

Evolution. 2013;4:914–9.

55. Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical

identification and removal of contaminant sequences in marker-gene and metagenomics data.

Microbiome. 2018;6:226.

56. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002;30:3059–66.

57. Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES Science Gateway for inference of

large phylogenetic trees. 2010 Gateway Computing Environments Workshop (GCE). 2010.

p. 1–8.

58. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees for

Large Alignments. PLoS One [Internet]. 2010 [cited 2020 Dec 7];5. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2835736/

59. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. PLoS One [Internet]. 2013 [cited 2020 Dec 7];8.

Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3632530/

60. Dixon P. VEGAN, a package of R functions for community ecology. Journal of Vegetation

Science. 2003;14:927–30.

61. Salazar G. EcolUtils: Utilities for community ecology analysis [Internet]. 2020 [cited 2020

Dec 7]. Available from: https://github.com/GuillemSalazar/EcolUtils

62. Knights D, Kuczynski J, Charlson ES, Zaneveld J, Mozer MC, Collman RG, et al. Bayesian

community-wide culture-independent microbial source tracking. Nature Methods. Nature

Publishing Group; 2011;8:761–3.

63. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biology. 2014;15:550.

64. Carey CJ, Beman JM, Eviner VT, Malmstrom CM, Hart SC. Soil microbial community

structure is unaltered by plant invasion, vegetation clipping, and nitrogen fertilization in

experimental semi-arid grasslands. Front Microbiol [Internet]. Frontiers; 2015 [cited 2020

Jun 1];6. Available from: https://www.frontiersin.org/articles/10.3389/fmicb.2015.00466/full

65. Dickens SJM, Allen EB, Santiago LS, Crowley D. Exotic annuals reduce soil heterogeneity

in coastal sage scrub soil chemical and biological characteristics. Soil Biology and

Biochemistry. 2013;58:70–81.

66. Kowalchuk GeorgeA, Buma DS, de Boer W, Klinkhamer PGL, van Veen JA. Effects of

above-ground plant species composition and diversity on the diversity of soil-borne

microorganisms. Antonie Van Leeuwenhoek. 2002;81:509.

67. Going BM, Hillerislambers J, Levine JM. Abiotic and biotic resistance to grass invasion in

serpentine annual plant communities. Oecologia. 2009;159:839–47.

68. Eskelinen A, Harrison S. Erosion of beta diversity under interacting global change impacts in

a semi-arid grassland. Journal of Ecology. 2015;103:397–407.

69. Wagg C, Schlaeppi K, Banerjee S, Kuramae EE, van der Heijden MGA. Fungal-bacterial

diversity and microbiome complexity predict ecosystem functioning. Nat Commun.

2019;10:4841.

70. Stinson KA, Campbell SA, Powell JR, Wolfe BE, Callaway RM, Thelen GC, et al. Invasive

Plant Suppresses the Growth of Native Tree Seedlings by Disrupting Belowground

Mutualisms. PLOS Biology. Public Library of Science; 2006;4:e140.

71. Batten KM, Scow KM, Espeland EK. Soil Microbial Community Associated with an

Invasive Grass Differentially Impacts Native Plant Performance. Microb Ecol. 2008;55:220–

8.

72. Ba L, Facelli E, Facelli JM. Plant-mycorrhizal fungi feedbacks: potential accomplices of

Avena barbata's high invasiveness. Plant Ecol. 2018;219:1045–52.

73. Drewniak L, Krawczyk PS, Mielnicki S, Adamska D, Sobczak A, Lipinski L, et al.

Physiological and Metagenomic Analyses of Microbial Mats Involved in Self-Purification of

Mine Waters Contaminated with Heavy Metals. Front Microbiol [Internet]. Frontiers; 2016

[cited 2020 Jul 2];7. Available from:

https://www.frontiersin.org/articles/10.3389/fmicb.2016.01252/full

74. Syranidou E, Thijs S, Avramidou M, Weyens N, Venieri D, Pintelon I, et al. Responses of

the Endophytic Bacterial Communities of Juncus acutus to Pollution With Metals, Emerging

Organic Pollutants and to Bioaugmentation With Indigenous Strains. Front Plant Sci

[Internet]. Frontiers; 2018 [cited 2020 Jul 2];9. Available from:

https://www.frontiersin.org/articles/10.3389/fpls.2018.01526/full

75. Trotsenko YuA, Ivanova EG, Doronina NV. Aerobic Methylotrophic Bacteria as

Phytosymbionts. Microbiology. 2001;70:623–32.

76. Fedorov DN, Doronina NV, Trotsenko YuA. Phytosymbiosis of aerobic methylobacteria:

New facts and views. Microbiology. 2011;80:443.

77. Macey MC, Pratscher J, Crombie AT, Murrell JC. Impact of plants on the diversity and

activity of methylotrophs in soil. Microbiome. 2020;8:31.

78. Fahey C, Koyama A, Antunes PM, Dunfield K, Flory SL. Plant communities mediate the

interactive effects of invasion and drought on soil microbial communities. The ISME Journal.

Nature Publishing Group; 2020;1–14.

79. Rubino F, Carberry C, M Waters S, Kenny D, McCabe MS, Creevey CJ. Divergent

functional isoforms drive niche specialisation for nutrient acquisition and use in rumen

microbiome. ISME J. 2017;11:932–44.

80. Ozbayram EG, Ince O, Ince B, Harms H, Kleinsteuber S. Comparison of Rumen and Manure

Microbiomes and Implications for the Inoculation of Anaerobic Digesters. Microorganisms.

Multidisciplinary Digital Publishing Institute; 2018;6:15.

81. Nuccio EE, Starr E, Karaoz U, Brodie EL, Zhou J, Tringe SG, et al. Niche differentiation is

spatially and temporally regulated in the rhizosphere. ISME J. 2020;1–16.

82. Dos Santos PC, Fang Z, Mason SW, Setubal JC, Dixon R. Distribution of nitrogen fixation

and nitrogenase-like sequences amongst microbial genomes. BMC Genomics. 2012;13:162.

83. Abdul Rahman N, Parks DH, Vanwonterghem I, Morrison M, Tyson GW, Hugenholtz P. A

Phylogenomic Analysis of the Bacterial Phylum Fibrobacteres. Front Microbiol [Internet].

2016 [cited 2020 Oct 7];6. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4704652/

84. Pepe-Ranney C, Koechli C, Potrafka R, Andam C, Eggleston E, Garcia-Pichel F, et al. Non-

cyanobacterial diazotrophs mediate dinitrogen fixation in biological soil crusts during early

crust formation. ISME J. 2016;10:287–98.

85. Eberl L, Vandamme P. Members of the genus Burkholderia: good and bad guys. F1000Res.

2016;5:1007.

86. Banerjee S, Schlaeppi K, Heijden MGA van der. Keystone taxa as drivers of microbiome

structure and functioning. Nature Reviews Microbiology. 2018;16:567.

87. Genin S, Boucher C. Lessons learned from the genome analysis of ralstonia solanacearum.

Annu Rev Phytopathol. Annual Reviews; 2004;42:107–34.

88. Mansfield J, Genin S, Magori S, Citovsky V, Sriariyanum M, Ronald P, et al. Top 10 plant

pathogenic bacteria in molecular plant pathology. Molecular Plant Pathology. 2012;13:614–

29.

89. Becker MH, Brucker RM, Schwantes CR, Harris RN, Minbiole KPC. The Bacterially

Produced Metabolite Violacein Is Associated with Survival of Amphibians Infected with a

Lethal Fungus. Appl Environ Microbiol. American Society for Microbiology; 2009;75:6635–

8.

90. Depoorter E, Bull MJ, Peeters C, Coenye T, Vandamme P, Mahenthiralingam E.

Burkholderia: an update on taxonomy and biotechnological potential as antibiotic producers.

Appl Microbiol Biotechnol. 2016;100:5215–29.

91. Shehata HR, Ettinger CL, Eisen JA, Raizada MN. Genes Required for the Anti-fungal

Activity of a Bacterial Endophyte Isolated from a Corn Landrace Grown Continuously by

Subsistence Farmers Since 1000 BC. Front Microbiol [Internet]. Frontiers; 2016 [cited 2020

Oct 8];7. Available from: https://www.frontiersin.org/articles/10.3389/fmicb.2016.01548/full

92. Estrada-de los Santos P, Palmer M, Chávez-Ramírez B, Beukes C, Steenkamp ET, Briscoe L,

et al. Whole Genome Analyses Suggests that Burkholderia sensu lato Contains Two

Additional Novel Genera (Mycetohabitans gen. nov., and Trinickia gen. nov.): Implications

for the Evolution of Diazotrophy and Nodulation in the Burkholderiaceae. Genes (Basel)

[Internet]. 2018 [cited 2020 Oct 7];9. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6116057/

93. Elliott GN, Chou J-H, Chen W-M, Bloemberg GV, Bontemps C, Martínez-Romero E, et al.

Burkholderia spp. are the most competitive symbionts of Mimosa, particularly under N-

limited conditions. Environmental Microbiology. 2009;11:762–78.

94. Kang Y, Carlson R, Tharpe W, Schell MA. Characterization of Genes Involved in

Biosynthesis of a Novel Antibiotic from Burkholderia cepacia BC11 and Their Role in

Biological Control of Rhizoctonia solani. Appl Environ Microbiol. 1998;64:3939–47.

95. Chapelle E, Mendes R, Bakker PAH, Raaijmakers JM. Fungal invasion of the rhizosphere

microbiome. The ISME Journal. Nature Publishing Group; 2016;10:265–8.

96. Talbot JM, Bruns TD, Taylor JW, Smith DP, Branco S, Glassman SI, et al. Endemism and

functional convergence across the North American soil mycobiome. PNAS. National

Academy of Sciences; 2014;111:6341–6.

97. Wolkovich EM, Lipson DA, Virginia RA, Cottingham KL, Bolger DT. Grass invasion

causes rapid increases in ecosystem carbon and nitrogen storage in a semiarid shrubland.

Glob Change Biol. 2010;16:1351–65.

98. Parmeter JR. Rhizoctonia Solani, Biology and Pathology. University of California Press;

1970.

99. Sen R, Hietala AM, Zelmer CD. Common anastomosis and internal transcribed spacer RFLP

groupings in binucleate Rhizoctonia isolates representing root endophytes of Pinus sylvestris,

Ceratorhiza spp. from orchid mycorrhizas and a phytopathogenic anastomosis group. New

Phytologist. 1999;144:331–41.

100. Veldre V, Abarenkov K, Bahram M, Martos F, Selosse M-A, Tamm H, et al. Evolution of

nutritional modes of Ceratobasidiaceae (Cantharellales, Basidiomycota) as revealed from

publicly available ITS sequences. Fungal Ecology. 2013;6:256–68.

101. Flory SL, Kleczewski N, Clay K. Ecological consequences of pathogen accumulation on an

invasive grass. Ecosphere. 2011;2:art120.

102. Anthony MA, Frey SD, Stinson KA. Fungal community homogenization, shift in dominant

trophic guild, and appearance of novel taxa with biotic invasion. Ecosphere. 2017;8:e01951.

103. Levine JM, Adler PB, Yelenik SG. A meta-analysis of biotic resistance to exotic plant

invasions. Ecology letters. 2004;7:975–89.

104. Facelli E, Smith SE, Facelli JM, Christophersen HM, Smith FA. Underground friends or

enemies: model plants help to unravel direct and indirect effects of arbuscular mycorrhizal

fungi on plant competition. New Phytologist. 2010;185:1050–61.

105. Emam T. Local soil, but not commercial AMF inoculum, increases native and non-native

grass growth at a mine restoration site. Restoration Ecology. 2016;24:35–44.

Figure 1. (a) Native annual forb species (from left to right, top to bottom): *Clarkia purpurea,*

Calycadenia pauciflora, Lasthenia californica, Agoseris heterophylla, Plantago erecta, and

Hemizonia congesta. (b) Invasive annual grasses (from left to right): *Bromus hordeaceus,*

Taeniatherum caput-medusae, and *Avena fatua.* (c) Shade-house set-up. Photos of *C.*

purpurea, P. erecta, H. congesta, C. pauciflora, B. hordeaceus, T. caput-medusae, and *A.*

fatua by Paul Aigner. Remaining photos by authors.

888   **Figure 2.** Bacterial microbiome structure differs between grasses and forbs, and shifts during

889   competition. Principal coordinates analysis (PCoA) visualization of weighted UniFrac distances

890   of bacterial communities associated with the rhizosphere. Points in the ordination are colored and

891   represented by shapes based on the rhizosphere of forbs grown alone (purple circles), grasses

892   grown alone (green triangles) and grass-forb pairs, i.e. competition (yellow squares).

  Figure 3. Shannon diversity of bacterial microbiomes is highest in forbs. Shannon diversity was

903   used to assess alpha diversity for bacterial microbiomes for each treatment (forbs grown alone,

904   grasses grown alone, competition). Comparisons that are significantly different from each other

905   are notated by different letters (e.g. a vs. b).

 Figure 4. Bacterial families that are differentially abundant between grass and forb microbiomes,

916   and between grass-forb pairs and either grass or forb microbiomes. Using DESeq2, bacterial

917   families were identified whose relative abundance differed significantly between treatments

918   (forbs grown alone, grasses grown alone, grown together in competition). Each plot shows the

919   relative abundance of families for each treatment and comparisons that are significant from each

920   other are notated by different letters (e.g. a vs. b).

928   **Figure 5.** Native forbs were more negatively affected by competition. Log response ratios for

929   forbs and grasses were calculated to determine the dominant competitor and were based on dried

930   aboveground biomass weight. Values below zero indicate competition with more negative values

931   signifying a stronger response to competition. Error bars here represent the standard error of the

932   mean.

942   **Figure 6.** Bacterial families contributing to competitive outcomes. Linear mixed effects models

943   assessing the contribution of normalized relative abundance of bacterial families to competitive

944   outcomes (log response ratios) for forbs (purple) or grasses (green). Solid line indicates

945   significance ($p < 0.05$), dashed line indicates nonsignificance.

Figure 7. Proportion of ASVs predicted to colonize from each source for each of the four

differentially abundant families that were correlated with competition. SourceTracker was used

to predict whether ASVs in competition treatments originated from grasses (dark green), forbs

(purple), the background soil mix used in the experiment (blue), or unknown habitats (light

green) and then the results were summarized by grouping ASVs by taxonomic family. Families

that were negatively correlated with forb competitive response and/or positively correlated with

grass competitive response were predominantly sourced from grass. Methylophilaceae, the one

family that showed a positive relationship with forb competitive response and a negative

relationship with grass competitive response, was predominantly sourced from forbs.