1 **FLAgellum Member 8 modulates extravascular trypanosome distribution in the**

2 **mammalian host**

4 Estefanía Calvo Alvarez[1]*, Aline Crouzols[1] and Brice Rotureau*[1]

6 [1] Trypanosome Transmission Group, Trypanosome Cell Biology Unit, Department of

7 Parasites and Insect Vectors and INSERM U1201, Institut Pasteur, Paris, France

13 *Corresponding   author: estefaniacalvoalvarez@gmail.com   (ECA),

14 brice.rotureau@pasteur.fr (BR)

Abstract

The African trypanosome flagellum is essential in multiple aspects of the parasite development. In the mammalian infective form of this protist, FLAgellar Member 8 (FLAM8) is a large protein distributed along the entire flagellum that is suspected to be involved in host-parasite interactions. Analyses of knockdown and knockout trypanosomes demonstrated that FLAM8 is not essential *in vitro* for survival, growth, motility and slender to stumpy differentiation. Functional investigations in experimental infections showed that *FLAM8*-deprived trypanosomes are able to establish and maintain the infection in the blood circulation, and to differentiate into transmissible stumpy forms. However, bioluminescence imaging revealed that *FLAM8*-null parasites exhibit an impaired dissemination in the extravascular compartment, especially in the skin, that is partially restored by the addition of a single rescue copy of *FLAM8*. To our knowledge, FLAM8 is the first example of a flagellar protein that modulates *T. brucei* parasite distribution in the host tissues, contributing to the maintenance of extravascular parasite populations in mammalian anatomical niches.

Author summary

African trypanosomes are blood and tissue-dwelling protists responsible for African trypanosomiases, a group of neglected tropical diseases in sub-Saharan Africa, including sleeping sickness in humans and nagana in cattle. Although underestimated for long, the importance of extravascular parasite niches has been recently re-discovered both in animal models and in infected individuals. The trypanosome flagellum is an essential organelle for the parasites, yet its contribution to host-parasite interactions, parasite tropism and spatiotemporal dissemination in the mammalian host remain scarce. To this end, we investigated the role of a flagellar protein termed FLAgellar Member 8 (FLAM8), present only at the distal tip of the flagellum in insect parasites, but differentially localized to the entire flagellum length in trypanosomes found in the vertebrate host. Our findings revealed that FLAM8 is dispensable *in vitro* for survival, growth, motility and differentiation of *T. brucei.* However, interestingly, FLAM8 modulated the extravascular dissemination of trypanosomes in the mammalian host, especially in the skin. FLAM8 is the first example of a flagellar protein that modulates *T. brucei* parasite distribution in extravascular host tissues.

Introduction

Trypanosoma brucei is an extracellular parasite responsible for African trypanosomiases in sub-Saharan Africa, a group of neglected tropical diseases including sleeping sickness in humans and nagana in cattle. African trypanosomes are blood and tissue-dwelling protists transmitted by the bite of the blood-feeding tsetse fly (*Glossina* genus). In the mammalian host, parasites have to face different micro-environments, including deadly challenges by multiple types of host immune responses and the variable availability of carbon sources. This requires major morphological and metabolic adaptations, driven by the activation of specific gene expression programs, that are critical for life-cycle progression [1-3]. Recently, the importance of extravascular tropism for *T. brucei* has been re-discovered and emphasized in animal models: in addition to the brain, parasites actually occupy most mammalian tissues, especially the skin and the adipose tissues [4-6]. However, when, where and how trypanosomes adapt to survive within these extravascular environments is not understood yet.

The trypanosome flagellum is an essential organelle anchored along the surface of the cell body and present in all stages of its development [7]. It is essential for parasite viability [8], cell division and morphogenesis [9], attachment to the tsetse salivary glands [10] and motility [11]. In the insect host, the flagellum remains at the forefront of the cell it tracts and is likely to be involved in sensory and signaling functions required for host-parasite interactions (reviewed in [12, 13]). In the mammalian host, flagellar motility was shown to be critical for establishment and maintenance of bloodstream infection [14]. Nevertheless, the overall contributions of the trypanosome

flagellum to parasite tropism and spatiotemporal dissemination dynamics in the

mammalian host are important aspects that still remained to be explored more in depth.

Our proteomic analysis of intact flagella purified from the insect stage of the parasite

identified a group of flagellar membrane and matrix proteins with unique patterns and

dynamics [15]. Amongst them, one large protein (3,075 amino acids) termed FLAgellar

Member 8 (FLAM8) is present only at the distal tip of the flagellum in the insect

procyclic form [15-17]. Interestingly, it is redistributed along the entire length of the

flagellum in mammalian bloodstream forms (BSF), including in stumpy transmissible

stages (Calvo-Álvarez *et al.*, submitted), which may imply a specific function for FLAM8

in the vertebrate host. Therefore, we hypothesized that the stage-specific redistribution

of FLAM8 could be involved in host-parasite interactions, possibly affecting parasite

homing and dynamics in host tissues. Here, we investigated the possible roles of

FLAM8 *in vitro* in terms of survival, proliferation, motility and differentiation. In addition,

experimental infections in mice monitored by bioluminescence imaging allowed us to

decipher the involvement of FLAM8 in parasite dissemination and spreading in both

the intra- and extra-vascular compartments of the mammalian host. Potential

consequences for parasite transmission along with cellular mechanisms of tissue

tropism are further discussed.

Methods

1. Strains, culture and *in vitro* differentiation

The AnTat 1.1E Paris pleomorphic clone of *Trypanosoma brucei brucei* was derived from a strain originally isolated from a bushbuck in Uganda in 1966 [27]. The monomorphic *T. brucei* strain Lister 427 [28] was also used. All bloodstream forms (BSF) were cultivated in HMI-11 medium supplemented with 10% (v/v) fetal bovine serum (FBS) [29] at 37°C in 5% CO_2. For *in vitro* slender to stumpy BSF differentiation, we used 8-pCPT-2′-O-Me-5′-AMP, a nucleotide analog of 5′-AMP from BIOLOG Life Science Institute (Germany). Briefly, $2x10^6$ cultured pleomorphic AnTat 1.1E slender forms were incubated with 8-pCPT-2′-O-Me-5′-AMP (5 µM) for 48 h [30]. For specific experiments, *in vitro* differentiation of BSF into PCF was performed by transferring freshly differentiated short stumpy forms into SDM-79 medium supplemented with 10% (v/v) FBS, 6 mM cis-aconitate and 20 mM glycerol at 27°C [31].

Monomorphic BSF "Single Marker" (SM) trypanosomes are derivatives of the Lister 427 strain, antigenic type MITat 1.2, clone 221a [32], and express the T7 RNA polymerase and tetracycline repressor. *FLAM8[RNAi]* cells express complementary single-stranded RNA corresponding to a fragment of the *FLAM8* gene from two tetracycline-inducible T7 promoters facing each other in the pZJM vector [33] integrated in the rDNA locus [34]. Addition of tetracycline (1 µg/mL) to the medium induces expression of sense and anti-sense RNA strands that can anneal to form double-stranded RNA (dsRNA) and trigger RNAi. For *in vivo* RNAi studies in mice, doxycycline hyclate (Sigma Aldrich) was added in sugared drinking water (0.2 g/L doxycycline hyclate combined with 50 g/L sucrose).

2. Generation of FLAM8 mutants

For the generation of the *FLAM8^RNAi* cell lines, a 380 bp (nucleotides 6665-7044) fragment of *FLAM8* (Tb927.2.5760), flanked by 5' HindIII and 3' XhoI restriction sites to facilitate subsequent cloning, was selected using the RNAit algorithm (http://trypanofan.bioc.cam.ac.uk/software/RNAit.html) in order to ensure that the targeted sequence was distinct from any other genes to avoid any cross-RNAi effects [35]. This *FLAM8* DNA fragment was synthesized by GeneCust Europe (Dudelange, Luxembourg) and inserted into the HindIII-XhoI digested pZJM vector [33]. The pZJM-FLAM8 plasmid was linearized with NotI prior to transfection using nucleofector technology (Lonza, Italy) as described previously [36]. The cell line was further engineered for endogenous tagging of *FLAM8* with an mNeonGreen (mNG) at its C-terminal end by using the p3329 plasmid [37], carrying a *FLAM8* gene fragment corresponding to *FLAM8* ORF nucleotides 8892-9391. Prior to nucleofection, NruI linearization of p3329-FLAM8-mNG plasmid was performed. For *in vivo* experiments in mice, *FLAM8^RNAi* parasites were finally modified by integrating a plasmid encoding for the chimeric triple reporter which combines the red-shifted firefly luciferase PpyREH9 and the tdTomato red fluorescent protein fused with a TY1 tag [18]. Transformants were selected with the appropriate antibiotic concentrations: phleomycin (1 μg/ml), blasticidin (5 μg/ml), G418 (2 μg/ml), and puromycin (0.1 μg/ml). Clonal populations were obtained by limiting dilution. Cell culture growth was monitored with an automatic Muse cell analyser (Merck Millipore, Paris).

For generating the *FLAM8* knockout and rescue cell lines, all insert templates were synthesized by GeneCust Europe (Dudelange, Luxembourg). For breaking the first allele, the 300 first nucleotides of the *FLAM8* gene flanking sequences were added on each side of a HYG resistance cassette (Fig. S2). For a complete disruption of the

FLAM8 locus, a second selectable marker (PAC) was flanked with the *FLAM8*-flanking

sequence at 5' and by 300 nucleotides of the *FLAM8* ORF (nucleotides 501-800) at 3'.

For generating an add-back rescue cell line, due to the important size of the *FLAM8*

ORF (9,228 nucleotides), the PAC selection marker was replaced by a BLE marker

cassette flanked by the 300 first nucleotides of the *FLAM8* 5' untranslated region (UTR)

and by the nucleotides 1 to 500 of the *FLAM8* ORF for reinsertion into the endogenous

locus of the knockout cell line. PCR amplifications of the DNA fragments bearing the

FLAM8 flanking sequences and the appropriate resistance markers were used for

nucleofection and generation of all cell lines. The primers used are listed below: 5'-

CATGACTTTACGTGTTTGGGCAC-3' (FW, located 82 bp upstream the flanking

5'UTR sequence); 5'-CTTGCTTGTTTCTGTTTCGCAAC-3' (RV, 130 bp downstream

the flanking 3'UTR sequence, used to replace one WT allele by HYG resistance

cassette); 5'- GCACACTAAAACTCATTGAAAGCC-3' (RV, 926 bp downstream the

ATG codon of *FLAM8*, used for second WT allele replacement by PAC cassette and

rescue line generation). All knockout and rescue cell lines were further transfected to

express the chimeric triple reporter protein PpyRE9H/TY1/tdTomato for multimodal *in*

vivo imaging approaches as described elsewhere [18]. Selection-marker recovery was

confirmed by screening individual clones in multi-well plates. Transformants were

selected with the appropriate antibiotic concentrations: phleomycin (1 µg/ml),

blasticidin (5 µg/ml), puromycin (0.1 µg/ml) and hygromycin (2.5 µg/ml). Clonal

populations were obtained by limiting dilution and cell culture growth was monitored

with an automatic Muse cell analyzer (Merck Millipore, Paris).

Knockout and rescue cell lines were validated by whole-genome sequencing (BGI,

Hong Kong). Briefly, genomic DNA from parental and mutant cell lines were sequenced

by the HiSeq4000 sequence system (Illumina), generating about 10 million 100-bp

reads and compared to that of the *T. brucei brucei* AnTat 1.1E Paris reference strain.

In addition, some validation of the construct integrations in mutants were performed by

PCR analysis according to standard protocols (Fig. S2 and Table 1).

Table 1. Oligonucleotides used for PCR validation of the Δ*FLAM8* knockout and

rescue cell lines.

Primer	Sequence	Target	
1	CATGACTTTACGTGTTTGGGCAC	*FLAM8* WT allele (82 nt upstream 5'UTR)	F
2	GCACACTAAAACTCATTGAAAGCC	*FLAM8* WT allele (926 nt downstream ATG)	R
3	CGTCCGAGGGCAAAGGAATAG	Hygromycin cassette	R
4	GACCGCGCACCTGGTGCATG	Puromycin cassette	R
5	GTGGCCGAGGAGCAGGACTGA	Phleomycin cassette	R

Orientation of primers: F, forward; R, reverse.

3. Motility analyses

In silico 2D tracking was performed as previously described [38]. For each BSF strain,

10 to 20 movies were recorded for 20 seconds (50 ms of exposure). Trypanosomes at

1×10^6 parasites/ml were maintained in matrix-dependent HMI-11 medium containing

0,5% methylcellulose at 37°C and were observed under the 10x objective of an

inverted DMI-4000B microscope (Leica) coupled to an ORCA-03G (Hamamatsu) or a

PRIM95B (Photometrics) camera. Movies were converted with the MPEG Streamclip

V.1.9b3 software (Squared 5) and analyzed with the medeaLAB CASA Tracking V.5.5

software (medea AV GmbH).

4. *In vitro* bioluminescence quantification and analysis

To perform the parasite density / bioluminescence intensity assay, BSF parasites were counted, centrifuged and resuspended in fresh HMI-11 medium. Then, 100 µl of this suspension containing 10^6 parasites were transferred into black clear-bottom 96-well plates and serial 2-fold dilutions were performed in triplicate adjusting the final volume to 200 µl with 300 µg/ml of beetle luciferin (Promega, France). Luciferase activity was quantified after 10 min of incubation with an IVIS Spectrum imager (PerkinElmer). Imaging data analysis was performed with the Living Image 4.3.1 software (PerkinElmer) by drawing regions of interest with constant size for well plate quantification. Total photon flux was calculated after removal of intensity values from WT parasites and / or parasite-free medium corresponding to the background noise. Results were analyzed as mean ± SD of three independent experiments.

5. Mouse infection and ethical statements

Seven-week-old male BALB/c mice were purchased from Janvier Laboratory (sub-strain BALB/cAnNRj) and used as models for experimental infection and monitoring of the bioluminescence signal with the IVIS Spectrum imager (PerkinElmer). This study was conducted in strict accordance with the recommendations from the Guide for the Care and Use of Laboratory Animals of the European Union (European Directive 2010/63/UE) and the French Government. The protocol was approved by the "Comité d'éthique en expérimentation animale de l'Institut Pasteur" CETEA 89 (Permit number: 2012-0043 and 2016-0017) and undertaken in compliance with the Institut Pasteur Biosafety Committee (protocol CHSCT 12.131). BR is authorized to perform experiments on vertebrate animals (license #A-75-2035) and is responsible for all the experiments conducted personally or under his supervision. For *in vivo* infections, groups of four and three animals (*FLAM8* knockdown and knockout-infected mice,

respectively) were injected intraperitoneally (IP) with 10^5 slender BSF parasites,

washed in TDB (Trypanosome Dilution Buffer: 5 mM KCl, 80 mM NaCl, 1 mM

$MgSO_4*7H_2O$, 20 mM Na_2HPO_4, 2 mM NaH_2PO_4, 20 mM glucose) and resuspended

in 100 μl of PBS prior animal inoculation.

6. *In vivo* bioluminescence imaging, analysis and parasitemia quantifications

Infection with bioluminescent parasites was monitored daily by detecting the

bioluminescence signal in whole animals with the IVIS Spectrum imager (PerkinElmer).

The equipment consists of a cooled charge-coupled camera mounted on a light-tight

chamber with a nose cone delivery device to keep the mice anaesthetized during

image acquisition with 1.5-2% isoflurane. A heated stage is comprised within the IVIS

Spectrum imager in order to maintain optimum body temperature. D-luciferin

potassium salt (Promega) stock solution was prepared in sterile PBS at 33.33 mg/ml,

and stored in a -20°C freezer. To produce bioluminescence, mice were inoculated by

the intraperitoneal route with 150 μl of D-luciferin stock solution (250 mg/Kg body

weight). After 10 minutes of incubation to allow substrate dissemination, all mice were

anaesthetized in an oxygen-rich induction chamber with 1.5-2% isoflurane, and images

were acquired by using automatic exposure (30 seconds to 5 minutes) depending on

signal intensity. Images were analyzed with the Living Image software version 4.3.1

(PerkinElmer). Data were expressed in total photons/second (p/s) corresponding to the

total flux of bioluminescent signal according to the selected area (regions of interest

with constant size covering the total body of the mouse). The background noise was

removed by subtracting the bioluminescent signal of the control mouse from the

infected ones for each acquisition. For parasite dissemination analyses, a minimum

value of photons/second (p/s) was set for all animals in every time point in order to

quantify the exact dissemination area (in cm^2) over the whole animal body. Parasitemia

was determined daily following tail bleeds and assayed by automated fluorescent cell

counting with a Muse cytometer (Merck-Millipore, detection limit at 10^2 parasites/ml)

according to the manufacturer's recommendations. The quantification of the total

intravascular parasite population was assessed by calculating the blood volume of all

animals according to their body weight and referring to daily parasitemia. To quantify

the number of extravascular parasites, a region of interest with a constant size was

used to obtain the total number of parasites within the entire animal bodies (total

bioluminescence). Subsequently, the total number of parasites present in the vascular

system was subtracted, resulting in estimating the total parasite population colonizing

the extravascular compartments at a given time point.

7. Immuno-fluorescence analysis (IFA)

Cultured parasites were washed twice in TDB and spread directly onto poly-L-lysine

coated slides. For methanol fixation, slides were air-dried for 10 min, fixed in methanol

at -20°C for 5 min and rehydrated for 20 min in PBS. For immunodetection, slides were

incubated for 1 h at 37°C with the appropriate dilution of the first antibody in 0.1% BSA

in PBS. After 3 consecutive 5 min washes in PBS, species and subclass-specific

secondary antibodies coupled to the appropriate fluorochrome (Alexa 488, Cy3, Cy5

Jackson ImmunoResearch) were diluted 1/400 in PBS containing 0.1% BSA and were

applied for 1 h at 37°C. After washing in PBS as indicated above, slides were finally

stained with 4',6-diamidino-2-phenylindole (DAPI, 1 µg/ml) for visualization of

kinetoplast and nuclear DNA content, and mounted under cover slips with ProLong

antifade reagent (Invitrogen), as previously described [7]. Slides were observed under

an epifluorescence DMI4000 microscope (Leica) with a 100x objective (NA 1.4), an

EL6000 (Leica) as light excitation source and controlled by the Micro-Manager V1.4.22

software (NIH), and images were acquired using an ORCA-03G (Hamamatsu) or a

PRIM95B (Photometrics) camera. Images were analyzed with ImageJ V1.8.0 (NIH).

The monoclonal antibody mAb25 (anti-mouse IgG2a, 1:10) was used as a flagellum

marker as it specifically recognizes the axoneme protein *Tb*SAXO1 [39]. FLAM8 was

detected using: i) a specific rabbit serum (1:500) kindly provided by Paul McKean

(University of Lancaster, UK), or ii) a monoclonal anti-mNeonGreen antibody (anti-

mouse IgG2c, 1:100, ChromoTek). Stumpy BSF were identified at the molecular level

with a rabbit polyclonal anti-PAD1 antibody (kindly provided by Keith Matthews,

University of Edinburgh; dilution 1:300) [40]. In the case of RNAi knockdown

experiments, IFA signals were normalized using the signal obtained in non-induced

controls as a reference.

8. Measurements, normalization and statistical analyses

Standardization of fluorescent signals was carried out by parallel setting of raw

integrated density signals in all the images to be compared in ImageJ V1.8.0 (NIH).

For clarity purposes, the brightness and contrast of several pictures were adjusted

after their analysis in accordance to editorial policies. Statistical analyses and plots

were performed with XLSTAT 2019.2.01 (Addinsoft) in Excel 2016 (Microsoft) or Prism

V8.2.1 (GraphPad). Statistical analyses include linear regression analyses for

bioluminescence / fluorescence intensity vs. parasite density, two-sided ANOVA /

ANCOVA tests for growth curves, with Tukey ad-hoc post-tests for inter-group

comparison, all at 95% confidence.

Results

1. *FLAM8* RNAi silencing does not affect parasite survival *in vitro* and *in vivo*.

The differential distribution of FLAM8 in the flagellum of the different trypanosome stages (Calvo-Álvarez *et al.*, submitted) raises the question of its specific functions during the parasite cycle. In order to investigate the potential role(s) of FLAM8 in the mammalian host, BSF were first engineered for inducible RNAi knockdown of *FLAM8* in a monomorphic strain expressing a mNeonGreen-tagged version of FLAM8. In order to monitor their behavior in mouse by whole-body imaging approaches, these *FLAM8::mNG FLAM8^{RNAi}* mutants were subsequently transformed to overexpress a chimeric triple reporter protein [18]. Upon RNAi induction with tetracycline for 72 h, *FLAM8* expression was reduced by 60% at the mRNA level (Fig. 1A) and became undetectable at the protein level by immunofluorescence (Fig. 1B). Parasite growth was monitored over 6 days upon induction of RNAi and no impact on proliferation was observed (Fig. 1C), which indicates that FLAM8 is not essential for survival of BSF parasites in cell culture conditions.

Figure 1. Characterization of the *FLAM8^{RNAi}* strain *in vitro*. A) Expression of *FLAM8* mRNA assessed by RT-PCR by the comparative C_T method in control, non-induced and induced *FLAM8^{RNAi}* parasites (72 h). **B)** Immunofluorescence pictures of non-induced (upper panels) and induced (bottom panels) *FLAM8^{RNAi} FLAM8::mNG* BSF during 72 h. Methanol-fixed trypanosomes were stained with an anti-mNG antibody (green) and DAPI for DNA content (blue). The scale bars represent 5 µm. **C)** Growth curves of control, non-induced and induced *FLAM8^{RNAi}* BSF parasites. All cell lines received 1 µg tetracycline during 6 days. Control parasites are Lister 427 "Single Marker" BSF parasites that do not bear the pZJM-FLAM8 plasmid for RNAi silencing.

Results represent the mean (± standard deviation, SD) of three independent
experiments.

Then, the linear correlation between the emitted bioluminescence and the number of
parasites was assessed in an IVIS Spectrum imager prior to *in vivo* challenge (Fig.
S1). In order to get insights into the function of FLAM8 in the mammalian host, groups
of male BALB/c mice were infected by the intraperitoneal route with 10^5 BSF parasites
of the parental, non-induced and induced cell lines (Fig. 2). *In vivo* RNAi silencing of
FLAM8 was maintained in mice by the addition of doxycycline in sugared drinking
water 48 h prior infection and until the end of the experiment. The course of the
infection was monitored daily by i) quantifying the parasitemia, and ii) acquiring the
bioluminescent signal emitted by the parasites in the entire organism with an IVIS
Spectrum imager. The number of parasites in the extravascular compartment at a
given timepoint can be extrapolated by subtracting the known number of trypanosomes
in the vascular system (parasitemia x blood volume, according to body weight) from
the total number of parasites in the organism (total bioluminescence). No differences
were detected neither in the establishment of the infection and the subsequent
variations in the number of intravascular parasites (Fig. 2A), nor in the number of
parasites occupying extravascular tissues (Fig. 2B) and the animal survival (not
shown) (Fig. 2, A-B). Similar population profiles were observed over the course of the
infection for both IV and EV parasites in each of the three independent groups of
infected mice (Fig. 2, C-E). Finally, in the extravascular compartment, no significant
differences were detected in terms of parasite dissemination over the entire animal
body in any of the groups tested (Fig. 2, F-G).

Figure 2. Functional investigations on *FLAM8^RNAi* parasites *in vivo* in the mammalian host. Groups of 4 BALB/c mice were injected intraperitoneally with either control, non-induced or induced *FLAM8^RNAi* BSF trypanosomes. One PBS-injected BALB/c animal was used as negative control. **A)** Number of parasites in the blood (intravascular, IV) of infected BALB/c mice during the course of the infection (5 days). **B)** Number of parasites in the extravascular compartment (extravascular, EV) of the same mice. **C, D, E)** Comparison of the number of trypanosomes present in the blood circulation (IV, continuous line) and in extravascular tissues (EV, dotted line) during the entire experimental infection. Significant differences are indicated with * ($p < 0.05$). **F)** Dissemination of control, non-induced and induced *FLAM8^RNAi* parasites, measured over the entire animal body (in cm^2) through the total bioluminescent surface, during the entire infection course. Results represent means ± standard deviation (SD). **G)** Representative normalized *in vivo* images of the bioluminescence radiance signal (in photons / second / cm^2 / steradian) emitted from BALB/c mice infected with control, non-induced and induced *FLAM8^RNAi* parasites 4 days post-infection (non-infected technical control mice were negative for bioluminescence, not shown).

2. *FLAM8* knockout affects trypanosome spreading in the mammalian host.

Considering that i) *FLAM8* RNAi silencing efficiency was only partial (40% *FLAM8* mRNA left after 72 h of induction), and that ii) the parental strain used for this first strategy was monomorphic (i.e. unable to differentiate into tsetse adapted stumpy stages), we reasoned that a gene knockout approach in a pleomorphic strain would be possibly more appropriate to evaluate the potential role(s) of FLAM8 during the mammalian host infection. Therefore, a Δ*FLAM8* knockout cell line was generated in pleomorphic BSF trypanosomes by homologous recombination. The replacement of

both *FLAM8* alleles by distinct resistance cassettes was verified by whole-genome

sequencing and PCRs (Fig. S2, A-B). The trypanosome cell lines generated were

further transformed to express the chimeric triple reporter construct as previously

described [18], and the linear correlation between the bioluminescence signal and the

total number of parasites was analyzed for all strains (Fig. S3). In Δ*FLAM8* knockout

parasites where a rescue copy of one *FLAM8* allele was added back into its

endogenous locus, the FLAM8 distribution was restored along the entire flagellum, as

assessed by immunofluorescence analysis (Fig. 3A). No impact on parasite growth in

culture conditions was observed (Fig. 3B), and the absence of *FLAM8* did not affect

parasite motility *in vitro* (Fig. 3C). We also studied whether the loss of FLAM8 could

have impacted the total length of the flagellum based on the measurement of the signal

obtained with the axonemal marker mAb25: no difference was observed among BSF

lines (Fig. 3D). Compatible with our previous observations after RNAi silencing, these

results show that pleomorphic trypanosomes tolerated the loss of *FLAM8 in vitro*.

Figure 3. Characterization of the Δ*FLAM8* null mutants *in vitro*. A)

Immunofluorescence pictures of parental, Δ*FLAM8* knockout and rescue pleomorphic

BSF parasites labelled with the anti-FLAM8 (green) and mAb25 (axoneme in magenta)

antibodies, DAPI staining for DNA content (blue). Scale bars show 5 µm. **B)** Growth

curve of one parental, three Δ*FLAM8* subclones and one rescue pleomorphic BSF

trypanosome cell lines over 6 consecutive days. **C)** Motility tracking analysis showing

the average speeds of BSF cell lines in matrix-dependent culture medium. No

statistical differences were observed. **D)** Measurements of the flagellum length based

on the axonemal marker mAb25 profiles in parental, three Δ*FLAM8* subclones and

rescue parasites. No statistical differences were found. The number of parasites

considered for quantifications (N) is indicated below graphs (C) and (D). Results

represent the mean ± standard deviation (SD) of three independent experiments.

Then, functional investigations in the mammalian host were performed by infecting

BALB/c mice either with the pleomorphic parental strain, three distinct Δ*FLAM8*

knockout subclones or one Δ*FLAM8* strain bearing a rescue copy of *FLAM8* (Fig. 4).

The infections were monitored daily during 4 weeks by quantifying the parasitemia and

the bioluminescence signals emitted from whole animals. Null mutant parasites were

able to establish an infection in the bloodstream as well as in the extravascular

compartment. In the blood circulation, the overall amounts of parasites throughout the

whole experimental infection were similar in all strains (Fig. 4A). The only significant

differences between the three Δ*FLAM8* subclones as compared to parental-infected

BALB/c mice were observed at days 5 and 6, coinciding with the first peak of

parasitemia (Fig. 4A). The absence of *FLAM8* did not prevent the *in vivo* differentiation

of proliferative slender into transmissible stumpy parasites in the intravascular

compartment (Fig. 4B), compatible with *in vitro* observations upon differentiation of all

mutant lines with no statistical differences (Fig. S4, A-B). In addition, freshly-

differentiated pleomorphic FLAM8-deprived parasites were able to further differentiate

and maintain as procyclic trypanosomes *in vitro* (Fig. S4C). On the other hand,

quantitative analyses of extravascular parasites showed a different scenario. Unlike in

the intravascular compartment, slightly yet significantly lower numbers of extravascular

trypanosomes were observed at days 6-12, 19 and 26 post-infection with Δ*FLAM8*

subclones as compared to mice infected with the parental strain (Fig. 4C). To note,

distribution profiles of all pleomorphic strains were different from those observed in

mice infected with monomorphic parasites: larger amounts of trypanosomes were seen

occupying the extravascular compartment reaching up to $5\text{-}8\text{x}10^9$ parasites, while the maximum amount found in the bloodstream never exceeded $7\text{x}10^7$ trypanosomes (Fig. S5).

Interestingly, the subsequent quantification of the parasite spreading over the whole animal bodies showed that the depletion of *FLAM8* resulted in a partially yet significantly impaired dissemination of all Δ*FLAM8* null mutant subclones in the extravascular compartment. This was mostly restored in trypanosomes bearing a rescue copy of *FLAM8* (Fig. 4D). The observation of whole-body bioluminescence images just after the first peak of parasitemia, when levels of parasitemia were low or undetectable in all mice, i.e. when the bioluminescence signal mostly came from parasites in the extravascular compartment, showed that this significant spreading impairment started to affect Δ*FLAM8* knockout parasites early during the infection course (Fig. 4E). Individual dissections at day 27 at the end of the experimental infection revealed that the impaired distribution of *FLAM8*-deprived trypanosomes in the extravascular compartment was not detectable in internal organs, with no differences in bioluminescence signal intensities among strains and / or organs (Fig. S6). These results indicate that FLAM8 is involved in the modulation of trypanosome distribution in the extravascular compartment of the mammalian host, especially in the skin.

Figure 4. Functional investigations on the Δ*FLAM8* null mutants *in vivo* in the mammalian host. Groups of 3 BALB/c mice were injected intraperitoneally with either one parental, three ΔF*LAM8* null subclones or one rescue strains. One PBS-injected BALB/c animal was used as negative control. **A)** Total number of parasites in the blood of infected mice (intravascular, IV) during the course of the infection (4 weeks).

Statistically significant differences between the parental strain and the three Δ*FLAM8*

subclones are indicated with *** p<0.0005. **B)** Representative immunofluorescence

pictures of transmissible short stumpy parasites found in the blood of BALB/c mice

infected with parental, Δ*FLAM8* and rescue parasites after the first peak of parasitemia.

Anti-PAD1 antibody was used as a surface-exposed differentiation marker (red), DAPI

staining for DNA content (blue). Scale bars represent 5 μm. **C)** Total number of

extravascular (EV) trypanosomes in the same mice. Statistically significant differences

between the parental strain and two or the three Δ*FLAM8* subclones are indicated with

** (p<0.005) or *** (p<0.0005), respectively. **D)** Dissemination of the parental, Δ*FLAM8*

subclones and rescue parasite strains, measured over the entire animal body (in cm^2)

through the total surface of bioluminescent signal, during the entire infection course.

Statistically significant differences between the parental strain and one, two or the

three Δ*FLAM8* subclones are indicated with * (p<0.05), ** (p<0.005) or *** (p<0.0005),

respectively. Results represent means ± standard deviation (SD). **E)** Normalized *in*

vivo images of the bioluminescence radiance intensity (in photons / second / cm^2 /

steradian) emitted 8 days post-infection in BALB/c mice infected with parental,

Δ*FLAM8* or rescue parasites (non-infected control mice were negative for

bioluminescence, C-).

Discussion

The differential localization of FLAM8 from the very distal tip in tsetse midgut procyclic parasites to the entire flagellum length in the mammalian-infectious stages (Calvo-Álvarez et al., submitted) prompted us to speculate that FLAM8 could play a specific role in the mammalian host. Here, we present for the first time a connection of a flagellar protein with the efficiency of trypanosomes to distribute in the host extravascular tissues. Quantitative analyses of experimental animal infections monitored by bioluminescence imaging showed that the absence of FLAM8 impaired parasite dissemination in the host extravascular compartment over the time of the infection, which was mostly recovered by the integration of a single rescue copy of the *FLAM8* gene in the endogenous locus.

1. FLAM8 and trypanosome transmission

Extravascular trypanosomes occupy the interstitial space of several organs, including the central nervous system, testes, adipose tissues and skin [4-6, 19-21]. The relevance of skin-dwelling trypanosomes in parasite transmission was demonstrated by xenodiagnosis experiments, early after the infective bite [22], or later in the infection [4], even in the absence of detectable parasitemia. More recently, the presence of extravascular trypanosomes was confirmed in the skin of confirmed and suspected cases of sleeping sickness [23]. Here, the impaired spreading of *FLAM8*-null parasites over the extravascular compartment, especially in the skin, would mathematically reduce the probability for parasites to be ingested by tsetse flies.

In the bloodstream, the balance between proliferative slender parasites and tsetse-adapted stumpy forms responds to a quorum sensing mechanism involving the production of oligopeptides and their reception through a specific transporter [24]. Although stumpy proportions were not evaluated over the entire course of the infection,

the absence of FLAM8 did not alter the ability of the parasites to differentiate into

transmissible stumpy forms at the peak of parasitemia, showing that the protein is most

probably not involved in this process. In addition, the ability of FLAM8-deprived stumpy

BSF to differentiate into procyclic *in vitro* suggests that stumpy cells present in the

blood and in the dermis could further develop upon ingestion by a tsetse fly.

2. On the possible cellular function(s) of FLAM8

Proliferative slender trypanosomes are highly mobile [25] and this motility was proved

to influence virulence *in vivo*. For instance, the lack of propulsive motility in flagellar

dynein *LC1* knockout mutants resulted in the inability of trypanosomes to establish an

infection in the bloodstream [14]. Here, *FLAM8* silencing in pleomorphic BSF

trypanosomes did not alter neither *in vitro* parasite growth nor cell motility in matrix-

dependent medium. However, quantitative analyses showed that *FLAM8*-null

trypanosomes were less numerous in extravascular tissues as compared to parental

controls. Assuming that parasite motility could be different in tissues and interstitial

spaces of extravascular niches with biophysical properties distinct from those in the

blood [25, 26], one cannot exclude that the motility of Δ*FLAM8* knockout parasites

might be somehow altered in the extravascular compartment. Intravital imaging for

motility analyses at the cell level would be needed in order to confirm this hypothesis.

Historically, the majority of studies on *T. brucei* virulence in experimental infections

have considered the blood circulation almost as the sole host compartment parasitized

by trypanosomes, whereas extravascular parasite niches and the underlying

exchanges between both compartments have been underestimated for long. The fact

that *FLAM8*-null trypanosomes were not able to disseminate properly over the

extravascular compartment could somehow imply defects in the way parasites sense

their microenvironment, resulting in an alteration of their extravascular tropism. For

example, FLAM8 could be involved in the parasite extravasation process and / or in

some subsequent host-parasite interactions promoting their maintenance and

development once they have reached the extravascular compartment.

To our knowledge, FLAM8 is the first flagellar component affecting parasite

dissemination in the extravascular host tissues *in vivo,* at least during the initial

dissemination step. Further investigations on the FLAM8 interactions with other

partners in the flagellum would help to unravel the roles of this fascinating and essential

organelle, especially regarding the modulation of trypanosome distribution in their

mammalian hosts and, therefore, parasite virulence.

Acknowledgements

We thank M. Bonhivers, D. Robinson, P. McKean, K. Matthews and K. Gull for providing various plasmids and antibodies. We acknowledge the IP BioImaging Plateforme for providing access to their equipment. We are grateful to P. Bastin for his strong scientific and human support. We warmly thank P. Bastin, M. Boshart and S. Bonnefoy for their critical reading of the manuscript.

References

1.   MacGregor P, Szoor B, Savill NJ, Matthews KR. Trypanosomal immune evasion, chronicity and transmission: an elegant balancing act. Nat Rev Microbiol. 2012;10(6):431-8. Epub 2012/05/01. doi: 10.1038/nrmicro2779. PubMed PMID: 22543519.

2.   Smith TK, Bringaud F, Nolan DP, Figueiredo LM. Metabolic reprogramming during the *Trypanosoma brucei* life cycle. F1000Res. 2017;6. Epub 2017/06/20. doi: 10.12688/f1000research.10342.2. PubMed PMID: 28620452; PubMed Central PMCID: PMCPMC5461901.

3.   Ooi CP, Bastin P. More than meets the eye: understanding *Trypanosoma brucei* morphology in the tsetse. Front Cell Infect Microbiol. 2013;3:71. Epub 2013/12/07. doi: 10.3389/fcimb.2013.00071. PubMed PMID: 24312899; PubMed Central PMCID: PMC3826061.

4.   Capewell P, Cren-Travaille C, Marchesi F, Johnston P, Clucas C, Benson RA, et al. The skin is a significant but overlooked anatomical reservoir for vector-borne African trypanosomes. Elife. 2016;5(September2016). Epub 2016/09/23. doi: 10.7554/eLife.17716. PubMed PMID: 27653219; PubMed Central PMCID: PMCPMC5065312.

5.   Caljon G, Van Reet N, De Trez C, Vermeersch M, Perez-Morga D, Van Den Abbeele J. The Dermis as a Delivery Site of *Trypanosoma brucei* for Tsetse Flies. PLoS Pathog. 2016;12(7):e1005744. doi: 10.1371/journal.ppat.1005744. PubMed PMID: 27441553; PubMed Central PMCID: PMCPMC4956260.

6.   Trindade S, Rijo-Ferreira F, Carvalho T, Pinto-Neves D, Guegan F, Aresta-Branco F, et al. *Trypanosoma brucei* Parasites Occupy and Functionally Adapt to the

Adipose Tissue in Mice. Cell Host Microbe. 2016. doi: 10.1016/j.chom.2016.05.002.

PubMed PMID: 27237364.

7.   Rotureau B, Subota I, Bastin P. Molecular bases of cytoskeleton plasticity during

the *Trypanosoma brucei* parasite cycle. Cellular microbiology. 2011;13(5):705-16.

Epub 2010/12/17. doi: 10.1111/j.1462-5822.2010.01566.x. PubMed PMID: 21159115.

8.   Broadhead R, Dawe HR, Farr H, Griffiths S, Hart SR, Portman N, et al. Flagellar

motility is required for the viability of the bloodstream trypanosome. Nature.

2006;440(7081):224-7. Epub 2006/03/10. doi: nature04541 [pii] 10.1038/nature04541.

PubMed PMID: 16525475.

9.   Kohl L, Robinson D, Bastin P. Novel roles for the flagellum in cell

morphogenesis and cytokinesis of trypanosomes. EMBO J. 2003;22(20):5336-46.

Epub 2003/10/09. doi: 10.1093/emboj/cdg518. PubMed PMID: 14532107.

10.   Tetley L, Vickerman K. Differentiation in *Trypanosoma brucei*: host-parasite cell

junctions and their persistence during acquisition of the variable antigen coat. J Cell

Sci. 1985;74:1-19. Epub 1985/03/01. PubMed PMID: 4030903.

11.   Langousis G, Hill KL. Motility and more: the flagellum of *Trypanosoma brucei*.

Nat Rev Microbiol. 2014;12(7):505-18. Epub 2014/06/17. doi: 10.1038/nrmicro3274.

PubMed PMID: 24931043.

12.   Rotureau B, Morales MA, Bastin P, Spath GF. The flagellum-MAP kinase

connection in Trypanosomatids: a key sensory role in parasite signaling and

development? Cellular microbiology. 2009;11(5):710-18. Epub 2009/02/12. doi:

CMI1295 [pii] 10.1111/j.1462-5822.2009.01295.x. PubMed PMID: 19207727.

13.   Roditi I, Schumann G, Naguleswaran A. Environmental sensing by African

trypanosomes. Curr Opin Microbiol. 2016;32:26-30. Epub 2016/05/01. doi:

10.1016/j.mib.2016.04.011. PubMed PMID: 27131101.

14.   Shimogawa MM, Ray SS, Kisalu N, Zhang Y, Geng Q, Ozcan A, et al. Parasite

motility is critical for virulence of African trypanosomes. Sci Rep. 2018;8(1):9122. Epub

2018/06/16. doi: 10.1038/s41598-018-27228-0. PubMed PMID: 29904094; PubMed

Central PMCID: PMCPMC6002391.

15.   Subota I, Julkowska D, Vincensini L, Reeg N, Buisson J, Blisnick T, et al.

Proteomic analysis of intact flagella of procyclic *Trypanosoma brucei* cells identifies

novel flagellar proteins with unique sub-localisation and dynamics. Mol Cell

Proteomics. 2014. Epub 2014/04/18. doi: 10.1074/mcp.M113.033357. PubMed PMID:

24741115.

16.   Fort C, Bonnefoy S, Kohl L, Bastin P. Intraflagellar transport is required for the

maintenance of the trypanosome flagellum composition but not its length. J Cell Sci.

2016;129(15):3026-41. Epub 2016/06/28. doi: 10.1242/jcs.188227. PubMed PMID:

27343245.

17.   Bertiaux E, Morga B, Blisnick T, Rotureau B, Bastin P. A Grow-and-Lock Model

for the Control of Flagellum Length in Trypanosomes. Curr Biol. 2018. Epub

2018/11/20. doi: 10.1016/j.cub.2018.10.031. PubMed PMID: 30449671.

18.   Calvo-Alvarez E, Cren-Travaillé C, Crouzols A, Rotureau B. A new chimeric

triple reporter fusion protein as a tool for in vitro and in vivo multimodal imaging to

monitor the development of African trypanosomes and *Leishmania* parasites. Infect

Genet Evol. 2018;63:391-403. Epub 2018/04/25. doi: 10.1016/j.meegid.2018.01.011.

PubMed PMID: 29339220.

19.   Goodwin LG. The pathology of African trypanosomiasis. Trans R Soc Trop Med

Hyg. 1970;64(6):797-817. Epub 1970/01/01. PubMed PMID: 5495630.

20.   Biteau N, Asencio C, Izotte J, Rousseau B, Fevre M, Pillay D, et al.

Trypanosoma brucei gambiense Infections in Mice Lead to Tropism to the

Reproductive Organs, and Horizontal and Vertical Transmission. PLoS Negl Trop Dis. 2016;10(1):e0004350. Epub 2016/01/07. doi: 10.1371/journal.pntd.0004350. PubMed PMID: 26735855; PubMed Central PMCID: PMCPMC4703293.

21. Carvalho T, Trindade S, Pimenta S, Santos AB, Rijo-Ferreira F, Figueiredo LM. *Trypanosoma brucei* triggers a marked immune response in male reproductive organs. PLoS Negl Trop Dis. 2018;12(8):e0006690. Epub 2018/08/16. doi: 10.1371/journal.pntd.0006690. PubMed PMID: 30110342; PubMed Central PMCID: PMCPMC6093638.

22. Caljon G, Van Den Abbeele J, Sternberg JM, Coosemans M, De Baetselier P, Magez S. Tsetse fly saliva biases the immune response to Th2 and induces anti-vector antibodies that are a useful tool for exposure assessment. Int J Parasitol. 2006;36(9):1025-35. Epub 2006/06/17. doi: 10.1016/j.ijpara.2006.05.002. PubMed PMID: 16777113.

23. Camara M, Soumah AM, Ilbouldo H, Travaille C, Clucas C, Cooper A, et al. Extravascular dermal trypanosomes in suspected and confirmed cases of gambiense Human African Trypanosomiasis. Clin Infect Dis. 2020. Epub 2020/07/09. doi: 10.1093/cid/ciaa897. PubMed PMID: 32638003.

24. Rojas F, Silvester E, Young J, Milne R, Tettey M, Houston DR, et al. Oligopeptide Signaling through TbGPR89 Drives Trypanosome Quorum Sensing. Cell. 2018. Epub 2018/12/07. doi: 10.1016/j.cell.2018.10.041. PubMed PMID: 30503212.

25. Bargul JL, Jung J, McOdimba FA, Omogo CO, Adung'a VO, Kruger T, et al. Species-Specific Adaptations of Trypanosome Morphology and Motility to the Mammalian Host. PLoS Pathog. 2016;12(2):e1005448. Epub 2016/02/13. doi: 10.1371/journal.ppat.1005448. PubMed PMID: 26871910; PubMed Central PMCID: PMCPMC4752354.

26. Sun SY, Kaelber JT, Chen M, Dong X, Nematbakhsh Y, Shi J, et al. Flagellum

couples cell shape to motility in *Trypanosoma brucei*. Proc Natl Acad Sci U S A.

2018;115(26):E5916-E25. Epub 2018/06/13. doi: 10.1073/pnas.1722618115. PubMed

PMID: 29891682; PubMed Central PMCID: PMCPMC6042131.

27. Le Ray D, Barry JD, Easton C, Vickerman K. First tsetse fly transmission of the

"AnTat" serodeme of *Trypanosoma brucei*. Ann Soc Belg Med Trop. 1977;57(4-5):369-

81. Epub 1977/01/01. PubMed PMID: 610616.

28. Bohringer S, Hecker H. Quantitative ultrastructural differences between strains

of the *Tryponasoma brucei* subgroup during transformation in blood. J Protozool.

1974;21(5):694-8. Epub 1974/11/01. doi: 10.1111/j.1550-7408.1974.tb03731.x.

PubMed PMID: 4449091.

29. Hirumi H, Hirumi K. Continuous cultivation of *Trypanosoma brucei* blood stream

forms in a medium containing a low concentration of serum protein without feeder cell

layers. J Parasitol. 1989;75(6):985-9. Epub 1989/12/01. PubMed PMID: 2614608.

30. Barquilla A, Saldivia M, Diaz R, Bart JM, Vidal I, Calvo E, et al. Third target of

rapamycin complex negatively regulates development of quiescence in *Trypanosoma*

brucei. Proc Natl Acad Sci U S A. 2012;109(36):14399-404. doi:

10.1073/pnas.1210465109. PubMed PMID: 22908264; PubMed Central PMCID:

PMCPMC3437835.

31. Czichos J, Nonnengaesser C, Overath P. *Trypanosoma brucei*: cis-aconitate

and temperature reduction as triggers of synchronous transformation of bloodstream

to procyclic trypomastigotes in vitro. Exp Parasitol. 1986;62(2):283-91. Epub

1986/10/01. doi: 10.1016/0014-4894(86)90033-0. PubMed PMID: 3743718.

32. Doyle JJ, Hirumi H, Hirumi K, Lupton EN, Cross GA. Antigenic variation in

clones of animal-infective *Trypanosoma brucei* derived and maintained *in vitro*.

Parasitology.   1980;80(2):359-69.   Epub   1980/04/01.   doi:
10.1017/s0031182000000810. PubMed PMID: 6102753.

33.   Wang Z, Morris JC, Drew ME, Englund PT. Inhibition of *Trypanosoma brucei*
gene expression by RNA interference using an integratable vector with opposing T7
promoters.   J   Biol   Chem.   2000;275(51):40174-9.   Epub   2000/10/03.   doi:
10.1074/jbc.M008405200. PubMed PMID: 11013266.

34.   Wirtz E, Leal S, Ochatt C, Cross GA. A tightly regulated inducible expression
system   for   conditional   gene   knock-outs   and   dominant-negative   genetics   in
Trypanosoma brucei. Mol Biochem Parasitol. 1999;99(1):89-101. Epub 1999/04/24.
PubMed PMID: 10215027.

35.   Redmond S, Vadivelu J, Field MC. RNAit: an automated web-based tool for the
selection   of   RNAi   targets   in   *Trypanosoma   brucei*.   Mol   Biochem   Parasitol.
2003;128(1):115-8. Epub 2003/04/23. doi: 10.1016/s0166-6851(03)00045-8. PubMed
PMID: 12706807.

36.   Burkard G, Fragoso CM, Roditi I. Highly efficient stable transformation of
bloodstream forms of *Trypanosoma brucei*. Mol Biochem Parasitol. 2007;153(2):220-
3.   Epub   2007/04/06.   doi: 10.1016/j.molbiopara.2007.02.008.   PubMed   PMID:
17408766.

37.   Kelly S, Reed J, Kramer S, Ellis L, Webb H, Sunter J, et al. Functional genomics
in *Trypanosoma brucei*: a collection of vectors for the expression of tagged proteins
from endogenous and ectopic gene loci. Mol Biochem Parasitol. 2007;154(1):103-9.
Epub   2007/05/22.   doi: S0166-6851(07)00099-0   [pii]
10.1016/j.molbiopara.2007.03.012. PubMed PMID: 17512617.

38.   Rotureau B, Ooi CP, Huet D, Perrot S, Bastin P. Forward motility is essential for

trypanosome infection in the tsetse fly. Cellular microbiology. 2014;16(3):425-33. Epub

2013/10/19. doi: 10.1111/cmi.12230. PubMed PMID: 24134537.

39.   Dacheux D, Landrein N, Thonnus M, Gilbert G, Sahin A, Wodrich H, et al. A

MAP6-related protein is present in protozoa and is involved in flagellum motility. PLoS

One. 2012;7(2):e31344. Epub 2012/02/23. doi: 10.1371/journal.pone.0031344.

PubMed PMID: 22355359; PubMed Central PMCID: PMCPMC3280300.

40.   Dean S, Marchetti R, Kirk K, Matthews KR. A surface transporter family conveys

the trypanosome differentiation signal. Nature. 2009;459(7244):213-7. Epub

2009/05/16. doi: nature07997 [pii] 10.1038/nature07997. PubMed PMID: 19444208.

Supporting information

Figure S1. Validation of the triple-reporter efficiency in monomorphic _FLAM8_[RNAi] **parasite lines.** Linear correlation between the number of parasites and the bioluminescence (in p/s) emitted by monomorphic _FLAM8_[RNAi] _FLAM8::mNG_ 427 BSF overexpressing the triple reporter chimeric protein [18] acquired by the IVIS Spectrum imager. Parasites without the RNAi plasmid (control), non-treated with tetracycline (non-induced) and treated with tetracycline (induced) are shown. Representative bioluminescent image of serial 1/2 dilutions performed in a 96-well plate (in photons / second / cm^2 / steradian). RNAi induction was triggered by the addition of 1 µg tetracycline and / or doxycycline for 72 h. Results represent the mean ± standard deviation (SD) of three independent experiments.

Figure S2. Molecular validation of the Δ_FLAM8_ null mutant cell lines. A) Whole-genome sequencing results showing _FLAM8_ wild-type allele (WT, upper panel), the loss of part of the 5' _FLAM8_ ORF in Δ_FLAM8_ knockout trypanosomes (middle panel) and the restoration of the full _FLAM8_ gene in rescue parasites (bottom panel), relative to the number of reads per 100-nt read length. The presence of the correct antibiotic cassettes is shown for knockout and rescue parasites (right middle and bottom panels). Δ_FLAM8_ knockout and rescue parasites are also bearing a construct for expression of a triple reporter (TR) as assessed by the detection of _bsd_ reads. **B)** Schemes showing the structure of the _FLAM8_ locus in wild-type parasites (upper scheme) and the integration plan of the different cassettes (lower _HYG-_, _PAC-_ and _BLE_-containing schemes). PCR confirmations of the successful integrations of all reporter cassettes (right panel). Primer pairs used for PCRs are indicated at the bottom of each line and

correspond to those drawn on the schemes. BSD: blasticidin; HYG: hygromycin; PAC:

puromycin; BLE: phleomycin.

Figure S3. Validation of the triple-reporter efficiency in pleomorphic Δ*FLAM8*

mutant cell lines. Linear correlation between the number of parasites and the

bioluminescence (in p/s) emitted by pleomorphic parental, Δ*FLAM8* knockout

subclones and rescue parasites overexpressing the triple reporter chimeric protein [18]

acquired by the IVIS Spectrum imager. Representative bioluminescent image of serial

1/2 dilutions performed in a 96-well plate (in photons / second / cm^2 / steradian).

Results represent the mean ± standard deviation (SD) of three independent

experiments.

Figure S4. The absence of *FLAM8* does not impact the parasite differentiation

into transmissible stumpy and insect procyclic forms *in vitro*. A) Representative

immunofluorescence pictures of methanol-fixed parental (left panel), Δ*FLAM8* null

subclones (middle panels) and rescue (right panel) stumpy trypanosomes, upon *in vitro*

treatment with a nucleotide 5'-AMP analog. Stumpy BSF were stained with anti-PAD1

antibody (red) and DAPI for DNA content (blue). Scale bars represent 5 μm. **B)**

Quantification of the proportion of stumpy trypanosomes in all pleomorphic cell lines

after *in vitro* differentiation in three independent experiments. No significant statistical

differences were found. The number of parasites considered for quantification (N) is

indicated above the graph. **C)** Representative pictures of methanol-fixed procyclic

parasites after *in vitro* differentiation from parental (left panel), Δ*FLAM8* null subclones

(middle panels) and rescue (right panel) procyclic parasites. Parasites were stained

with the anti-axonemal mAb25 (magenta) and anti-FLAM8 (green) antibodies, DAPI

staining for DNA content (blue). Scale bars represent 5 µm.

Figure S5. Comparison of intra- and extra-vascular trypanosome populations

during the experimental infection with the ΔFLAM8 null mutants. A-E) Total

numbers of parasites present in the bloodstream (IV, continuous line) and in the

extravascular tissues (EV, dotted line) of BALB/c mice infected with pleomorphic

parental (A), rescue (B) and three ΔFLAM8 null subclones (C, D, E) during the whole

course of the infection (27 days). Statistically significant differences were found in all

parasite strains during the entire infection period. The presence of parasites by

bioluminescence imaging was evident at day 1 post-infection, while blood parasites

were first detected at day 3, in agreement with previous observations [4].

Figure S6. Functional investigations on the ΔFLAM8 null mutants in dissected

internal organs. A) Ratios between the total photon fluxes quantified *ex vivo* in

dissected internal organs and the corresponding values of parasitemia at day 27 post

infection in BALB/c mice infected with parental, three ΔFLAM8 null subclones or rescue

pleomorphic BSF parasites. No statistical differences were observed. **B)** Normalized

representative pictures of the bioluminescent signals emitted by parasites in dissected

internal organs arranged in the same organization for quantification purposes.

Radiance intensity scale is provided in photons / second / cm^2 / steradian. a: liver; b:

spleen; c: kidneys; d: seminal vesicles; e: testes and epididymis; f: heart; g: lungs; h:

lymph nodes; i: stomach and intestine.

Figure 1

Figure 2

Figure 3

Figure 4

N=3 mice/group   Radiance (p/sec/cm²/sr, 10⁹)