Clr4[SUV39H1] and Bdf2[BRD4] ubiquitination mediate transcriptional silencing via heterochromatic phase transitions

Hyun-Soo Kim[1], Benjamin Roche[1], Sonali Bhattacharjee[1], Leila Todeschini[2], An-Yun Chang[1], Chris Hammell[1], André Verdel[2], Robert A. Martienssen[1,3,4*]

[1] Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 11724, USA.

[2] Institut for Advanced Biosciences, UMR InsermU1209/CNRS5309/UGA, University of Grenoble Alpes, Grenoble, France.

[3] Howard Hughes Medical Institute, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA

[4] Lead Contact

* Correspondence: martiens@cshl.edu

Abstract

Heterochromatin has crucial roles in genome stability by silencing repetitive DNA and by

suppressing recombination[1,2]. In the fission yeast *Schizosaccharomyces pombe*, Clr4

methylates histone H3 lysine 9 (H3K9) to maintain centromeric heterochromatin during

the cell cycle, guided by RNA interference (RNAi)[3,4]. Clr4 interacts with a specialized

ubiquitination complex CLRC, which is recruited in part by small RNA and the RNAi

transcriptional silencing complex (RITS) but otherwise, the role of CLRC is unknown[5-8].

Here we show that the E2 ubiquitin conjugating enzyme Ubc4 interacts with CLRC and

targets Clr4 for mono-ubiquitination at an intrinsically disordered region. Mono-

ubiquitination is essential for the transition from co-transcriptional (H3K9me2) to

transcriptional (H3K9me3) gene silencing by transiently releasing Clr4 from centromeric

heterochromatin, and is reversed by the deubiquitinating enzyme Ubp3. Addition of

ubiquitin changes the liquid-liquid phase separation (LLPS) of Clr4 to enhance its

dynamic activity. Ubc4-CLRC also poly-ubiquitinates Bdf2, the homologue of BET family

double bromodomain protein BRD4 for degradation, which otherwise accumulates at

centromeres to promote transcription and production of small RNAs.

Main text

Co-transcriptional gene silencing (CTGS) depends on the production of centromeric

sRNAs by RNA polymerase II and RNAi, which guides H3K9me2 restoration during S

phase of cell cycle. The subsequent transition from H3K9me2 to H3K9me3 and binding

of HP1 proteins Swi6 and Chp2 to heterochromatin, suppresses RNA Polymerase II-

dependent transcription by transcriptional gene silencing (TGS)[2,9-11]. However, the

mechanism that promotes the transition from RNAi-dependent CTGS to TGS is unclear.

Ubiquitination is a post-translational modification which requires ubiquitin, E1 ubiquitin

activating enzyme, E2 ubiquitin conjugating enzyme and E3 ubiquitin ligase[12]. We

previously showed that a specific mutation in the E2 ubiquitin conjugating enzyme Ubc4

abolishes heterochromatic silencing at centromeres, mating type loci and telomeres[13]

(Extended Data Fig 1b, d). Overexpression of Ubc4 also causes chromosome mis-

segregation[14]. We found that the *ubc4-G48D* (*ubc4-1*) mutation results in Ubc4 protein

instability (Extended Data Fig. 1c, e). Replacement of *ubc4* with the mouse homologue

UBE2D2 fully complemented *ubc4-1* and *UBE2D2-G48D* had similar silencing defects

(Extended Data Fig. 1b). A spontaneous second-site suppressor, S137T (see Material &

Methods), stabilizes Ubc4-1, restoring the silencing defects (Extended Data Fig. 1d-f).

Ubc4 and Cul4 regulate H3K9 methylation through RNAi

The E3 ubiquitin ligase, Cul4 is also involved in heterochromatin silencing and is part of

the CLRC complex[5-8]. *ubc4-1* and the *cul4-GFP* fusion (*cul4-1*) mutants have very

similar phenotypes, including loss of silencing of *ade6+* and *ura4+* reporter genes in

centromeric heterochromatin (Extended Data Fig. 1b), as well as increased binding of

Rpb1, the catalytic component of RNA polymerase II (Extended Data Fig. 2c).

Interestingly, while H3K9me3 was lost from both *ubc4-1* and *cul4-1* mutants at all

heterochromatic loci tested, H3K9me2 had increased occupancy at *dg* and *dh* repeats.

Increased H3K9me2 was also observed at the centromeric *ade6+* reporter gene, which

accumulates small RNA, but not at the *ura4+* gene, which does not (Fig. 1a and

Extended Data Fig. 2a, 2b, 3a). Swi6[HP1] protein binding was present at intermediate

levels, presumably because it recognizes both H3K9me2 and H3K9me3 (Extended

Data Fig. 2e)[9,15]. To determine whether RNAi is important for maintaining H3K9me2 in

ubc4-1 mutant cells, we introduced the *ubc4-1* mutation in *dcr1Δ* mutant cells and found

that the *dcr1Δ ubc4-1* double mutant had greatly reduced H3K9me2, and was more

sensitive to microtubule destabilizing agent thiabendazole (TBZ), showing that

centromeric heterochromatin was completely lost (Extended Data Fig. 3b-e). The more

severe alleles *ubc4-3xHA* and *cul4Δ* abolished H3K9me2 altogether (Extended Data

Fig. 4a-c). Live imaging revealed that Ubc4 and Cul4 proteins localize to nuclear

periphery (Extended Data Fig. 4d), and they form a stable complex by co-

immunoprecipitation (Co-IP) (Extended Data Fig. 4e). Ubc4 also forms a complex with

Raf1 and Raf2, two other components of CLRC (Extended Data Fig. 4f). These results

clearly indicate that Ubc4 and Cul4 form the E2/E3 complex by which CLRC targets

substrates important for heterochromatin assembly.

Ubc4-CLRC mono-ubiquitinates Clr4

Next, we set out to identify the substrates of this novel E2/E3 complex, Ubc4-CLRC.

Interestingly, Clr4 co-purifies with the CLRC complex *in vivo*, but does not behave like a

structural component of CLRC *in vitro*[5-8]. The Clr4 human homolog, SETDB1 is mono-

ubiquitinated at its SET-Insertion domain and this modification is important for its H3K9

methyltransferase activity[16]. To test whether Clr4 is a substrate of CLRC, we purified

3xFlag-Clr4 protein from *S. pombe* cells (Methods) and detected a single higher

molecular weight form that reacted with ubiquitin antibodies in a Ubc4-dependent

fashion (Fig. 1c, Extended Data Fig. 5a, b). A pulldown of ubiquitinated proteins using

Ubiquitination Affinity Beads (UBA) revealed a similar higher molecular weight Clr4

which is also dependent on Ubc4 (Extended Data Fig. 5c). In addition to Ubc4, the

presence of this higher molecular weight Clr4 depends on Cul4, Rik1, Raf1 and Raf2,

the other components of CLRC but not on the proteasome inhibitor MG-132 (Extended

Data Fig. 5d, e). These results strongly suggest that Ubc4 together with CLRC mono-

ubiquitinates Clr4 *in vivo*. As the abundance of Clr4-Ub protein is too low to detect

ubiquitinated residues by Mass Spectrometry, we ubiquitinated 6xHis-Clr4 recombinant

protein purified from *E. coli* cells *in vitro*. This approach uncovered many possible

ubiquitination sites, which mostly reside in an intrinsically disordered region (IDR)

between the chromodomain (CD) and PreSET domain (Extended Data Fig. 5g-i) which

is involved in nucleosome binding[17]. To identify which of these were functional *in vivo*,

36 lysines of Clr4 were arranged into 6 subgroups, and mutated into arginines (R). We

also included a mutant of all 36 lysines (*AllKtoR*). The all-lysine mutant (*AllKtoR*) and

the second block mutant (*2-8KtoR*), corresponding to the middle of the IDR, completely

abolished ubiquitination (Fig. 1e and Extended Data Fig. 6b) consistent with results of

Mass Spectrometry. Although no single lysine to arginine mutant lost ubiquitination

(Extended Data Fig. 6c) we could narrow down the possible sites to 4 lysines (*clr4-2-*

4KtoR; K109, K110, K113, and K114) (Fig. 1e and Extended Data Fig. 6d-f). ChIP-Seq

revealed that *clr4-AllKtoR* lost both H3K9me2 and H3K9me3, resembling strong alleles

of *ubc4* and *cul4*, while *clr4-2-4KtoR* had the same increase of H3K9me2 and decrease

of H3K9me3 seen in *ubc4-1* and *cul4-1* hypomorphs (Fig. 1g and Extended Data Fig.

6g). Thus mono-ubiquitination of Clr4 by Ubc4-CLRC is required for H3K9 tri-

methylation at centromeric heterochromatin.

Clr4 ubiquitination induces its dissociation from chromatin

Clr4 ubiquitination does not change the histone methylation activity of Clr4 (Extended

Data Fig. 7a) but binding to chromatin was increased in *ubc4-1*, *cul4-1*, *clr4-2-8KtoR* or

clr4-AllKtoR mutants, especially at *dh* where sRNAs are produced (Fig. 2a and

Extended Data Fig. 7b). To further explore the effect of Clr4 mono-ubiquitination on

chromatin binding, we carried out nucleosome binding assays using *in vitro*

ubiquitinated 6xHis-Clr4 recombinant protein or 3xFlag-Clr4 purified from *S. pombe*

cells with tethered nucleosomes (Extended Data Fig. 7c). In contrast to unmodified Clr4

protein which had strong binding affinity to the nucleosome, especially when H3K9me3

was present, ubiquitinated Clr4 did not bind to the nucleosome (Fig. 2b Extended Data

Fig. 7d). These results are consistent with the increased binding *in vivo* in the absence

of mono-ubiquitination in *ubc4-1* or *cul4-1*. Binding in the absence of ubiquitin depended

on the Clr4 chromodomain (Fig. 2c and Extended Data Fig. 7e)[9,18]. These results

indicate that Clr4 mono-ubiquitination reduces the binding activity of Clr4 to chromatin

and that this is important for the transition from H3K9 di- to tri-methylation.

Ubp3 deubiquitinates Clr4

Ubiquitination is a reversible, highly dynamic and transient modification, and its removal

is catalyzed by deubiquitinating enzymes[12]. A deletion mutant of deubiquitinating

enzyme Ubp3 rescues heterochromatic silencing defects of *ago1Δ,* and *ubp3Δ* shows

positive epistatic genetic interactions with CLRC component mutants[19,20]. While deletion

of Ubp3 did not increase Clr4 mono-ubiquitination (Extended Data Fig. 8a),

overexpression had a silencing defect at the centromeric *ura4*[+] reporter gene and

reduced the mono-ubiquitination of Clr4 (Fig. 1f and Extended Data Fig. 8b). The

overexpression of Ubp3 also resulted in an increase of H3K9me2 and decrease of

H3K9me3 as well as increased RNA Pol II binding, noncoding RNA transcription and

Clr4 binding at centromeric heterochromatin (Fig. 1g and Extended Data Fig. 8c-f).

These results indicate that Ubp3 counteracts Clr4 mono-ubiquitination activity of Ubc4-

CLRC and reverses Clr4 activity.

Clr4^{IDR} displays ubiquitin-dependent phase separation properties.

Swi6 forms 4 foci in the nucleus corresponding to the chromocenter and the three

telomeric clusters, all of which depend on Clr4 (Fig. 3a and Extended Data Fig. 9a). In

ubc4-1 mutants, the telomeric foci disappear and the chromocenter is retained, while in

dcr1Δ mutants the telomeric foci are retained and the chromocenter is reduced (Fig. 1b

and 3a and Extended Data Fig. 3c, 3d and 9a). Along with HP1 proteins from human,

Swi6 has been shown to undergo liquid-liquid phase separation (LLPS) which

contributes to the highly dynamic features of heterochromatin[21,22]. Clr4 protein has an

intrinsically disordered region (IDR) or low-complexity domain (LCD) between its

chromodomain (CD) and PreSET domain[17], and IDR/LCD are domains typically

responsible for phase separation (Extended Data Fig. 9b)[23]. We found that recombinant

GFP-Clr4^{Full} and GFP-Clr4^{IDR} (51-220 amino acids) proteins form liquid droplets *in vitro*

at low salt conditions even without addition of crowding agents like PEG8000 (Fig. 3b

and Extended Data Fig. 9c, d). Next, we treated the proteins with 1,6-Hexanediol which

dissolves liquid droplets by disrupting hydrophobic interaction of protein-protein or

protein-RNA[24]. We found that the formation of liquid droplets of GFP-Clr4^{IDR} was greatly

reduced by 1,6-haxanediol treatment indicating that these liquid droplets are formed by

LLPS (Extended Data Fig. 9d). Clr4^{IDR} binds ubiquitin[25] and we added concentrated

ubiquitin to GFP-Clr4^{IDR} in order to mimic the effect of ubiquitination of Clr4. Strikingly,

we found that adding ubiquitin greatly reduced the size of LLPS of GFP-Clr4^{IDR} (Fig. 3b,

c). Furthermore, fluorescence recovery after photobleaching (FRAP) of the liquid

droplets showed a faster recovery rate of GFP-Clr4^{IDR} with ubiquitin than GFP-Clr4^{IDR}

alone (Fig. 3d, Extended Data Fig. 9e) indicating that increased dynamics of GFP-

Clr4^{IDR} by addition of ubiquitin confer reduced size of liquid droplets and escape from

the LLPS.

Ubc4 and Cul4 regulate transcription of sRNA through Bdf2^{BRD4}

In *ubc4-1* and *cul4-1* mutants, we observed an almost 2-fold increase of centromeric

sRNA, as previously observed for *cul4* (Fig. 4a, Extended Data Fig. 10a)[18],

accompanied by increased chromatin binding of Ago1 and Chp1, two other components

of the RNA-induced transcriptional gene silencing (RITS) complex (Extended Data Fig.

10c). The putative H3K9 demethylase, Epe1, localizes to centromeric heterochromatin

in a Swi6-dependent manner and contributes to sRNA production[26,27]. The BET family

double bromodomain protein Bdf2 binds to the centromeric heterochromatin boundary

in Epe1-dependent manner and interacts with the TFIID transcription factor

complex[28,29]. *bdf2Δ* suppressed the increased sRNA production in *ubc4-1* and *cul4-1*

mutant cells (Fig. 4a). The chromatin binding of Bdf2 at centromeric heterochromatin

was greatly increased in both *ubc4-1* and *cul4-1* (Fig. 4b) and at euchromatic genes in

ubc4-1 (Extended Data Figure 10d, e), while the protein level was increased in *ubc4-1*

only (Fig. 4c). These results suggest that Ubc4 is a general regulator for Bdf2, while

Cul4 regulates Bdf2 specifically at centromeric heterochromatin. Bdf2 protein purified

from cells treated with proteasome inhibitor MG-132 accumulated multiple ubiquitinated

bands dependent on Ubc4, suggesting that Ubc4 poly-ubiquitinates Bdf2 for its

degradation (Fig. 4d). These results indicate that Ubc4 and Cul4 regulates the protein

level and chromatin binding of Bdf2 by poly-ubiquitination, which is important for sRNA

production by recruiting TFIID and RNA Pol II to centromeric heterochromatin.

Discussion

We demonstrate that Ubc4-CLRC is essential for the switch between RNAi-dependent

CTGS and RNAi-independent TGS (Fig. 4e). During S phase, centromeric

heterochromatin is transcribed, and CLRC is recruited to centromeric heterochromatin

by the RITS complex and small RNA[10,11,30]. Clr4 di-methylates newly deposited H3K9,

and has strong binding affinity to existing H3K9me3. Subsequent mono-ubiquitination of

Clr4 by Ubc4-CLRC releases it from H3K9me3, allowing tri-methylation of adjacent

H3K9me1 or -me2 nucleosomes independent of RNAi[31]. Mono-ubiquitination also

triggers dissolution of subcellular phase separated droplets, allowing replacement by

Swi6 condensates that also bind H3K9me3 and silence transcription[22]. Poly-

ubiquitination of Bdf2 further reduces transcription and small RNA production

completing the transition to TGS. De-ubiquitination by Ubp3 reverses the process and

allows Clr4 recruitment in the next cell cycle (Fig. 4e).

In *ubc4-1* and *cul4-1* mutants, or in mutants of Clr4 that can no longer be ubiquitinated,

CTGS cannot transition to TGS, leading to increased sRNA production, and failure to

release Clr4. In contrast, strong *ubc4-3xHA* and *cul4Δ* mutants completely abolish

centromeric heterochromatin resembling *clr4Δ*, *rik1Δ*, *raf1Δ* and *raf2Δ* (Extended Data

Fig. 4a-c), because Clr4 is no longer recruited to heterochromatin by CLRC and RNAi in

the first place (Extended Data Fig. 5f). Ubc4-CLRC may possess additional targets,

such as H3K14, which was recently shown to be a prerequisite for H3K9 methylation *in*

vitro[25]. Bdf2, like its human homologue BRD4, has IDRs which may induce LLPS of

acetylated chromatin (Extended Data Fig. 9f)[32]. Poly-ubiquitination of Bdf2 may similarly

release Bdf2 from the heterochromatin phase for subsequent degradation[33]. The

ubiquitin-related protein modifier SUMO contributes to small RNA-dependent

localization of the Clr4 homologue SETDB1 to chromatin in Drosophila[34], and is

required for heterochromatic silencing in *S. pombe*[35], which we speculate may also be

mediated by a similar mechanism as reported here[36]. The existence of IDR/LCD

between chromo and PreSET domains of mammalian SUV39H1 (Extended Data Fig.

9g) and the interaction of human BRD4 with CUL4B[37] suggest that parallel mechanisms

might regulate transcriptional silencing in higher eukaryotes.

References

Janssen, A., Colmenares, S. U. & Karpen, G. H. Heterochromatin: Guardian of the Genome. *Annu Rev Cell Dev Biol* **34**, 265-288, doi:10.1146/annurev-cellbio-100617-062653 (2018).

Castel, S. E. & Martienssen, R. A. RNA interference in the nucleus: roles for small RNAs in transcription, epigenetics and beyond. *Nat Rev Genet* **14**, 100-112, doi:10.1038/nrg3355 (2013).

Volpe, T. A. *et al.* Regulation of heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. *Science* **297**, 1833-1837, doi:10.1126/science.1074973 (2002).

Verdel, A. *et al.* RNAi-mediated targeting of heterochromatin by the RITS complex. *Science* **303**, 672-676, doi:10.1126/science.1093686 (2004).

Horn, P. J., Bastie, J. N. & Peterson, C. L. A Rik1-associated, cullin-dependent E3 ubiquitin ligase is essential for heterochromatin formation. *Genes Dev* **19**, 1705-1714, doi:10.1101/gad.1328005 (2005).

Hong, E. J., Villén, J., Gerace, E. L., Gygi, S. P. & Moazed, D. A cullin E3 ubiquitin ligase complex associates with Rik1 and the Clr4 histone H3-K9 methyltransferase and is required for RNAi-mediated heterochromatin formation. *RNA Biol* **2**, 106-111, doi:10.4161/rna.2.3.2131 (2005).

Jia, S., Kobayashi, R. & Grewal, S. I. Ubiquitin ligase component Cul4 associates with Clr4 histone methyltransferase to assemble heterochromatin. *Nat Cell Biol* **7**, 1007-1013, doi:10.1038/ncb1300 (2005).

215   8   Kuscu, C. *et al.* CRL4-like Clr4 complex in Schizosaccharomyces pombe

216   depends on an exposed surface of Dos1 for heterochromatin silencing. *Proc Natl*

217   *Acad Sci U S A* **111**, 1795-1800, doi:10.1073/pnas.1313096111 (2014).

218   9   Jih, G. *et al.* Unique roles for histone H3K9me states in RNAi and heritable

219   silencing of transcription. *Nature* **547**, 463-467, doi:10.1038/nature23267 (2017).

220   10   Kloc, A., Zaratiegui, M., Nora, E. & Martienssen, R. RNA interference guides

221   histone modification during the S phase of chromosomal replication. *Curr Biol* **18**,

222   490-495, doi:10.1016/j.cub.2008.03.016 (2008).

223   11   Chen, E. S. *et al.* Cell cycle control of centromeric repeat transcription and

224   heterochromatin assembly. *Nature* **451**, 734-737, doi:10.1038/nature06561

225   (2008).

226   12   Hochstrasser, M. Origin and function of ubiquitin-like proteins. *Nature* **458**, 422-

227   429, doi:10.1038/nature07958 (2009).

228   13   Irvine, D. V. *et al.* Mapping epigenetic mutations in fission yeast using whole-

229   genome next-generation sequencing. *Genome Res* **19**, 1077-1083,

230   doi:10.1101/gr.089318.108 (2009).

231   14   Javerzat, J. P., Cranston, G. & Allshire, R. C. Fission yeast genes which disrupt

232   mitotic chromosome segregation when overexpressed. *Nucleic Acids Res* **24**,

233   4676-4683, doi:10.1093/nar/24.23.4676 (1996).

234   15   Schalch, T. *et al.* High-affinity binding of Chp1 chromodomain to K9 methylated

235   histone H3 is required to establish centromeric heterochromatin. *Mol Cell* **34**, 36-

236   46, doi:10.1016/j.molcel.2009.02.024 (2009).

16   Sun, L. & Fang, J. E3-Independent Constitutive Monoubiquitination

Complements Histone Methyltransferase Activity of SETDB1. *Mol Cell* **62**, 958-

966, doi:10.1016/j.molcel.2016.04.022 (2016).

17   Akoury, E. *et al.* Disordered region of H3K9 methyltransferase Clr4 binds the

nucleosome and contributes to its activity. *Nucleic Acids Res* **47**, 6726-6736,

doi:10.1093/nar/gkz480 (2019).

18   Zhang, K., Mosch, K., Fischle, W. & Grewal, S. I. Roles of the Clr4

methyltransferase complex in nucleation, spreading and maintenance of

heterochromatin. *Nat Struct Mol Biol* **15**, 381-388, doi:10.1038/nsmb.1406

(2008).

19   Reyes-Turcu, F. E., Zhang, K., Zofall, M., Chen, E. & Grewal, S. I. Defects in

RNA quality control factors reveal RNAi-independent nucleation of

heterochromatin. *Nat Struct Mol Biol* **18**, 1132-1138, doi:10.1038/nsmb.2122

(2011).

20   Ryan, C. J. *et al.* Hierarchical modularity and the evolution of genetic

interactomes across species. *Mol Cell* **46**, 691-704,

doi:10.1016/j.molcel.2012.05.028 (2012).

21   Larson, A. G. *et al.* Liquid droplet formation by HP1α suggests a role for phase

separation in heterochromatin. *Nature* **547**, 236-240, doi:10.1038/nature22822

(2017).

22   Sanulli, S. *et al.* HP1 reshapes nucleosome core to promote phase separation of

heterochromatin. *Nature* **575**, 390-394, doi:10.1038/s41586-019-1669-2 (2019).

23   Kato, M. *et al.* Cell-free formation of RNA granules: low complexity sequence

domains form dynamic fibers within hydrogels. *Cell* **149**, 753-767,

doi:10.1016/j.cell.2012.04.017 (2012).

24   Kroschwald, S. *et al.* Promiscuous interactions and protein disaggregases

determine the material state of stress-inducible RNP granules. *Elife* **4**, e06807,

doi:10.7554/eLife.06807 (2015).

25   Oya, E. *et al.* H3K14 ubiquitylation promotes H3K9 methylation for

heterochromatin assembly. *EMBO Rep* **20**, e48111,

doi:10.15252/embr.201948111 (2019).

26   Zofall, M. & Grewal, S. I. Swi6/HP1 recruits a JmjC domain protein to facilitate

transcription of heterochromatic repeats. *Mol Cell* **22**, 681-692,

doi:10.1016/j.molcel.2006.05.010 (2006).

27   Trewick, S. C., Minc, E., Antonelli, R., Urano, T. & Allshire, R. C. The JmjC

domain protein Epe1 prevents unregulated assembly and disassembly of

heterochromatin. *Embo j* **26**, 4670-4682, doi:10.1038/sj.emboj.7601892 (2007).

28   Wang, J. *et al.* Epe1 recruits BET family bromodomain protein Bdf2 to establish

heterochromatin boundaries. *Genes Dev* **27**, 1886-1902,

doi:10.1101/gad.221010.113 (2013).

29   Kim, H. S. *et al.* Identification of a BET family bromodomain/casein kinase II/TAF-

containing complex as a regulator of mitotic condensin function. *Cell Rep* **6**, 892-

905, doi:10.1016/j.celrep.2014.01.029 (2014).

30   Bayne, E. H. *et al.* Stc1: a critical link between RNAi and chromatin modification

required for heterochromatin integrity. *Cell* **140**, 666-677,

doi:10.1016/j.cell.2010.01.038 (2010).

31   Al-Sady, B., Madhani, H. D. & Narlikar, G. J. Division of labor between the

chromodomains of HP1 and Suv39 methylase enables coordination of

heterochromatin spread. *Mol Cell* **51**, 80-91, doi:10.1016/j.molcel.2013.06.013

(2013).

32   Gibson, B. A. *et al.* Organization of Chromatin by Intrinsic and Regulated Phase

Separation. *Cell* **179**, 470-484.e421, doi:10.1016/j.cell.2019.08.037 (2019).

33   Yasuda, S. *et al.* Stress- and ubiquitylation-dependent phase separation of the

proteasome. *Nature* **578**, 296-300, doi:10.1038/s41586-020-1982-9 (2020).

34   Ninova, M. *et al.* Su(var)2-10 and the SUMO Pathway Link piRNA-Guided Target

Recognition to Chromatin Silencing. *Mol Cell* **77**, 556-570.e556,

doi:10.1016/j.molcel.2019.11.012 (2020).

35   Xhemalce, B., Seeler, J. S., Thon, G., Dejean, A. & Arcangioli, B. Role of the

fission yeast SUMO E3 ligase Pli1p in centromere and telomere maintenance.

Embo j **23**, 3844-3853, doi:10.1038/sj.emboj.7600394 (2004).

36   Shin, J. A. *et al.* SUMO modification is involved in the maintenance of

heterochromatin stability in fission yeast. *Mol Cell* **19**, 817-828,

doi:10.1016/j.molcel.2005.08.021 (2005).

37   Liu, H. *et al.* Inflammation-dependent overexpression of c-Myc enhances

CRL4(DCAF4) E3 ligase activity and promotes ubiquitination of ST7 in colitis-

associated cancer. *J Pathol* **248**, 464-475, doi:10.1002/path.5273 (2019).

Materials and Methods

Yeast strains and plasmids manipulation

Yeast strains, oligonucleotide sequences and plasmids are listed in Supplementary

Information Table S1-3. All yeast strains were constructed by PCR-based targeting

method[38] and lithium acetate transformation method[39]. For N-terminus tagging of 3xFlag

to *clr4*, *rik1* and *raf1*, endogenous promoters and terminators of each gene were used.

For some 3xFlag-Clr4 lysine mutants, *adh1* terminator with Kan resistant (Kan^R) gene

was inserted at the end of *clr4* open reading frames (ORF). For overexpression of *ubp3*

and *ubp8*, 500 bp of *adh1* promoter was inserted between their promoter and ORF. To

construct pET-14b-Clr4, the coding DNA sequence of full-length Clr4 (amino acids 1-

490) was amplified from *S. pombe* genomic DNA, digested with *NdeI* and cloned into

NdeI-digested pET-14b. For pHis-parallel-GFP-Clr4^{Full} or pHis-parallel-GFP-Clr4^{IDR}

construction, the coding DNA sequence of full-length Clr4^{Full} (amino acids 1-490) or

Clr4^{IDR} (amino acids 51-220) were amplified from *S. pombe* genomic DNA, digested

with *BamHI*/*XhoI* and cloned into *BamHI*/*XhoI*-digested pHis-parallel-GFP.

Chromatin Immunoprecipitation (ChIP)

Chromatin Immunoprecipitation (ChIP) experiments were performed as previously

described[29]. Briefly, freshly grown 40 ml cells in YES were fixed with 1% formaldehyde

from 37% stock (Sigma F8775) for 20 min at room temperature for H3K9me2,

H3K9me3, RNA Pol II, Ago1 and Chp1 ChIP. For Swi6 and 3xFlag-Clr4 ChIP, 40 ml

cells were pre-incubated at 18 °C for 2 h and fixed with 1.45% Formaldehyde from 16%

stock (ThermoFisher 28908) for 30 min at 18 °C. Cells were quenched with 360 mM

glycine and 2.4 mM Tris for 5 min. Whole-cell extracts (WCE) were prepared using

FastPrep-24 and sonicated using Covaris. 1~2 μg of antibodies like anti-H3K9me2

(Abcam ab1220), anti-H3K9me3 (Absolute Antibody Ab00700-1.26), anti-RNA

polymerase II (BioLegend 904001), Swi6 (Abcam ab188276), Ago1 (Abcam Ab18190),

Chp1 (Abcam ab18191) and anti-Flag (Sigma F1804) were pre-incubated with 50 μl

Protein A and Protein G sepharose (GE 17513801 and 17061801) mix for 3~4 h and

then incubated with sheared chromatin overnight at 4 °C. After washing, bead-antibody-

chromatin mixture were treated with Proteinase K (ThermoFisher 100005393) for 1 h

ant 42 °C and incubated 5~6 h at 65 °C to reverse cross-links. DNA was cleaned up

using ChIP DNA Clean & Concentrator (Zymo Research D5205) and subsequently used

for qPCR or ChIP-Seq.

ChIP-Seq

ChIP-Seq libraries were constructed using NEBNext Ultra II DNA Library Prep Kit for

Illumina (NEB E7645) and barcoded multiplex libraries were sequenced using Illumina

MiSeq platform, which generates paired-end 101 bp reads. Raw reads were trimmed

using Trimmomatic, mapped to *S. pombe* genome using Bowtie2 and visualized in IGV

genome browser. Total ChIP-seq read counts for defined region were analyzed using

MultiBigwigSummary or MultiBamSummary.

Co-Immunoprecipitation (Co-IP) and Immunoprecipitation (IP) for ubiquitinated

Clr4

Whole-cell extracts (WCE) were prepared from freshly grown yeast cells in lysis buffer

(40mM Tris-HCl, pH 7.5, 1 mM EDTA, 10% Glycerol, 0.1% NP-40, 300 mM NaCl) with

Protease inhibitor cocktail (Roche 11836170001). For Co-IP of Cul4-GFP with Ubc4, 2

µg of GFP antibody (Roche 11814460001) was pre-incubated with 50 µl Protein A and

Protein G Sepharose (GE Healthcare 17513801 and 17061801) mix for 3~4 h and then

incubated with WCE overnight at 4 °C. After washing, purified proteins were loaded on

4–20% Mini-PROTEAN TGX Gel (Bio-rad 456-1093) and Western blotting was

performed with anti-GFP (Abcam 290) and anti-mUBE2D3 (Proteintech 11677-1-AP)

antibodies to detect Cul4-GFP and Ubc4 using ECL (Thermo Fisher 32209) assay

according to the manufacturer's instruction. For Co-IP of Flag-Rik1, Flag-Dos1 or Dos2-

Flag with Ubc4, 50 µl of Anti-Flag M2 Affinity Gel (Sigma A2220) was incubated with

WCE overnight at 4 °C. For detection of Flag tagged protein and Ubc4, anti-Flag (Sigma

F7425) and anti-mUBE2D3 (Proteintech 11677-1-AP) antibodies were used. For

Immunoprecipitation of ubiquitinated Clr4, WCE prepared from freshly grown cells in

lysis buffer (50mM Hepes, pH 7.6, 150mM NaCl, 300mM KAc, 5mM MgAc, 10%

Glycerol, 20mM β-GP, 1mM EDTA, 1mM EGTA, 0.1% NP-40, 1mM DTT, 250U/ml

Dnase I (Roche 04716728001), 25~50 µg/ml Rnase A (Sigma R4642), 1mM PMSF,

0.3mM PR-619 (Sigma 662141)) with Protease inhibitor cocktail (Roche 11836170001)

and De-ubiquitination/sumoylation Inhibitor (Cytoskeleton NEM09BB) was incubated

with Anti-Flag M2 Affinity Gel or Ubiquitination Affinity Beads (UAB, Cytoskeleton

UBA01- Beads) overnight at 4 °C. For detection of ubiquitinated Clr4 protein, anti-Flag

(Sigma F7425) or anti-ubiquitin (Cell Signalling 3936T) antibodies were used. Cells

were also treated with proteasome inhibitor 100 µM MG-132 (APExBIO A2585) for 4 h

before harvesting.

Quantitative reverse transcription PCR (RT-qPCR)

Total RNAs were purified from freshly grown yeast cells using *Quick*-RNA

Fungal/Bacterial Miniprep Kit (Zymoresearch R2014). Genomic DNAs were removed

using DNase I (Roche 04716728001) and RNAs were cleaned using RNA Clean &

Concentrator (Zymoresearch R1013). Reverse transcription was carried out using

SuperScript IV First-Strand Synthesis System (ThermoFisher 18091050) with random

hexamer or oligo dT as primers. For quantitative PCR (qPCR), iQ SYBR Green

Supermix (Bio-rad 1708882) was used with the primers listed in Supplementary

Information Table 2 and samples were run on CFX96 Real-Time PCR Detection System

(Bio-rad).

sRNA-Seq

Total RNAs were purified from freshly grown yeast cells using *Quick*-RNA

Fungal/Bacterial Miniprep Kit (Zymoresearch R2014) and sRNA were enriched using

RNA Clean & Concentrator (Zymoresearch R1013). sRNA libraries were constructed

using NEBNext Small RNA Library Prep Set for Illumina (NEB E7330) and barcoded

multiplex libraries were sequenced using Illumina MiSeq platform, which generates

single-end 36 nt reads. Reads were adapter-trimmed using FASTX-Toolkit, keeping a

minimum read size of 15 nt, and mapped to the *S. pombe* genome using Bowtie[40]

allowing up to 1 mismatch. The strand-specific genomic coverage was calculated using

Bedtools[41] and represented in 20 nt windows.

Silencing spot assay

Cells grown freshly on YES plate were picked and resuspended in water to make a concentration of ~5 × 10^4 cells per µl. 8 µl of 10-fold serial dilutions were spotted on appropriate plates like YES, YE, EMM-Ura, YES+FOA, YES+TBZ, and incubated 3 days at 30 °C for photography. *ura4*+ reporter gene silencing was also tested on 5-fluoro-orotic acid (FOA) plates, which kills cells expressing Ura4.

Iodine staining

Freshly grown yeast cells were spotted on SPA plates to induce sporulation, incubated at room temperature for 3 days, and then stained with iodine vapor for 2-3 min. Staining of starch which is abundant in mature spore produces a dark brown color.

EMS mutagenesis and whole-genome sequencing for *ubc4-G48D suppressor*

EMS mutagenesis was carried out as previously described[42]. Briefly, freshly grown 1 x 10^9 yeast cells were washed with distilled water, resuspended in 1.5 ml of 0.1 M sodium phosphate buffer (pH 7.0) and treated with 50 µl EMS (Sigma M0880) for 1h at 30 °C. 0.2 ml of treated cells was moved to 8 ml of sterile 5% sodium thiosulfate to inactivate EMS. Cells were washed, resuspended in sterile water and plated on YES plate. Single colonies from plates were patched on SPA plates to induce sporulation, incubated at room temperature for 3 days, and then stained with iodine vapor for 2-3 min. For whole-genome sequencing for *ubc4-G48D* suppressors, genomic DNA was purified from freshly grown yeast cells using Genomic-tip 20/G column (QIAGEN 10223). DNA library was constructed using the TruSeq DNA PCR-Free Library Prep Kit (Illumina 20015962).

Barcoded DNA library was sequenced using Illumina MiSeq platform, which generates

paired-end 151 nt reads and analyzed as previously described[43]. Briefly, reads were

adapter-trimmed and quality-filtered using Sickle (paired-end mode), then mapped to

the *S. pombe* genome using Bowtie2[44]. Duplicate reads were removed using Samtools

(≤ 0.14% per library). SNPs were called using FreeBayes to find SNPs present in

suppressor strain but absent in the parental strain[45]. This uncovered a G→C

transversion at position chr2:716192, a second-site mutation in the *ubc4* gene. The

mutation was verified by Sanger sequencing and confirmed by constructing fresh same

mutation.

Live cell imaging

Images of freshly growing cells were collected using Zeiss Axio Imager M2 with DIC and

EGFP channels and processed using Zeiss Zen software.

CLRC complex purification from *S. pombe*

Protein purifications were performed as described previously[4]. Briefly, Rik1-TAP were

purified from 6 to 12 g of cells. Cells were lysed in 1 volume of lysis buffer (50 mM

HEPES-KOH, pH 7.6, 300 mM KAc, 10% glycerol, 1 mM EGTA, 1 mM EDTA, 0.1% NP-

40, 1 mM DTT, 5 mM MgAc, 1 mM NaF, 20 mM β-GP, 1 mM PMSF, 1 mM Benzamidine

and 1 µg/ml of Leupeptin, Aprotinin, Bestatin and Pepstatin). 1 volume lysis buffer was

added and the lysate was centrifuged at 16,500 rpm for 25 min at 4 °C in a SA600 rotor.

The first affinity purification was performed using Ig-G Sepharose beads at 4 °C for 2 h.

The beads were washed with lysis buffer and TEV-C buffer (10 mM Tris-HCl, pH 8.0,

150 mM KAc, 0.1% NP-40, 0.5 mM EDTA, 1 mM DTT) and bound protein was eluted

using a GST-TEV protease. The eluate was diluted in 2 volumes of CAM-B buffer (10

mM Tris-HCL, pH 8.0, 150 mM NaCl, 1 mM MgAc, 1 mM Imidazole, 2 mM $CaCl_2$, 10

mM βME) and the second affinity purification was performed by adding Calmodulin-

Sepharose at 4 °C for 1 h. The beads lysate was washed with CAM-B buffer containing

0.1 % NP-40 and bound protein was eluted with CAM-E buffer (10 mM Tris-HCL, pH

8.0, 150 mM NaCl, 1 mM MgAc, 1 mM Imidazole, 10 mM EGTA, 10 mM βME).

Recombinant protein purification from *E. coli*, *in vitro* ubiquitination, Histone

Methylation assay and nucleosome binding assay

BL21-CodonPlus (DE3)-RIL strains (Agilent Technologies 230245) that express 6xHis-

Clr4, 6xHis-GFP-Clr4[Full] or 6xHis-GFP-Clr4[IDR] were grown with 0.4 mM IPTG for 20 h at

18 °C. Induced proteins were purified on HisPur Ni-NTA Superflow Agarose

(ThermoFisher 25214) and eluted with elution buffer containing 150 mM Imidazole. For

in vitro ubiquitination, purified 6xHis-Clr4 protein was mixed with human Ubiquitin-

activating Enzyme (UBE1, R&D systems E-305), human UbcH5c/UBE2D3 (R&D

systems E2-627), human Ubiquitin (R&D systems U-100H) or human HA-Ubiquitin

(R&D systems U-110), Mg-ATP (R&D systems B-20) and 10x E3 Ligase Reaction

Buffer (R&D systems B-71). Addition of Rik1-TAP complex purified from *S. pombe* also

added to the reaction gave similar results. For Histone Methylation assay, ubiquitinated

or unmodified Clr4 proteins were mixed with Recombinant Mononucleosomes (H3.1,

Active Motif 31467) or Recombinant Mononucleosomes H3K9me3 (H3.2, Active Motif

31555) and *S*-(5′-Adenosyl)-*L*-methionine chloride dihydrochloride (SAM, Sigma 7007)

in HMT buffer (50mM Tris, pH 8.0, 10 mM MgCl$_2$, 1 mM DTT, 5% Glycerol and 1 mM

PMSF). Samples were boiled and loaded on 4–20% Mini-PROTEAN TGX Gel (Bio-rad

456-1093) and Western blotting was performed with anti-His (ThermoFisher MA1-135),

anti-HA (Roche 11583816001), anti-H3 (Abcam ab1791), and anti-H3K9me3 (Abcam

ab8898) using ECL (Thermo Fisher 32209) assay according to the manufacturer's

instruction. For nucleosome binding assay, biotin labeled Recombinant

Mononucleosomes (H3.1, Active Motif 31467) or Recombinant Mononucleosomes

H3K9me3 (H3.2, Active Motif 31555) was pre-incubated with Streptavidin beads (GE

Healthcare 17-5113-01) and incubated in vitro ubiquitinated Clr4 or Clr4 proteins

purified from *S. pombe* cells. Samples were boiled and loaded on 4–20% Mini-

PROTEAN TGX Gel (Bio-rad 456-1093) and Western blotting was performed with anti-

Flag (Abcam ab1791), anti-H3K9me3 (Abcam ab8898) and anti-H3 (Abcam ab1791),

and using ECL (Thermo Fisher 32209) assay according to the manufacturer's

instruction.

Mass Spectrometry

The gel band sample was reduced and alkylated with *N*-ethylmaleimide (NEM),

digested with trypsin for 1.5 h at 37 °C and washed with ammonium

bicarbonate/acetonitrile to remove stain, SDS and other reagents. Peptides were

extracted from the gel pieces, dried down and re-dissolved in 2.5% acetonitrile, 0.1%

formic acid. Each digest was run by nanoLC-MS/MS using a 2 h gradient on a 0.075

mm x 250 mm C18 column feeding into a Q-Exactive HF mass spectrometer. All

MS/MS samples were analyzed using Mascot (Matrix Science, London, UK; version

2.6.1). Mascot was set up to search the cRAP_20150130.fasta;

custom4_20190215.fasta; SwissProt_2019_02 database (selected for

Schizosaccharomyces pombe, unknown version, 5269 entries) assuming the digestion

enzyme trypsin. Mascot was searched with a fragment ion mass tolerance of 0.060 Da

and a parent ion tolerance of 10.0 PPM. Deamidated of asparagine and glutamine,

oxidation of methionine, phospho of serine, threonine and tyrosine, GG of lysine, N-

ethylmaleimide of cysteine were specified in Mascot as variable modifications. Scaffold

(version Scaffold_4.8.9, Proteome Software Inc., Portland, OR) was used to validate

MS/MS based peptide and protein identifications. Peptide identifications were accepted

if they could be established at greater than 80.0% probability by the Peptide Prophet

algorithm[46] with Scaffold delta-mass correction. Protein identifications were accepted if

they could be established at greater than 99.0% probability and contained at least 2

identified peptides. Protein probabilities were assigned by the Protein Prophet

algorithm[47]. Proteins that contained similar peptides and could not be differentiated

based on MS/MS analysis alone were grouped to satisfy the principles of parsimony.

Proteins sharing significant peptide evidence were grouped into clusters.

Liquid-Liquid Phase Separation (LLPS)

LLPS of GFP-Clr4[Full] and GFP-Clr4[IDR] proteins were induced by diluting protein to 12

µM and 16 µM respectively in 20 mM Tris, pH 7.5, 30 mM NaCl, 0.5 mM MgAc, 1%

Glycerol and 15 mM Imidazole. To observe the effect of ubiquitin on GFP-Clr4[IDR] LLPS,

0.9 µg/µl of human ubiquitin (R&D systems U-100H) or buffer only (10 mM Hepes, pH

7.6) were added to the GFP-Clr4[IDR]. To observe the effect of 1,6-Hexanediol (Sigma

H11807) on GFP-Clr4^{IDR} LLPS, GFP-Clr4^{IDR} protein was diluted to 44.8 µM in 56 mM

Tris, pH 7.5, 84 mM NaCl, 1.4 mM MgAc, 2.8% Glycerol, 42 mM Imidazole and 3.6%

PEG8000 with or without 7.2% 1,6-Hexanediol. LLPS was imaged on Zeiss Axio Imager

M2 with differential interference contrast (DIC) and EGFP channels and processed

using Zeiss Zen software.

FRAP experiments

FRAP experiments were performed as previously described[48]. Briefly, GFP-Clr4^{IDR}

protein were diluted to 44.8 µM in 56 mM Tris, pH 7.5, 84 mM NaCl, 1.4 mM MgAc,

2.8% Glycerol, 42 mM Imidazole and 3.6% PEG8000 with or without 0.65 µg/µl

ubiquitin. FRAP experiments were carried out using Zeiss LSM710 Confocal Laser

Scanning Microscope with 63x objective. All Bleaching experiments were performed

with 488 nm argon laser at 100% intensity. Images before and after photobleaching

were taken at 5 s interval and analyzed using standard methods.

Data availability

Genome-wide datasets are deposited in the Gene Expression Omnibus (GEO) under

the accession number GSE156069. The token for private access is **ojipouckhladzeh.**

Additional references for Materials and Methods

4   Verdel, A. *et al.* RNAi-mediated targeting of heterochromatin by the RITS

518   complex. *Science* **303**, 672-676, doi:10.1126/science.1093686 (2004).

29   Kim, H. S. *et al.* Identification of a BET family bromodomain/casein kinase II/TAF-

containing complex as a regulator of mitotic condensin function. *Cell Rep* **6**, 892-

905, doi:10.1016/j.celrep.2014.01.029 (2014).

38   Bähler, J. *et al.* Heterologous modules for efficient and versatile PCR-based

gene targeting in Schizosaccharomyces pombe. *Yeast* **14**, 943-951,

doi:10.1002/(sici)1097-0061(199807)14:10<943::Aid-yea292>3.0.Co;2-y (1998).

39   Murray, J. M., Watson, A. T. & Carr, A. M. Transformation of

Schizosaccharomyces pombe: Lithium Acetate/ Dimethyl Sulfoxide Procedure.

Cold Spring Harb Protoc **2016**, pdb.prot090969, doi:10.1101/pdb.prot090969

(2016).

40   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. *Genome Biol*

10, R25, doi:10.1186/gb-2009-10-3-r25 (2009).

41   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**, 841-842, doi:10.1093/bioinformatics/btq033

(2010).

42   Winston, F. EMS and UV mutagenesis in yeast. *Curr Protoc Mol Biol* **Chapter**

13, Unit 13.13B, doi:10.1002/0471142727.mb1303bs82 (2008).

43   Roche, B., Arcangioli, B. & Martienssen, R. A. RNA interference is essential for

cellular quiescence. *Science* **354**, doi:10.1126/science.aah5651 (2016).

44   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*

Methods **9**, 357-359, doi:10.1038/nmeth.1923 (2012).

45   Garrison, E. & Marth, G. Haplotype-based variant detection from short-read

sequencing. *arXiv e-prints*, arXiv:1207.3907 (2012).

<https://ui.adsabs.harvard.edu/abs/2012arXiv1207.3907G>.

46   Keller, A., Nesvizhskii, A. I., Kolker, E. & Aebersold, R. Empirical statistical model

to estimate the accuracy of peptide identifications made by MS/MS and database

search. *Anal Chem* **74**, 5383-5392, doi:10.1021/ac025747h (2002).

47   Nesvizhskii, A. I., Keller, A., Kolker, E. & Aebersold, R. A statistical model for

identifying proteins by tandem mass spectrometry. *Anal Chem* **75**, 4646-4658,

doi:10.1021/ac0341261 (2003).

48   Wang, H. *et al.* Rubisco condensate formation by CcmM in β-carboxysome

biogenesis. *Nature* **566**, 131-135, doi:10.1038/s41586-019-0880-5 (2019).

Acknowledgement

We thank all Martienssen lab members for valuable discussions and suggestions. This

work was supported by the Howard Hughes Medical Institute-Gordon and Betty Moore

Foundation (GMBF3033) and a grant from the National Institute of Health (GM076396)

(to R.A.M.).

Author contributions

H.S.K. performed experiments and bioinformatics. B.R. contributed sRNA-seq

experiments and bioinformatics. S.B. contributed to IP for ubiquitinated Clr4 and *in vitro*

protein purification. A.Y.C. contributed to EMS mutagenesis, C.H. contributed to LLPS

562   and LT and AV contributed to Rik1-TAP purification. H.S.K. and R.A.M. designed the

563   experiments and wrote the manuscript.

Competing interests

565   The authors declare no competing interests.

Supplementary Information Table 1. *S. pombe* strains used in this study.

Number	Name	Genotype	Reference
FY2002	WT	*h⁺ otrIR(SphI)::ade6 ImrIL(NcoI)::leuI-32 ade6-DN/N ura4-DS/E leu1-32*	R. Allshire
HSK29	*ubc4-G48D (ubc4-1)*	FY2002_*ubc4-G48D-KanMX6*	This study
HSK335	*cul4-GFP (cul4-1)*	FY2002_*cul4-GFP-KanMX6*	This study
HSK351	*clr4Δ*	FY2002_*clr4Δ::KanMx6*	This study
HSK770	*dcr1Δ*	FY2002_*dcr1Δ::NatMx6*	This study
HSK790	*dcr1Δ ubc4-1*	FY2002_*dcr1Δ::NatMx6 ubc4-G48D-KanMX6*	This study
HSK606	*3xFlag-Clr4*	FY2002_*HphMX6-3xFlag-Clr4*	This study
HSK618	*3xFlag-Clr4 ubc4-1*	FY2002_*HphMX6-3xFlag-Clr4 ubc4-G48D-KanMX6*	This study
HSK1118	*3xFlag-Clr4-AllKtoR*	FY2002_*HphMX6-3xFlag-Clr4-AllKtoR*	This study
HSK971	*3xFlag-Clr4-2-8KtoR*	FY2002_*HphMX6-3xFlag-Clr4-2-8KtoR*	This study
HSK1121	*3xFlag-Clr4-2-4KtoR*	FY2002_*HphMX6-3xFlag-Clr4-2-4KtoR*	This study
HSK918	*3xFlag-Clr4 ubp3-OE*	FY2002_*HphMX6-3xFlag-Clr4 NatMX6-adh1p-ubp3*	This study
HSK1049	*3xFlag-Clr4-W31G*	FY2002_*HphMX6-3xFlag-Clr4-W31G*	This study
HSK1054	*3xFlag-Clr4-W31G ubc4-1*	FY2002_*HphMX6-3xFlag-Clr4-W31G ubc4-G48D-KanMX6*	This study
HSK679	*GFP-Swi6*	FY2002_*HphMX6-GFP-Swi6*	This study
HSK778	*GFP-Swi6 clr4Δ*	FY2002_*HphMX6-GFP-Swi6 clr4Δ::KanMX6*	This study
HSK700	*GFP-Swi6 ubc4-1*	FY2002_*HphMX6-GFP-Swi6 ubc4-G48D-KanMX6*	This study
HSK776	*GFP-Swi6 dcr1Δ*	FY2002_*HphMX6-GFP-Swi6 dcr1Δ::NatMX6*	This study
HSK808	*GFP-Swi6 dcr1Δ ubc4-1*	FY2002_*HphMX6-GFP-Swi6 dcr1Δ::NatMX6 ubc4-G48D-KanMX6*	This study
HSK518	*bdf2Δ*	FY2002_*bdf2Δ::KanMX6*	This study
HSK543	*ubc4-1 bdf2Δ*	FY2002_*ubc4-G48D-KanMX6 bdf2Δ::NatMX6*	This study
HSK545	*cul4-1 bdf2Δ*	FY2002_*cul4-GFP-KanMX6 bdf2Δ::NatMX6*	This study
HSK524	*Bdf2-3xFlag*	FY2002_*Bdf2-3xFlag-HphMX6*	This study
HSK571	*Bdf2-3xFlag ubc4-1*	FY2002_*Bdf2-3xFlag-HphMX6 ubc4-G48D-KanMX6*	This study
HSK573	*Bdf2-3xFlag cul4-1*	FY2002_*Bdf2-3xFlag-HphMX6 cul4-GFP-KanMX6*	This study
HSK583	*Bdf2-3xFlag ddb1Δ*	FY2002_*Bdf2-3xFlag-HphMX6 ddb1Δ::KanMX6*	This study
HSK575	*Bdf2-3xFlag clr3Δ*	FY2002_*Bdf2-3xFlag-HphMX6 clr3Δ::KanMX6*	This study
HSK458	*cul4-1 ubc4-1*	FY2002_*cul4-GFP-KanMX6 ubc4-G48D-NatMX6*	This study
HSK608	*ubc4::mUBE2D2*	FY2002_*KanMX6-ubc4::mUBE2D2*	This study
HSK610	*ubc4::mUBE2D2-G48D*	FY2002_*KanMX6-ubc4::mUBE2D2-G48D*	This study
KFP26	*h⁻*	*h⁻ ura4-D18 leu1-32 ade6-M210 his3-D1*	Lab stock
SPG18	*h⁹⁰*	*h⁹⁰ ura4-D18 leu2-32 ade6 -M216 his2*	Lab stock
DI304	*h⁹⁰ ubc4-G48D*	*h⁹⁰ leu1-32 ubc4-G48D-KanMX6*	Irvine et al 2009
AY709	*h⁹⁰ubc4-G48D/S137T*	*h⁹⁰ leu1-32 ubc4-G48D-KanMX6 ubc4-S137T* (EMS mutagenesis)	This study
HSK33	*raf1Δ*	FY2002_*raf1Δ::KanMx6*	This study
HSK307	*raf2Δ*	FY2002_*raf2Δ::KanMx6*	This study
HSK355	*rik1Δ*	FY2002_*rik1Δ::KanMx6*	This study
HSK351	*clr3Δ*	FY2002_*clr3Δ::KanMx6*	This study
HSK644	*ubc4-3xHA*	FY2002_*ubc4-3xHA-KanMX6*	This study
HSK646	*cul4Δ*	FY2002_*cul4Δ::NatMx6*	This study
HSK716	*GFP-Swi6 ubc4-3xHA*	FY2002_*HphMX6-GFP-Swi6 ubc4-3xHA-KanMX6*	This study
HSK717	*GFP-Swi6 cul4Δ*	FY2002_*HphMX6-GFP-Swi6 cul4Δ::NatMX6*	This study
HSK642	*ubc4-GFP*	FY2002_*ubc4-GFP-KanMX6*	This study
HSK393	*3xFlag-Rik1*	FY2002_*HphMX6-3xFlag-Rik1*	This study
HSK389	*3xFlag-Raf1*	FY2002_*HphMX6-3xFlag-Raf1*	This study
HSK274	*Raf2-3xFlag*	FY2002_*Raf2-3xFlag-HphMX6*	This study
HSK620	*3xFlag-Clr4 cul4-1*	FY2002_*HphMX6-3xFlag-Clr4 cul4-GFP-KanMX6*	This study
HSK920	*3xFlag-Clr4 rik1Δ*	FY2002_*HphMX6-3xFlag-Clr4 rik1Δ::NatMx6*	This study
HSK921	*3xFlag-Clr4 raf1Δ*	FY2002_*HphMX6-3xFlag-Clr4 raf1Δ::NatMx6*	This study

Number	Name	Genotype	Reference
HSK922	*3xFlag-Clr4 raf2∆*	FY2002_*HphMX6-3xFlag-Clr4 raf2∆::NatMx6*	This study
HSK1125	*3xFlag-Clr4-1-6KtoR*	FY2002_*HphMX6-3xFlag-Clr4-1-6KtoR*	This study
HSK1127	*3xFlag-Clr4-3-6KtoR*	FY2002_*HphMX6-3xFlag-Clr4-3-6KtoR*	This study
HSK1129	*3xFlag-Clr4-4-6KtoR*	FY2002_*HphMX6-3xFlag-Clr4-4-6KtoR*	This study
HSK1131	*3xFlag-Clr4-5-5KtoR*	FY2002_*HphMX6-3xFlag-Clr4-5-5KtoR*	This study
HSK1132	*3xFlag-Clr4-6-5KtoR*	FY2002_*HphMX6-3xFlag-Clr4-6-5KtoR*	This study
HSK1140	*3xFlag-Clr4-K87R*	FY2002_*HphMX6-3xFlag-Clr4-K87R*	This study
HSK1142	*3xFlag-Clr4-K94R*	FY2002_*HphMX6-3xFlag-Clr4-K94R*	This study
HSK1144	*3xFlag-Clr4-K109R*	FY2002_*HphMX6-3xFlag-Clr4-K109R*	This study
HSK1146	*3xFlag-Clr4-K110R*	FY2002_*HphMX6-3xFlag-Clr4-K110R*	This study
HSK1148	*3xFlag-Clr4-K113R*	FY2002_*HphMX6-3xFlag-Clr4-K113R*	This study
HSK1150	*3xFlag-Clr4-K114R*	FY2002_*HphMX6-3xFlag-Clr4-K114R*	This study
HSK1152	*3xFlag-Clr4-K122R*	FY2002_*HphMX6-3xFlag-Clr4-K122R*	This study
HSK1154	*3xFlag-Clr4-K127R*	FY2002_*HphMX6-3xFlag-Clr4-K127R*	This study
HSK1013	*3xFlag-clr4-K109/110/113/114R*	FY2002_*HphMX6-3xFlag-Clr4-K109/110/113/114R-KanMX6*	This study
HSK999	*3xFlag-Clr4-K87/94/109/110/113/114R*	FY2002_*HphMX6-3xFlag-Clr4-K87/94/109/110/113/114R-KanMX6*	This study
HSK1000	*3xFlag-Clr4-K109/110/113/114/122/127R*	FY2002_*HphMX6-3xFlag-Clr4-K109/110/113/114/122/127R-KanMX6*	This study
HSK1015	*3xFlag-Clr4-K110/113/114R*	FY2002_*HphMX6-3xFlag-Clr4-110/113/114R-KanMX6*	This study
HSK1017	*3xFlag-Clr4-K109/113/114R*	FY2002_*HphMX6-3xFlag-Clr4-109/113/114R-KanMX6*	This study
HSK1019	*3xFlag-Clr4-K109/110/114R*	FY2002_*HphMX6-3xFlag-Clr4-109/110/114R-KanMX6*	This study
HSK1021	*3xFlag-Clr4-K109/110/113R*	FY2002_*HphMX6-3xFlag-Clr4-109/110/113R-KanMX6*	This study
HSK1082	*3xFlag-Clr4-K109/110R*	FY2002_*HphMX6-3xFlag-Clr4-109/110R-KanMX6*	This study
HSK1062	*3xFlag-Clr4-K109/113R*	FY2002_*HphMX6-3xFlag-Clr4-109/113R-KanMX6*	This study
HSK1064	*3xFlag-Clr4-K109/114R*	FY2002_*HphMX6-3xFlag-Clr4-109/114R-KanMX6*	This study
HSK1086	*3xFlag-Clr4-K110/113R*	FY2002_*HphMX6-3xFlag-Clr4-110/113R-KanMX6*	This study
HSK1065	*3xFlag-Clr4-K110/114R*	FY2002_*HphMX6-3xFlag-Clr4-110/114R-KanMX6*	This study
HSK1084	*3xFlag-Clr4-K113/114R*	FY2002_*HphMX6-3xFlag-Clr4-113/114R-KanMX6*	This study
HSK919	*3xFlag-Clr4 ubp3∆*	FY2002_*HphMX6-3xFlag-Clr4 ubp3∆::NatMx6*	This study
HSK395	*ubp3-OE***	FY2002_*HphMX6-adh1p-ubp3* (*ubp3* overexpression)	This study
HSK401	*ubp8-OE***	FY2002_*HphMX6-adh1p-ubp8* (*ubp8* overexpression)	This study

Supplementary Information Table 2. Oligonucleotides used in this study.

Name	Description	Purpose	Sequence
p30-qPCR-F	*dg*	ChIP-qPCR, RT-qPCR	CCATATCAATTTCCCATGTTCC
p30-qPCR-R	*dg*	ChIP-qPCR, RT-qPCR	CATCAAGCGAGTCGAGATGA
p33-qPCR-F	*dh*	ChIP-qPCR, RT-qPCR	TATCCTGCGTCTCGGTATCC
p33-qPCR-R	*dh*	ChIP-qPCR, RT-qPCR	CTGTTCGTGAATGCTGAGAAAG
act1-qPCR-F	*act1*	ChIP-qPCR, RT-qPCR	TGCACCTGCCTTTTATGTTG
act1-qPCR-F	*act1*	ChIP-qPCR, RT-qPCR	TGGGAACAGTGTGGGTAACA
ade6-qPCR-F	*ade6*	ChIP-qPCR	TTCCTACTGCCATCAAAGCA
ade6-qPCR-R	*ade6*	ChIP-qPCR	GCACGCTGTTGAATTGAGAA
ura4-qPCR-F	*ura4*	ChIP-qPCR	TACCTTTGGGACGTGGTCTC
ura4-qPCR-R	*ura4*	ChIP-qPCR	CCCGTCTCCTTTAACATCCA
irc3-R-qPCR-F	*irc-3R*	ChIP-qPCR	GGCAACCTTTCAAGTTCTCG
irc3-R-qPCR-R	*irc-3R*	ChIP-qPCR	ATTCTTCCCATCTGCGTTTG
tlh1-R-qPCR-F	*tlh1*	ChIP-qPCR	GTTCTCCATGCTCCACTGGT
tlh1-R-qPCR-R	*tlh1*	ChIP-qPCR	CCAACAGTGGTCTTCGGTTT
ubp3-qPCR-F	*ubp3*	RT-qPCR	GCCAAATTTTTGGTGGTCAG
ubp3-qPCR-R	*ubp3*	RT-qPCR	TGAATGTCGAGTTGCAGAGG
Clr4(His)-NdeI-F	6xHis-Clr4	Plasmid construction	CCGCGCGGCAGCCATATGTCGCCTAAACAAGAGGAGTATG
Clr4(His)-NdeI-R	6xHis-Clr4	Plasmid construction	CGGATCCTCGAGCATATGTTAACCGAAAAGCCAGCCACGACAA
Clr4-BamHI-F	6xHis-GFP-Clr4	Plasmid construction	CGGGTACCATGGATCCGATGTCGCCTAAACAAGAGGAGTATG
Clr4-XhoI-R	6xHis-GFP-Clr4	Plasmid construction	GGTGGTGGTGCTCGAGTTAACCGAAAAGCCAGCCACGACAA
Clr4-IDR-BamHI-F	6xHis-GFP-IDR	Plasmid construction	CGGGTACCATGGATCCGTCTGCCGTTTTAGCAGAATGGAAAA
Clr4-IDR-XhoI-R	6xHis-GFP-IDR	Plasmid construction	GGTGGTGGTGCTCGAGCTATTCGGGTCCTTCAATTTCCCGC

Supplementary Information Table 3. Plasmids used in this study.

Name	Description	Application
pET-14b-Clr4	6xHis-Clr4	Recombinant 6xHis-Clr4 protein purification from *E. coli*
pHis-parellel-GFP-Clr4^{Full}	6xHis-GFP-Clr4^{Full}	Recombinant 6xHis-GFP-Clr4^{Full} protein purification from *E. coli*
pHis-parellel-GFP-Clr4^{IDR}	6xHis-GFP-Clr4^{IDR}	Recombinant 6xHis-GFP-Clr4^{IDR} protein purification from *E. coli*

Figure 1

Figure 1. Mono-ubiquitination of Clr4 by Ubc4-CLRC is required for H3K9 trimethylation.

a, ChIP-qPCR assays showing enrichment of H3K9me2 (left) and H3K9me3 (right) at centromeric *dg* locus in the indicated strains. *clr4* deletion mutant (*clr4Δ*) shows complete loss of H3K9me2 and H3K9me3. Error bars, standard deviation (S.D., 3 biological replicates).

b, ChIP-qPCR assays showing enrichment of H3K9me2 at centromeric *dg* locus in the indicated strains. *dcr1Δ ubc4-1* double mutant shows complete abolishment of H3K9me2. Error bars, S.D. (3 biological replicates).

c, Western blot (WB) analyses of immunoprecipitated 3xFlag-Clr4 (3xFlag IP) from WT and *ubc4-1* mutant cells treated with proteasome inhibitor MG-132 (100μM) for 4 h before harvest. Bottom, WB analysis of 3xFlag-Clr4 from whole cell extract (WCE).

d, Schematic diagram of domain structure of Clr4 protein and the position of lysine residues. CD: chromodomain, PS: PostSET. Ovals show the position of all 36 lysines of Clr4, which are subgrouped to 6 groups. Second group (2-8K) is highlighted and amino acid sequences are shown. 2-4K region covers K109-K114.

e, WB analyses same as **c**, but showing WT, all-lysine mutant (*AllKtoR*), second block mutant (*2-8KtoR*) and mutant of 4 lysines (K109, K110, K113 and K114) of second block (*2-4KtoR*).

f, Western blot (WB) analysis of immunoprecipitated 3xFlag-Clr4 from WT and *ubp3*-overexpression *(ubp3-OE)* strains.

g, ChIP-seq reads of H3K9me2 and H3K9me3 mapped to chromosome 3 centromere right arm in the indicated strains. Relat. enrich., relative enrichemnt. Total count of ChIP-seq reads from different strains were divided by the total count of WT to show relative enrichment. Bottom, schematic diagram of chromosome 3 centromere right arm showing innermost repeat (*imr*), outermost repeats (*dg*, *dh*) and boundary region (*irc*) between heterochromatin and euchromatin. Vertical red lines and pink boxes represent tRNA and euchromatic genes respectively.

Figure 2. **Clr4 mono-ubiquitination induces its dissociation from chromatin.**

a, ChIP-seq reads of 3xFlag-Clr4 mapped to chromosome 3 centromere (*cnt3*) right arm in indicated strains. Relat. enrich., relative enrichemnt. Bottom, schematic diagram of chromosome 3 centromere (*cnt3*) right arm.

b, *In vitro* protein binding assay of *in vitro* ubiquitinated Clr4 protein with nucleosomes. His-Clr4 purified from *E. coli* cells was ubiquitinated with human E1, E3 (UBE2D3) and HA-ubiquitin and incubated with nucleosomes with unmodified histone H3 (me0) or H3K9me3 histone (me3) anchored to streptavidin beads. Input and precipitated proteins were analyzed by Western blot (WB) using antibodies as shown.

c, ChIP-seq reads of 3xFlag-Clr4 mapped to chromosome 3 centromere (*cnt3*) right arm in indicated strains. Relat. enrich., relative enrichment. Bottom, schematic diagram of chromosome 3 centromere (*cnt3*) right arm.

d, Schematic diagram showing the presumptive subunit arrangement in the complex of Ubc4-CLRC and its substrate Clr4. Red circle (U), ubiquitin. The arrow shows the mono-ubiquitination activity of Ubc4-CLRC toward Clr4.

Figure 3. Ubiquitin reduces GFP-Clr4^{IDR} LLPS activity.

a. Live cell imaging showing GFP-Swi6 heterochromatic foci in indicated strains. In WT 4 foci represent three telomeres and the centromeric chromocenter, which disappears in *dcr1Δ*. Only the chromocenter focus is retained in *ubc4-1* which loses H3K9me from telomeres. The foci are completely abolished in *clr4Δ* and *dcr1Δ ubc4-1* mutants. Scale bar, 2.5μm. Dashed lines show the outline of each *S. pombe* cells.

b, Reduced size of LLPS of GFP-Clr4^{IDR} (amino acid 51-220) by addition of ubiquitin. Buffer or ubiquitin was added to GFP-Clr4^{IDR} protein and the formation of LLPS was monitored. Scale bar, 10μm.

c, Reduced size of LLPS of GFP-Clr4^{IDR} by addition of ubiquitin (**b**). Top 30 LLPS droplets are measured and size distribution is shown. Black bar, average. ***P value=1.1X10^{-5}.

d, FRAP of GFP-Clr4^{IDR} by addition of ubiquitin before and after bleaching. The change of fluorescence was analyzed and indicated in mean and S.D. (n=7 and n=9 for Buffer and Ubiquitin, respectively). Arrow shows the time of bleaching (~40s).

Figure 4. **Ubc4-CLRC poly-ubiquitinates Bdf2 and regulates sRNA production.**

a, sRNA-seq reads mapped to chromosome 3 centromere right arm. The increased sRNA production in *ubc4-1* and *cul4-1* mutants is suppressed by the deletion mutation of *bdf2*. sRNA at boundary region (*irc*) were abolished in all mutants. Remaining sRNA peaks in *dcr1Δ* are tRNA-related products. Relat. enrich., relative enrichment of sRNA without tRNA. Bottom, schematic diagram of chromosome 3 centromere right arm.

b, ChIP-seq reads of Bdf2-3xFlag mapped to chromosome 3 centromere (*cnt3*) right arm in indicated strains. Relat. enrich., relative enrichment. Bottom, schematic diagram of chromosome 3 centromere right arm.

c, Western blot (WB) analyses for Bdf2-3xFlag in indicated strains. Ponceau S staining of membrane is shown for equal loading of protein for WB.

d, Western blot (WB) analyses of immunoprecipitated Bdf2-3xFlag (Flag IP) from WT and *ubc4-1* mutant cells treated with proteasome inhibitor MG-132 (100μM) for 4 h before harvest (left). Bdf2-3xFlag enrichment by ubiquitination affinity beads (UAB IP) are shown on right.

e, Model for the function of Ubc4-CLRC-mediated Clr4 mono-ubiquitination and Bdf2 poly-ubiquitination. Left, Bdf2 recruits RNA Pol II complex to produce centromeric ncRNAs, which are converted to sRNAs by the sequential action of RNAi machineries and these sRNAs reinforce the Clr4 activity for H3K9me2. Ubc4-CLRC poly-ubiquitinates Bdf2 for its degradation and reduces recruitment of RNA Pol II. Right, Clr4 converts H3K9me2 to H3K9me3 but because of its strong binding affinity to H3K9me3, it remains to bind nucleosome with H3K9me3. Clr4 mono-ubiquitination by Ubc4-CLRC dissociates Clr4 from chromatin and Ubp3 removes the mono-ubiquitin from Clr4. Unmodified Clr4 binds other nucleosomes with H3K9me3 and converts RNAi-independent H3K9me1 or -2 nucleosomes nearby to H3K9me3. Both reduction of sRNA and conversion of H3K9me2 to H3K9me3 promote the transition from CTGS to TGS.

Extended Data Figure 1. Analysis of *ubc4* mutants and its protein stability.

a, Ubc4 shows strong amino acid sequence identity and similarity with its mouse homologues, mUBE2D2 and mUBE2D3. Glycine 48 of Ubc4 is conserved in mouse homologues but Serine 137 is not. Multiple sequence alignment of Ubc4 and its homologues was generated using Clustal Omega.

b, *ubc4-G48D* (*ubc4-1*) and *cul4-GFP* (*cul4-1*) mutants decrease heterochromatin silencing. Swapping *S. pombe ubc4* gene with mouse UBE2D2 gene complements essential function of Ubc4 and *mUBE2D2-G48D* also shows silencing defect. Left, silencing assay of *imr1L::ura4*[+] and *otr1R::ade6*[+] on no Uracil (-Ura) and low-adenine (Low ade) media to assess *ura4*[+] and *ade6*[+] expression and TBZ-sensitivity assay on 15μg/ml microtubule-destabilizing agent, thiabendazole (TBZ) in the indicated strains. NS, non-selective medium. Right, RT-qPCR analysis for centromeric *dg* transcription (normalized to *act1*). Error bars, S.D. (3 biological replicates).

c, Ubc4 protein is greatly reduced in *ubc4-G48D* mutant. For Western blot analysis of Ubc4 protein in the indicated strains, antibody for mUBE2D3 was used to detect Ubc4. Tubulin was used as loading control.

d, Mating-type switching defect in *ubc4-G48D* mutant was suppressed by the secondary mutation, *S137T*. Non-switchable *h⁻* and WT and *ubc4-G48D* mutants in switchable strain (h^{90}) were plated and incubated on sporulation medium and spores were stained with iodine vapors. The enrichment of starch in spore enables staining with iodine and dark staining indicates the strain is switch competent and can make spores but light staining indicates the strain is switching defective and cannot make spores efficiently. *ubc4-S137T* mutation which was identified by EMS mutagenesis of *ubc4-G48D*, iodine staining and whole genome sequencing suppresses the mating switching defect of *ubc4-G48D* mutant (ubc4-*G48D/S137T*).

e, *ubc4-S137T* mutation restores decreased protein level of *ubc4-G48D* mutant. Western blot analysis of Ubc4 protein and tubulin as control. Middle panel shows the longer exposure of upper panel for Ubc4 protein.

f, *ubc4-S137T* mutation suppresses the silencing defect of *ubc4-G48D* mutant. Silencing assay is same with **b,** but FOA containing (FOA) media, which kills *ura4⁺* expressing cells, was also used to assess *ura4⁺* expression.

Extended Data Figure 2. Ubc4 and Cul4 regulate H3K9me status.

a-c, ChIP-qPCR assays showing enrichment of H3K9me2 (**a**), H3K9me3 (**b**) and RNA Pol II (Rpb1, **c**) at centromeric *dh*, *ade6+* and *ura4+* reporter genes at centromere, boundary region between heterochromatin and euchromatin of centromere 3 (*irc3-R*) and telomeric *tlh1* locus. Error bars, S.D. (3 biological replicates).

d, ChIP-seq reads of H3K9me2 (left) and H3K9me3 (right) mapped to telomere 1 left region. Relat. enrich., relative enrichment. Bottom, schematic diagrams of *tlh1* gene locus.

e, ChIP-qPCR assays showing slightly decreased enrichment of HP1 protein, Swi6 at centromeric *dg* locus in *ubc4-1* and *cul4-1* strains. Error bars, S.D. (2 biological replicates).

Extended Data Figure 3. Ubc4 and Cul4 regulate H3K9 methylation through RNAi

a, ChIP-seq reads of H3K9me2 and H3K9me3 mapped to chromosome 1 centromere (*cnt1*) in the indicated strains. Dashed lines indicates the loci where reporter genes, *ura4+* and *ade6+* are inserted at centromere 1. The change of H3K9me2 and H3K9me3 on *ura4+* and *ade6+* reporter genes are shown (top). Arrow in the box shows the direction of gene. sRNA-seq reads of WT are also shown. Relat. enrich., relative enrichment. Bottom, schematic diagram of chromosome 1 centromere (*cnt1*) showing innermost repeat (*imr*), outermost repeats (*dg*, *dh*) and centromere core (*cen*). Vertical red lines and pink boxes represent tRNA and euchromatic genes respectively.

b, TBZ-sensitivity assay on media containing 15μg/ml thiabendazole (TBZ) and silencing assay of *otr1R::ade6+* on low-adenine (Low ade) media in the indicated strains.

c, ChIP-qPCR assays showing enrichment of H3K9me2 at telomeric *tlh1* loci in the indicated strains. Error bars, S.D. (2 biological replicates).

d, ChIP-qPCR assays showing enrichment of H3K9me3 at centromeric *dg* and telomeric *tlh1* loci in the indicated strains. Error bars, S.D. (2 biological replicates).

e, ChIP-qPCR assays showing enrichment of RNA Pol II (Rpb1) at centromeric *dg* (left) and telomeric *tlh1* (right) loci in the indicated strains. Error bars, S.D. (2 biological replicates).

Extended Data Figure 4. Ubc4 and Cul4 localize to nuclear periphery and Ubc4 forms complex with CLRC.

a, *ubc4* mutant with triple HA tag at carboxyl-terminus (*ubc4-3xHA*) and *cul4* deletion mutant (*cul4Δ*) are viable but show severe growth defect. Big cells in each picture show WT cells. *ubc4-3xHA* cells were grown on YES media and *cul4Δ* cells were grown on low-adenine media.

b, ChIP-qPCR assays showing enrichment of H3K9me2 and H3K9me3 at centromeric *dg* locus in the indicated strains. Error bars, S.D. (2 biological replicates).

c. Live cell imaging showing the localization of GFP-Swi6 in indicated strains. The localization of GFP-Swi6 is completely abolished in *ubc4-3xHA* and *cul4Δ* mutants, which is consistent with H3K9me2 and H3K9me3 ChIP-qPCR results (**b**). Scale bar, 2.5μm.

d, Live cell imaging showing that Cul4-GFP and Ubc4-GFP localize to nuclear periphery. The strain without GFP tag (untag) was used as negative control. Scale bar, 2.5μm.

e, Co-immunoprecipitation (Co-IP) of Cul4-GFP with Ubc4. Whole-cell extracts (WCE) were immunoprecipitated with GFP antibody and Western blot analyses were carried out with GFP and mUBE2D3 antibodies.

f, Co-immunoprecipitation (Co-IP) of 3xFlag-Rik1, 3xFlag-Raf1 and Raf2-3xFlag with Ubc4. Whole-cell extracts (WCE) were immunoprecipitated with Flag antibody and Western blot analyses were carried out with Flag and mUBE2D3 antibodies.

Extended Data Figure 5. Characterization of ubiquitination sites of Clr4 by Mass Spectrometry analysis.

a, same as Fig. 1c**,** but shows extended exposure of WB analysis for immunoprecipitated 3xFlag-Clr4 from WT and _ubc4-1_ mutant cells. Asterisk indicates possible di- or multi-ubiquitin chain which is also largely dependent on Ubc4.

b, WB analysis for 3xFlag-Clr4 enriched by α-Flag affinity beads and WB analysis using ubiquitin antibody.

c, WB analyses same as Fig. 1c, but showing 3xFlag-Clr4 enrichment by ubiquitination affinity beads (UAB IP).

d, Proteasome pathway-independent Clr4 ubiquitination. Western blot (WB) analysis of immunoprecipitated 3xFlag-Clr4 from wild type (WT) and second block mutant (_2-8KtoR_) with or without treatment of proteasome inhibitor MG-132 for 4h before harvest.

e, WB analyses same as Fig. 1c, but showing also CLRC mutants like _cul4-1_, _rik1Δ_, _raf1Δ_ and _raf2Δ_.

f, ChIP-qPCR assay showing enrichment of 3xFlag-Clr4 at centromeric _dh_ locus in the indicated strains. Error bars, S.D. (2 biological replicates).

g, 6xHis-Clr4 purified from *E. coli* cells were *in vitro* ubiquitinated with human E1, E3 (UBE2D3) and ubiquitin and analyzed by Mass Spectrometry. The peptides with possible ubiquitination sites were shown. Di-glycine-Lysine (GG) remnants after trypsin digestion indicate possible ubiquitination sites. Prob., probability. Mass Spectrometry results were analyzed using Scaffold program.

h, Amino acid sequence of 6xHis-Clr4 protein. Sequence coverage by Mass Spectrometry analysis is highlighted in yellow and amino acids with post-translational modifications (PTM) are shown in green. Possible ubiquitination sites are highlighted with a red box.

i, Schematic diagram of domain structure of Clr4 protein and the position of lysine residues same as Fig. 1d but also shows possible ubiquitination sites characterized by Mass Spectrometry analysis on top.

Extended Data Figure 6. Characterization of ubiquitination sites of Clr4 by genetic analysis.

a, Schematic diagram of domain structure of Clr4 protein and position of lysine residues, same as Fig. 1d.

b, Western blot (WB) analysis of immunoprecipitated 3xFlag-Clr4 from all-lysine mutant (*AllKtoR*) and 6 block mutants.

c-f, Western blot (WB) analysis of immunoprecipitated 3xFlag-Clr4 from 8 individual lysine to arginine mutants (**c**), 4 or 6 lysines to arginines mutants (**d**), triple combination of K109, K110, K113 and K114 to arginines mutants (**e**) or double combination of K109, K110, K113 and K114 to arginines mutants (**f**) of second block (2-8K) of Clr4.

g, ChIP-qPCR assays showing enrichment of H3K9me2 (left), H3K9me3 (right) at centromeric *dh* locus in the indicated strains. Error bars, S.D. (3 biological replicates).

Extended Data Figure 7. Increased enrichment of Clr4 protein at centromeric heterochromatin in *ubc4-1, cul4-1*, second block mutant (*2-8KtoR*) and all-lysine mutant (*AllKtoR*).

a, Histone methylation assay. Nucleosomes with unmodified histone H3 (H3K9me0) or H3K9me3 histone (H3K9me3) were incubated with unmodified (ubiquitin, -) or *in vitro* ubiquitinated Clr4 proteins (ubiquitin, +) for the indicated times. Western blot (WB) analyses were carried out using H3K9me2 and H3K9me3 antibodies.

b, ChIP-qPCR assay showing enrichment of 3xFlag-Clr4 at centromeric *dh* locus in the indicated strains. Error bars, S.D. (3 biological replicates).

c, Schematic diagram for *in vitro* protein binding assay with nucleosomes. First, nucleosomes were anchored to Streptavidin beads using Biotin moiety at the end of DNA. Unmodified or ubiquitinated Clr4 proteins are added to reaction mixture and bound proteins were analyzed further by WB analyses.

d, *In vitro* protein binding assay of Clr4 protein from *S. pombe* cells with nucleosomes. 3xFlag-Clr4 purified from *S. pombe* cells were incubated with nucleosomes with unmodified histone H3 (me0) or H3K9me3 histone (me3) anchored to streptavidin beads. Input and precipitated proteins were analyzed by WB using antibodies as shown. Similar to the results of Fig. 2b, ubiquitinated Clr4 did not bind to the nucleosome.

e, ChIP-qPCR assay showing enrichment of 3xFlag-Clr4 at centromeric *dh* locus in the indicated strains. Error bars, S.D. (3 biological replicates).

Extended Data Figure 8. The role of Ubp3 for de-ubiquitination of Clr4

a, Western blot (WB) analysis of immunoprecipitated 3xFlag-Clr4 shows that *ubp3* deletion does not cause clear change of Clr4 ubiquitination.

b, Silencing assay of centromeric *ade6⁺* gene (*otr1R::ade6⁺*) on low-adenine (Low ade) and *ura4⁺* gene (*imr1L::ura4⁺*) on no Uracil (-Ura) and 5'-fluroorotic acid (FOA) media for count selection of *ura4⁺* expression in Ubp3- or Ubp8-overexpression (-OE) cells. TBZ-sensitivity assay on media containing 15μg/ml thiabendazole (TBZ) media.

c. RT-qPCR analysis for *ubp3*, centromeric *dh* and *act1* transcripts in the indicated strains. Error bars, S.D. (3 biological replicates).

d, ChIP-qPCR assays show enrichment of H3K9me2 (left) and H3K9me3 (right) at centromeric *dg* locus in the indicated strains. Error bars, S.D. (3 biological replicates).

e, ChIP-qPCR assays showing enrichment of RNA Pol II (Rpb1) at centromeric *dg* locus in the indicated strains. Error bars, S.D. (3 biological replicates).

f, ChIP-qPCR assays show enrichment of 3xFlag-Clr4 at centromeric *dg* locus in the indicated strains. Error bars, S.D. (3 biological replicates).

Extended Data Figure 9. LLPS of GFP-Clr4 proteins and disorder prediction

a. Live cell imaging showing the localization of GFP-Swi6 in indicated strains, same as Fig. 3a but showing GFP and DIC toghether. Scale bar, 2.5μm.

b, Schematic diagram of domain structure of Clr4 protein. Disorder prediction (PONDR) shows that the hinge region between chromodomain (CD) and PreSET of Clr4 is highly disordered. Red bar below the Clr4 schematic diagram shows the location of amino acids which is used for GFP-Clr4^{IDR} construct (51-220 amino acids).

c, LLPS of GFP-Clr4^{Full} protein. DIC; Differential interference contrast. Scale bar, 10μm.

d, The effect of 7.2% (w/v) 1,6-Hexanediol on LLPS of GFP-Clr4^{IDR} protein. Scale bar, 5μm.

e, FRAP of GFP-Clr4^{IDR} by addition of ubiquitin before and after bleaching. Representative droplests from Buffer and Ubiquitin with or without bleach were shown. Arrow shows the time of bleaching (~40s). Scale bar, 2μm

f, Schematic diagram of domain structure of Bdf2 protein (top). Disorder prediction (PONDR) shows that two Bromo and BET domains are ordered but the other regions are highly disordered.

g, Schematic diagram of domain structure of mouse SUV39H1 protein (top). Disorder prediction (PONDR) shows that the hinge region between chromodomain (CD) and PreSET of mSUV39H1 is highly disordered.

Extended Data Figure 10. Increased centromeric sRNA in *ubc4-1* and *cul4-1* mutants.

a, sRNA-seq reads mapped to chromosome 1 centromere. *ura4+* reporter gene is inserted in left *imr* (innermost repeats) just next to where sRNAs are produced. *ade6+* is located in right *otr* (outer repeats) where sRNA is hugely produced especially in *ubc4-1* and *cul4-1* mutants. Relat. enrich., relative enrichment of sRNA without tRNA.

b, Increased sRNA production at *ade6+* reporter gene in *ubc4-1* and *cul4-1* mutants but not at *ura4+*. sRNA-seq reads of *ura4+* and *ade6+* reporter genes inserted at centromere 1 heterochromatin in the indicated strains. Relat. enrich., relative enrichment of sRNA.

c, ChIP-qPCR assays showing enrichment of Ago1 and Chp1 at centromeric *dh* locus in the indicated strains. Error bars, S.D. (3 biological replicates).

d, ChIP-qPCR assays showing enrichment of Bdf2-3xFlag at centromeric *dg* (left) and euchromatic *act1* (right) loci in the indicated strains. Error bars, S.D. (2 biological replicates).

e, ChIP-seq reads of Bdf2-3xFlag mapped to euchromatic loci on chromosome 1 right arm and rDNA loci on chromosome 3 right end in indicated strains. Relat. enrich., relative enrichment. Bottom, schematic diagram of chromosomal loci.