Transmission of one predicts another: Apathogenic proxies for transmission dynamics

of a fatal virus

Marie L.J. Gilbertson[1†], Nicholas M. Fountain-Jones[2]*, Jennifer L. Malmberg[3,4]*, Roderick B.

Gagne[3,5], Justin S. Lee[3], Simona Kraberger[6], Sarah Kechejian[3], Raegan Petch[3], Elliott Chiu[3],

Dave Onorato[7], Mark W. Cunningham[8], Kevin R. Crooks[9], W. Chris Funk[10], Scott Carver[2], Sue

VandeWoude[3], Kimberly VanderWaal[1], Meggan E. Craft[1,11]

[†]Corresponding author: jone1354@umn.edu

*These authors contributed equally

[1]Department of Veterinary Population Medicine, University of Minnesota, St Paul, MN 55108.

[2]School of Natural Sciences, University of Tasmania, Hobart Australia 7001.

[3]Department of Microbiology, Immunology, and Pathology, Colorado State University, Fort

Collins, CO 80523

[4]Department of Veterinary Sciences, University of Wyoming, Laramie, Wyoming 82071.

[5]Wildlife Futures Program, Department of Pathobiology, University of Pennsylvania School of

Veterinary Medicine, Philadelphia, PA 19104

[6]The Biodesign Center for Fundamental and Applied Microbiomics, Arizona State University,

Tempe, Arizona, AZ 85287, USA

[7]Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission,

Naples, FL 34114.

[8]Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission,

Gainesville, FL 32601.

[9]Department of Fish, Wildlife, and Conservation Biology, Colorado State University, Fort Collins,

CO 80523.

[10]Department of Biology, Graduate Degree Program in Ecology, Colorado State University, Fort

Collins, CO 80523.

[11] Department of Ecology, Evolution and Behavior, University of Minnesota, St Paul, MN 55108.

Abstract

Identifying drivers of transmission prior to an epidemic—especially of an emerging pathogen—is

a formidable challenge for proactive disease management efforts. We tested a novel approach

in the Florida panther, hypothesizing that apathogenic feline immunodeficiency virus (FIV)

transmission could predict transmission dynamics for pathogenic feline leukemia virus (FeLV).

We derived a transmission network using FIV whole genome sequences, and used exponential

random graph models to determine drivers structuring this network. We used these drivers to

predict FeLV transmission pathways among panthers and compared predicted outbreak

dynamics against empirical FeLV outbreak data. FIV transmission was primarily driven by

panther age class and distances between panther home range centroids. Prospective FIV-

based modeling predicted FeLV dynamics at least as well as simpler, often retrospective

approaches, with evidence that FIV-based predictions captured the spatial structuring of the

observed FeLV outbreak. Our finding that an apathogenic agent can predict transmission of an

analogously transmitted pathogen is an innovative approach that warrants testing in other host-

pathogen systems to determine generalizability. Use of such apathogenic agents holds promise

for improving predictions of pathogen transmission in novel host populations, and could thereby

provide new strategies for proactive pathogen management in human and animal systems.

Keywords

50
51

Introduction

Infectious disease outbreaks can have profound impacts on conservation, food security, and global health and economics. Mathematical models have proven a vital tool for understanding transmission dynamics of pathogens [1], but struggle to predict the dynamics of novel or emerging agents [2]. This is at least partially due to the challenges associated with characterizing contacts relevant to transmission processes. Common modeling approaches that assume all hosts interact and transmit infections to the same degree ignore key drivers of transmission. Such drivers can include specific transmission-relevant behaviors including grooming or fighting in animals [3], concurrent sexual partnerships in humans [4], or homophily [5], and result in flawed epidemic predictions [6,7]. Further, identifying drivers of transmission and consequent control strategies for any given pathogen is typically done reactively or retrospectively in an effort to stop or prevent further outbreaks or spatial spread (e.g. [8,9]). These constraints limit the ability to perform prospective disease management planning tailored to a given target population, increasing the risk of potentially catastrophic pathogen outbreaks, as observed in humans [10], domestic animals [11], and species of conservation concern (e.g., [12–14]).

A handful of studies have evaluated whether common infectious agents present in the healthy animal microbiome or virome can indicate contacts between individuals that may translate to interactions promoting pathogen transmission [15–22]. Such an approach circumvents some of the uncertainties associated with more traditional approaches to contact detection [6]. In these cases, genetic evidence from the transmissible agent itself is used to define between-individual interactions for which contact was sufficient for transmission to occur. Results of such studies show mixed success [15–18]. For example, members of the same household [19,20] or animals with close social interactions [21,22] have been found to share microbiota, but disentangling social mechanisms of this sharing is complicated by shared diets, environments, and behaviors [23].

These studies have, however, revealed ideal characteristics of non-disease inducing

infectious agents (hereafter, *apathogenic agents*) for use as markers of transmission-relevant

interactions. Such apathogenic agents should have rapid mutation rates to facilitate discernment

of transmission relationships between individuals over time [24,25]. Furthermore, these agents

should be relatively common and well-sampled in a target population, have a well-characterized

mode of transmission that is similar to the pathogen of interest, and feature high strain alpha-

diversity (local diversity) and high strain turnover [25,26]. RNA viruses align well with these

characteristics [27] such that apathogenic RNA viruses could act as "proxies" of specific modes

of transmission (i.e., direct transmission) and indicate which drivers underlie transmission

processes. Such drivers, including but not limited to host demographics, relatedness, specific

behaviors, or space use, would subsequently allow prediction of transmission dynamics of

pathogenic agents with the same mode of transmission [25].

Here, we test the feasibility of this approach using a naturally occurring host-pathogen

system to test if an apathogenic RNA virus can act as a proxy for direct transmission processes

and subsequently predict transmission of a pathogenic RNA virus. Florida panthers (*Puma*

concolor coryi) are an endangered subspecies of puma found only in southern Florida. We have

documented that this population is infected by several feline retroviruses relevant to our study

questions [28,29]. Feline immunodeficiency virus (FIVpco; hereafter, FIV) occurs in

approximately 50% of the population and does not appear to cause significant clinical disease

[28]. FIV is transmitted by close contact (i.e., fighting and biting), generally has a rapid mutation

rate (intra-individual evolution rate of 0.00129 substitutions/site/year; [30]), and, as a chronic

retroviral infection, can be persistently detected after the time of infection. Panthers are infected

with feline leukemia virus (FeLV), also a retrovirus, which caused a well documented, high

mortality outbreak among panthers in 2002-2004 [29]. FeLV infrequently spills over into

panthers following exposure to infected domestic cats [31]. Once spillover occurs, FeLV is

transmitted between panthers by close contact and results in one of three infection states:

progressive, regressive, or abortive infection [29]. Progressive cases are infectious and result in

mortality; regressive infections are unlikely to be infectious—though this is unclear in panthers—

and recover [29,32,33]. Abortive cases clear infection and are not themselves infectious [32].

The objectives of this study were therefore: (1) to determine which drivers shape FIV

transmission in Florida panthers, and (2) test if these drivers can predict transmission dynamics

of analogously transmitted FeLV in panthers. Success of this approach in our model system

would pave the way for testing similar apathogenic agents in other host-pathogen systems,

thereby improving our understanding of drivers of individual-level heterogeneity in transmission,

and consequently our ability to predict transmission dynamics of novel agents in human and

animal populations.

Methods

Dataset assembly

We assembled an extensive dataset covering almost 40 years of Florida panther

research and including panther sex and age class. A subset of the population is monitored using

very high frequency (VHF) telemetry collars, with relocations determined via aircraft typically

three times per week. Previous panther research has generated a microsatellite dataset for

monitored panthers [34], and a dataset of 60 full FIV genomes (proviral DNA sequenced within

a tiled amplicon framework in [35]). In addition, to augment observations from the 2002-04 FeLV

outbreak [29], we leveraged an FeLV database which documents FeLV status (positive and

negative) for 31 panthers from 2002-04 as determined by qPCR.

FIV transmission inference

To determine drivers of FIV transmission, we first generated a "who transmitted to

whom" transmission network using 60 panther FIV genomes collected from 1988 to 2011

(average of minimum annual panther counts across this period was 62.3 panthers; [36]). We

used the program Phyloscanner [37] (see Figure 1 for workflow across all analyses), which

assumes both within- and between-host evolution when inferring transmission relationships

between sampled and even unsampled hosts [37]. Phyloscanner operates in a two step

process, first inferring within- and between-host phylogenies in windows along the FIV genome.

Then, using the within-host viral diversity gleaned from deep sequencing, Phyloscanner

functionally performs ancestral state reconstruction to infer transmission relationships between

hosts, outputting transmission trees or networks. For Phyloscanner's step one, we used 150bp

windows, allowing 25bp overlap between windows. To test sensitivity to this choice, we

separately ran a full Phyloscanner analysis with 150bp windows, but without overlap between

windows (supplementary methods). The tiled amplicon PCR approach used to generate our FIV

genomic data biases for detection of one known variant, such that we did not expect detectable

superinfections. In the second step of Phyloscanner, we therefore held the parameter which

penalizes within-host diversity (k) equal to 0. We used a patristic distance threshold of 0.05 and

allowed missing and more complex transmission relationships. Because we had uneven read

depth across FIV genomes, we downsampled to a maximum of 200 reads per host. The output

of the full Phyloscanner analysis was a single transmission network (hereafter, *main FIV*

network), but see supplementary methods for details regarding analysis of the sensitivity of our

results to variations in and summary across multiple transmission networks.

Figure 1: *Conceptual workflow across all analysis steps. Processes are shown on the left in*

blue; specific outcomes are shown on the right in green; the final analysis outcome is in yellow

at the bottom right. Solid lines show direct flows or outcomes. Dashed lines show processes

acting on or in concert with prior outcomes: for example, exponential random graph modeling

(ERGM) was performed using the FIV transmission network, and the combination of the two

produced the ERGM coefficients outcome.

Statistical analysis of FIV transmission networks

Phyloscanner transmission tree output suggests direction of transmission, but in our

case, these results were often uncertain (see Results). To avoid putting undue emphasis on an

uncertain direction of transmission, we simplified the transmission tree output to undirected,

unweighted (binary) networks and performed statistical analysis of these networks using

exponential random graph models (ERGMs; [38]). ERGMs model the edges in networks, with

explanatory variables representing the potential structural drivers of the observed network [38].

By including network structural variables, ERGMs account for the inherent non-independence of

network data. As such, we modeled "transmission relationships" (i.e., being connected in the

transmission network) as a function of network structural variables and transmission variables

we a priori expected to influence direct transmission processes in panthers. We considered

several structural variables: an intercept-like edges term [38]; geometrically weighted edgewise

shared partner distribution (*gwesp*; representation of network triangles); alternating k-stars

(*altkstar*; representation of star structures); and 2-paths (2 step paths from *i* to *k* via *j*; [39]). In

addition, we considered a suite of transmission variables (see supplementary methods for

additional variable details): panther sex; age class (subadult or adult); pairwise genetic

relatedness (panther microsatellite data from [34]); position of panther home range centroid

(95% minimum convex polygon) or capture location (hereafter, *centroid*) relative to the major I-

75 freeway (locations could be north or south of this east-west freeway); distance from centroid

to nearest urban area (in km; USA Urban Areas layer, ArcGIS; [40]); pairwise geographic

distance between centroids (log-transformed; Figure S1); and pairwise home range overlap

(utilization distribution overlap indices of 95% bivariate normal home range kernels; [41,42]).

Because ERGMs are prone to degeneracy with increasing complexity [38], we first

performed forward selection for network structural variables, followed by forward selection of

dyad-independent variables, while controlling for network structure. Model selection was based

on AIC and goodness of fit, and MCMC diagnostics were assessed for the final model

(supplementary methods). ERGMs were fit with the *ergm* package [43] in R (v3.6.3; [44]).

Panther population simulations

We next simulated FeLV transmission through a network representing panthers during

the 2002-04 FeLV outbreak where network edges represented likely transmission pathways

based on ERGM-identified predictors of FIV transmission (*FIV-based model*)*.* Hereafter, a *full-*

simulation includes both simulation of the panther population with its likely transmission

pathways and simulation of FeLV transmission within that population. Below, we describe the

process for a single simulation, but these procedures were repeated for each full simulation.

We first based the simulated population size on the range of empirical estimates from

2002-2004 (Table S1; [36]). Additional characteristics of the simulated population included those

identified as significant variables in the ERGM analysis: age category and pairwise geographic

distances between panther home range centroids (see Results). We randomly assigned age

categories to the simulated population based on the proportion of adults versus subadults. Age

proportions were based on age distributions in the western United States [45], which

qualitatively align with the historically elevated mean age of the Florida panther population [46].

Pairwise geographic distances for the simulated population were generated by randomly

assigning simulated home range centroids based on the distribution of observed centroids on

the landscape (supplementary methods).

We then used ERGM coefficients to generate network edges among the simulated

panther population using the *ergm* package in R [43]. The FIV transmission network spanned 15

years of observations and represents a subset of the actual contact network, as it includes only

those interactions that resulted in successful transmission. We therefore had a high degree of

uncertainty regarding the appropriate network density for our simulations. To manage this

uncertainty, we constrained density (ratio of existing edges to all possible edges) in our network

simulations across a range of parameter space (net_dens, Table S1).

Simulation of FeLV transmission on FIV-based networks

The next step in each full simulation was to model FeLV transmission through the

network generated from FIV predictors of transmission. FeLV transmission was based on a

stochastic chain binomial process on the simulated network, following a modified SIR

compartmental model (Figure 2). Simulations were initiated with one randomly selected

infectious individual and proceeded in weekly time steps. Transmission simulations lasted until

no infectious individuals remained or until 2.5 years, whichever came first.

Figure 2: *Diagram of flows of individuals between compartments in transmission model. Virus*

icons indicate infectious states, with the regressive infection icon darkened to represent reduced

or uncertain infectiousness of this class. Note: a vaccination process was also included in the

transmission model, but is not shown for simplicity. With vaccination, susceptibles could be

vaccinated, and vaccinated individuals subsequently infected as with susceptibles, but with an

additional probability of (1-ve). See Table S1 for definitions of parameters.

Transmission was dependent on the following (Figure 2; see Table S1 for parameter

definitions): (1) existence of an edge between two individuals, (2) the dyad in question involving

a susceptible and infectious individual, and (3) a random binomial draw based on the probability

of transmission given contact (β). In addition, *Puma concolor* generally have low expected

weekly contact rates [47]; we therefore included a weekly contact probability, represented as a

random binomial draw for contact in a given week (ω).

Upon successful transmission, infectious individuals were randomly assigned to one of

three outcomes of FeLV infection [29]. *Progressive* infections (probability P) are infectious (β),

develop clinical disease, and die due to infection (μ). *Regressive* infections (also probability P)

recover from infection ($K^*\mu$, where K is a constant) and, having entered a state of viral latency,

are not considered at risk of FeLV reinfection [29,48]. Using model assumptions derived from

known patterns of FeLV infection in domestic cats, regressive individuals are not infectious [29],

but given ongoing uncertainty, we included some transmission from regressives ($C^*\beta$, where C

is a constant). *Abortive* infections (probability 1-2P) are never infectious, clearing infection and

joining the recovered class. While the duration of immunity in abortive cases has not been

studied in panthers, because abortive cases clear infection through a strong immune response

and develop anti-FeLV antibodies, reinfection with FeLV is considered extremely unlikely [48].

A vaccination process was included in simulations as panthers were vaccinated against

FeLV during the historical FeLV outbreak starting in 2003. Vaccination occurred at a rate, τ, and

applied to the whole population, as wildlife managers are unlikely to know if a panther is

susceptible at the time of capture or darting. However, only susceptible individuals transitioned

to the vaccinated class (i.e. vaccination failed in non-susceptibles). Because panthers were

vaccinated in the empirical outbreak with a domestic cat vaccine with unknown efficacy in

panthers, we allowed vaccinated individuals to become infected in transmission simulations by

including a binomial probability for vaccine failure (1-vaccine efficacy, *ve*, Table S1).

The panther population size remained roughly static through the course of the FeLV

outbreak [36]. We therefore elected not to include background mortality, but did include

infection-induced mortality. To maintain a consistent population size, we therefore included a

birth/recruitment process. Because FIV-based simulated networks drew edges based on

population characteristics, we treated births as a "respawning" process, in which territories

vacated due to mortality were reoccupied by a new susceptible at rate, v. This approach allowed

us to maintain the ERGM-based network structure and is biologically reasonable, as vacated

panther territories are unlikely to remain unoccupied for long.

Comparison of simulation predictions to observed FeLV outbreak

To evaluate the performance of our FIV-based model, we also predicted FeLV

transmission dynamics using three alternative models: random networks, home range overlap-

based networks, and a well-mixed model. The random networks model used Erdős-Rényi

random networks, matching network densities from the FIV-based model (Table S1), but

otherwise allowing edges to occur between any pairs of individuals. Overlap-based networks

were generated using the degree distributions of panther home range overlap networks from

2002-2004 and simulated annealing with the R package *statnet* ([49]; supplementary methods).

For both random and overlap-based networks, FeLV transmission was simulated as in the FIV-

based simulations. The well-mixed model was a stochastic, continuous time compartmental

model (Gillespie algorithm), with rate functions aligning with the chain binomial FeLV

transmission probabilities (see supplementary methods).

We performed transmission simulations for all *model types* (FIV-based, overlap-based,

random, and well-mixed) across a range of reasonable parameter space (Table S1), using a

Latin hypercube design (LHS) to generate 150 parameter sets that efficiently sampled

parameter space [50,51]. For each parameter set and model type, we performed 50 simulations

(30,000 total). In each simulation, we recorded the number of mortalities and the duration of

outbreaks, which were each summarized (medians) across each parameter set. For each model

type, we determined if each parameter set's predicted median mortalities, duration of outbreaks,

and abortive cases were within a reasonable range based on the observed FeLV outbreak (5-20

mortalities, 78-117 week duration, at least 5 abortive infections; [29]). If so, a parameter set was

deemed "feasible" for that model type. Ranges were used to account for uncertainty in

observations and population size in this cryptic species (supplementary methods). To test for

differences in the frequency of feasible FeLV predictions between model types, we fit a binomial

generalized linear mixed model (GLMM), assuming a logistic regression with "feasible" (vs

"unfeasible") as the outcome, model type as a predictor variable, and a random intercept for

parameter set.

We tested for spatial clustering of cases in the observed FeLV outbreak by leveraging

our database of qPCR-based FeLV status. We performed a local spatial clustering analysis of

FeLV cases and controls using SaTScan (50% maximum, circular window; [52]). A SaTScan

analysis seeks to identify clusters of cases in which the observed cases within a particular

cluster exceed random expectation; this analysis reports the observed/expected ratio and radius

of any significant clusters. In addition, we performed a global cluster analysis with Cuzick and

Edward's test (global cluster detection with case-control data) in the R package *smacpod* (1, 3,

5, 7, 9, and 11 nearest neighbors; 999 iterations; [53–55]). To determine if simulated FeLV

cases likewise demonstrated spatial clustering, we repeated SaTScan local cluster analysis and

Cuzick and Edward's tests (at 3, 5, and 7 nearest neighbors) with feasible FIV-based simulation

results. To determine if detected clustering in FIV-based simulations was simply based on our

respawning protocol, we also performed both spatial analyses with feasible overlap-based

simulation results as a "negative control." Because the overlap-based model was not spatially

explicit, we assigned the same geographic locations to nodes in the overlap-based networks

from the corresponding FIV-based networks.

303   To determine if certain transmission parameters were important for feasible outcomes,

304 we performed post hoc random forest variable importance analyses for each of the four model

305 types with "feasible" as the binary response variable (using the R package randomForest

306 [56,57]; see supplementary results).

308 **Results**

309 *FIV transmission network analysis*

310   In the main FIV network, Phyloscanner inferred 42 potential transmission relationships

311 (edges) between 19 individuals (nodes; network density = 0.25), after removing 9 edges that

312 were between individuals known not to be alive at the same time (Figure 3). Panther FIV

313 genomes missing from the transmission network were those for whom transmission

314 relationships could not be inferred by Phyloscanner (see Discussion). ERGM results for the

315 main FIV network identified triangle (*gwesp*) and star structures (*altkstar*) as key structural

316 variables, and age category and log transformed pairwise geographic distance as key

317 transmission variables (Tables 1, S2). Though altkstar was not statistically significant, inclusion

318 of this variable contributed to improved AIC and goodness of fit outcomes. Adults were more

319 likely to be involved in transmission events (but see discussion of sample size limitations) and

320 inferred transmission events were more likely between individuals which were geographically

321 closer to each other. The fitted model showed reasonable goodness of fit (Figures S2-3).

Figure 3: *Phyloscanner-derived main FIV transmission network. Node shape indicates panther*

age class (square = subadult; circle = adult). Node color indicates panther sex (blue = male;

red = female). Edge weight represents Phyloscanner tree support for each edge (thicker edge =

increased support); for visualization purposes, edges are displayed as the inverse of the

absolute value of the log of these support values. While pictured as a directed and weighted

network, statistical analyses used binary, undirected networks.

Table 1: Main FIV transmission network exponential random graph model results

Variable	Estimate	SE	*p*-value
Edges (intercept)	-2.56	1.33	0.055
gwesp	0.98	0.26	<0.001
altkstar	-0.70	0.96	0.47
Age (Adult)	0.93	0.44	0.03
Log pairwise distance	-0.45	0.21	0.03

Note: "gwesp" is geometrically weighted edgewise shared partner distribution (a representation

of triangle structures) and "altkstar" is alternating k-stars (a representation of star structures).

Age classes were subadult and adult, with subadults the reference level; pairwise distances

were between home range centroids and log-transformed. Only those variables from the final

model are shown. Estimates shown are untransformed; SE represents standard error; p-*values*

less than 0.05 were considered statistically significant.

FeLV simulations

About 9% of parameter sets across all model types were classified as feasible (Figures

S6-S7). The FIV-based model had the highest odds of feasibility, though this difference did not

achieve statistical significance (Table 2). SaTScan analysis of observed FeLV status found

weak evidence of local spatial clustering (two clusters detected, but not statistically significant

with *p*=0.165 and 0.997, respectively; Figure S5). Cuzick and Edward's tests found evidence of

global clustering at 3, 5, and 7 nearest neighbor levels (test statistic T_k where *k* is number of

nearest neighbors considered: $T_3 = 20$, *p* = 0.049; $T_5 = 32$, *p* = 0.028; $T_7 = 43$, *p* = 0.023).

Feasible parameter sets from both the FIV-based and overlap-based models produced some

evidence of local and global spatial clustering of simulated FeLV cases (Figures 4, S8).

However, the FIV-based model better captured the size and strength of predicted local clusters

(SaTScan radius and observed/expected cases, respectively; Figure 4) and was moderately

better at capturing global spatial patterns (Figure S8).

The post-hoc random forest analyses typically showed poor balanced accuracy and area

under the curve (AUC) results. However, the parameter shaping transmission from regressively

infected individuals (C), consistently showed support for weak to moderate transmission from

regressives (i.e., C = 0.1 or 0.5; Figure S11).

Table 2: Fixed effects results from model-type performance GLMM*

Variable	Estimate	SE	*p*-value
Intercept	0.055	0.40	<0.001
FIV-based network model	1.55	0.42	0.30
Random network model	1.32	0.43	0.52
Overlap-based network model	1.21	0.44	0.66

*Note: estimates provided are exponentiated; the well-mixed model was the reference group

and none of the model-type results achieved statistical significance.

Figure 4: *SaTScan cluster analysis for feasible FIV-based and overlap-based network*

simulations show stronger agreement for the FIV-based model, compared to the overlap-based

model, between empirical observations (red horizontal lines) and model predictions for (A) FeLV

cluster size and (B) Observed/Expected FeLV cases associated with the top detected cluster.

Shown are feasible simulation results in which at least one cluster was detected with p-*values*

less than or equal to 0.1; further, only the results from the top cluster are shown.

Discussion

In this study we develop a new approach whereby we leverage knowledge of

transmission dynamics of a common apathogenic agent to prospectively predict dynamics of an

uncommon and virulent pathogen. Our approach was distinctly different from simpler models we

tested, as the apathogenic (FIV)-based approach focused on underlying drivers or mechanisms

of transmission and could be used to prospectively identify predictors of transmission and

develop disease control plans prior to an outbreak of a virulent pathogen (FeLV). We found that

FIV transmission in panthers is primarily driven by distance between home range centroids and

age class, and that our prospective FIV-based approach predicted FeLV transmission dynamics

at least as well as simpler or more reactive approaches. While we do not propose that this

apathogenic agent approach could accurately predict exactly when, where, and to whom

transmission might occur, our results support the role of apathogenic agents as novel tools for

prospectively determining sources of individual-level heterogeneity in transmission and

consequently improving proactive disease management.

An FIV-based model captures FeLV transmission dynamics

We found that our network model based on drivers of FIV transmission produced FeLV

outbreak predictions consistent with the observed FeLV outbreak. The FIV-based approach

performed at least as well as simpler models, per our GLMM analysis, with evidence that FIV

better predicted the observed spatial dynamics for FeLV transmission. A key difference between

the FIV-based approach and other spatially explicit methods is that FIV allowed us to determine

the importance of spatial dynamics prospectively and then translate to predictions of FeLV

transmission, rather than relying on retrospective FeLV spatial analyses. Furthermore, while

more complex potential drivers of transmission (e.g., host relatedness or assortative mixing by

age or sex) were not found to be important for FIV transmission, these may yet be key for

structuring transmission in other systems. Simpler model types like random networks or

metapopulation models may struggle to make transmission predictions that incorporate these

drivers of transmission-relevant contact. The predictive power we observed here using an

apathogenic virus could thus open new opportunities to determine behavioral and ecological

drivers of individual-level heterogeneity in the context of pathogen transmission, and even

shape proactive epidemic management strategies for pathogens such as FeLV.

Our network statistical analysis (ERGMs) determined that pairwise geographic distances

and age category structure FIV transmission in the Florida panther. These findings are well

supported by panther and FIV biology, providing confidence in the functioning of our

apathogenic agent workflow. For example, panthers are wide-ranging animals but maintain

home ranges, and this appears to translate to increased transmission between individuals that

are close geographically. This finding is supported by the tendency for FIV phylogenies to show

distinct broad [58] and fine scale [59] geographic clustering in *Puma concolor*. Further,

specifically among Florida panthers, spatial autocorrelation of FIV exposure status was

previously found to approach statistical significance [60]. The wide-ranging nature of puma

appears to limit geographic clustering of many infectious agents [60], with FIV a notable

exception to this pattern. In addition, because FIV is a persistent infection, we would expect

cumulative risk of transmission to increase over an individual's lifetime and adults would

consequently be involved in more transmission events. The low number of subadult individuals

in our dataset, however, means that this finding must be interpreted with some caution.

With these ERGM results in mind, key components of the success of our apathogenic

agent approach are likely that (1) FIV is a largely species-specific virus with transmission

pathways closely matching intraspecific transmission of FeLV, and (2) both FIV and FeLV,

perhaps unusually for infectious agents of puma, display spatial clustering of infection. Here,

FIV fundamentally acted as a proxy for close, direct contact in panthers, and could consequently

determine drivers of such contacts. If, for example, FIV also exhibited strong vertical or

environmental transmission, we would no longer expect the predictive success for FeLV we

observed here. This consideration highlights the importance of careful apathogenic agent

selection when attempting to predict pathogen transmission, a workflow which must now be

tested in additional host-pathogen systems. For example, the mixed results when using

commensal agents to identify close social relationships in other systems [15–18,21,22]

highlights that some host-apathogenic agent combinations will work better than others for

determining drivers of transmission. Within our study, Phyloscanner struggled to elucidate

transmission relationships between many of our FIV genomes, likely due to unusually low

genetic diversity among our FIV isolates, or our use of proviral DNA (which has lower diversity

than circulating RNA) [35]. While the drivers of transmission we identified are biologically

reasonable, we may have lacked the power to identify more complex relationships (e.g.,

homophily) due to the low number of individuals in our transmission network.

Elaborating on agent selection suggestions outlined in the introduction, we propose that

apathogenic agent selection should carefully consider agent genetic diversity within a target

population—not just expected diversity based on typical mutation rates, as in our case—and

favor those agents with high diversity to facilitate transmission inference. We also propose that

apathogenic agents should represent the timescales of transmission for the pathogen of

interest. For example, FeLV produces long, slow epidemics, such that FIV transmission

relationships may be most representative across the longer timescales we evaluated here. In

contrast, short, acute pathogen epidemics would likely best be represented by apathogenic

agent transmission over shorter timescales. Our results highlight that, perhaps most importantly,

an apathogenic agent should have a well characterized mode of transmission that closely

matches transmission of the pathogen of interest, as this was likely key to our success with FIV

and FeLV. Future research should determine how divergent an apathogenic agent may be from

a pathogen of interest while still predicting transmission dynamics.

While few parameter sets in our simulations were classified as feasible, this appears to

be predominantly the result of the wide range of parameter space explored through our LHS

sampling design. This limitation was fundamentally due to uncertainties in FeLV transmission

parameters, and is representative of the uncertainties experienced in predicting transmission of

emerging or understudied pathogens. Infectiousness of regressives, the number of introductions

of FeLV to the panther population, and the duration of FeLV infection in panthers were all

important sources of uncertainty in our models. All three of these dynamics can have significant

impacts on the duration of a simulated epidemic, allowing an epidemic to continue to "stutter"

along at low levels [61], much as was observed in the empirical FeLV outbreak. Our *post hoc*

random forest analysis provided some evidence of weak transmission from regressive

individuals, but this finding would need to be validated with additional research, as it is in stark

contrast to FeLV dynamics in domestic cats. Reducing uncertainties in these three key

dynamics would significantly narrow the range of our predictions, and even assist in ongoing

management efforts for FeLV in endangered panthers. The effect of transmission parameter

uncertainties underscores the importance of linking laboratory and model-based research to

generate more accurate transmission forecasts [62].

Limitations and future directions

The suite of tools for inferring transmission networks from infectious agent genomes is

rapidly expanding [24]. In this study, we used the program Phyloscanner as it maximized the

information from our deep sequencing viral data. However, our FIV sequences were generated

within a tiled amplicon framework [35,63], which biases intrahost diversity and limits viral

haplotypes [64]. Phyloscanner was originally designed to analyze RNA from virions and not

proviral DNA, as we have done here. We have attempted to mitigate the effects of these

limitations by analyzing several different Phyloscanner outputs to confirm consistency in our

results, and by using only binary networks to avoid putting undue emphasis on transmission

network edge probabilities, as these are likely uncertain. Further, our primary conclusions from

the transmission networks—that age and pairwise distance are important for transmission—are

biologically plausible and supported by other literature, as discussed above. Nevertheless,

future work should evaluate additional or alternative transmission network inference platforms.

In addition, ERGMs assume the presence of the "full network" and it is as yet unclear

how missing data may affect transmission inferences [38]. ERGMs are also prone to

degeneracy with increased complexity and do not easily capture uncertainty in transmission

events, as most weighted network ERGM (or generalized ERGM) approaches have been

tailored for count data (e.g., [65]). ERGMs may therefore not be the ideal solution for identifying

drivers of transmission networks in all systems. Alternatives may include advancing dyad-based

modeling strategies [66], which may more easily manage weighted networks and instances of

missing data.

Our FIV-based approach required extensive field sampling, and many disciplines from

viral genomics through simulation modeling. However, with increasing availability of virome data

and even field-based sequencing technology, our approach may become more accessible with

time. Further, the predictive benefits seen here, while needing further testing and validation,

could become a key strategy for proactive pathogen management in species of conservation

concern, populations of high economic value (e.g. production animals), populations with

infrequent pathogen outbreaks that make targeted surveillance more difficult, or populations at

high risk of spillover, all of which may most benefit from rapid, efficient epidemic responses.

Conclusions

Here, we integrated genomic and network approaches to identify drivers of FIV

transmission in the Florida panther. This apathogenic agent acted as a marker of close, direct

contact transmission, and was subsequently successful in predicting the observed transmission

dynamics of the related pathogen, FeLV. Further testing of apathogenic agents as markers of

transmission and their ability to predict transmission of related pathogens is needed, but holds

promise for revolutionizing proactive epidemic management across host-pathogen systems.

Acknowledgements

Thanks to M. Michalska-Smith, K. Worsley-Tonks, J. Mistrick, and S. N. Hart for key feedback.

This research was supported by the National Science Foundation (DEB-1413925, 1654609, and

2030509). MLJG was supported by the Office of the Director, National Institutes of Health (NIH

T32OD010993), the University of Minnesota Informatics Institute MnDRIVE program, and the

Van Sloun Foundation. JLM was supported by the ACVP/STP Coalition for Veterinary Pathology

Fellows and the Linda Munson Fellowship for Wildlife Pathology Research. The content is solely

the responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health. Puma icon by Freepik at Flaticon.com.

Data Availability

Data and R code for replication of simulations and analyses is available on GitHub

(https://github.com/mjones029/FIV_FeLV_Transmission) and upon acceptance will be archived

at Zenodo.

Works Cited

1. Anderson RM, May RM. 1991 *Infectious Diseases of Humans: Dynamics and Control*. OUP Oxford.

2. Metcalf CJE, Lessler J. 2017 Opportunities and challenges in modeling emerging infectious diseases. *Science* **357**, 149–152.

3. Drewe JA. 2010 Who infects whom? Social networks and tuberculosis transmission in wild meerkats. *Proc. Biol. Sci.* **277**, 633–642.

4. Morris M, Kretzschmar M. 1995 Concurrent partnerships and transmission dynamics in networks. *Soc. Networks* **17**, 299–318.

5. Cauchemez S, Bhattarai A, Marchbanks TL, Fagan RP, Ostroff S, Ferguson NM, Swerdlow D, Pennsylvania H1N1 working group. 2011 Role of social networks in shaping disease transmission during a community outbreak of 2009 H1N1 pandemic influenza. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 2825–2830.

6.   Craft ME, Caillaud D. 2011 Network models: an underutilized tool in wildlife epidemiology?
Interdiscip. Perspect. Infect. Dis. **2011**, 676949.

7.   Keeling MJ, Eames KTD. 2005 Networks and epidemic models. *J. R. Soc. Interface* **2**, 295–
307.

8.   Keeling MJ, Woolhouse MEJ, May RM, Davies G, Grenfell BT. 2003 Modelling vaccination
strategies against foot-and-mouth disease. *Nature* **421**, 136–142.

9.   Lloyd-Smith JO, George D, Pepin KM, Pitzer VE, Pulliam JRC, Dobson AP, Hudson PJ,
Grenfell BT. 2009 Epidemic dynamics at the human-animal interface. *Science* **326**, 1362–
1367.

10. Dobson AP *et al.* 2020 Ecology and economics for pandemic prevention. *Science* **369**,
379–381.

11. Knight-Jones TJD, Rushton J. 2013 The economic impacts of foot and mouth disease--
What are they, how big are they and where do they occur? *Prev. Vet. Med.* **112**, 161–173.

12. Williams ES, Thorne ET, Appel MJ, Belitsky DW. 1988 Canine distemper in black-footed
ferrets (*Mustela nigripes*) from Wyoming. *J. Wildl. Dis.* **24**, 385–398.

13. Roelke-Parker ME *et al.* 1996 A canine distemper virus epidemic in Serengeti lions
(*Panthera leo*). *Nature* **379**, 441–445.

14. Sillero-Zubiri C, King AA, Macdonald DW. 1996 Rabies and mortality in Ethiopian wolves
(*Canis simensis*). *J. Wildl. Dis.* **32**, 80–86.

15. Bull CM, Godfrey SS, Gordon DM. 2012 Social networks and the spread of Salmonella in a
sleepy lizard population. *Mol. Ecol.* **21**, 4386–4392.

16. Blasse A, Calvignac-Spencer S, Merkel K, Goffe AS, Boesch C, Mundry R, Leendertz FH.
2013 Mother-offspring transmission and age-dependent accumulation of simian foamy virus
in wild chimpanzees. *J. Virol.* **87**, 5193–5204.

17. Chiyo PI, Grieneisen LE, Wittemyer G, Moss CJ, Lee PC, Douglas-Hamilton I, Archie EA.
2014 The influence of social structure, habitat, and host traits on the transmission of
Escherichia coli in wild elephants. *PLoS One* **9**, e93408.

18. Blyton MDJ, Banks SC, Peakall R, Gordon DM. 2013 High temporal variability in
commensal *Escherichia coli* strain communities of a herbivorous marsupial. *Environ.*
Microbiol. **15**, 2162–2172.

19. Lax S *et al.* 2014 Longitudinal analysis of microbial interaction between humans and the
indoor environment. *Science* **345**, 1048–1052.

20. Song SJ *et al.* 2013 Cohabiting family members share microbiota with one another and with
their dogs. *Elife* **2**, e00458.

21. VanderWaal KL, Atwill ER, Isbell LA, McCowan B. 2014 Linking social and pathogen
transmission networks using microbial genetics in giraffe (*Giraffa camelopardalis*). *J. Anim.*
Ecol. **83**, 406–414.

22. Springer A, Mellmann A, Fichtel C, Kappeler PM. 2016 Social structure and Escherichia coli
sharing in a group-living wild primate, Verreaux's sifaka. *BMC Ecol.* **16**, 6.

23. Archie EA, Tung J. 2015 Social behavior and the microbiome. *Current Opinion in
Behavioral Sciences* **6**, 28–34.

24. Hall MD, Woolhouse MEJ, Rambaut A. 2016 Using genomics data to reconstruct
transmission trees during disease outbreaks. *Rev. Sci. Tech.* **35**, 287–296.

25. Gilbertson MLJ, Fountain-Jones NM, Craft ME. 2018 Incorporating genomic methods into
contact networks to reveal new insights into animal behaviour and infectious disease
dynamics. *Behaviour* **155**, 759–791.

26. Blyton MDJ, Banks SC, Peakall R, Lindenmayer DB, Gordon DM. 2014 Not all types of host
contacts are equal when it comes to E. coli transmission. *Ecol. Lett.* **17**, 970–978.

27. Archie EA, Luikart G, Ezenwa VO. 2009 Infecting epidemiology with genetics: a new
frontier in disease ecology. *Trends Ecol. Evol.* **24**, 21–30.

28. Carver S *et al.* 2016 Pathogen exposure varies widely among sympatric populations of wild
and domestic felids across the United States. *Ecol. Appl.* **26**, 367–381.

29. Cunningham MW *et al.* 2008 Epizootiology and management of feline leukemia virus in the
Florida puma. *J. Wildl. Dis.* **44**, 537–552.

30. Krakoff E, Gagne RB, VandeWoude S, Carver S. 2019 Variation in Intra-individual Lentiviral
Evolution Rates: a Systematic Review of Human, Nonhuman Primate, and Felid Species. *J.
Virol.* **93**. (doi:10.1128/JVI.00538-19)

31. Brown MA, Cunningham MW, Roca AL, Troyer JL, Johnson WE, O'Brien SJ. 2008 Genetic
characterization of feline leukemia virus from Florida panthers. *Emerg. Infect. Dis.* **14**, 252–
259.

32. Greene CE. 2012 *Infectious diseases of the dog and cat*. 4th ed.. St. Louis, Mo.:
Elsevier/Saunders.

33. Hartmann K. 2012 Clinical aspects of feline retroviruses: a review. *Viruses* **4**, 2684–2710.

34. Van De Kerk M, Onorato DP, Hostetler JA, Bolker BM, Oli MK. 2019 Dynamics,
persistence, and genetic management of the endangered florida panther population.
Wildlife Monogr. **203**, 3–35.

35. Malmberg JL *et al.* 2019 Altered lentiviral infection dynamics follow genetic rescue of the
Florida panther. *Proc. Biol. Sci.* **286**, 20191689.

36. McClintock BT, Onorato DP, Martin J. 2015 Endangered Florida panther population size
determined from public reports of motor vehicle collision mortalities. *J. Appl. Ecol.* **52**, 893–
901.

37. Wymant C *et al.* 2018 PHYLOSCANNER: Inferring Transmission from Within- and
Between-Host Pathogen Genetic Diversity. *Mol. Biol. Evol.* **35**, 719–733.

38. Silk MJ, Fisher DN. 2017 Understanding animal social structure: exponential random graph

models in animal behaviour research. *Anim. Behav.* **132**, 137–146.

39. Morris M, Handcock MS, Hunter DR. 2008 Specification of Exponential-Family Random
Graph Models: Terms and Computational Aspects. *J. Stat. Softw.* **24**, 1548–7660.

40. Esri. USA Urban Areas (FeatureServer). See
https://services.arcgis.com/P3ePLMYs2RVChkJx/arcgis/rest/services/USA_Urban_Areas/F
eatureServer.

41. Fieberg J, Kochanny CO, Lanham. 2005 Quantifying home-range overlap: the importance
of the utilization distribution. *J. Wildl. Manage.* **69**, 1346–1359.

42. Calenge C. 2006 The package adehabitat for the R software: tool for the analysis of space
and habitat use by animals. *Ecological Modelling.* **197**, 1035.

43. Hunter DR, Handcock MS, Butts CT, Goodreau SM, Morris M. 2008 ergm: A Package to
Fit, Simulate and Diagnose Exponential-Family Models for Networks. *J. Stat. Softw.* **24**,
nihpa54860.

44. R Core Team. 2018 R: A Language and Environment for Statistical Computing.

45. Logan KA, Sweanor LL. 2001 *Desert Puma: Evolutionary Ecology And Conservation Of An*
Enduring Carnivore. Island Press.

46. Johnson WE *et al.* 2010 Genetic restoration of the Florida panther. *Science* **329**, 1641–
1645.

47. Elbroch LM, Quigley H. 2016 Social interactions in a solitary carnivore. *Curr. Zool.* **63**, 357–
362.

48. Hofmann-Lehmann R, Hartmann K. 2020 Feline leukaemia virus infection: A practical
approach to diagnosis. *J. Feline Med. Surg.* **22**, 831–846.

49. Handcock MS, Hunter DR, Butts CT, Goodreau SM, Morris M. 2008 statnet: Software tools
for the representation, visualization, analysis and simulation of network data. *J. Stat. Softw.*
24, 1548.

50. Marino S, Hogue IB, Ray CJ, Kirschner DE. 2008 A methodology for performing global
uncertainty and sensitivity analysis in systems biology. *J. Theor. Biol.* **254**, 178–196.

51. Carnell R. 2012 lhs: Latin hypercube samples. *R package version 0.10, URL http://CRAN.*
R-project.org/package=lhs

52. Kulldorff M. 1997 A spatial scan statistic. *Communications in Statistics - Theory and*
Methods **26**, 1481–1496.

53. Cuzick J, Edwards R. 1990 Spatial clustering for inhomogeneous populations. *J. R. Stat.*
Soc. **52**, 73–96.

54. French J. 2020 smacpod: Statistical Methods for the Analysis of Case-Control Point Data.

55. Kanankege KST, Alvarez J, Zhang L, Perez AM. 2020 An Introductory Framework for
Choosing Spatiotemporal Analytical Tools in Population-Level Eco-Epidemiological

638   Research. *Front Vet Sci* **7**, 339.

639 56. Liaw A, Wiener M. 2002 Classification and Regression by randomForest. *R News*. **2**, 18–
640   22.

641 57. White LA, VandeWoude S, Craft ME. 2020 A mechanistic, stigmergy model of territory
642   formation in solitary animals: Territorial behavior can dampen disease prevalence but
643   increase persistence. *PLoS Comput. Biol.* **16**, e1007457.

644 58. Lee JS *et al.* 2014 Evolution of puma lentivirus in bobcats (*Lynx rufus*) and mountain lions
645   (*Puma concolor*) in North America. *J. Virol.* **88**, 7727–7737.

646 59. Fountain-Jones NM *et al.* 2021 Host relatedness and landscape connectivity shape
647   pathogen spread in the puma, a large secretive carnivore. *Commun Biol* **4**, 12.

648 60. Gilbertson MLJ, Carver S, VandeWoude S, Crooks KR, Lappin MR, Craft ME. 2016 Is
649   pathogen exposure spatially autocorrelated? Patterns of pathogens in puma (*Puma
650   concolor*) and bobcat (*Lynx rufus*). *Ecosphere* **7**, e01558.

651 61. Blumberg S, Lloyd-Smith JO. 2013 Inference of R(0) and transmission heterogeneity from
652   the size distribution of stuttering chains. *PLoS Comput. Biol.* **9**, e1002993.

653 62. Plowright RK, Sokolow SH, Gorman ME, Daszak P, Foley JE. 2008 Causal inference in
654   disease ecology: investigating ecological drivers of disease emergence. *Front. Ecol.
655   Environ.* **6**, 420–429.

656 63. Quick J *et al.* 2017 Multiplex PCR method for MinION and Illumina sequencing of Zika and
657   other virus genomes directly from clinical samples. *Nat. Protoc.* **12**, 1261–1276.

658 64. Grubaugh ND *et al.* 2019 An amplicon-based sequencing framework for accurately
659   measuring intrahost virus diversity using PrimalSeq and iVar. *Genome Biol.* **20**, 8.

660 65. Krivitsky PN. 2012 Exponential-family random graph models for valued networks. *Electron.
661   J. Stat.* **6**, 1100–1128.

662 66. Yang A *et al.* 2021 Effects of social structure and management on risk of disease
663   establishment in wild pigs. *J. Anim. Ecol.* **90**, 820-833. (doi:10.1111/1365-2656.13412)