CORTICAL MECHANISMS OF SACCADIC SUPPRESSION

Sachira Denagamage[1,2], Mitchell P. Morton[1,2], John H. Reynolds[4], Monika P. Jadi[1,2,3,*] and

Anirvan S. Nandy[1,2,*]

[1]Department of Neuroscience, Yale University, New Haven, CT 06511

[2]Interdepartmental Neuroscience Program, Yale University, New Haven, CT 06511

[3]Department of Psychiatry, Yale University, New Haven, CT 06511

[4]Systems Neurobiology Laboratories, The Salk Institute for Biological Studies, La Jolla, CA 92037

[*]co-senior and corresponding authors

Correspondence: monika.jadi@yale.edu, anirvan.nandy@yale.edu

ABSTRACT

Saccades are a ubiquitous and crucial component of our visual system, allowing for the efficient

deployment of the fovea and its accompanying neural resources. Initiation of a saccade is known

to cause saccadic suppression, a temporary reduction in visual sensitivity[1,2] and visual cortical

firing rates[3-6]. While saccadic suppression has been well characterized at the level of perception

and single neurons, relatively little is known about the visual cortical networks governing this

phenomenon. Here we examine the effects of saccadic suppression on distinct neural

subpopulations within area V4. We find cortical layer- and cell type-specific differences in the

magnitude and timing of peri-saccadic modulation. Neurons in the input layer show changes in

firing rate and inter-neuronal correlations prior to saccade onset, suggesting that this layer receives

information about impending saccades. Putative inhibitory interneurons in the input layer elevate

their firing rate during saccades and may play a role in suppressing the activity of other cortical

subpopulations. A computational model of this circuit recapitulates our empirical observations and

demonstrates that an input layer-targeting pathway could initiate saccadic suppression by

enhancing local inhibitory activity. Together, our results provide a mechanistic understanding of

how eye movement signaling interacts with cortical circuitry to enforce visual stability.

INTRODUCTION

As we observe the world around us, our eyes dart from point to point. Each of these shifts in gaze

is a saccade – a ballistic movement of both eyes. Saccades significantly improve the efficiency of

the primate visual system by allowing us to flexibly deploy our fovea, the dedicated high acuity

zone at the center of the retina, towards objects of interest in our environment. However, saccades

also pose a substantial challenge for ongoing visual processing, as each saccade produces abrupt

and rapid motion across the retina. Therefore, saccades are also accompanied by a temporary

reduction in visual sensitivity, termed saccadic suppression, that serves to blunt our perception of

this motion[1,2]. Perhaps reflecting the diminished sensory processing during saccades, the firing

rates of neurons in many regions of the visual cortex are also transiently suppressed[3-6]. Yet, despite

extensive psychophysical characterization and single neuron electrophysiological analysis,

relatively little is known about the circuit level mechanisms underlying saccadic suppression in

the visual cortex. One prevailing hypothesis suggests that saccadic suppression is mediated by a

corollary discharge signal originating in the brain regions responsible for initiating eye

movements[7-9]. Given the synaptic and motor delays accompanying the execution of the saccadic

motor command, a coincident corollary discharge to the visual cortex could trigger compensatory

mechanisms prior to the start of the eye movement. Consistent with this idea, changes in neural

activity before saccade onset have been observed in several visual cortical areas[5]. However, in

spite of clear neural evidence that the visual cortex receives information about upcoming saccades,

the pathway by which a saccadic signal arrives in the visual cortex is unknown, and how that signal

interacts with local circuitry to mediate saccadic suppression remains unclear.

We examined these questions within area V4, a critical hub in the visual processing stream

responsible for object recognition[10]. V4 is known to exhibit saccadic suppression in both humans[4]

and macaques[6], and undergoes dynamic shifts in stimulus sensitivity during the preparation of

saccades[11]. To investigate the underlying neural mechanisms, we studied saccadic suppression in

the context of the laminar cortical circuit, which is composed of six layers with highly stereotyped

patterns of intra- and inter-laminar connectivity[12,13]. In higher-order visual areas, such as V4, layer

IV (the *input* layer) is the primary target of projections carrying visual information from lower-

order areas, such as V1, V2, and V3[14,15]. After arriving in the input layer, visual information is

then processed by local neural subpopulations as it is sent to layers II/III (the *superficial* layer) and

layers V/VI (the *deep* layer), which serve as output nodes in the laminar circuit[12,16]. In V4, the

superficial and deep layers feed information forward to upstream visual areas, such as ST[17], TEO[18],

and TE[19]. Alongside feedforward input from lower-order visual areas, V4 also receives projections

from other cortical regions, such as the prefrontal cortex[15], as well as from subcortical structures[20].

Accordingly, several pathways targeting V4 could be responsible for relaying information about

upcoming saccades: 1) projections from lower visual areas (V1/V2/V3), which predominantly

target the input layer[14,15], 2) a projection from the frontal eye fields (FEF), which predominantly

targets the superficial and deep layers[21,22], and 3) a projection from the pulvinar, which

predominantly targets the superficial and input layers[23-25]. Neurons in V1[26,27], FEF[28-30], and

pulvinar[31-33] are known to respond to saccades, and the corresponding pathways are therefore

strong candidates for initiating saccadic suppression in V4.

Given the organization of cortical circuitry, we reasoned that a saccadic signaling pathway

should target the site of entry for visual information, the input layer, to most efficiently suppress

visual processing. Furthermore, in light of the similarities between peri-saccadic stimulus

processing and general reductions in visual gain[6,26], we considered whether saccadic suppression

might be mediated in a manner comparable to visual gain modulation. Specifically, we

hypothesized that an elevation in local inhibitory activity, a common mechanism for gain control[34-

36], functionally suppresses neural processing in V4 during saccades. We sought to confirm our

predictions by utilizing a cortical layer-specific recording approach in combination with

electrophysiological cell type classification to simultaneously record from six well-defined neural

subpopulations in V4.

RESULTS

Dissociating the Effects of Vision and Eye Movements

Past approaches for studying saccadic modulation have typically involved cued saccade

tasks[4,6,26,37], which risk conflating the neural effects of visual stimulus processing with the neural

effects of saccades. To dissociate saccadic signaling from visually induced activity in area V4, we

performed a series of spontaneous recordings while two rhesus macaques were seated in the dark.

We found that despite their inability to observe specific objects in the environment, both animals

continued to make saccades freely under these conditions. We identified individual saccades by

estimating the onset and offset times of ballistic eye movements (see Methods). In order to remove

fixational eye movements (microsaccades), we discarded saccades with amplitudes less than 0.7

degrees of visual angle. Additionally, to prevent the effects of one eye movement from overlapping

with those of the next, we only included saccades that were separated from one another by at least

500 ms. This approach provided us with a set of 4420 saccades that were used for all subsequent

analyses. We evaluated the quality of our eye movement dataset by examining amplitude and

velocity traces from individual saccades, as well as by characterizing the main sequence. The main

sequence is the approximately linear amplitude-velocity relationship observed across all

saccades[38], which we found held true for our data (Figure 1A).

While the monkeys were making spontaneous saccades, we simultaneously recorded

neural activity from well-isolated single-units (n = 211), multi-units (n = 110), and local field

potentials (LFPs) using linear array probes in area V4. The use of recording chambers containing

an optically-transparent artificial dura allowed us to make electrode penetrations perpendicular to

the cortical surface, and therefore in good alignment with individual cortical columns (see

Methods). The alignment of each penetration was confirmed by mapping receptive fields along

the depth of cortex, with overlapping receptive fields indicating proper alignment (Figure 1B).

Laminar boundaries were identified with current source density (CSD) analysis[39,40]. The CSD,

defined as the second spatial derivative of the LFP, maps the location of current sources and sinks

along the depth of the probe. Each layer of the visual cortex produces a characteristic current

source-sink pattern in response to visual stimulation, with the input layer producing a current sink

followed by a current source, and the superficial and deep layers producing the opposite pattern of

activity (Figure 1C). By identifying transitions between current sources and sinks, we assigned

laminar identities to each of our recording sites (Figure 1D – LFP traces colored by layer). We

also separated well-isolated single-units into two populations on the basis of their waveform

duration: narrow-spiking units, corresponding to putative inhibitory interneurons, and broad-

spiking units, corresponding to putative excitatory neurons (Figure 1D – spikes colored by unit

type)[41-43]. The three cortical layers and two unit types thus gave us six distinct neural

subpopulations to consider in subsequent analyses.

Narrow-Spiking Input Layer Units Show Positive Peri-Saccadic Modulation

To evaluate neural subpopulation-specific responses, we first estimated the peri-saccadic firing

rate of well-isolated single-units with a kernel-based approach, in which each spike is convolved

with a gaussian kernel[44] (Figure S1). We selected the kernel bandwidth on a unit-by-unit basis at

each point in time using an optimization algorithm that minimizes mean integrated squared error[45].

When looking at units grouped by cell type, narrow-spiking units exhibit less peri-saccadic

suppression than broad-spiking units (Figure 2A). The cause of this difference became clear when

the units were further separated by layer. All laminar subpopulations of broad-spiking units

displayed peri-saccadic suppression, as did narrow-spiking units in the superficial and deep layers.

In contrast, narrow-spiking input layer units elevated their firing rate in response to a saccade

(Figure 2B-C; see Figure S2 for additional single-unit examples).

To quantify these observations, we calculated a modulation index that represents the degree

to which the firing rate of each unit deviated from its pre-saccadic baseline activity (see Methods).

We found that narrow-spiking input layer units were the only group with a positive modulation

index, which was significantly different from most other subpopulations (Figure 2D). We also

noted that at the single-unit level, each of the six subpopulations had some proportion of positively

modulated units (Figure S3). Therefore, the difference in subpopulation level firing rates could be

caused by narrow-spiking input layer neurons having a greater average magnitude of positive

modulation, or by having a greater proportion of positively modulated units. We determined both

to be true. Narrow-spiking input layer neurons displayed a larger magnitude of positive modulation

(Figure 2E), and were also more likely to be positively modulated (Figure 2F). We found it

particularly striking that, in accordance with our hypothesis, only narrow-spiking input layer units

(i.e. putative inhibitory interneurons) showed an enhancement in firing. This led us to believe that

an elevation in input layer inhibitory activity could be responsible for the peri-saccadic suppression

of firing within the visual cortex reported previously[3-6].

Input Layer Units Respond Prior to Saccades

In order to gate out visual signals received during a saccade, neurons in the cortical layer initiating

suppression would need to be activated prior to the start of the eye movement. To confirm that this

was true for the input layer, we performed a timing analysis to determine whether any of the

recorded subpopulations show significant changes in firing rate prior to saccade onset. For each

single-unit, we found the time at which their activity first deviated outside a 95% confidence

interval calculated from their pre-saccadic baseline activity, an event that was marked as the time

of first significant response. The results of this approach for six example units (the same example

units as in Figure S1) are illustrated in Figure 3A. While there was response variability at the level

of single-units, the input layer subpopulations (both narrow- and broad-spiking) were the only ones

that responded prior to saccade onset (Figure 3B). This provides strong evidence that the input

layer receives information about upcoming saccades with sufficient time to enact changes in

activity throughout the cortical column.

Saccades Increase Correlated Variability in the Input Layer

If input layer neurons receive pre-saccadic excitation from a common source, that signaling should

also manifest itself as an increase in correlated variability among pairs of input layer units prior to

saccade onset. To test this, we computed spike count correlations (SCC) between pairs of

simultaneously recorded single- and multi-units as a function of time, and pooled these results into

six groups on the basis of their pairwise laminar locations. All pair types showed a substantial

increase in SCC around the time of saccades (Figure S4A). We quantified the precise time at which

correlations first began to rise with a bootstrapping analysis (see Methods). SCC began increasing

prior to saccade onset in input-input pairs (Figure 4A), but after saccade onset in all other pairs

(Figure 4B). This pattern of activity is consistent with the pre-saccadic arrival of a signal that is

shared among input layer units. The delayed response of all other pair types suggests that they

inherit their activity from the input layer. Additionally, the precise time at which input-input

correlations began to rise (~40 ms before saccade onset) is remarkably similar to the time at which

input layer units began to show firing rate modulation (also ~40 ms before saccade onset), further

indicating that these two observations have a common cause: the activation of a saccadic signaling

pathway.

Alongside the expected rise in correlations, we also found a reduction in correlations that

began ~250 ms prior to saccade onset among input-input pairs. Because the timing of this drop

precedes saccadic initiation, which begins ~ 150 ms before saccade onset[46,47], it could not be the

result of a corollary discharge signal. Instead, it may reflect the shift in internal state that occurs

before saccadic initiation as the subject shifts their attention towards a potential target location[48].

In accordance with this possibility, pupil diameter, a well-established correlate of internal state[49,50],

begins to ramp several hundred milliseconds prior to saccade onset (Figure S4B)[51]. Similar shifts

in internal state are known to be accompanied by comparable reductions in correlated

variability[50,52-54].

To further characterize changes in coordinated peri-saccadic activity, we also examined

differences in spike-spike coherence (SSC) before and after saccade onset. Spike-spike coherence

is a measure of the degree to which the activity of two signals, in this case a pair of simultaneously

recorded single- and/or multi-units, fluctuates together across a range of frequencies. We found

that saccade onset increases wide-band coherence most prominently in the input layer, with

weaker, but still significant, effects in the superficial and deep layers (Figure 4C). When quantified

as a modulation index (Figure 4D), we observed significant modulation in all layers at all

frequencies, with the exception of the 30-100 Hz band in the superficial layer. The input layer

exhibits larger increases in coherence, particularly at higher frequencies, while the deep layer

experiences smaller increases, particularly at lower frequencies. Collectively, these data suggest

that saccade onset leads to a temporary elevation of correlated activity, which manifests itself most

prominently in the input layer. The timing and magnitude of changes in correlated variability,

measured with both SCC and SSC, further suggests that an external signal arrives prior to saccade

onset within the input layer before then propagating to the other layers.

Saccades Increase the Strength of Low-Frequency Wide-Band Activity

Increases in coordinated activity at the level of single neurons can result in an elevation of

population-level wide-band activity. To determine whether this held true for saccades, we

calculated the spectral power, a frequency-resolved measure of signal strength, of the LFP around

the time of saccade onset. Saccade onset was associated with a large elevation in low frequency

power, and a slight reduction in high frequency power (Figure 5A). Comparing the power spectrum

before and after saccade onset, we found that these differences were significant in all three

frequency bands tested, with an elevation in the 0-12 Hz and 15-25 Hz bands and a reduction in

the 30-80 Hz band after saccade onset (Figure 5B). Next, we sought to investigate whether the

increased strength of low-frequency wide-band activity entrains spiking units, as reported

previously in other contexts[55-57]. We found that saccade onset significantly increases low-

frequency spike-LFP phase locking (Figure 5C), as measured by pairwise phase consistency

(PPC). These results indicate that saccade onset increases the strength of population-level low-

frequency wide-band activity, which could in turn entrain the activity of local neurons.

A Computational Model Demonstrates that Activation of an Input Layer-Targeting Projection is Sufficient for Inducing Saccadic Suppression

Given that input layer units exhibit significant changes in firing properties prior to saccade onset,

we hypothesized that a projection targeting this layer could serve as the initiator of saccadic

suppression within area V4. To determine whether such a projection would be sufficient for

initiating suppression across all cortical layers, as well as to gain further insight into the associated

circuit-level mechanisms, we developed a computational model of peri-saccadic activity changes

within a cortical column. Our model draws inspiration from previous cortical circuit models that

describe population-specific firing rate changes in response to the activation of an external input[58].

To more faithfully simulate cortical circuit-level dynamics, our model consists of six

interconnected populations across three layers of the cortex (Figure 6A, left). Each cortical layer

contains an excitatory (E) and inhibitory (I) population, which project to each other as well as to

themselves. Given evidence of multiple operating regimes for a local E-I network in the visual

cortex, connections within a layer were tuned to allow the local E-I network to switch in and out

of an inhibition-stabilized network (ISN) regime[59,60]: the baseline regime of the network is non-

ISN, but direct external input to the E population switches it to ISN (see Methods; Figure S5A-C).

Connections between layers exist in the form of projections from E neurons in the source layer to

both E and I neurons in the target layer, with E-E and E-I connections tuned to affect a net increase

in E activity in the target layer in response to an increase in E activity in the source layer. In our

model, the input layer projects to the superficial layer, which then projects to the deep layer; this

connectivity motif highlights the primary pathway for information flow within a visual cortical

column[13,61]. In light of our empirical firing rate analysis, which found that both broad- and narrow-

spiking units in the input layer show responses prior to saccade onset (Figure 3B), we chose to

model the input layer as the primary recipient of the 'saccadic signal' input. We tuned the strength

of the excitatory 'saccadic signal' pathway to the local E and I populations in this layer such that

its activation affected a net suppression of the E activity.

In response to the activation of a saccadic signal of sufficient strength slightly before the

time of saccade onset, the E input layer population rate was suppressed. In addition, our model

simultaneously recapitulated our experimental findings in the input layer I population

(enhancement), as well as the E/I populations in other layers (suppression) by virtue of the inter-

and intra-layer connectivity (Figure 6B, S5B). Since local E activity is a major source of excitation

to the I population within a layer, we next explored the role of the saccadic signal in sustaining

input layer I activity despite a concurrent decline in local E activity. We explored three scenarios

with saccadic signals of varying magnitudes ramping up and down over a fixed period (Figure 6C,

top). We found that the increase in I activity in the input layer indeed depended on the strength of

the saccadic signal and was sustained at sufficiently high magnitudes. Examination of the

contribution of different excitatory pathways to the I population showed that sufficiently high

saccadic signal magnitude led to an increase in net excitation onto the I population that exceeded

the loss in excitation resulting from the reduction of local E firing rates (Figure 6C, middle and

bottom). Our model thus demonstrates that an input layer-specific saccadic signal of adequate

strength can increase local I activity, which is sufficient for inducing suppression across the depth

of the cortex. These findings correspond closely to our empirical observations. The model further

predicted that the time course of the increase in input layer I activity was dependent on the baseline

operating regime of the E-I network. When the network was robustly within the non-ISN regime,

I activity increased immediately following saccadic signal onset. However, when the network was

close to the switching point between the two regimes, the I activity showed little change below a

saccadic signal of sufficient magnitude, even when the local E activity showed significant

suppression (Figure S5E). This latter scenario corresponds closely to our empirical observations

in the input layer (Figure 2B-C), suggesting that the baseline state of the cortical network lies close

to the boundary between regimes.

Our experimental paradigm examined peri-saccadic cortical dynamics in the absence of

visual input. While this allowed us to investigate the cortical dynamics that are purely a result of

the saccadic signal and avoid potential stimulus-induced artifacts in our recordings, it leaves open

the question of how activity patterns might change in the presence of visual stimulation. To explore

this further, we added a second external excitatory input ('visual signal') to the input layer E

population (Figure 6A, right), which is the major target of feedforward visual input into V4[14]. The

addition of this second input shifted the local E-I network into an ISN regime[59], in which E and I

activity increase or decrease concurrently. As a result, simultaneous activation of the visual signal

led to a net reduction in the firing rate of both E and I input layer populations (Figure 6D, S5C).

This produced a similar reduction in excitatory drive to the superficial and deep layers, where

firing was again suppressed. These results are consistent with prior reports of suppression from

uncategorized neurons in the visual cortex in the presence of visual stimulation[3,5,6].

DISCUSSION

Our results provide evidence that the input layer of area V4 receives the earliest information about upcoming saccades. Units in the input layer showed changes in firing rate prior to saccade onset, while units in other layers did not. In addition, elevated spike-count correlations, a common byproduct of activating a shared input to a neural population[62-65], were observed earliest among pairs of input layer neurons. Consistent with a single input source (i.e. projection) producing both phenomena, changes in the firing rate and correlational structure of input layer units were initiated with similar timing, approximately 40 ms before saccade onset. These convergent results indicate that the saccadic signaling pathway in question predominantly targets the input layer, which thereby allows us to infer its source. Given the lack of pre-saccadic modulation in the superficial and deep layers, it is unlikely that a projection from the FEF is responsible for the observed suppression. Likewise, it is doubtful that suppression is fed forward by projections from lower visual areas (V1/V2/V3), as such a mechanism would not explain the enhancement of narrow-spiking input layer firing rates in V4. Furthermore, to our knowledge, there have been no clear reports of saccadic suppression in V2 or V3, and while V1 is known to be suppressed during saccades, it does not exhibit pre-saccadic modulation[26], and could therefore not be responsible for the changes in activity described here. Thus, we propose that saccadic suppression in V4 is likely initiated via thalamo-cortical projections from the pulvinar.

Several lines of research are consistent with this view. Peri-saccadic modulation has been observed in the pulvinar of macaques, where neurons respond to saccades made in the dark without visual stimulation[32,33]. Neurons in the pulvinar of cats are able to distinguish between internally generated saccades and simulated saccades produced with image motion, demonstrating that they encode the underlying motor command[31]. Furthermore, the pulvinar is known to play a crucial role

in spatial attention[66-68], which is necessary for the proper execution of saccades towards target

stimuli; indeed, temporary inactivation of the pulvinar results in significant deficits in saccadic

target selection and execution[69,70]. These findings indicate that neurons in the pulvinar would be

able to provide a meaningful saccadic signal to their efferent targets.

We also found that narrow-spiking units in the input layer showed a peri-saccadic

enhancement in firing. Narrow-spiking units are thought to correspond to parvalbumin-expressing

inhibitory interneurons[71,72], and are known to receive strong thalamocortical excitation.[73-75] This

led us to speculate that an elevated level of inhibition within the input layer could suppress local

excitatory neurons, as well as neurons in other cortical layers (Figure 6A). To explore this

hypothesis, we developed a six-population firing rate model of the visual cortex and its modulation

during saccades. The model demonstrated that pre-saccadic activation of an input layer-targeting

projection is sufficient for suppressing signaling within a cortical column. A compelling aspect of

our model is its simplicity – with straightforward inter-population connectivity rules, we were able

to both replicate our empirical findings as well as reconcile the opposing patterns of activity

observed in the input layer subpopulations. The model also reproduces trends in our experimental

observations on the relative time course of broad and narrow spiking activity under a set of

conditions wherein the input layer operates on the cusp of ISN and non-ISN operating regimes.

Given these results, we propose that inhibitory input layer neurons initiate saccadic suppression

and gate the processing of visual information entering the local cortical network.

We also observed increases in correlated activity across all cortical layers during saccades,

in the form of elevated spike-count correlations, spike-spike coherence, low-frequency power, and

low frequency spike-phase locking. Comparable increases in neural correlation have been

associated with diminished stimulus sensitivity in multiple model organisms across a variety of

sensory modalities[50,76-79]. This may be a result of heightened correlations functionally limiting the

ability of a neural population to encode information, as suggested by previous experimental and

theoretical work[80-83]. Similarly, saccadic suppression at the level of neurons and circuits is also

accompanied by a behaviorally significant reduction in stimulus detection capabilities[1,2],

indicating that the increases in correlated variability reported here may reflect or contribute to that

deficit.

In summary, we conclude that an input layer-targeting projection, likely originating in the

pulvinar, initiates saccadic suppression within V4 by enhancing the firing of local inhibitory

interneurons. Future work could extend this line of subpopulation-specific, circuit-level analysis

to examine activity changes during saccades in the presence of visual stimulation. Our model

provides a prediction for such experiments, suggesting that the peri-saccadic activity of inhibitory

input layer neurons would also be suppressed when the cortex is in a state of active information

processing. The absence of any reports of subpopulation-specific differences in peri-saccadic

modulation during previous studies employing cued saccade tasks is consistent with this view.

METHODS

Surgical Procedures

Surgical procedures have been described in detail previously[84,85]. In brief, an MRI-compatible, low-profile titanium chamber was placed over the pre-lunate gyrus on the basis of preoperative MRI imaging in two rhesus macaques (right hemisphere in monkey A, left hemisphere in monkey C). The native dura mater was then removed, and a silicone-based, optically clear artificial dura (AD) was inserted, resulting in an optical window over dorsal V4. All procedures were approved by the Institutional Animal Care and Use Committee and conformed to NIH guidelines.

Electrophysiology

At the beginning of each recording session, a plastic insert with an opening for targeting electrodes was lowered into the chamber and secured. This served to stabilize the recording site against cardiac pulsations. Neurons were recorded from cortical columns in dorsal V4 using 16-channel linear array electrodes (''laminar probes''; Plexon, Plexon V-probe). The laminar probes were mounted on adjustable x-y stages attached to the recording chamber and positioned over the center of the pre-lunate gyrus under visual guidance through a microscope (Zeiss). This ensured that the probes were maximally perpendicular to the surface of the cortex and thus had the best possible trajectory to make a normal penetration down a cortical column. Across recording sessions, the probes were positioned over different sites along the center of the gyrus in the parafoveal region of V4 with receptive field eccentricities between 2 and 7 degrees of visual angle. Care was taken to target cortical sites with no surface micro-vasculature and, in fact, the surface micro-vasculature was used as reference so that the same cortical site was not targeted across recording sessions. The probes were advanced using a hydraulic microdrive (Narishige) to first

penetrate the AD and then through the cortex under microscopic visual guidance. Probes were

advanced until the point that the top-most electrode (toward the pial surface) registered LFP

signals. At this point, the probe was retracted by about 100–200 mm to ease the dimpling of the

cortex due to the penetration. This procedure greatly increased the stability of the recordings and

also increased the neuronal yield in the superficial electrodes.

The distance from the tip of the probes to the first electrode contact was either 300 mm or

700 mm. The inter-electrode distance was 150 mm, thus negating the possibility of recording the

same neural spikes in adjacent recording channels. Neuronal signals were recorded extra-

cellularly, filtered, and stored using the Multi-channel Acquisition Processor system (Plexon).

Neuronal signals were classified as either multi-unit clusters or isolated single-units using the

Plexon Offline Sorter program. Single-units were identified based on two criteria: (1) if they

formed an identifiable cluster, separate from noise and other units, when projected into the

principal components of waveforms recorded on that electrode and (2) if the inter-spike interval

(ISI) distribution had a well-defined refractory period. Single-units were classified as either

narrow-spiking (putative interneurons) or broad-spiking (putative pyramidal cells) based on

methods described in detail previously[41]. Specifically, only units with waveforms having a clearly

defined peak preceded by a trough were potential candidates. Units with trough-to-peak duration

less than 225 ms were classified as narrow-spiking units; units with trough- to-peak duration

greater than 225 ms were classified as broad-spiking units.

Data were collected over 17 sessions (8 sessions in monkey A and 9 in monkey C), yielding

a total of 211 single-units (99 narrow-spiking, 112 broad-spiking) and 110 multi-unit clusters.

Recording

Stimuli were presented on a computer monitor placed 57 cm from the eye. Eye position

was continuously monitored with an infrared eye tracking system (ISCAN ETL-200). Trials were

aborted if eye position deviated more than 1 degree of visual angle from fixation. Experimental

control was handled by NIMH Cortex software (http://www.cortex.salk.edu/).

Receptive Field Mapping

At the beginning of each recording session, neuronal RFs were mapped using subspace

reverse correlation in which Gabor (eight orientations, 80% luminance contrast, spatial frequency

1.2 cycles/degree, Gaussian half-width 2 degrees) or ring (80% luminance contrast) stimuli

appeared at 60 Hz while monkeys fixated. Each stimulus appeared at a random location selected

from an 11 x 11 grid with 1 degree spacing in the appropriate visual quadrant. Spatial receptive

maps were obtained by applying reverse correlation to the evoked LFP signal at each recording

site. For each spatial location in the 11 x 11 grid, we calculated the time-averaged power in the

stimulus-evoked LFP (0–200 ms after each stimulus flash) at each recording site. The resulting

spatial map of LFP power was taken as the spatial RF at the recording site. For the purpose of

visualization, the spatial RF maps were smoothed using spline interpolation and displayed as

stacked contours plots of the smoothed maps (Figure 1B). All RFs were in the lower visual

quadrant (lower left in monkey A and lower right in monkey C) and with eccentricities between 2

and 7 degrees of visual angle.

Current Source Density Mapping

In order to estimate the laminar identity of each recording channel, we used a CSD mapping

procedure[39]. Monkeys maintained fixation while 100% luminance contrast ring stimuli were

flashed (30 ms), centered at the estimated RF overlap region across all channels. The size of the

ring was scaled to about three-quarters of the estimated diameter of the RF. CSD was calculated

as the second spatial derivative of the flash-triggered LFPs. The resulting time-varying traces of current across the cortical layers can be visualized as CSD maps (Figure 1C). Red regions depict current sinks in the corresponding region of the cortical laminae; blue regions depict current sources. The input layer (Layer 4) was identified as the first current sink followed by a reversal to current source. The superficial (Layers 1–3) and deep (Layers 5 and 6) layers had opposite sink-source patterns. LFPs and spikes from the corresponding recording channels were then assigned to one of three layers: superficial, input, or deep.

Spontaneous Recordings

In the main experiment, all monitors and lights were turned off in the recording room and adjacent control room to ensure that the environment was as dark as experimentally feasible. Eye tracker and electrophysiological data were recorded while the animals executed eye-movements freely in the dark.

Data Analysis

Saccade Identification

To identify saccade onset and offset times from the raw eye-tracker data, we used the Cluster Fix algorithm[86]. Cluster Fix performs k-means clustering on four parameters (distance, velocity, acceleration, and angular velocity) to find natural partitions in the eye-tracker data and identify saccades. To ensure that our set of identified eye movements was 'clean,' we imposed several additional criteria: 1) to avoid adjacent saccades affecting our analysis, we only considered saccades separated from neighboring saccades by at least 500 ms, 2) to exclude microsaccades, we only considered saccades that were greater in amplitude than 0.7 degrees of visual angle, 3) to exclude slower eye movements that Cluster Fix had mislabeled as saccades, we only considered eye movements less than 200 ms in duration, and 4) to exclude periods of data erroneously labeled

as saccades due to blinking or drowsiness, we identified those periods using pupil diameter

measurements and excluded nearby saccades.

Saccadic eye movements are known to have an approximately linear relationship between

amplitude and peak velocity that is referred to as the main sequence[38]. To ensure that our approach

identified a set of eye movements that quantitatively resembled the properties of saccades, we

calculated the amplitude and peak velocity for each of the identified eye movements and found an

approximately linear relationship (Figure 1A).

Firing Rate Estimation and Modulation Index

To estimate the peri-saccadic firing rate, spikes were extracted for each saccade beginning

500 ms prior to saccade onset and ending 500 ms after saccade onset. To obtain a continuous

estimate of the firing rate, each spike was convolved with a gaussian kernel and the average firing

rate across saccades was calculated for each unit (Figure S1; Figure S2). The kernel bandwidth

was selected on a unit-by-unit basis at each point in time using a bandwidth optimization algorithm

that minimized the mean integrated squared error[45]. In order to obtain reliable firing rate estimates,

only units that fired at least 75 spikes within 500 ms of saccade onset (before or after) were

included.

A modulation index was calculated for each unit to estimate the degree to which their

activity deviated from baseline following saccade onset (Figure 2D, E; Figure S3). The modulation

index was defined as $FR_s/FR_b - FR_b$, where FR_s is the maximum firing rate in the 200 ms after

saccade onset and FR_b is the baseline firing rate of the unit, calculated by averaging the firing rate

for each unit in a time window 350 to 250 ms before saccade onset.

Spike Count Correlations

To obtain a continuous estimate of spike count correlations, we calculated the Pearson

correlation of spike counts across saccades at each time bin for every pair of simultaneously

recorded single- and multi-units (Figure 4A, Figure S4A). The time window was 201 ms in width,

and was shifted by 1 ms steps. Spike counts were z-scored to control for firing rate differences

across time. Traces were smoothed with a 101 ms Gaussian weighted moving average. To perform

the timing analysis, we took the local minimum in correlation as the time at which the rise begins.

Bootstrapping was performed for each pair type to obtain confidence intervals on the estimate

(Figure 4B).

Spike-Spike Coherence

We computed the coherence between simultaneously recorded, multi-unit pairs within a

cortical layer using multi-taper methods[87] for two different windows: the 200 ms before and after

saccade onset (Figure 4C). Spike trains were tapered with a single Slepian taper (TW = 3, K = 5).

Magnitude of coherence estimates depend on the number of spikes used to create the estimates[88].

To control for differences in firing rate across the two windows, we adopted a rate-matching

procedure[53]. In order to obtain a baseline for the coherence expected solely due to trends in firing

time-locked to saccade onset, we also computed coherence in which saccade identities were

randomly shuffled. Only units that elicited at least 50 spikes from each neuron cumulatively across

all saccades in both time windows were considered. A modulation index, defined as

$(SSC_{after} - SSC_{before})/(SSC_{after} + SSC_{before})$, was calculated for each pair of units at each

frequency band (Figure 4D).

Power

Local field potential power was calculated with multi-tapered methods[87]. Signals were

tapered with a single Slepian taper (TW = 3, K = 5). Spectrograms (Figure 5A) were generated by

sliding a 201 ms window by increments of 1 ms. Power spectra before (-200 to 0 ms) and after (0

to 200 ms) saccade onset were calculated with static variants of the same methods (Figure 5B). No

layer-specific differences were found, so results shown here are averaged across all recording

channels.

Pairwise Phase Consistency

The pairwise phase consistency (PPC) is an estimate of spike-field locking that is not

biased by spike count or firing rate[89]. Two segments of data were analyzed: the 200 ms before and

after saccade onset. In order to obtain reliable estimates, only units that fired at least 50 spikes

cumulatively across all saccades during both time windows were considered. The spectro-temporal

representation of the local field potential signal was generated through a continuous wavelet

transform with a family of complex Morlet wavelets (spanning frequencies from 2 to 100 Hz, with

6 cycles at each frequency). Phase information was extracted at the time of each spike, and the

PPC was then calculated for each unit with each LFP channel in the laminar probes (Figure 5C).

We observed minimal change in the PPC as a function of LFP layer, so the results presented here

were averaged across all recording channels.

Computational Model

The firing rate or mean field model consists of a network of six populations representing

excitatory (E) and inhibitory (I) populations in the superficial (L1), input (L2), and deep (L3) layers

of the cortex. The firing rate model describes temporal changes in the firing rates (r) of each

population as a function of the activity of two external inputs (i_{sacc}, the saccadic signal, and i_{stim},

the visual signal) and the activity of other populations in the network. The connectivity between

populations is depicted in Figure 6A. The network consists of a coupled E-I network in each layer

and captures the core aspect of connectivity between cortical layers: information flow from input

to superficial to deep layer. The saccadic signal is sent to both E and I populations in the input

layer, while the visual signal is sent only to the E population in the input layer. The saccadic signal

is described by a ramping function (Figure 6C, top). The visual signal is only activated in the

second set of simulations, where it is held constant to replicate sustained visual input. The six-

population rate model is described by the following equations:

498
$$\tau_E \frac{dr_{E_{L1}}}{dt} = -r_{E_{L1}} + G_E\left(W_{EE} \cdot r_{E_{L1}} - W_{EI} \cdot r_{I_{L1}} + W_{EE_{L12}} \cdot r_{E_{L2}}\right) \tag{1}$$

499
$$\tau_I \frac{dr_{I_{L1}}}{dt} = -r_{I_{L1}} + G_I\left(W_{IE} \cdot r_{E_{L1}} - W_{II} \cdot r_{I_{L1}} + W_{IE_{L12}} \cdot r_{E_{L2}}\right) \tag{2}$$

500
$$\tau_E \frac{dr_{E_{L2}}}{dt} = -r_{E_{L2}} + G_E\left(W_{EE} \cdot r_{E_{L2}} - W_{EI} \cdot r_{I_{L2}} + W_{EE_{SS}} \cdot i_{sacc} + i_{stim}\right) \tag{3}$$

501
$$\tau_I \frac{dr_{I_{L2}}}{dt} = -r_{I_{L2}} + G_I\left(W_{IE} \cdot r_{E_{L2}} - W_{II} \cdot r_{I_{L2}} + W_{IE_{SS}} \cdot i_{sacc}\right) \tag{4}$$

502
$$\tau_E \frac{dr_{E_{L3}}}{dt} = -r_{E_{L3}} + G_E\left(W_{EE} \cdot r_{E_{L3}} - W_{EI} \cdot r_{I_{L3}} + W_{EE_{L32}} \cdot r_{E_{L1}}\right) \tag{5}$$

503
$$\tau_I \frac{dr_{I_{L3}}}{dt} = -r_{I_{L3}} + G_I\left(W_{IE} \cdot r_{E_{L3}} - W_{II} \cdot r_{I_{L3}} + W_{IE_{L32}} \cdot r_{E_{L1}}\right) \tag{6}$$

504
$$G(x) = \begin{cases} 0 & \text{for } x < \theta \\ m(x - \theta)^{2.5} & \text{for } \theta < x < \theta + 1/m \\ 1 & \text{for } x > \theta + 1/m \end{cases} \tag{7}$$

τ_E and τ_I are the rates at which the excitatory and inhibitory populations approach their steady

states. G_E and G_I are the population response functions, described by equation (7), for the

excitatory and inhibitory populations that transform the given inputs into a firing response. θ is

the threshold input and m is the rate of the response function. Parameters tuned in the model to

recapitulate the population level observations in experimental data: $W_{EE} = 25$, $W_{II} = 4$, $W_{EI} =$

$25, W_{IE} = 20, W_{EE_{L12}} = 20, W_{IE_{L12}} = 2, W_{EE_{SS}} = 0.5, W_{EI_{SS}} = 3, W_{EE_{L32}} = 30, W_{IE_{L32}} = 4, \tau_E =$

$0.008, \tau_I = 0.004, m_E = 0.001, m_I = 0.005, \theta_{E_{L1}} = -3, \theta_{E_{L2}} = -5, \theta_{E_{L3}} = -5, \theta_{I_{L1}} = 0,$

$\theta_{I_{L2}} = 8, \theta_{I_{L3}} = 8, i_{stim} = 10.$

A further tuning was applied to the model to switch each E-I population between an

inhibitory stabilized network (ISN) regime and a non-ISN one[60]. Experimental data from visual

cortex suggest that a cortical network switches to an ISN regime during stimulus processing[59],

hence we included this feature to generate model predictions for saccadic suppression-related

laminar activity during stimulus presentation. Previous modeling work has shown that a firing-

rate E-I network model can switch between these regimes either by modification of connection

weights (W), change in slope or rate of response functions (m) or a combination of both[60]. We

implemented the simplest mechanism for the network to switch between ISN and non-ISN

regime: nonlinear response functions. When visual stimulation (modeled as an excitatory input to

the input layer E population) is applied to the model cortical column, the network moves to a

higher slope region of its response functions, and switches to the ISN operating regime. A

phase-plane analysis of the E-I network model illustrates how this results in a distinct response

of the network to exactly the same saccadic signal input (Figure S5).

ACKNOWLEDGEMENTS

This research was supported by NIH R01 EY021827 to JHR and ASN, NARSAD Young

Investigator Grant, Ziegler Foundation Grant and Yale Orthwein Scholar Funds to ASN, NIH

R00 EY025026 to MPJ, and by NEI core grants for vision research P30 EY019005 to the Salk

Institute and P30 EY026878 to Yale University. We would like to thank Catherine Williams and

Mat LeBlanc for excellent animal care.

AUTHOR CONTRIBUTIONS

SD & ASN conceptualized the project. ASN collected the data and supervised the project. SD

analyzed the data, with assistance from MPM and ASN. MPJ developed the computational model.

SD, ASN, MPJ, and JHR wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

Matin, E. Saccadic suppression: a review and an analysis. *Psychol Bull* **81**, 899-917, doi:10.1037/h0037368 (1974).

Zuber, B. L. & Stark, L. Saccadic suppression: elevation of visual threshold associated with saccadic eye movements. *Exp Neurol* **16**, 65-79, doi:10.1016/0014-4886(66)90087-2 (1966).

Thiele, A., Henning, P., Kubischik, M. & Hoffmann, K. P. Neural mechanisms of saccadic suppression. *Science* **295**, 2460-2462, doi:10.1126/science.1068788 (2002).

Kleiser, R., Seitz, R. J. & Krekelberg, B. Neural correlates of saccadic suppression in humans. *Curr Biol* **14**, 386-390, doi:10.1016/j.cub.2004.02.036 (2004).

Bremmer, F., Kubischik, M., Hoffmann, K. P. & Krekelberg, B. Neural dynamics of saccadic suppression. *J Neurosci* **29**, 12374-12383, doi:10.1523/JNEUROSCI.2908-09.2009 (2009).

Zanos, T. P., Mineault, P. J., Guitton, D. & Pack, C. C. Mechanisms of Saccadic Suppression in Primate Cortical Area V4. *J Neurosci* **36**, 9227-9239, doi:10.1523/JNEUROSCI.1015-16.2016 (2016).

Wurtz, R. H., Joiner, W. M. & Berman, R. A. Neuronal mechanisms for visual stability: progress and problems. *Philos Trans R Soc Lond B Biol Sci* **366**, 492-503, doi:10.1098/rstb.2010.0186 (2011).

Sommer, M. A. & Wurtz, R. H. A pathway in primate brain for internal monitoring of movements. *Science* **296**, 1480-1482, doi:10.1126/science.1069590 (2002).

Sommer, M. A. & Wurtz, R. H. Influence of the thalamus on spatial visual processing in frontal cortex. *Nature* **444**, 374-377, doi:10.1038/nature05279 (2006).

Roe, A. W. *et al.* Toward a unified theory of visual area V4. *Neuron* **74**, 12-29, doi:10.1016/j.neuron.2012.03.011 (2012).

Han, X., Xian, S. X. & Moore, T. Dynamic sensitivity of area V4 neurons during saccade preparation. *Proc Natl Acad Sci U S A* **106**, 13046-13051, doi:10.1073/pnas.0902412106 (2009).

Hirsch, J. A. & Martinez, L. M. Laminar processing in the visual cortical column. *Curr Opin Neurobiol* **16**, 377-384, doi:10.1016/j.conb.2006.06.014 (2006).

Douglas, R. J. & Martin, K. A. Neuronal circuits of the neocortex. *Annu Rev Neurosci* **27**, 419-451, doi:10.1146/annurev.neuro.27.070203.144152 (2004).

Felleman, D. J. & Van Essen, D. C. in *Cereb cortex.*   (Citeseer).

15   Ungerleider, L. G., Galkin, T. W., Desimone, R. & Gattass, R. Cortical connections of area
V4 in the macaque. *Cereb Cortex* **18**, 477-499, doi:10.1093/cercor/bhm061 (2008).

16   Rockland, K. S. & Pandya, D. N. Laminar origins and terminations of cortical connections
of the occipital lobe in the rhesus monkey. *Brain Res* **179**, 3-20, doi:10.1016/0006-
8993(79)90485-2 (1979).

17   Boussaoud, D., Ungerleider, L. G. & Desimone, R. Pathways for motion analysis: cortical
connections of the medial superior temporal and fundus of the superior temporal visual
areas in the macaque. *J Comp Neurol* **296**, 462-495, doi:10.1002/cne.902960311 (1990).

18   Distler, C., Boussaoud, D., Desimone, R. & Ungerleider, L. G. Cortical connections of
inferior temporal area TEO in macaque monkeys. *J Comp Neurol* **334**, 125-150,
doi:10.1002/cne.903340111 (1993).

19   Borra, E., Ichinohe, N., Sato, T., Tanifuji, M. & Rockland, K. S. Cortical connections to
area TE in monkey: hybrid modular and distributed organization. *Cereb Cortex* **20**, 257-
270, doi:10.1093/cercor/bhp096 (2010).

20   Gattass, R., Galkin, T. W., Desimone, R. & Ungerleider, L. G. Subcortical connections of
area V4 in the macaque. *J Comp Neurol* **522**, 1941-1965, doi:10.1002/cne.23513 (2014).

21   Anderson, J. C., Kennedy, H. & Martin, K. A. Pathways of attention: synaptic relationships
of frontal eye field to V4, lateral intraparietal cortex, and area 46 in macaque monkey. *J
Neurosci* **31**, 10872-10881, doi:10.1523/JNEUROSCI.0622-11.2011 (2011).

22   Stanton, G. B., Bruce, C. J. & Goldberg, M. E. Topography of projections to posterior
cortical areas from the macaque frontal eye fields. *J Comp Neurol* **353**, 291-305,
doi:10.1002/cne.903530210 (1995).

23   Benevento, L. A. & Rezak, M. The cortical projections of the inferior pulvinar and adjacent
lateral pulvinar in the rhesus monkey (Macaca mulatta): an autoradiographic study. *Brain
Res* **108**, 1-24, doi:10.1016/0006-8993(76)90160-8 (1976).

24   Shipp, S. The functional logic of cortico-pulvinar connections. *Philos Trans R Soc Lond B
Biol Sci* **358**, 1605-1624, doi:10.1098/rstb.2002.1213 (2003).

25   Rockland, K. S. Distinctive Spatial and Laminar Organization of Single Axons from
Lateral Pulvinar in the Macaque. *Vision* **4**, 1 (2020).

26   McFarland, J. M., Bondy, A. G., Saunders, R. C., Cumming, B. G. & Butts, D. A. Saccadic
modulation of stimulus processing in primary visual cortex. *Nat Commun* **6**, 8110,
doi:10.1038/ncomms9110 (2015).

27   Wurtz, R. H. Response of striate cortex neurons to stimuli during rapid eye movements in
the monkey. *J Neurophysiol* **32**, 975-986, doi:10.1152/jn.1969.32.6.975 (1969).

28   Bizzi, E. Discharge of frontal eye field neurons during saccadic and following eye
609   movements in unanesthetized monkeys. *Exp Brain Res* **6**, 69-80, doi:10.1007/BF00235447
610   (1968).

611   29   Schall, J. D. & Hanes, D. P. Neural basis of saccade target selection in frontal eye field
612   during visual search. *Nature* **366**, 467-469, doi:10.1038/366467a0 (1993).

613   30   Everling, S. & Munoz, D. P. Neuronal correlates for preparatory set associated with pro-
614   saccades and anti-saccades in the primate frontal eye field. *J Neurosci* **20**, 387-400 (2000).

615   31   Robinson, D. L., McClurkin, J. W., Kertzman, C. & Petersen, S. E. Visual responses of
616   pulvinar and collicular neurons during eye movements of awake, trained macaques. *J*
617   *Neurophysiol* **66**, 485-496, doi:10.1152/jn.1991.66.2.485 (1991).

618   32   Berman, R. A. & Wurtz, R. H. Signals conveyed in the pulvinar pathway from superior
619   colliculus to cortical area MT. *J Neurosci* **31**, 373-384, doi:10.1523/JNEUROSCI.4738-
620   10.2011 (2011).

621   33   Sudkamp, S. & Schmidt, M. Response characteristics of neurons in the pulvinar of awake
622   cats to saccades and to visual stimulation. *Exp Brain Res* **133**, 209-218,
623   doi:10.1007/s002210000374 (2000).

624   34   Katzner, S., Busse, L. & Carandini, M. GABAA inhibition controls response gain in visual
625   cortex. *J Neurosci* **31**, 5931-5941, doi:10.1523/JNEUROSCI.5753-10.2011 (2011).

626   35   Wilson, N. R., Runyan, C. A., Wang, F. L. & Sur, M. Division and subtraction by distinct
627   cortical inhibitory networks in vivo. *Nature* **488**, 343-348, doi:10.1038/nature11347
628   (2012).

629   36   Mitchell, S. J. & Silver, R. A. Shunting inhibition modulates neuronal gain during synaptic
630   excitation. *Neuron* **38**, 433-445, doi:10.1016/s0896-6273(03)00200-9 (2003).

631   37   Berman, R. A., Cavanaugh, J., McAlonan, K. & Wurtz, R. H. A circuit for saccadic
632   suppression in the primate brain. *J Neurophysiol* **117**, 1720-1735,
633   doi:10.1152/jn.00679.2016 (2017).

634   38   Bahill, A. T., Clark, M. R. & Stark, L. The main sequence, a tool for studying human eye
635   movements. *Mathematical Biosciences* **24**, 191-204 (1975).

636   39   Mitzdorf, U. Current source-density method and application in cat cerebral cortex:
637   investigation of evoked potentials and EEG phenomena. *Physiol Rev* **65**, 37-100,
638   doi:10.1152/physrev.1985.65.1.37 (1985).

639   40   Nandy, A. S., Nassi, J. J. & Reynolds, J. H. Laminar Organization of Attentional
640   Modulation in Macaque Visual Area V4. *Neuron* **93**, 235-246,
641   doi:10.1016/j.neuron.2016.11.029 (2017).

41   Mitchell, J. F., Sundberg, K. A. & Reynolds, J. H. Differential attention-dependent
response modulation across cell classes in macaque visual area V4. *Neuron* **55**, 131-141,
doi:10.1016/j.neuron.2007.06.018 (2007).

42   Tamura, H., Kaneko, H., Kawasaki, K. & Fujita, I. Presumed inhibitory neurons in the
macaque inferior temporal cortex: visual response properties and functional interactions
with adjacent neurons. *J Neurophysiol* **91**, 2782-2796, doi:10.1152/jn.01267.2003 (2004).

43   Hasenstaub, A. *et al.* Inhibitory postsynaptic potentials carry synchronized frequency
information in active cortical networks. *Neuron* **47**, 423-435,
doi:10.1016/j.neuron.2005.06.016 (2005).

44   Kass, R. E., Ventura, V. & Cai, C. Statistical smoothing of neuronal data. *Network* **14**, 5-
15, doi:10.1088/0954-898x/14/1/301 (2003).

45   Shimazaki, H. & Shinomoto, S. Kernel bandwidth optimization in spike rate estimation. *J
Comput Neurosci* **29**, 171-182, doi:10.1007/s10827-009-0180-4 (2010).

46   Westheimer, G. Mechanism of saccadic eye movements. *AMA Arch Ophthalmol* **52**, 710-
724, doi:10.1001/archopht.1954.00920050716006 (1954).

47   Pare, M. & Munoz, D. P. Saccadic reaction time in the monkey: Advanced preparation of
oculomotor programs is primarily responsible for express saccade occurrence. *Journal of
Neurophysiology* **76**, 3666-3681 (1996).

48   Deubel, H. & Schneider, W. X. Saccade target selection and object recognition: evidence
for a common attentional mechanism. *Vision Res* **36**, 1827-1837, doi:10.1016/0042-
6989(95)00294-4 (1996).

49   Gilzenrat, M. S., Nieuwenhuis, S., Jepma, M. & Cohen, J. D. Pupil diameter tracks changes
in control state predicted by the adaptive gain theory of locus coeruleus function. *Cogn
Affect Behav Neurosci* **10**, 252-269, doi:10.3758/CABN.10.2.252 (2010).

50   Reimer, J. *et al.* Pupil fluctuations track fast switching of cortical states during quiet
wakefulness. *Neuron* **84**, 355-362, doi:10.1016/j.neuron.2014.09.033 (2014).

51   Jainta, S., Vernet, M., Yang, Q. & Kapoula, Z. The pupil reflects motor preparation for
saccades-even before the eye starts to move. *Frontiers in human neuroscience* **5**, 97 (2011).

52   Doiron, B., Litwin-Kumar, A., Rosenbaum, R., Ocker, G. K. & Josic, K. The mechanics
of state-dependent neural correlations. *Nat Neurosci* **19**, 383-393, doi:10.1038/nn.4242
(2016).

53   Mitchell, J. F., Sundberg, K. A. & Reynolds, J. H. Spatial attention decorrelates intrinsic
activity fluctuations in macaque area V4. *Neuron* **63**, 879-888,
doi:10.1016/j.neuron.2009.09.013 (2009).

54   Cohen, M. R. & Maunsell, J. H. Attention improves performance primarily by reducing
interneuronal correlations. *Nat Neurosci* **12**, 1594-1600, doi:10.1038/nn.2439 (2009).

55   Anastassiou, C. A., Perin, R., Markram, H. & Koch, C. Ephaptic coupling of cortical
neurons. *Nat Neurosci* **14**, 217-223, doi:10.1038/nn.2727 (2011).

56   Fries, P. Rhythms for Cognition: Communication through Coherence. *Neuron* **88**, 220-235,
doi:10.1016/j.neuron.2015.09.034 (2015).

57   Sirota, A. *et al.* Entrainment of neocortical neurons and gamma oscillations by the
hippocampal theta rhythm. *Neuron* **60**, 683-697, doi:10.1016/j.neuron.2008.09.014 (2008).

58   Veit, J., Hakim, R., Jadi, M. P., Sejnowski, T. J. & Adesnik, H. Cortical gamma band
synchronization through somatostatin interneurons. *Nature neuroscience* **20**, 951 (2017).

59   Ozeki, H., Finn, I. M., Schaffer, E. S., Miller, K. D. & Ferster, D. Inhibitory stabilization
of the cortical network underlies visual surround suppression. *Neuron* **62**, 578-592,
doi:10.1016/j.neuron.2009.03.028 (2009).

60   Tsodyks, M. V., Skaggs, W. E., Sejnowski, T. J. & McNaughton, B. L. Paradoxical effects
of external modulation of inhibitory interneurons. *J Neurosci* **17**, 4382-4388 (1997).

61   Douglas, R. J. & Martin, K. A. Mapping the matrix: the ways of neocortex. *Neuron* **56**,
226-238, doi:10.1016/j.neuron.2007.10.017 (2007).

62   Bair, W., Zohary, E. & Newsome, W. T. Correlated firing in macaque visual area MT: time
scales and relationship to behavior. *J Neurosci* **21**, 1676-1697 (2001).

63   Kohn, A. & Smith, M. A. Stimulus dependence of neuronal correlation in primary visual
cortex of the macaque. *J Neurosci* **25**, 3661-3673, doi:10.1523/JNEUROSCI.5106-
04.2005 (2005).

64   Shadlen, M. N. & Newsome, W. T. The variable discharge of cortical neurons: implications
for connectivity, computation, and information coding. *J Neurosci* **18**, 3870-3896 (1998).

65   Kriener, B., Tetzlaff, T., Aertsen, A., Diesmann, M. & Rotter, S. Correlations and
population dynamics in cortical networks. *Neural Comput* **20**, 2185-2226,
doi:10.1162/neco.2008.02-07-474 (2008).

66   Saalmann, Y. B., Pinsk, M. A., Wang, L., Li, X. & Kastner, S. The pulvinar regulates
information transmission between cortical areas based on attention demands. *Science* **337**,
753-756, doi:10.1126/science.1223082 (2012).

67   Zhou, H., Schafer, R. J. & Desimone, R. Pulvinar-Cortex Interactions in Vision and
Attention. *Neuron* **89**, 209-220, doi:10.1016/j.neuron.2015.11.034 (2016).

68 Petersen, S. E., Robinson, D. L. & Morris, J. D. Contributions of the pulvinar to visual
spatial attention. *Neuropsychologia* **25**, 97-105, doi:10.1016/0028-3932(87)90046-7
(1987).

69 Wilke, M., Turchi, J., Smith, K., Mishkin, M. & Leopold, D. A. Pulvinar inactivation
disrupts selection of movement plans. *J Neurosci* **30**, 8650-8659,
doi:10.1523/JNEUROSCI.0953-10.2010 (2010).

70 Wilke, M., Kagan, I. & Andersen, R. A. Effects of pulvinar inactivation on spatial decision-
making between equal and asymmetric reward options. *J Cogn Neurosci* **25**, 1270-1283,
doi:10.1162/jocn_a_00399 (2013).

71 Kawaguchi, Y. & Kubota, Y. Correlation of physiological subgroupings of nonpyramidal
cells with parvalbumin- and calbindinD28k-immunoreactive neurons in layer V of rat
frontal cortex. *J Neurophysiol* **70**, 387-396, doi:10.1152/jn.1993.70.1.387 (1993).

72 Galarreta, M. & Hestrin, S. A network of fast-spiking cells in the neocortex connected by
electrical synapses. *Nature* **402**, 72-75, doi:10.1038/47029 (1999).

73 Cruikshank, S. J., Urabe, H., Nurmikko, A. V. & Connors, B. W. Pathway-specific
feedforward circuits between thalamus and neocortex revealed by selective optical
stimulation of axons. *Neuron* **65**, 230-245, doi:10.1016/j.neuron.2009.12.025 (2010).

74 Gibson, J. R., Beierlein, M. & Connors, B. W. Two networks of electrically coupled
inhibitory neurons in neocortex. *Nature* **402**, 75-79, doi:10.1038/47035 (1999).

75 Agmon, A. & Connors, B. W. Correlation between Intrinsic Firing Patterns and
Thalamocortical Synaptic Responses of Neurons in Mouse Barrel Cortex. *Journal of
Neuroscience* **12**, 319-329 (1992).

76 Poulet, J. F. & Petersen, C. C. Internal brain state regulates membrane potential synchrony
in barrel cortex of behaving mice. *Nature* **454**, 881-885, doi:10.1038/nature07150 (2008).

77 Vinck, M., Batista-Brito, R., Knoblich, U. & Cardin, J. A. Arousal and locomotion make
distinct contributions to cortical activity patterns and visual encoding. *Neuron* **86**, 740-754,
doi:10.1016/j.neuron.2015.03.028 (2015).

78 Arroyo, S., Bennett, C. & Hestrin, S. Correlation of Synaptic Inputs in the Visual Cortex
of Awake, Behaving Mice. *Neuron* **99**, 1289-1301 e1282,
doi:10.1016/j.neuron.2018.08.008 (2018).

79 John, E. R. *et al.* Invariant reversible QEEG effects of anesthetics. *Conscious Cogn* **10**,
165-183, doi:10.1006/ccog.2001.0507 (2001).

80 Zohary, E., Shadlen, M. N. & Newsome, W. T. Correlated neuronal discharge rate and its
implications for psychophysical performance. *Nature* **370**, 140-143,
doi:10.1038/370140a0 (1994).

Abbott, L. F. & Dayan, P. The effect of correlated variability on the accuracy of a population code. *Neural Comput* **11**, 91-101, doi:10.1162/089976699300016827 (1999).

Bartolo, R., Saunders, R. C., Mitz, A. R. & Averbeck, B. B. Information-Limiting Correlations in Large Neural Populations. *J Neurosci* **40**, 1668-1678, doi:10.1523/JNEUROSCI.2072-19.2019 (2020).

Averbeck, B. B., Latham, P. E. & Pouget, A. Neural correlations, population coding and computation. *Nat Rev Neurosci* **7**, 358-366, doi:10.1038/nrn1888 (2006).

Nassi, J. J., Avery, M. C., Cetin, A. H., Roe, A. W. & Reynolds, J. H. Optogenetic Activation of Normalization in Alert Macaque Visual Cortex. *Neuron* **86**, 1504-1517, doi:10.1016/j.neuron.2015.05.040 (2015).

Ruiz, O. *et al.* Optogenetics through windows on the brain in the nonhuman primate. *J Neurophysiol* **110**, 1455-1467, doi:10.1152/jn.00153.2013 (2013).

Konig, S. D. & Buffalo, E. A. A nonparametric method for detecting fixations and saccades using cluster analysis: removing the need for arbitrary thresholds. *J Neurosci Methods* **227**, 121-131, doi:10.1016/j.jneumeth.2014.01.032 (2014).

Mitra, P. P. & Pesaran, B. Analysis of dynamic brain imaging data. *Biophys J* **76**, 691-708, doi:10.1016/S0006-3495(99)77236-X (1999).

Zeitler, M., Fries, P. & Gielen, S. Assessing neuronal coherence with single-unit, multi-unit, and local field potentials. *Neural Comput* **18**, 2256-2281, doi:10.1162/neco.2006.18.9.2256 (2006).

Vinck, M., Battaglia, F. P., Womelsdorf, T. & Pennartz, C. Improved measures of phase-coupling between spikes and the Local Field Potential. *J Comput Neurosci* **33**, 53-75, doi:10.1007/s10827-011-0374-4 (2012).

Figure 1. Laminar Recordings and Saccade Identification

(A) The amplitude-velocity relationship ('main sequence') for all saccades (*n* = 4420) in the dataset. An approximately linear trend is expected for saccades, as is found here (*R²* = 0.555). dva = degrees of visual angle.

(B) Stacked contour plot showing spatial receptive fields (RFs) along the laminar probe from an example session. The RFs are well aligned, indicating perpendicular penetration down a cortical column. Zero depth represents the center of the input layer as estimated with current source density (CSD) analysis.

(C) CSD displayed as a colored map. The x-axis represents time from stimulus onset; the y-axis represents cortical depth. The CSD map has been spatially smoothed for visualization.

(D) Example of continuous eye-tracker and electrophysiological recordings. Eye traces are plotted at the top in grey. LFP traces are plotted by depth below, and spikes are overlaid on the corresponding channel. LFP traces have been color coded by layer identity; spikes have been color coded by cell type.

Figure 2. Narrow-Spiking Input Layer Units Display Peri-Saccadic Enhancement of Firing

(A) Average peri-saccadic normalized firing rate for broad- (n = 102 units) and narrow-spiking (n = 95 units) populations. Thin lines indicate standard error of the mean.

(B) As in (A), but including only broad-spiking units from the superficial (n = 33), input (n = 35), and deep (n = 34) layer populations.

(C) As in (A), but including only narrow-spiking units from the superficial (n = 25 units), input (n = 24 units), and deep (n = 46 units) layer populations.

(D) Average peri-saccadic modulation index for the six neural subpopulations. Two-way ANOVA (F_{layer} = 1.16, P = 0.3151; $F_{cell\ type}$ = 1.84, P = 0.1765; $F_{interaction}$ = 5.98, P = 0.0030) with Tukey's multiple comparison tests (narrow-spiking superficial vs narrow-spiking input, *P = 0.0423; narrow-spiking input vs narrow-spiking deep, *P = 0.0495; narrow-spiking input vs broad-spiking input, **P = 0.0070; narrow-spiking input vs broad-spiking deep, #P = 0.0817). Error bars indicate standard error of the mean.

(E) Average peri-saccadic modulation index for positively modulated units from the six neural subpopulations. Two-way ANOVA (F_{layer} = 1.87, P = 0.1666; $F_{cell\ type}$ = 2.32, P = 0.1355; $F_{interaction}$ = 5.51, P = 0.0077) with Tukey's multiple comparison tests (narrow-spiking superficial vs narrow-spiking input, #P = 0.0833; narrow-spiking input vs narrow-spiking deep, **P = 0.0073; narrow-spiking input vs broad-spiking input, *P = 0.0106; narrow-spiking input vs broad-spiking deep, *P = 0.0384). Error bars indicate standard error of the mean.

(F) Proportion of units in each neural subpopulation with a positive peri-saccadic modulation index.

Figure 3. Input Layer Units Display Modulation Prior to Saccade Onset

(A) Six example single-units demonstrating the approach for identifying the time of significant peri-saccadic firing rate modulation. 95% confidence intervals were calculated for the baseline activity of each unit (dashed grey lines) and the first deviation outside of those bounds was marked as the time of first significant response (red line). The example units shown are the same as those from Figure S1.

(B) Average timing of first significant peri-saccadic modulation for the six neural subpopulations. Input layer units, both broad- and narrow-spiking, show modulation prior to saccade onset. Two-tailed one-sample t-test (broad-spiking input layer, **P = 0.0049; narrow-spiking input layer, *P = 0.0492). Error bars indicate standard error of the mean.

Figure 4. Saccade Onset Increases Correlated Variability Most Prominently in the Input Layer

(A) Spike count correlations as a function of time relative to saccade onset for input-input unit pairs (n = 240 pairs). Correlations were calculated with a sliding 201 ms window. Thin lines indicate bootstrapped 95% confidence intervals.

(B) Average start time of spike count correlation rise relative to saccade onset. Y-axis labels indicate layer identity of unit pairs (S - superficial; I - input; D - deep). Error bars represent bootstrapped 95% confidence intervals. Input-input pairs begin to rise before saccade onset and before all other pair combinations. All other pairs respond after saccade onset.

(C) Average spike-spike coherence between pairs of simultaneously recorded units (single-units and multi-units) before (-200 to 0 ms) and after (0 to 200 ms) saccade onset as a function of frequency. Superficial layer, n = 299 pairs; input layer, n = 191 pairs; deep layer, n = 604 pairs. Thin lines indicate standard error of the mean. Dashed lines indicate spike-spike coherence for shuffled data.

(D) Same data as in (C), but averaged across three frequency bands and plotted as a modulation index: (SSC_{after} - SSC_{before})/(SSC_{after} + SSC_{before}). Two-tailed one-sample t-tests (Superficial 0-12 Hz, ***P = 1.64×10^{-10}; Superficial 15-25 Hz, **P = 0.0027; Superficial 30-100 Hz, P = 0.5468; Input 0-12 Hz, ***P = 2.2131×10^{-7}; Input 15-25 Hz, ***P = 5.5458×10^{-4}; Input 30-100 Hz, ***P = 1.6503×10^{-5}; Deep 0-12 Hz, **P = 0.0013; Deep 15-25 Hz, *P = 0.0116; Deep 30-100 Hz, **P = 0.0050). Error bars indicate standard error of the mean.

Figure 5. Saccade Onset Increases Low Frequency Power and Spike-LFP Locking

(A) Time-frequency representation (spectrogram) of LFP power around saccade onset. Signal strength is represented as percent change in power (i.e. the raw power at each timepoint is divided by the average power in the baseline period, defined here as 250 to 150 ms before saccade onset).

(B) Top: Average power spectra before (-200 to 0 ms) and after (0 to 200 ms) saccade onset. Thin lines indicate standard error of the mean. Bottom: Same data as in top, but averaged across three frequency bands and normalized to data before saccade onset for visualization. Two tailed paired-sample t-tests (0-12 Hz, ***P = 3.4669×10^{-92}; 15-25 Hz, ***P = 9.3690×10^{-74}; 30-100 Hz, ***P = 2.7322×10^{-9}). Error bars indicate standard error of the mean.

(C) Top: Average spike-LFP locking spectra before (-200 to 0 ms) and after (0 to 200 ms) saccade onset. Thin lines indicate standard error of the mean. Bottom: Same data as in top, but averaged across three frequency bands. Two tailed paired-sample t-tests (0-12 Hz, ***P = 1.7147×10^{-19}; 15-25 Hz, P = 0.1598; 30-100 Hz, P = 0.1091). Error bars indicate standard error of the mean.

Figure 6. A Computational Model Confirms that Activation of an Input Layer-Targeting Projection Induces Saccadic Suppression

(A) Connectivity between populations in our computational model. Excitatory (E) and inhibitory (I) populations within each layer project to each other as well as themselves. The input layer excitatory population projects to the superficial layer, while the superficial excitatory population projects to the deep layer. In our model of spontaneous saccades (left), we activate an external excitatory projection that selectively targets the input layer. In our model of saccades executed in the presence of visual stimuli (right), we add an additional input that selectively targets input layer excitatory neurons.

(B) Normalized peri-saccadic firing rate of simulated neural subpopulations during spontaneously executed saccades (*n* = 110 simulated saccades). The input layer inhibitory subpopulation shows enhancement of firing, while all other subpopulations show suppression.

(C) Further dissection of input layer activity. (Top) The saccadic signal arriving in the input layer was simulated with a ramping function. Represented here are saccadic signals of three different strengths. From weakest to strongest, these are represented by dotted, continuous, and dashed lines, both here and in the following subpanels. (Middle) The excitatory and inhibitory input layer subpopulation firing rates in response to three saccadic signals of varying strength. (Bottom) Net excitatory drive onto the inhibitory input layer subpopulation in response to three saccadic signals of varying strength. The excitatory drive is the sum of drive from the external saccadic input and drive from the local excitatory population.

(D) Normalized peri-saccadic firing rate of simulated neural subpopulations during saccades executed in the presence of visual stimulation (*n* = 102 simulated saccades). Visual stimulation is represented by the activation of a second input, which selectively targets the input layer excitatory subpopulation. Here, all subpopulations within the cortical column show peri-saccadic suppression.

Figure S1. Peri-Saccadic Single-Unit Firing Rate Estimation

Peri-saccadic single-unit firing rates were estimated by smoothing individual spikes with gaussian kernels. The kernel bandwidth was selected for each unit at each point in time using an optimization algorithm that minimizes the mean integrated squared error[45]. The firing rate estimates produced by this approach are shown for six representative example units, one from each recorded neural subpopulation. The smoothed firing rates (black) are overlaid on the raster plots showing raw spikes (blue). In the raster plot, each row represents a single saccade. Thin lines are boot-strapped 95% confidence intervals of the firing rate estimate.

Figure S2. Additional Narrow-Spiking Input Layer Single-Unit Examples
Additional example narrow-spiking input layer single-units that display peri-saccadic enhancement of firing. Firing rates were estimated using a kernel smoothing approach and bandwidth optimization algorithm[45]. The smoothed firing rates (black) are overlaid on the spike raster plots (blue). Thin lines are bootstrapped 95% confidence intervals of the firing rate estimate.

Figure S3. Single-Unit Modulation Indices
Single-unit modulation indices as a function of spike waveform duration. Neural populations from each layer are displayed in different subplots. The dashed horizontal line represents a modulation index of zero (no firing rate change), while the dashed vertical line represents the threshold separating narrow- and broad- spiking units.

Figure S4. Input Layer Pairs Show Increases in Spike Count Correlations Prior to Saccade Onset

(A) Spike count correlations as a function of time relative to saccade onset for all combinations of layer-wise unit pairs. Superficial-superficial, *n* = 442 pairs; superficial-input, *n* = 558 pairs; superficial-deep, *n* = 755 pairs; input-input, *n* = 240 pairs; input-deep, *n* = 501 pairs; deep-deep, *n* = 680 pairs. Correlations were calculated with a sliding 201 ms window. Thin lines indicate bootstrapped 95% confidence intervals.

(B) Normalized pupil diameter around the time of saccade onset (*n* = 4420 saccades). Thin lines indicate standard error of the mean.

A

Saccadic signal

I   E   Visual stimulation

B

Spontaneous (visual stim. = 0)

Excitatory firing rate (norm)

baseline
peak sacc. signal
baseline
peak sacc. signal

Inhibitory firing rate (norm)

C

Visual stimulation

Excitatory firing rate (norm)

Inhibitory firing rate (norm)

D

Saccadic signal

Time relative to saccade onset (ms)

E

$W_{EE_A} < W_{EE_B} < W_{EE_C}$

Excitatory firing rate (norm)

W_{EE_A} W_{EE_B} W_{EE_C}

Increasing saccadic signal

Inhibitory firing rate (norm)

Figure S5. Phase Plane Analysis of the Input Layer E-I System

(A) The E-I network in the input layer.

(B) The nullclines and stable activity levels of the input layer E and I populations during simulation of spontaneous activity. Equations and parameters are the same as those described in the Methods and used in Figure 6. The orange and blue curves show the E and I nullclines, respectively, of the E-I network in (A). Each point on a nullcline indicates the steady state firing rate of the E or I population when the activity of the other population is fixed, i.e. the rate of change in equations (3) or (4) is zero. The ISN and non-ISN regimes of the network are demarcated by the switch between positive and negative slopes along the E nullclines. This ability to switch is a consequence of strong E-E connectivity, and is absent when the E-E connectivity strength is either weak or zero. The thin nullclines depict firing rates when saccadic input is at baseline; the thick nullclines depict firing rates when saccadic input increases by a non-zero value. The intersections of the E and I nullclines indicate the steady state firing rates at which the network stabilizes if allowed to seek equilibrium (indicated by black and pink dots at baseline and during peak saccadic suppression, respectively). As illustrated here, the nullclines for our model intersect in the non-ISN regime in the absence of visual input, causing E and I activity to shift in opposite directions in response to the saccadic signal.

(C) Same as in (B), but during simulated visual stimulation. The addition of excitatory input to the E population shifts the E nullclines to the right, causing them to intersect with the I nullclines in the ISN regime. As illustrated here, this causes the E and I populations to shift in the same direction in response to the saccadic signal.

(D) The approximate level of saccadic signal strength that corresponds to the E (orange) and I (blue) nullcline sets shown in (B) and (C). The phase plane analysis is most applicable to constant levels of saccadic signal, and is only an approximate depiction of the scenario of a ramp signal shown in (D), and as used in the model simulation in Figure 6.

(E) Effect of E-E connection strength on inhibitory firing rate changes in response to the saccadic signal. Each set of E nullclines corresponds to a given E-E connection strength (W_{EE}), and illustrates the effects of raising saccadic signal magnitude (indicated by greater line thickness), causing either increases or decreases in steady state firing rates (black dots), as determined by their intersection with the I nullclines. This phase plane analysis approximately predicts three ways in which I firing rate can shift in response to increasing saccadic input: 1) it increases quickly in response to a small increase in saccadic signal (W_{EE_A}), 2) it increases minimally until a threshold level of saccadic signal is reached (W_{EE_B}), or 3) it first decreases then increases with rising saccadic signal strength (W_{EE_C}). The operating regime shown in the simulation results (Figure 6) that recapitulate experimental observations corresponds to the E-E connection strength illustrated by W_{EE_B}. In this case, the network is maintained at the boundary between ISN (positive E nullcline slope) and non-ISN (negative E nullcline slope) regimes in the absence of the saccadic signal.