# A non-adaptive demographic mechanism for genome expansion in *Streptomyces*

Mallory J Choudoir[a†], Marko J Järvenpää[b‡], Pekka Marttinen[b], and *Daniel H Buckley[a]

[a]School of Integrative Plant Science, Cornell University, Ithaca, NY, USA

[b]Department of Computer Science, Helsinki Institute for Information Technology HIIT, Aalto University, Espoo, Finland

[†]Present address: Department of Microbiology, University of Massachusetts Amherst, Amherst, MA, USA

[‡]Present address: Department of Biostatistics, University of Oslo, Oslo, Norway

*Corresponding author:     Daniel H Buckley
                           School of Integrative Plant Science, Bradfield Hall
                           Cornell University, Ithaca, NY 14850
                           dhb28@cornell.edu, (607) 255 1716

**Running title:** Non-adaptive mechanism for genome expansion

**Abstract**

The evolution of microbial genome size is driven by gene acquisition and loss events that occur

at scales from individual genomes to entire pangenomes. The equilibrium between gene gain and

loss is shaped by evolutionary forces, including selection and drift, which are in turn influenced

by population demographics. There is a well-known bias towards deletion in microbial genomes,

which promotes genome streamlining. Less well described are mechanisms that promote genome

expansion, giving rise to the many microbes, such as *Streptomyces*, that have unusually large

genomes. We find evidence of genome expansion in *Streptomyces* sister-taxa, and we

hypothesize that a recent demographic range expansion drove increases in genome size through a

non-adaptive mechanism. These *Streptomyces* sister-taxa, NDR (northern-derived) and SDR

(southern-derived), represent recently diverged lineages that occupy distinct geographic ranges.

Relative to SDR genomes, NDR genomes are larger, have more genes, and their genomes are

enriched in intermediate frequency genes. We also find evidence of relaxed selection in NDR

genomes relative to SDR genomes. We hypothesize that geographic range expansion, coupled

with relaxed selection, facilitated the introgression of non-adaptive horizontally acquired genes,

which accumulated at intermediate frequencies through a mechanism known as genome surfing.

We show that similar patterns of pangenome structure and genome expansion occur in a

simulation that models the effects of population expansion on genome dynamics. We show that

non-adaptive evolutionary phenomena can explain expansion of microbial genome size, and

suggest that this mechanism might explain why some bacteria with large genomes can be found

in soil.

**Introduction**

Microbial genomes are extraordinarily dynamic. Genome size varies considerably, and gene

content in strains of the same species can differ dramatically, giving rise to the pangenome. The

pangenome concept has transformed our understanding of evolutionary processes in diverse taxa

[1–4]. The pangenome is the entire collection of genes in a microbial species, and is subdivided

into core genes present in all strains, dispensable or accessory genes present in some strains, and

strain-specific or unique genes [5, 6]. Rates of gene acquisition and gene loss determine the

individual genome size, and consequently, pangenome composition is shaped by evolutionary

mechanisms that alter gene frequencies in microbial populations [7–9].

Genome size varies by four orders of magnitude ($10^4$–$10^7$ kb) in eukaryotic organisms and two

orders of magnitude in prokaryotic organisms (from less than 150 kb in certain endosymbionts to

over 10 Mb for some free-living bacteria) [10]. Unlike eukaryotes, whose genomes contain large

portions of non-coding DNA, prokaryotic gene content is directly related to genome size because

bacterial and archaeal taxa have high coding density [11, 12]. While microbial genomes are

constantly in flux, deletion rates are approximately three-fold greater than rates of gene

acquisition [13]. Multiple factors contribute to the strong deletion bias in microbial genomes,

including selection for efficiency, "use it or lose it" purging of nonessential genes, and genetic

drift [14–17].

Because of this tendency towards deletion, microbial genome reduction has been examined in

greater detail than genome expansion. For example, the evolutionary mechanisms driving

genome reduction in obligate pathogens like *Rickettsia* and symbionts like *Buchnera* in aphids

are well described [17, 18]. The transition from a free-living to a host-associated lifestyle

involves substantial loss of superfluous genes, and generations of vertical transmission in small

asexual populations leads to gene inactivation and deletion accelerated by genetic drift (16).

Alternatively, genome streamlining leads to reduction of both genome and cell size through

selection for increased metabolic efficiency in free-living microbes with large populations [15].

Genome streamlining is historically associated with marine oligotrophic *Pelagibacter* [14, 19]

but has more recently been described for soil-dwelling *Verrucomicrobia* [20].

Large genomes are frequent among terrestrial free-living microbes, and must be the product of

evolutionary forces that drive genome expansion. A common, though relatively untested,

hypothesis to explain large genomes is that high environmental heterogeneity (a characteristic of

terrestrial habitats) selects for metabolic versatility afforded by gene gain, and thereby drives

genome expansion [21, 22]. For example, massive gene acquisition and adaptation to alkaline

conditions caused genome expansion in the myxobacterium *Sorangium cellulosum*, which at 15

Mb is one of the largest known bacterial genomes [23]. Mechanisms of gene gain include

duplication or horizontal gene transfer (HGT), and large genomes are enriched in functional

genes acquired from phylogenetically distant origins [24]. Much of the evolution of gene

families can be attributed to HGT rather than duplication events [25, 26], and HGT is a major

driver of genome expansion [27, 28]. While HGT-mediated gene acquisition occurs with great

frequency, microbial genomes remain relatively small, and genome size tends to be fairly

conserved within a species.

Gene frequencies at the population-level are governed by selection and drift, and these

evolutionary forces determine whether a newly acquired gene will be purged from the

pangenome or whether it will sweep to fixation. The strength of selection and drift varies

inversely, and their relative contributions are determined by a gene's selection coefficient and

effective population size ($N_e$) [29, 30]. Drift can exert large effects on populations with small $N_e$,

but these effects decline as $N_e$ increases and selection intensifies. Our ability to disentangle the

contributions of selection and drift to pangenome dynamics are complicated by the fact that it

remains difficult to estimate microbial $N_e$ [31, 32] and to delimit microbial population and

species boundaries [33–35]. Another complication is that demographic models often include the

simplifying expectation that $N_e$ is invariable over time.

Rapid changes in population size are typical in the evolutionary histories of many microbial

species, and fluctuations in $N_e$ such as population bottlenecks or expansions can have profound

impacts on contemporary patterns of genomic diversity. For example, the population structure

for many pathogenic bacterial lineages is exemplified by episodes of rapid expansion of clonal

complexes repeated across space and time [36–38]. Microbial population expansions can also be

linked to ecological or geographical range expansions [39–42]. For instance, demographic

expansion in the oral bacteria *Streptococcus mutans* coincides with the origin of human

agricultural practices [41].

We find evidence for post-glacial range expansion in the genus *Streptomyces*, and these species

exhibit several of the genetic characteristics described in plant and animal species whose

biogeography was influenced by Pleistocene glaciation [43, 44]. By examining *Streptomyces*

isolated from sites across North America, we observed genetic evidence for dispersal limitation,

a latitudinal gradient in taxonomic richness, and a latitudinal gradient in genetic diversity [45,

46]. We also identified recently diverged sister-taxa comprising a more genetically diverse

southern-derived (SDR) clade and a more homogenous northern-derived (NDR) clade, which

occupied discrete geographic ranges spanning the boundary of glaciation [47]. We further

observed larger genomes in the northern clade compared to the southern clade.

We hypothesize that genome expansion in NDR is a consequence of demographic change driven

by post-Pleistocene range expansion. Here, we evaluate the effects of historical range expansion

on lineage divergence, genome size, and pangenome structure, and assess these data in the

context of the genome surfing hypothesis. Genome surfing is a non-adaptive mechanism which

describes the introgression of horizontally acquired genes facilitated by relaxed selection and

amplified by geographic expansion [48]. We hypothesize that range expansion, coupled with

relaxed selection, dampened gene loss thereby facilitating an increase in non-adaptive,

intermediate frequency genes in the NDR pangenome. We infer gene gain and loss dynamics by

evaluating patterns of shared gene content between strains. We predict that the contribution of

drift is greater in NDR compared to SDR, and determine the relative strength of selection by

comparing genome-wide rates of amino acid substitution between clades. Finally, we evaluate

our hypothesis by modeling population expansion under a regime of relaxed selection and ask

whether these demographic conditions increase retention of horizontally acquired genes at

intermediate frequencies, ultimately causing genome expansion.

**Material and Methods**

*Streptomyces isolation and genomic DNA extraction*

The strains in this study belong to a larger culture collection of *Streptomyces* isolated from

surface soils (0–5 cm) spanning sites across the United States (see [45, 46]) (Table S1). To

minimize the effects of environmental filtering in driving patterns of microbial diversity, we

selected sample locations with similar ecologies including meadow, pasture, or native grasslands

dominated by perennials and with moderately acidic to neutral soils (pH $6.0 \pm 1.0$, mean $\pm$ SD).

*Streptomyces* strains were isolated by plating air-dried soils on glycerol-arginine agar (pH 8.7)

plus cycloheximide and Rose Bengal [49, 50] as previously described [51]. Genomic DNA was

extracted with a standard phenol/chloroform/isoamyl alcohol protocol from 72 h liquid cultures

grown at 30˚C with shaking in yeast extract-malt extract medium (YEME) + 0.5% glycine [52].

*Genome sequencing, assembly, and annotation*

Genome sequencing, assembly, and annotation is previously described (see [47]). Briefly, we

used the Nextera DNA Library Preparation Kit (Illumina, San Diego, CA, USA) to prepare

sequencing libraries. Genomes were sequenced on an Illumina HiSeq2500 instrument with

paired-end reads (2 x 100 bp). Genomes were assembled with the A5 pipeline [53] and annotated

with RAST [54]. This generated high quality draft genome assemblies with over 25X coverage

and estimated completeness > 99% as assessed with CheckM [55]. We used ITEP and MCL

clustering (inflation value = 2.0, cutoff = 0.04, maxbit score) [56] to identify orthologous

protein-coding gene clusters (i.e., genes). Genome sequences are available through NCBI under

BioProject PRJNA401484 accession numbers SAMN07606143–SAMN07606166.

*Phylogeny*

Phylogenetic relationships were reconstructed from whole genome alignments. We used Mugsy

[57] to generate multiple genome nucleotide alignments and trimAl v1.2 [58] for automatic

trimming of poorly aligned regions. Maximum likelihood (ML) trees were built using the

generalized time reversible nucleotide substitution model [59] with gamma distributed rate

heterogeneity among sites (GTRGAMMA) in RAxML v7.3.0[60], and bootstrap support was

determined following 20 ML searches with 100 inferences using the RAxML rapid bootstrapping

algorithm [61]. Average nucleotide identity (ANI) was calculated from whole genome nucleotide

alignments using mothur [62].

*Pangenome and population genetics analyses*

The pangenome was determined from the gene content of 24 *Streptomyces* genomes (Table S2).

Strains in this collection were initially chosen for whole genome sequencing based on their

genetic similarity at house-keeping loci (see [46]). Subsequent analyses focused on  recently

diverged sister-taxa clades of 10 genomes each, the northern-derived (NDR) and southern-

derived (SDR) lineages. Gene content patterns between strains and pangenome gene frequency

distributions were determined from gene presence/absence data.

Gene-level attributes across gene pools were determined from the average of all nucleotide

sequences within an orthologous protein-coding gene cluster (see above). GC content was

calculated for each gene using the R package Biostrings [63]. Codon usage bias was calculated

for each gene using the R package cordon [64]. Clade-level population genetic traits were

evaluated using 2,778 single-copy genes conserved across all 24 genomes. For each core gene,

nucleotide sequences were aligned using MAFFT v.7 [65], and Gblocks [66] removed poorly

aligned positions. PAL2NAL [67] generated codon alignments, and SNAP [68] calculated intra-

clade non-synonymous ($K_A$) and synonymous ($K_S$) substitution rates (values > 2 were filtered

prior to plotting and statistical analysis).

*Demographic simulation*

We assumed that the SDR pangenome approximates the gene frequency distribution of the last

common ancestor of NDR and SDR. For the starting generation 0, we used the model from

Marttinen *et al.* [69] to simulate a population of sequences and learn parameter values for rates

of gene acquisition and deletion that produced the frequency distribution for SDR. To model

range expansion demographics (i.e., severe bottleneck followed by exponential growth), we

sampled 5 strains from generation 0 as the founding population for the subsequent generation,

and simulated this for 100 generations. The simulated population had a growth rate of 5% per

generation until a maximum of 100 individuals was reached. We varied the initial sizes of the

founding population as well as the growth rate, and observed qualitatively similar results.

The model included gene acquisition events and deletion events similar to Marttinen *et al.* [69]

but modified to allow for multiple changes. Instead of acquisitions/deletions happening

independently, there were k=20 simultaneous acquisitions/deletions per strain per generation.

The previous model [69] included a multiplicative fitness penalty of 0.99 for each gene

exceeding a pre-specified genome size threshold. During the expansion, we relaxed the penalty

for excess genes to $0.99^{\text{(current size/max size)}}$ allowing for genome size variation.

**Results**

*Streptomyces sister-taxa*

We sequenced the genomes of 20 *Streptomyces* strains isolated from ecologically similar

grasslands sites across the United States (Table S1, Table S2). These genomes derive from sister-

taxa comprising a northern-derived (NDR) and southern-derived clade (SDR), which originate

from sites spanning the historical extent of glaciation (Figure S1, see [45]). These sister-taxa

represent closely related but genetically distinct microbial species. Genomes within NDR share

$97.8 \pm 1.3\%$ (mean $\pm$ SD) ANI and those within SDR share $97.6 \pm 0.1\%$ (mean $\pm$ SD) ANI,

while inter-clade genomic ANI is $93.0 \pm 0.14\%$ (mean $\pm$ SD). An ANI of 93–96% is typically

indicative of taxonomic species boundaries [70, 71]. For comparative purposes, we also

sequenced the genomes of four strains that co-localized with the sister-taxa. The closest

taxonomic neighbor to these 24 strains is *Streptomyces griseus* subsp. *griseus* NBRC 13350,

although all strains share < 95% ANI with this type strain (Figure S1).

*Genomic attributes and gene content*

NDR genomes are larger ($8.70 \pm 0.23$ Mb, mean $\pm$ SD) than SDR genomes ($7.87 \pm 0.19$ Mb,

mean $\pm$ SD), and this difference is significant (Mann Whitney U test; $P < 0.0001$) (Figure 1a).

NDR genomes also have also have more orthologous protein-coding gene clusters (hereby

referred to as genes) ($7,775 \pm 196$ genes, mean $\pm$ SD) than SDR genomes ($7,093 \pm 205$ genes,

mean $\pm$ SD), and this difference is also significant (Mann Whitney U test; $P < 0.0001$) (Figure

1b). As expected, there is a strong positive correlation between genome size and gene content

($R^2 = 0.95$, $P < 0.0001$), but coding density did not differ between clades (Figure S2). SDR

genomes are more genetically diverse than NDR. Nucleotide diversity ($\pi$) across conserved,

single-copy core genes is greater in SDR than NDR, and this difference is significant (Mann

Whitney U test; $P < 0.0001$) (see [47]). Finally, NDR genomes have slightly lower genome-wide

GC content ($71.50 \pm 0.087\%$, mean $\pm$ SD) than SDR genomes ($71.62 \pm 0.11\%$, mean $\pm$ SD), and

this difference is significant (Mann Whitney U test; $P = 0.017$) (Figure 1c). Shared gene content

between strains correlates with genomic similarity as measured by ANI (NDR: $R^2 = 0.82$, $P <$

$0.0001$; SDR: $R^2 = 0.64$, $P < 0.0001$) (Figure 2). However, gene content varies more in NDR

than in SDR, and there is a significant interaction between genomic similarity and clade with

respect to gene content shared between strains (Table S3). This interaction comes from shared

gene content between strains increasing more rapidly over recent phylogenetic timescales in

NDR compared to SDR (Figure 3).

*Pangenome structure and dynamics*

The 24 *Streptomyces* genomes (Table S2) contain 22,055 total orthologous protein-coding gene

clusters (i.e., genes), and 42% (9,285 genes) are strain-specific. All 24 genomes share 3,234

(2,778 single-copy) genes, which represent 40–48% of the total gene content per strain. While

NDR has a smaller core genome than SDR (4,234 and 4,400 genes, respectively), its pangenome

is larger (13,681 genes in NDR versus 12,259 genes in SDR) and contains a greater number of

clade-specific genes (5,647 genes unique to NDR versus 4,308 genes unique to SDR) (Figure 3,

Figure 4, Table S4).

For most microbial species, pangenome frequency distributions are U-shaped, reflecting high

proportions of both strain-specific genes and core genes [72]. While the pangenome structures of

our *Streptomyces* sister-taxa generally conform to this shape, the NDR pangenome is enriched in

intermediate frequency accessory genes relative to SDR (Figure 4). The proportion of

intermediate-low frequency (i.e., present in 3–5 strains) accessory genes is higher in NDR than

in SDR (19% of total genes for NDR versus 9.2% of total genes for SDR) (Table S4), and this

difference is statistically significant (two proportion z-test; $P < 0.0001$). Conversely, the

proportion of intermediate-high frequency (i.e., present in 6–8 strains) accessory genes is

equivalent (6.9% of total genes for NDR versus 7.2% of total genes for SDR; two proportion z-

test; $P = 0.26$) (Table S4).

Next, we determined if genes across different gene pools, binned according to their pangenome

frequencies, differed in genetic attributes including per-gene GC content and codon usage bias.

GC content differs between gene pools for both NDR and SDR pangenomes (ANOVA; $F_{3, 25932}$

$= 267.5$, $P$-value $< 0.0001$) (Figure S3). In general, GC content is greater in high frequency and

core genes compared to rare and intermediate frequency genes for both sister-taxa. Codon usage

bias as measured by the effective number of codons (ENC) [73] also differs between gene pools

for both NDR and SDR pangenomes (ANOVA; $F_{3, 21624} = 1862.7$, $P$-value $< 0.0001$) (Figure

S4). Rare and intermediate frequency genes exhibit less overall codon bias compared to high

frequency and core genes, which tend to use codons more preferentially.

*Historical population demography*

Due to founder effects occurring at the edge of an expanding population, $N_e$ is dramatically

reduced during geographic range expansion [74]. Consequently, relaxed selection will

accompany range expansion since the contribution of selection scales directly with $N_e$. Based on

the theory of neutral molecular evolution, which states that selection on synonymous sites is

negligible [75], the ratio of non-synonymous to synonymous amino acid substitutions ($K_A/K_S$)

253 reflects the relative strength of selection acting on a sequence. When assessed at the level of

254 single-copy genes conserved between the sister taxa (2,444 genes), we observe that genome-wide

255 $K_A/K_S$ tends to be higher in NDR than in SDR (Figure 5), and this difference is significant

256 (Mann-Whitney U test; $P < 0.0001$). This result indicates that selection is weaker and genetic

257 drift stronger in NDR relative to SDR.

259 We used a population model (modified from [69]) to determine whether demographic expansion

260 could produce increased intermediate gene frequencies and result in genome expansion. We

261 simulated gene gain and loss events in a population undergoing exponential growth over 100

262 generations, and determined changes in pangenome structure and genome size. To approximate

263 relaxed selection during the population expansion, we imposed a fitness penalty for newly

264 acquired genes that scaled inversely with population size. At the beginning of expansion, most

265 genes were present at high frequencies due to strong founder effects (Figure S5, top and middle).

266 During the expansion, we observed a transient enrichment of intermediate frequency genes

267 within the pangenome (Figure S5, top and middle). Total gene content also increased during

268 population expansion due to relaxed selection pressure when $N_e$ was small, which allowed for

269 the persistence of newly HGT-acquired genes. Genome size stabilized when $N_e$ reached

270 maximum size, and selection pressure balanced HGT-mediated gene gain with simultaneous

271 gene loss (Figure S5, bottom).

273 **Discussion**

274 We have hypothesized that the biogeography of our *Streptomyces* sister-taxa is explained by

275 historical demographic change driven by geologic and climatic events that occurred in the late

Pleistocene [46, 47]. Following the last glacial maxima, North American plant and animal

species rapidly colonized glacial retreat zones, and the genetic consequences of post-glacial

expansion are well documented and include northern-ranged populations with low diversity that

established vast geographic extent [43, 44]. We hypothesize that the recent common ancestor of

NDR and SDR inhabited southern glacial refugia prior to the last glacial maxima (LGM). Post

glaciation, NDR dispersed northward and colonized the latitudinal range it occupies today (see

[46]). We previously described patterns of gene flow, genomic diversity, and ecological

adaptation in these sister-taxa, with both adaptive and non-adaptive processes likely reinforcing

lineage divergence [47]. Here, we evaluate the outcomes of historical range expansion on sister-

taxa pangenome structure and genome size.

Expanding populations experience repeated founder effects as individuals along the leading edge

disperse and colonize new landscapes, creating spatial patterns of genetic diversity akin to

genetic drift [74]. Allele surfing, or gene surfing, is a non-adaptive mechanism that propagates

rare alleles along an expanding edge such that neutral, or even deleterious, variants 'surf' to

higher frequencies than would be expected under population equilibrium [76–78]. When applied

to expanding microbial populations, gene surfing can facilitate genome surfing, a neutral

mechanism acting at the pangenome level that causes rare genes to surf to higher frequencies

independent of natural selection [48]. Below, we outline how historical range expansion and

genome surfing could give rise to genome expansion in *Streptomyces*.

Genome surfing is most likely to occur in microbial populations with intermediate levels of

dispersal and in taxa capable of HGT. Bacteria in the genus *Streptomyces* are ubiquitous in soil

and produce desiccation and starvation resistant spores which are easily disseminated [52],

making them ideal for studying patterns of biogeography dependent on dispersal limitation.

Rates of HGT in *Streptomyces* are among the highest estimated across a range of bacterial

species [51, 79, 80]. In many instances, HGT events occurred in ancestral lineages creating

patterns of shared genetic ancestry and reticulate evolution in many extant *Streptomyces* species

[81]. We previously observed a distance decay relationship between sites up to 6,000 km apart,

indicative of dispersal limitation at intermediate spatial scales that allows detection of geographic

patterns of diversity across the sampled range [45, 46]. We also found evidence of restricted

gene flow between the core genomes of NDR and SDR [47].

Since NDR and SDR sister-taxa share a recent common ancestor (Figure S1), they must also

share a common ancestral genome size. Hence, differences in genome size accompanying

lineage divergence resulted from either genome expansion in NDR or genome reduction in SDR.

Given that changes in genome size are ultimately the result of gene gain and loss, we first

evaluated differences in shared gene content between NDR and SDR strains. We find greater

variability in shared gene content in NDR compared to SDR (Figure 2, Figure 3). This result

suggests relative gene content stability for SDR and gene content instability for NDR, most

notably in recent phylogenetic history (Figure 3). Likewise, the pangenome of NDR exceeds that

of SDR by over 1,000 genes. Evidence suggests that during range expansion, founders at the

expansion edge disperse into new habitats and acquire genes from local gene pools

asymmetrically at unequal rates, and gene flow is almost exclusively from local to invading

genomes [82]. These data are consistent with the observation that that NDR has a larger, more

diverse, and more dynamic pangenome than SDR due to introgression from local gene pools.

Regardless of their origin, most novel horizontally-acquired genes are neutral or nearly neutral

[83]. In most situations, selection will balance gene gain with gene deletion, and genome size

will remain relatively constant.

Genetic diversity in individuals at the leading edge of an expanding population is dramatically

reduced, and their genomes experience relaxed selection pressure due to consecutive population

bottlenecks and low $N_e$ [84]. We find that NDR has lower genetic diversity [47] and higher rates

of $K_A/K_S$ across its core genome relative to SDR (Figure 5), which is consistent with the

prediction that NDR has experienced a period of relaxed selection relative to SDR. A positive

correlation is observed between GC content and selection pressure on microbial genomes [85,

86], and genome expansion in *Chlamydia* has been linked to relaxed selection resulting in a

decrease in genome-wide GC content [87]. We likewise observe a decrease in genome-wide GC

content in NDR relative to SDR (Figure 1). Relaxed selection pressure in NDR would mitigate

the natural bias towards deletion and permit genes acquired by HGT to persist in the genome,

regardless of their adaptive coefficient. Microbial sectoring that accompanies geographic range

expansion [88] would then allow these newly acquired genes to accumulate at intermediate

frequencies in the pangenome. The fact that NDR has larger overall genome size and that relative

selection pressure is lower in NDR than SDR, is contrary to the predictions of the metabolic

versatility hypothesis of large genomes.

We hypothesize that relaxed selection and drift caused genome expansion in NDR. While these

same mechanisms are known to promote genome reduction in endosymbionts and obligate

pathogens [17, 18], it is important to recognize that these outcomes are not contradictory (Figure

6). Genome size is regulated by rates of gene gain and loss, the selective coefficient for each

gene in the genome, and the strength of selection. Endosymbionts and obligate intracellular

pathogens have small population sizes and accordingly, relaxed selection and stronger drift.

Relaxed selection pressure should lessen deletion bias. But under these conditions, host

compensation for microbial gene function radically alters selective coefficients of core genes,

thereby favoring genome reduction, and slightly deleterious mutations accumulate over time via

Muller's ratchet [89, 90]. In addition, rates of HGT from non-host sources are essentially zero,

since there is little opportunity for endosymbionts to interact with other microbial cells, resulting

in a one way track to genome erosion. In contrast, for free-living microbes relaxed selection

pressure should bring about genome expansion by shifting the selective coefficients of accessory

genes towards neutral. For example, genome expansion in *Chlamydia* was driven by relaxed

selection, recombination, and introgression [87]. In this way small population size can favor

genome erosion in endosymbionts, while also favoring genome expansion in free-living

organisms (Figure 6). Meanwhile, free-living organisms that have large population sizes and

high selection pressure will experience high rates of deletion that purge unnecessary genes in

order to promote genome streamlining [14, 15].

Newly acquired genes tend to occur at low frequency in a population unless they provide an

adaptive benefit [91], while adaptive genes will increase rapidly in frequency to join the core

genome. These dynamics are believed to explain the characteristic U-shape of pangenome gene

frequency distributions [72, 92]. Deviations from U-shape expectations, including increased

intermediate frequency genes, can result from changes in selection coefficients of genes or under

conditions where HGT exceeds deletion rates [93]. Alternatively, negative frequency dependent

selection can cause highly beneficial genes to occur at low and intermediate frequencies [94, 95].

A large portion of rare genes in microbial pangenomes are hypothetical proteins or genes of

unknown function acquired through HGT [96, 97]. For both NDR and SDR, approximately 60%

of unique-rare genes (i.e., present in 1–2 strains) are annotated as hypothetical proteins. Nearly

half of the 2,596 genes in NDR's intermediate-low frequency gene pool (i.e., present in 3–5

strains) are also hypothetical genes. Furthermore, intermediate-low frequency genes are similar

to rare frequency genes in regards to GC content (Figure S3) and codon usage (Figure S4). These

data are consistent with our hypotheses that NDR intermediate frequency genes represent

evolutionarily recent HGT-gene acquisitions, which increased in frequency as a result of genome

surfing.

HGT-mediated genome expansion supplies a reservoir of novel genetic material for the evolution

of gene families [25, 26], biosynthetic pathways [98], and formation of new metabolic networks

[99]. Hence, the metabolic versatility of large genomes might be a classic example of an

evolutionary spandrel [100], an adaptive trait associated with large genomes that originated not

because of selection for versatility, but rather because the acquisition of diverse metabolic

pathways is a byproduct of non-adaptive evolutionary process that cause genome expansion.

We show that pangenome analysis of *Streptomyces* sister-taxa verifies several predictions of the

hypothesis that genome expansion within this clade was enabled by non-adaptive evolutionary

processes, most likely driven by late Pleistocene demography. We hypothesize that small

effective population size and relaxed selection, a consequence of geographic range expansion,

allowed for genes newly acquired by HGT to increase in frequency within the NDR pangenome

as a result of genome surfing. Further amplifying this effect is introgression of genes from local

gene pools encountered following dispersal into new environments. Non-adaptive genome

expansion is inherently a non-equilibrium process driven by a transient period of relaxed

selection, and population stabilization will re-impose selection pressures that favor deletion. At

this point, intermediate frequency genes will either be lost to deletion or fixed if they provide

adaptive benefits, and these processes will shift the pangenome structure back to U-shaped

expectations. These insights highlight the importance of considering population demography and

the profound influence of historical contingency on contemporary patterns of microbial genome

diversity.

**Data Availability**

*Streptomyces* genome sequences are available through NCBI under BioProject PRJNA401484

accession numbers SAMN07606143–SAMN07606166.

**Acknowledgements**

This work was supported by the National Science Foundation under Grant No. DEB-1456821

awarded to Daniel H Buckley.

**Competing Interests**

The authors claim no conflicts of interest nor have competing interests.

**References**

1.    Vernikos G, Medini D, Riley DR, Tettelin H. Ten years of pan-genome analyses. *Curr Opin*

414  *Microbiol* 2015; **23**: 148–154.

415 2. Rasko DA, Rosovitz MJ, Myers GSA, Mongodin EF, Fricke WF, Gajer P, et al. The

416  pangenome structure of *Escherichia coli*: comparative genomic analysis of *E. coli*

417  commensal and pathogenic isolates. *J Bacteriol* 2008; **190**: 6881–6893.

418 3. Reno ML, Held NL, Fields CJ, Burke PV, Whitaker RJ. Biogeography of the *Sulfolobus*

419  *islandicus* pan-genome. *Proc Natl Acad Sci U S A* 2009; **106**: 8605–8610.

420 4. Lefébure T, Bitar PDP, Suzuki H, Stanhope MJ. Evolutionary dynamics of complete

421  *Campylobacter* pan-genomes and the bacterial species concept. *Genome Biol Evol* 2010; **2**:

422  646–655.

423 5. Medini D, Donati C, Tettelin H, Masignani V, Rappuoli R. The microbial pan-genome.

424  *Curr Opin Genet Dev* 2005; **15**: 589–594.

425 6. Tettelin H, Masignani V, Cieslewicz MJ, Donati C, Medini D, Ward NL, et al. Genome

426  analysis of multiple pathogenic isolates of *Streptococcus agalactiae*: Implications for the

427  microbial 'pan-genome'. *Proc Natl Acad Sci U S A* 2005; **102**: 13950–13955.

428 7. McInerney JO, McNally A, O'Connell MJ. Why prokaryotes have pangenomes. *Nature*

429  *Microbiology* 2017; **2**: 17040.

430 8. Brockhurst MA, Harrison E, Hall JPJ, Richards T, McNally A, MacLean C. The ecology

431  and evolution of pangenomes. *Curr Biol* 2019; **29**: R1094–R1103.

432 9. Azarian T, Huang I-T, Hanage WP. Structure and Dynamics of Bacterial Populations:

433  Pangenome Ecology. In: Tettelin H, Medini D (eds). *The Pangenome: Diversity, Dynamics*

434  *and Evolution of Genomes*. 2020. Springer, Cham (CH).

435 10. Lynch M. Streamlining and simplification of microbial genome architecture. *Annu Rev*

436  *Microbiol* 2006; **60**: 327–349.

11. Fraser CM, Eisen J, Fleischmann RD, Ketchum KA, Peterson S. Comparative genomics and

understanding of microbial biology. *Emerg Infect Dis* 2000; **6**: 505–512.

12. Kuo C-H, Moran NA, Ochman H. The consequences of genetic drift for bacterial genome

complexity. *Genome Res* 2009; **19**: 1450–1454.

13. Puigbò P, Lobkovsky AE, Kristensen DM, Wolf YI, Koonin EV. Genomes in turmoil:

quantification of genome dynamics in prokaryote supergenomes. *BMC Biol* 2014; **12**: 66.

14. Viklund J, Ettema TJG, Andersson SGE. Independent genome reduction and phylogenetic

reclassification of the oceanic SAR11 clade. *Mol Biol Evol* 2012; **29**: 599–615.

15. Giovannoni SJ, Cameron Thrash J, Temperton B. Implications of streamlining theory for

microbial ecology. *ISME J* 2014; **8**: 1553–1565.

16. Mira A, Ochman H, Moran NA. Deletional bias and the evolution of bacterial genomes.

*Trends Genet* 2001; **17**: 589–596.

17. Moran NA. Microbial minimalism: Genome reduction in bacterial pathogens. *Cell* 2002;

**108**: 583–586.

18. McCutcheon JP, Moran NA. Extreme genome reduction in symbiotic bacteria. *Nat Rev*

*Microbiol* 2011; **10**: 13–26.

19. Giovannoni SJ, Tripp HJ, Givan S, Podar M, Vergin KL, Baptista D, et al. Genome

streamlining in a cosmopolitan oceanic bacterium. *Science* 2005; **309**: 1242–1245.

20. Brewer TE, Handley KM, Carini P, Gilbert JA, Fierer N. Genome reduction in an abundant

and ubiquitous soil bacterium '*Candidatus* Udaeobacter copiosus'. *Nat Microbiol* 2016; **2**:

16198.

21. Konstantinidis KT, Tiedje JM. Trends between gene content and genome size in prokaryotic

species with larger genomes. *Proc Natl Acad Sci U S A* 2004; **101**: 3160–3165.

22.  Dini-Andreote F, Andreote FD, Araújo WL, Trevors JT, van Elsas JD. Bacterial genomes:

habitat specificity and uncharted organisms. *Microb Ecol* 2012; **64**: 1–7.

23.  Han K, Li Z-F, Peng R, Zhu L-P, Zhou T, Wang L-G, et al. Extraordinary expansion of a

*Sorangium cellulosum* genome from an alkaline milieu. *Sci Rep* 2013; **3**: 2101.

24.  Cordero OX, Hogeweg P. The impact of long-distance horizontal gene transfer on

prokaryotic genome size. *Proc Natl Acad Sci U S A* 2009; **106**: 21748–21753.

25.  Lerat E, Daubin V, Ochman H, Moran NA. Evolutionary origins of genomic repertoires in

bacteria. *PLoS Biol* 2005; **3**: e130.

26.  Treangen TJ, Rocha EPC. Horizontal transfer, not duplication, drives the expansion of

protein families in prokaryotes. *PLoS Genet* 2011; **7**: e1001284.

27.  Bohlin J, Brynildsrud OB, Sekse C, Snipen L. An evolutionary analysis of genome

expansion and pathogenicity in *Escherichia coli*. *BMC Genomics* 2014; **15**: 882.

28.  Tsai Y-M, Chang A, Kuo C-H. Horizontal gene acquisitions contributed to genome

expansion in insect-Symbiotic *Spiroplasma clarkii*. *Genome Biol Evol* 2018; **10**: 1526–

1532.

29.  Wright S. Evolution in Mendelian populations. *Genetics* 1931; **16**: 97–159.

30.  Kimura M. Evolutionary rate at the molecular level. *Nature* 1968; **217**: 624–626.

31.  Bobay L-M, Ochman H. The Evolution of Bacterial Genome Architecture. *Front Genet*

2017; **8**: 72.

32.  Rocha EPC. Neutral theory, microbial practice: Challenges in bacterial population genetics.

*Mol Biol Evol* 2018; **35**: 1338–1347.

33.  Roselló-Mora R, Amann R. The species concept for prokaryotes. *FEMS Microbiol Rev*

2001; **25**: 39–67.

34.    Achtman M, Wagner M. Microbial diversity and the genetic nature of microbial species.

484           *Nat Rev Microbiol* 2008; **6**: 431–440.

485    35.    Shapiro BJ. What microbial population genomics has taught us about speciation. In: Polz

486           MF, Rajora OP (eds). *Population Genomics: Microorganisms*. 2019. Springer International

487           Publishing, Cham, pp 31–47.

488    36.    Smith NH, Dale J, Inwald J, Palmer S, Gordon SV, Hewinson RG, et al. The population

489           structure of *Mycobacterium bovis* in Great Britain: clonal expansion. *Proc Natl Acad Sci U*

490           *S A* 2003; **100**: 15271–15275.

491    37.    Achtman M. Population structure of pathogenic bacteria revisited. *Int J Med Microbiol*

492           2004; **294**: 67–73.

493    38.    Nübel U, Dordel J, Kurt K, Strommenger B, Westh H, Shukla SK, et al. A timescale for

494           evolution, population expansion, and spatial spread of an emerging clone of methicillin-

495           resistant *Staphylococcus aureus*. *PLoS Pathog* 2010; **6**: e1000855.

496    39.    Wirth T, Hildebrand F, Allix-Béguec C, Wölbeling F, Kubica T, Kremer K, et al. Origin,

497           spread and demography of the *Mycobacterium tuberculosis* complex. *PLoS Pathog* 2008; **4**:

498           e1000160.

499    40.    Takuno S, Kado T, Sugino RP, Nakhleh L, Innan H. Population genomics in bacteria: a case

500           study of *Staphylococcus aureus*. *Mol Biol Evol* 2012; **29**: 797–809.

501    41.    Cornejo OE, Lefebure T, Pavinski 2. Paulina, Lang P, Richards 2. Vincent P., Eilertson K,

502           et al. Evolutionary and population genomics of the cavity causing bacteria *Streptococcus*

503           *mutans*. *Evolution*  ; **30**: 881–893.

504    42.    Montano V, Didelot X, Foll M, Linz B, Reinhardt R, Suerbaum S, et al. Worldwide

505           population structure, long-term demography, and local adaptation of *Helicobacter pylori*.

*Genetics* 2015; **200**: 947–963.

43.　Hewitt G. Some genetic consequences of ice ages, and their role in divergence and

speciation. *Biological Journal of the Linnean Society* 1996; **58**: 247–276.

44.　Hewitt GM. Genetic consequences of climatic oscillations in the Quaternary. *Philos Trans*

*R Soc Lond B Biol Sci* 2004; **359**: 183–195.

45.　Andam CP, Doroghazi JR, Campbell AN, Kelly PJ, Choudoir MJ, Buckley DH. A

latitudinal diversity gradient in terrestrial bacteria of the genus *Streptomyces*. *MBio* 2016; **7**:

e02200–15.

46.　Choudoir MJ, Doroghazi JR, Buckley DH. Latitude delineates patterns of biogeography in

terrestrial *Streptomyces*. *Environ Microbiol* 2016; **18**: 4931–4945.

47.　Choudoir MJ, Buckley DH. Phylogenetic conservatism of thermal traits explains dispersal

limitation and genomic differentiation of *Streptomyces* sister-taxa. *ISME J* 2018; **12**: 2176–

2186.

48.　Choudoir MJ, Panke-Buisse K, Andam CP, Buckley DH. Genome surfing as driver of

microbial genomic diversity. *Trends Microbiol* 2017; **25**: 624–636.

49.　El-Nakeeb MA, Lechevalier HA. Selective isolation of aerobic Actinomycetes. *Appl*

*Microbiol* 1963; **11**: 75–77.

50.　Ottow JCG. Rose Bengal as a selective aid in the isolation of fungi and actinomycetes from

natural sources. *Mycologia* 1972; **64**: 304.

51.　Doroghazi JR, Buckley DH. Widespread homologous recombination within and between

*Streptomyces* species. *ISME J* 2010; **4**: 1136.

52.　Kieser, T, Bibb, MJ, Buttner, MJ, Charter, KF, Hopwood, DA. Practical *Streptomyces*

Genetics. 2000. John Innes Foundation, Norwich, UK.

53. Tritt A, Eisen JA, Facciotti MT, Darling AE. An integrated pipeline for *de novo* assembly

of microbial genomes. *PLoS One* 2012; **7**: e42304.

54. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST server:

Rapid annotations using subsystems technology. *BMC Genomics* 2008; **9**: 75.

55. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes.

*Genome Res* 2015; **25**: 1043–1055.

56. Benedict MN, Henriksen JR, Metcalf WW, Whitaker RJ, Price ND. ITEP: an integrated

toolkit for exploration of microbial pan-genomes. *BMC Genomics* 2014; **15**: 8.

57. Angiuoli SV, Salzberg SL. Mugsy: fast multiple alignment of closely related whole

genomes. *Bioinformatics* 2011; **27**: 334–342.

58. Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009; **25**: 1972–

1973.

59. Tavare S. Some probabilistic and statistical problems in the analysis of DNA sequences.

Lectures Math Life Sci 17: 57-86 58. Warscheid T, Braams J (2000) Biodeterioration of

stone: a review. *Int Biodeterior Biodegradation* 1986; **46**: 343–368.

60. Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with

thousands of taxa and mixed models. *Bioinformatics* 2006; **22**: 2688–2690.

61. Stamatakis A, Hoover P, Rougemont J, Renner S. A rapid bootstrap algorithm for the

RAxML web servers. *Syst Biol* 2008; **57**: 758–771.

62. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing

mothur: Open-source, platform-independent, community-supported software for describing

and comparing microbial communities. *Appl Environ Microbiol* 2009; **75**: 7537–7541.

63.  Pagès HA, Gentleman P, DebRoy R. Biostrings: Efficient manipulation of biological

strings. *R package version 2.59*. 2020.

64.  Elek A, Kuzman M, Vlahoviček K. coRdon: codon usage analysis and prediction of gene

expressivity. *R package version 1.8.0.* 2019.

65.  Katoh K, Standley DM. MAFFT Multiple sequence alignment software version 7:

Improvements in performance and usability. *Mol Biol Evol* 2013; **30**: 772–780.

66.  Talavera, Gerard, Castresana, Jose. Improvement of phylogenies after removing divergent

and ambiguously aligned blocks from protein sequence alignments. *Syst Biol* 2007; **56**:

564–577.

67.  Suyama M, Torrents D, Bork P. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. *Nucleic Acids Res* 2006; **34**: W609–

12.

68.  Korber BT. HIV Signature and Sequence Variation Analysis. In: Allen G. Rodrigo and

Gerald H. Learn (ed). *Computational Analysis of HIV Molecular Sequences*. 2000.

Dordrecht, Netherlands: Kluwer Academic Publishers, p Chapter 4, pages 55–72.

69.  Marttinen P, Croucher NJ, Gutmann MU, Corander J, Hanage WP. Recombination

produces coherent bacterial species clusters in both core and accessory genomes. *Microbial*

*Genomics* 2015; **1**.

70.  Kim M, Oh H-S, Park S-C, Chun J. Towards a taxonomic coherence between average

nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of

prokaryotes. *Int J Syst Evol Microbiol* 2014; **64**: 346–351.

71.  Ciufo S, Kannan S, Sharma S, Badretdin A, Clark K, Turner S, et al. Using average

nucleotide identity to improve taxonomic assignments in prokaryotic genomes at the NCBI.

*Int J Syst Evol Microbiol* 2018; **68**: 2386–2392.

72.  Haegeman B, Weitz JS. A neutral theory of genome evolution and the frequency

distribution of genes. *BMC Genomics* 2012; **13**: 1.

73.  Wright F. The 'effective number of codons' used in a gene. *Gene* 1990; **87**: 23–29.

74.  Slatkin M, Excoffier L. Serial founder effects during range expansion: a spatial analog of

genetic drift. *Genetics* 2012; **191**: 171–181.

75.  Kimura M. The Neutral Theory of Molecular Evolution. 1983. Cambridge University Press.

76.  Edmonds CA, Lillie AS, Cavalli-Sforza LL. Mutations arising in the wave front of an

expanding population. *Proc Natl Acad Sci U S A* 2004; **101**: 975–979.

77.  Travis JMJ, Münkemüller T, Burton OJ, Best A, Dytham C, Johst K. Deleterious mutations

can surf to high densities on the wave front of an expanding population. *Mol Biol Evol*

2007; **24**: 2334–2343.

78.  Chuang A, Peterson CR. Expanding population edges: theories, traits, and trade offs. *Glob*

*Chang Biol* 2016; **2**: 494–512.

79.  Doroghazi JR, Buckley DH. Intraspecies comparison of *Streptomyces pratensis* genomes

reveals high levels of recombination and gene conservation between strains of disparate

geographic origin. *BMC Genomics* 2014; **15**: 970.

80.  Cheng K, Rong X, Huang Y. Widespread interspecies homologous recombination reveals

reticulate evolution within the genus *Streptomyces*. *Mol Phylogenet Evol* 2016; **102**: 246–

254.

81.  Andam CP, Choudoir MJ, Vinh Nguyen A, Sol Park H, Buckley DH. Contributions of

ancestral inter-species recombination to the genetic diversity of extant *Streptomyces*

lineages. *ISME J* 2016; **10**: 1731–1741.

82. Currat M, Ruedi M, Petit RJ, Excoffier L. The hidden side of invasions: massive

introgression by local genes. *Evolution* 2008; **62**: 1908–1920.

83. Gogarten JP, Townsend JP. Horizontal gene transfer, genome innovation and evolution. *Nat*

*Rev Microbiol* 2005; **3**: 679–687.

84. Excoffier L, Foll M, Petit RJ. Genetic consequences of range expansions. *Annu Rev Ecol*

*Evol Syst* 2009; **40**: 481–501.

85. Hildebrand F, Meyer A, Eyre-Walker A. Evidence of selection upon genomic GC-content

in bacteria. *PLoS Genet* 2010; **6**: e1001107.

86. Raghavan R, Kelkar YD, Ochman H. A selective force favoring increased G+C content in

bacterial genes. *Proc Natl Acad Sci U S A* 2012; **109**: 14504–14507.

87. Bohlin J. Genome expansion in bacteria: the curios case of *Chlamydia trachomatis. BMC*

*Res Notes* 2015; **8**: 512.

88. Hallatschek O, Hersen P, Ramanathan S, Nelson DR. Genetic drift at expanding frontiers

promotes gene segregation. *Proc Natl Acad Sci U S A* 2007; **104**: 19926–19930.

89. Moran NA. Accelerated evolution and Muller's rachet in endosymbiotic bacteria. *Proc Natl*

*Acad Sci U S A* 1996; **93**: 2873–2878.

90. Rispe C, Moran NA. Accumulation of deleterious mutations in endosymbionts: Muller's

ratchet with two levels of selection. *Am Nat* 2000; **156**: 425–441.

91. Kuo C-H, Ochman H. The fate of new bacterial genes. *FEMS Microbiol Rev* 2009; **33**: 38–

43.

92. Lobkovsky AE, Wolf YI, Koonin EV. Gene frequency distributions reject a neutral model

of genome evolution. *Genome Biol Evol* 2013; **5**: 233–242.

93. Domingo-Sananes MR, McInerney JO. Selection-based model of prokaryote pangenomes.

*bioRxiv*. 2019; doi:10.1101/782573.

94. Cordero OX, Ventouras L-A, DeLong EF, Polz MF. Public good dynamics drive evolution

of iron acquisition strategies in natural bacterioplankton populations. *Proc Natl Acad Sci U*

*S A* 2012; **109**: 20059–20064.

95. McNally A, Kallonen T, Connor C, Abudahab K, Aanensen DM, Horner C, et al.

Diversification of colonization factors in a multidrug-resistant *Escherichia coli* lineage

evolving under negative frequency-dependent selection. *MBio* 2019; **10**: e00644–19.

96. Mira A, Klasson L, Andersson SGE. Microbial genome evolution: sources of variability.

*Curr Opin Microbiol* 2002; **5**: 506–512.

97. Mira A, Martín-Cuadrado AB, D'Auria G, Rodríguez-Valera F. The bacterial pan-genome:

a new paradigm in microbiology. *Int Microbiol* 2010; **13**: 45–57.

98. Boucher Y, Doolittle WF. The role of lateral gene transfer in the evolution of isoprenoid

biosynthesis pathways. *Mol Microbiol* 2000; **37**: 703–716.

99. Pál C, Papp B, Lercher MJ. Adaptive evolution of bacterial metabolic networks by

horizontal gene transfer. *Nat Genet* 2005; **37**: 1372–1375.

100. Gould SJ, Lewontin RC. The spandrels of San Marco and the Panglossian paradigm: a

critique of the adaptationist programme. *Proc R Soc Lond B Biol Sci* 1979; **205**: 581–598.

 **Figure Legends**

**Figure 1.** Genomic attributes of NDR and SDR sister-taxa. NDR genomes are larger, have more

genes, and have lower GC content compared to SDR genomes. Plots show the distributions of

genome size in Mb (a), number of genes (b), and genome-wide GC content (%) (c) for

*Streptomyces* sister-taxa. Boxplots show the clade-level medians, interquartile ranges, and 1.5

times interquartile ranges. Colored circles illustrate the values for individual genomes belonging

to the NDR clade (blue) or the SDR clade (green).

**Figure 2.** Genomic similarity versus shared gene content for NDR and SDR. Differences in

shared gene content across increasing average nucleotide identity (ANI) are greater within the

NDR clade compared to the SDR clade (Table S3). Circles show pairwise comparisons of the

number of shared genes between two strains versus ANI and are colored by clade according to

the legend. Dashed lines show linear regressions, and the shaded area is the 95% confidence

interval.

**Figure 3.** Presence/absence of genes across phylogeny. Gene content changes more rapidly

across ancestral phylogenetic nodes for NDR genomes compared to SDR genomes. Tree is made

from whole genome nucleotide alignments, and the scale bar shows nucleotide substitutions per

site (see Figure S1). Branch colors reflect clade membership. Phylogenetic nodes are labeled

with the number of genes conserved in all members of descendent nodes. Gray pie charts at tree

tips show the portion of total genes per genome that are strain-specific (black slice). Right panel

plots the differences in gene content across the phylogeny beginning at the shared ancestral node

and ending with extant taxa at the terminal tips for NDR (blue-solid) and SDR (green-dashed)

lineages. Multiple lines represent monophyletic lineages.

**Figure 4.** Pangenome gene frequency distributions. NDR genomes are enriched in intermediate

frequency genes. Plots show the pangenome gene frequency distributions for NDR (left) and

SDR (right). Bars show the population-level sums of genes present in 1–10 genomes. See Table

S3 for raw values and proportions.

**Figure 5.** $K_A/K_S$ values between the NDR and SDR sister-taxa core genome. NDR core genes

have, on average, greater rates of non-synonymous to synonymous amino acid substitutions

compared to SDR core genes. Circles plot clade-level rates of non-synonymous to synonymous

amino acid substitutions ($K_A/K_S$) for each of 2,444 single-copy core genes for NDR (y-axis) and

SDR (x-axis). Axes are logarithmic scale. The black dashed line is a slope of 1, and points along

this line are genes with equal $K_A/K_S$ mean values in both clades. $K_A/K_S$ is proportional to the

relative strength of genetic drift and inversely proportional to the relative strength of selection.

**Figure 6.** Conceptual overview of the evolutionary processes and demographic conditions that

support changes in genome size. Genome erosion (left) in endosymbionts is the result of small

$N_e$ and strong genetic drift, with host compensation lowering costs of deletion while restricting

gene flow (HGT). Genome streamlining (middle) in free-living microbes with large populations

like *Pelagibacter* involves strong selection and elimination of non-adaptive genes. Genome

expansion (right) in *Streptomyces* is facilitated by high rates of HGT and relaxed selection,

allowing for the accumulation of non-adaptive genes and ultimately larger genomes.

## Genome Erosion
(e.g. Endosymbionts)

Host environment restricts HGT, lowers cost of deletion

**deletion >> acquisition**

Elimination of redundant genes

Negligible HGT, small $N_e$, strong genetic drift, favors **genome reduction**

## Genome Streamlining
(e.g. *Pelagibacter*)

HGT

**deletion ≥ acquisition**

Elimination of non-adaptive genes

HGT, large $N_e$, strong selection, favors **genome reduction**

## Genome Expansion
(e.g. *Streptomyces*)

Relaxed selection lowers deletion rates

HGT

**deletion < acquisition**

Accumulation of horizontally acquired genes

High HGT, small Ne, weak selection, favors **genome expansion**

Supplementary Information for

**A non-adaptive demographic mechanism for genome expansion in *Streptomyces***

MJ Choudoir, MJ Järvenpää, P Marttinen, and DH Buckley

**Table S1.** *Streptomyces* strains were isolated from 11 grassland sites across the United States. Strain names begin with the site code referencing their isolation location. Mean annual temperature (MAT) reflects the 30-year average reported by NOAA.

| Sample Site Location | Code | Latitude (˚N) | Longitude (˚W) | MAT (˚C) |
|---|---|---|---|---|
| Manley Hot Springs, AK | man | 63.9 | -149.0 | -3.9 |
| Bothell, WA | wa | 47.7 | -122.2 | 11.4 |
| Astoria, OR | or | 46.2 | -123.9 | 10.8 |
| Rhinelander, WI | rh | 45.6 | -89.3 | 5.4 |
| Bear Creek, WI | sk | 43.4 | -90.1 | 7.6 |
| Brookfield, WI | b | 43.1 | -88.1 | 7.3 |
| Palo Alto, CA | st | 37.4 | -122.2 | 14.5 |
| Greensboro, NC | gb | 36.1 | -79.9 | 15.1 |
| Starkville, MS | ms | 33.5 | -88.8 | 17.2 |
| Austin, TX | t | 30.2 | -97.8 | 21.0 |
| Fort Pierce, FL | f | 27.5 | -80.4 | 22.9 |

**Table S2.** Descriptive attributes for 24 *Streptomyces* genomes (previously described in (1)). NCBI accession numbers are associated with BioProject PRJNA401484. Strain names reflect sample sites (see Table S1). Clade membership includes the northern-derived (NDR) and southern-derive (SDR) clades, and the remaining four strains belong to independent (IND) lineages. Genome size in Mb. Genome-wide GC content (%). Number of open reading frames (ORFs). Protein-coding orthologous gene clusters (Genes).

| Strain | NCBI Accession | Clade | Size (Mb) | GC (%) | ORFs | Genes |
|--------|----------------|-------|-----------|--------|------|-------|
| b62 | SAMN07606143 | SDR | 7.82 | 71.63 | 7073 | 6568 |
| b84 | SAMN07606144 | NDR | 8.71 | 71.49 | 7657 | 7092 |
| b94 | SAMN07606145 | IND | 8.03 | 72.56 | 6939 | 6502 |
| f150 | SAMN07606147 | SDR | 8.06 | 71.60 | 7320 | 6794 |
| f51 | SAMN07606146 | SDR | 7.72 | 71.77 | 6851 | 6405 |
| gb14 | SAMN07606148 | SDR | 7.88 | 71.67 | 6998 | 6489 |
| man185 | SAMN07606149 | IND | 8.07 | 71.60 | 7244 | 6701 |
| ms115 | SAMN07606150 | SDR | 7.50 | 71.74 | 6776 | 6322 |
| ms184 | SAMN07606151 | SDR | 7.86 | 71.68 | 6958 | 6491 |
| or20 | SAMN07606153 | NDR | 9.08 | 71.54 | 8087 | 7474 |
| or3 | SAMN07606152 | NDR | 8.93 | 71.70 | 7956 | 7329 |
| rh195 | SAMN07606155 | NDR | 8.63 | 71.49 | 7804 | 7245 |
| rh206 | SAMN07606156 | NDR | 8.62 | 71.48 | 7817 | 7262 |
| rh207 | SAMN07606175 | NDR | 8.62 | 71.48 | 7825 | 7268 |
| rh34 | SAMN07606154 | NDR | 8.22 | 71.54 | 7392 | 6897 |
| sk226 | SAMN07606158 | IND | 7.96 | 72.51 | 6930 | 6505 |
| st115 | SAMN07606160 | IND | 8.35 | 71.75 | 7498 | 6966 |
| st140 | SAMN07606161 | SDR | 7.96 | 71.50 | 7184 | 6635 |

| st170 | SAMN07606162 | SDR | 8.08 | 71.42 | 7351 | 6801 |
|-------|--------------|-----|------|-------|------|------|
| st77 | SAMN07606159 | SDR | 8.15 | 71.54 | 7346 | 6784 |
| t99 | SAMN07606163 | SDR | 7.78 | 71.67 | 7076 | 6607 |
| wa1002 | SAMN07606164 | NDR | 8.63 | 71.48 | 7577 | 7038 |
| wa1063 | SAMN07606165 | NDR | 8.84 | 71.39 | 7879 | 7351 |
| wa1071 | SAMN07606166 | NDR | 8.74 | 71.39 | 7755 | 7208 |

**Table S3.** Linear model summary. Table reports coefficient, standard error, t statistic, and *P*-value for explanatory variables. See Figure 2.

|  | **B** | **S.E.** | ***t*** | ***P*-value** |
|--|------|---------|--------|--------------|
| (Constant) | -30696 | 2022 | -15.18 | < 0.001 |
| ANI | 37173 | 2066 | 17.99 | < 0.001 |
| Clade | 21220 | 3370 | 6.30 | < 0.001 |
| ANI x Clade | -21906 | 3448 | -6.35 | < 0.001 |

*$R^2$ = 0.82, Adjusted $R^2$ = 0.81*

**Table S4.** Pangenome frequency distributions for NDR and SDR clades. Table reports the number of genes (n) and the proportion (prop) of the total pangenome across frequencies for NDR and SDR. Frequency refers to the number of genomes a gene is present in, ranging from 1–10. Gene pools are categorized by gene frequencies. For example, intermediate-low genes are present in 3–4 strains, and intermediate-high genes are present in 6–8 strains. *P*-values are from a two proportion z-test, with Bonferrori adjustment for multiple comparisons, evaluating the null hypothesis that the proportion of genes at each frequency is the same for NDR and SDR. Significant *P*-values (< 0.0001) are in bold italics.

| | | NDR | | SDR | | |
| :---: | :---: | :---: | :---: | :---: | :---: | :---: |
| **Gene Pool** | **Frequency** | **n** | **prop** | **n** | **prop** | ***P*-value** |
| Unique | 1 | 4132 | 0.302 | 4188 | 0.342 | ***9.8E-11*** |
| Rare | 2 | 926 | 0.068 | 1097 | 0.089 | ***7.3E-10*** |
| Intermediate - *Low* | 3 | 1182 | 0.086 | 494 | 0.040 | ***3.3E-50*** |
| | 4 | 881 | 0.064 | 377 | 0.031 | ***3.3E-35*** |
| | 5 | 533 | 0.039 | 260 | 0.021 | ***1.5E-15*** |
| Intermediate - *High* | 6 | 430 | 0.031 | 294 | 0.024 | 0.0032 |
| | 7 | 230 | 0.017 | 289 | 0.024 | 0.0012 |
| | 8 | 279 | 0.020 | 303 | 0.025 | 0.21 |
| Near Core | 9 | 854 | 0.062 | 557 | 0.045 | ***2.0E-08*** |
| Core | 10 | 4234 | 0.309 | 4400 | 0.359 | ***3.7E-16*** |

**Figure S1.** Whole genome phylogeny. *Streptomyces* northern-derived (NDR) and southern-derived (SDR) clades are recently diverged and originate from discrete latitudinal geographic ranges. Phylogenetic relationships were reconstructed from whole genome alignments using maximum likelihood and a GTRGAMMA model of evolution. The scale bar indicates nucleotide substitution per site, and nodes are labeled with bootstrap values. Strains are named according to their sample location (see Table S1). NDR clade branches are blue, and SDR clade branches are green. *Streptomyces griseus* NBRC 13350 is included as the closest taxonomic neighbor. The tree was rooted with *Streptomyces hydroscopicus* and *Streptomyces coelicolor* A3(2). Map shows the sample locations which are labeled with site codes (see Table S1) and colored according to the extent of historical glaciation (see (2)).

**Figure S2.** Genome size correlates positively with gene content. Plots show the relationship between genome size in Mb and number of genes (i.e., protein-coding orthologous gene clusters) (a) and open reading frames (ORFs) (b). Circles show values for individual *Streptomyces* genomes (NDR in blue and SDR in green). Lines show the linear regression, and the shaded area is the 95% confidence interval.

**Figure S3.** Per-gene GC content. For each gene frequency pool, plots illustrate the distributions of GC skew for NDR (top) and SDR (bottom). Circles show the average GC content (%) for each gene, and boxplots show the medians, interquartile ranges, and 1.5 times interquartile ranges. Gene pools are defined by frequency and include unique-rare (UR) (present in 1–2 strains), intermediate-low (IL) (present in 3–5 strains), intermediate-high (IH) (present in 6–8), and core-near-core (CNC) (present in 9–10 strains) (Table S4).

**Figure S4.** Per-gene codon usage bias. For each gene frequency pool, plots illustrate the distributions of codon usage bias as measured by the effective number of codons (ENC) (3) for NDR (top) and SDR (bottom). Circles show the average ENC for each gene, and boxplots show the medians, interquartile ranges, and 1.5 times interquartile ranges. ENC values range from 61 (indicating no codon usage bias, or all synonymous codons are used in equal frequency), to 2 (indicating extreme codon bias, or extreme favoring of certain codons). Gene pools are defined by frequency and include unique-rare (UR) (present in 1–2 strains), intermediate-low (IL) (present in 3–5 strains), intermediate-high (IH) (present in 6–8), and core-near-core (CNC) (present in 9–10 strains) (Table S4).

**Figure S5.** Demographic simulation. We simulated a population range expansion and modeled pangenome dynamics and genome size for 100 generations. *Top:* Pangenome gene frequency distributions from a single randomly selected simulation. Generation 0 shows the frequency distribution of the initial population before the bottleneck. Generation 1 shows the frequency distribution right after the bottleneck. Generations 50 and 100 show the frequency distributions at the subsequent generations. *Middle:* Trajectories of gene frequencies across 100 generations for 5 independent simulations. Colored lines show the proportion of genes within the pangenome present at different frequencies as according to legend. For example, blue lines are the proportions

of total genes present in 2–5 strains. Dashed horizontal lines are the observed proportions of the SDR clade (see Table S3). The vertical gray line indicates the generation when the population reached the maximum size. *Bottom:* Genome size. Lines show the average genome size for 5 independent simulations.

**Supplementary References**

1.   Choudoir MJ, Buckley DH. 2018. Phylogenetic conservatism of thermal traits explains dispersal limitation and genomic differentiation of *Streptomyces* sister-taxa. ISME J 12:2176–2186.

2.   Andam CP, Doroghazi JR, Campbell AN, Kelly PJ, Choudoir MJ, Buckley DH. 2016. A Latitudinal Diversity Gradient in Terrestrial Bacteria of the Genus *Streptomyces*. MBio 7:e02200–15.

3.   Wright F. 1990. The "effective number of codons" used in a gene. Gene 87:23–29.