Climate warming changes synchrony of plants and pollinators

Running head: Plant pollinator shifts

Jonas Freimuth[1], Oliver Bossdorf[1], J. F. Scheepens[2], Franziska M. Willems[1,3]

[1]Plant Evolutionary Ecology, Institute of Evolution and Ecology, University of Tübingen, Auf

der Morgenstelle 5, 72076 Tübingen, Germany

[2]Plant Evolutionary Ecology, Institute of Ecology, Evolution and Diversity, Goethe

University Frankfurt, Max-von-Laue-Str. 13, 60438 Frankfurt am Main, Germany

[3] Conservation Biology, University of Marburg, Marburg, Germany (present address)

Correspondence: Franziska Willems, email: franziska.willems@biologie.uni-marburg.de,

phone: +49 6421 2823374

OrcIDs:

Jonas Freimuth: https://orcid.org/0000-0002-9259-3696

Oliver Bossdorf: https://orcid.org/0000-0001-7504-6511

J.F. Scheepens: https://orcid.org/0000-0003-1650-2008

Franziska Willems: https://orcid.org/0000-0002-5481-3686

Abstract

Climate warming changes the phenology of many species. When interacting organisms

respond differently, climate change may disrupt their interactions and affect the stability of

ecosystems. Here, we used GBIF occurrence records to examine phenology trends in plants

and their associated insect pollinators in Germany since the 1980s. We found strong

phenological advances in plants, but differences in the extent of shifts among pollinator

groups. The temporal trends in plant and insect phenologies were generally associated with

interannual temperature variation, and thus likely driven by climate change. The phenological

advancement of plants depended on their level of pollinator dependence: pollinator-

independent plants generally showed weaker phenological shifts than pollinator-dependent

plants. When examining the temporal co-occurrence of plant-pollinator pairs from 1980

onwards, the temporal trends in their synchrony also differed among pollinator groups, and

tended to become more synchronized, mainly because the phenology of plants responded

more strongly to climate change and plants caught up with the pollinators. However, if the

observed trends continue, many interactions may become more asynchronous again in the

future. Our study suggests that climate change affects the phenologies of interacting groups of

organisms, and that it also influences their synchrony.

Keywords: asynchrony, GBIF, mismatch, phenology, pollination mode

Introduction

Phenological events are periodically occurring events in the life cycle of organisms. The timing of these events often depends on environmental factors such as temperature or photoperiod, and it is well known that climate change affects some of these and thus changes the phenologies of many organisms [1]. With such phenology shifts, there is increasing risk of phenological mismatches between interacting organisms, potentially exceeding the natural resilience of ecosystems [2]. Climate change-induced phenological shifts have been documented extensively for individual species [3], but we still know much less about how these shifts affect ecological interactions. Kharouba et al. [4] recently reviewed 54 published interaction studies across ecosystems and interaction types and found no clear general trend, with about half of the studied interactions becoming more asynchronous but the other half becoming even more synchronized through climate change.

Plant-pollinator systems are among the biotic interactions expected to suffer most from a mismatch of phenological events [5]. Several previous studies have observed mismatches [6,7], but in others, pollinators and plants seemed to be able to keep up with each other [8]. An interesting question in this context is also which of the two partners is advancing faster if there is an increasing mismatch. So far, the evidence here is also mixed. For instance, Gordo & Sanz [9] found pollinators to advance faster than trees, and Parmesan [3] that butterflies advanced faster than herbaceous plants, but in a study by Kudo & Ida [10] it was the plants – spring ephemerals – that advanced faster than their bee pollinators.

Mismatches of plant-pollinator interactions can have negative consequences for both partners. For the pollinators, this can include lower survival rates, a decreased overall fitness and higher parasite loads [11]. Moreover, mismatches might also impact pollinator demography, the body sizes [6] and frequencies of sexes, and thus population viability [11]. On the plant side, desynchronized pollinator interactions are mainly expected to impact plant

fitness and thus long-term population growth and survival. For instance, Kudo & Ida [10]

found that seed counts were reduced in early-flowering spring ephemerals after

desynchronization with their bee pollinators. However, in another study fly-pollinated plants

did not show similar responses [12].

Plants differ in their level of dependence on plant pollinators, and an intriguing question

therefore is to what extent phenology responses to climate change are linked to the pollinator

dependence of plants. Bond [13] theorized that wind-pollinated plants might experience little

negative consequences of climate change as they do not depend on interactions with animals.

Conversely, insect-pollinated plants may be subject to strong selection toward phenologies

that are in synchrony with their pollinators. This hypothesis was later corroborated in an

empirical study by Fitter & Fitter [14]. A more recent study on orchids [15] found that

pollination mode influenced the degree of plant advances in flowering phenology, indicating

that self-pollinating and thus pollinator-independent plants were not constrained by pollinator

phenology. The main idea of these previous studies is that all else being equal, pollinator-

independent plants should exhibit stronger phenological shifts in response to the same climate

changes.

Testing hypotheses about plant-pollinator responses to climate change is not trivial.

Since changes in phenology take place on the scale of decades [16], we need long-term data.

A possible source of long-term data on plant phenology are herbarium specimens [17,18],

which can indicate the day of year that a specific species was flowering in a given location

and year. Herbarium data provide unique historical depth, but they need to be treated with

caution because of the sampling biases associated with them [19,20]. In recent years the

digitization of herbaria as well as other collections and observation data, including on other

taxa such as pollinating insects, e.g., from long-term monitoring networks, is creating an

increasing number of public data bases that contain vast amounts of natural history data of

large spatial and temporal scales [21]. These data are increasingly used for analyses of broad

ecological trends and global changes [20,22]. One of the largest and most important hubs of

large-scale and long-term ecological data sets is the Global Biodiversity Facility (GBIF), an

intergovernmental initiative and public data base that provides access to biodiversity data

compiled from various individual sources like academic institutions, government agencies or

independent collections (GBIF, 2019).

Another matter is finding a measure for changes in phenology. Primack et al. [23]

demonstrated that the collection dates of herbarium specimens of a plant species in a year can

be used as a proxy for flowering time in that year. The same approach of using occurrence

records in natural history collections or other data bases can in principle be used to estimate

the activity times of other groups of organisms such as insects (4 and references therein). For

instance, analyses of natural history collections in the UK have demonstrated phenology

changes in bees [7] and butterflies [24]. Thus, the occurrences of plants and insects in GBIF

may be used to estimate activity shifts of different groups, as well as their synchrony. When

we use the term 'activity' in this paper, we refer to the period in an organism's life when it

can interact with its ecological partner. For plants this is the period of flowering, for insect

pollinators the period of flight.

We used data from GBIF to study phenological shifts of plants and pollinators in

Germany, at the level of taxonomic groups as well as individual interactions. We asked the

following questions: (i) Are there long-term trends in the phenology of plants and pollinators?

(ii) If yes, are phenology trends related to climate change? (iii) How are phenological changes

of plants related to their pollinator dependencies? (iv) How does climate change affect the

synchrony of plant-pollinator interactions?

Methods

Phenology data

We worked with occurrence records of plants and insects available from the GBIF database [25–28]. For the plants, we restricted ourselves to species covered by the BioFlor database of plant traits [29], because we needed to be able to classify plants by their level of pollinator dependence (see below). For the insects we restricted ourselves to beetles (Coleoptera), flies (Diptera), bees (Hymenoptera) as well as butterflies and moths (Lepidoptera), as these groups contain most insect pollinators [30]. We used the R package *rgbif* [31] to download all available records of the above taxa from GBIF. Our basic criteria for including records were that they originated from Germany, and that they referred to either a living specimen (e.g., a captured insect), a human observation, just an observation (i.e., when the exact type of observation was not clear), or a preserved specimen (e.g., an herbarium record). If names of plant species were not accepted names, we used the R package *taxsize* [32] to check the names against the GBIF backbone taxonomy and determine the actual accepted name.

Prior to the data analyses, we subjected the data to several steps of quality control. First, we removed all records from before 1980 as these turned out to be too inconsistent, with few records per year and large gaps with years without records. We also removed the records from 2021 as the year had not been complete at the time of our analysis. Second, we removed all records from the days of year (DOY) 1, 95, 121, 163, 164, 166 and 181, and in particular DOY 365 and 366 from the natural history museum of Luxemburg, because the high numbers of records on these days indicated that either records without collecting date had been assigned these by default, or the dates were used by BioBlitz events where very large numbers of records are taken on a specific day of the year. Including these data would have strongly biased the intra-annual distributions of our records. Finally, we removed some records for which no elevation or temperature data could be obtained (see below).

To ensure reasonable coverage of the studied time interval, we then restricted the

records to species which had at least 10 occurrence records in every decade covered (2020

included in the last decade).

After these data curation steps, we maintained just above 12 million occurrence records

that covered altogether 1,764 species, with 11.4 million records of 1.438 plant species, around

590.000 records of 207 species of butterflies and moths, some 76.000 records of 20 bee

species and 30.000 records of 22 fly species, and almost 25.000 records of 77 species of

beetles (**Table 1**). There were large differences between plants and insects not only in the

numbers of records but also in their temporal distribution across the studied period (**Figure S

1**). While plants had relatively even record numbers across years, the insect groups, in

particular flies and bees, were strongly underrepresented in the earlier decades, and record

numbers increased rapidly only in the last 20 years, probably due to the advent of platforms

like iNaturalist.org and naturgucker.de, which allow recording of species occurrences by

citizen naturalists, and which made up most of the insect occurrence data for Germany in

GBIF.

Temperature and elevation data, pollinator dependence, and individual interactions

Besides the main phenology data from GBIF, we obtained several other data sets required for

our analyses. To test for associations with climate, we used temperature data from the Climate

Research Unit (CRU, https://crudata.uea.ac.uk) at the University of East Anglia, specifically

the Time-Series dataset version 4.05 [33], which contains gridded temperature data from

1901-2020 at a resolution of 0.5° longitude by 0.5° latitude. From this dataset we extracted

the monthly mean temperatures and averaged them to obtain the annual mean temperatures at

the sites of occurrence records. To be able to control for elevation at the locations of

occurrence records, we used elevation data at a 90 m resolution from the NASA Shuttle Radar

Topography Mission (SRTM), obtained from the SRTM 90m DEM Digital Elevation

Database [34] and accessed through the *raster* package [35] in R.

To classify plants by their level of pollinator dependence we used plant trait data from

BioFlor [29]. A species was assigned as pollinator-dependent when it was either known to be

self-incompatible and pollinated by an insect, dioecious and pollinated by an insect, or

protogynous/protandrous while also being pollinated by an insect. In contrast, species that

were pollinated abiotically or through selfing, that exclusively reproduced vegetatively, or

were apomicts, were classified as pollinator independent. If none of the above applied, we

assigned an intermediate pollinator dependence. If part of the information above was missing,

no pollinator dependence was determined, and the species was excluded from the analyses

involving pollinator dependence.

Finally, we obtained data on individual plant-pollinator interactions from a UK database

on plant-pollinator interactions hosted by the Centre for Ecology and Hydrology (CEH). This

database included all known interactions between plants and flower-visiting bees, butterflies,

and hoverflies (but unfortunately neither beetles nor moths) in the UK, a country similar to

Germany in terms of climate and species composition. While these interaction data are

unlikely to represent all possible species interactions in Germany, we could not find similar

data for our study area.

Data analysis

All data wrangling and analysis was done in R [36]. Before analysing species reactions, we

wanted to check for consistent patterns of climate change in Germany. For this, we fitted a

linear model to the unique annual mean temperature values at the sites of the records (i.e., the

$0.5° * 0.5°$ grid cells) in our data over time.

To understand phenology changes in plants and insects, we first estimated the

phenological shifts in each group as the slopes of linear regressions linking activity, i.e., the

DOY of a record to its year of observation. We estimated group-specific phenological shifts

in two linear mixed-effect models: one that estimated shifts over time and one that related

phenology variation to temperature. Both models included the DOY of a record as the

response variable, and the latitude, longitude, and elevation of a record as fixed effects. The

temporal-change model additionally included the year of a record as a fixed effect and as a

random effect across species; the temperature-change model instead included the annual mean

temperature at the site of a record as a fixed effect and as random effect across species. We

used the *lme4* package [37] in R to fit these models and assessed model fits by visually

inspecting the relationships between residuals and fitted values, and between residuals and

covariates (**Supplementary material 1 and 2**). Although there were some systematic patterns

in the residuals (for example residuals of low fitted values tended to be higher than residuals

of high fitted values), the random effects from the models agreed well enough with more

complex generalized additive models, so we considered our linear model reasonably robust.

After estimating average group-level phenological shifts, we compared these between the

different insect orders and plants through ANOVAs, using Tukey post-hoc tests for pairwise

tests of differences. We assessed the normal distribution of the slopes through boxplots and

checked potential outliers for their plausibility by examining the data from which these slopes

were estimated. The same procedure was used in all subsequent ANOVAs. To test for an

association between temporal shifts and temperature change, we calculated Pearson

correlations between the slopes of time- and temperature-relationships within each group.

Next, we tested whether phenology trends differed between plant groups with different

levels of pollinator dependence. We performed an ANOVA testing for differences in the

average shifts of plants of different levels of pollinator dependence, and then determined

pairwise differences between groups with a Tukey post-hoc test. Again, we did this both for

temporal and temperature-related shifts.

Finally, we analysed the synchrony of plants and pollinators using the data on

individual plant-pollinator interactions. For each plant-insect pollinator pair we calculated the

differences of the slopes and intercepts estimated for plants and pollinators to obtain a linear

equation describing the change in their asynchrony. To test for differences in the average shift

between groups of interactions, we ran ANOVAs on both the temporal and temperature-

related asynchrony shift data and used Tukey post-hoc tests to examine pairwise differences

between the groups. As analyses of the directions and magnitudes of asynchrony shifts do not

provide any information about the actual state of asynchrony in an interaction, we further

calculated the estimated values of the regressions of asynchrony for each unique year or

temperature point in our dataset. We used these data to then calculate and visualize the

percentages of interactions per group and year/temperature point in which insects were active

earlier than plants.

Results

We observed a strong overall trend of climate warming. Across all records, the annual mean

temperature increased by 0.49 °C per decade ($F_{1, 8477} = 2038$, $R^2_{adj} = 0.19$, $P < 0.001$ for the

linear model), with a total increase of around 2 °C across the entire study period 1980-2020

(**Figure S 2**).

Temporal trends in plant and insect phenology

The phenology of plants advanced on average more strongly than that of any of the insect

pollinators (**Table 1, Figure 1A, C, Figure S 3A**). Among the insect groups, the flies showed

the strongest average temporal shifts, while changes were weaker in butterflies/moths and

bees, and beetles even slightly delayed their activity by almost one day per decade. ANOVA

indicated significant differences among the five studied taxonomic groups ($F_{4, 1759} = 33.44$, P

< 0.001), with plants showing significantly stronger temporal shifts than butterflies/moths,

bees, and beetles, respectively (Tukey post-hoc test significant at $\alpha = 0.05$), and both flies and

butterflies/moths advancing more strongly than beetles.

Temperature sensitivity of plant and insect phenology

In all studied taxonomic groups except for the beetles, the climate sensitivities of phenology

mirrored their temporal trends (**Table 1, Figure 1B, C, Figure S 3B**), again with plants

showing the strongest negative association between temperature and activity, followed by the

flies, then butterflies/moths and bees. Surprisingly, despite their lack of temporal shifts, the

beetles also showed a similar temperature as the butterflies/moths and bees. ANOVA

confirmed that there were significant differences among the studied taxonomic groups ($F_{4, 1759}$

$= 22.19$, $P < 0.001$), although not as clear as in the shifts over time. Here, the only clear

differences were between plants and bees, and between plans and butterflies/moths (Tukey

post-hoc test, significant at $\alpha = 0.05$).

Pollinator dependence

The phenology of plants, and its temporal trends, differed only moderately among plants of

different levels of pollinator dependence (**Table 2**, **Figure S 4**). In all three groups, 85-90% of

the plants advanced their flowering over time, but ANOVA indicated differences among the

groups ($F_{2, 1039} = 7.20$, $P < 0.001$), with slightly weaker temporal shifts in the insect-

independent plants, and significant pairwise differences for the comparison between insect-

independent and intermediate plants as well as between insect-independent and -dependent

plants (Tukey post-hoc tests, significant at $\alpha = 0.05$). The results were similar for the

temperature sensitivities, with generally high fractions (81-92%) of plants showing negative

temperature-phenology relationships in all three groups, but groups also differing from each

other ($F_{2, 1039} = 10.24$, $P < 0.001$). This was mainly driven by weaker average temperature

sensitivities of the pollinator-independent plants, which differed significantly from the insect-

dependent and intermediate plants, respectively (Tukey post-hoc tests, significant at $\alpha = 0.05$).

Synchrony of plant-pollinator interactions

When examining the synchrony of individual plant-pollinator interactions, we found that the

three pollinator groups differed in the magnitude and direction of synchrony changes (**Figure**

2A, B), but that except in hoverflies interactions with plants did not become more

asynchronous but rather more synchronized during the last decades (**Figure 2C**). The

magnitudes of temporal changes were generally greatest for plant-bee interactions, and plant-

butterfly interactions were shifting less than plant-hoverfly interactions (ANVOA: $F_{2, 3985} =$

326.8, $P < 0.001$; significant post-hoc tests for pairwise comparisons between all groups;

Table 3; **Figure 2C**). Moreover, in all three pollinator groups, the insects tended to be the

earlier partners in interactions, at least historically. However, in all three groups there was

also a trend towards a decreased fraction of earlier insects, with plants catching up over time

in plant-butterfly interactions, or even 'overtaking' the insects in plant-hoverfly interactions

(**Figure 2 B, C**). The results are similar when we examined plant-pollinator synchrony in

relation to temperature changes. Here, too, the average changes in synchrony were strongest

in the plant-bee interactions (**Table 3;** ANOVA: $F_{2, 3985} = 87.32$, $P < 0.001$ for differences

between the three groups), and the phenological advances of insects generally decreased with

increasing temperatures, or even turned around at high temperatures in the case of plant-

hoverfly interactions (**Figure S 5**).

Discussion

In this study, we took advantage of large collections of occurrence records to examine phenological trends of flowering plants and insect pollinators in Germany. We asked whether phenology changes affected the synchrony of plants and insects, and whether observed changes in phenology, and variation therein, were related to the different taxonomic groups' responses to climate warming. We also examined whether the phenology responses of plants depended on their levels of pollinator dependence. Our results showed that the phenological shifts of plants and insects indeed differed, with plants shifting most strongly, but also substantial variation among the different insect groups. These changes took place across a period in which the mean annual temperature showed a marked increase in Germany. As peak flowering historically occurred after peak insect activity, these trends led to an increase in plant-pollinator synchrony during the last decades but could potentially lead to future desynchronization if climate change continues.

Plants and insects also differed in their overall temperature sensitivity. While plants shifted on average by ca. five days per degree of warming, even the most sensitive insect groups, the flies, shifted much less with ca. one day per degree of warming. The largest difference between plants and an insect group, the beetles, was almost three and a half days per degree of warming.

Even though groups did not always exhibit temperature sensitivities that reflected their phenology shifts over time, there were clear correlations of individual species shifts over time with their temperature sensitivities. Therefore, it seems likely that the two are causally related, i.e., that anthropogenic climate warming is responsible for the observed phenology shifts.

Lastly, we found an interesting, albeit unexpected, difference between pollinator-dependent and pollinator-independent plants, with pollinator-independent plants showing

much weaker phenology shifts than both intermediately and entirely pollinator-dependent

plants.

Caveats

When interpreting the results of our study, it is important to consider some caveats of the

collections data and occurrence records we used, as the majority of them did not originate

from scientific recording schemes. For instance, the spatial distribution of collections data is

usually quite heterogeneous, and so was our data (**Figure S 1**). We attempted to correct for

spatial heterogeneity in our models by including geographical information. Also, it is likely

that not all of the recorded plants were in bloom at the time of recording, or that not all insects

recorded were imagines.

Phenological shifts over time

The general differences between plants and insects in their advancement of phenology seem

to indicate a shift in the synchrony between plants and their pollinators, with plants advancing

faster than insects. The phenology advances of plants in our data are generally stronger than

reports from previous publications. Fitter & Fitter [14] performed a long-term observation

study of changes in first flowering dates of hundreds of plant species in England, comparing

flowering during 1991-2000 with 1954-1990, and found an average advancement of -4.5

days. This is much less than the -7.2 days per decade we found over the period from 1980 to

2020. A more recent long-term analysis by CaraDonna et al. [38] of phenology changes in

subalpine meadow plants in the Rocky Mountains found an even stronger average

advancement of first flowering of -6.4 days per decade on the level of the entire community

during the study period from 1974 to 2012. Since CaraDonna et al. [38] also analysed peak

floral abundance, their data should be particularly comparable to our estimation of peak

flowering through the DOY of peak occurrence. They found a rate of advancement of -5.3

days per decade in spring peak abundance but only -3.3 days for the summer peak floral

abundance. Both less than the advancement we found.

For insects, previous studies seem to be less consistent, with widespread but not

universal advances in springtime phenology (mostly associated with warming) over the last

decades [39]. For butterflies, long-term records showed that their times of first flights

(correlated with peak appearance) advanced on average by -3.7 days per decade in the 2000s

compared to the previous decades in England [40], and by -7.7 days per decade in California

[41]. The magnitude of the shifts observed in England is similar to what we estimated for

butterflies/moths in Germany (on average -4.0 days per decade). Bartomeus et al. [42] found

that bee phenology in New York advanced by -1.9 days per decade during 1965-2011, which

is also less than the -2.6 days per decade we found in our data, which might be explained by

the additional nine years of data we analysed. Data on the phenology shifts of hoverflies are

scarce [43], but Olsen et al. [44] found an advancement of -6.2 days of the 10th percentile of

collection day (their measure of first flight) during 2000 to 2018, which would be stronger

than the -5.0 we found. Together with previous results, our data indicate that phenological

changes of insects are accelerating during the last decades, but that spring phenology probably

shifts more strongly than peak phenology.

Temperature sensitivities of plant and insects

Although the temperature sensitivities of groups did not entirely mirror their shifts over time,

we found strong correlations between temporal shifts and temperature sensitivities at the

species level. This strongly suggests a link between phenology shifts and climate change,

supporting previous studies such as CaraDonna et al. [38] and Song et al. [45]. We found that

temperature sensitivities, i.e., associations between temperature and phenology, strongly

differed among taxonomic groups. Plants were generally more sensitive to temperature than

insect pollinators. Previous studies on insect phenology in the temperate zone (reviewed in

[39]) have shown that increased spring temperatures are often associated with earlier insect

emergence, but that this pattern cannot be generalized as easily as for plants, as the

temperature–phenology relationships of insects are more complex. While many insects are

able to plastically respond to warmer temperatures by speeding up their rates of development

(and thus potentially emerge earlier), others have been found to not respond, or even to delay

their phenology. This might be due to the dependence of insect development on other cues

such as rainfall [46], due to cold period requirements of insects during their diapause (climate

warming can cause a loss or reduction of this chilling period, and this tends to increase the

amount of warming required for subsequent emergence; [39]), or because species overwinter

in diapause states in which they are not temperature-sensitive [47]. Fründ et al. [47] showed

that bees overwintering in larval stages responded to higher winter temperatures with delayed

emergence, while bees overwintering as adults showed advanced emergence (but had greater

weight losses during overwintering). We did see some species with negative temperature

sensitivity in our data which delayed their phenology. This also connects well to some of the

findings reviewed by Forrest [48], for instance that during winter above-ground nesting bees

experience different temperatures than the plants they feed on during the summer. Such

microclimate differences between insects and plants during overwintering may sometimes

explain contrasting climate responses. In other cases, delays in the first appearance of adults

may result from longer growing seasons. For example, longer growing seasons have reduced

selection for rapid development in some high-elevation grasshoppers, in such a way that they

reach maturity later — but at a larger size — than in the past [49]. Furthermore, warming can

change the number of generations per year (voltinism; [39]). All the above-mentioned

mechanisms can cause variation in the phenology shifts of insects with climate warming and

may therefore explain why climate change is not always accompanied by phenological

advances but might also cause delays – as we observed for the beetles.

Another interesting idea is that the phenological advancement of the plants itself could

cause delayed phenology of some pollinators. Wallisdevries & van Swaay [50] found that

advanced plant growth led to delayed development of butterflies since the cooling created by

shading leaves worsened foraging conditions for the larvae. However, in our study we did not

see this effect for butterflies/moths as their phenology shifts closely tracked the shifts of

plants, perhaps because of the high levels of specialization of many butterfly larvae [51].

Pollinator-dependence of plants

We found that pollinator dependence influenced the plants' reaction to climate. However,

counterintuitively, pollinator-dependent plants shifted more strongly than pollinator-

independent plants. Our finding seems to confirm Fitter & Fitter [14] who also found

significant differences between insect-pollinated plants (-4.8 days shift in day of first

flowering) versus wind-pollinated plants (-3.5 days shift) and suggested this was because

shifting pollinator activity forced plants to flower earlier. Pollinator-dependent plants

advancing faster than pollinator-independent plants would also be in accordance with our

observation that pollinators historically are the earlier partner in interactions. Following this

idea, and of course ignoring the many other factors influencing plant phenological changes,

one could conclude that the strong shifts plants exhibited over time and the large temperature

sensitivities they show, allow plants to catch up with their pollinators.

Changes in plant-pollinator synchrony

When we analysed the synchrony of plant-pollinator interactions, we found clear trends in

shifting synchrony, but they varied among insect pollinator groups. Since the phenology of

plants generally advanced faster than that of the insects during the last decades, but plants had

generally been the later partner in most plant-pollinator interactions, these shifts lead to

greater synchrony overall. However, if the observed trends continue, then many of the studied

interactions will soon reach points of perfect synchrony, and after that the interactions may

become more asynchronous again, albeit in the other direction. For plant-hoverfly interactions

this point has already been reached. If interactions will become more asynchronous again in

the future, then resilience of pollinator networks, in particular through pollinator generalism,

could buffer some of the impacts of phenological mismatches [6] However, while generalist

pollinators make up the larger part of the interactions in most pollination networks, some

plant-pollinator interactions are highly specialized, and these might be the ones suffering most

from future mismatches [52].

Acknowledgements

We are grateful to the two reviewers of a previous submission, whose thorough and

constructive feedback substantially improved our manuscript. This work has been supported

by the DFG Priority Program 1374 "Infrastructure-Biodiversity-Exploratories" (DFG project

BO 3241/7-1 to OB).

Authors contributions

JF and FMW conceived the study; JF and FMW collected analysed the data, JF analysed it,

and wrote the first draft of the manuscript, with guidance from FMW. JFS and OB provided

input to data analysis and manuscript writing. All authors read and approved the final

manuscript.

Data availability

The R code used to conduct the analysis is available at

https://github.com/jonasfreimuth/Phenological-shifts-germany.

References

[1] Cleland EE., Chuine I, Menzel A, Mooney HA. & Schwartz MD. 2007 Shifting plant

phenology in response to global change. *Trends in Ecology & Evolution* **22**, 357–365. DOI:

10.1016/j.tree.2007.04.003.

[2] Memmott J, Craze PG., Waser NM. & Price MV. 2007 Global warming and the disruption

of plant-pollinator interactions. *Ecology letters* **10**, 710–717. DOI: 10.1111/j.1461-

0248.2007.01061.x.

[3] Parmesan C. 2007 Influences of species, latitudes and methodologies on estimates of

phenological response to global warming. *Global Change Biol* **13**, 1860–1872. DOI:

10.1111/j.1365-2486.2007.01404.x.

[4] Kharouba HM., Ehrlén J, Gelman A, Bolmgren K, Allen JM., Travers SE. & Wolkovich

EM. 2018 Global shifts in the phenological synchrony of species interactions over recent

decades. *Proceedings of the National Academy of Sciences of the United States of America*

115, 5211–5216. DOI: 10.1073/pnas.1714511115.

[5] Scheffers BR., Meester L de, Bridge TC. L., Hoffmann AA., Pandolfi JM., Corlett RT.,

Butchart SH. M., Pearce-Kelly P, Kovacs KM. & Dudgeon D *et al.* 2016 The broad

footprint of climate change from genes to biomes to people. *Science (New York, N.Y.)* **354**.

DOI: 10.1126/science.aaf7671.

[6] Miller-Rushing AJ., Høye TT., Inouye DW. & Post E. 2010 The effects of phenological

mismatches on demography. *Philosophical transactions of the Royal Society of London.*

Series B, Biological sciences **365**, 3177–3186. DOI: 10.1098/rstb.2010.0148.

[7] Robbirt KM., Roberts DL., Hutchings MJ. & Davy AJ. 2014 Potential disruption of

pollination in a sexually deceptive orchid by climatic change. *Current biology : CB* **24**,

2845–2849. DOI: 10.1016/j.cub.2014.10.033.

[8] Bartomeus I, Ascher JS., Wagner DL., Danforth BN., Colla S, Kornbluth S & Winfree R.

2011 Climate-associated phenological advances in bee pollinators and bee-pollinated

plants. *Proceedings of the National Academy of Sciences of the United States of America*

108, 20645–20649. DOI: 10.1073/pnas.1115559108.

[9] Gordo O; Sanz JJ. 2006 Temporal trends in phenology of the honey bee Apis mellifera (L.)

and the small white Pieris rapae (L.) in the Iberian Peninsula (1952-2004). *Ecological*

Entomology **31**, 261–268. DOI: 10.1111/j.1365-2311.2006.00787.x.

[10] Kudo G; Ida TY. 2013 Early onset of spring increases the phenological mismatch between

plants and pollinators. *Ecology* **94**, 2311–2320. DOI: 10.1890/12-2003.1.

[11] Schenk M, Krauss J & Holzschuh A. 2018 Desynchronizations in bee-plant interactions

cause severe fitness losses in solitary bees. *The Journal of animal ecology* **87**, 139–149.

DOI: 10.1111/1365-2656.12694.

[12] Kudo G, Nishikawa Y, Kasagi T & Kosuge S. 2004 Does seed production of spring

ephemerals decrease when spring comes early? *Ecological Research* **19**, 255–259. DOI:

10.1111/j.1440-1703.2003.00630.x.

[13] Bond WJ. 1995 Effects of Global Change on Plant—Animal Synchrony: Implications for

Pollination and Seed Dispersal in Mediterranean Habitats. In *Global Change and*

Mediterranean-Type Ecosystems (ed. J. M. Moreno & W. C. Oechel), vol 117, pp. 181–

202. New York, NY: Springer New York. DOI: 10.1007/978-1-4612-4186-7_9.

[14] Fitter AH; Fitter RS. R. 2002 Rapid changes in flowering time in British plants. *Science*

(New York, N.Y.) **296**, 1689–1691. DOI: 10.1126/science.1071617.

[15] Molnár A, Tökölyi J, Végvári Z, Sramkó G, Sulyok J & Barta Z. 2012 Pollination mode

predicts phenological response to climate change in terrestrial orchids: a case study from

central Europe. *Journal of Ecology* **100**, 1141–1152. DOI: 10.1111/j.1365-

2745.2012.02003.x.

[16] Parmesan C. 2006 Ecological and Evolutionary Responses to Recent Climate Change.

Annu. Rev. Ecol. Evol. Syst. **37**, 637–669. DOI:

10.1146/annurev.ecolsys.37.091305.110100.

[17] Jones CA; Daehler CC. 2018 Herbarium specimens can reveal impacts of climate change

on plant phenology; a review of methods and applications. *PeerJ* **6**, e4576. DOI:

10.7717/peerj.4576.

[18] Lang PL. M., Willems FM., Scheepens JF., Burbano HA. & Bossdorf O. 2019 Using

herbaria to study global environmental change. *The New phytologist* **221**, 110–122. DOI:

10.1111/nph.15401.

[19] Daru BH., Park DS., Primack RB., Willis CG., Barrington DS., Whitfeld TJ. S., Seidler

TG., Sweeney PW., Foster DR. & Ellison AM. *et al.* 2018 Widespread sampling biases in

herbaria revealed from large-scale digitization. *The New phytologist* **217**, 939–955. DOI:

10.1111/nph.14855.

[20] Maldonado C, Molina CI., Zizka A, Persson C, Taylor CM., Albán J, Chilquillo E, Rønsted

N & Antonelli A. 2015 Estimating species diversity and distribution in the era of Big Data:

to what extent can we trust public databases? *Global ecology and biogeography : a journal*

of macroecology **24**, 973–984. DOI: 10.1111/geb.12326.

[21] Newbold T. 2010 Applications and limitations of museum data for conservation and

ecology, with particular attention to species distribution models. *Progress in Physical*

Geography: Earth and Environment **34**, 3–22. DOI: 10.1177/0309133309355630.

[22] Chapman, A. D. 2005 *Uses of primary species-occurrence data, version 1.0.* Copenhagen.

[23] Primack D, Imbres C, Primack RB., Miller-Rushing AJ. & Del Tredici P. 2004 Herbarium
specimens demonstrate earlier flowering times in response to warming in Boston.
American journal of botany **91**, 1260–1264. DOI: 10.3732/ajb.91.8.1260.

[24] Brooks SJ., Self A, Toloni F & Sparks T. 2014 Natural history museum collections provide
information on phenological change in British butterflies since the late-nineteenth century.
International Journal of Biometeorology **58**, 1749–1758.

[25] occdownload gbif.org. 2021 *Occurrence Download:* The Global Biodiversity Information
Facility. DOI: 10.15468/dl.t2x9cj.

[26] occdownload gbif.org. 2021 *Occurrence Download:* The Global Biodiversity Information
Facility. DOI: 10.15468/dl.z9wz76.

[27] occdownload gbif.org. 2021 *Occurrence Download:* The Global Biodiversity Information
Facility. DOI: 10.15468/dl.3anj3a.

[28] occdownload gbif.org. 2021 *Occurrence Download:* The Global Biodiversity Information
Facility. DOI: 10.15468/dl.hb2fgs.

[29] Klotz S, Kühn I & Durka W. 2002 BIOLFLOR – Eine Datenbank zu Biologisch-
Ökologischen Merkmalen der Gefäßpflanzen in Deutschland. In *Schriftenreihe für*
Vegetationskunde, vol 38, pp. 1–333.

[30] Kevan PG; Baker HG. 1983 Insects as Flower Visitors and Pollinators. *Annu. Rev.*
Entomol. **28**, 407–453. DOI: 10.1146/annurev.en.28.010183.002203.

[31] Chamberlain, S. A. & Boettiger, C. 2017 *R Python, and Ruby clients for GBIF species*
occurrence data. DOI: 10.7287/peerj.preprints.3304v1.

[32] Chamberlain SA; Szöcs E. 2013 taxize: taxonomic search and retrieval in R.
F1000Research **2**, 191. DOI: 10.12688/f1000research.2-191.v2.

[33] University of East Anglia Climatic Research Unit, Harris, I. C., Jones, P. D. & Osborn, T.
2021 *CRU TS4.05. Climatic Research Unit (CRU) Time-Series (TS) version 4.05 of high-*

resolution gridded data of month-by-month variation in climate (Jan. 1901- Dec. 2020).

https://catalogue.ceda.ac.uk/uuid/c26a65020a5e4b80b20018f148556681.   Accessed   1

September 2021.

[34] Jarvis, A., Reuter, H. I., Nelson, A. & Guevara, E. 2021 *Hole-filled seamless SRTM data*

V4. https://srtm.csi.cgiar.org.

[35] Hijmans, R. J. 2020 raster: Geographic Data Analysis and Modeling. https://CRAN.R-

project.org/package=raster.

[36] R Core Team. 2008 R: A language and environment for statistical computing. Vienna,

Austria. http://www.r-project.org./.

[37] Bates D, Mächler M, Bolker B & Walker S. 2015 Fitting Linear Mixed-Effects Models

Using lme4. *J. Stat. Soft.* **67**. DOI: 10.18637/jss.v067.i01.

[38] CaraDonna PJ., Iler AM. & Inouye DW. 2014 Shifts in flowering phenology reshape a

subalpine plant community. *Proceedings of the National Academy of Sciences of the*

United States of America **111**, 4916–4921. DOI: 10.1073/pnas.1323073111.

[39] Forrest JR. K. 2016 Complex responses of insect phenology to climate change. *Current*

opinion in insect science **17**, 49–54. DOI: 10.1016/j.cois.2016.07.002.

[40] Roy DB; Sparks TH. 2000 Phenology of British butterflies and climate change. *Global*

Change Biol **6**, 407–416. DOI: 10.1046/j.1365-2486.2000.00322.x.

[41] Forister ML; Shapiro AM. 2003 Climatic trends and advancing spring flight of butterflies

in lowland California. *Global Change Biol* **9**, 1130–1135. DOI: 10.1046/j.1365-

2486.2003.00643.x.

[42] Bartomeus I, Park MG., Gibbs J, Danforth BN., Lakso AN. & Winfree R. 2013

Biodiversity ensures plant-pollinator phenological synchrony against climate change. *Ecol*

Letters **16**, 1331–1338. DOI: 10.1111/ele.12170.

[43] Hassall C, Owen J & Gilbert F. 2017 Phenological shifts in hoverflies (Diptera:

Syrphidae): linking measurement and mechanism. *Ecography* **40**, 853–863. DOI:

10.1111/ecog.02623.

[44] Olsen K, Holm TE., Pape T & Simonsen TJ. 2020 Natural history museum collection and

citizen science data show advancing phenology of Danish hoverflies (Insecta: Diptera,

Syrphidae) with increasing annual temperature. *PloS one* **15**, e0232980. DOI:

10.1371/journal.pone.0232980.

[45] Song Z, Fu YH., Du Y, Li L, Ouyang X, Ye W & Huang Z. 2020 Flowering phenology of

a widespread perennial herb shows contrasting responses to global warming between

humid and non-humid regions. *Funct Ecol* **34**, 1870–1881. DOI: 10.1111/1365-

2435.13634.

[46] Bonal R, Hernández M, Espelta JM., Muñoz A & Aparicio JM. 2015 Unexpected

consequences of a drier world: evidence that delay in late summer rains biases the

population sex ratio of an insect. *Royal Society open science* **2**, 150198. DOI:

10.1098/rsos.150198.

[47] Fründ J, Zieger SL. & Tscharntke T. 2013 Response diversity of wild bees to overwintering

temperatures. *Oecologia* **173**, 1639–1648. DOI: 10.1007/s00442-013-2729-1.

[48] Forrest JR. K. 2015 Plant-pollinator interactions and phenological change: what can we

learn about climate impacts from experiments and observations? *Oikos* **124**, 4–13. DOI:

10.1111/oik.01386.

[49] Buckley LB., Nufio CR., Kirk EM. & Kingsolver JG. 2015 Elevational differences in

developmental plasticity determine phenological responses of grasshoppers to recent

climate warming. *Proceedings. Biological sciences* **282**, 20150441. DOI:

10.1098/rspb.2015.0441.

[50] Wallisdevries MF; van Swaay CA. M. 2006 Global warming and excess nitrogen may

induce butterfly decline by microclimatic cooling. *Global Change Biol* **12**, 1620–1626.

DOI: 10.1111/j.1365-2486.2006.01202.x.

[51] Gilbert LE; Singer MC. 1975 Butterfly ecology. *Annual Review of Ecology and*

Systematics 6, 365–395.

[52] Bascompte J; Jordano P. 2007 Plant-Animal Mutualistic Networks: The Architecture of

Biodiversity. Annu. Rev. Ecol. Evol. Syst. 38, 567–593. DOI:

10.1146/annurev.ecolsys.38.091206.095818.

Table 1 Sizes of the studied taxonomic groups their numbers of records, their ranges of records per species, the extents of their average temporal

shifts (± 1 S.E.) of phenology, average temperature sensitivities (± 1 S.E.), and the respective fractions of species that advance over time, or with

increasing temperatures.

Group	# Species	# Records	# Records / Species			Temporal shifts		Temperature sensitivity	
			Min	Median	Max	Days / decade	% Advancing	Days / 1 °C	% Advancing
Beetles	77	24,758	55	196.0	1,879	0.93 ± 0.94	44%	-1.68 ± 0.52	73%
Flies	22	30,927	166	674.5	9,194	-5.03 ± 0.88	91%	-3.88 ± 0.71	91%
Bees	20	76,438	91	599.0	15,059	-2.61 ± 2.04	65%	-2.01 ± 1.3	60%
Butterflies/ Moths	207	589,970	69	803.0	33,228	-4.02 ± 0.32	86%	-1.94 ± 0.25	74%
Plants	1,438	11,357,981	62	2,280.0	121,612	-7.24 ± 0.19	86%	-5.15 ± 0.15	84%

599 **Table 2** Average temporal shifts and temperature sensitivities (each ± 1 S.E.) of plants with different levels of pollinator dependence, and the

600 respective fractions of species that advance over time, or with increasing temperatures.

Insect Dependence	# Species	Temporal shifts		Temperature sensitivity	
		Days / decade	% Advancing	Days / 1 °C	% Advancing
Pollinator-dependent	148	-8.56 ± 0.61	90%	-5.76 ± 0.45	91%
Intermediate	665	-7.59 ± 0.27	87%	-5.62 ± 0.23	85%
Pollinator-independent	229	-5.95 ± 0.43	85%	-3.7 ± 0.35	81%

602 **Table 3** Changes in asynchrony of plant-pollinator interactions, separately by pollinator groups. Temporal changes (± 1 S.E.) are in days /

603 decade, and temperature sensitivities (± 1 S.E.) are expressed as days / °C.

Interaction group	# Pairwise interactions	# Plant species	# Pollinator species	Temporal shifts [days / decade]	Temperature sensitivities [days / 1 °C]
Bee - Plant	924	230	12	-11.38 ± 0.41	-5.64 ± 0.26
Butterfly - Plant	1,782	227	39	-2.1 ± 0.2	-2.59 ± 0.17
Hoverfly - Plant	1,282	194	17	-3.47 ± 0.21	-1.84 ± 0.17

Figure 1. Temporal trends (A) and temperature sensitivities (B) and their correlation (C) of

the phenology (peak flowering/activity) of plants and pollinator groups, with the coloured

dots indicating the averages for each group. Black dots indicate individual species slopes

confidence intervals. Given above the dots are group size (*n*), Pearson correlation coefficient

(*r*), and the significance level of the correlation (***: $P < 0.001$).

Figure 2. Asynchrony of individual plant-pollinator interactions, and their temporal trends,

separated by pollinator groups. (A) Average decadal asynchrony changes of individual

interactions (grey dots), and the means for each group (coloured dots and 95% CI whiskers).

(B) Group average changes of asynchrony (black dots: individual interactions yearly

asynchrony, grey contours: indication of point density, coloured lines: linear regression for

each group). (C) Fraction of interactions with earlier insect activity per group.