Single-cell RNA-seq analysis reveals penaeid shrimp hemocyte subpopulations and cell differentiation process

Keiichiro Koiwai [a, *], Takashi Koyama [b, c], Soichiro Tsuda [d], Atsushi Toyoda [e], Kiyoshi Kikuchi [b], Hiroaki Suzuki [f], Ryuji Kawano [a]

[a] Department of Biotechnology and Life Science, Tokyo University of Agriculture and Technology, Koganei-shi, Tokyo 184-8588, Japan

[b] Fisheries Laboratory, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Hamamatsu, Shizuoka 431-0214, Japan

[c] Graduate School of Fisheries and Environmental Sciences, Nagasaki University, Bunkyo-machi, Nagasaki-shi, Nagasaki 852-8521, Japan

[d] bitBiome Inc., Waseda University Incubation Center, Nishiwaseda, Shinjuku-ku, Tokyo 169-0051, Japan

[e] Advanced Genomics Center, National Institute of Genetics, Mishima, Shizuoka 411-8540, Japan

[f] Department of Precision Mechanics, Faculty of Science and Engineering, Chuo University, Bunkyo-ku, Tokyo 112-8551, Japan

* Keiichiro Koiwai

Email: koiwai@go.tuat.ac.jp

Author Contributions: K. Koiwai designed the experiments; K. Koiwai, T.K., S.T., and A.T. performed the experiments; K. Koiwai analyzed the data; K. Kikuchi, H.S., and R.K. supervised the research; K. K. wrote the paper.

Keywords: scRNA-seq; non-model; hemocytes; cell differentiation; crustacean; shrimp

This PDF file includes:

Main Text
Figures 1 to 7

Abstract

Crustacean aquaculture is expected to be a major source of fishery commodities in the near future.

Hemocytes are key players of the immune system in shrimps; however, their classification, maturation, and

differentiation are still under debate. To date, only discrete and inconsistent information on the classification

of shrimp hemocytes has been reported, showing that the morphological characteristics are not sufficient to

resolve their actual roles. Our present study using single-cell RNA sequencing, revealed nine types of

hemocytes of *Marsupenaeus japonicus* based on their transcriptional profiles. We identified markers of each

subpopulation and the differentiation pathways involved in their maturation. We also discovered cell growth

factors that might play crucial roles in hemocyte differentiation. Different immune roles among these

subpopulations were suggested from the analysis of differentially expressed immune-related genes. These

results provide a unified classification of shrimp hemocytes, which improves the understanding of its immune

system.

Introduction

Aquaculture is an important source of animal protein and is considered one of the most important

long-term growth areas of food production, providing 60% of fish for human consumption (1)

(http://www.fao.org/fishery/statistics/en). However, crustaceans that lack an adaptive immune system (2-4)

are vulnerable to pathogens. This means that ordinal vaccination is not applicable to crustaceans, unlike in

fish aquaculture. Shrimp is the main target species for crustacean aquaculture. Therefore, an immune priming

system for shrimp, which is entirely different from conventional vaccines, needs to be developed to control

the infection of pathogens. However, little is known about the immune system of crustaceans due to the lack

of biotechnological tools, such as uniform antibodies and other biomarkers (5).

Hemocytes, which are immune cells of crustaceans, are traditionally divided into three

morphological types based on the dyeing of intracellular granules, which was established by Bauchau and

colleagues (6-8). However, there have been additional reports on the classification of the hemocytes of

shrimp; they were classified into four, eight, and five types based on electron microscopy (9), another dyeing

method (10), and iodixanol density gradient centrifugation (11), respectively. As the morphology and dye

staining properties of shrimp hemocytes are not absolute indicators, no unified understanding of their role

has been established yet. Molecular markers, such as specific mRNAs, antibodies, or lectins, are usually

available for characterizing the subpopulations of cells in model organisms, but this is not often the case for

non-model organisms. Although monoclonal antibodies have been developed for some hemocytes of shrimp

(12-18), their number is lower than that of humans, and their correspondence to the cell type, as well as their

differentiation stage are under debate.

Recently, single-cell mRNA sequencing (scRNA-seq) techniques have dramatically changed this

scene, allowing researchers to annotate non-classified cells solely based on the mRNA expression patterns

of each cell. In particular, droplet-based mRNA sequencing, such as Drop-seq, developed by Macosko, Basu

(19), has gained popularity for classifying cells and identifying new cell types. The enormous amount of
biological data obtained from scRNA-seq leads us to classify cells into specific groups, analyze their
heterogeneity, predict the functions of single-cell populations based on the gene expression profiles, and
determine the cell proliferation or development pathways based on the pseudo-time ordering of a single cell
(20, 21). More recently, hemocytes of invertebrate, fly, and mosquito models have been subjected to these
types of microfluidic-based scRNA-seq to reveal their functions (22, 23).

Here, we performed scRAN-seq analysis on *Marsupenaeus japonicus* hemocytes to classify the
hemocyte types and to characterize their functions using the custom-built Drop-seq platform. To perform
scRNA-seq, a high-quality gene reference is essential; however, such reference genomes are scarce for
crustaceans because of the extremely high proportion of simple sequence repeats (5). We circumvented this
problem by preparing reference genomes using hybrid *de novo* assembly of short- and long-read RNA
sequencing results. The sequences obtained from the scRNA-seq were mapped onto the reference genes
successfully. Our scRNA-seq uncovered the transcriptional profiles of a few thousand *M. japonicus*
hemocytes. We identified the markers of each population and the differentiation pathways associated with
their maturation. We also discovered the cell growth factors that might play crucial roles in hemocyte
differentiation. Different immune roles among these subpopulations were also suggested from the analysis
of differentially expressed immune-related genes. Our results present a unified classification of shrimp
hemocytes and a deeper understanding of the immune system of shrimp.

Results
scRNA-seq clustering of *Marsupenaeus japonicus* hemocytes
Our study utilized scRNA-seq to determine the cellular subtypes with a distinct transcriptional
expression (Fig. 1 A). To map the scRNA-seq sequences from *M. japonicus* hemocytes, we first prepared *de*
novo assembly of the reference genes using hybrid assembly of short- and long-read RNA sequencing results.
Then, by using self-built Drop-seq microfluidic chips, single hemocytes were captured and their mRNA was
barcoded using the droplet-based strategy. This process was performed in triplicates for three shrimp
individuals. Following library preparation and sequencing, the transcriptomes obtained from scRNA-seq
were mapped against the reference genes to discover the cell types.

Using the Drop-seq procedure, we profiled a total of 2,704 cells and obtained a median value of 718
unique molecular identifiers (UMIs) and 334 genes per cell across three replicates (Fig. S1 A and B).
Approximately 300 genes were detected, and the total number of mRNAs expressed among individual cells
varied between 100 and 1,400 (Fig. S1 C and D). There was some transcriptional variability, which may have
resulted from the artifacts of the Drop-seq system, because it is consistent with the original Drop-seq paper
(19) and with recent findings in other organisms, such as fish (24), flies (23), and mosquitoes (22, 25).

Applying the SCTransform batch correction method integrated into the Seurat package allowed us
to remove the individual differences. SCTransform successfully integrated all three shrimp datasets, among

which we identified a total of nine clusters (Fig. 1 B) and obtained 3,334 commonly expressed genes. Each
cluster contained the following number of cells: Hem 1, 65 cells (2.4%); Hem 2, 461 cells (17.0%); Hem 3,
313 cells (11.6%); Hem 4, 524 cells (19.4%); Hem 5, 92 cells (3.4%); Hem 6, 553 cells (20.5%); Hem 7, 135
cells (5.0%); Hem 8, 170 cells (6.3%); and Hem 9, 391 cells (14.5%), respectively.

99
100 **Figure 1.** scRNA-seq of penaeid shrimp *M. japonicus* hemocytes. (A) Schematics of the microfluidics-based
101 scRNA-seq, preparation of the *de novo* assembled gene list, *in silico* analysis workflow and morphology-
102 based cell classification. (B) UMAP (uniform manifold approximation and projection) plot of SCTransform
103 batch corrected and integrated of hemocytes from three shrimps (*n* = 2,704).

Cluster specific markers and their functional prediction

A total of 40 cluster-specific markers were predicted using the Seurat FindMarkers tool (Fig. 2,
Dataset S1, and Dataset S2). For each cluster, seven (Hem 1), six (Hem 2), one (Hem 3), four (Hem 4), eight
(Hem 5), five (Hem 7), one (Hem 8), and eight (Hem 9) markers were selected. Their functions were then
annotated using BLASTX searching for penaeid shrimp identical proteins downloaded from a public
database.

Hem 1 specific markers, *histone acetyltransferase* lysine acetyltransferase 6A (*HAT KAT6A)*,
gamma-aminobutyric acid (*GABA*) *transporter*, and *polypyrimidine tract-binding protein 1* (*PTBP1*) are
genes related to cell proliferation, cell migration, and colony formation in human tumor studies. HAT

KAT6A is known to be a chromatin regulator that controls fundamental cellular processes and is implicated

in regulating tumor progression (26). Autocrine/paracrine signaling via GABA receptors negatively controls

ES cell and peripheral neural crest stem cell proliferation (27). This GABA signaling pathway critically

regulates proliferation independently of differentiation, apoptosis, and overt damage to DNA (27). PTBP1 is

a multi-functional RNA-binding protein that is overexpressed in glioma, a type of tumor that occurs in the

brain, and a decreased expression of PTBP inhibits cell migration and increases the adhesion of cells to

fibronectin and vitronectin (28, 29). PTBP has been shown to be involved in germ cell differentiation in

Drosophila melanogaster and is essential for the development of *Xenopus laevis* (30). These pieces of

evidence strongly suggest that these Hem 1 markers are related to cell proliferation in shrimp hemocytes.

Among the six markers found in cluster Hem 2, three markers were annotated with hemocyte

transglutaminase (HemTGase), and one marker showed high similarity with von Willebrand factor D and

epidermal growth factor (EGF) domain-containing protein (VWDE). Interestingly, both Hem 1 and Hem 2

showed a high expression of TGase, an immature hemocyte marker of crayfish and shrimp. When the

extracellular TGase is digested, hemocytes start to differentiate into mature hemocytes (31-33). The high

expression of TGase suggests that both Hem 1 and Hem 2 are in the early stage of hemocytes. In cluster Hem

3, the only identified marker showed high similarity with tubulointerstitial nephritis antigen (TINAGL). In

cluster Hem 4, all markers showed similarity with the hypothetical protein.

In clusters Hem 5 to Hem 9, many of the cluster-specific markers showed similarity with immune-

related genes of penaeid shrimp, and very few genes were unknown. We postulate that this is partly because

the public database is rich in immune-related genes because of their importance. In cluster Hem 5, markers

showed a high similarity with c-type lysozyme, viral responsive protein (VRP), fibrous sheath CABYR-

binding (FSCB)-like, chitin binding-like protein, anti-lipopolysaccharide factor (ALF)-A1, and

PDGF/VEGF-related factor 1. In cluster Hem 7, markers showed high similarity with selenium-dependent

glutathione peroxidase (Se-GPX), BigPEN, ALF-C1, and hemocyte Kunitz protease inhibitor (KPI). In

cluster Hem 8, the marker showed high similarity with KPI. In cluster Hem 9, the markers showed similarity

with insulin-like growth factor-binding protein (IGFBP)-related protein 1, hemocyte KPI, crustacean

hematopoietic factor (CHF)-like protein, single whey acidic protein domain (SWD)-containing protein, and

penaeidin-II.

Altogether, only 70% (28/40) of markers were annotated from the BLAST searches on the penaeid

shrimp identical protein data. We also performed GO annotation using eggNOG-mapper (34, 35)

(http://eggnog-mapper.embl.de/) to predict the functions of contigs, but only about 10% of the contigs were

annotated, and we could not reach GO analysis.

From the results of this cluster-specific marker analysis, we found that Hem 1 and Hem 2 are

immature hemocytes. Therefore, we analyzed the whole maturation process of hemocytes using pseudo-

temporal analysis. Additionally, cell growth-related genes, such as *VWDE*, *TINGAL*, *IGFBP-related protein*,

and *CHF*, were identified as cluster-specific markers. We predicted their functions to pursue the linkage

between these genes and the hemocyte differentiation process. Finally, the immune-related genes, which

were enriched in Hem 5 to 9, were analyzed to predict the immune function of each cluster.

152
153
154   **Figure 2.** Dot plot representing the marker genes per cluster based on the average expression predicted using

155   the Seurat FindMarker tool. Color gradient of the dot represents the expression level, while the size represents

156   the percentage of cells expressing any gene per cluster.

158   **Pseudo temporal ordering of cells delineates hemocyte lineages**

159   To investigate the dynamics of hemocyte differentiation, we performed lineage-tree reconstruction

using the monocle3 learn_graph function. The differentiation and proliferation of hemocytes in shrimp and

other crustaceans are still under debate (36). Since the cell cycle- and hemocyte-type-specific markers are

better studied in *Drosophila*, we checked the commonly expressed genes of *M. japonicus* that are similar to

the markers of *Drosophila* to determine whether they are present in shrimp. In the search for cell cycle-

specific markers (Dataset S3), a large portion of cells in Hem 1 expressed the G2/M-related genes of

Drosophila: *heterochromatin protein 1b* (*HP1b*); Mj-13141, *CCCTC-binding factor* (*CTCF*); Mj-14904,

suppressor of variegation 205 (*Su(var)205*); Mj-27796 (Fig. 3 A and Fig. S2). *Drosophila HP1* and

Su(var)205 are known to be essential for the maintenance of the active transcription of the euchromatic genes

functionally involved in cell-cycle progression, including those required for DNA replication and mitosis

(37, 38). CTCF has zinc finger domains and plays an important role in the development and cell division of

fly and mammalian cells (39, 40). In the BLAST search of two genes related to cell-cycle progression, that

is, Mj-13141 and Mj-27796, on identical proteins of penaeid shrimp, they showed high similarity with the

chromobox protein homolog 1-like (Dataset S1). A chromobox family protein contributes to

lymphomagenesis by enhancing stem cell self-renewal and/or by increasing the replicative potential of cancer

stem cells in human tumor cells (41). BLAST search of Mj-14904 on identical proteins of penaeid shrimp

showed similarity with the zinc finger protein that contains the same domain of CTCF (Dataset S1). These

findings suggested that hemocytes grouped as Hem 1 are tightly regulated by these G2/M phase-related genes

to promote cell division.

Among the specific markers of four types of *Drosophila* hemocytes, four genes in prohemocytes,

11 genes in plasmatocytes, 11 genes in lamellocytes, and 6 genes in crystal cells showed similarity with the

shrimp genes (Dataset S4). However, no genes were significantly expressed in certain clusters (Fig. S3).

Therefore, we concluded that these *Drosophila* hemocyte markers cannot be adapted as markers for the

hemocytes of shrimp.

We considered Hem 1 to be the initial state of hemocytes and set it as the starting point in the

differentiation process because Hem 1 expressed cell proliferation-related genes, TGase (an immature

hemocyte marker), and G2/M phase-related genes (37-42). From this pseudo temporal ordering analysis, we

found four main lineages starting from Hem 1 to Hem 5, Hem 7, Hem 8, and Hem 9 at the endpoints (Fig. 3

B and Fig. S4 A-D). In crayfish, hematopoietic stem cells are present in hematopoietic tissue (HPT), and two

types of hemocyte lineages starting from a hematopoietic stem cell exist (43). In *Penaeus monodon*, hyaline

cells (i.e., agranulocytes) are considered as the young and immature hemocytes of two types of matured

hemocytes (9). Our pseudo temporal ordering analysis revealed that the hemocytes of *M. japonicus*

differentiate from a single subpopulation into four major populations. The differentiation process of *M.*

japonicus hemocytes was continuous, not discrete, which was in agreement with previous arguments on the

crustacean hematopoiesis mechanism (9, 43, 44).

194
195 **Figure 3.** Pseudo temporal ordering of hemocyte lineages. (A) Violin plots displaying normalized expression
196 levels of each cell cycle related genes across all clusters. (B) Visualization of clusters (from Figure 1B) onto
197 the pseudo time map using monocle 3.

Expression of cell growth-related genes

The exploration of cluster-specific markers revealed that cell growth-related genes were specifically
expressed in certain clusters. Therefore, we highlighted five genes that are predicted to be involved in cell
growth and differentiation: *VWDE-like*, *TINAGL*, *PDGF/VEGF-related factor 1*, *IGFBP-related protein 1*,
and *CHF-like* (Fig. 4, Fig. S5, and Dataset S5).

VWDE-like was highly expressed in clusters Hem 1 to Hem 3 (Fig. 4 A and B). The expression of
VWDE is a common feature of blastemas, which are capable of regenerating limbs and fins in a variety of
highly regenerative species (45). VWDE contains several epidermal growth factor (EGF) domains and is
expected to be a downstream effector once a blastema has been established, but it is not a driver of blastema
formation (45). VWDE-like of *M. japonicus* was expressed more strongly in Hem 2 than in Hem 1, indicating
that VWDE-like also works as a downstream effector against Hem 1, which is predicted to be composed of
undifferentiated hemocytes.

TINAGL, which was expressed in Hem 3 and 4, is a secreted extracellular protein that is essential
for early angiogenesis in developing zebrafish embryos (46) and humans (47). TINAGL of *M. japonicus* has
been proposed to participate in angiogenic cell differentiation. TINAGL is also known as a suppressor of
cancer progression and metastasis by binding to receptors of EGF in humans (48). Interestingly, TINAGL of
M. japonicus was expressed in Hem 3 and Hem 4, in which the expression of VWDE-like was weakened

(Fig. 4 A and B). This suggests that TINAGL might suppress the differentiation function of VWDE-like in Hem 3 and 4 in shrimp.

Cells expressing PDGF/VEGF-related factors were dominant in Hem 4 and Hem 5 (Fig. 4 A and B). The vascular endothelial growth factor (VEGF) signaling pathway is essential for vasculogenesis, cell proliferation, and tumor migration in mammals (49, 50). Furthermore, in *Drosophila*, VEGF homologs control the number of circulating hemocytes (51). The high expression of PDGF/VEGF-related factors in Hem 4 and Hem 5 showed that Hem 3 would differentiate into Hem 5 through Hem 4 via the VEGF signaling pathway.

Hem 6 to Hem 9 expressed insulin-like growth factor (IGF) binding protein (IGFBP) (Fig. 4 A and B). IGFBP delivers IGFs to the target cells in mammal studies and is essential for cell growth or differentiation (52). The high expression of a receptor of the insulin-like peptide at mature hemocytes in the mosquito suggests that the insulin signaling pathway regulates hemocyte proliferation (53). The silencing and overexpression of IGFBP caused a decrease and increase in the growth of hemocytes, respectively, in the loss and gain function study of abalone *Haliotis diversicolor* (54). These studies indicate that the IGFBP-related insulin signaling pathway is important for hemocyte proliferation and differentiation in invertebrates. IGFBP might play an essential role in the differentiation of hemocytes from Hem 4 to Hem 6–9 in shrimp. The high expression of IGFBP is determined in the brain and gonads of *Litopenaeus vannamei* (55). This fact also suggests that IGFBP plays a possible role in organ growth and maturation in shrimp.

Up to this point, the analyzed genes were proliferation- and differentiation-promoting, but there was also the specific expression of the hemocyte homeostasis regulatory gene, crustacean hematopoietic factor (CHF) (43) (Fig. 4 A and B). CHF is a hematopoietic factor of crayfish, and the silencing of CHF leads to an increase in the apoptosis of cells in HPT, and a reduction in the number of circulating hemocytes (33). Additionally, the silencing of laminin, a receptor of CHF, reduces the number of circulating hemocytes by decreasing the number of agranulocytes, as opposed to granulocytes, in *P. vannamei* (55). CHF-like was expressed at cluster Hem 9 (Fig. 4 A and B), in which hemocytes are predicted to be matched here. Taken together, CHF-like expressed from matured hemocytes, Hem 9, might work as a hematopoietic factor against agranulocytes or regulate the homeostasis of agranulocytes.

Figure 4. Cell growth-related gene expression across all clusters. (A) Expression pattern of the cell growth-related genes across cell clusters. (B) Violin plots displaying normalized expression levels of each cell growth-related genes across all clusters.

Expression of immune-related genes in single cells

Hemocytes of shrimp play key roles in their immunity; therefore, we selected immune-related genes from 3,334 commonly expressed genes, and then analyzed their expression levels to deduce the detailed immune functions of each cluster.

Antimicrobial peptides (AMPs) play the most important role in the immunity of shrimps and are well known to be stored in granulocytes (56, 57). The expression patterns of AMPs revealed that the major AMPs of penaeid shrimp were expressed in clusters Hem 5 to Hem 9 (Fig. 5 A-F, Fig. S6 and Dataset S6). Therefore, Hem 5 to Hem 9 were predicted to be granulocytes. These AMP-expressing clusters were broadly classified into two groups. One group was Hem 5, which expressed several AMPs, such as ALF-A1 and c-type lysozyme (Fig. 5 D and F), but also expressed other immune-related genes different from AMPs: *chitin-binding protein* and *virus responsible protein* (*VRP*) (Fig. 2). Chitin-binding protein is located on the cell surface and interacts with virus envelope proteins in shrimp (58). VRP is distributed in granulocytes, and infection by pathogenic viruses causes an increase in VRP transcripts (59). The other group is Hem 6 to 9, which strongly expresses major AMPs, such as penaeidin, stylicin, and SWD, suggesting that this group corresponds to the most studied immune-related cell subpopulation to date. The level of penaeidin-positive hemocytes is increased after bacterial infection (60), but is reduced after a virus infection (61). Stylicin and SWD mRNA expression was decreased after virus infection in *M. japonicus* and *P. monodon*.

Together, the functions of these two groups can be characterized as follows. In Hem 5, VRP, whose level is known to be increased by viral infection, was specifically expressed, while AMPs were downregulated. Thus, Hem 5 is predicted to be the group that contributes to the immune response against viruses. Additionally, a virus infection causes an increase in the expression of both VEGF and its receptors

in hemocytes to regulate a downstream signaling pathway (62-64). The high expression of PDGF/VEGF-

related factors at Hem 5 (Fig. 2 and 4) supports our prediction that Hem 5 plays an essential role in shrimp

biodefense against viruses. Previously, a subtype of granulocytes, termed semi-granular cells (SGC), were

isolated from *L. vannamei*, in which lysozyme was highly expressed. It is possible that Hem 5 in *M. japonicus*

corresponds to SGC found in *L. vannamei* because c-type lysozyme was strongly expressed in Hem 5, in our

study. Conversely, the AMPs specifically expressed at Hem 6–9 were upregulated by bacterial infection.

Therefore, these clusters contribute to bacterial defense. Different types of ALF play different roles in shrimp

immunity, which improves the synergism in shrimp antimicrobial defenses (65). The expression patterns of

ALFs were also different in each cluster (Fig. 5 C and D, Fig. S6 and Dataset S6), suggesting that different

clusters have different immunological functions.

Some of the hemocyte-type-specific markers related to their immune function have been studied in

crayfish: prophenoloxidase (proPO) in matured hemocytes, copper/zinc superoxide dismutase (SOD) in

SGC, and Kazal-type proteinase inhibitor (KPI) in granular cells (GC), and transglutaminase (TGase) in

immature cells (66). We BLAST searched homologies between these genes and common expressed genes to

check their potential as specific markers. As a result, 31 genes were found to have homology with the genes

listed above (Fig. S6 and Dataset S6). TGase was strongly expressed in immature clusters Hem 1 and Hem

2, and can, therefore, be used as a marker of immature hemocytes (Fig. 5 I and Fig. S6). ProPOs were not

highly expressed in the present single-cell study. Both KPI and SOD were mostly expressed at both Hem 8

and Hem 9, and their expression levels were similar between these clusters (Fig. 5 G and H and Fig. S6).

These results suggest that the functional segregation of hemocytes in shrimp is different from that in crayfish,

in which these molecular markers are expressed distinctly between GC and SGC. The KPI of *M. japonicus*

is expressed in only some hemocytes in healthy shrimps, and bacterial infection causes an increase in KPI

expression (67). Therefore, it can be predicted that the number of KPI-positive hemocytes in Hem 8 and 9

would be increased upon a pathogen infection, in shrimp.

In the mosquito, scRNA-seq revealed a new subpopulation called "antimicrobial granulocytes" that

expressed characteristic AMPs (22). Similarly, in *M. japonicus*, the expression patterns of immune-related

genes were also different among certain clusters, suggesting that shrimp hemocytes are more heterogeneous

than previously thought. It is anticipated that the class of granulocytes discussed in previous studies is actually

a mixture of clusters exhibiting different roles.

298
Figure 5. Expression pattern of the immune-related genes across cell clusters. (A) Penaeidin-II, (B) single
WAP domain containing protein, (C) ALF-CA, (D) ALF-A1, (E) stylicin, (F) c-type lysozyme, (G) KPI, (H)
SOD, and (I) TGase.

Validation of marker genes and the relationship between clusters and morphology

Our scRNA-seq results revealed nine major subpopulations and their marker genes, and the possible
differentiation trajectory of *M. japonicus* hemocytes. Next, we examined the correlation between the
morphology and expression of marker genes. Two major populations of hemocytes were sorted based on the
forward versus side scatter plot obtained using microfluidic-based fluorescence-activated cell sorter (FACS).
The sorted populations were observed using microscopy. FACS was able to separate hemocytes into two
morphologically different populations (Fig. 6 A–E): smaller cells with low internal complexity in region 1
(R1) (50.7% ± 5.2%) and larger cells with high internal complexity in region 2 (R2) (47.7% ± 3.4%). From
DIC and dye staining imaging (Fig. 6 A and B), we observed that cells in the R1 region contained no or few
granules in the cytoplasm. The nucleus occupied a large portion of the volume in these cells (Fig. 6 C).
Conversely, those in the R2 region had many granules in the cytoplasmic region, which occupied a large
portion of cells (Fig. 6 D).

We conducted qRT-PCR analysis to determine the expression of some representative genes in these
populations. The results showed that the ΔΔCt values of the transcripts of HemTGase were higher in R1
hemocytes than in R2 hemocytes, while the values of penaeidin, crustin, stylicin, proPO, and Cu/ZnSOD2
were higher in R2 hemocytes (Fig. 6 F). The ΔΔCt values of transcripts of c-type lysozyme and VRP were

similar in R1 and R2 hemocytes. The combination of FACS and qRT-PCR results confirmed that the gene

expression of the two populations in shrimp hemocytes was roughly divided based on morphology, that is,

agranulocytes and granulocytes, which is consistent with our scRNA-seq analysis. Granulocytes found in the

R2 region expressed major AMPs, such as penaeidin, crustin, and stylicin, suggesting that they consist of

clusters Hem 6 to Hem 9. However, c-type lysozyme and VRP, which are markers of Hem 5 (Fig. 2), were

expressed in cells in both R1 and R2 regions, indicating that Hem 5 exists in both populations and is

indistinguishable from the morphological characteristics. The number of cells occupying R1 and R2 was

50.7% ± 5.2% and 47.7% ± 3.4% (from $n = 4$), while the total number of cells classified as Hem 1 − 4 and

Hem 5 − 9 in our single-cell analysis was 50.4% (1,363 cells) and 49.6% (1,341 cells), respectively.

Figure 6. Morphological analysis of hemocytes and transcript profiles based on morphology. (A) DIC

(Differential Interference Contrast) image of unsorted total hemocytes. (B) Dye stained total hemocytes. (C)

DIC imaging and dye staining of R1 sorted hemocytes. (D) DIC imaging and dye staining of R2 sorted

hemocytes. (E) FACS (fluorescence-activated cell sorting) analysis of hemocytes. Based on the FSC (forward

scatter) and SSC (side scatter) two-dimensional space, two regions (R1 and R2) were obtained. (F)

Differential gene expression analysis between R1 and R2 of hemocytes sorted using FACS. ΔΔCt values

were analyzed using qRT-PCR. Higher ΔΔCt values indicate a higher accumulation of mRNA transcripts.

Discussion

Our single-cell transcriptome analysis revealed that there are nine subpopulations of hemocytes in

shrimp. This result differs from any previous classification strategy based on various approaches, such as

simple staining (7-9), monoclonal antibodies (18), flow cytometry (68), and lectin-binding profiles (69). We

hope that future studies will clarify the relationships between these phenotypic features and our
classification to comprehend the role of each cluster in the immune system of shrimp.

The cluster-specific markers and cell proliferation-related genes found here helped us to understand
how shrimp hemocytes differentiate. A strong expression of TGase, cell proliferation, and G2/M state-related
genes in Hem 1 suggested that hemocytes in this cluster are oligopotent and located upstream in the
differentiation process. In crustaceans, especially shrimp and crayfish, it is known that hemocytes are
produced in HPT, and that differentiated hematopoietic cells from HPT circulate in the body fluid (9, 36,
66). However, G2/M state hemocytes of *M. japonicus* exist in the hemolymph and account for only 0.63% ±
0.28% of the circulating hemocytes (70). Hem 1 accounts for only 2.4% of the analyzed cells. This similarity
in the fraction also indicates that only a very small fraction of oligopotent or initial state hemocytes exist
among the circulating hemocytes.

Our results also revealed that the growth-related genes were expressed at specific clusters (Fig. 4 A
and B). From these results, we propose that shrimp hemocytes differentiate as follows (Fig. 7 A and B): 1)
Hem 1 is the initial state of circulating hemocytes and has an oligopotent ability, which leaked out from HPT;
2) Hem 3 is differentiated from Hem 1 by VWDE-like through Hem 2; 3) Hem 4 is differentiated from Hem
3 by TINAGL; 4) Hem 5 is differentiated from Hem 4 by PDGF/VEGF-related factor; 5) Hem 7 to Hem 9
are differentiated from Hem 4 by IGFBP-related protein through Hem 6; 6) CHF-like is expressed in Hem 9
to maintain the immature hemocytes, Hem 1 to Hem 4. Loss and gain function studies of these differentiating
factors are necessary to prove the full differentiation process of penaeid shrimp hemocytes. Since we
identified a group of proliferating cells and differentiation factors here, the next step is to establish a way to
isolate and cultivate them. The markers of each cluster identified here will be good tools for isolating specific
cell types.

Crayfish's hemocytes and *Drosophila* hemocytes have been classified into three and four major
types according to the marker genes, respectively (23, 36). However, these crayfish or *Drosophila* marker
genes were found to be inadequate as cluster-specific markers for penaeid shrimp in our study. Insects and
crustaceans are thought to become independent about 500 million years ago (71, 72), and shrimp and crayfish
are thought to have become evolutionarily independent about 450 million years ago (73). Thus, it is
straightforward to reason that the functions of these genes have changed during the evolutionary process.
Therefore, we should not simply argue that the morphological and functional similarities between shrimp
and *Drosophila*/crayfish hemocytes are the same.

Most of the markers that were characteristically expressed in Hem 1 and Hem 4 could not be
functionally predicted by the BLAST search (Fig. 2). We speculate that the characteristic genes of Hem 1 are
associated with cell duplication and that Hem 4 is associated with the regulation of cell differentiation. The
division and differentiation mechanisms of shrimp hemocytes are still largely unknown, and no techniques
on culture shrimp hemocytes *in vitro* have been reported. The analysis of these unknown gene characteristics

may reveal these mechanisms. Notably, no specific marker gene was found in Hem 6, probably because these
cells are yet to be specialized, unlike those in Hem 7 to Hem 9.

In conclusion, we succeeded in classifying shrimp hemocytes into nine subpopulations based on
their transcriptional profiles, while they were only classified into two groups using FACS. Furthermore, our
results imply that hemocytes differentiate from a single initial population. Although we have not yet
successfully cultured crustacean hemocytes in passaging cultures, information on these subpopulations and
marker genes will provide a foothold for hemocyte culture studies. Despite our success in the classification
of hemocytes, we have not yet been able to fully understand the functions of each hemocyte group in detail.
One reason for this is that the functions of some marker genes are still unknown. The present single-cell
transcriptome data serves as a platform providing the necessary information for the continuous study of
shrimp genes and their functions. Additionally, we have only determined the subpopulations of hemocytes
in the normal state, so that our next goal will be to analyze the hemocytes in the infected state of certain
diseases at different levels of cell maturation. In this way, we will be able to identify the major subpopulations
that may work against the infectious agent. Likewise, it will be interesting to see how hemocytes from
fertilized eggs mature. Additionally, single-cell analysis of hematopoietic tissues can also be expected to
reveal more detailed differentiation mechanisms. Unlike terrestrial invertebrates, such as *Drosophila* and
mosquitoes, shrimps are creatures that live in the ocean and are evolutionarily distant from each other.
Understanding the immune system of shrimp species will require continuous effort, but we believe that it
will provide efficient solutions to aquaculture problems.

395
Figure 7. Model of the roles of cell growth-related genes in hemocyte differentiation. (A) Shrimp hemocyte
differentiation process, 1) Hem 1 is the initial state of circulating hemocytes, which have oligopotent ability,

which leaked out from the hematopoietic tissue (HPT); 2) Hem 3 is differentiated from Hem 1 by VWDE-

like through Hem 2; 3) Hem 4 is differentiated from Hem 3 by TINAGL; 4) Hem 5 is differentiated from

Hem 4 by PDGF/VEGF-related factor; 5) Hem 7 to Hem 9 are differentiated from Hem 4 by IGFBP-related

protein through Hem 6; 6) CHF-like is expressed in Hem 9 to maintain the immature hemocytes, Hem 1 to

Hem 4. (B) Schematics of cell growth or homeostasis-related gene expression across clusters. The gene

expression distribution was cluster-specific.

Materials and Methods

Shrimp and cell preparation

Twenty-three female kuruma shrimp, *Marsupenaeus japonicus*, with an average weight of 20 g,

were purchased from a local distributor and maintained in artificial seawater with a 34 ppt salinity with a

recirculating system at 20°C. Hemolymph was collected using an anticoagulant solution suitable for penaeid

shrimp (7) from an abdominal site. The collected hemolymph was centrifuged at 800 x *g* for 10 min to collect

the hemocytes, which were then washed twice with PBS and the osmolarity was adjusted to kuruma shrimp

(KPBS: 480 mM NaCl, 2.7 mM KCl, 8.1 mM $Na_2HPO_4·12H_2O$, 1.47 mM KH_2PO_4, pH 7.4).

Preparation of expressing gene list of hemocytes

De novo assembled transcript data were prepared as a reference for mapping Drop-seq data, because

the genome sequence of *M. japonicus* is still unknown. To improve the quality of the *de novo* assembled

transcript sequences, we prepared long-read mRNA sequences using MinION (Oxford Nanopore

Technologies) direct RNA sequencing to conduct hybrid *de novo* assembly. Poly(A) tailed RNA was purified

from 58 μg of total RNA from the hemocytes of 16 shrimp using Dynabeads Oligo(dT)$_{25}$ (Thermo Fisher

Scientific), and 500 ng of poly-(A) RNA was ligated to adaptors using a direct RNA sequencing kit (Oxford

Nanopore Technologies) according to the manufacturer's manual version DRS_9080_v2_revL_14Aug2019.

Finally, 44 ng of the library was obtained and sequenced using MinION, by using a MinION flow cell R9.4.1

(Oxford Nanopore Technologies). All sequencing experiments were performed using MinKNOW v3.6.5

without base calling. Raw sequence data were then base-called using Guppy v3.6.1. Once the raw signal from

the MinION fast5 files was converted into fastq files, the sequencing errors were corrected using TALC v1.01

(74)(https://gitlab.igh.cnrs.fr/lbroseus/TALC) by using the Illumina short reads sequence of kuruma shrimp

hemocytes (DDBJ Sequence Read Archive (DRA) accession number DRA004781). The corrected long-read

sequences from MinION and short-read sequences from Illumina Miseq were hybrid *de novo* assembled

using rnaSPAdes v3.14.1 (75)(https://cab.spbu.ru/software/rnaspades/) and Trinity 2.10.0

(76)(https://github.com/trinityrnaseq/trinityrnaseq/wiki). All assembled *de novo* transcripts were merged and

subjected to the EvidentialGene program v2022.01.20 (http://arthropods.eugenes.org/EvidentialGene/) to

remove similar sequences with a default parameter. The remaining sequences were renamed as Mj-XXX and

used as a hemocyte-expressing gene list. The assembled sequences and code used to perform base-calling

and *de novo* assembly are available on GitHub at https://github.com/KeiichiroKOIWAI/Drop-
seq_on_shrimp.

Single-cell and single-bead encapsulation by a microfluidic device and exonuclease and reverse
transcribe reaction on a bead

439   The drop-seq procedure was used to encapsulate single hemocytes and single mRNA capture beads
together into fL-scale microdroplets, as previously described (19). The following steps were performed in
triplicates for three shrimp individuals. Briefly, the self-built Drop-seq microfluidic device was prepared by
molding polydimethylsiloxane (PDMS; Sylgard 184, Dow Corning Corp.) from the microchannel structure
formed by the negative photoresist (SU-8 3050, Nippon Kayaku Co.). Using this device, droplets containing
a cell and a Barcoded Bead SeqB (ChemGenes Corporation) were produced up to 2 mL per sample using a
pressure pump system (Droplet generator, On-chip Biotechnologies Co., Ltd.). During the sample
introduction, the vial bottles containing cells and beads were shaken using a vortex mixer to prevent
sedimentation and aggregation (77). Droplets were collected from the channel outlet into the 50-mL corning
tube and incubated at 80°C for 10 min in a water bath to promote hybridization of the poly(A) tail of mRNA
and oligo d(T) on beads. After incubation, droplets were broken promptly and barcoded beads with captured
transcriptomes were reverse transcribed using Maxima H Minus Reverse Transcriptase (Thermo Fisher
Scientific) at RT for 30 min, then at 42°C for 90 min. Then, the beads were treated with Exonuclease I (New
England Biolabs) to obtain single-cell transcriptomes attached to microparticles (STAMP). The first-strand
cDNAs on beads were amplified using PCR. The beads obtained above were distributed throughout PCR
tubes (1,000 beads per tube), wherein 1× KAPA HiFi HS Ready Mix (KAPA Biosystems) and 0.8 μM 1st
PCR primer were included in a 25 μL reaction volume. PCR amplification was achieved using the following
program: initial denaturation at 95°C for 3 min; 4 cycles at 98°C for 20 s, 65°C for 45 s, and 72°C for 6 min;
12 cycles of 98°C for 20 s, 67°C for 20 s, and 72°C for 6 min; and a final extension at 72°C for 5 min. The
amplicons were pooled, double-purified with 0.9× AMPure XP beads (Beckman Coulter), and eluted in 100
μL of ddH$_2$O. Sequence-ready libraries were prepared according to Picelli, Bjorklund (78). A total of 1 ng of
each cDNA library was fragmented using home-made Tn5 transposome in a solution containing 10 mM
TAPS-NaOH (pH 8.5), 5 mM MgCl$_2$, and 10% dimethylformamide at 55°C for 10 min. The cDNA fragments
were purified using a DNA Clean & Concentrator Kit (Zymo Research) and eluted in 25 μL of ddH$_2$O. The
index PCR reaction was performed by adding 12 μL of the elute to a mixture consisting of 1× Fidelity Buffer,
0.3 mM dNTPs, 0.5 U KAPA HiFi DNA polymerase (KAPA Biosystems), 0.2 μM P5 universal primer, and
0.2 μM i7 index primer. Each reaction was achieved as follows: initial extension and subsequent denaturation
at 72°C for 3 min and 98°C for 30 s; 12 cycles of 98°C for 10 s, 63°C for 30 s, and 72°C for 30 s; and a final
extension at 72°C for 5 min. The amplified library was purified using 0.9× AMPure XP beads and sequenced
(paired-end) on an Illumina NextSeq 500 sequencer (NextSeq 500/550 High Output v2 kit (75 cycles); 20
cycles for read1 with custom sequence primer, 8 cycles for index read, 64 cycles for read2. Before performing

the Drop-seq on kuruma shrimp hemocytes, we validated the protocol by performing the same procedure

using a mixture of HEK293 and NIH3T3 cells and sequencing the test library using a Miseq Reagent Kit v3

(150 cycles).

Analysis of single-cell data

Paired-end reads were processed and mapped to the reference *de novo* assembled gene list of

hemocytes following the Drop-seq Core Computational Protocol version 2.0.0, and the corresponding Drop-

seq tools v2.3.0 (https://github.com/broadinstitute/Drop-seq) provided by McCarroll Lab

(http://mccarrolllab.org/dropseq/). The Picard suite (https://github.com/broadinstitute/picard) was used to

generate the unaligned bam files. The steps included the detection of barcode and UMI sequences, filtration

and trimming of low-quality bases and adaptors or poly(A) tails, and the alignment of reads using bowtie2

v2.4.1 (79) (http://bowtie-bio.sourceforge.net/bowtie2/index.shtml). The cumulative distribution of reads

from the aligned bam files was obtained using BAMTagHistogram, and the number of cells was inferred

using Drop-seq tools.

Data integration

After digital expression data from 3 shrimps were read using Seurat v3.2.1 (80,

81)(https://satijalab.org/seurat/), SCTransform (81) was performed to remove the technical variation and to

select common expressing genes, while retaining biological heterogeneity. We ran a PCA using the

expression matrix of the top 3,000 most variable genes. The total number of principal components (PCs)

required to compute and store was 50. The UMAP was then performed using the following parameters:

n.neighbors, min.dist, and n.components were 10L, 0.1, and 2, respectively, to visualize the data in the two-

dimensional space, and then the clusters were predicted with a resolution of 0.5.

Functional prediction of commonly expressed genes

The predicted functions of selected assembled sequences as common expressing genes across three

replicates were searched using BLAST program v2.2.31 (82, 83)

(https://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/LATEST/) on penaeid shrimp identical proteins

(downloaded from NCBI Identical Protein Groups on 19th of August, 2020) with the blastx parameter of e-

value as 0.0001 and num_alignments as 3. Then, the functions of each cluster were predicted based on the

marker genes. Marker genes were predicted using the Seurat FindAllMarkers tool with the following

parameters: min.pct as 0.5, logfc.threshold as 1, and test.

Visualization of genes with distinctive functions on single hemocytes

To visualize the genes with distinctive functions of shrimp, we extracted the distinct sequences from

commonly expressed genes based on their blastx results against penaeid shrimp identical proteins. Here, we

focused on the cell growth-related genes listed in Dataset S5 and on immune-related genes, such as antimicrobial peptides (AMPs), transglutaminase, copper/zinc superoxide dismutase, and Kazal-type proteinase inhibitors listed in Dataset S6. From these genes that showed a characteristic expression among single-cell data, dot plots, violin plots, or feature plot visualizations were applied using Seurat functions.

Comparison with *Drosophila* marker genes

To check whether the *Drosophila* marker genes are applicable to shrimp, we performed a BLAST search on *the Drosophila* cell cycle and cell type markers (https://github.com/hbc/tinyatlas). Common genes among the three shrimp species were tblastx searched for *Drosophila melanogaster* genes (dmel-all-gene-r6.34.fasta; downloaded from FlyBase https://flybase.org/) with the parameters of e-value as 0.0001 and num_alignments as 3.

Pseudo temporal ordering of cells using Monocle 3

The integrated data of Seurat were transferred to Monocle 3 (84) (https://github.com/cole-trapnell-lab/monocle3) to calculate a cell trajectory using the learn_graph function. We assigned the start point based on the expression of cell proliferation-related genes and *Drosophila* marker genes of the cell cycle.

Cell sorting of hemocytes and qRT-PCR of marker genes

To validate the Drop-seq results on hemocytes, populations of hemocytes in the forward scatter (FSC) and side scatter (SSC) two-dimensional space were sorted using a microfluidic cell sorter (On-chip sort, On-chip Biotechnologies Co., Ltd.) from four shrimp individuals (Fig. S7). In the FSC/SSC two-dimensional space, two main populations were predicted as Region1 (R1): small/simple and Region2 (R2): large/complexity populations, which were defined as agranulocytes and granulocytes, respectively. After sorting, some sorted hemocytes were immediately fixed in 2% formalin in KPBS and stained with a May-Grunwald and Giemsa staining solution to observe the cellular components. Non-stained and stained hemocytes were subjected to microscopy IX71 (Olympus Corporation) to observe their structures.

Total RNA was also collected from sorted cells and pre-sorted cells. The concentration of RNA was measured using a nanodrop, and cDNA was transcribed using a High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific). Constructed cDNA was diluted five times with TE buffer and subjected to qRT-PCR using KOD SYBR qPCR (TOYOBO Co. Ltd.), following the manufacturer's protocol. The expression of each gene was calculated using the ΔΔCT method against elongation factor-1 alpha and total hemocytes.

Data files and analysis code

The raw sequence data of newly sequenced *M. japonicus* transcriptomic reads were archived in the DDBJ Sequence Read Archive (DRA) of the DNA Data Bank of Japan as follows: MinION mRNA direct

sequencing: DRA010948; Drop-seq shrimp rep1: DRA010950; shrimp rep2: DRA010951; shrimp rep3:

DRA010952; mixture sample of HEK293 and 3T3 cells: DRA010949. The Seurat digital expression data

were archived in the Genomic Expression Archive of the DNA Data Bank of Japan: E-GEAD-403. Fast5

data of MinION direct RNA sequencing will be made available upon request from the authors. The code used

to perform *de novo* assembly, clustering, and marker analysis is available on GitHub at

https://github.com/KeiichiroKOIWAI/Drop-seq_on_shrimp. The key resources are listed in Dataset S7.

Acknowledgments

We would like to thank Fumiko Sunaga for her technical support in performing cell sorting and

analysis, and Editage (www.editage.com) for English language editing. This work was supported by JSPS

KAKENHI (JP19J00539 and JP20K15603) to K. Koiwai; a Grant-in-Aid for Scientific Research on

Innovative Areas (17H06425) to K. Kikuchi.

Competing Interest Statement: The authors declare no conflict of interest.

References

1. FAO. Fishery and aquaculture statistics 2018/FAO annuaire. Rome: FAO; 2020.

2. Jiravanichpaisal P, Lee BL, Söderhäll K. Cell-mediated immunity in arthropods: hematopoiesis,
coagulation, melanization and opsonization. Immunobiology. 2006;211(4):213-36.

3. Tassanakajon A, Somboonwiwat K, Supungul P, Tang S. Discovery of immune molecules and their
crucial functions in shrimp immunity. Fish Shellfish Immunol. 2013;34(4):954-67.

4. Flegel TW. A future vision for disease control in shrimp aquaculture. Journal of the World Aquaculture
Society. 2019;50(2):249-66.

5. Zhang X, Yuan J, Sun Y, Li S, Gao Y, Yu Y, et al. Penaeid shrimp genome provides insights into
benthic adaptation and frequent molting. Nat Commun. 2019;10(1):356.

6. Bauchau AG. Cruataceans. Invertabrate blood cells. 1981;2:385-420.

7. Söderhäll K, Smith VJ. Separation of the haemocyte populations of *Carcinus maenas* and other marine
decapods, and prophenoloxidase distribution. Dev Comp Immunol. 1983;7(2):229-39.

8. Johansson MW, Keyser P, Sritunyalucksana K, Söderhäll K. Crustacean haemocytes and
haematopoiesis. Aquaculture. 2000;191(1-3):45-52.

9. van de Braak CB, Botterblom MH, Liu W, Taverne N, van der Knaap WP, Rombout JH. The role of the
haematopoietic tissue in haemocyte production and maturation in the black tiger shrimp (*Penaeus*
monodon). Fish Shellfish Immunol. 2002;12(3):253-72.

10. Kondo M, Tomonaga S, Takahashi Y. Granulocytes with cytoplasmic deposits of kuruma prawn.
Aquaculture Science. 2012;60(1):151-2.

11. Dantas-Lima JJ, Tuan VV, Corteel M, Grauwet K, An NTT, Sorgeloos P, et al. Separation of *Penaeus*
vannamei haemocyte subpopulations by iodixanol density gradient centrifugation. Aquaculture.
2013;408-409:128-35.

12. Rodriguez J, Boulo V, Mialhe E, Bachère E. Characterisation of shrimp haemocytes and plasma
components by monoclonal antibodies. J Cell Sci. 1995;108:1043-50.

13. van de Braak CB, Taverne N, Botterblom MH, van der Knaap WP, Rombout JH. Characterisation of
different morphological features of black tiger shrimp (*Penaeus monodon*) haemocytes using
monoclonal antibodies. Fish Shellfish Immunol. 2000;10(6):515-30.

14. Sung HH, Wu PY, Song YL. Characterisation of monoclonal antibodies to haemocyte subpopulations
of tiger shrimp (*Penaeus monodon*): immunochemical differentiation of three major haemocyte types.
Fish Shellfish Immunol. 1999;9(3):167-79.

15. Sung HH, Sun R. Use of monoclonal antibodies to classify hemocyte subpopulations of tiger shrimp
(*Penaeus monodon*). Journal of Crustacean Biology. 2002;22(2):337-44.

16. Winotaphan P, Sithigorngul P, Muenpol O, Longyant S, Rukpratanporn S, Chaivisuthangkura P, et al.
Monoclonal antibodies specific to haemocytes of black tiger prawn *Penaeus monodon*. Fish Shellfish
Immunol. 2005;18(3):189-98.

17. Lin Y, Zhan W, Li Q, Zhang Z, Wei X, Sheng X. Ontogenesis of haemocytes in shrimp
(*Fenneropenaeus chinensis*) studied with probes of monoclonal antibody. Dev Comp Immunol.
2007;31(11):1073-81.

18. Xing J, Chang Y, Tang X, Sheng X, Zhan W. Separation of haemocyte subpopulations in shrimp
Fenneropenaeus chinensis by immunomagnetic bead using monoclonal antibody against granulocytes.
Fish Shellfish Immunol. 2017;60:114-8.

19. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly parallel genome-wide
expression profiling of individual cells using nanoliter droplets. Cell. 2015;161(5):1202-14.

20. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol. 2014;32(4):381-6.

21. Soneson C, Robinson MD. Bias, robustness and scalability in single-cell differential expression analysis. Nat Methods. 2018;15(4):255-61.

22. Raddi G, Barletta ABF, Efremova M, Ramirez JL, Cantera R, Teichmann SA, et al. Mosquito cellular immunity at single-cell resolution. Science. 2020;369(6507):1128-32.

23. Tattikota SG, Cho B, Liu Y, Hu Y, Barrera V, Steinbaugh MJ, et al. A single-cell survey of *Drosophila* blood. eLife. 2020;9:e54818.

24. Carmona SJ, Teichmann SA, Ferreira L, Macaulay IC, Stubbington MJ, Cvejic A, et al. Single-cell transcriptome analysis of fish immune cells provides insight into the evolution of vertebrate immune cell types. Genome Res. 2017;27(3):451-61.

25. Severo MS, Landry JJM, Lindquist RL, Goosmann C, Brinkmann V, Collier P, et al. Unbiased classification of mosquito blood cells by single-cell genomics and high-content imaging. Proc Natl Acad Sci U S A. 2018;115(32):E7568-E77.

26. Lv D, Jia F, Hou Y, Sang Y, Alvarez AA, Zhang W, et al. Histone acetyltransferase KAT6A upregulates PI3K/AKT signaling through TRIM24 binding. Cancer Res. 2017;77(22):6190-201.

27. Andang M, Hjerling-Leffler J, Moliner A, Lundgren TK, Castelo-Branco G, Nanou E, et al. Histone H2AX-dependent GABA(A) receptor regulation of stem cell proliferation. Nature. 2008;451(7177):460-4.

28. Cheung HC, Hai T, Zhu W, Baggerly KA, Tsavachidis S, Krahe R, et al. Splicing factors PTBP1 and PTBP2 promote proliferation and migration of glioma cell lines. Brain. 2009;132(Pt 8):2277-88.

29. Shibayama M, Ohno S, Osaka T, Sakamoto R, Tokunaga A, Nakatake Y, et al. Polypyrimidine tract-binding protein is essential for early mouse development and embryonic stem cell proliferation. FEBS J. 2009;276(22):6658-68.

30. Jiao H, Dong P, Yan L, Yang Z, Lv X, Li Q, et al. TGF-beta1 Induces polypyrimidine tract-binding protein to alter fibroblasts proliferation and fibronectin deposition in keloid. Sci Rep. 2016;6:38033.

31. Lin X, Söderhäll K, Söderhäll I. Transglutaminase activity in the hematopoietic tissue of a crustacean, *Pacifastacus leniusculus*, importance in hemocyte homeostasis. BMC Immunol. 2008;9:58.

32. Huang CC, Sritunyalucksana K, Söderhäll K, Song YL. Molecular cloning and characterization of tiger shrimp (*Penaeus monodon*) transglutaminase. Dev Comp Immunol. 2004;28(4):279-94.

33. Lin X, Söderhäll K, Söderhäll I. Invertebrate hematopoiesis: an astakine-dependent novel hematopoietic factor. J Immunol. 2011;186(4):2073-9.

34. Huerta-Cepas J, Szklarczyk D, Heller D, Hernandez-Plaza A, Forslund SK, Cook H, et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 2019;47(D1):D309-D14.

35. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, et al. Fast genome-wide functional annotation through orthology assignment by eggNOG-Mapper. Mol Biol Evol. 2017;34(8):2115-22.

36. Söderhäll I. Crustacean hematopoiesis. Dev Comp Immunol. 2016;58:129-41.

37. De Lucia F, Ni JQ, Vaillant C, Sun FL. HP1 modulates the transcription of cell-cycle regulators in *Drosophila melanogaster*. Nucleic Acids Res. 2005;33(9):2852-8.

38. Paro R, Hogness DS. The polycomb protein shares a homologous domain with a heterochromatin-associated protein of *Drosophila*. Proc Natl Acad Sci U S A. 1991;88(1):263-7.

39. Mohan M, Bartkuhn M, Herold M, Philippen A, Heinl N, Bardenhagen I, et al. The *Drosophila*
insulator proteins CTCF and CP190 link enhancer blocking to body patterning. EMBO J.
2007;26(19):4203-14.

40. Rasko JE, Klenova EM, Leon J, Filippova GN, Loukinov DI, Vatolin S, et al. Cell growth inhibition by
the multifunctional multivalent zinc-finger factor CTCF. Cancer Res. 2001;61(16):6002-7.

41. Scott CL, Gil J, Hernando E, Teruya-Feldstein J, Narita M, Martinez D, et al. Role of the chromobox
protein CBX7 in lymphomagenesis. Proc Natl Acad Sci U S A. 2007;104(13):5389-94.

42. Junkunlo K, Söderhäll K, Noonin C, Söderhäll I. PDGF/VEGF-related receptor affects
transglutaminase activity to control cell migration during crustacean hematopoiesis. Stem Cells Dev.
2017;26(20):1449-59.

43. Lin X, Söderhäll I. Crustacean hematopoiesis and the astakine cytokines. Blood. 2011;117(24):6417-
24.

44. Noonin C, Lin X, Jiravanichpaisal P, Söderhäll K, Söderhäll I. Invertebrate hematopoiesis: an anterior
proliferation center as a link between the hematopoietic tissue and the brain. Stem Cells Dev.
2012;21(17):3173-86.

45. Leigh ND, Sessa S, Dragalzew AC, Payzin-Dogru D, Sousa JF, Aggouras AN, et al. von Willebrand
factor D and EGF domains is an evolutionarily conserved and required feature of blastemas capable of
multitissue appendage regeneration. Evol Dev. 2020;22(4):297-311.

46. Brown LJ, Alawoki M, Crawford ME, Reida T, Sears A, Torma T, et al. Lipocalin-7 is a matricellular
regulator of angiogenesis. PLoS One. 2010;5(11):e13905.

47. Mary S, Kulkarni MJ, Mehendale SS, Joshi SR, Giri AP. Tubulointerstitial nephritis antigen-like 1
protein is downregulated in the placenta of pre-eclamptic women. Clin Proteomics. 2017;14:8.

48. Shen M, Jiang YZ, Wei Y, Ell B, Sheng X, Esposito M, et al. Tinagl1 suppresses triple-negative breast
cancer progression and metastasis by simultaneously inhibiting integrin/FAK and EGFR signaling.
Cancer Cell. 2019;35(1):64-80.

49. Neufeld G, Cohen T, Gengrinovitch S, Poltorak Z. Vascular endothelial growth factor (VEGF) and its
receptors. The FASEB Journal. 1999;13(1):9-22.

50. Peichev M, Naiyer AJ, Pereira D, Zhu Z, Lane WJ, Williams M, et al. Expression of VEGFR-2 and
AC133 by circulating human CD34(+) cells identifies a population of functional endothelial precursors.
Blood. 2000;95(3):952-8.

51. Munier AI, Doucet D, Perrodou E, Zachary D, Meister M, Hoffmann JA, et al. PVF2, a PDGF/VEGF-
like growth factor, induces hemocyte proliferation in *Drosophila* larvae. EMBO Rep. 2002;3(12):1195-
200.

52. Varma Shrivastav S, Bhardwaj A, Pathak KA, Shrivastav A. Insulin-like growth factor binding protein-
3 (IGFBP-3): unraveling the role in mediating IGF-independent effects within the cell. Front Cell Dev
Biol. 2020;8:286.

53. Castillo J, Brown MR, Strand MR. Blood feeding and insulin-like peptide 3 stimulate proliferation of
hemocytes in the mosquito *Aedes aegypti*. PLoS Pathog. 2011;7(10):e1002274.

54. Wang G, Li N, Zhang L, Zhang L, Zhang Z, Wang Y. IGFBP7 promotes hemocyte proliferation in
small abalone *Haliotis diversicolor*, proved by dsRNA and cap mRNA exposure. Gene.
2015;571(1):65-70.

55. Charoensapsri W, Sangsuriya P, Lertwimol T, Gangnonngiw W, Phiwsaiya K, Senapin S. Laminin
receptor protein is implicated in hemocyte homeostasis for the whiteleg shrimp *Penaeus (Litopenaeus)*
vannamei. Dev Comp Immunol. 2015;51(1):39-47.

56. Rosa RD, Barracco MA. Antimicrobial peptides in crustaceans. Isj-Invert Surviv J. 2010;7(2):262-84.

57. Bachère E, Gueguen Y, Gonzalez M, de Lorgeril J, Garnier J, Romestand B. Insights into the anti-
microbial defense of marine invertebrates: the penaeid shrimps and the oyster *Crassostrea gigas*.
Immunol Rev. 2004;198:149-68.

58. Chen KY, Hsu TC, Huang PY, Kang ST, Lo CF, Huang WP, et al. *Penaeus monodon* chitin-binding
protein (PmCBP) is involved in white spot syndrome virus (WSSV) infection. Fish Shellfish Immunol.
2009;27(3):460-5.

59. Elbahnaswy S, Koiwai K, Zaki VH, Shaheen AA, Kondo H, Hirono I. A novel viral responsive protein
(MjVRP) from *Marsupenaeus japonicus* haemocytes is involved in white spot syndrome virus
infection. Fish Shellfish Immunol. 2017;70:638-47.

60. Munoz M, Vandenbulcke F, Saulnier D, Bachère E. Expression and distribution of penaeidin
antimicrobial peptides are regulated by haemocyte reactions in microbial challenged shrimp. Eur J
Biochem. 2002;269(11):2678-89.

61. Zhang K, Koiwai K, Kondo H, Hirono I. White spot syndrome virus (WSSV) suppresses penaeidin
expression in *Marsupenaeus japonicus* hemocytes. Fish & Shellfish Immunol. 2018;78:233-7.

62. Li S, Wang Z, Li F, Yu K, Xiang J. A novel vascular endothelial growth factor receptor participates in
white spot syndrome virus infection in *Litopenaeus vannamei*. Front Immunol. 2017;8:1457.

63. Wang Z, Li S, Li F, Xie S, Xiang J. Identification and function analysis of a novel vascular endothelial
growth factor, LvVEGF3, in the Pacific whiteleg shrimp *Litopenaeus vannamei*. Dev Comp Immunol.
2016;63:111-20.

64. Wang Z, Li S, Yu Y, Yu K, Zhang X, Xiang J, et al. Identification and characterization of two novel
vascular endothelial growth factor genes in *Litopenaeus vannamei*. Fish Shellfish Immunol.
2019;84:259-68.

65. Rosa RD, Vergnes A, de Lorgeril J, Goncalves P, Perazzolo LM, Sauné L, et al. Functional divergence
in shrimp anti-lipopolysaccharide factors (ALFs): from recognition of cell wall components to
antimicrobial activity. PLoS One. 2013;8(7):e67937.

66. Söderhäll I. Recent advances in crayfish hematopoietic stem cell culture: a model for studies of
hemocyte differentiation and immunity. Cytotechnology. 2013;65(5):691-5.

67. Mai HN, Nguyen HT, Koiwai K, Kondo H, Hirono I. Characterization of a Kunitz-type protease
inhibitor (MjKuPI) reveals the involvement of MjKuPI positive hemocytes in the immune responses of
kuruma shrimp *Marsupenaeus japonicus*. Dev Comp Immunol. 2016;63:121-7.

68. Koiwai K, Alenton RRR, Shiomi R, Nozaki R, Kondo H, Hirono I. Two hemocyte sub-populations of
kuruma shrimp *Marsupenaeus japonicus*. Mol Immunol. 2017;85:1-8.

69. Koiwai K, Kondo H, Hirono I. Isolation and molecular characterization of hemocyte sub-populations in
kuruma shrimp *Marsupenaeus japonicus*. Fisheries Science. 2019;85(3):521-32.

70. Sequeira T, Tavares D, Arala-Chaves M. Evidence for circulating hemocyte proliferation in the shrimp
Penaeus japonicus. Dev Comp Immunol. 1996;20(2):97-104.

71. Rota-Stabelli O, Daley AC, Pisani D. Molecular timetrees reveal a Cambrian colonization of land and a
new scenario for ecdysozoan evolution. Curr Biol. 2013;23(5):392-8.

72. Thomas GWC, Dohmen E, Hughes DST, Murali SC, Poelchau M, Glastad K, et al. Gene content
evolution in the arthropods. Genome Biol. 2020;21(1):15.

73. Wolfe JM, Breinholt JW, Crandall KA, Lemmon AR, Lemmon EM, Timm LE, et al. A phylogenomic
framework, evolutionary timeline and genomic resources for comparative studies of decapod
crustaceans. Proc Biol Sci. 2019;286(1901):20190079.

74. Broseus L, Thomas A, Oldfield AJ, Severac D, Dubois E, Ritchie W. TALC: Transcript-level Aware
Long-read Correction. Bioinformatics. 2020.

75. Bushmanova E, Antipov D, Lapidus A, Prjibelski AD. rnaSPAdes: a *de novo* transcriptome assembler
and its application to RNA-Seq data. Gigascience. 2019;8(9):1-13.

76. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length transcriptome
assembly from RNA-Seq data without a reference genome. Nat Biotechnol. 2011;29(7):644-52.

77. Biočanin M, Bues J, Dainese R, Amstad E, Deplancke B. Simplified Drop-seq workflow with
minimized bead loss using a bead capture and processing microfluidic chip. Lab Chip.
2019;19(9):1610-20.

78. Picelli S, Bjorklund AK, Reinius B, Sagasser S, Winberg G, Sandberg R. Tn5 transposase and
tagmentation procedures for massively scaled sequencing projects. Genome Research.
2014;24(12):2033-40.

79. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9(4):357-9.

80. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic data across
different conditions, technologies, and species. Nat Biotechnol. 2018;36(5):411-20.

81. Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data using
regularized negative binomial regression. Genome Biol. 2019;20(1):296.

82. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture
and applications. BMC Bioinformatics. 2009;10:421.

83. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. Journal of
Molecular Biology. 1990;215(3):403-10.

84. Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, et al. The single-cell transcriptional
landscape of mammalian organogenesis. Nature. 2019;566(7745):496-502.