1   **Theoretical and Applied Genetics**

3   **SPECIAL ISSUE: Breeding towards Agricultural Sustainability - invited**
4   **ORIGINAL ARTICLE**

7   **Genetic diversity, distribution and domestication history of the neglected**

8   **GGA^{t}A^t genepool of wheat[#]**

Ekaterina D. Badaeva[1,2,*], Fedor A. Konovalov[3,4], Helmut Knüpffer[4], Agostino Fricano[5],

Alevtina S. Ruban[4,6], Zakaria Kehel[7], Svyatoslav A. Zoshchuk[2], Sergei A. Surzhikov[2], Kerstin

Neumann[4], Andreas Graner[4], Karl Hammer[4], Anna Filatenko[4,8], Amy Bogaard[9], Glynis Jones[10],

Hakan Özkan[11], Benjamin Kilian[4,12]

1 – N.I. Vavilov Institute of General Genetics, Russian Academy of Sciences, Moscow, **Russia**
2 – Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow, **Russia**
3 – Independent Clinical Bioinformatics Laboratory, Moscow, **Russia**
4 – Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Gatersleben, **Germany**
5 – Council for Agricultural Research and Economics – Research Centre for Genomics &
Bioinformatics, Fiorenzuola d'Arda (PC), **Italy**
6 – KWS SAAT SE & Co. KGaA, Einbeck, **Germany**
7 – International Center for the Agricultural Research in the Dry Areas (ICARDA), Rabat,
Morocco
8 – Independent Researcher, St. Petersburg, **Russia**
9 – School of Archaeology, Oxford, **United Kingdom**
10 – Department of Archaeology, University of Sheffield, **United Kingdom**
11 – Department of Field Crops, Faculty of Agriculture, University of Çukurova, Adana, **Turkey**
12 – Global Crop Diversity Trust, Bonn, **Germany**

*Correspondence: Ekaterina D. Badaeva; ORCID: https://orcid.org/0000-0001-7101-9639;
e-mail: katerinabadaeva@gmail.com

Key message **We present a comprehensive survey of cytogenetic and genomic diversity of**
the GGA^{t}A^t genepool of wheat, thereby unlocking these plant genetic resources for wheat
improvement.

Abstract
Wheat yields are stagnating around the world and new sources of genes for resistance or
tolerances to abiotic traits are required. In this context, the tetraploid wheat wild relatives are
among the key candidates for wheat improvement. Despite of its potential huge value for wheat
breeding, the tetraploid GGA^{t}A^t genepool is largely neglected. Understanding the population
structure, native distribution range, intraspecific variation of the entire tetraploid GGA^{t}A^t
genepool and its domestication history would further its use for wheat improvement. We report
the first comprehensive survey of genomic and cytogenetic diversity sampling the full breadth
and depth of the tetraploid GGA^{t}A^t genepool. We show that the extant GGA^{t}A^t genepool
consists of three distinct lineages. We provide detailed insights into the cytogenetic composition
of GGA^{t}A^t wheats, revealed group-, and population-specific markers and show that
chromosomal rearrangements play an important role in intraspecific diversity of *T. araraticum.*
We discuss the origin and domestication history of the GGA^{t}A^t lineages in the context of state-
of-the-art archaeobotanical finds. We shed new light on the complex evolutionary history of the
GGA^{t}A^t wheat genepool. We provide the basis for an increased use of the GGA^{t}A^t wheat
genepool for wheat improvement. The findings have implications for our understanding of the
origins of agriculture in southwest Asia.

Keywords: *Triticum araraticum*, *Triticum timopheevii*, Crop Wild Relatives, Chromosomal
passports, Fertile Crescent, New glume wheat

Introduction
The domestication of plants since the Neolithic Age resulted in the crops that feed the world
today. However, successive rounds of selection during the history of domestication led to a
reduction in genetic diversity, which now limits the ability of the crops to further evolve
(Tanksley and McCouch 1997; van Heerwaarden et al. 2010). This is exacerbated by the demand
for high crop productivity under climate change. Crop wild relatives (CWR) represent a large
pool of beneficial allelic variation and are urgently required to improve the elite genepools
(Dempewolf et al. 2017; Kilian et al. 2021). Bread wheat and durum wheat are the staple crops

for about 40% of the world's population. But as wheat yields are stagnating around the world

(Iizumi et al. 2017; Ray et al. 2013; Ray et al. 2012); new sources of genes for resistance or

tolerances to abiotic traits such as drought and heat are required. In this context, the wheat wild

relatives are among the key sources for bread wheat (*T. aestivum* L., 2n = 6x = 42, BBAADD)

and durum wheat (*T. durum* Desf., 2n = 4x = 42, BBAA) improvement (Dante et al. 2013; Zhang

et al. 2016).

However, in nature, no wild hexaploid wheat has ever been found. Only two wild tetraploid

wheat species (2n = 4x = 28) were discovered, namely, (i) wild emmer wheat *T. dicoccoides*

(Körn. ex Asch. et Graebn.) Körn. ex Schweinf. [syn. *T. turgidum* subsp. *dicoccoides* (Körn. ex

Asch. & Graebn.) Thell.] and (ii) Armenian, or Araratian emmer *T. araraticum* Jakubz. (syn. *T.*

timopheevii (Zhuk.) Zhuk. subsp. *armeniacum* (Jakubz.) van Slageren). Morphologically, both

species are very similar but differ in their genome constitution (Zohary et al. 2012). *Triticum*

dicoccoides has the genome formula BBAA and *T. araraticum* has GGA^tA^t (Jiang and Gill

1994).

The wheat section Timopheevii mainly consists of wild tetraploid *Triticum araraticum*

(GGA^tA^t), domesticated tetraploid *T. timopheevii* (Zhuk.) Zhuk. (Timopheev's wheat, GGA^tA^t),

and hexaploid *T. zhukovskyi* Menabde et Ericzjan (2n = 6x= 42, $GGA^tA^tA^mA^m$) (Dorofeev et al.

1979; Goncharov 2012).

Wild *T. araraticum* was first collected by M.G. Tumanyan and A.G. Araratyan during 1925–28

southeast of Erevan, Armenia (Nazarova 2007), soon after the discovery of domesticated *T.*

timopheevii by P.M. Zhukovsky (Zhukovsky 1928) (Supplementary Material S1). Subsequently,

T. araraticum was found in several other locations in Armenia and Azerbaijan (Dorofeev et al.

1979; Jakubziner 1933, 1959), as well as in Iran, Iraq and Turkey. Single herbarium specimens

resembling *T. araraticum* have been sporadically recorded among *T. dicoccoides* accessions

collected from the Fertile Crescent (Jakubziner 1932; Sachs 1953). However, only botanical

expeditions from the University of California at Riverside (USA) to Turkey in 1965, to the

Fertile Crescent in 1972–73 (Johnson and Hall 1967; Johnson and Waines 1977) and the

Botanical Expedition of Kyoto University to the Northern Highlands of Mesopotamia in 1970

(Tanaka and Ishii 1973; Tanaka and Kawahara 1976) significantly expanded our understanding

of the natural distribution of *T. araraticum*. More recently *T. araraticum* was found in north-

western Syria (Valkoun et al. 1998). Especially in south-eastern Anatolia, Turkey, the

distribution area of *T. araraticum* overlaps with the distribution range of *T. dicoccoides*. From

the western to eastern Fertile Crescent, it is assumed that *T. araraticum* gradually substitutes

T. dicoccoides (Johnson 1975), and *T. dicoccoides* is absent from Transcaucasia (Özkan et al.

2011). In most habitats, *T. araraticum* grows in patches and in mixed stands with other wild

cereals (Troitzky 1932; Tumanyan 1930).

It is difficult to distinguish *T. araraticum* from *T. dicoccoides* by morphology under field

conditions (Dagan and Zohary 1970; Tanaka and Sakamoto 1979). However, both species can

easily be differentiated based on biochemical, immunological, cytological and molecular

markers (Badaeva et al. 1994; Gill and Chen 1987; Jiang and Gill 1994; Kawahara and Tanaka

1977; Konarev et al. 1976; Lilienfeld and Kihara 1934). From an archaeobotanical perspective,

both species can be reliably identified based on several characteristics of charred spikelets (Jones

et al. 2000). In blind tests, it was possible to distinguish modern representatives of the two

species with a c. 90% accuracy, on the basis of the primary keel of the glume, which arises just

below the rachis disarticulation scar, and the prominent vein on the secondary keel (observable at

the base of the glume, which is the part of spikelet most commonly preserved by charring in

archaeological material) (Jones et al. 2000).

According to cytogenetic and molecular analyses, *T. araraticum* originated as a result of

hybridization between *Aegilops speltoides* Tausch (2n = 2x= 14, SS) and *Triticum urartu*

Thumanjan ex Gandilyan (2n = 2x= 14, AA) independently from *T. dicoccoides* (Dvořák et al.

1988; Rodríguez et al. 2000a,b). Similarity of the cytoplasmic genomes of *Ae. speltoides* and *T.*

araraticum indicated that *Ae. speltoides* was the maternal parent of *T. araraticum* (Tsunewaki

1996). Hybrids between *T. araraticum* × *T. dicoccoides* were reported as sterile because of

meiotic disturbances and gene interactions (Makushina 1938; Svetozarova 1939; Tanaka and

Ishii 1973; Wagenaar 1961), although a few authors (Noda and Ge 1989; Sachs 1953; Tanaka

and Ichikawa 1972; Tanaka and Kawahara 1976) reported relatively good chromosome pairing

in the F_1 hybrids in some *T. araraticum* × *T. dicoccoides* combinations.

Triticum dicoccoides is considered to be the older species than *T. araraticum* (Gornicki et al.

2014; Huang et al. 2002), which is supported by higher similarity of the S-G genomes compared

to the S-B genomes (Jiang and Gill 1993; Kilian et al. 2007; Rodríguez et al. 2000a).

Cytogenetic and molecular data showed that the speciation of *T. araraticum* was accompanied

by complex species-specific translocations involving chromosomes 1G-6A[t]-4G and 3A[t]-4A[t]

(Chen and Gill 1984; Jiang and Gill 1993; Rodríguez et al. 2000b; Salina et al. 2006) as well as

with mutations of the primary DNA structure causing the divergence of homoeologous

chromosomes (e.g., chromosomes 3A-3A[t]) (Dobrovolskaya et al. 2009). These changes of

karyotype structure are specific for the whole section Timopheevii as compared to the emmer

wheat lineage (BBAA, BBAADD) (Badaeva et al. 1986; Hutchinson et al. 1982; Zhang et al.

2013).

Intraspecific diversity of *T. araraticum* was detected by karyotype analysis (Badaeva et al. 1990,

1994; Kawahara et al. 1996; Kawahara and Tanaka 1977) and using nuclear (Nave et al. 2021;

Shcherban et al. 2016) and chloroplast DNA markers (Mori et al. 2009).

Most recent phylogenetic studies based on whole chloroplast genome sequences, genome-wide

sequence information and enlarged taxon sampling provided increased resolution of the

evolutionary history within the Triticeae tribe, thereby shedding also new light on the GGA^{t}A^t

wheat genepool (Bernhardt et al. 2017; Gornicki et al. 2014).

Research and pre-breeding activities have focused on *T. dicoccoides* because it gave rise to the

economically most important wheats, *T. durum* Desf. and *T. aestivum* L. (Avni et al. 2017; El

Haddad et al. 2021). However, *T. araraticum* and *T. timopheevii* have also contributed to bread

wheat improvement. Important genes were transferred such as genes controlling resistance

against stem rust, leaf rust, powdery mildew or wheat leaf blotch (Allard and Shands 1954;

Brown-Guedira et al. 1996, 2003; Dyck 1992; McIntosh and Gyarfas 1971). Cytoplasmic male

sterility (CMS) induced by *T. timopheevii* cytoplasm showed great potential for heterotic hybrid

technology (Maan and Lucken 1972; Mikó et al. 2011; Würschum et al. 2017). However, despite

of its potential huge value for bread and durum wheat improvement, only a comparatively small

number of genes was transferred from *T. timopheevii* (even less from *T. araraticum*). Most of the

gene contributions originate from only one line (D-357-1) bred by R. Allard at the University of

Wisconsin in 1948 (Martynov et al. 2018).

Understanding the population structure and the intraspecific variation of the entire tetraploid

GGA^{t}A^t genepool would further its use for wheat improvement. In this study, we report the first

comprehensive survey of cytogenetic and genomic diversity sampling the full breadth and depth

of the tetraploid GGA^{t}A^t genepool. We provide new insights into the genetic relationships

among GGA^{t}A^t wheats and its domestication, taking into account the state-of-the-art

archaeobotanical finds.

Materials and methods

Germplasm

A comprehensive germplasm collection of tetraploid wheats was established comprising 862

genebank accessions (1–5 genotypes per accession number): 450 *T. araraticum,* 88 *T.*

timopheevii, three *T. militinae* Zhuk. et Migusch.*,* one *T. zhukovskyi,* 307 *T. dicoccoides,* eight *T.*

dicoccon, and five *T. durum*. Additionally, the following materials were included: (i) four

samples of *T. araraticum* collected by Dr. Nelli Hovhannisyan in Armenia; (ii) 17 samples of *T.*

araraticum and 12 samples of *T. dicoccoides* collected by Dr. H. Özkan in Turkey; and (iii) 26

samples of *T. dicoccoides* collected by Drs. E. Badaeva, O.M. Raskina and A. Belyayev in Israel.

Altogether, 921 accessions of wild and domesticated tetraploid wheats were examined

(Supplementary Table S2).

A subset of 787 tetraploid wheat genotypes representing 765 genebank accessions was examined

using Sequence-Specific Amplification Polymorphism (SSAP) markers. The subset included 360

genotypes of *T. araraticum*, 76 *T. timopheevii* (including two *T. militinae*), while 351 genotypes

of *T. dicoccoides* were considered as an outgroup. Of them, 243 (67%) *T. araraticum,* 139

(39.6%), and nine *T. timopheevii* genotypes (including one *T. zhukovsky*) were shared with C-

banding analysis. The whole collection was single-seed descended (SSD) at least twice under

field conditions (2009–2012) and taxonomically re-identified in the field at IPK, Gatersleben in

2011 (Supplementary Table S2).

Based on the results of the SSAP analysis, a subset of 103 genotypes, including 37 *T. araraticum*

collected from different geographic regions and representing all genetic groups, one *T. militinae*,

14 *T. timopheevii*, 38 *T. dicoccoides,* 9 *T. dicoccon,* and four *T. durum* was selected for a

complementary analysis using Amplified Fragment Length Polymorphism (AFLP) markers to

infer the population structure of GGA^{t}A^t wheats (Supplementary Table S2). Forty-seven *T.*

araraticum and *T. timopheevii* accessions (88.7%) were shared with SSAP and 39 (73.6%) with

C-banding analyses.

Karyotype diversity of 370 *T. araraticum* accessions was assessed by C-banding and

Fluorescence *in situ* hybridization (FISH) in comparison with 17 *T. timopheevii* and one *T.*

zhukovskyi genotypes (Supplementary Table S2). According to the C-banding analysis, most *T.*

araraticum accessions (353 of 370) were karyotypically uniform and were treated as single

genotype each. Seventeen accessions were heterogeneous and consisted of two (13 accessions)

or even three (four accessions) cytogenetically distinct genotypes, which were treated as

different entities (genotypes). In order to infer the population structure of GGA^{t}A^t wheats, 265

typical genotypes of *T. araraticum* were selected representing all karyotypic variants, seven *T.*

timopheevii, one *T. zhukovskyi.* A total of 87 *T. araraticum* genotypes representing all

geographic regions and chromosomal groups were selected for Fluorescence i*n situ* hybridization

(FISH) analysis.

Molecular analysis

SSAP analysis

Evolutionary relationships among the comprehensive collection of wild and domesticated

tetraploid wheat taxa were first inferred based on polymorphic retrotransposon insertions. For

this, the highly multiplex genome fingerprinting method SSAP was implemented based on

polymorphic insertions of retrotransposon families *BARE-1* and *Jeli* spread across the wheat

chromosomes. DNA was isolated from freeze-dried leaves of 787 SSD plants, using the Qiagen

DNeasy Kit (Hilden, Germany). The SSAP protocol was based on Konovalov et al. (2010) with

further optimizations for capillary-based fragment detection (Supplementary Material S3). In

total, 656 polymorphic markers were generated for BBAA- and GGA^{t}A^t-genome wheats by

amplification of multiple retrotransposon insertion sites. Data analysis was performed in

SplitsTree 4.15.1 (Huson and Bryant 2006). NeighborNet planar graphs of Dice distances (Dice

1945) were constructed based on presence/absence of SSAP bands in the samples.

AFLP analysis

The AFLP protocol, as described by Zabeau and Vos (1993), was performed with minor

modifications according to Altıntas□ et al. (2008) and Alsaleh et al. (2015). In total, six AFLP

primer combinations were used to screen the collection of 103 lines (Supplementary Material

S3). Neighbor-Joining (NJ) trees were computed based on Jaccard distances (Jaccard 1908;

Perrier et al. 2003). NeighborNet planar graphs were generated based on Hamming distances

(Huson and Bryant 2006). Genetic diversity parameters and genetic distances were calculated

using Genalex 6.5 (Peakall and Smouse 2012).

Cytogenetic analysis

C-banding

Chromosomal preparation and C-banding procedure followed the protocol published by Badaeva

et al. (1994). The A^t- and G-genome chromosomes were classified according to the nomenclature

proposed by Badaeva et al. (1991) except for chromosomes 3A^t and 4A^t. Based on meiotic

analysis of the F$_1$ *T. timopheevii* × *T. turgidum* hybrids (Rodríguez et al. 2000b) and considering

karyotype structure of the synthetic wheat *T. × soveticum* (Zhebrak) (Mitrofanova et al. 2016),

the chromosomes 3A^t and 4A^t were exchanged. Population structure of *T. araraticum* and the

phylogenetic relationship with *T. timopheevii* were inferred based on chromosomal passports

compiled for 265 genotypes representing all geographic regions and chromosomal groups (247

T. araraticum, 17 *T. timopheevii*, one *T. zhukovskyi*). Chromosomal passports were constructed

by comparing the karyotype of the particular accession with the generalized idiogram of A^t- and

G-genome chromosomes (Fig. 1; Supplementary Material S3), as described for *T. dicoccon*

Schrank (Badaeva et al. 2015b).

Fluorescence *in situ* hybridization (FISH)

Two polymorphic G-genome specific DNA probes, Spelt-1 (Salina et al. 1998) and Spelt-52

(Salina et al. 2004), were used for screening all 95 (87 *T. araraticum,* seven *T. timopheevii* and

one *T. zhukovskyi*) genotypes considered for FISH analysis. Additionally, 26 of the 95 genotypes

were analyzed using the probe pAesp_SAT86 (Badaeva et al. 2015a), which also showed

differences between the accessions. The probes pSc119.2 (Bedbrook et al. 1980), GAA_n

(Pedersen et al. 1996), and pTa-535 (Komuro et al. 2013) were subsequently hybridized to the

same chromosomal spread to allow chromosome identification. Classification of pSc119.2- and

pTa-535-labeled chromosomes followed the nomenclature of (Badaeva et al. 2016; Jiang and

Gill 1993). Chromosomes hybridized with the GAA_{10} microsatellite sequence were classified

according to nomenclature suggested for C-banded chromosomes.

Results

Genetic diversity and population structure of the GGA^tA^t genepool

Simultaneous amplification of multiple retrotransposon insertion sites using eight Long Terminal

Repeat (LTR) primer combinations generated 656 polymorphic Sequence-Specific Amplification

Polymorphism (SSAP) markers: 255 markers were obtained for *Jeli* insertions and 401 markers

for *BARE-1* insertions. According to our previous study (Konovalov et al. 2010), *Jeli* targets

mostly the A genome, while *BARE-1* is distributed between A- and B/G-genome chromosomes.

Altogether, 787 wheat genotypes representing 753 genebank accessions were considered for data

analysis, after excluding apparently misidentified taxa and several cases of low-yield DNA

extraction (Supplementary Table S2, column 6). Several major observations were made: (i) the

extant GGA^tA^t genepool consists of three distinct lineages (two *T. araraticum* lineages and one

of *T. timopheevii*, TIM). Surprisingly, wild *T. araraticum* consists of two major genetic lineages,

preliminarily designated as 'ARA-0' and 'ARA-1' (Fig. 2; Supplementary Figure S4;

Supplementary Figure S5); (ii) while ARA-0 was found to be geographically widespread, ARA-

1 was only found in south-eastern Turkey (Adiyaman, Kahramanmaraş, Gaziantep, Kilis) and in

north-western Syria, where the distribution ranges of *T. araraticum* and of *T. dicoccoides*

overlap (Fig. 3); (iii) as expected, *T. dicoccoides* is genetically more diverse, supporting a more

recent origin of *T. araraticum* (Supplementary Figure S4; Supplementary Table S6); (iv) among

the GGA^tA^t lineages, ARA-0 harbors more genetic diversity (Supplementary Table S6); (v)

differences between A^t- and A^t+G genome diversity patterns based on *Jeli* and *BARE-1* were

discovered, respectively (Supplementary Table S6); and (vi) Nei's genetic distance between

lineages based on all 656 SSAP markers or considering only the *BARE-1* markers revealed that

ARA-0 is phylogenetically more closely related to TIM than ARA-1. However, considering only

the *Jeli* markers, ARA-1 was more closely related to TIM. *Triticum dicoccoides* (DIC) was

genetically related most closely to the ARA-1 lineage (Supplementary Table S6).

A carefully selected subset of 103 genotypeswas used for Amplified Fragment Length
Polymorphism (AFLP) analysis (Supplementary Table S2, column 8). A screen using the six
most promising AFLP primer combinations uncovered a total of 146 polymorphic markers
across all genotypes. Major findings can be summarized as: (i) clustering of genotypes based on
AFLP markers corroborates the existence of the three major and genetically distinct lineages of
GGA^{t}A^t wheats (Supplementary Figure S7). One lineage comprised all *T. timopheevii* (and its
derived mutant *T. militinae* Zhuk. et Migusch.) genotypes. Importantly, the two lineages of *T.*
araraticum and their distribution ranges were verified; (ii) the summary statistics highlight that,
in contrast to the SSAP analysis, ARA-1 was more diverse than TIM for all parameters
(Supplementary Table S8); (iii) ARA-1 was found to be genetically most closely related to TIM
(Supplementary Table S8); (iv) ARA-1 was related most closely to DIC Supplementary Table
S8); (v) potential hybridization signals between ARA-0 and ARA-1 lineages were identified
(Supplementary Figure S7, purple split); (vi) ARA-1 lines collected around Kilis,
Kahramanmaraş and Gaziantep in Turkey were genetically closest to TIM (Supplementary
Figure S7, purple split); and (vii) shared splits between ARA-0 genotypes collected in Armenia
and Azerbaijan with TIM (Supplementary Figure S7, purple split) indicate the potential
contribution of ARA-0 to the formation of TIM or probably hybridization of ARA-0 with TIM.
The genetically closest ARA-0 genotype to TIM was collected from the present Ararat province
of Armenia.

In total, 248 genotypes of *T. araraticum* and 17 of *T. timopheevii* collected across the entire
distribution range and representing all karyotypic variants (Supplementary Table S2, column 12)
were selected to infer the population structure based on chromosomal passports (Badaeva et al.
2015b). The results based on 96 informative C-bands supported the molecular findings using *Jeli*
markers and AFLP markers and were congruent with the AFLP marker results as the closest
ARA-1 genotypes to TIM were collected near Gaziantep (61 km SE from Türkoğlu, SW of
Karadağ), Turkey (Fig. 3; Supplementary Figure S9; Supplementary Table S10).
Fluorescence *in situ* hybridization (FISH) using six DNA probes was carried out for 95
genotypes (Supplementary Table S2, column 13), which, in turn, corroborates the findings
obtained using *Jeli* markers, AFLP and C-banding markers (Supplementary Table S11).
296
297 ***Intraspecific genetic diversity of T. araraticum based on karyotype structure and C-banding***
298 ***patterns***
299 Cytogenetic analysis of 391 *T. araraticum* genotypes in comparison with 17 *T. timopheevii*
300 genotypes provided detailed insights into the genetic composition of GGA^{t}A^t wheats. First,

cytogenetic analysis highlighted significant differences between wild emmer *T. dicoccoides* and *T. araraticum/T. timopheevii* in karyotype structure and C-banding patterns (Fig. 4). Second, we revealed high diversity of the C-banding patterns and broad translocation polymorphisms within *T. araraticum* (Figs. 4, 5). The karyotype lacking chromosomal rearrangements was defind as 'normal' (N) and was the most frequent karyotype variant shared by *T. timopheevii* and *T. araraticum*. The 'normal' karyotype was found in 175 of 391 *T. araraticum* genotypes (44.6%) (more specifically: 155 of 342 ARA-0 = 45.32%; 20 of 49 ARA-1 = 40.81%). These 175 genotypes differed from each other only in the presence/absence or size of one to several C-bands. The ratio of karyotypically normal genotypes decreased from 86.7% in Azerbaijan (excluding Nakhichevan), to 60.0% in Turkey, 46.5% in Iraq, 30.6% in Armenia, 20.0% in Nakhichevan, Azerbaijan, 16.7% in Syria, to 9.1% in Iran (Supplementary Figure S12). Local populations differed in the ratio of normal/rearranged genotypes. For example, some populations from Dahuk, Iraq, possessed only karyotypically normal genotypes, while in others all genotypes possessed chromosomal rearrangements. Similarly, in Turkey the frequency of karyotypically normal genotypes varied from 100% (Mardin) to 0% (Kilis) (Supplementary Table S13).

C-banding patterns of *T. araraticum* were highly polymorphic. Based on the presence or absence of particular C-bands (Figs. 1, 4), all genotypes were divided into two groups. The first, larger group (ARA-0) comprised of 342 genotypes (Supplementary Figures S14 - S19). The second group (ARA-1) included 49 genotypes (Supplementary Figure S18 *h1–h5; h6–h9*; Supplementary Figure S20). Interestingly, genebank accession TA1900 presumably collected 32 km S of Denizli, Dulkadiroğlu district, Kahramanmaraş province in the Taurus Mountain Range of Turkey in 1959 shared karyotypic features of *T. timopheevii* and ARA-0 (Fig. 4) and, probably, it is a natural or artificially produced hybrid of *T. timopheevii* with an unknown genotype of *T. araraticum.*

Six chromosomes carried diagnostic C-bands for the ARA-1 lineage at the following positions: 1A^{t}L3, 4A^{t}S7, 5A^{t}L3, 1GL5, 2GL7, 3GL7+L11 (Fig. 1). All these diagnostic C-bands were present in all ARA-1 genotypes, both with normal and rearranged karyotypes (Supplementary Figure S18, Supplementary Fig. S20). Some C-bands were also common for both ARA-0 and ARA-1 (e.g., 2A^{t}L3 and 2GS15, Fig. 4), but their size was larger in ARA-1 genotypes.

Only few C-bands were characteristic for the ARA-0 lineage (Figs. 1, 4). Three C-bands appeared with a frequency of over 95%, and two of them, 6A^{t}L3 and 5GL15, occurred only in the ARA-0 group. The third C-band, 2GL13, was detected in 97% of ARA-0, but also in few ARA-1 genotypes.

'Region-specific' C-bands (in terms of highest frequency) (Badaeva et al. 1994) were detected for ARA-0. For example, (i) the band 4A^{t}L7 dominated in Turkey and Transcaucasia

(Supplementary Figure S14, Supplementary Figure S19); (ii) a medium to large $3A^tS7$ band

was frequently observed in Iran and Sulaymaniyah (Iraq); (ii) one distinct 5GL13 band was

frequently detected in genotypes from Erbil, Iraq (Supplementary Figure S15, Supplementary

Figure S16). The specificity of C-banding patterns in genotypes originating from the same

geographic region was not only determined by single bands, but usually by a particular

combination of C-bands on several chromosomes. These region-specific banding patterns were

observed for both, normal and translocated forms. For example, the unique banding pattern of

chromosome 7G, lacking the marker C-band 7GS11, but carrying large bands for 7GS13, 7GS17

and 7GS21 on the short arm, and 7GL13 and 7GL15 on the long arm was common in

Transcaucasia (Supplementary Figure S14). The chromosome 6G lacking telomeric C-bands on

the long arm was frequent in genotypes from Dahuk and Sulaymaniyah (both Iraq), and also

occurred in ARA-1 genotypes from Kahramanmaraş, Turkey (Supplementary Figure S20).

Chromosomal rearrangements play an important role in intraspecific diversity of T. araraticum

In total, 216 out of 391 (55.4%) *T. araraticum* accessions carried a translocated karyotype.

Seventy-six variants of chromosomal rearrangements including single and multiple

translocations, paracentric and pericentric inversions were identified (Figs. 5, 6). Novel

rearrangements were represented by 44 variants, while 32 variants were described earlier

(Badaeva et al. 1990, 1994; Kawahara et al. 1996; Kawahara and Tanaka 1977, 1981). One-

hundred-forty-seven genotypes differed from the 'normal' karyotype by one, 45 genotypes by

two (double translocations), 21 genotypes by three (triple translocations) and three genotypes –

by four chromosomal rearrangements (quadruple translocations) (Supplementary Table S13).

Altogether, we revealed 52 (33 novel) variants of single chromosomal rearrangements (Figs. 5,

6). They included paracentric (one variant) and pericentric inversions (seven variants) and 44

single translocations involving A^t-A^t, A^t-G, or G-G-genome chromosomes. Double

rearrangements were represented by 16 independent and three cyclic translocations; among them

10 were novel. Triple translocations were represented by three variants, two of which – $T2A^t$:7G

+ $T6A^t$:5G:6G and T3G:7G:$7A^t$ + $T6A^t$:6G – were found here for the first time. Both variants of

quadruple translocations have been identified earlier in Transcaucasia (Badaeva et al. 1990,

1994).

A translocation between two chromosomes could give rise to different products depending on the

breakpoint position and arm combination in rearranged chromosomes. For example, a

centromeric translocation between 1G and 2G resulted in two translocation variants which were

distinct in arm combinations (S:S *vs.* S:L). Three translocation variants involving chromosomes

3G and 4G differed from each other in arm combination and breakpoint position. To discriminate

different translocation variants involving same chromosomes, they were designated as T1G:2G-1

and T1G:2G-2, etc. (Supplementary Table S13).

Most variants of chromosomal rearrangements were unique and identified in one or few

genotypes, and only four variants were relatively frequent. These were a triple translocation

$T2A^t$:4G:7G + $T4A^t$:$7A^t$, *per*Inv$7A^t$-1, T6G:7G, and T2G:4G:6G (21, 16, 12, and 13 genotypes,

respectively). Taken together, these four variants accounted for approximately 16% of the whole

materials we studied. Genotypes carrying the same rearrangement usually had similar C-banding

patterns and were collected from the same, or closely located geographic regions. For example,

(i) all genotypes with $T2A^t$:4G:7G + $T4A^t$:$7A^t$ originated from Nakhichevan, Azerbaijan; (ii)

T6G:7G and T2G:4G:6G were found in Erbil, Iraq; and (iii) *per*Inv$7A^t$-1 was collected in Iran

and in the neighboring region of Sulaymaniyah, Iraq.

Other frequent translocations had a more restricted distribution and usually occurred in a single

population. Only few genotypes carrying the same translocations were identified in spatially

separated populations. These genotypes differed in their C-banding patterns, for example: (i)

T2G:4G was found in four genotypes from Erbil, Iraq and in two genotypes from Siirt, Turkey;

(ii) T1G:3G was identified in five ARA-0 genotypes from Dahuk, Iraq, one ARA-1 from Turkey

and one ARA-1 genotype of a mixed accession IG 117895 collected in Syria; and (iii) T1G:5G

was identified not only in three cytogenetically distinct *T. araraticum* ARA-0 genotypes from

Armenia and Azerbaijan and ARA-1 from Turkey (Fig. 6; Supplementary Figure S14,

Supplementary Figure S20), but also in *T. timopheevii* (Badaeva et al. 2016).

Most populations consisted of both genotypes with 'normal' karyotype and genotypes with one

to several variants of chromosomal rearrangements. However, their ratio and spectra differed

between regions (Fig. 6; Supplementary Figure S12; Supplementary Material S21).

Cytogenetic diversity of GGA^{t}A^t wheats assessed using FISH markers

To further investigate the intraspecific diversity of *T. araraticum* and to assess their phylogenetic

relationships with *T. timopheevii*, we carried out FISH using six DNA probes. The probes pTa-

535, pSc119.2 and GAA_n ensured chromosome identification (Badaeva et al. 2016) whereas

Spelt-1, Spelt-52 and pAesp_SAT86 were used to estimate intra- and interspecific variation (Fig.

7).

The distribution of pTa-535 was monomorphic among *T. araraticum* and *T. timopheevii,* while

the pSc119.2 site on $1A^tL$ discriminated ARA-1 and TIM from ARA-0 (Fig. 7*a*).

The variability of pAesp_SAT86 hybridization patterns was analyzed in four *T. timopheevii* and 26 *T. araraticum* genotypes, of them seven were from ARA-1 and 19 from ARA-0 lineages (Supplementary Table S2). Distribution of the pAesp_SAT86 probe in all *T. timopheevii* genotypes was similar except for chromosome $7A^t$ in genotype K-38555, which was modified due to a paracentric inversion or insertion of an unknown chromosomal fragment. Labeling patterns, however, were highly polymorphic for *T. araraticum* (Fig. 7; Supplementary Figure S22). Large pAesp_SAT86 sites were found only on some G-genome chromosomes. The A^t-genome chromosomes possessed several small, but genetically informative polymorphic sites. Some of these sites were lineage-specific. Most obvious differences were observed for $3A^t$, 4G and 7G chromosomes (Fig. 1). Thus, all TIM and ARA-1 genotypes carried the pAesp_SAT86 signal in the middle of $3A^tS$, while for ARA-0 it was located sub-terminally on the long arm. One large pAesp_SAT86 cluster was present on the long arm of 4G in ARA-0, but on the short arm in ARA-1 and TIM. Two large and adjacent pAesp_SAT86 clusters were detected on 7GS in ARA-1 and TIM, but they were split between opposite chromosome arms in all ARA-0 genotypes. Differences between ARA-0 and ARA-1 in pAesp_SAT86 cluster position on 4G and 7G could be caused by pericentric inversions. Some other pAesp_SAT86 sites were identified in either one of the three groups. ARA-1 exhibited the largest polymorphism of pAesp_SAT86 labeling patterns among all groups (Supplementary Figure S22).

FISH with Spelt-1 and Spelt-52 probes on chromosomes of 87 *T. araraticum* genotypes from different chromosomal groups and of different geographic origin, seven *T. timopheevii*, and one *T. zhukovskyi* genotypes revealed high intraspecific diversity of *T. araraticum* and low polymorphism in *T. timopheevii* (Supplementary Figure S23). The broadest spectra of labeling patterns were found in genotypes from Dahuk and Sulaymaniyah (Iraq) and in the ARA-1 group from Turkey, while material from Transcaucasia exhibited the lowest variation. The polymorphism was due to variation in the number of Spelt-1 and Spelt-52 sites, their size and chromosomal location (Supplementary Table S24). The Spelt-1 signals in various combinations appeared in sub-telomeric regions of either one or both arms of $2A^t$, $6A^t$, and all G-genome chromosomes. The Spelt-52 signals were observed in various combinations on $2A^tS$, 1GS, and 6GL chromosomes. Among them, two Spelt-1 sites (6GL and 7GL) and one Spelt-52 locus (2GS) were novel. The patterns of Spelt-1 and Spelt-52 repeats varied across geographic regions and between chromosomal groups (Supplementary Material S25; Supplementary Table S26).

Discussion

We present the most comprehensive survey of cytogenetic and genomic diversity of GGA^tA^t wheats. We describe the composition, distribution and characteristics of the GGA^tA^t genepool. Building on our results, the latest published complementary genomics studies and state-of-the-art archaeobotanical evidence we revisit the domestication history of the GGA^tA^t wheats. We arrived at the following four key findings:

1. The GGA^tA^t genepool consists of three distinct lineages

We sampled the full breadth and depth of GGA^tA^t wheat diversity and discovered a clear genetic and geographic differentiation among extant GGA^tA^t wheats. Surprisingly, and supported by all marker types, three clearly distinct lineages were identified. The first lineage is comprised of all *T. timopheevii* genotypes (and the derived *T. militinae* and *T. zhukovskyi*; note that all *T. militinae* and all *T. zhukovskyi* accessions maintained *ex situ* in genebanks are each derived from only one original genotype). Interestingly, *T. araraticum* consists of two lineages that we preliminarily describe as 'ARA-0' and 'ARA-1'. This finding is in contrast to Kimber and Feldman (1987) who concluded that *T. araraticum* does not contain cryptic species, molecularly distinct from those currently recognized. Based on passport data, ARA-0 was found across the whole predicted area of species distribution. ARA-1 was only detected in south-eastern Turkey and in neighboring north-western Syria. It is interesting to note that only in this part of the Fertile Crescent, the two wild tetraploid wheat species, *T. dicoccoides* and *T. araraticum*, grow in abundance in mixed stands.

Based on Figure 3, collecting gaps are evident for *T. araraticum*, and future collecting missions should focus on four specific regions: (i) between the Euphrates river in the west and the Elazığ – Silvan – Mardin transect region in the east. Interestingly, so far only *T. dicoccoides* was reported from this region; (ii) between Bitlis (Turkey) – Amadiyah (Iraq) in the west and the Armenian border in the east including north-western Iran; (iii) between Adıyaman – Silvan in the south and Tunceli in the north; and (iv) between Hama in the south and west/north-west of Aleppo in Syria.

We found only two *T. araraticum* populations which contained representatives of both ARA lineages: (i) 45 km south-east of Kahramanmaraş to Gaziantep, and (ii) 4 km north of St. Simeon on the road to Afrin in Syria. However, in both cases, the ARA-1 lineage was significantly more frequent than ARA-0. This suggests at least a certain level of taxon boundary between ARA-0 and ARA1 lineages and should be investigated in the future.

Independent support for the existence of two wild *T. araraticum* lineages and their distribution comes from Mori et al. (2009) based on 13 polymorphic chloroplast microsatellite markers (cpSSR) (Supplementary Table S2, column 15). The 'plastogroup G-2' was distributed in south-eastern Turkey and northern Syria and was closely related to *Triticum timopheevii* (Mori et al. 2009). However, Gornicki et al. (2014), based on whole chloroplast genome sequence information and sufficient taxon sampling (13 Triticeae species and 1127 accessions; 163 accessions in common with our study, Supplementary Table S2, column 17), provided increased resolution of the chloroplast genome phylogeny and showed that the *T. timopheevii* lineage possibly originated in northern Iraq (*and thus according to our data, belong to the ARA-0 lineage as no ARA-1 occurs in Iraq*). This was supported by Bernhardt et al. (2017), who, based on re-sequencing 194 individuals at the chloroplast locus *ndhF* (2232 bp) and on whole genome chloroplast sequences of 183 individuals representing 15 Triticeae genera, showed that some ARA-0 and TIM genotypes are most closely related. All GGA^tA^t wheats re-sequenced by Bernhardt et al. (2017) were considered in our study (Supplementary Table S2, column 16). Haplotype analysis of the *Brittle rachis 1* (*BTR1-A*) gene in a set of 32 *T. araraticum* in comparison with two *T. timopheevii* accessions (Nave et al. 2021) also showed closer relationships of domesticated *T. timopheevii* to wild *T. araraticum* from Iraq. That is more, one of these accessions, TA102 (PI 538461, 1 km NE of Salahaddin) shared the same haplotype with *T. timopheevii* and it was assigned to ARA-0 group by our study (Supplementary Table S2, column 18).

It is important to note that our results (i.e., the characteristics, composition and geographical distribution of ARA-0 and ARA-1 lineages) are not in agreement with the latest comprehensive taxonomical classification of wheat by Dorofeev et al. (1979), who divided *T. araraticum* into two subspecies: subsp. *kurdistanicum* Dorof. et Migusch. and subsp. *araraticum* (Supplementary Table S2, column 2). We propose to re-classify the GGA^tA^t genepool taxonomically in the future.

2. The karyotypic composition of GGA^tA^t wheats is as complex as the phylogenetic history of the GGA^tA^t genepool

Based on karyotype analyses, translocation spectra and distribution of DNA probes, *T. araraticum* populations from Dahuk and Sulaymaniyah (both Iraq) harbored the highest karyotypic diversity among all *T. araraticum* populations studied. We consider the region around Dahuk in Northern Iraq as the center of diversity of *T. araraticum*, and this is probably the region where *T. araraticum* originated. This is supported by Nave et al. (2021), who found the highest haplotype diversity among *T. araraticum* from Iraq, and by Bernhardt et al. (2017) and

Gornicki et al. (2014) who traced chloroplast haplotypes from *Aegilops speltoides* growing in

Iraq via *T. araraticum* (ARA-0) to *T. timopheevii* and *T. zhukovskyi*.

The karyotype 'similar' to those in 'normal' *T. timopheevii* was found in 44.6% of all *T.*

araraticum genotypes. This is the group of candidates, in which the closest wild relative(s) to *T.*

timopheevii is (are) expected. The frequency of the normal karyotype varied among countries

and between populations (Supplementary Figure S12). It is interesting to note that the Samaxi-

Akhsu population in Azerbaijan and some populations near Dahuk (Iraq) possessed mostly

karyotypically normal genotypes. Diagnostic C-bands for the ARA-1 lineage, both with normal

and rearranged karyotypes, were 1A^{t}L3, 4A^{t}S7, 5A^{t}L3, 1GL5, 2GL7, 3GL7+L11 (Fig. 1). As

expected, the number of C-bands characteristic for ARA-0 was smaller (due to the wide

geographical distribution) and only two C-bands were lineage-specific and found in normal as

well as translocated genotypes: 6A^{t}L3 and 5GL15.

However, some FISH patterns suggested that *T. timopheevii* probably originated in Turkey and

probably from ARA-1 (or, ARA-1 and TIM may have originated from a common ancestor, but

then diverged). This is supported by the following observations: (i) TIM and ARA-1 carry the

pSc119.2 signal in the middle of 1A^t long arm, while this site was absent from ARA-0; (ii) all

ARA-0 and most ARA-1 possessed the Spelt-52 signal on 6GL, but it is absent in all TIM and

five ARA-1 genotypes from Gaziantep-Kilis, Turkey. The distribution of Spelt-1 and Spelt-52

probes on chromosomes of these five genotypes was similar to, and in accession IG 116165

(ARA-1 from Gaziantep) almost identical with TIM; (iii) the pAesp_SAT86 patterns on

chromosomes 3A^t, 4G, and 7G are similar in TIM and ARA-1 but differed from ARA-0.

Differences between ARA-1 and TIM based on FISH patterns of some other chromosomes as

well as the results of C-banding and molecular analyses suggest that extant ARA-1 genotypes are

not the direct progenitors of TIM but that the ARA-1 lineage is most closely related to it.

Based on AFLP, C-banding, FISH and *Jeli* retrotransposon markers, TIM was genetically most

closely related to ARA-1. Additional evidence for the close relationship between TIM and ARA-

1 lineages comes from allelic variation at the *VRN-1* locus of genome A^t (Shcherban et al. 2016).

This analysis revealed a 2.7 kb deletion in intron 1 of *VRN-A1* in three *T. timopheevii* and four *T.*

araraticum accessions, which, according to our data, belong to the ARA-1 lineage. However, at

Vrn-G1, TIM from Kastamonu in Turkey (PI 119442) shared the same haplotype (*Vrn1Ga*) with

ARA-1 samples, while TIM from Georgia harbored haplotype *VRN-G1* as found in ARA-0.

These results suggest multiple introgression events and incomplete lineage sorting as suggested

by Bernhardt et al. (2017, 2020).

Regular chromosome pairing observed in the F$_1$ hybrids of lines with 'normal' karyotypes

(Kawahara et al. 1996), identified in our study as ARA-0 × ARA-1 (Supplementary Table S2,

column 14), suggested that karyotypic differences between ARA-0 and ARA-1 lineages are not
associated with structural chromosomal rearrangements such as large translocations or
inversions.

The emergence or loss of most lineage-specific Giemsa C-bands (Fig. 3) or FISH loci
(Supplementary Figure 22, Supplementary Figure 23) could be due to heterochromatin re-
patterning: amplification, elimination or transposition of repetitive DNA sequences. Wide
hybridization can also induce changes in C-banding and FISH patterns of *T. araraticum*
chromosomes. Changes in pAesp_SAT86 hybridization patterns on 4G and 7G chromosomes,
however, are likely to be caused by pericentric inversions, which are also frequent in common
wheat (Qi et al. 2006). The role of inversions in inter- and intraspecific divergence is probably
underestimated. In our case, it seems possible that divergence between ARA-1/TIM (two
inversions) from ARA-0 (no inversion) was associated with at least two pericentric inversions.

We did not find any genotype harboring both ARA-0 and ARA-1 specific FISH sites, although
ARA-0 and ARA-1 genotypes co-existed in two populations in Turkey and Syria. However,
based on FISH (Spelt-1 site on chromosome 6GL and 7GS, respectively), hybridization between
certain ARA-1 and ARA-0 lines can be predicted.

Iran occupies a marginal part of the distribution range of *T. araraticum*. An abundance of the
pericentric inversion of the $7A^t$ chromosome in the Iranian group indicates that it is derived from
Iraq. The karyotypically 'normal' genotype was probably introduced to Transcaucasia via
Western Azerbaijan (Iran). The low diversity of FISH patterns and the low C-banding
polymorphism of *T. araraticum* from Transcaucasia indicate that *T. araraticum* was introduced
as a single event. Interestingly, the AFLP data suggested some similarity between ARA-0 from
Armenia and Azerbaijan and *T. timopheevii*.

We hypothesize that homoploid hybrid speciation (HHS) (Abbott et al. 2010; Nieto Feliner et al.
2017; Soltis and Soltis 2009) and incomplete lineage sorting may be the possible mechanisms
explaining the origin of the ARA-1 lineage. Although this assumption was not experimentally
supported, it is favored by some indirect evidence. ARA-1 grows in sympatry and in mixed
populations with *T. dicoccoides* (Fig. 3) and is phylogenetically most closely related to *T.
dicoccoides* (Supplementary Table S6; Supplementary Table S8). ARA-1 is morphologically
more similar to *T. dicoccoides*. Thus, five of 10 misclassified *T. araraticum* accessions belonged
to ARA-1 group (Supplementary Table S2), two of which, PI 656871 and IG 116176, were the
mix of *T. dicoccoides* and ARA-1 . Five misclassified ARA-0 accessions from USDA-ARS
collection were from Siirt, Turkey, however in other gene bank three of these accessions were
treated as *T. araraticum*. Relatively good chromosome pairing was observed in the F_1 hybrids of
some *T. araraticum x T. timopheevii* combinations (Tanaka and Ichikawa 1972), however, pollen

fertility of such hybrids was very low (0.3 – 5.4%). ARA-1 could be derived from ancient

hybridization of *T. timopheevii* × *T. dicoccoides*; or alternatively, ARA-1 and TIM could be

derived from the hybridization ARA-0 × *T. dicoccoides*.

3. Does the *T. timopheevii* population found in western Georgia represent the last remnant

of a widespread ancient cultivation area of GGA^{t}A^t wheats?

Wild emmer *T. dicoccoides* belongs to the first cereals to be domesticated by humans in the

Fertile Crescent and the evolution and domestication history of *T. dicoccoides* are relatively well

studied (Badaeva et al. 2015b; Civáň et al. 2013; Özkan et al. 2011). Domesticated emmer

Triticum dicoccon Schrank was a staple crop of Neolithic agriculture, was widely cultivated for

over 10,000 years and harbored impressive genetic diversity (Nesbitt and Samuel 1996; Szabo

and Hammer 1996; Zaharieva et al. 2010). The domestication of *T. dicoccoides* provided the key

for durum wheat (Maccaferri et al. 2019) and bread wheat evolution (Pont et al. 2019).

Much less is known about the domestication history of *T. timopheevii*. It is believed that *T.*

timopheevii is the domesticated form of *T. araraticum* (Dorofeev et al. 1979; Jakubziner 1932).

In contrast to *T. dicoccon*, *T. timopheevii* was, since its discovery, considered as a

'*monomorphous narrowly endemic species*' (Dekaprelevich and Menabde 1932) cultivated in

few villages of western Georgia (Stoletova 1924-25; Zhukovsky 1928) (Supplementary Material

S27). Dekaprelevich and Menabde (1932) noticed that the area of cultivation had probably been

larger in the past. The last plants of *T. timopheevii in situ* were found by the expedition of the

N.I. Vavilov Institute of Plant Genetic Resources (VIR, Russia) in 1983 near the village of

Mekvena (Tskhaltubo, Georgia) and deposited in the VIR genebank under accession number K-

56422 [E.V. Zuev, personal communication]. Today, the widespread view is that the cultivation

area of *T. timopheevii* was restricted to Georgia in the (recent) past (Feldman 2001; Mitrofanova

et al. 2016, Zohary et al. 2012).

However, hulled tetraploid wheat morphologically similar to *T. timopheevii* was identified at

three Neolithic sites and one Bronze Age site in northern Greece and described by Jones et al.

(Jones et al. 2000) as a **'**New' Glume Wheat (new glume wheat, NGW). The glume bases of

these archaeological finds morphologically resemble *T. timopheevii* more than any other extant

domesticated wheat (Jones et al. 2000). After these finds of NGW in Greece, this wheat was also

identified at Neolithic and Bronze Age sites in Turkey, Bulgaria, Romania, Hungary, Slovakia,

Austria, Italy, Poland, Germany and France (Bieniek 2002, 2007; Bogaard et al. 2007, 2013;

Ergun 2018; Fairbairn et al. 2002; Fiorentino and Ulaş 2010; Fischer and Rösch 2004; Hajnalová

2007; Kenéz et al. 2015; Kohler-Schneider 2003; Kreuz and Boenke 2002; Perego 2017; Rottoli

and Pessina 2007; Toulemonde et al. 2015; Ulaş and Fiorentino 2020; Valamoti and Kotsakis

2007). Earlier finds of an 'unusual' glume wheat in Serbia (Borojevic 1991) and Turkey (de

Moulins 1997) have subsequently been recognized as NGW (Kenéz et al. 2015; Jones et al.

2000; Kroll 2016). Criteria were also established for distinguishing the grains of NGW (Kohler-

Schneider 2003).

At some sites, NGW appeared as a minor component and may have been part of the

accompanying weed flora of cereal fields (Kenéz et al. 2014; Ulaş and Fiorentino 2020). In other

cases, it was probably cultivated in a mix with einkorn (Jones et al. 2000; Kohler-Schneider

2003) and/or emmer. The recovery of large quantities in storage deposits of whole spikelets at

Çatalhöyük in Turkey, caryopses and spikelet bases at the early Bronze Age settlement of

Clermont-Ferrand in France, and rich deposits including whole spikelets at bronze age sites in

Italy, demonstrated that, at least in some places, NGW was a major crop in itself (Bogaard et al.

2013, 2017; Ergun 2018; Kenéz et al. 2014; Perego 2017; Toulemonde et al. 2015).

Based on intensive archaeobotanical investigations at Çatalhöyük in central Anatolia, for

example, NGW was the predominant hulled wheat, overtaking emmer wheat around 6500 cal BC

and remaining dominant until the site's abandonment c. 5500 cal BC. The finds suggested that

this wheat was a distinct crop, processed, stored, and presumably grown, separately from other

glume wheats. NGW formed part of a diverse plant food assemblage at Neolithic

Çatalhöyük, including six cereals, five pulses and a range of fruits, nuts and other plants,

which enabled this early farming community to persist for 1500 years (c. 7100 to 5500 cal BC)

(Bogaard et al. 2013, 2017).

Recently, polymerase chain reactions specific for the wheat B and G genomes, and extraction

procedures optimized for retrieval of DNA fragments from heat-damaged charred material, have

been used to identify archaeological finds of NGW (Czajkowska et al. 2020). DNA sequences

from the G genome were detected in two of these samples, the first comprising grain from the

mid 7[th] millennium BC at Çatalhöyük in Turkey, and the second made up of glume bases from

the later 5[th] millennium BC site of Miechowice 4 in Poland. These results provide evidence that

NGW is indeed a cultivated member of the GGA^{t}A^t genepool (Czajkowska et al. 2020). As

NGW is a recognized wheat type across a broad geographic area in prehistory, dating back to the

9[th] millennium BC in SW Asia, this indicates that *T. timopheevii* (*sensu lato, s.l. = domesticated*

GGA^{t}A^t wheat in general), was domesticated from *T. araraticum* during early agriculture, and

was widely cultivated in the prehistoric past (Czajkowska et al. 2020).

This raises the question of whether the few populations of *T. timopheevii* (*sensu stricto, s.str.*)

found in western Georgia were the last remnants of a wider GGA^{t}A^t wheat cultivation or whether

the *T. timopheevii* of Georgia was a local domestication independent of the domestication of *T.*

araraticum in SW Asia. To answer this question, sequence information for NGW, the Georgian

T. timopheevii, and the two lineages of *T. araraticum* (ARA-0, ARA-1) need to be compared.

Vavilov (1935) suggested that *T. timopheevii* of western Georgia was probably originally

introduced from north-eastern Turkey. As cited by Dorofeev et al. (1979), Menabde and Ericzjan

(1942) associated the origin of *T. timopheevii* with the region of the ancient kingdom of Urartu,

whence immigrant ancestors of modern-day Georgians introduced it into western Georgia.

Certainly, the possibility of introduction of Timofeev's wheat into Georgia from the south should

not be rejected (Dorofeev et al. 1979). Is the domestication history of *T. timopheevii s.str.*

connected with other endemic wheats of Georgia, such as *T. karamyschevii* Nevski (*T.*

georgicum Dekapr. et Menabde or *T. paleocolchicum* Menabde) and *T. carthlicum* Nevski,

which were cultivated by Mingrelians in Western Georgia (Jorjadze et al. 2014; Mosulishvili et

al. 2017)?

4. Was the cultivation range of *T. timopheevii* (*s.str.*) wider in the recent past?

We screened all available passport data and found three cases, which could potentially help to

reconstruct the recent past cultivation range of *T. timopheevii s.str*: Interestingly, two *T.*

timopheevii accessions maintained in two *ex situ* genebanks are reported to originate from

Turkey (Supplementary Table S2) [https://www.genesys-pgr.org/]: (i) ATRI 3433 (TRI 3433)

conserved in the Federal *ex situ* Genebank of Germany hosted at the Institute of Plant Genetics

and Crop Plant Research, IPK, Gatersleben. This *T. timopheevii* line was most likely collected by

E. Baur in Turkey in 1926 (Schiemann 1934); and (ii) PI 119442 identified among a barley

sample obtained from a market in Araç, near Kastamonu, Turkey (Fig. 3) in 1936 by Westover

and Wellmann, and maintained at the National Plant Germplasm System, USDA-ARS, USA.

Both accessions harbor the 'normal' karyotype of *T. timopheevii*, both were characterized in our

studies and confirmed as a typical *T. timopheevii*. Additionally, the accession TA1900,

presumably collected 32 km south of Denizli near Kahramanmaraş in Turkey on the 14th of

August 1959 and maintained in the wheat germplasm collection of the Wheat Genetics Resource

Center, Kansas State University, U.S.A., is interesting because it shared karyotypic features of *T.*

timopheevii and the ARA-0 lineage (Fig. 4; Supplementary Figure S19, *i31*). However, we are

not fully convinced that this line is a true natural hybrid. Based on the passport data, this

accession could potentially have escaped from an experimental field or a breeding station, or

received introgression(s) during *ex situ* maintenance (Zencirci et al. 2018). Assuming that the

passport data is correct, we could speculate that *T. timopheevii* may have been cultivated in

Turkey during the first half of the 20th century. However, is this realistic option?

We believe not. As reported by Stoletova (1924-25), Dekaprelevich and Menabde (1929, 1932),

Menabde (1948), Dekaprelevich (1954), *T. timopheevii s.str.* was part of the spring landrace

Zanduri (mixture of *T. timopheevii s.str.* and *T. monococcum*) and well adapted to the historical

provinces Lechkhumi and Racha of Georgia (Supplementary Material S28). The Zanduri

landrace was cultivated in the '*humid and moderately cool climate zone 400–800*' m above the

sea level (Dorofeev et al. 1979). Martynov et al. (2018) reported that *T. timopheevii* potentially

has a '*low potential for plasticity*' and is not drought tolerant. Climate at origin based on

bioclimatic variables (Fick et al. 2017; R Core Team 2017) clearly differs between the regions of

Western Georgia where the Zanduri landrace grew till the recent past, and both Kastamonu and

Kahramanmaraş regions in Turkey (Supplementary Material S28). We speculate that the three *T.*

timopheevii accessions which were collected in Turkey were probably introduced from

Transcaucasia or elsewhere and may have been left over from unsuccessful cultivation or

breeding experiments of *T. timopheevii s.str.* in the recent historical past. Also, based on

botanical records, *T. timopheevii (s.str.)* has only been identified in Georgia, but not in Turkey or

elsewhere (Davis 1965-1988; Hanelt 2001). From this we conclude that the cultivation range of

T. timopheevii (*s str.*) was not wider in the recent past.

Conclusions

The evolutionary history of the GGA^{t}A^t wheat genepool is complex. However, some pieces of

the puzzle are clearly recognizable: the region around Dahuk in northern Iraq can be considered

the center of origin, but also the center of diversity of *T. araraticum.*

The origin of *T. timopheevii s.str.* remains unclear, but we speculate that it was probably

introduced from Turkey, on the grounds that wild *T. araraticum* does not grow in Georgia and

that *T. timopheevii s.str.* is more closely related to *T. araraticum* from Turkey or northern Iraq

than to the Transcaucasian types.

Based on bioclimatic variables, we predict that *T. timopheevii s.str.* is maladapted to the climate

outside Western Georgia. If this speculation is correct, it suggests a sister-group relationship

between (i) the Georgian *T. timopheevii* (*s.str.*) and both *T. araraticum* lineages (ARA-0, ARA-

1), but also between (ii) *T. timopheevii s.str.* and the prehistoric SW Asian *T. timopheevii s.l.*

The oldest known records of prehistoric *T. timopheevii s.l.* are of Turkish origin: (i) Aşıklı

Höyük in Cappadocia, and (ii) Cafer Höyük (just inside the Fertile Crescent and potentially older

than Aşıklı Höyük but less thoroughly dated) (Cauvin et al. 2011; Quade et al. 2018), though

more prehistoric finds may be identified in the future. Emphasis for further archaeobotanical

research should be given to the central part of the Fertile Crescent, including northern Iraq

because this is probably the region where wild *T. araraticum* originated.

The unexpected discovery of the ARA-1 lineage is exciting and may have implications for our

understanding of the origins of agriculture in southwest Asia. Was ARA-1 involved in the

formation of *T. timopheevii s.l.* (= NGW)?

The distribution area of the ARA-1 lineage requires our attention. It is interesting to note that the

ARA-1 lineage (GGA^{t}A^t) and *T. dicoccoides* (BBAA) grow in sympatry and in mixed

populations only in a peculiar geographical area in the Northern Levant. This specific area is

located between Kahramanmaraş in the north, Gaziantep in the east, Aleppo in the south and the

eastern foothills of the Nur Dağlari mountains in the west. It is interesting that the closest extant

wild relatives of domesticated einkorn, barley and emmer were also collected here: (i) the beta

race of wild einkorn (Kilian et al. 2007) and the closest wild relatives to einkorn *btr1* type

(Pourkheirandish et al. 2018); (ii) the closest wild barley to *btr2* barley (Pourkheirandish et al.

2015) and to *btr1b* barley (Civáň and Brown 2017). This specific region was among the areas

predicted with high probability as potential refugia for wild barley during the Last Glacial

Maximum (LGM) by Jakob et al. (2014); and (iii) wild emmer subgroup II of the central-eastern

race (Özkan et al. 2011). On the other hand, some genetic research has suggested that

domesticated emmer wheat and barley received substantial genetic input from other regions of

the Fertile Crescent, resulting in hybridized populations of different wild lineages indicating a

mosaic of ancestry from multiple source populations (Civáň et al. 2013; Oliveira et al. 2020;

Poets et al. 2015). It will be interesting to see whether further genetic and archaeobotanical

research on *T. araraticum* lineages and *T. timopheevii s.l.* can help to resolve this issue.

Finally, and fortunately, 1294 accessions of *T. araraticum* and 590 accessions of *T. timopheevii*

s.str. are stored in 24 and 37 *ex situ* genebank repositories, respectively (Knüpffer 2009). Our

study provides the basis for a more efficient use of *T. araraticum* and *T. timopheevii* materials

for crop improvement.

Acknowledgements

We thank the following for providing seeds, DNA and passport data: N. I. Vavilov Institute of
Plant Genetic Resources (VIR, Russia), the Federal *ex situ* Genebank of Germany, Gatersleben
(IPK, Germany), the International Center for Agricultural Research in the Dry Areas (ICARDA),
the Max Planck Institute for Plant Breeding Research (MPIPZ, Germany), Kyoto University
(National Bioresource Project, NBRP, Japan), the John Innes Centre (JIC, UK), Wheat Genetics
Resource Centre, Kansas State University (WGRC, USA), Centre for Genetic Resources (WUR,
CGN, Netherlands), the Australian Winter Cereal Collection Tamworth (AWCC Australia), the
National Plant Germplasm System (USDA-ARS, USA), Eastern Cereal and Oilseed Research

Centre, Agriculture and Agri-Food Canada (Canada), Botanical Institute Tbilisi (Georgia), Tbilisi Agricultural Institute (Georgia), Institute of Cytology and Genetics (Siberian Branch of the Russian Academy of Sciences, Russia), and the Russian State Agrarian University Moscow, Timiryazev Agricultural Academy (Russia). We also thank Drs. P.A. Gandilyan, M. Nazarova and I.G. Gukasyan (all Botanical Institute, Erevan, Armenia), Drs. I.D.O. Mustafaev and N.Kh. Aminov (both Institute of Genetic Resources of Azerbaijan, Baku, Azerbaijan) for providing materials from their collection missions. We thank the following for excellent technical assistance: Sigi Effgen (MPIPZ), Christiane Kehler, Birgit Dubsky, Heike Harms, Ute Krajewski, Marita Nix, Kerstin Wolf, Jürgen Marlow, Michael Grau, Peter Schreiber (all IPK), the 'Experimental Fields and Nurseries' and 'Genome Diversity' groups at IPK. We thank Dr. E.V. Zuev for providing us the data on collection sites of *T. araraticum* and *T. timopheevii* accessions from VIR collection. We are greatly indebted to Maarten Koornneef and George Willcox for discussions and support. We thank two anonymous reviewers, who helped us to improve an earlier version of this manuscript.

 Legends to Figures

Fig. 1 Generalized idiogram and nomenclature of the A^t- and G-genome chromosomes. The
C-banding pattern is shown on the left, the pSc119.2 (red) and pAesp_SAT86 (green) pattern on
the right side of each chromosome. 1–7 – homoeologous groups; S – short arm, L – long arm.
The numerals on the left-hand side designate putative positions of C-bands/FISH sites that can
be detected on the chromosome arm; C-bands specific for the ARA-1 group are shown with pink
numerals, C-bands specific for the ARA-0 group are indicated by green numerals. Red asterisks
on the right-hand side indicate C-bands that were considered for the "chromosomal passport".

Fig. 2 Genetic relationships between GGA^{t}A^t and BBAA wheats. NeighborNet planar graph
of Dice distances representing the diversity of 787 GGA^{t}A^t and BBAA tetraploid wheat
genotypes based on 656 SSAP markers.

Fig. 3 Natural geographical distribution of wild tetraploid *T. araraticum* and *T. dicoccoides*.
Green dots correspond to collection sites of ARA-0 accessions, pink dots to ARA-1, and dark
blue dots to *T. dicoccoides* (DIC). The collection sites of *T. timopheevii* and *T. zhukovskyi* are
shown with turquoise and yellow dots, respectively. Key excavation sites in Turkey where NGW
was identified are indicated with red triangles.

Fig. 4 Comparison of the C-banding patterns of *T. dicoccoides* (DIC), *T. timopheevii* (TIM,
***a–c,* normal karyotypes), *T. araraticum* ARA-1 (*d–f*) and ARA-0 (*g–t*).** DIC (IG 117174,
Gaziantep), *a* – KU-1818 (Georgia); *b* – PI 119442; *c* – TA1900; *d* – IG 116165; *e* – PI 654340;
f – KU-1950; *g* – CItr 17677; *h* – KU-8917; *i* – KU-8909; *j* – KU-1933 (all from Turkey); *k* –
CItr 17680 (Iran); *l* – PI 427381 (Erbil, Iraq); *m* – PI 538518*; n* – PI 427425 (Dahuk, Iraq); *o* –
KU-8705; *p* – KU-8695 (Shaqlawa, Erbil, Iraq); *q* – KU-8451; *r* – KU-8774 (Sulaymaniyah,
Iraq); *s* – TRI 11945 (Nakhichevan); *t* – KU-1901 (Armenia). 1–7 – homoeologous groups. C-
bands typical for ARA-1 are indicated with blue arrows, for ARA-0 – with green arrows, and C-
bands characteristic for *T. timopheevii* – with red arrows. Black arrows point to rearranged
chromosomes in genotypes.

Fig. 5 Chromosomal rearrangements identified in *T. araraticum*. The number of
translocation variant corresponds to the number of the respective variant in Supplementary Table
S13. Novel variants are designated with black numbers, and already known variants by red
numbers.

Fig. 6 Intraspecific divergence of *T. araraticum* and *T. timopheevii*. Combinations of
chromosome arms in rearranged chromosomes are designated. Line colors mark the different
groups: ARA-0 (green), ARA-1 (pink) and *T. timopheevii* (black). Solid arrows designate novel
rearrangements; arrows with asterisk designate previously described rearrangements (Badaeva et
al. 1990, 1994). The numerals above/next to the arrows indicate the number of accessions
carrying the respective translocation.

Fig. 7 Distribution of different families of tandem repeats on chromosomes of *T.
***timopheevii* and *T. araraticum*.** *Triticum timopheevii,* KU-107 (*a*), and *T. araraticum,* CItr
17680, ARA-0 (*b*), KU-8944, ARA-0 (*c*), KU-1984B, ARA-1 (*d*), PI 427364, ARA-0 (*e*), and
2630, ARA-1 (*f*). The following probe combinations were used: *a, b* – pSc119.2 (green) + pTa-
535 (red); *d, e* – pAesp_SAT86 (red) + GAA$_n$ (green); *c, f* – Spelt-1 (red) + Spelt-52 (green). The
position of pSc119.2 site on 1A^t chromosome typical for *T. timopheevii* and ARA-1 is shown

with an arrow (*a*). Translocated chromosomes (*c, d*) are arrowed. Chromosomes are designated
according to genetic nomenclature; the A^t-genome chromosomes are designated with yellow
numerals and the G-genome chromosomes with white numerals. Scale bar, 10 μm.

Electronic supplementary material

The online version of this article (https:// doi. org/ xxx) contains supplementary material, which
is available to authorized users.
801

Supplementary information

Supplementary information accompanies this paper at <mark>http xxx</mark>

Declarations

Funding

This work was supported by the European Community's Seventh Framework Programme (FP7,
2007–2013) under the grant agreement n°FP7-613556, Whealbi project
[http://www.whealbi.eu/project/], the German Research Foundation (DFG) (FKZ KI 1465/1-1
and FKZ KI 1465/5-1), the Federal Office for Agriculture and Food (BLE) of Germany (FKZ
01/12-13-KAD), the Leibniz Institute of Plant Genetics and Crop Research (IPK, Gatersleben,
Germany), the Max Planck Institute for Plant Breeding Research (MPIPZ, Cologne, Germany)
and the Crop Wild Relatives Project (*Adapting Agriculture to Climate Change: Collecting,*
Protecting and Preparing Crop Wild Relatives) which is supported by the Government of
Norway and managed by the Global Crop Diversity Trust [https://www.cwrdiversity.org/]. EDB
acknowledges support from the Russian State Foundation for Basic Research (projects 20-04-
00284, 17-04-00087, 14-04-00247), State Budget Project and the Program "Dynamics and
Preservation of Gene Pools" from the Presidium of the Russian Academy of Sciences.

Conflicts of interest/ Competing interests
The authors declare that they have no conflict of interest.

Availability of data and material

All data sets supporting the conclusions of this article are available in the Electronic
supplementary material and from the corresponding author Ekaterina D. Badaeva
(katerinabadaeva@gmail.com). The acquisition of plant material used in this study complies
with institutional, national, and international guidelines.

Code availability

Not applicable

Authors' contributions

EB conceived, designed and conducted the cytogenetic experiments and analyzed data, wrote the first manuscript version.

FK conducted the SSAP experiments and analyzed the data.

HK verified genebank information concerning the origin of the material, translated texts from Russian into English, and contributed to the discussions and editing of the manuscript.

AF, supported the molecular data analysis.

AR contributed to the FISH analysis.

ZK conducted the simulation of regional agro-ecological variation based on bioclimatic variables.

SZ contributed to the work on mapping of Spelt-1 and Spelt-52 probes on *T. timopheevii* and *T. araraticum* genotypes.

SS designed and synthesized oligo-probes for FISH analyses.

KN contributed to data analysis, supported the field trials and edited the manuscript.

AG contributed to the discussions and editing of the manuscript.

AF phenotyped and taxonomically re-identified the collection, contributed to the discussions and edited the manuscript.

KH phenotyped and taxonomically re-identified the collection, contributed to the discussions and edited the manuscript.

ABcontributed to the discussions and editing of the manuscript.

GJ contributed to the discussions and editing of the manuscript.

HÖ conducted the AFLP experiments, analyzed the data,contributed to the discussions and edited the manuscript.

BK conceived the project, established the germplasm collection, analyzed data, interpreted results, wrote and edited the manuscript.

Ethics approval
The authors declare that the experiments comply with the current laws of Germany.

Consent to participate
Not applicable

Consent for publication
Not applicable

Open Access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative

References

Abbott RJ, Hegarty MJ, Hiscock SJ, Brennan AC (2010) Homoploid hybrid speciation in action.
TAXON 59:1375-1386. https://doi.org/10.1002/tax.595005

Allard RW, Shands RG (1954) Inheritance of resistance to stem rust and powdery mildew in
cytologically stable spring wheats derived from *Triticum timopheevi*. Phytopathol 44:266-
274.

Alsaleh A, Baloch FS, Derya M, Azrak M, Kilian B, Özkan H, Nachit M (2015) Genetic linkage
map of Anatolian durum wheat derived from a cross of Kunduru-1149 × Cham1. Plant Mol
Biol Rep 33:209-220. https://doi.org/10.1007/s11105-014-0749-6

Altıntaş S, Toklu F, Kafkas S, Kilian B, Brandolini A, Özkan H (2008) Estimating genetic
diversity in durum and bread wheat cultivars from Turkey using AFLP and SAMPL
markers. Plant Breed 127:9-14. https://doi.org/10.1111/j.1439-0523.2007.01424.x

Avni R, Nave M, Barad O, Baruch K, Twardziok SO, Gundlach H, Hale I, Mascher M, Spannagl
M, Wiebe K, Jordan KW, Golan G, Deek J, Ben-Zvi B, Ben-Zvi G, Himmelbach A,
MacLachlan RP, Sharpe AG, Fritz A, Ben-David R, Budak H, Fahima T, Korol A, Faris
JD, Hernandez A, Mikel MA, Levy AA, Steffenson B, Maccaferri M, Tuberosa R,
Cattivelli L, Faccioli P, Ceriotti A, Kashkush K, Pourkheirandish M, Komatsuda T, Eilam
T, Sela H, Sharon A, Ohad N, Chamovitz DA, Mayer KFX, Stein N, Ronen G, Peleg Z,
Pozniak CJ, Akhunov ED, Distelfeld A (2017) Wild emmer genome architecture and
diversity elucidate wheat evolution and domestication. Science (New York) 357:93-97.
https://doi.org/10.1126/science.aan0032

Badaeva ED, Amosova AV, Goncharov NP, Macas J, Ruban AS, Grechishnikova IV, Zoshchuk
SA, Houben A (2015a) A set of cytogenetic markers allows the precise identification of all
A-genome chromosomes in diploid and polyploid wheat. Cytogenet Genome Res 146:71-
79. https://doi.org/10.1159/000433458

Badaeva ED, Badaev NS, Gill BS, Filatenko AA (1994) Intraspecific karyotype divergence in
Triticum araraticum (Poaceae). Plant Syst Evol 192:117-145.
https://doi.org/10.1007/BF00985912

Badaeva ED, Boguslavsky RL, Badaev NS, Zelenin AV (1990) Intraspecific chromosomal
polymorphism of *Triticum araraticum* (Poaceae) detected by C-banding technique. Plant
Syst Evol 169:13-24. https://doi.org/10.1007/BF00935980

Badaeva ED, Budashkina EB, Badaev NS, Kalinina NP, Shkutina FM (1991) General features of chromosome substitutions in *Triticum aestivum x T. timopheevii* hybrids. Theor Appl Genet 82:227-232. https://doi.org/10.1007/BF00226218

Badaeva ED, Keilwagen J, Knüpffer H, Waßermann L, Dedkova OS, Mitrofanova OP, Kovaleva ON, Liapunova OA, Pukhalskiy VA, Özkan H, Graner A, Willcox G, Kilian B (2015b) Chromosomal passports provide new insights into diffusion of emmer wheat. PLoS ONE 10:e0128556. https://doi.org/10.1371/journal.pone.0128556

Badaeva ED, Ruban AS, Zoshchuk SA, Surzhikov SA, Knüpffer H, Kilian B (2016) Molecular cytogenetic characterization of *Triticum timopheevii* chromosomes provides new insight on genome evolution of *T. zhukovskyi*. Plant Syst Evol:1-14. https://doi.org/10.1007/s00606-016-1309-3

Badaeva ED, Shkutina FM, Bogdevich IN, Badaev NS (1986) Comparative study of *Triticum aestivum* and *T. timopheevii* genomes using C-banding technique. Plant Syst Evol 154: 183-194. https://doi.org/10.1007/BF00990122

Bedbrook RJ, Jones J, O'Dell M, Thompson RJ, Flavell RB (1980) A molecular description of telomeric heterochromatin in *Secale* species. Cell 19:545-560. https://doi.org/10.1016/0092-8674(80)90529-2

Bernhardt N, Brassac J, Dong X, Willing E-M, Poskar CH, Kilian B, Blattner FR (2020) Genome-wide sequence information reveals recurrent hybridization among diploid wheat wild relatives. Plant J 102: 493-506. https://doi.org/10.1111/tpj.14641

Bernhardt N, Brassac J, Kilian B, Blattner FR (2017) Dated tribe-wide whole chloroplast genome phylogeny indicates recurrent hybridizations within Triticeae. BMC Evolut Biol 17:141. https://doi.org/10.1186/s12862-017-0989-9

Bieniek A (2002) Archaeobotanical analysis of some early Neolithic settlements in the Kujawy region, central Poland, with potential plant gathering activities emphasised. Veget Hist Archaeobot 11:33-40. https://doi.org/10.1007/s003340200004

Bieniek A (2007) Neolithic plant husbandry in the Kujawy region of central Poland. In: Colledge S, Conolly J (eds) The Origin and Spread of Domestic Plants in Southwest Asia and Europe. Left Coast Press, Walnut Creek, pp 327–342

Bogaard A, Bending J, Jones G (2007) Archaeobotanical evidence for plant husbandry and use. In: Whittle AWR (ed) The Early Neolithic on the Great Hungarian Plain Investigations of

the Körös culture site of Ecsegfalva 23, County Békés. Institute of Archaeology,
Hungarian Academy of Sciences, Budapest, pp 421–445

Bogaard A, Charles M, Livarda A, Ergun M, Filipovic D, Jones G (2013) The archaeobotany of
mid-later Neolithic occupation levels at Çatalhöyük. In: Hodder I (ed) Humans and
Landscapes of Çatalhöyük: Reports from the 2000-2008 seasons. Monographs of the
Cotsen Institute of Archaeology, University of California at Los Angeles, Los Angeles, pp
93-128

Bogaard A, Filipović D, Fairbairn A, Green L, Stroud E, Fuller D, Charles M (2017)
Agricultural innovation and resilience in a long-lived early farming community: the 1,500-
year sequence at Neolithic to early Chalcolithic Çatalhöyük, central Anatolia. Anatolian
Studies 67:1-28

Borojevic K (1991) Emmer aus Feudvar. In: Hänsel B, Medović P (eds) Vorbericht über die
jugoslawisch-deutschen Ausgrabungen in der Siedlung von Feudvar bei Mosorin (Gem
Titel, Vojvodina) von 1986–1990: Bronzezeit–Vorrömische Eisenzeit. Verlag Philipp von
Zabern, Darmstadt, pp 171–177

Brown-Guedira G, Badaeva ED, Gill BS, Cox TS (1996) Chromosome substitutions of *Triticum*
timopheevii in common wheat and some observations on the evolution of polyploid wheat
species. Theor Appl Genet 93:1291- 1298. https://doi.org/10.1007/BF00223462

Brown-Guedira GL, Singh S, Fritz AK (2003) Performance and mapping of leaf rust resistance
transferred to wheat from *Triticum timopheevii* subsp. *armeniacum*. Phytopathol 93:784-
789

Cauvin J, Aurenche O, Cauvin M-C, Balkan-Atlı N (2011) The Pre-Pottery site of Cafer Höyük.
In: Özdogan M, Başgelen N, Kuniholm P (eds) The Neolithic in Turkey. Archaeology and
Art Publications, Istanbul, pp 1-40

Chen PD, Gill BS (1984) The origin of chromosome 4A, and genomes B and G of tetraploid
wheats. In: Sakamoto S (ed) Proceedings of the 6[th] International Wheat Genetics
Symposium, November 28 - December 3 1983. Plant Germ-Plasm Inst., Kyoto University,
Kyoto, Japan, pp 39-48

Civáň P, Brown TA (2017) A novel mutation conferring the nonbrittle phenotype of cultivated
barley. New Phytol 214:468-472. https://doi.org/10.1111/nph.14377

Civáň P, Ivaničová Z, Brown TA (2013) Reticulated origin of domesticated emmer wheat

supports a dynamic model for the emergence of agriculture in the Fertile Crescent. PLoS

ONE 8:e81955. https://doi.org/10.1371/journal.pone.0081955

Czajkowska BI, Bogaard A, Charles M, Jones G, Kohler-Schneider M, Mueller-Bieniek A,

Brown TA (2020) Ancient DNA typing indicates that the "new" glume wheat of early

Eurasian agriculture is a cultivated member of the *Triticum timopheevii* group. J Arch Sci

123:105258. https://doi.org/10.1016/j.jas.2020.105258

Dagan J, Zohary D (1970) Wild tetraploid wheats from West Iran cytogenetically identical with

Israeli *T. dicoccoides*. Wheat Inf Serv 31:15-17

Dante FP, Campbell MT, Folsom JJ., Cui X, Kruger GR., P. Baenziger S, Walia H (2013).

Introgression of novel traits from a wild wheat relative improves drought adaptation in

wheat. Plant Physiol, 161 (4) 1806-1819; DOI: 10.1104/pp.113.214262

Davis PH (1965-1988) Flora of Turkey and the East Aegean Islands. Edinburgh University

Press, Edinburgh

de Moulins D (1997) Agricultural changes at Euphrates and steppe sites in the mid-8[th] to the 6[th]

millennium B.C. Archaeopress, Oxford.

Dekaprelevich LL (1954) Species, varieties and cultivars of Georgian wheat. Tr Inst Polevodstva

AN GSSR 8:3-61 (in Russian)

Dekaprelevich LL, Menabde VL (1929) Regarding the investigation of crop plants in western

Georgia. Zap Nauchno-prikl otdelov Tiflisskogo Bot Sada 6:219–254 (in Georgian)

Dekaprelevich LL, Menabde VL (1932) Hulled wheat in Western Georgia. Bull Appl Bot, Genet

Plant Breed 5:1-46 (in Russian)

Dempewolf H, Baute G, Anderson J, Kilian B, Smith C, Guarino L (2017) Past and future use of

wild relatives in crop breeding. Crop Sci 57:1070-1082.

https://doi.org/10.2135/cropsci2016.10.0885

Dice LR (1945) Measures of the amount of ecologic association between species. Ecology

26:297-302. https://doi.org/10.2307/1932409

Dobrovolskaya O, Sourdille P, Bernard M, Salina E (2009) Chromosome synteny of the a

genome of two evolutionary wheat lines. Rus J Genet 45:1368-1375.

https://doi.org/10.1134/S1022795409110118

Dorofeev VF, Filatenko AA, Migushova EF, Udachin RA, Jakubziner MM (1979) Cultivated
flora of the USSR. 1. Wheat. Kolos, Leningrad (in Russian)

Dvořák J, Mcguire PE, Cassidy B (1988) Apparent sources of the A genomes of wheats inferred
from polymorphism in abundance and restriction fragment length of repeated nucleotide
sequences. Genome 30:680-689. http://doi.org/10.1139/g88-115

Dyck PL (1992) Transfer of a gene for stem rust resistance from *Triticum araraticum* to
hexaploid wheat. Genome 35:788-792. https://doi.org/10.1139/g92-120

El Haddad N, Kabbaj H, Zaïm M, El Hassouni K, Tidiane Sall A, Azouz M, Ortiz R, Baum M,
Amri A, Gamba F, Bassi FM (2021) Crop wild relatives in durum wheat breeding: Drift or
thrift? Crop Sci 61: 37-54. https://doi.org/10.1002/csc2.20223

Ergun M (2018) "Where the wild things are". Contextual insights into wild plant exploitation at
Aceramic Neolithic Aşıklı Höyük, Turkey. Paléorient 44:9-28

Fairbairn A, Asouti E, Near J, Martinoli D (2002) Macro-botanical evidence for plant use at
Neolithic Çatalhöyük south-central Anatolia, Turkey. Veget Hist Archaeobot 11:41-54.
https://doi.org/10.1007/s003340200005

Feldman M (2001) Origin of cultivated wheat. In: Bonjean AP, Angus WJ (eds) The world
wheat book A history of wheat breeding. Tec & Doc/Intersept Ltd., Londres-Paris-New
York, pp 3-56

Fick S E and Hijmans R (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for
global land areas (pp.1-14). Int J Climatol. http://dx.doi.org/10.1002/joc.5086

Fiorentino G, Ulaş B (2010) A recent attestation of new glume wheat type: biometrical and
morphological studies of forks from the Neolithic levels in Mersin-Yumuktepe (Turkey).
In: Bittmann F (ed) 15th Conference of the International Workgroup for
Pelaeoethnobotany, Wilhelmshaven, Germany, May 31 –June 5, 2010. Terra Nostra,
Wilhelmshaven, Germany, v. 2, p 183.

Fischer E, Rösch A (2004) Archäobotanische Untersuchungen. In: Schier W, Draşovean F (eds)
Praehistorische Zeitschrift Vorbericht über die rumänisch-deutschen Prospektionen und
Ausgrabungen in der befestigten Tellsiedlung von Uivar, jud Timiş, Rumänien (1998–
2002), pp 209–220

Gill BS, Chen PD (1987) Role of cytoplasm-specific introgression in the evolution of the
polyploid wheats. Proc Natl Acad Sci USA 84:6800-6804.
https://doi.org/10.1073/pnas.84.19.6800

Goncharov NP (2012) Comparative genetics of wheats and their related species 2edn. Academic
Publishing Group @Geo@, Novosibirsk

Gornicki P, Zhu H, Wang J, Challa GS, Zhang Z, Gill BS, Li W (2014) The chloroplast view of
the evolution of polyploid wheat. New Phytol: 204:704-714.
https://doi.org/10.1111/nph.12931

Hajnalová M (2007) Early farming in Slovakia: an archaeobotanical perspective. In: Colledge S,
Connolly J (eds) The origins and spread of domestic plants in Southwest Asia and Europe.
Left Coastal Press, Walnut Creek, pp 295–314

Hanelt P (2001) Mansfeld's encyclopedia of agricultural and horticultural crops. In: Hanelt P (ed)
Mansfeld's encyclopedia of agricultural and horticultural crops.

Huang S, Sirikhachornkit A, Su X, Faris J, Gill B, Haselkorn R, Gornicki P (2002) Genes
encoding plastid acetyl-CoA carboxylase and 3-phosphoglycerate kinase of the
Triticum/Aegilops complex and the evolutionary history of polyploid wheat. Proc Natl
Acad Sci USA 99:8133-8138. https://doi.org/10.1073/pnas.072223799

Huson DH, Bryant D (2006) Application of phylogenetic networks in evolutionary studies. Mol
Biol Evol 23:254-267. https://doi.org/10.1093/molbev/msj030

Hutchinson J, Miller TE, Jahier J, Shepherd KW (1982) Comparison of the chromosomes of
Triticum timopheevi with related wheats using the techniques of C-banding and *in situ*
hybridization. Theor Appl Genet 64:31-40. https://doi.org/10.1007/BF00303647

Iizumi T, Furuya J, Shen Z, Kim W, Okada M, Fujimori S, Hasegawa T, Nishimori M (2017)
Responses of crop yield growth to global temperature and socioeconomic changes. Sci Rep
7:7800. https://doi.org/10.1038/s41598-017-08214-4

Jaccard P (1908) Nouvelles recherches sur la distribution florale. Bull Soc Vaud Sci Nat 44:223-
270

Jakob SS, Rödder D, Engler JO, Shaaf S, Özkan H, Blattner FR, Kilian B (2014) Evolutionary
history of wild barley (*Hordeum vulgare* subsp. *spontaneum*) analyzed using multilocus
sequence data and paleodistribution modeling. Genome Biol Evol 6:685-702.
https://doi.org/10.1093/gbe/evu047

Jakubziner MM (1932) Contribution to the knowledge of wild wheat in Transcaucasia. Bull Appl
Bot, Gen Plant Breed 5:147-198 (in Russian)

Jakubziner MM (1933) The results of the VIR expedition in Transcaucasia. Plant Industry in
USSR, Series A 5-6:216 (in Russian)

Jakubziner MM (1959) New wheat species. In: Jenkins BP (ed) Proceedings of the 1st
1056   International Wheat Genetics Symposium, August 11-15 1959. Winnipeg Public Press,
1057   Winnipeg, Canada, pp 207-220

1058   Jiang J, Gill BS (1993) Sequential chromosome banding and *in situ* hybridization analysis.
1059   Genome 36:792-795. https://doi.org/10.1139/g93-104

1060   Jiang J, Gill BS (1994) Different species-specific chromosome translocations in *Triticum*
1061   *timopheevii* and *T. turgidum* support the diphyletic origin of polyploid wheats. Chrom Res
1062   2:59-64. https://doi.org/10.1007/BF01539455

1063   Johnson BL (1975) Identification of the apparent B-genome donor of wheat. Can J Genet Cytol
1064   17:21-39. https://doi.org/10.1139/g75-004

1065   Johnson BL, Hall O (1967) Report of a joint expedition throughTurkey to collect species of the
1066   Triticinae under the auspices ofthe University of California, U.S.A., Kungl. Skogs-och
1067   Lantbruksakademien of Sweden and the Food and Agriculture Organization of theUnited
1068   Nations. Wheat Inf Serv 25:19-29

1069   Johnson BL, Waines JG (1977) Use of wild wheat resources. California Agriculture
1070   September:8-9

1071   Jones GH, Valamoti S, Charles M (2000) Early crop diversity: A "new" glume wheat from
1072   northern Greece. Veget Hist Archaebot 9:133-146. https://doi.org/10.1007/BF01299798

1073   Jorjadze M, Berishvili T, Shatberashvili E (2014) The ancient wheats of Georgia and their
1074   traditional use in the southern part of the country. Emir J Food Agric 26:192-202.
1075   https://doi.org/10.9755/ejfa.v26i2.17522

1076   Kawahara T, Badaeva ED, Badaev NS, Gill BS (1996) Spontaneous translocations in *Triticum*
1077   *araraticum* Jakubz. . Wheat Inf Serv 83:7-14

1078   Kawahara T, Tanaka M (1977) Six chromosome types in *Triticum araraticum* Jakubz. different
1079   with reciprocal translocations. Jpn J Genet 52:261-267. https://doi.org/10.1266/jjg.52.261

1080   Kawahara T, Tanaka M (1981) Intraspecific differentiation in chromosome structure in the wild
1081   tetraploid wheats. Wheat Inf Serv 52:33

1082   Kenéz Á, Pető Á, Gyulai F (2014) Evidence of 'new glume wheat' from the Late Neolithic
1083   (Copper Age) of south-eastern Hungary (4th millennium cal. b.c.). Veget Hist Archaeobot
1084   23:551-566. https://doi.org/10.1007/s00334-013-0405-3

Kilian B, Dempewolf H, Guarino L, Werner P, Coyne C, Warburton ML (2021) Crop Science
special issue: Adapting agriculture to climate change: a walk on the wild side. Crop Sci
61:32-36. https://doi.org/10.1002/csc2.20418

Kilian B, Ozkan H, Deusch O, Effgen S (2007) Independent wheat B and G genome origins in
outcrossing *Aegilops* progenitor haplotypes. Mol Biol Evol 24.
https://doi.org/10.1093/molbev/msl151

Kimber G, Feldman M (1987) Wild wheat, an Introduction. College of Agriculture University of
Missouri, Columbia, Mo, U.S.A.

Knüpffer H (2009) Triticeae genetic resources in *ex situ* genebank collections. In: Muehlbauer
GJ, Feuillet C (eds) Genetics and Genomics of the Triticeae, First edn. Springer, New
York, pp 31-79

Kohler-Schneider M (2003) Contents of a storage pit from late Bronze Age Stillfried, Austria:
another record of the "new" glume wheat. Veget Hist Archaeobot 12:105–111.
https://doi.org/10.1007/s00334-003-0010-y

Komuro S, Endo R, Shikata K, Kato A (2013) Genomic and chromosomal distribution patterns
of various repeated DNA sequences in wheat revealed by a fluorescence in situ
hybridization procedure. Genome 56:131-137. https://doi.org/10.1139/gen-2013-0003

Konarev VG, Gavriliuk EP, Penyova TI, Kovnarev AV, Khakimova AG, Migushova EF (1976)
On the nature and origin of wheat genomes by the data of biochemistry and
immunochemistry of seed proteins. Agricult Biol 11:656-665 (in Russian)

Konovalov F, Goncharov N, Goryunova S, Shaturova A, Proshlyakova T, Kudryavtsev A (2010)
Molecular markers based on LTR retrotransposons BARE-1 and Jeli uncover different
strata of evolutionary relationships in diploid wheats. Mol Genet Genom 283:551-563.
https://doi.org/10.1007/s00438-010-0539-2

Kreuz A, Boenke N (2002) The presence of two-grained einkorn at the time of the Bandkeramik
culture. Veget Hist Archaeobot 11:233-240. https://doi.org/10.1007/s003340200026

Kroll H (2016) Die Pflanzenfunde von Feudvar. In: Kroll H, Reed R (eds) Feudvar III: Die
Archäobotanik. Wützberg University Press, Wützberg. pp 37-194

Lilienfeld F, Kihara H (1934) Genomanalyze bei *Triticum* und *Aegilops*. V. *Triticum timopheevi*
Zhuk. . Cytologia (Tokyo) 6:87-122. https://doi.org/10.1508/cytologia.6.87

Maan SS, Lucken KA (1972) Interacting male sterility-male fertility restoration systems for hybrid wheat research. Crop Sci 12:360-364. https://doi.org/10.2135/cropsci1972.0011183X001200030031x

Maccaferri M, Harris NS, Twardziok SO, Pasam RK, Gundlach H, Spannagl M, Ormanbekova D, Lux T, Prade VM, Milner SG, Himmelbach A, Mascher M, Bagnaresi P, Faccioli P, Cozzi P, Lauria M, Lazzari B, Stella A, Manconi A, Gnocchi M, Moscatelli M, Avni R, Deek J, Biyiklioglu S, Frascaroli E, Corneti S, Salvi S, Sonnante G, Desiderio F, Marè C, Crosatti C, Mica E, Özkan H, Kilian B, De Vita P, Marone D, Joukhadar R, Mazzucotelli E, Nigro D, Gadaleta A, Chao S, Faris JD, Melo ATO, Pumphrey M, Pecchioni N, Milanesi L, Wiebe K, Ens J, MacLachlan RP, Clarke JM, Sharpe AG, Koh CS, Liang KYH, Taylor GJ, Knox R, Budak H, Mastrangelo AM, Xu SS, Stein N, Hale I, Distelfeld A, Hayden MJ, Tuberosa R, Walkowiak S, Mayer KFX, Ceriotti A, Pozniak CJ, Cattivelli L (2019) Durum wheat genome highlights past domestication signatures and future improvement targets. Nature Genet 51:885-895. https://doi.org/10.1038/s41588-019-0381-3

Makushina EN (1938) New wheat species *Titicum armeniacum* (Jakubz.) sp n Proc USSR Acad Sci (Doklady Academii Nauk SSSR) 21:350-353 (in Russian)

Martynov SP, Dobrotvorskaya TV, Krupnov VA (2018) Analysis of the distribution of *Triticum timopheevii* Zhuk. genetic material in common wheat varieties (*Triticum aestivum* L.). Russ J Genet 54:166-175. https://doi.org/10.1134/S1022795418020126

McIntosh RA, Gyarfas J (1971) *Triticum timopheevii* as a source of resistance to wheat stem rust. Z Pflanzenzucht 66:240-248.

Menabde VL (1948) The Georgian wheats. Izdatelstvo Akademii Nauk Gruzinskoy SSR, Tbilisi (in Georgian)

Miko P, Megyeri M, Kovacs G. Characterization of *Triticum timopheevii* gene bank accessions to gain useful materials for organic wheat breeding. In: Veisz O (ed) Agrisafe Final Conference (March 21-23, 2011). Climate Change: Challenges and Opportunities in Agriculture. Agricultural Research Institute of the Hungarian Academy of Sciences, Budapest, Hungary, 2011. p 90-93

Mitrofanova O, Badaeva E, Salina EA (2016) *T. timopheevii* and *T. zhukovskii*, the bread wheat cousins and their contribution to T. aestivum improvement. In: Bonjean AP, Angus WJ, VanGinkel M (eds) The World Wheat Book: A History of Wheat Breeding (v3). Doc Lavoisier, France

Mori N, Kondo Y, Ishii T, Kawahara T, Valkoun J, Nakamura C (2009) Genetic diversity and origin of timopheevi wheat inferred by chloroplast DNA fingerprinting. Breed Sci 59:571-578. https://doi.org/10.1270/jsbbs.59.571

Mosulishvili M, Bedoshvili D, Maisaia I (2017) A consolidated list of Triticum species and varieties of Georgia to promote repatriation of local diversity from foreign genebanks. Ann Agrar Sci 15:61-70. https://doi.org/10.1016/j.aasci.2017.02.006

Nave M, Ta M, Raupp J, Tiwari VK., Ozkan H, Poland J, Hale I, Komatsuda T, Distelfeld A (2021) The independent domestication of Timopheev's wheat: Insights from haplotype analysis of the *Brittle rachis 1* (*BTR1-A*) gene. Gene 12, 338. https://doi.org/10.3390/genes12030338

Nazarova EA (2007) To the taxonomy and nomenclature of *Triticum araraticum* Jakubz. (Poaceae). Flora of Plant Resources of Armenia 16:32-34

Nesbitt M, Samuel D (1996) From staple crop to extinction? The archaeology and history of the hulled wheats In: Padulosi S, Hammer K, Heller J (eds) Hulled wheats. IPGRI, Castelvecchio Pascoli, Tuscany, Italy, pp 41-100

Nieto Feliner G, Álvarez I, Fuertes-Aguilar J, Heuertz M, Marques I, Moharrek F, Piñeiro R, Riina R, Rosselló JA, Soltis PS, Villa-Machío I (2017) Is homoploid hybrid speciation that rare? An empiricist's view. Heredity 118:513. https://doi.org/10.1038/hdy.2017.7

Noda K, Ge K (1989) Structural differentiation of chromosomes between *Triticum dicoccoides* Korn. and *T. araraticum* Jakubz., showing high meiotic pairing homology. Jpn J Genet 64:191-198. https://doi.org/10.1266/jjg.64.191

Oliveira HR, Jacocks L, Czajkowska BI, Kennedy SL, Brown TA (2020) Multiregional origins of the domesticated tetraploid wheats. PLOS ONE 15:e0227148. https://doi.org/10.1371/journal.pone.0227148

Özkan H, Willcox G, Graner A, Salamini F, Kilian B (2011) Geographic distribution and domestication of wild emmer wheat (*Triticum dicoccoides*). Genet Res Crop Evol 58:11-53. https://doi.org/10.1007/s10722-010-9581-5

Peakall R, Smouse PE (2012) GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research – an update. Bioinformatics 28:2537–2539

Pedersen C, Rasmussen SK, Linde-Laursen I (1996) Genome and chromosome identification in cultivated barley and related species of the *Triticeae* (Poaceae) by *in situ* hybridization with the GAA-satellite sequence. Genome 39:93-104. https://doi.org/10.1139/g96-013

Perego R (2017) Contribution to the development of the Bronze Age plant economy in the
surrounding of the Alps: an archaeobotanical case study of two Early and Middle Bronze
Age sites in northern Italy (Lake Garda region). University of Basel, Switzerland

Perrier X, Flori A, Bonnot F (2003) Data analysis methods. In: Hamon P, Seguin M, Perrier X,
Glaszmann JC (eds) Genetic diversity of cultivated tropical plants. Enfield, Science
Publishers, Montpellier, pp 43-76

Placido DF, Campbell MT, Folsom JJ, Cui X, Kruger GR, Baenziger PS, Walia H (2013)
Introgression of novel traits from a wild wheat relative improves drought adaptation in
wheat. Plant Physiol 161:1806. https://doi.org/10.1104/pp.113.214262

Poets A, Fang Z, Clegg M, Morrell P (2015) Barley landraces are characterized by
geographically   heterogeneous   genomic   origins.   Genome   Biol   16:173.
https://doi.org/10.1186/s13059-015-0712-3

Pont C, Leroy T, Seidel M, Tondelli A, Duchemin W, Armisen D, Lang D, Bustos-Korts D,
Goué N, Balfourier F, Molnár-Láng M, Lage J, Kilian B, Özkan H, Waite D, Dyer S,
Letellier T, Alaux M, Russell J, Keller B, van Eeuwijk F, Spannagl M, Mayer KFX,
Waugh R, Stein N, Cattivelli L, Haberer G, Charmet G, Salse J, Wheat, Barley Legacy for
Breeding Improvement c (2019) Tracing the ancestry of modern bread wheats. Nature
Genet 51:905-911. https://doi.org/10.1038/s41588-019-0393-z

Pourkheirandish M, Dai F, Sakuma S, Kanamori H, Distelfeld A, Willcox G, Kawahara T,
Matsumoto T, Kilian B, Komatsuda T (2018) On the origin of the non-brittle rachis trait of
domesticated   einkorn   wheat.   Front   Plant   Sci   8:2031.
https://doi.org/10.3389/fpls.2017.02031

Pourkheirandish M, Hensel G, Kilian B, Senthil N, Chen G, Sameri M, Azhaguvel P, Sakuma S,
Dhanagond S, Sharma R, Mascher M, Himmelbach A, Gottwald S, Nair SK, Tagiri A,
Yukuhiro F, Nagamura Y, Kanamori H, Matsumoto T, Willcox G, Middleton CP, Wicker
T, Walther A, Waugh R, Fincher GB, Stein N, Kumlehn J, Sato K, Komatsuda T (2015)
Evolution   of   the   grain   dispersal   system   in   barley.   Cell   162:527-539.
https://doi.org/10.1016/j.cell.2015.07.002

Qi L, Friebe B, Gill BS (2006) Complex genome rearrangements reveal evolutionary dynamics
of   pericentromeric   regions   in   the   Triticeae.   Genome   49:1628-1639.
https://doi.org/10.1139/G06-123

Quade J, Stiner MC, Copeland A, Clarke AE, Özbaşaran M (2018) Summary of carbon-14
dating of the cultural levels of Aşıklı Ho□yük. In: Özbaşaran M, Duru G, Stiner MC (eds)
The early settlement at Aşıklı Ho□yük: Essays in honor of Ufuk Esin. Ege Yayinlari,
Istanbul, pp 43–56

R Core Team (2016) R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Ray DK, Mueller ND, West PC, Foley JA (2013) Yield trends are insufficient to double global
crop   production   by   2050.   PLOS   ONE   8:e66428.
https://doi.org/10.1371/journal.pone.0066428

Ray DK, Ramankutty N, Mueller ND, West PC, Foley JA (2012) Recent patterns of crop yield
growth and stagnation. Nature Commun 3:1293. https://doi.org/10.1038/ncomms2296

Rodríguez S, Maestra B, Perera E, Díez M, Naranjo T (2000a) Pairing affinities of the B- and G-
genome chromosomes of polyploid wheats with those of *Aegilops speltoides*. Genome
43:814–819. https://doi.org/10.1139/gen-43-5-814

Rodríguez S, Perera E, Maestra B, Díez M, Naranjo T (2000b) Chromosome structure of
Triticum   timopheevii   relative   to   *T.   turgidum*.   Genome   43:923-930.
https://doi.org/10.1139/gen-43-6-923

Rottoli M, Pessina A (2007) Neolithic agriculture in Italy. In: Colledge S, Conolly J (eds) The
origin and spread of domestic plants in South- west Asia and Europe. Left Coast Press,
Walnut Creek, pp 141-154

Sachs L (1953) Chromosome behavior in species hybrids with *Triticum timopheevi*. Heredity
7:49-58. https://doi.org/10.1038/hdy.1953.4

Salina EA, Leonova IN, Efremova TT, Roder MS (2006) Wheat genome structure: translocations
during   the   course   of   polyploidization.   Funct   Integr   Gen   6:71-80.
https://doi.org/10.1007/s10142-005-0001-4

Salina EA, Numerova OM, Ozkan H, Feldman M (2004) Alterations in subtelomeric tandem
repeats   during   early   stages   of   allopolyploidy   in   wheat.   Genome   47:860-867.
https://doi.org/10.1139/g04-044

Salina EA, Pestsova EG, Adonina IG, Vershinin AV (1998) Identification of a new family of
tandem   repeats   in   Triticeae   genomes.   Euphytica   100:231-237.
https://doi.org/10.1023/A:1018360324242

Schiemann E (1934) Erwin Baur. Berichte der Deutschen Botanischen Gesellschaft 52:51-114

Shcherban AB, Schichkina AA, Salina EA (2016) The occurrence of spring forms in tetraploid
Timopheevi wheat is associated with variation in the first intron of the VRN-A1 gene.
BMC Plant Biol 16:107-118. https://doi.org/10.1186/s12870-016-0925-y

Soltis PS, Soltis DE (2009) The role of hybridization in plant speciation. Ann Rev Plant Biol
60:561-588. https://doi.org/10.1146/annurev.arplant.043008.092039

Stoletova EA (1924-25) Emmer (An assay of study of one of the vanishing crops) Bull Appl Bot
Plant Breed 14:27-111 (in Russian)

Svetozarova BB (1939) On the second genome of *Triticum timopheevii* Zhuk. Proc Acad Sci
USSR (Doklady Academii Nauk SSSR) 23:472-476 (in Russian)

Szabo AT, Hammer K (1996) Notes on the taxonomy of farro: *Triticum monococcum, T.*
dicoccon and *T. spelta*. In: Padulosi S, K. H, Heller J (eds) Hulled Wheat. IPGRI, Rome,
Italy, pp 2-40

Tanaka M, Ichikawa S (1972) Cytogenetical relationships of two types of *Trtiticum araraticum*
Jakubz. to other tetraploid wheat species. Jpn J Genet 47:103-114.
https://doi.org/10.1266/jjg.47.103

Tanaka M, Ishii H (1973) Cytogenetic evidence on the speciation of wild tetraploid wheats
collected in Iraq, Turkey, and Iran. In: Sears ER, Sears LMS (eds) 4[th] International Wheat
Genetics Symposium. Mo. Agr. Exp. Stn, Columbia, Mo, pp 115-121

Tanaka M, Kawahara T (1976) Wild tetraploid wheats from Northern Iraq cytogenetically
closely related to each other. Wheat Inf Serv 43:3

Tanaka M, Sakamoto S (1979) Morphological and physiological variations in wild tetraploid
wheats collected from the Zagros mountains. In: Tanaka M (ed) A preliminary report of
studies on the differentiation of tetraploid wheats from different biological levels. Report
of the Germ-plasm Institute, Kyoto, Japan, pp 12-17

Tanksley SD, McCouch SR (1997) Seed banks and molecular maps: Unlocking genetic potential
from the wild. Science (New York) 277:1063.
https://doi.org/10.1126/science.277.5329.1063

Toulemonde F, Durand F, Berrio L, Bonnaire E, Daoulas G, Wiethold J (2015) Records of
"new" glume wheat in France: a review. Veget Hist Archaeobot 24:197-206.
https://doi.org/10.1007/s00334-014-0479-6

Troitzky NA (1932) Botanical characteristic of the habitats of wild wheats in Armenia (Erevan).
Bull Appl Bot, Genet Plant Breed 5:53-68 (in Russian)

Tsunewaki K (1996) Plasmon analysis as the counterpart of genome analysis. In: Jauhar PP (ed)
Methods of Genome Analysis in Plant: Their Merrits and Piffals. CRC Press, Boca Ration,
pp 271-299

Tumanyan MG (1930) Wild eincorns and emmers in Armenia. Bull Appl Bot (Trudy po
Prikladnoi Botanike) 24:1-14 (in Russian)

Ulaş B, Fiorentino G (2020) Recent attestations of "new" glume wheat in Turkey: a reassessment
of its role in the reconstruction of Neolithic agriculture. Veget Hist Archaeobot.
https://doi.org/10.1007/s00334-020-00807-w

Valamoti SM, Kotsakis K (2007) Transitions to agriculture in the Aegean: the archaeobotanical
evidence. In: Colledge S, Conolly J (eds) The origins and spread of domestic plants in
Southwest Asia and Europe. Left Coast Press, Walnut Creek, pp 75-92

Valkoun J, Giles WJ, Konopka JP (1998) Part 6. Conservation of wild progenitors. Current
geographical distribution and habitat of wild wheats and barley. In: Damania AB, Valkoun
J, Willcox G, Qualset CO (eds) The Origins of Agriculture and Crop Domestication.
ICARDA, Aleppo, Syria,, pp 293-299

van Heerwaarden J, Doebley J, Briggs WH, Glaubitz JC, Goodman MM, de Jesus Sanchez
Gonzalez J, Ross-Ibarra J (2010) Genetic signals of origin, spread, and introgression in a
large   sample   of   maize   landraces.   Proc   Natl   Acad   Sci:201013011.
https://doi.org/10.1073/pnas.1013011108

Vavilov NI (1935) Botaniko-geograficheskie osnovy selektsii [Botanical-geographical
foundations of breeding]. Teoreticheskie osnovy selektsii rastenii [Theoretical foundations
of plant breeding], Moscow, Leningrad, pp 17-75 (in Russian)

Wagenaar EB (1961) Studies on the genome constitution of *Triticum timopheevi* Zhuk. I.
Evidence for genetic control of meiotic irregularities in tetraploid hybrids. Can J Genet
Cytol 3:47-60. https://doi.org/10.1139/g61-010

Würschum T, Leiser WL, Weissmann S, Maurer HP (2017) Genetic architecture of male fertility
restoration   of   *Triticum*   *timopheevii*.   Theor   Appl   Genet   130:1253-1266.
https://doi.org/10.1007/s00122-017-2885-5

Zabeau M, Vos P (1993) Selective restriction fragment amplification: a general method for DNA
fingerprinting. European patent application number 92402629.7; publication number
0534858 Al

Zaharieva M, Ayana N, Hakimi A, Misra S, Monneveux P (2010) Cultivated emmer wheat
(*Triticum dicoccon* Schrank), an old crop with promising future: a review. Genet Res Crop
Evol 57:937-962. https://doi.org/10.1007/s10722-010-9572-6

Zencirci N, Yılmaz H, Garaybayova N, Karagöz A, Kilian B, Özkan H, Hammer K, Knüpffer H
(2018) Mirza (Hacızade) Gökgöl (1897–1981): the great explorer of wheat genetic
resources in Turkey. Genet Res Crop Evol 65:693-711. https://doi.org/10.1007/s10722-
018-0606-9

Zhang H, Bian Y, Gou X, Dong Y, Rustgi S, Zhang B, Xu C, Li N, Qi B, Han F, von Wettstein
D, Liu B (2013) Intrinsic karyotype stability and gene copy number variations may have
laid the foundation for tetraploid wheat formation. Proc Natl Acad Sci 110:19466-19471.
https://doi.org/10.1073/pnas.1319598110

Zhang H, Mittal N, Leamy LJ, Barazani O, Song B-H (2017) Back into the wild—Apply
untapped genetic diversity of wild relatives for crop improvement. Evolut Appl 10:5-24.
https://doi.org/10.1111/eva.12434

Zhukovsky PM (1928) The new wheat species Bull Appl Bot (Trudy po Prikladnoi Botanike) 19:
59-66 (in Russian)

Zohary D, Hopf M, Weiss E (2012) Domestication of plants in the Old World, 4[th] edition.
Oxford Univ. Press, Oxford

Zoshchuk S, Badaeva E, Zoshchuk N, Adonina I, Shcherban' A, Salina E (2007) Intraspecific
divergence in wheats of the Timopheevii group as revealed by *in situ* hybridization with
tandem repeats of the Spelt1 and Spelt52 families. Russ J Genet 43: 636-645.
https://doi.org/10.1134/S1022795407060063

permanent, invariable C-bands

permanent C-bands, variable in size

inconsistent C-bands

► C-bands/ pAesp_SAT86 sites typical for ARA-1 group

hybridization sites of pAesp_SAT86

hybridization sites of pSc119.2

overlaping sites

► C-bands/ pAesp_SAT86 sites typical for ARA-0 group

🔵	*T. dicoccoides*
🟢	*T. araraticum*, ARA-0
🔴	*T. araraticum*, ARA-1
🔵	*T. timopheevii*