1 **BMC Biology**

3 **Research Article**

5 **Genetic diversity, distribution and domestication history of the neglected**

6 **GGAᵗAᵗ genepool of wheat[#]**

Ekaterina D. Badaeva[1,2,*], Fedor A. Konovalov[3,4], Helmut Knüpffer[4], Agostino Fricano[5], Alevtina S.

Ruban[4,6], Zakaria Kehel[7], Svyatoslav A. Zoshchuk[2], Sergei A. Surzhikov[2], Kerstin Neumann[4],

Andreas Graner[4], Karl Hammer[4], Anna Filatenko[4,8], Amy Bogaard[9], Glynis Jones[10], Hakan Özkan[11],

Benjamin Kilian[4,12]

1 – N.I. Vavilov Institute of General Genetics, Russian Academy of Sciences, Moscow, **Russia**
2 – Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow, **Russia**
3 – Independent Clinical Bioinformatics Laboratory, Moscow, **Russia**
4 – Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Gatersleben, **Germany**
5 – Council for Agricultural Research and Economics – Research Centre for Genomics &
Bioinformatics, Fiorenzuola d'Arda (PC), **Italy**
6 – KWS SAAT SE & Co. KGaA, Einbeck, **Germany**
7 – International Center for the Agricultural Research in the Dry Areas (ICARDA), Rabat, **Morocco**
8 – N.I. Vavilov Institute of Plant Genetic Resources (VIR), St. Petersburg, **Russia**
9 – School of Archaeology, Oxford, **United Kingdom**
10 – Department of Archaeology, University of Sheffield, **United Kingdom**
11 – Department of Field Crops, Faculty of Agriculture, University of Çukurova, Adana, **Turkey**
12 – Global Crop Diversity Trust, Bonn, **Germany**

*Correspondence: katerinabadaeva@gmail.com

Abstract

Background: Wheat yields are stagnating around the world and new sources of genes for resistance or tolerances to abiotic traits are required. In this context, the tetraploid wheat wild relatives are among the key candidates for wheat improvement. Despite of its potential huge value for wheat breeding, the tetraploid GGA^{t}A^t genepool is largely neglected. Understanding the population structure, native distribution range, intraspecific variation of the entire tetraploid GGA^{t}A^t genepool and its domestication history would further its use for wheat improvement.

Results: We report the first comprehensive survey of genomic and cytogenetic diversity sampling the full breadth and depth of the tetraploid GGA^{t}A^t genepool. We show that the extant GGA^{t}A^t genepool consists of three distinct lineages. We provide detailed insights into the cytogenetic composition of GGA^{t}A^t wheats, revealed group-, and population-specific markers and show that chromosomal rearrangements play an important role in intraspecific diversity of *T. araraticum*. We discuss the origin and domestication history of the GGA^{t}A^t lineages in the context of state-of-the-art archaeobotanical finds.

Conclusions: We shed new light on the complex evolutionary history of the GGA^{t}A^t wheat genepool. We provide the basis for an increased use of the GGA^{t}A^t wheat genepool for wheat improvement. The findings have implications for our understanding of the origins of agriculture in southwest Asia.

Keywords: *Triticum araraticum*, *Triticum timopheevii*, Crop Wild Relatives, Genetic diversity, Chromosomal passports, Domestication history, Fertile Crescent, Archaeobotany, New glume wheat

Background

The domestication of plants since the Neolithic Age resulted in the crops that feed the world today. However, successive rounds of selection during the history of domestication led to a reduction in genetic diversity, which now limits the ability of the crops to further evolve [Tanksley and McCouch 1997, van Heerwarden et al. 2011]. This is exacerbated by the demand for high crop productivity under climate change. Crop wild relatives (CWR) represent a large pool of beneficial

allelic variation and are urgently required to improve the elite genepools [Dempewolf et al. 2017;

Kilian et al. 2021]. Bread wheat and durum wheat are the staple crops for about 40% of the world's

population. But as wheat yields are stagnating around the world [Ray et al. 2012, Ray et al. 2013,

Lizumi et al. 2017]; new sources of genes for resistance or tolerances to abiotic traits such as

drought and heat are required. In this context, the wheat wild relatives are among the key sources for

bread wheat (*T. aestivum* L., 2n = 6x= 42, BBAADD) and durum wheat (*T. durum* Desf., 2n = 4x =

42, BBAA) improvement [Dante et al. 2013, Zhang et al. 2016].

However, in nature, no wild hexaploid wheat has ever been found. Only two wild tetraploid wheat

species (2n = 4x = 28) were discovered, namely, (i) emmer wheat *T. dicoccoides* (Körn. ex Asch. et

Graebn.) Körn. ex Schweinf., and (ii) Armenian, or Araratian emmer *T. araraticum* Jakubz. (syn. *T.*

timopheevii (Zhuk.) Zhuk. subsp. *armeniacum* (Jakubz.) van Slageren). Morphologically, both

species are very similar but differ in their genome constitution [Zohary et al. 2012]. *Triticum*

dicoccoides has the genome formula BBAA and *T. araraticum* has GGA^tA^t.

The wheat section Timopheevii mainly consists of wild tetraploid *Triticum araraticum* (GGA^tA^t),

domesticated tetraploid *T. timopheevii* (Zhuk.) Zhuk. (Timopheev's wheat, GGA^tA^t), and hexaploid

T. zhukovskyi Menabde et Ericzjan (2n = 6x= 42, $GGA^tA^tA^mA^m$) [Dorofeev et al. 1979, Goncharov

2012].

Wild *T. araraticum* was first collected by M.G. Tumanyan and A.G. Araratyan during 1925–28

southeast of Erevan, Armenia [Nazarova 2007], and thus after the discovery of domesticated *T.*

timopheevii by P.M. Zhukovsky in Georgia in 1922 [Zhukovsky 1928] (Additional file 1: Note 1).

Subsequently, *T. araraticum* was found in several other locations in Armenia and Azerbaijan

[Dorofeev et al. 1979, Jakubziner 1933, Jakubziner 1959], as well as in Iran, Iraq and Turkey. Single

herbarium specimens resembling *T. araraticum* have been sporadically recorded among *T.*

dicoccoides accessions collected from the Fertile Crescent [Jakubziner 1932, Sachs 1953]. However,

only botanical expeditions from the University of California at Riverside (USA) to Turkey in 1965,

to the Fertile Crescent in 1972–73 [Johnson and Hall 1967, Johnson and Waines 1977] and the

Botanical Expedition of Kyoto University to the Northern Highlands of Mesopotamia in 1970

[Tanaka and Ishii 1973, Tanaka and Kawahara 1976] significantly expanded our understanding of

the natural distribution of *T. araraticum*. More recently *T. araraticum* was found in north-western

Syria [Valkoun et al. 1998]. Especially in south-eastern Anatolia, Turkey, the distribution area of *T.*

araraticum overlaps with the distribution range of *T. dicoccoides*. From the western to eastern

Fertile Crescent, it is assumed that *T. araraticum* gradually substitutes *T. dicoccoides* [Johnson

1975], and *T. dicoccoides* is absent from Transcaucasia [Özkan et al. 2011]. In most habitats, *T.*
araraticum grows in patches and in mixed stands with other wild cereals [Tumanyan 1930, Troitzky
1932].

It is difficult to distinguish *T. araraticum* from *T. dicoccoides* by morphology under field conditions
[Dagan and Zohary 1970, Tanaka and Sakamoto 1979]. However, both species can easily be
differentiated based on biochemical, immunological, cytological and molecular markers [Badaeva et
al. 1994, Gill and Chen 1987, Jiang and Gill 1994, Kawahara and Tanaka 1977, Konarev et al. 1976,
Lilienfeld and Kihara 1934]. From an archaeobotanical perspective, both species can be reliably
identified based on several characteristics of charred spikelets [Jones et al. 2000]. In blind tests, it
was possible to distinguish modern representatives of the two species with a c. 90% accuracy, on the
basis of the primary keel of the glume, which arises just below the rachis disarticulation scar, and
the prominent vein on the secondary keel (observable at the base of the glume, which is the part of
spikelet most commonly preserved by charring in archaeological material) [Jones et al. 2000].

According to cytogenetic and molecular analyses, *T. araraticum* originated as a result of
hybridization between *Aegilops speltoides* Tausch (2n = 2x= 14, SS) and *Triticum urartu*
Thumanjan ex Gandilyan (2n = 2x= 14, AA) independently from *T. dicoccoides* [Dvořák et al. 1988,
Rodríguez et al. 2000a]. Similarity of the cytoplasmic genomes of *Ae. speltoides* and *T. araraticum*
indicated that *Ae. speltoides* was the maternal parent of *T. araraticum* [Tsunewaki 1996]. Hybrids
between *T. araraticum* × *T. dicoccoides* were reported as sterile because of meiotic disturbances and
gene interactions [Makushina 1938, Svetozarova 1939, Tanaka and Ishii 1973, Wagenaar 1961],
although a few authors [Noda and Ge 1989a, Sachs 1953, Tanaka and Kawahara 1976] reported
relatively good chromosome pairing in the F_1 hybrids in some *T. araraticum* × *T. dicoccoides*
combinations.

Triticum dicoccoides is considered to be the older species than *T. araraticum* [Gornicki et al. 2014,
Huang et al. 2002], which is supported by higher similarity of the S-G genomes compared to the S-B
genomes [Jiang and Gill 1993, Kilian et al. 2007, Rodríguez et al. 2000a]. Cytogenetic and
molecular data showed that the speciation (tetraploidization) of *T. araraticum* was accompanied by
complex species-specific translocations involving chromosomes 1G-6A^t-4G and 3A^t-4A^t [Chen and
Gill 1984, Jiang and Gill 1993, Maestra and Naranjo 1999, Salina et al. 2006] as well as with
mutations of the primary DNA structure causing the divergence of homoeologous chromosomes
(e.g., chromosomes 3A-3A^t) [Dobrovolskaya et al. 2009]. These changes of karyotype structure are

specific for the whole section Timopheevii as compared to the emmer wheat lineage (BBAA,

BBAADD) [Badaeva et al. 1986, Hutchinson et al. 1982, Zhang et al. 2013].

Intraspecific diversity of *T. araraticum* was detected by karyotype analysis [Badaeva et al. 1994,

Kawahara et al. 1996, Kawahara and Tanaka 1981]. Most recent phylogenetic studies based on

whole chloroplast genome sequences, genome-wide sequence information and enlarged taxon

sampling provided increased resolution of the evolutionary history within the Triticeae tribe, thereby

shedding also new light on the GGA^{t}A^t wheat genepool [Gornicki et al. 2014, Bernhardt et al. 2017].

Research and pre-breeding activities have focused on *T. dicoccoides* because it gave rise to the

economically most important wheats, *T. durum* Desf. and *T. aestivum* L. [Avni et al. 2017, El

Haddad et al. 2021]. However, *T. araraticum* and *T. timopheevii* have also contributed to bread

wheat improvement. Important genes were transferred such as genes controlling resistance against

stem rust, leaf rust, powdery mildew or wheat leaf blotch [Allard and Shands 1954, McIntosh and

Gyarfas 1971, Dyck 1992, Brown-Guedira et al. 1996, Brown-Guedira et al. 2003]. Cytoplasmic

male sterility (CMS) induced by *T. timopheevii* cytoplasm showed great potential for heterotic

hybrid technology [Maan and Lucken 1972, Mikó et al. 2011, Würschum et al. 2017]. However,

despite of its potential huge value for bread and durum wheat improvement, only a comparatively

small number of genes was transferred from *T. timopheevii* (even less from *T. araraticum*). Most of

the gene contributions originate from only one line (D-357-1) bred by R. Allard at the University of

Wisconsin in 1948 [Martynov et al. 2018].

Understanding the population structure and the intraspecific variation of the entire tetraploid

GGA^{t}A^t genepool would further its use for wheat improvement. In this study, we report the first

comprehensive survey of cytogenetic and genomic diversity sampling the full breadth and depth of

the tetraploid GGA^{t}A^t genepool. We provide new insights into the genetic relationships among

GGA^{t}A^t wheats and its domestication, taking into account the state-of-the-art archaeobotanical

finds.

Results

Genetic diversity and population structure of the GGA^{t}A^t genepool

Simultaneous amplification of multiple retrotransposon insertion sites using eight Long Terminal

Repeat (LTR) primer combinations generated 656 polymorphic Sequence-Specific Amplification

Polymorphism (SSAP) markers: 255 markers were obtained for *Jeli* insertions and 401 markers for

BARE-1 insertions. According to our previous study [Konovalov et al. 2010], *Jeli* targets mostly the

A genome, while *BARE-1* is distributed between A- and B/G-genome chromosomes. Altogether, 787 wheat genotypes representing 753 genebank accessions were considered for data analysis, after excluding apparently misidentified taxa and several cases of low-yield DNA extraction (Additional file 2: Table S1). Several major observations were made: (i) the extant GGA^tA^t genepool consists of three distinct lineages (two *T. araraticum* lineages and one of *T. timopheevii*, TIM). Surprisingly, wild *T. araraticum* consists of two major genetic lineages, preliminarily designated as 'ARA-0' and 'ARA-1' (Fig. 1; Additional file 3: Figs. S1, S2); (ii) while ARA-0 was found to be geographically widespread, ARA-1 was only found in south-eastern Turkey (Adiyaman, Kahramanmaraş, Gaziantep, Kilis) and in north-western Syria, where the distribution ranges of *T. araraticum* and of *T. dicoccoides* overlap (Fig. 2); (iii) as expected, *T. dicoccoides* is genetically more diverse than the GGA^tA^t genepool, supporting a more recent origin of *T. araraticum*; (iv) among the GGA^tA^t lineages, ARA-0 harbors most genetic diversity; (v) differences between A^t- and A^t+G genome diversity patterns were discovered; and (vi) Nei's genetic distance between lineages based on all 656 SSAP markers or considering only the *BARE-1* markers revealed that ARA-0 is phylogenetically more closely related to TIM than ARA-1. However, considering only the *Jeli* markers, ARA-1 was more closely related to TIM. *Triticum dicoccoides* (DIC) was genetically related most closely to the ARA-1 lineage (Additional file 4: Table S2).

A carefully selected subset of 103 genotypes was used for Amplified Fragment Length Polymorphism (AFLP) analysis (Additional file 2: Table S1). A screen using the six most promising AFLP primer combinations uncovered a total of 146 polymorphic markers across all genotypes. Major findings can be summarized as: (i) clustering of genotypes based on AFLP markers corroborates the existence of the three major and genetically distinct lineages of GGA^tA^t wheats (Additional file 3: Fig. S3a). One lineage comprised all *T. timopheevii* (and its derived mutant *T. militinae* Zhuk. et Migusch.) genotypes. Importantly, the two lineages of *T. araraticum* and also their distribution ranges were verified; (ii) the summary statistics highlight that, in contrast to the SSAP analysis, ARA-1 was more diverse than TIM for all parameters (Additional file 4: Table S3); (iii) ARA-1 was found to be genetically most closely related to TIM; (iv) ARA-1 was related most closely to DIC (Additional file 4: Table S3); (v) potential hybridization signals between ARA-0 and ARA-1 lineages were identified (Additional file 3: Fig. S3b, purple split); (vi) ARA-1 lines collected around Kilis, Kahramanmaraş and Gaziantep in Turkey were genetically closest to TIM (Additional file 3: Fig. S3c, light blue split in ARA-1); and (vii) shared splits between ARA-0 genotypes collected in Armenia and Azerbaijan with TIM (light blue split in Additional file 3: Fig.

S3b; and brown split in Additional file 3: Fig. S3c) indicate the potential contribution of ARA-0 to

the formation of TIM or probably hybridization of ARA-0 with TIM. The genetically closest ARA-0

genotype to TIM was collected from the present Ararat province of Armenia.

In total, 248 genotypes of *T. araraticum* and 17 of *T. timopheevii* collected across the entire

distribution range and representing all karyotypic variants (Additional file 2: Table S1) were

selected to infer the population structure based on chromosomal passports [Badaeva et al. 2015a].

The results based on 96 informative C-bands supported the molecular findings using SSAP and

AFLP markers and were congruent with the AFLP marker results as the closest ARA-1 genotypes to

TIM were collected near Gaziantep (61 km SE from Türkoğlu, SW of Karadağ), Turkey (Fig. 2;

Additional file 3: Fig. S4; Additional file 4: Table S4).

Fluorescence *in situ* hybridization (FISH) using six DNA probes was carried out for 95 genotypes,

which in turn corroborates the findings obtained using AFLP and C-banding markers (Additional

file 4: Table S5).

Intraspecific genetic diversity of T. araraticum based on karyotype structure and C-banding

patterns

Cytogenetic analysis of 391 *T. araraticum* genotypes representing 370 genebank accessions in

comparison with 17 *T. timopheevii* genotypes provided detailed insights into the genetic

composition of GGA^tA^t wheats. First, cytogenetic analysis highlighted significant differences

between wild emmer *T. dicoccoides* and *T. araraticum/T. timopheevii* in karyotype structure and C-

banding patterns (Fig. 3). Second, we revealed high diversity of the C-banding patterns and broad

translocation polymorphisms within *T. araraticum* (Fig. 3; Fig. 4; Additional file 3: Fig. S5–S12).

The most frequent karyotype variant shared by *T. timopheevii* and *T. araraticum* was defined as

'normal' (N) and it was found in 175 of 391 *T. araraticum* genotypes (44.6%) (more specifically:

155 of 342 ARA-0 = 45.32%; 20 of 49 ARA-1 = 40.81%; Additional file 2: Table S1). These 175

genotypes differed from each other only in the presence/absence or size of one to several C-bands.

The ratio of karyotypically normal genotypes decreased from 86.7% in Azerbaijan (excluding

Nakhichevan), to 60.0% in Turkey, 46.5% in Iraq, 30.6% in Armenia, 20.0% in Nakhichevan,

Azerbaijan, 16.7% in Syria, to 9.1% in Iran (Additional file 3: Fig. S5). Local populations differed

in the ratio of normal/rearranged genotypes. For example, some populations from Dahuk, Iraq,

possessed only karyotypically normal genotypes, while in others all genotypes possessed

chromosomal rearrangements. Similarly, in Turkey the frequency of karyotypically normal

genotypes varied from 100% (Mardin) to 0% (Kilis) (Additional file 2: Table S1; Additional file 4:
Table S6).

C-banding patterns of *T. araraticum* were highly polymorphic. Based on the presence or absence of
particular C-bands (Figs. 3, 5), all genotypes were divided into two groups. The first, larger group
(ARA-0) comprised of 342 genotypes (Additional file 3: Figs. S6–S11). The second group (ARA-1)
included 49 genotypes (Additional file 3: Fig. S10, *h1–h5; h6–h9;* Fig. S12). Interestingly,
genebank accession TA1900 presumably collected 32 km S of Denizli, Dulkadiroğlu district,
Kahramanmaraş province in the Taurus Mountain Range of Turkey in 1959 (Fig. 3c; Additional file
2: Table S1; Additional file 3: Fig. S11, *i31*) shared karyotypic features of *T. timopheevii* and ARA-
0, and, probably, it is a natural or artificially produced hybrid of *T. timopheevii* with an unknown
genotype of *T. araraticum.*

Six chromosomes carried diagnostic C-bands for the ARA-1 lineage at the following positions:
1A^{t}L3, 4A^{t}S7, 5A^{t}L3, 1GL5, 2GL7, 3GL7+L11 (Fig. 5; Additional file 3: Fig. S10, *h1–h5; h6–h9;*
Fig. S11). All these diagnostic C-bands were present in all ARA-1 genotypes, both with normal and
rearranged karyotypes. Some C-bands were also common for both ARA-0 and ARA-1 (Fig. 3), but
their size was larger in ARA-1 genotypes (2A^{t}L3, 2GS15).

Only few C-bands were characteristic for the ARA-0 lineage (Fig. 3, Fig. 5). Three C-bands
appeared with a frequency of over 95%, and two of them, 6A^{t}L3 and 5GL15, occurred only in the
ARA-0 group. The third C-band, 2GL13, was detected in 97% of ARA-0, but also in few ARA-1
genotypes.

'Region-specific' C-bands (in terms of highest frequency) [Badaeva et al. 1994] were detected for
ARA-0. For example, (i) the band 4A^{t}L7 dominated in Turkey and Transcaucasia (Additional file 3:
Figs. S7, S11); (ii) a medium to large 3A^{t}S7 band was frequently observed in Iran and
Sulaymaniyah (Iraq); (iii) one distinct 5GL13 band was frequently detected in genotypes from Erbil,
Iraq (Additional file 3: Figs. S9, S10). The specificity of C-banding patterns in genotypes
originating from the same geographic region was not only determined by single bands, but usually
by a particular combination of C-bands on several chromosomes. These region-specific banding
patterns were observed for both, normal and translocated forms. For example, the unique banding
pattern of chromosome 7G, lacking the marker C-band 7GS11, but carrying large bands for 7GS13,
7GS17 and 7GS21 on the short arm, and 7GL13 and 7GL15 on the long arm was common in
Transcaucasia (Additional file 3: Fig. S6). The chromosome 6G lacking telomeric C-bands on the
long arm was frequent in genotypes from Dahuk and Sulaymaniyah (both Iraq) (Additional file 3:

Figs. S7, *c16–c20*, S10, *f9–f15*), and also occurred in ARA-1 genotypes from Kahramanmaraş,

Turkey (Additional file 3: Fig. S12, *j2–j9*).

Chromosomal rearrangements play an important role in intraspecific diversity of T. araraticum

In total, 216 out of 391 (55.4%) *T. araraticum* accessions carried a translocated karyotype. Seventy-

six variants of chromosomal rearrangements including single and multiple translocations,

paracentric and pericentric inversions were identified (Fig. 4; Additional file 2: Table S1; Additional

file 4: Table S6). Novel rearrangements were represented by 44 variants, while 32 variants were

described earlier [Badaeva et al. 1994, Kawahara et al. 1996, Kawahara and Tanaka 1977, Kawahara

and Tanaka 1981]. One-hundred-forty-seven genotypes differed from the 'normal' karyotype by

one, 45 genotypes by two (double translocations), 21 genotypes by three (triple translocations) and

three genotypes – by four chromosomal rearrangements (quadruple translocations).

Altogether, we revealed 52 (33 novel) variants of single chromosomal rearrangements (Fig. 4;

Additional file 4: Table S6). They included paracentric (one variant) and pericentric inversions

(seven variants) and 44 single translocations involving A^t-A^t, A^t-G, or G-G-genome chromosomes.

Double rearrangements were represented by 16 independent and three cyclic translocations; among

them 10 were novel. Triple translocations were represented by three variants, two of which –

$T2A^t$:7G + $T6A^t$:5G:6G and T3G:7G:$7A^t$ + $T6A^t$:6G – were found here for the first time. Both

variants of quadruple translocations have been identified earlier in Transcaucasia [Badaeva et al.

1994, Badaeva et al. 1990].

A translocation between two chromosomes could give rise to different products depending on the

breakpoint position and arm combination in rearranged chromosomes. For example, a centromeric

translocation between 1G and 2G resulted in two translocation variants which were distinct in arm

combinations (S:S *vs.* S:L). Three translocation variants involving chromosomes 3G and 4G differed

from each other in arm combination and breakpoint position. To discriminate different translocation

variants involving same chromosomes, they were designated as T1G:2G-1 and T1G:2G-2, etc.

(Additional file 4: Table S6).

Most variants of chromosomal rearrangements were unique and identified in one or few genotypes,

and only four variants were relatively frequent. These were a triple translocation $T2A^t$:4G:7G +

$T4A^t$:$7A^t$, *per*Inv7A^t-1, T6G:7G, and T2G:4G:6G (21, 16, 12, and 13 genotypes, respectively).

Taken together, these four variants accounted for approximately 16% of the whole materials we

studied. Genotypes carrying the same rearrangement usually had similar C-banding patterns and

were collected from the same, or closely located geographic regions. For example, (i) all genotypes with $T2A^t{:}4G{:}7G + T4A^t{:}7A^t$ originated from Nakhichevan, Azerbaijan; (ii) $T6G{:}7G$ and $T2G{:}4G{:}6G$ were found in Erbil, Iraq; and (iii) *per*Inv$7A^t$-1 was collected in Iran and in the neighboring region of Sulaymaniyah, Iraq.

Other frequent translocations had a more restricted distribution and usually occurred in a single population. Only few genotypes carrying the same translocations were identified in spatially separated populations. These genotypes differed in their C-banding patterns, for example: (i) $T2G{:}4G$ was found in four genotypes from Erbil, Iraq and in two genotypes from Siirt, Turkey; (ii) $T1G{:}3G$ was identified in five ARA-0 genotypes from Dahuk, Iraq, one ARA-1 from Turkey and one ARA-1 genotype of a mixed accession IG 117895 collected in Syria; and (iii) $T1G{:}5G$ was identified not only in three cytogenetically distinct *T. araraticum* ARA-0 genotypes from Armenia and Azerbaijan (Additional file 3: Fig. S6, *b10*) and ARA-1 from Turkey (Additional file 3: Fig. S12, *k2*), but also in *T. timopheevii* [Badaeva et al. 2016].

Most populations consisted of both genotypes with 'normal' karyotype and genotypes with one to several variants of chromosomal rearrangements. However, their ratio and spectra differed between regions (Fig. 6; Additional file 1: Note 2; Additional file 3: Fig. S5).

Cytogenetic diversity of GGA^{t}A^t wheats assessed using FISH markers

To further investigate the intraspecific diversity of *T. araraticum* and to assess their phylogenetic relationships with *T. timopheevii*, we carried out FISH using six DNA probes. The probes pTa-535, pSc119.2 and GAA_n ensured chromosome identification [Badaeva et al. 2016] whereas Spelt-1, Spelt-52 and pAesp_SAT86 were used to estimate intra- and interspecific variation (Fig. 7).

The distribution of pTa-535 was monomorphic among *T. araraticum* and *T. timopheevii* (Fig. 7*a, b*). Only the pSc119.2 site located on $1A^tL$ discriminated ARA-1 and TIM from ARA-0 (Fig. 7*a*).

The variability of pAesp_SAT86 hybridization patterns was analyzed in four *T. timopheevii* and 26 *T. araraticum* genotypes, of them seven were from ARA-1 and 19 from ARA-0 lineages (Additional file 2: Table S1; Additional file 3: Fig. S13). Distribution of the pAesp_SAT86 probe in all *T. timopheevii* genotypes was similar except for chromosome $7A^t$ in genotype K-38555, which was modified due to a paracentric inversion or insertion of an unknown chromosomal fragment. Labeling patterns, however, were highly polymorphic for *T. araraticum* (Fig. 7; Additional file 3: Fig. S13). Large pAesp_SAT86 sites were found only on some G-genome chromosomes. The A^t-genome chromosomes possessed several small, but genetically informative polymorphic sites. Some of these sites were lineage-specific. Most obvious differences were observed for $3A^t$, 4G and 7G

chromosomes (Fig. 5). Thus, all TIM and ARA-1 genotypes carried the pAesp_SAT86 signal in the middle of 3A^{t}S, while for ARA-0 it was located sub-terminally on the long arm (Additional file 3: Fig. S13). One large pAesp_SAT86 cluster was present on the long arm of 4G in ARA-0, but on the short arm in ARA-1 and TIM. Two large and adjacent pAesp_SAT86 clusters were detected on 7GS in ARA-1 and TIM, but they were split between opposite chromosome arms in all ARA-0 genotypes. Differences between ARA-0 and ARA-1 in pAesp_SAT86 cluster position on 4G and 7G could be caused by pericentric inversions. Some other pAesp_SAT86 sites were identified in either one of the three groups. ARA-1 exhibited the largest polymorphism of pAesp_SAT86 labeling patterns among all groups (Additional file 3: Fig. S13).

FISH with Spelt-1 and Spelt-52 probes on chromosomes of 87 *T. araraticum* genotypes from different chromosomal groups and of different geographic origin, seven *T. timopheevii,* and one *T. zhukovskyi* genotypes revealed high intraspecific diversity of *T. araraticum* and low polymorphism in *T. timopheevii* (Additional file 3: Fig. S14). The broadest spectra of labeling patterns were found in genotypes from Dahuk and Sulaymaniyah (Iraq) and in the ARA-1 group from Turkey (Additional file 3: Fig. S14), while material from Transcaucasia exhibited the lowest variation. The polymorphism was due to variation in the number of Spelt-1 and Spelt-52 sites, their size and chromosomal location (Additional file 4: Table S7). The Spelt-1 signals in various combinations appeared in sub-telomeric regions of either one or both arms of 2A^t, 6A^t, and all G-genome chromosomes. The Spelt-52 signals were observed in various combinations on 2A^{t}S, 1GS, and 6GL chromosomes (Additional file 3: Fig. S14; Additional file 4: Tables S7–S8). Among them, two Spelt-1 sites (6GL and 7GL) and one Spelt-52 locus (2GS) were novel. The patterns of Spelt-1 and Spelt-52 repeats varied across geographic regions and between chromosomal groups (Additional file 1: Note 3; Additional file 4: Table S8).

Discussion

We present the most comprehensive survey of cytogenetic and genomic diversity of GGA^{t}A^t wheats. We describe the composition, distribution and characteristics of the GGA^{t}A^t genepool. Building on our results, the latest published complementary genomics studies and state-of-the-art archaeobotanical evidence we revisit the domestication history of the GGA^{t}A^t wheats. We arrived at the following four key findings:

1. The GGA^{t}A^t genepool consists of three distinct lineages

We sampled the full breadth and depth of GGA^{t}A^t wheat diversity and discovered a clear genetic

and geographic differentiation among extant GGA^{t}A^t wheats. Surprisingly, and supported by all

marker types, three clearly distinct lineages were identified. The first lineage is comprised of all *T.*

timopheevii genotypes (and the derived *T. militinae* and *T. zhukovskyi*; note that all *T. militinae* and

all *T. zhukovskyi* accessions maintained *ex situ* in genebanks are each derived from only one original

genotype). Interestingly, *T. araraticum* consists of two lineages that we preliminarily describe as

'ARA-0' and 'ARA-1'. This finding is in contrast to Kimber and Feldman [Kimber and Feldman

1987] who concluded that *T. araraticum* does not contain cryptic species, molecularly distinct from

those currently recognized. ARA-0 was found across the whole predicted area of species

distribution. ARA-1 was only detected in south-eastern Turkey and in neighboring north-western

Syria. It is interesting to note that only in this part of the Fertile Crescent, the two wild tetraploid

wheat species, *T. dicoccoides* and *T. araraticum*, grow in abundance in mixed stands.

Based on Figure 2, collecting gaps are evident for *T. araraticum*, and future collecting missions

should focus on four specific regions: (i) between the Euphrates river in the west and the Elazığ –

Silvan – Mardin transect region in the east. Interestingly, so far only *T. dicoccoides* was reported

from this region; (ii) between Bitlis (Turkey) – Amadiyah (Iraq) in the west and the Armenian

border in the east including north-western Iran; (iii) between Adıyaman – Silvan in the south and

Tunceli in the north; and (iv) between Hama in the south and west/north-west of Aleppo in Syria.

We found only two *T. araraticum* populations which contained representatives of both ARA

lineages: (i) 45 km south-east of Kahramanmaraş to Gaziantep, and (ii) 4 km north of St. Simeon on

the road to Afrin in Syria. However, in both cases, the ARA-1 lineage was significantly more

frequent than ARA-0. This suggests at least a certain level of taxon boundary between ARA-0 and

ARA1 lineages and should be investigated in the future.

Independent support for the existence of two wild *T. araraticum* lineages and their distribution

comes from Mori et al. [Mori et al. 2009] based on 13 polymorphic chloroplast microsatellite

markers (cpSSR). The 'plastogroup G-2' was distributed in south-eastern Turkey and northern Syria

and was closely related to *Triticum timopheevii* [Mori et al. 2009]. However, Gornicki et al.

[Gornicki et al. 2014], based on whole chloroplast genome sequence information and sufficient

taxon sampling (13 Triticeae species and 1127 accessions; 163 accessions in common with our

study), provided increased resolution of the chloroplast genome phylogeny and showed that the *T.*

timopheevii lineage possibly originated in northern Iraq (*and thus according to our data, belong to*

the ARA-0 lineage as no ARA-1 occurs in Iraq). This was supported by Bernhardt et al. [Bernhardt

et al. 2017], who, based on re-sequencing 194 individuals at the chloroplast locus *ndhF* (2232 bp) and on whole genome chloroplast sequences of 183 individuals representing 15 Triticeae genera, showed that some ARA-0 and TIM genotypes are most closely related. All GGA^{t}A^t wheats re-sequenced by Bernhardt et al. [Bernhardt et al. 2017] were considered in our study (3x ARA-0, 2x ARA-1, 1x TIM, 1x *T. militinae*) (Additional file 2: Table S1).

It is important to note that our results (i.e., the characteristics, composition and geographical distribution of ARA-0 and ARA-1 lineages) are not in agreement with the latest comprehensive taxonomical classification of wheat by Dorofeev et al. [Dorofeev et al. 1979], who divided *T. araraticum* into two subspecies: subsp. *kurdistanicum* Dorof. et Migusch. and subsp. *araraticum* (Additional file 2: Table S1). We propose to re-classify the GGA^{t}A^t genepool taxonomically in the future.

2. The karyotypic composition of GGA^{t}A^t wheats is as complex as the phylogenetic history of the GGA^{t}A^t genepool

Based on karyotype analyses, translocation spectra and distribution of DNA probes, *T. araraticum* populations from Dahuk and Sulaymaniyah (both Iraq) harbored the highest karyotypic diversity among all *T. araraticum* populations studied. We consider the region around Dahuk in Northern Iraq as the center of diversity of *T. araraticum*, and this is probably the region where *T. araraticum* originated. This is supported by Bernhardt et al. [Bernhardt et al. 2017] and Gornicki et al. [Gornicki et al. 2014] who traced chloroplast haplotypes from *Aegilops speltoides* growing in Iraq via *T. araraticum* (ARA-0) to *T. timopheevii* and *T. zhukovskyi*.

The karyotype 'similar' to those in 'normal' *T. timopheevii* was found in 44.6% of all *T. araraticum* genotypes. This is the group of candidates, in which the closest wild relative(s) to *T. timopheevii* is (are) expected. The frequency of the normal karyotype varied among countries and between populations. It is interesting to note that the Samaxi-Akhsu population in Azerbaijan and some populations near Dahuk (Iraq) possessed mostly karyotypically normal genotypes. Diagnostic C-bands for the ARA-1 lineage, both with normal and rearranged karyotypes, were 1A^{t}L3, 4A^{t}S7, 5A^{t}L3, 1GL5, 2GL7, 3GL7+L11. As expected, the number of C-bands characteristic for ARA-0 was smaller (due to the wide geographical distribution) and only two C-bands were lineage-specific and found in normal as well as translocated genotypes: 6A^{t}L3 and 5GL15.

Some FISH patterns suggested that *T. timopheevii* probably originated in Turkey and probably from ARA-1 (or, ARA-1 and TIM may have originated from a common ancestor, but then diverged). This

is supported by the following observations: (i) TIM and ARA-1 carry the pSc119.2 signal in the
middle of $1A^t$ long arm, while this site was absent from ARA-0; (ii) all ARA-0 and most ARA-1
possessed the Spelt-52 signal on 6GL, but it is absent in TIM and five ARA-1 genotypes from
Gaziantep-Kilis, Turkey. The distribution of Spelt-1 and Spelt-52 probes on chromosomes of these
five genotypes was similar to, and in accession IG 116165 (ARA-1 from Gaziantep) almost identical
with TIM; (iii) the pAesp_SAT86 patterns on chromosomes $3A^t$, 4G, and 7G are similar in TIM and
ARA-1 but differed from ARA-0. Differences between ARA-1 and TIM based on FISH patterns of
some other chromosomes as well as the results of C-banding and molecular analyses suggest that
extant ARA-1 genotypes are not the direct progenitors of TIM but that the ARA-1 lineage is most
closely related to it.

Based on AFLP, C-banding, FISH and *Jeli* retrotransposon markers, TIM was genetically most
closely related to ARA-1. Additional evidence for the close relationship between TIM and ARA-1
lineages comes from allelic variation at the *VRN-1* locus of genome A^t [Shcherban et al. 2016]. This
analysis revealed a 2.7 kb deletion in intron 1 of *VRN-A1* in three *T. timopheevii* and four *T.*
araraticum accessions, which, according to our data, belong to the ARA-1 lineage. However, at
Vrn-G1, TIM from Kastamonu in Turkey (PI 119442) shared the same haplotype (*Vrn1Ga*) with
ARA-1 samples, while TIM from Georgia harbored haplotype *VRN-G1* as found in ARA-0. These
results suggest multiple introgression events and incomplete lineage sorting as suggested by
Bernhardt et al. [Bernhardt et al. 2017] and Bernhardt et al. [Bernhardt et al. 2019].

Regular chromosome pairing observed in the F_1 hybrids of lines with 'normal' karyotypes
[Kawahara et al. 1996], identified in our study as ARA-0 × ARA-1 (Additional file 2: Table S1),
suggested that karyotypic differences between ARA-0 and ARA-1 lineages are not associated with
structural chromosomal rearrangements such as large translocations or inversions.

The emergence or loss of most lineage-specific Giemsa C-bands (Fig. 3) or FISH loci (Additional
file 3: Fig. S13, S14) could be due to heterochromatin re-pattering: amplification, elimination or
transposition of repetitive DNA sequences. Wide hybridization can also induce changes in C-
banding and FISH patterns of *T. araraticum* chromosomes. Changes in pAesp_SAT86 hybridization
patterns on 4G and 7G chromosomes, however, are likely to be caused by pericentric inversions,
which are also frequent in common wheat [Qi et al. 2006]. The role of inversions in inter- and
intraspecific divergence is probably underestimated. In our case, it seems possible that divergence
between ARA-1/TIM (two inversions) from ARA-0 (no inversion) was associated with at least two
pericentric inversions.

We did not find any genotype harboring both ARA-0 and ARA-1 specific FISH sites, although
ARA-0 and ARA-1 genotypes co-existed in two populations in Turkey and Syria. However, based
on FISH (Spelt-1 site on chromosome 6GL and 7GS, respectively), hybridization between certain
ARA-1 and ARA-0 lines can be predicted.

Iran occupies a marginal part of the distribution range of *T. araraticum*. An abundance of the
pericentric inversion of the $7A^t$ chromosome in the Iranian group indicates that it is derived from
Iraq. The karyotypically 'normal' genotype was probably introduced to Transcaucasia via Western
Azerbaijan (Iran). The low diversity of FISH patterns and the low C-banding polymorphism of *T.*
araraticum from Transcaucasia indicate that *T. araraticum* was introduced as a single event.
Interestingly, the AFLP data suggested some similarity between ARA-0 from Armenia and
Azerbaijan and *T. timopheevii*.

We hypothesize that homoploid hybrid speciation (HHS) [Abbott et al. 2010, Nieto Feliner et al.
2017, Soltis and Soltis 2009] and incomplete lineage sorting may be the possible mechanisms
explaining the origin of the ARA-1 lineage. ARA-1 grows in sympatry and in mixed populations
with *T. dicoccoides* and is phylogenetically most closely related to *T. dicoccoides*. ARA-1 could be
derived from the hybridization of *T. timopheevii* × *T. dicoccoides*; or alternatively, ARA-1 and TIM
could be derived from the hybridization ARA-0 × *T. dicoccoides*.

3. Does the *T. timopheevii* population found in western Georgia represent the last remnant of a widespread ancient cultivation area of GGA^tA^t wheats?

Wild emmer *T. dicoccoides* belongs to the first cereals to be domesticated by humans in the Fertile
Crescent and the evolution and domestication history of *T. dicoccoides* are relatively well studied
[Badaeva et al. 2015a, Civáň et al. 2013, Özkan et al. 2011]. Domesticated emmer *Triticum*
dicoccon Schrank was a staple crop of Neolithic agriculture, was widely cultivated for over 10,000
years and harbored impressive genetic diversity [Nesbitt and Samuel 1996, Szabó and Hammer
1996, Zaharieva et al. 2010]. The domestication of *T. dicoccoides* provided the key for durum wheat
[Maccaferri et al. 2019] and bread wheat evolution [Pont et al. 2019].

Much less is known about *T. timopheevii*. It is believed that *T. timopheevii* is the domesticated form
of *T. araraticum* [Jakubziner 1932, Dorofeev et al. 1979]. In contrast to *T. dicoccon*, *T. timopheevii*
was, since its discovery, considered as a '*monomorphous narrowly endemic species*' [Dekaprelevich
and Menabde 1932] cultivated in few villages of western Georgia [Stoletova 1924–25, Zhukovsky
1928; Additional file 1: Note 4]. Dekaprelevich and Menabde [Dekaprelevich and Menabde 1932]

noticed that the area of cultivation had probably been larger in the past. The last plants of *T.*
timopheevii in situ were found by the expedition of the N.I. Vavilov Institute of Plant Genetic
Resources (VIR, Russia) in 1983 near the village of Mekvena (Zskaltubo, Georgia) and deposited in
the VIR genebank under accession number K-56422 [E.V. Zuev, personal communication]. Today,
the widespread view is that the cultivation area of *T. timopheevii* was restricted to Georgia in the
(recent) past [Feldman 2001, Zohary et al. 2012].

However, hulled tetraploid wheat morphologically similar to *T. timopheevii* was identified at three
Neolithic sites and one Bronze Age site in northern Greece and described by Jones et al. [Jones et al.
2000] as a '**New**' Glume Wheat (NGW). The glume bases of these archaeological finds
morphologically resemble *T. timopheevii* more than any other extant domesticated wheat [Jones et
al. 2000]. After these finds of NGW in Greece, this wheat was also identified at Neolithic and
Bronze Age sites in Turkey, Bulgaria, Romania, Hungary, Slovakia, Austria, Italy, Poland, Germany
and France [Bieniek 2002, Bieniek 2007, Fairbairn et al. 2002, Kohler-Schneider 2003, Fischer and
Rösch 2004, Hajnalová 2007, Bogaard et al. 2007, Bogaard 2013, Valamoti and Kotsakis 2007,
Kreuz et al. 2005, Rotolli and Pessina 2007, Ulaş and Fiorentino in press]. Earlier finds of an
'unusual' glume wheat in Serbia [Borojevic 1991] and Turkey [de Moulins 1997] have subsequently
been recognized as NGW [Jones et al. 2000, Kroll 2016,]. Criteria were also established for
distinguishing the grains of NGW [Kohler-Schneider 2003].

At some sites, NGW appeared as a minor component and may have been part of the accompanying
weed flora of cereal fields [Kenéz et al. 2014, Ulaş and Fiorentino in press]. In other cases, it was
probably cultivated in a mix with einkorn [Jones et al. 2000, Kohler-Schneider 2003] and/or emmer.
The recovery of large quantities in storage deposits of whole spikelets at Çatalhöyük in
Turkey, caryopses and spikelet bases at the early Bronze Age settlement of Clermont-Ferrand in
France, and rich deposits including whole spikelets at bronze age sites in Italy, demonstrated that, at
least in some places, NGW was a major crop in itself [Bogaard et al. 2013, Kenéz et al. 2014
Toulemonde et al. 2015, Bogaard et al. 2017, Perego 2017, Ergun 2018].

Based on intensive archaeobotanical investigations at Çatalhöyük in central Anatolia, for
example, NGW was the predominant hulled wheat, overtaking emmer wheat around 6500 cal BC
and remaining dominant until the site's abandonment c. 5500 cal BC. The finds suggested that this
wheat was a distinct crop, processed, stored, and presumably grown, separately from other glume
wheats. NGW formed part of a diverse plant food assemblage at Neolithic Çatalhöyük,
including six cereals, five pulses and a range of fruits, nuts and other plants, which enabled this

early farming community to persist for 1500 years (c. 7100 to 5500 cal BC) [Bogaard et al. 2013,

Bogaard et al. 2017].

Recently, polymerase chain reactions specific for the wheat B and G genomes, and extraction

procedures optimized for retrieval of DNA fragments from heat-damaged charred material, have

been used to identify archaeological finds of NGW [Czajkowska et al. 2020]. DNA sequences from

the G genome were detected in two of these samples, the first comprising grain from the mid 7th

millennium BC at Çatalhöyük in Turkey, and the second made up of glume bases from the later 5th

millennium BC site of Miechowice 4 in Poland. These results provide evidence that NGW is indeed

a cultivated member of the GGA^{t}A^t genepool [Czajkowska et al. 2020]. As NGW is a recognized

wheat type across a broad geographic area in prehistory, dating back to the 9th millennium BC in

SW Asia, this indicates that *T. timopheevii* (*sensu lato, s.l. = domesticated GGA^{t}A^t wheat in*

general), was domesticated from *T. araraticum* during early agriculture, and was widely cultivated

in the prehistoric past [Czajkowska et al. 2020].

This raises the question of whether the few populations of *T. timopheevii* (*sensu stricto*, *s.str.*) found

in western Georgia were the last remnants of a wider GGA^{t}A^t wheat cultivation or whether the *T.*

timopheevii of Georgia was a local domestication independent of the domestication of *T. araraticum*

in SW Asia. To answer this question, sequence information for NGW, the Georgian *T. timopheevii*,

and the two lineages of *T. araraticum* (ARA-0, ARA-1) need to be compared.

Vavilov [Vavilov 1935] suggested that *T. timopheevii* of western Georgia was probably originally

introduced from north-eastern Turkey. Menabde and Ericzjan [Menabde and Ericzjan 1942 as cited

by Dorofeev et al. 1979] associated the origin of *T. timopheevii* with the region of the ancient

kingdom of Urartu, whence immigrant ancestors of modern-day Georgians introduced it into

western Georgia. Certainly, the possibility of introduction of Timofeev's wheat into Georgia from

the south should not be rejected [Dorofeev et al. 1979]. Is the domestication history of *T.*

timopheevii s.str. connected with other endemic wheats of Georgia, such as *T. karamyschevii* Nevski

(*T. georgicum* Dekapr. et Menabde or *T. paleocolchicum* Menabde) and *T. carthlicum* Nevski,

which were cultivated by Mingrelians in Western Georgia [Jorjadze et al. 2014, Mosulishvili et al.

2017]?

4. Was the cultivation range of *T. timopheevii* (*s.str.*) wider in the recent past?

We screened all available passport data and found three interesting cases, which could potentially

help to reconstruct the recent past cultivation range of *T. timopheevii s.str*: Interestingly, two *T.*

timopheevii accessions maintained in two *ex situ* genebanks are reported to originate from Turkey

(Additional file 2: Table S1) [https://www.genesys-pgr.org/]: (i) ATRI 3433 (TRI 3433) conserved

in the Federal *ex situ* Genebank of Germany hosted at the Institute of Plant Genetics and Crop Plant

Research, IPK, Gatersleben. This *T. timopheevii* line was most likely collected by E. Baur in Turkey

in 1926 [Schiemann 1934]; and (ii) PI 119442 identified among a barley sample obtained from a

market in Araç, near Kastamonu, Turkey (Fig. 2) in 1936 by Westover and Wellmann, and

maintained at the National Plant Germplasm System, USDA-ARS, USA. Both accessions harbor the

'normal' karyotype of *T. timopheevii*, both were characterized in our studies and confirmed as a

typical *T. timopheevii*. Additionally, the accession TA1900, presumably collected 32 km south of

Denizli near Kahramanmaraş in Turkey on the 14th of August 1959 and maintained in the wheat

germplasm collection of the Wheat Genetics Resource Center, Kansas State University, U.S.A., is

interesting because it shared karyotypic features of *T. timopheevii* and the ARA-0 lineage (Fig. 3;

Additional file 3: Fig. S11). However, we are not fully convinced that this line is a true natural

hybrid. Based on the passport data, this accession could potentially have escaped from an

experimental field or a breeding station, or received introgression(s) during *ex situ* maintenance

[Zencirci et al. 2018]. Assuming that the passport data is correct (Additional file 2: Table S1), we

could speculate that *T. timopheevii* may have been cultivated in Turkey during the first half of the

20th century. However, is this a realistic option?

We believe not. As reported by Stoletova 1924–1925, Dekaprelevich and Menabde 1929,

Dekaprelevich and Menabde 1932, Menabde 1948, Dekaprelevich 1954, *T. timopheevii s.str.* was

part of the spring landrace Zanduri (mixture of *T. timopheevii s.str.* and *T. monococcum*) and well

adapted to the historical provinces Lechkhumi and Racha of Georgia (Additional file 1: Note 4). The

Zanduri landrace was cultivated in the '*humid and moderately cool climate zone 400–800*' m above

the sea level [Dorofeev et al. 1979]. Martynov et al. (2018) reported that *T. timopheevii* potentially

has a '*low potential for plasticity*' and is not drought tolerant. Climate at origin based on bioclimatic

variables clearly differs between the regions of Western Georgia where the Zanduri landrace grew

till the recent past, and both Kastamonu and Kahramanmaraş regions in Turkey (Additional file 1:

Note 5). We speculate that the three *T. timopheevii* accessions which were collected in Turkey were

probably introduced from Transcaucasia or elsewhere and may have been left over from

unsuccessful cultivation or breeding experiments of *T. timopheevii s. str.* in the recent historical

past. Also, based on botanical records, *T. timopheevii (s.str.)* has only been identified in Georgia, but

not in Turkey or elsewhere [Davis 1965-1988, Hanelt 2001]. From this we conclude that the

cultivation range of *T. timopheevii* (*s str.*) was not wider in the recent past.

Conclusions

The evolutionary history of the $GG A^t A^t$ wheat genepool is complex. However, some pieces of the

puzzle are clearly recognizable: the region around Dahuk in northern Iraq can be considered the

center of origin, but also the center of diversity of *T. araraticum.*

The origin of *T. timopheevii s.str.* remains unclear, but we speculate that it was probably introduced

from Turkey, on the grounds that wild *T. araraticum* does not grow in Georgia and that *T.*

timopheevii s.str. is more closely related to *T. araraticum* from Turkey or northern Iraq than to the

Transcaucasian types.

Based on bioclimatic variables, we predict that *T. timopheevii s.str.* is maladapted to the climate

outside Western Georgia. If this speculation is correct, it suggests a sister-group relationship

between (i) the Georgian *T. timopheevii* (*s.str.*) and both *T. araraticum* lineages (ARA-0, ARA-1),

but also between (ii) *T. timopheevii s.str.* and the prehistoric SW Asian *T. timopheevii s.l.*

The oldest known records of prehistoric *T. timopheevii s.l.* are of Turkish origin: (i) A□ıklı Höyük

in Cappadocia, and (ii) Cafer Höyük (just inside the Fertile Crescent and potentially older than

A□ıklı Höyük but less thoroughly dated) [Cauvin et al. 2011; Quade et al. 2018], though more

prehistoric finds may be identified in the future. Emphasis for further archaeobotanical research

should be given to the central part of the Fertile Crescent, including northern Iraq because this is

probably the region where wild *T. araraticum* originated.

The unexpected discovery of the ARA-1 lineage is exciting and may have implications for our

understanding of the origins of agriculture in southwest Asia. Was ARA-1 involved in the formation

of *T. timopheevii s.l.* (= NGW)?

The distribution area of the ARA-1 lineage requires our attention. It is interesting to note that the

ARA-1 lineage ($GG A^t A^t$) and *T. dicoccoides* (BBAA) grow in sympatry and in mixed populations

only in a peculiar geographical area in the Northern Levant. This specific area is located between

Kahramanmaraş in the north, Gaziantep in the east, Aleppo in the south and the eastern foothills of

the Nur Dağlari mountains in the west. It is interesting that the closest extant wild relatives of

domesticated einkorn, barley and emmer were also collected here: (i) the beta race of wild einkorn

[Kilian et al. 2007] and the closest wild relatives to einkorn *btr1* type [Pourkheirandish et al. 2018];

(ii) the closest wild barley to *btr2* barley [Pourkheirandish et al. 2015] and to *btr1b* barley [Civán☐

and Brown 2017]. This specific region was among the areas predicted with high probability as

potential refugia for wild barley during the Last Glacial Maximum (LGM) by Jakob et al. [Jakob et

al. 2014]; and (iii) wild emmer subgroup II of the central-eastern race [Özkan et al. 2011]. On the

other hand, some genetic research has suggested that domesticated emmer wheat and barley

received substantial genetic input from other regions of the Fertile Crescent, resulting in hybridized

populations of different wild lineages indicating a mosaic of ancestry from multiple source

populations [Civán☐ et al. 2013, Poets 2015, Oliveira et al. 2020]. It will be interesting to see

whether further genetic and archaeobotanical research on *T. araraticum* lineages and *T. timopheevii*

s.l. can help to resolve this issue.

Finally, and fortunately, 1294 accessions of *T. araraticum* and 590 accessions of *T. timopheevii*

s.str. are stored in 24 and 37 *ex situ* genebank repositories, respectively [Knüpffer 2009]. Our study

provides the basis for a more efficient use of *T. araraticum* and *T. timopheevii* materials for crop

improvement.

Methods

Germplasm

A comprehensive germplasm collection of tetraploid wheats was established comprising 862

genebank accessions (1–5 genotypes per accession number): 450 *T. araraticum,* 88 *T. timopheevii,*

three *T. militinae* Zhuk. et Migusch., one *T. zhukovskyi,* 307 *T. dicoccoides,* eight *T. dicoccon,* and

five *T. durum.* Additionally, the following materials were included: (i) four samples of *T.*

araraticum collected by Dr. Nelli Hovhannisyan in Armenia; (ii) 17 samples of *T. araraticum* and

12 samples of *T. dicoccoides* collected by Dr. H. Özkan in Turkey; and (iii) 26 samples of *T.*

dicoccoides collected by Drs. E. Badaeva, O.M. Raskina and A. Belyayev in Israel. Altogether, 921

accessions of wild and domesticated tetraploid wheats were examined (Additional file 2: Table S1).

A subset of 787 tetraploid wheat genotypes representing 765 genebank accessions was examined

using Sequence-Specific Amplification Polymorphism (SSAP) markers. The subset included 360

genotypes of *T. araraticum*, 76 *T. timopheevii* (including two *T. militinae*), while 351 genotypes of

T. dicoccoides were considered as an outgroup. The whole collection was single-seed descended

(SSD) at least twice under field conditions (2009–2012) and taxonomically re-identified in the field

at IPK, Gatersleben in 2011 (Additional file 2: Table S1).

A subset of 103 genotypes, including 38 *T. araraticum*, two *T. militinae*, 13 *T. timopheevii*, 40 *T.*
dicoccoides, 10 *T. dicoccon,* and four *T. durum* was selected for a complementary analysis using
Amplified Fragment Length Polymorphism (AFLP) markers to infer the population structure of
GGA^{t}A^t wheats (Additional file 2: Table S1).

Karyotype diversity of 370 *T. araraticum* accessions was assessed by C-banding and Fluorescence
in situ hybridization (FISH) in comparison with 17 *T. timopheevii* and one *T. zhukovskyi* genotypes.
According to C-banding analysis, most *T. araraticum* accessions (353 of 370) were karyotypically
uniform and were treated as single genotype each. Seventeen accessions were heterogeneous and
consisted of two (13 accessions) or even three (four accessions) cytogenetically distinct genotypes,
which were treated as different entities (genotypes). Altogether 391 distinct genotypes of *T.*
araraticum were distinguished. In total, 243 genotypes were shared with SSAP and 33 with AFLP
analyses (Additional file 2: Table S1). In order to infer the population structure of GGA^{t}A^t wheats,
265 typical genotypes were selected representing all karyotypic variants.

Seven *T. timopheevii,* one *T. zhukovskyi,* and 87 *T. araraticum* genotypes representing all
chromosomal groups were selected for Fluorescence i*n situ* hybridization (FISH) analysis
(Additional file 2: Table S1).

Molecular analysis

SSAP analysis

Evolutionary relationships among wild and domesticated tetraploid wheat taxa were first inferred
based on polymorphic retrotransposon insertions. For this, the highly multiplex genome
fingerprinting method SSAP was implemented based on polymorphic insertions of retrotransposon
families *BARE-1* and *Jeli* spread across wheat chromosomes. DNA was isolated from freeze-dried
leaves of 787 SSD plants, using the Qiagen DNeasy Kit (Hilden, Germany). SSAP protocol was
based on Konovalov et al. [Konovalov et al. 2010] with further optimizations for capillary-based
fragment detection (Additional file 1: Note 6). In total, 656 polymorphic markers were generated for
BBAA- and GGA^{t}A^t-genome wheats by amplification of multiple retrotransposon insertion sites.
Data analysis was performed in SplitsTree 4.15.1 [Huson and Bryant 2006]. NeighborNet planar
graphs of Dice distances [Dice 1945] were constructed based on presence/absence of SSAP bands in
the samples.

AFLP analysis

The AFLP protocol, as described by Zabeau and Vos [Zabeau and Vos 1993], was performed with

minor modifications according to Altıntas̗ et al. [Altıntas̗ et al. 2008] and Alsaleh et al. [Alsaleh

et al. 2015]. In total, six AFLP primer combinations were used to screen the collection of 103 lines

(Additional file 1: Note 7). Neighbor-Joining (NJ) trees were computed based on Jaccard distances

[Jaccard 1908, Perrier et al. 2003]. NeighborNet planar graphs were generated based on Hamming

distances [Huson and Bryant 2006]. Genetic diversity parameters and genetic distances were

calculated using Genalex 6.5 [Peakall and Smouse 2012].

Cytogenetic analysis

C-banding

Chromosomal preparation and C-banding procedure followed the protocol published by [Badaeva et

al. 1994]. The A^t- and G-genome chromosomes were classified according to nomenclature proposed

by [Badaeva et al. 1991] except for chromosomes $3A^t$ and $4A^t$. Based on meiotic analysis of the F_1

T. timopheevii $\times$ *T. turgidum* hybrids [Rodríguez et al. 2000b] and considering karyotype structure

of the synthetic wheat *T.* $\times$ *soveticum* (Zhebrak) [Mitrofanova et al. 2016], the chromosomes $3A^t$

and $4A^t$ were exchanged. Population structure of *T. araraticum* and the phylogenetic relationship

with *T. timopheevii* were inferred based on chromosomal passports compiled for 265 genotypes with

high-quality C-banding patterns (247 *T. araraticum*, 17 *T. timopheevii*, one *T. zhukovskyi*).

Chromosomal passports were constructed by comparing the karyotype of the particular accession

with the generalized idiogram of A^t- and G-genome chromosomes (Fig. 5), as described for *T.*

dicoccon Schrank [Badaeva et al. 2015a] (Additional file 1: Note 8).

Fluorescence *in situ* hybridization (FISH)

Two polymorphic G-genome specific DNA probes, Spelt-1 [Salina et al. 1998] and Spelt-52 [Salina

et al. 2004], were used for all 95 (87 *T. araraticum,* seven *T. timopheevii* and one *T. zhukovskyi*)

genotypes considered for FISH analysis. Additionally, 26 of the 95 genotypes (Additional file 2:

Table S1) were analyzed using the probe pAesp_SAT86 [Badaeva et al. 2015b], which also showed

differences between the accessions. The probes pSc119.2 [Bedbrook et al. 1980], GAA_n [Pedersen

et al. 1996], and pTa-535 [Komuro et al. 2013] were subsequently hybridized to the same

chromosomal spread to allow chromosome identification. Classification of pSc119.2- and pTa-535-

labeled chromosomes followed the nomenclature of [Badaeva et al. 2016, Jiang and Gill 1993].

Chromosomes hybridized with the GAA_{10} microsatellite sequence were classified according to

nomenclature suggested for C-banded chromosomes (Additional file 1: Note 8).

Legends to Figures

Fig. 1 Genetic relationships between GGA^{t}A^t and BBAA wheats. NeighborNet planar graph of Dice distances representing the diversity of 787 GGA^{t}A^t and BBAA tetraploid wheat genotypes based on 656 SSAP markers.

Fig. 2 Natural geographical distribution of wild tetraploid *T. araraticum* and *T. dicoccoides*. Green dots correspond to collection sites of ARA-0 accessions, pink dots to ARA-1, and dark blue dots to *T. dicoccoides* (DIC). The collection sites of *T. timopheevii* and *T. zhukovskyi* are shown with turquoise and yellow dots, respectively. Key excavation sites in Turkey where NGW was identified are indicated with red triangles.

Fig. 3 Comparison of the C-banding patterns of *T. dicoccoides* (DIC), *T. timopheevii* (TIM, *a–c*, normal karyotypes), *T. araraticum* ARA-1 (*d–f*) and ARA-0 (*g–t*). DIC (IG 117174, Gaziantep), *a* – KU-1818 (Georgia); *b* – PI 119442; *c* – TA1900; *d* – IG 116165; *e* – PI 654340; *f* – KU-1950; *g* – CItr 17677; *h* – KU-8917; *i* – KU-8909; *j* – KU-1933 (all from Turkey); *k* – CItr 17680 (Iran); *l* – PI 427381 (Erbil, Iraq); *m* – PI 538518*; n* – PI 427425 (Dahuk, Iraq); *o* – KU-8705; *p* – KU-8695 (Shaqlawa, Erbil, Iraq); *q* – KU-8451; *r* – KU-8774 (Sulaymaniyah, Iraq); *s* – TRI 11945 (Nakhchivan); *t* – KU-1901 (Armenia). 1–7 – homoeologous groups. C-bands typical for ARA-1 are indicated with blue arrows, for ARA-0 – with green arrows, and C-bands characteristic for *T. timopheevii* – with red arrows. Black arrows point to rearranged chromosomes in genotypes.

Fig. 4 Chromosomal rearrangements identified in *T. araraticum*. The number of translocation variant corresponds to the number of the respective variant in Additional file 4: Table S6. Novel variants are designated with black numbers, and already known variants by red numbers.

Fig. 5 Generalized idiogram and nomenclature of the A^t- and G-genome chromosomes. The C-banding pattern is shown on the left, the pSc119.2 (red) and pAesp_SAT86 (green) pattern on the right side of each chromosome. 1–7 – homoeologous groups; S – short arm, L – long arm. The numerals on the left-hand side designate putative positions of C-bands/FISH sites that can be detected on the chromosome arm; C-bands specific for the ARA-1 group are shown with pink numerals, C-bands specific for the ARA-0 group are indicated by green numerals. Red asterisks on the right-hand side indicate C-bands that were considered for the "chromosomal passport".

Fig. 6 Intraspecific divergence of *T. araraticum* and *T. timopheevii*. Combinations of chromosome arms in rearranged chromosomes are designated. Line colors mark the different groups: ARA-0 (green), ARA-1 (pink) and *T. timopheevii* (black). Solid arrows designate novel rearrangements; arrows with asterisk designate previously described rearrangements [Badaeva et al. 1990, Badaeva et al. 1994]. The numerals above/next to the arrows indicate the number of accessions carrying the respective translocation.

Fig. 7 Distribution of different families of tandem repeats on chromosomes of *T. timopheevii* and *T. araraticum*. *Triticum timopheevii,* KU-107 (*a*), and *T. araraticum,* CItr 17680, ARA-0 (*b*), KU-8944, ARA-0 (*c*), KU-1984B, ARA-1 (*d*), PI 427364, ARA-0 (*e*), and 2630, ARA-1 (*f*). The following probe combinations were used: *a, b* – pSc119.2 (green) + pTa-535 (red); *d, e* –

722 pAesp_SAT86 (red) + GAA$_n$ (green); *c, f* – Spelt-1 (red) + Spelt-52 (green). The position of
723 pSc119.2 site on 1A^t chromosome typical for *T. timopheevii* and ARA-1 is shown with an arrow (*a*).
724 Translocated chromosomes (*c, d*) are arrowed. Chromosomes are designated according to genetic
725 nomenclature; the A^t-genome chromosomes are designated with yellow numerals and the G-genome
726 chromosomes with white numerals. Scale bar, 10 μm.

Supplementary information

728 **Supplementary information** accompanies this paper at <mark>http xxx</mark>

Additional file 1:

Note 1. Regarding the first finds of *T. timopheevii*. **Note 2.** Region-specific chromosomal
rearrangements. **Note 3.** Additional findings of the FISH analysis. **Note 4.** Information extracted and
translated from Dorofeev et al. (1979), regarding the cultivation area and geographical distribution
of *Triticum timopheevii*, and the "Zanduri" wheat complex. Translation by H. Knüpffer and
colleagues , February 2020. **Note 5.** Was the cultivation range of *T. timopheevii* (*s.str.*) wider in the
recent past? **Note 6.** Details about the SSAP protocol and analysis. **Note 7.** Details about the AFLP
protocol and analysis. **Note 8.** Details about the C-banding and FISH protocols and analyses.

Additional file 2:

Table S1. List of materials investigated. Column 1: Species; taxonomically misclassified accessions
are shown in red cells; Column 2: Subspecies (when available); Column 3: Accession number;
Column 4: Seed/ DNA sources; Column 5: Other identifiers; Column 6: SSAP code; Column 7:
SSAP-NNet group; Column 8: AFLP-code; Column 9: AFLP-NNet group; Column 10: group of
GGA^{t}A^t wheat defined visually based on C-banding-analysis; Column 11: Translocation type -
codes of chromosomal rearrangements identified in T. araraticum; Column 12: PASSP -
"chromosome passport" codes; Column 13: FISH - combination of probes used in FISH analysis;
Column 14: Accession number considered by Kawahara et al. (1996): translocation type according
to Kawahara et al. (1996); Column 15: Accession number considered by Mori et al. (2009): Column
16: Haplotype of T. araraticum/ T. dicoccoides according to Mori et al. (2009); Column 17:
Accession number considered by Gornicki et al. (2014); Column 18: Country of origin; Column 19:
Collection site.

Additional file 3:

Fig. S1. NeighborNet phylogenetic networks constructed for BBAA and GGA^{t}A^t-genome wheat
genotypes based on Dice genetic distances and SSAP markers. **Fig. S2.** NeighborNet phylogenetic

networks constructed for GGA^{t}A^t-genome wheat genotypes only based on Dice genetic distances
and SSAP markers. **Fig. S3. (a)** NJ tree for 103 genotypes based on Jaccard distances; **(b)**
NeighborNet (NNet) planar graph based on Hamming distances for 59 genotypes. **(c)** NNet planar
graph based on Hamming distances for 52 genotypes. **Fig. S4.** Neighbor-Joining (NJ) tree for 265
genotypes of *T. araraticum* and *T. timopheevii* using 379 C-banding markers based on Jaccard
distances. **Fig. S5.** The ratios of chromosomal rearrangements among geographic populations of *T.*
araraticum. **Fig. S6.** Polymorphism of the C-banding patterns of *T. araraticum* from Transcaucasia.
Fig. S7. Polymorphism of the C-banding patterns of *T. araraticum* from Iraq (I). **Fig. S8.**
Polymorphism of the C-banding patterns of *T. araraticum* from Erbil, Iraq (II). **Fig. S9.**
Polymorphism of the C-banding patterns of *T. araraticum* from Sulaymaniyah, Iraq (III). **Fig. S10.**
Polymorphism of the C-banding patterns of *T. araraticum* from Iran (*g1–g18*) and Syria (*h1–h9*).
Fig. S11. Polymorphism of the C-banding patterns of *T. araraticum* from Turkey (ARA-0). **Fig.**
S12. Polymorphism of the C-banding patterns of *T. araraticum* from Turkey (ARA-1). **Fig. S13.**
Comparison of the distribution of pAesp_SAT86 probe on chromosomes of *T. dicoccoides* (DIC), *T.*
timopheevii TIM (T1–T4), *T. araraticum* ARA-1 (A1–A7) and ARA-0 (A8–A24). **Fig. S14.**
Distribution of Spelt-1 (red) and Spelt-52 (green) probes on chromosomes of *T. timopheevii* (T) and
T. araraticum representing ARA-1 (A1, A2, A4–A10) and ARA-0 (A3, A11–A29) groups.

Additional file 4:

Table S2. Summary statistics and genetic distances based on SSAP markers. **Table S3.** Summary
statistics and genetic distances based on AFLP markers. **Table S4.** Summary statistics and genetic
distances based on C-bands. **Table S5.** Summary statistics and genetic distances based on FISH
markers. **Table S6.** Variants of chromosomal rearrangements in *T. araraticum.* **Table S7.**
Distribution of Spelt-1 and Spelt-52 loci over chromosomes. **Table S8.** Distribution of Spelt-1 and
Spelt-52 loci in geographic/chromosomal groups in comparison with *T. timopheevii.*

List of abbreviations

AFLP, Amplified Fragment Length Polymorphism
ARA-0, *T. araraticum* lineage 0 (wild, widespread)
ARA-1, *T. araraticum* lineage 1 (wild, native only to south-eastern Turkey (Adiyaman,
Kahramanmaraş, Gaziantep, Kilis) and in north-western Syria)
DIC, *T. dicoccoides*
FISH, Fluorescence *in situ* hybridization
LTR, Long Terminal Repeat
NGW, new glume wheat
SSAP, Sequence-Specific Amplification Polymorphism
SSD, single seed descent
TIM, *T. timopheevii* (domesticated)

Acknowledgements

We thank the following for providing seeds, DNA and passport data: N. I. Vavilov Institute of Plant Genetic Resources (VIR, Russia), the Federal *ex situ* Genebank of Germany, Gatersleben (IPK, Germany), the International Center for Agricultural Research in the Dry Areas (ICARDA), the Max Planck Institute for Plant Breeding Research (MPIPZ, Germany), Kyoto University (National Bioresource Project, NBRP, Japan), the John Innes Centre (JIC, UK), Wheat Genetics Resource Centre, Kansas State University (WGRC, USA), Centre for Genetic Resources (WUR, CGN, Netherlands), the Australian Winter Cereal Collection Tamworth (AWCC Australia), the National Plant Germplasm System (USDA-ARS, USA), Eastern Cereal and Oilseed Research Centre, Agriculture and Agri-Food Canada (Canada), Botanical Institute Tbilisi (Georgia), Tbilisi Agricultural Institute (Georgia), Institute of Cytology and Genetics (Siberian Branch of the Russian Academy of Sciences, Russia), and the Russian State Agrarian University Moscow, Timiryazev Agricultural Academy (Russia). We also thank Drs. P.A. Gandilyan, M. Nazarova and I.G. Gukasyan (all Botanical Institute, Erevan, Armenia), Drs. I.D.O. Mustafaev and N.Kh. Aminov (both Institute of Genetic Resources of Azerbaijan, Baku, Azerbaijan) for providing materials from their collection missions. We thank the following for excellent technical assistance: Sigi Effgen (MPIPZ), Christiane Kehler, Birgit Dubsky, Heike Harms, Ute Krajewski, Marita Nix, Kerstin Wolf, Jürgen Marlow, Michael Grau, Peter Schreiber (all IPK), the 'Experimental Fields and Nurseries' and 'Genome Diversity' groups at IPK. We thank Dr. E.V. Zuev for providing us the data on collection sites of *T. araraticum* and *T. timopheevii* accessions from VIR collection. We are greatly indebted to Maarten Koornneef and George Willcox for discussions and support. We thank two anonymous reviewers, who helped us to improve an earlier version of this manuscript.

Authors' contributions

Ekaterina D. Badaeva conceived, designed and conducted the cytogenetic experiments and analyzed data, wrote the first manuscript version.

Fedor Konovalov conducted the SSAP experiments and analyzed the data.

Helmut Knüpffer verified genebank information concerning the origin of the material, translated texts from Russian into English, and contributed to the discussions and editing of the manuscript.

Agostino Fricano, supported the molecular data analysis.

Alevtina S. Ruban contributed to the FISH analysis.

Zakaria Kehel conducted the simulation of regional agro-ecological variation based on bioclimatic variables.

Svyatoslav A. Zoshchuk contributed to the work on mapping of Spelt-1 and Spelt-52 probes on *T. timopheevii* and *T. araraticum* genotypes.

Sergey A. Surzhikov designed and synthesized oligo-probes for FISH analyses.

Kerstin Neumann contributed to data analysis, supported the field trials and edited the manuscript.

Andreas Graner contributed to the discussions and editing of the manuscript.

Anna Filatenko phenotyped and taxonomically re-identified the collection, contributed to the
discussions and edited the manuscript.
Karl Hammer phenotyped and taxonomically re-identified the collection, contributed to the
discussions and edited the manuscript.
Amy Bogaard contributed to the discussions and editing of the manuscript.
Glynis Jones contributed to the discussions and editing of the manuscript.
Hakan Özkan conducted the AFLP experiments, analyzed the data and edited the manuscript.
Benjamin Kilian conceived the project, established the germplasm collection, analyzed data,
interpreted results, wrote and edited the manuscript.

Funding

This work was supported by the European Community's Seventh Framework Programme (FP7, 2007–2013) under the grant agreement n°FP7-613556, Whealbi project [http://www.whealbi.eu/project/], the German Research Foundation (DFG) (FKZ KI 1465/1-1 and FKZ KI 1465/5-1), the Federal Office for Agriculture and Food (BLE) of Germany (FKZ 01/12-13-KAD), the Leibniz Institute of Plant Genetics and Crop Research (IPK, Gatersleben, Germany), the Max Planck Institute for Plant Breeding Research (MPIPZ, Cologne, Germany) and the Crop Wild Relatives Project (*Adapting Agriculture to Climate Change: Collecting, Protecting and Preparing Crop Wild Relatives*) which is supported by the Government of Norway and managed by the Global Crop Diversity Trust [https://www.cwrdiversity.org/]. EDB acknowledges support from the Russian State Foundation for Basic Research (projects 20-04-00284, 17-04-00087, 14-04-00247), State Budget Project and the Program "Dynamics and Preservation of Gene Pools" from the Presidium of the Russian Academy of Sciences.

Availability of data and materials

All data sets supporting the conclusions of this article are available in the additional files and from the corresponding author Ekaterina D. Badaeva (katerinabadaeva@gmail.com). The acquisition of plant material used in this study complies with institutional, national, and international guidelines.

Ethics approval

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

References

Abbott RJ, Hegarty MJ, Hiscock SJ, Brennan AC. Homoploid hybrid speciation in action. Taxon.
2010; doi: 10.1002/tax.595005

Allard RW, Shands RG (1954) Inheritance of resistance to stem rust and powdery mildew in
cytologically stable spring wheats derived from *Triticum timopheevii*. Phytopathology 44:266-
274

Alsaleh A, Baloch FS, Derya M, Azrak M, Kilian, B, Nachit MM, Özkan H. Genetic linkage map of
Anatolian durum wheat derived from a cross of Kunduru-1149 x Cham1. Plant Molecular
Biology Reporter. 2015; 33 (2), 209-220.

Altıntas□ S, Toklu F, Kafkas S, Kilian B, Brandolini A, Özkan H. Estimating genetic diversity in
durum and bread wheat cultivars from Turkey using AFLP and SAMPL markers. Plant Breed.
2008; doi:10.1111/j.1439-0523.2007.01424.x

Avni R., Nave M., Barad O. et al. Wild emmer genome architecture and diversity elucidate wheat
evolution and domestication. Science. 2017; doi: 10.1126/science.aan0032.

Badaeva ED, Shkutina FM, Bogdevich IN, Badaev NS. Comparative study of *Triticum aestivum* and
T. timopheevii genomes using C-banding technique. Plant Syst Evol. 1986; doi:
10.1007/BF00990122

Badaeva ED, Boguslavsky RL, Badaev NS, Zelenin AV. Intraspecific chromosomal polymorphism
of *Triticum araraticum* (Poaceae) detected by C-banding technique. Plant Syst Evol. 1990;
doi: 10.1007/BF00935980.

Badaeva ED, Budashkina EB, Badaev NS, Kalinina NP, Shkutina FM. General features of
chromosome substitutions in *Triticum aestivum* x *T. timopheevii* hybrids. Theor Appl Genet.
1991; doi: 10.1007/BF00226218.

Badaeva ED, Badaev NS, Gill BS, Filatenko AA. Intraspecific karyotype divergence in *Triticum*
araraticum (Poaceae). Plant Syst Evol. 1994; doi: 10.1007/BF00985912.

Badaeva ED, Keilwagen J, Knüpffer H, Waßermann L, Dedkova OS, Mitrofanova OP, et al.
Chromosomal passports provide new insights into diffusion of emmer wheat. PLoS ONE.
2015a; doi: 10.1371/journal.pone.0128556

Badaeva ED, Amosova AV, Goncharov NP, Macas J, Ruban AS, Grechishnikova IV, Zoshchuk SA,
Houben A. A set of cytogenetic markers allows the precise identification of all A-genome

chromosomes in diploid and polyploid wheat. Cytogenet Genome Res. 2015b; doi:
10.1159/000433458.
Badaeva ED, Ruban AS, Zoshchuk SA, Surzhikov SA, Knüpffer H, Kilian B. Molecular cytogenetic
characterization of *Triticum timopheevii* chromosomes provides new insight on genome
evolution of *T. zhukovskyi*. Plant Syst Evol. 2016; doi: 10.1007/s00606-016-1309-3
Bedbrook RJ, Jones J, O'Dell M, Thompson RJ, Flavell RB. A molecular description of telomeric
heterochromatin in *Secale* species. Cell. 1980; doi: 10.1016/0092-8674(80)90529-2
Bernhardt N., Brassac J., Kilian B., Blattner FR. Dated tribe-wide whole chloroplast genome
phylogeny indicates recurrent hybridizations within Triticeae. BMC Evol Biol 17, 141 (2017).
https://doi.org/10.1186/s12862-017-0989-9
Bernhardt N., Brassac J., Dong X., Willing E. M., Poskar C. H., Kilian B., Blattner F. R. (2019)
Genome-wide sequence information reveals recurrent hybridization among diploid wheat wild
relatives. Plant J. doi: 10.1111/tpj.14641
Bieniek A., 2002. Archaeobotanical analysis of some early Neolithic settlements in the Kujawy
region, central Poland, with potential plant gathering activities emphasized. Veget. Hist.
Archaeobot. 11: 33–40.
Bieniek A., 2007. Neolithic plant husbandry in the Kujawy region of central Poland. In: Colledge,
S., Conolly, J. (Eds.), The Origin and Spread of Domestic Plants in Southwest Asia and
Europe. Left Coast Press, Walnut Creek, pp. 327–342.
Bogaard A, Bending J, Jones G, 2007. Archaeobotanical evidence for plant husbandry and use. In:
Whittle A, (ed) The Early Neolithic on the Great Hungarian Plain. Investigations of the Körös
culture site of Ecsegfalva 23, County Békés. Varia Archaeologica Hungarica 21:421–445.
Bogaard A, Charles M, Livarda A, Ergun M, Filipovic D and Jones G, 2013. The archaeobotany of
mid-later Neolithic occupation levels at Çatalhöyük. In I. Hodder (ed.), Humans and
Landscapes of Çatalhöyük: Reports from the 2000-2008 seasons. Los Angeles: Monographs of
the Cotsen Institute of Archaeology, University of California at Los Angeles, pp. 93-128.
Bogaard, A., Filipović, D., Fairbairn, A., Green, L., Stroud, E., Fuller, D. & Charles, M. 2017.
Agricultural innovation and resilience in a long-lived early farming community: the 1500-year
sequence at Neolithic-early Chalcolithic Çatalhöyük, central Anatolia. Anatolian Studies 67:
1-28
Borojevic, K., 1991. Emmer aus Feudvar. In: Hänsel, B., Medović, P. (Eds.), Vorbericht über die
jugoslawisch-deutschen Ausgrabungen in der Siedlung von Feudvar bei Mosorin (Gem. Titel,
Vojvodina) von 1986–1990: Bronzezeit–Vorrömische Eisenzeit. Verlag Philipp von Zabern,
Darmstadt, pp. 171–177.
Brown-Guedira GL, Gill BS, Bockus WW, Cox ST, Hatchett JH, Leath S, Peterson CJ, Thomas JB,
Zwer PK. Evaluation of a collection of wild timopheevii wheat for resistance to disease and
arthropod pests. Plant Disease, 1996; 80:928-933.
Brown-Guedira GL, Singh S, Fritz AK. Performance and mapping of leaf rust resistance transferred
to wheat from *Triticum timopheevii* subsp. *armeniacum*. Phytopathology, 2003; 93:784-789.
Chen PD, Gill BS. The origin of chromosome 4A, and genomes B and G of tetraploid wheats. In:
Sakamoto S (ed) Proceedings of the 6th International Wheat Genetics Symposium, November

28 – December 3 1983. Plant Germ-Plasm Inst., Kyoto, Japan: Kyoto University; 1984. p 39-48.

Civán, P., and Brown, T. A. (2017). A novel mutation conferring the nonbrittle phenotype of cultivated barley. New Phytol. 214, 468–472. doi: 10.1111/nph.14377

Civán, P., Ivanicová, Z., Brown, T.A., 2013. Reticulated origin of domesticated emmer wheat supports a dynamic model for the emergence of agriculture in the Fertile Crescent. PLoS ONE 8(11), e81955. doi: 10.1371/journal.pone.0081955

Cauvin, J., O. Aurenche, M.-C. Cauvin and N. Balkan-Atlı 2011. The Pre-Pottery Site of Cafer Höyük. The Neolithic in Turkey, vol 1. M. Özdogan, N. Başgelen and P. Kuniholm. Istanbul, Archaeology and Art Publications, pp. 1-40.

Czajkowska, B.I., Bogaard, A., Charles, M., Jones, G., Kohler-Schneider, M., Mueller-Bieniek, A, Brown, T.A. 2020. Ancient DNA typing indicates that the "new" glume wheat of early Eurasian agriculture is a cultivated member of the *Triticum timopheevii* group. J. Arch. Sci.

Dagan, J., and Zohary, D. 1970. Wild tetraploid wheats from West Iran cytogenetically identical with Israeli *T. dicoccoides*. Wheat Inform. Serv. 31: 15-17.

Dante F. Placido, Malachy T. Campbell, Jing J. Folsom, Xinping Cui, Greg R. Kruger, P. Stephen Baenziger, Harkamal Walia. 2013. Introgression of Novel Traits from a Wild Wheat Relative Improves Drought Adaptation in Wheat. Plant Physiology, 161 (4) 1806-1819; DOI: 10.1104/pp.113.214262

Davis PH, ed. (1965-1988) Flora of Turkey and the East Aegean Islands, 10 vols. Edinburgh: University Press.

Dekaprelevich LL, Menabde VL (1929) Regarding the investigation of crop plants in western Georgia. Zap. Nauchno-prikl. otdelov Tiflisskogo Bot. Sada 6:219–254 (in Russian with English summary)

Dekaprelevich LL, Menabde VL (1932). Hulled wheat in Western Georgia. Tr. prikl. bot. gen. i sel. 5,1: 1–46 (in Russian)

Dekaprelevich LL (1954) Species, varieties and cultivars of Georgian wheat. Tr. Inst. Polevodstva AN GSSR 8:3–61 (in Russian)

Dempewolf, H., Baute, G., Anderson, J., Kilian, B., Smith, C., Guarino, L. (2017) Past and future use of wild relatives in crop breeding. Crop Science 57: 1070-1082.

de Moulins, D., 1997. Agricultural changes at Euphrates and steppe sites in the mid-8[th] to the 6[th] millennium B.C. British Archaeological Reports International Series 683, Archaeopress, Oxford.

Dice LR. 1945. Measures of the amount of ecologic association between species. Ecology. 26:297–302.

Dobrovolskaya O, Sourdille P, Bernard M, Salina E (2009) Chromosome synteny of the A genome of two evolutionary wheat lines. Russ J Genet. 2009; doi: 10.1134/S1022795409110118.

Dorofeev VF, Filatenko AA, Migushova EF, Udachin RA, Jakubziner MM. Cultivated flora of the USSR. I. Wheat. Leningrad: Kolos; 1979.

Dvořák J, Mcguire PE, Cassidy B (1988) Apparent sources of the A genomes of wheats inferred from polymorphism in abundance and restriction fragment length of repeated nucleotide sequences. Genome 30:680-689

Dyck PL (1992) Transfer of a gene for stem rust resistance from *Triticum araraticum* to hexaploid
wheat. Genome 35:788-792

El Haddad, N., Kabbaj, H., Zaïm, M., El Hassouni, K., Tidiane Sall, A., Azouz, M., … Bassi, F.
(2021) Crop wild relatives in durum wheat breeding: Drift or thrift? *Crop Science* 61(1)
https://doi.org/10.1002/csc2.20223

Ergun, M. (2018). "Where the wild things are". Contextual insights into wild plant exploitation at
Aceramic Neolithic Aşıklı Höyük, Turkey Paléorient 44: 9-28.

Fairbairn A, Asouti E, Near J, Martinoli D. Macro-botanical evidence for plant use at Neolithic
Çatalhöyük south-central Anatolia, Turkey. Vegetation History and Archaeobotany. 2002: doi:
https://doi.org/10.1007/s003340200005

Feldman M. Origin of cultivated wheat. In: Bonjean AP, Angus WJ, editors. The world wheat book
A history of wheat breeding. Londres-Paris-New York: Tec & Doc/Intersept Ltd; 2001. p 3-
56.

Fiorentino G, Ulaş B (2010) A recent attestation of new glume wheat type: biometrical and
morphological studies of forks from the Neolithic levels in Mersin-Yumuktepe (Turkey). In:
Bittmann F (ed) 15th Conference of the International Workgroup for Pelaeoethnobotany,
Wilhelmshaven, Germany, May 31 –June 5, 2010. Terra Nostra, Wilhelmshaven, Germany, v.
2, p 183.

Fischer E, Rösch A (2004) Archäobotanische Untersuchungen. In: Schier W, Drasovean F (eds)
Vorbericht über die rumänisch-deutschen Prospektionen und Ausgrabungen in der befestigten
Tellsiedlung von Uivar, jud. Timiş, Rumänien (1998–2002). Prähist Z 79: 209–220 DOI:
https://doi.org/10.1515/prhz.2004.79.2.145

Gill BS, Chen PD. Role of cytoplasm-specific introgression in the evolution of the polyploid wheats.
Proc Natl Acad Sci USA. 1987; doi:10.1073/pnas.84.19.6800

Goncharov NP. Comparative genetics of wheats and their related species 2nd ed. Novosibirsk:
Academic Publishing Group @Geo@; 2012.

Gornicki P, Zhu H, Wang J, Challa GS, Zhang Z, Gill BS, Li W. The chloroplast view of the
evolution of polyploid wheat. New Phytologist. 2014; doi: 10.1111/nph.12931

Hajnalová M (2007) Early farming in Slovakia: an archaeobotanical perspective. In: Colledge S,
Conolly J (eds) The origins and spread of domestic plants in Southwest Asia and Europe. Left
Coast Press, Walnut Creek, pp 295–314.

Hanelt P, ed. (2001) Mansfeld's encyclopedia of agricultural and horticultural crops. Volumes 1-6
URL: http://mansfeld.ipk-gatersleben.de/pls/htmldb_pgrc/f?p=185:3:2422827336895397#

Huang S, Sirikhachornkit A, Su X, Faris J, Gill B, Haselkorn R, Gornicki P. Genes encoding plastid
acetyl-CoA carboxylase and 3-phosphoglycerate kinase of the *Triticum/Aegilops* complex and
the evolutionary history of polyploid wheat. Proc Natl Acad Sci USA. 2002; doi:
10.1073/pnas.072223799.

Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. Mol Biol Evol.
2006; doi: 10.1093/molbev/msj030

Hutchinson J, Miller TE, Jahier J, Shepherd KW. Comparison of the chromosomes of *Triticum*
timopheevii with related wheats using the techniques of C-banding and in situ hybridization.
Theor Appl Genet. 1982; doi: 10.1007/BF00303647.

Jaccard P. 1908. Nouvelles recherches sur la distribution florale. Bull Soc Vaud. Sci Nat. 44:223–
270.

Jakob S, Rödder D, Engler J, Shaaf S, Özkan H, Blattner F, Kilian B (2014) Evolutionary history of
wild barley (*Hordeum vulgare* subsp. *spontaneum*) analyzed using multilocus sequence data
and paleodistribution modeling. Genome biology and evolution 6. doi: 10.1093/gbe/evu047

Jakubziner MM. Contribution to the knowledge of wild wheat in Transcaucasia. Bull Appl Bot,
Genet Plant Breed. 1932; 5:147-198 (in Russian).

Jakubziner MM. The results of the VIR expedition in Transcaucasia. Plant Industry in USSR, Series
A. 1933; 5-6:216 (in Russian).

Jakubziner MM. New wheat species. In: Jenkins BP (ed) Proceedings of the 1st International Wheat
Genetics Symposium, August 11-15 1959. Winnipeg, Canada: Winnipeg Public Press; 1959. p
207-220.

Jiang J, Gill BS. Sequential chromosome banding and in situ hybridization analysis. Genome. 1993;
doi: 10.1139/g93-104.

Jiang J, Gill BS. Different species-specific chromosome translocations in *Triticum timopheevii* and
T. turgidum support the diphyletic origin of polyploid wheats. Chrom Res. 1994; doi:
10.1007/BF01539455.

Johnson BL. Identification of the apparent B-genome donor of wheat. Can J Genet Cytol. 1975; doi:
10.1139/g75-004.

Johnson BL, Hall O. Report of a joint expedition through Turkey to collect species of the Triticinae
under the auspices of the University of California, U.S.A., Kungl. Skogs- och
Lantbruksakademien of Sweden and the Food and Agriculture Organization of the United
Nations. Wheat Inf Serv. 1967;25:19-29.

Johnson BL, Waines JG. Use of wild wheat resources. California Agriculture. 1977. September:8-9

Jones GH, Valamoti S, Charles M. Early crop diversity: A "new" glume wheat from northern
Greece. Veget Hist Archaeobot. 2000; doi: 10.1007/BF01299798

Jorjadze M, Berishvili T, Shatberashvili E (2014) The ancient wheats of Georgia and their
traditional use in the southern part of the country. Emir. J. Food Agric. 26(2): 192-202, doi:
10.9755/ejfa.v26i2.17522

Kawahara T, Tanaka M. Six chromosome types in *Triticum araraticum* Jakubz. different with
reciprocal translocations. Jap J Genet. 1977; doi: 10.1266/jjg.52.261.

Kawahara T, Tanaka M. Intraspecific differentiation in chromosome structure in the wild tetraploid
wheats. Wheat Inf Serv. 1981;52:33.

Kawahara T, Badaeva ED, Badaev NS, Gill BS. Spontaneous translocations in *Triticum araraticum*
Jakubz. Wheat Inf Serv. 1996;83:7-14

Kenéz Á, Pető Á, Gyulai F (2014) Evidence of 'new glume wheat' from the Late Neolithic (Copper
Age) of south-eastern Hungary (4th millennium cal BC). Vegetation History and
Archaeobotany. 2014; doi: 10.1007/s00334-013-0405-3

Kilian B, Özkan H, Deusch O, Effgen S, Brandolini A, Kohl J, Martin W, Salamini F. Independent
wheat B and G genome origins in outcrossing *Aegilop*s progenitor haplotypes. Mol Biol Evol.
2007; doi: 10.1093/molbev/msl151.

Kilian B, Dempewolf H, Guarino L, Werber P, Coyne C, Warburton ML (2021) Adapting agriculture to climate change: A walk on the wild side. *Crop Science* 61(1); https://doi.org/10.1002/csc2.20418

Kimber G, Feldman M. Wild wheat, an introduction. Columbia, Mo, U.S.A.: College of Agriculture University of Missouri; 1987.

Knüpffer H. Triticeae genetic resources in ex situ genebank collections. In: Muehlbauer GJ, Feuillet C, editors. Genetics and Genomics of the Triticeae. 1st ed. Heidelberg, London, New York: Springer; 2009. p 31–79.

Kohler-Schneider M. 2003. Contents of a storage pit from late Bronze Age Stillfried, Austria: another record of the "new" glume wheat. Vegetation History and Archaeobotany. 12: 105–111.

Komuro, S., Endo, R., Shikata, K., and Kato, A. (2013). Genomic and chromosomal distribution patterns of various repeated DNA sequences in wheat revealed by a fluorescence in situ hybridization procedure. Genome 56, 131-137. doi: 10.1139/gen-2013-0003

Konarev VG, Gavriliuk EP, Penyova TI, Konarev AV, Khakimova AG, Migushova EF. On the nature and origin of wheat genomes by the data of biochemistry and immunochemistry of seed proteins. Agricultural Biology. 1976; 11:656-665 (in Russian).

Konovalov FA, Goncharov NP, Goryunova SV, Shaturova A, Proshlyakova T, Kudriavtsev AM. Molecular markers based on LTR retrotransposons *BARE-1* and *Jeli* uncover different strata of evolutionary relationships in diploid wheats. Mol Genet Genom. 2010; https://doi.org/10.1007/s00438-010-0539-2

Kreuz A, Marinova E, Schäfer E, Wiethold J (2005) A comparison of early Neolithic crop and weed assemblages from the Linear-bandkeramik and the Bulgarian Neolithic cultures: differences and similarities. Veget Hist Archaeobot 14:237–258.

Kroll, H., 2016. Die Pflanzenfunde von Feudvar. In: Kroll, H., Reed, R. (Eds.), Feudvar III: Die Archäobotanik. Wützberg University Press, Wützberg, pp. 37–194.

Lilienfeld F, Kihara H. Genomanalyse bei *Triticum* und *Aegilops*. V. *Triticum timopheevii* Zhuk. Cytologia. 1934; doi: 10.1508/cytologia.6.87

Lizumi, T., Furuya, J., Shen, Z. et al. Responses of crop yield growth to global temperature and socioeconomic changes. Sci Rep 7, 7800 (2017). https://doi.org/10.1038/s41598-017-08214-4

Maan SS, Lucken KA. Interacting male sterility-male fertility restoration systems for hybrid wheat research. Crop Sci. 1972; doi: 10.2135/cropsci1972.0011183X001200030031x

Maccaferri, M., Harris, N. S., Twardziok, S. O., Pasam, R. K. et al. (2019) Durum wheat genome highlights past domestication signatures and future improvement targets. Nat. Genet. 51: 885-895. https://doi.org/10.1038/s41588-019-0381-3

Maestra B, Naranjo T. Structural chromosome differentiation between *Triticum timopheevii* and *T. turgidum* and *T. aestivum*. Theor Appl Genet. 1999; doi: 10.1007/s001220051130.

Makushina EN. New wheat species *Triticum armeniacum* (Jakubz.) sp. n. Proc Acad Sci USSR. 1938; 21:350-353 (in Russian).

Martynov, S.P., Dobrotvorskaya, T.V. & Krupnov, V.A. Analysis of the distribution of *Triticum timopheevii* Zhuk. genetic material in common wheat varieties (*Triticum aestivum* L.). Russ J Genet 54, 166–175 (2018). https://doi.org/10.1134/S1022795418020126

Menabde VL (1948) The Georgian wheats (in Russian). Izdatelstvo Akademii Nauk Gruzinskoy SSR. Tbilisi

McIntosh RA, Gyarfas J. *Triticum timopheevii* as a source of resistance to wheat stem rust. Z Pflanzenzücht. 1971; 66: 240-248.

Mikó P, Megyeri M, Kovács G. Characterization of *Triticum timopheevii* gene bank accessions to gain useful materials for organic wheat breeding. In: Veisz O (ed) Agrisafe Final Conference (March 21-23, 2011). Climate Change: Challenges and Opportunities in Agriculture. Agricultural Research Institute of the Hungarian Academy of Sciences, Budapest, Hungary, 2011. p 90-93.

Mitrofanova, O., Badaeva, E., and Salina, E. (2016). *Triticum timopheevii, T. araraticum* and *T. zhukovskyi,* bread and durum wheat relatives carrying the G genome. *In* "The World Wheat Book" (A. Bonjean, W. Angus and J. M. van Ginkel, eds.), Vol. 3, pp. 1167-1228. Lavoisier Tec&Doc.

Mori N, Kondo Y, Ishii T, Kawahara T, Valkoun J, Nakamura C. Genetic diversity and origin of timopheevii wheat inferred by chloroplast DNA fingerprinting. Breed Sci. 2009; doi: 10.1270/jsbbs.59.571.

Mosulishvili M, Bedoshvili D, Maisaida I (2017) A consolidated list of *Triticum* species and varieties of Georgia to promote repatriation of local diversity from foreign genebanks. Annals of Agrarian Science 15, 61-70, http://dx.doi.org/10.1016/j.aasci.2017.02.006

Nazarova EA. On the taxonomy and nomenclature of *Triticum araraticum* Jakubz. (Poaceae). Flora, Vegetation, and Plant Resources of Armenia. 2007;16: 32-34 (in Russian).

Nesbitt M, Samuel D. From staple crop to extinction? The archaeology and history of the hulled wheats. In: Padulosi S, Hammer K, Heller J (eds) Hulled wheats. Tuscany, Italy: IPGRI; 1996. p 41-100.

Nieto Feliner G, Álvarez I, Fuertes-Aguilar J, Heuertz M, Marques I, Moharrek F, Piñeiro R, Riina R, Rosselló JA, Soltis PS, Villa-Machío I. Is homoploid hybrid speciation that rare? An empiricist's view. Heredity. 2017; doi: 10.1038/hdy.2017.7

Noda K, Ge K. Structural differentiation of chromosomes between *Triticum dicoccoides* Körn. and *T. araraticum* Jakubz., showing high meiotic pairing homology. Jap J Genet. 1989; doi: 10.1266/jjg.64.191.

Oliveira, H.R., Jacocks, L., Czajkowska, B.I., Kennedy, S.L. and Brown, T.A. (2020) Multiregional origins of the domesticated tetraploid wheats. PLoS ONE 15(1): e0227148.

Özkan H, Willcox G, Graner A, Salamini F, Kilian B. (2011) Geographic distribution and domestication of wild emmer wheat (*Triticum dicoccoides*). Genet Res Crop Evol. 2011; doi: 10.1007/s10722-010-9581-5.

Peakall R, Smouse PE. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research – an update. Bioinformatics. 2012; 28: 2537–2539.

Pedersen C, Rasmussen SK, Linde-Laursen I. Genome and chromosome identification in cultivated barley and related species of the Triticeae (Poaceae) by *in situ* hybridization with the GAA-satellite sequence. Genome. 1996; doi: 10.1139/g96-013.

Perego, R., 2017. Contribution to the development of the Bronze Age plant economy in the surrounding of the Alps: an archaeobotanical case study of two Early and Middle Bronze Age sites in northern Italy (Lake Garda region). Unpublished PhD Thesis, University of Basel.

Poets, A.M., Fang, Z., Clegg, M.T., Morrell, P.L., 2015. Barley landraces are characterized by geographically heterogeneous genomic origins. Genome Biol. 16: 173.

Pont C, Leroy T, Seidel M, Tondelli A, Duchemin W, Armisen D, Lang D, Bustos-Korts D, Goué N, Balfourier F, Molnár-Láng M, Lage J, Kilian B, Özkan H, Waite D, Dyer S, Letellier T, Alaux M, Russell J, Keller B, van Eeuwijk F, Spannagl M, Mayer KFX, Waugh R, Stein N, Cattivelli L, Haberer G, Charmet G, Salse J, Wheat, barley legacy for breeding improvement. Tracing the ancestry of modern bread wheats. Nature Genetics. 2019; doi: 10.1038/s41588-019-0393-z

Pourkheirandish, M., Hensel, G., Kilian, B., Senthil, N., Chen, G., Sameri, M., et al. (2015). Evolution of the grain dispersal system in barley. Cell 162, 527–539. doi: 10.1016/j.cell.2015.07.002

Pourkheirandish M, Dai F, Sakuma S, Kanamori H, Distelfeld A, Willcox G, Kawahara T, Matsumoto T, Kilian B and Komatsuda T (2018) On the Origin of the Non-brittle Rachis Trait of Domesticated Einkorn Wheat. *Front. Plant Sci.* 8:2031. doi: 10.3389/fpls.2017.02031

Quade, J., Stiner, M.C., Copeland, A., Clarke, A.E., Özbaşaran, M., 2018. Summary of carbon-14 dating of the cultural levels of Aşıklı Höyük. In: Özbaşaran, M., Duru, G., Stiner, M.C. (Eds.), The Early Settlement at Aşıklı Höyük: Essays in Honor of Ufuk Esin. Istanbul, Ege Yayinlari, pp. 43–56.

Qi L, Friebe B, Gill BS. Complex genome rearrangements reveal evolutionary dynamics of pericentromeric regions in the Triticeae. Genome. 2006; doi: 10.1139/G06-123.

Ray, D., Ramankutty, N., Mueller, N. et al. Recent patterns of crop yield growth and stagnation. Nat Commun 3, 1293 (2012). https://doi.org/10.1038/ncomms2296

Ray DK, Mueller ND, West PC, Foley JA (2013) Yield Trends Are Insufficient to Double Global Crop Production by 2050. PLoS ONE 8(6): e66428. https://doi.org/10.1371/journal.pone.0066428

Rodríguez S, Maestra B, Perera E, Díez M, Naranjo T. Pairing affinities of the B- and G-genome chromosomes of polyploid wheats with those of *Aegilops speltoides*. Genome. 2000a; doi: 10.1139/gen-43-5-814

Rodríguez S, Perera E, Maestra B, Díez M, Naranjo T. Chromosome structure of *Triticum timopheevii* relative to *T. turgidum*. Genome. 2000b; doi: 10.1139/gen-43-6-923.

Rottoli M, Pessina A (2007) Neolithic agriculture in Italy. In: Colledge S, Conolly J (eds) The origin and spread of domestic plants in South- west Asia and Europe. Left Coast Press, Walnut Creek, pp 141–154

Sachs L. Chromosome behavior in species hybrids with *Triticum timopheevii*. Heredity. 1953; doi: 10.1038/hdy.1953.4

Salina EA, Pestsova EG, Adonina IG, Vershinin AV. Identification of a new family of tandem repeats in Triticeae genomes. Euphytica. 1998; doi: 10.1023/A:1018360324242.

Salina EA, Numerova OM, Özkan H, Feldman M. Alterations in subtelomeric tandem repeats during early stages of allopolyploidy in wheat. Genome. 2004; doi: 10.1139/g04-044.

Salina EA, Leonova IN, Efremova TT, Röder MS. Wheat genome structure: translocations during the course of polyploidization. Funct Integr Genomics. 2006; doi: 10.1007/s10142-005-0001-4.

Schiemann E: *Erwin Baur*. In: *Berichte der Deutschen Botanischen Gesellschaft.* Bd. 52, 1934, 2. Generalversammlungs-Heft, ISSN 0011-9970, S. 51–114

Shcherban AB, Schichkina AA, Salina EA. The occurrence of spring forms in tetraploid Timopheevi wheat is associated with variation in the first intron of the *VRN-A1* gene. BMC Plant Biology. 2016; doi: 10.1186/s12870-016-0925-y

Soltis PS, Soltis DE. The role of hybridization in plant speciation. Ann Rev Plant Biol. 2009; doi:10.1146/annurev.arplant.043008.092039

Stoletova EA (1924–1925) Emmer, *Triticum dicoccum* Schrank – An assay of study of one of the vanishing crops. Tr. prikl. bot. sel. 14,1:27–111 (in Russian).

Svetozarova BB. On the second genome of *Triticum timopheevii* Zhuk. Proc Acad Sci of the USSR. 1939;23: 472-6 (in Russian).

Szabó AT, Hammer K (1996) Notes on the taxonomy of farro: *Triticum monococcum, T. dicoccon* and *T. spelta.* In: Padulosi S, K. H, Heller J (eds) Hulled Wheat. IPGRI, Rome, Italy, pp 2-40

Tanaka M, Ishii H. Cytogenetic evidence on the speciation of wild tetraploid wheats collected in Iraq, Turkey, and Iran. In: Sears ER, Sears LMS, editors. 4th International Wheat Genetics Symposium. Columbia, Mo, USA: Mo. Agr. Exp. Stn; 1973. p 115-21.

Tanaka M, Kawahara T (1976) Wild tetraploid wheats from Northern Iraq cytogenetically closely related to each other. Wheat Inf Serv 43:3

Tanaka M, Sakamoto S (1979) Morphological and physiological variations in wild tetraploid wheats collected from the Zagros mountains. In: Tanaka M (ed) A preliminary report of studies on the differentiation of tetraploid wheats from different biological levels (Report of the Germ-plasm Institute 4). Faculty of Agriculture, Kyoto University, Kyoto, 12–17.

Tanksley, S.D., and S.R. McCouch (1997) Seed banks and molecular maps: Unlocking genetic potential from the wild. Science 277:1063–1066. doi:10.1126/science.277.5329.1063

Troitzky NA. Botanical characteristic of the habitats of wild wheats in Armenia (Erevan). Bull Appl Bot, Genet Plant Breed. 1932; 5:53-68 (in Russian).

Tsunewaki K. Plasmon analysis as the counterpart of genome analysis. In: Jauhar PP, editor. Methods of Genome Analysis in Plant: Their Merits and Pitfalls. Boca Raton: CRC Press; 1996., p 271-99.

Tumanyan MG. Wild einkorns and emmers in Armenia. Bulletin of Applied Botany (Trudy po Prikladnoi Botanike). 1930; 24:1-14 (in Russian).

Toulemonde F, Durand F, Berrio L, Bonnaire E, Daoulas G, Wiethold J. Records of "new" glume wheat in France: a review. Veget Hist Archaeobot. 2015; doi: 10.1007/s00334-014-0479-6

Ulaş B and Fiorentino G (2020) Recent attestations of "new" glume wheat in Turkey: a reassessment of its role in the reconstruction of Neolithic agriculture. Vegetation History and Archaeobotany, https://doi.org/10.1007/s00334-020-00807-w

Valamoti SM and Kotsakis K 2007. Transitions to agriculture in the Aegean: the archaeobotanical evidence. In: Colledge S, Conolly J (eds) The origins and spread of domestic plants in Southwest Asia and Europe. Left Coast Press, Walnut Creek, pp 75–92.

Valkoun J, Giles WJ, Konopka JP. Part 6. Conservation of wild progenitors. Current geographical
distribution and habitat of wild wheats and barley. In: Damania AB, Valkoun J, Willcox G,
Qualset CO, editors. The Origins of Agriculture and Crop Domestication. Aleppo, Syria:
ICARDA; 1998. p 293-299.

van Heerwaarden, J., J. Doebley, W.H. Briggs, J.C. Glaubitz, M.M. Goodman, J. de Jesus Sanchez
Gonzalez, and J. Ross-Ibarra (2011) Genetic signals of origin, spread, and introgression in a
large sample of maize landraces. Proc. Natl. Acad. Sci. USA 108:1088–1092.
doi:10.1073/pnas.1013011108

Vavilov, N. I. 1935. Botaniko-geograficheskie osnovy selektsii [Botanical-geographical foundations
of breeding]. In: Teoreticheskie osnovy selektsii rastenii [Theoretical foundations of plant
breeding], t. 1, Moscow; Leningrad, pp. 17–75.

Wagenaar EB. Studies on the genome constitution of *Triticum timopheevii* Zhuk. I. Evidence for
genetic control of meiotic irregularities in tetraploid hybrids. Can J Genet Cytol. 1961; doi:
10.1139/g61-010.

Würschum T, Leiser WL, Weissmann S, Maurer HP. Genetic architecture of male fertility
restoration of *Triticum timopheevii*. Theor Appl Genet. 2017; doi: 10.1007/s00122-017-2885-5

Zabeau M, Vos P. Selective restriction fragment amplification: a general method for DNA
fingerprinting. European patent application number 92402629.7; publication number 0534858
Al; 1993.

Zaharieva M, Ayana N, Hakimi A, Misra S, Monneveux P. Cultivated emmer wheat (*Triticum
dicoccon* Schrank), an old crop with promising future: a review. Genet Res Crop Evol. 2010;
doi: 10.1007/s10722-010-9572-6.

Zencirci, N., Yılmaz, H., Garaybayova, N., Karagöz, A., Kilian, B., Özkan, H., Hammer, K.,
Knüpffer, H. (2018). Mirza Hacızade Gökgöl 1897–1981: The great explorer of wheat genetic
resources in Turkey. Genet Resour Crop Evol 65:693□711.

Zhang H, Bian Y, Gou X, Dong Y, Rustgi S, Zhang B, Xu C, Li N, Qi B, Han F, von Wettstein D,
Liu B. Intrinsic karyotype stability and gene copy number variations may have laid the
foundation for tetraploid wheat formation. Proc Natl Acad Sci U.S.A. 2013; doi:
.1073/pnas.1319598110.

Zhang, H., Mittal, N., Leamy, L. J., Barazani, O., and Song, B.-H. (2016). Back into the wild-Apply
untapped genetic diversity of wild relatives for crop improvement. Evolut. Appl. 10, 5–24.
doi: 10.1111/eva.12434

Zhukovsky PM. The new wheat species. Bulletin of Applied Botany (Trudy po Prikladnoi
Botanike). 1928; 19:59-66 (in Russian).

Zohary D, Hopf M, Weiss E. Domestication of plants in the Old World, 4th edition. Oxford: Oxford

●	*T. dicoccoides*
●	*T. araraticum*, ARA-0
●	*T. araraticum*, ARA-1
●	*T. timopheevii*

DIC	TIM	ARA-1	← Turkey →	← Iran	← Iraq →	Caucasus →	DIC	TIM	ARA-1	← Turkey →	← Iran	← Iraq →	← Caucasus →

1 — T1GL:5GS

2 — T2A'L:6A'L · T2A'L:7GS · T2GL:6GL

3

4 — T4A'S:7A'S · T5GL:1GS · inv4G · T4GS:2A'S

5

6 — T6A'S:2A'S · T6GS:2GS

7 — inv7A'-1 · inv7A'-2 · T7A'L:4A'L · T7GL:4GL

a b c d e f g h i j k l m n o p q r s t

A^t genome   G-genome

■ permanent, invariable C-bands	■ hybridization sites of pAesp_SAT86
▨ permanent C-bands, variable in size	■ hybridization sites of pSc119.2
▦ inconsistent C-bands	▥ overlaping sites
▶ C-bands/ pAesp_SAT86 sites typical for ARA-1 group	▶ C-bands/ pAesp_SAT86 sites typical for ARA-0 group