BMC Biology

Research Article

Genetic diversity, distribution and domestication history of the neglected GGA^{t}A^t genepool of wheat[#]

[#]We dedicate this article to our visionary colleagues Moshe Feldman and Francesco Salamini

Ekaterina D. Badaeva[1,2,*], Fedor A. Konovalov[3,4], Helmut Knüpffer[4], Agostino Fricano[5], Alevtina S. Ruban[4,6], Zakaria Kehel[7], Svyatoslav A. Zoshchuk[2], Sergei A. Surzhikov[2], Kerstin Neumann[4], Andreas Graner[4], Karl Hammer[4], Anna Filatenko[4,8], Amy Bogaard[9], Glynis Jones[10], Hakan Özkan[11], Benjamin Kilian[4,12]

– N.I. Vavilov Institute of General Genetics, Russian Academy of Sciences, Moscow, **Russia**
– Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow, **Russia**
– Independent Clinical Bioinformatics Laboratory, Moscow, **Russia**
– Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Gatersleben, **Germany**
– Council for Agricultural Research and Economics – Research Centre for Genomics & Bioinformatics, Fiorenzuola d'Arda (PC), **Italy**
– KWS SAAT SE & Co. KGaA, Einbeck, **Germany**
– International Center for the Agricultural Research in the Dry Areas (ICARDA), Rabat, **Morocco**
– Independent Researcher, St. Petersburg, **Russia**
– School of Archaeology, Oxford, **United Kingdom**
– Department of Archaeology, University of Sheffield, **United Kingdom**
– Department of Field Crops, Faculty of Agriculture, University of Çukurova, Adana, **Turkey**
– Global Crop Diversity Trust, Bonn, **Germany**

*Correspondence: katerinabadaeva@gmail.com

Abstract

Background: Wheat yields are stagnating around the world and new sources of genes for resistance or tolerances to abiotic traits are required. In this context, the tetraploid wheat wild relatives are among the key candidates for wheat improvement. Despite of its potential huge value for wheat breeding, the tetraploid GGA^{t}A^t genepool is largely neglected. Understanding the population structure, native distribution range, intraspecific variation of the entire tetraploid GGA^{t}A^t genepool and its domestication history would further its use for wheat improvement.

Results: We report the first comprehensive survey of genomic and cytogenetic diversity sampling the full breadth and depth of the tetraploid GGA^{t}A^t genepool. We show that the extant GGA^{t}A^t genepool consists of three distinct lineages. We provide detailed insights into the cytogenetic composition of GGA^{t}A^t wheats, revealed group-, and population-specific markers and show that chromosomal rearrangements play an important role in intraspecific diversity of *T. araraticum*. We discuss the origin and domestication history of the GGA^{t}A^t lineages in the context of state-of-the-art archaeobotanical finds.

Conclusions: We shed new light on the complex evolutionary history of the GGA^{t}A^t wheat genepool. We provide the basis for an increased use of the GGA^{t}A^t wheat genepool for wheat improvement. The findings have implications for our understanding of the origins of agriculture in southwest Asia.

Keywords: *Triticum araraticum*, *Triticum timopheevii*, Crop Wild Relatives, Genetic diversity, Chromosomal passports, Domestication history, Fertile Crescent, Archaeobotany, New glume wheat

Background

The domestication of plants since the Neolithic Age resulted in the crops that feed the world today. However, successive rounds of selection during the history of domestication led to a reduction in genetic diversity, which now limits the ability of the crops to further evolve [1, 2]. This is exacerbated by the demand for high crop productivity under climate change. Crop wild relatives (CWR) represent a large pool of beneficial allelic variation and are urgently required to improve the elite genepools [3, 4]. Bread wheat and durum wheat are the staple crops for about

40% of the world's population. But as wheat yields are stagnating around the world [5-7]; new sources of genes for resistance or tolerances to abiotic traits such as drought and heat are required. In this context, the wheat wild relatives are among the key sources for bread wheat (*T. aestivum* L., 2n = 6x= 42, BBAADD) and durum wheat (*T. durum* Desf., 2n = 4x = 42, BBAA) improvement [8, 9].

However, in nature, no wild hexaploid wheat has ever been found. Only two wild tetraploid wheat species (2n = 4x = 28) were discovered, namely, (i) emmer wheat *T. dicoccoides* (Körn. ex Asch. et Graebn.) Körn. ex Schweinf., and (ii) Armenian, or Araratian emmer *T. araraticum* Jakubz. (syn. *T. timopheevii* (Zhuk.) Zhuk. subsp. *armeniacum* (Jakubz.) van Slageren). Morphologically, both species are very similar but differ in their genome constitution [10] *Triticum dicoccoides* has the genome formula BBAA and *T. araraticum* has GGA^{t}A^t.

The wheat section Timopheevii mainly consists of wild tetraploid *Triticum araraticum* (GGA^{t}A^t), domesticated tetraploid *T. timopheevii* (Zhuk.) Zhuk. (Timopheev's wheat, GGA^{t}A^t), and hexaploid *T. zhukovskyi* Menabde et Ericzjan (2n = 6x= 42, GGA^{t}A^{t}A^{m}A^m) [11, 12].

Wild *T. araraticum* was first collected by M.G. Tumanyan and A.G. Araratyan during 1925–28 southeast of Erevan, Armenia [13], and thus after the discovery of domesticated *T. timopheevii* by P.M. Zhukovsky in Georgia in 1922 [14] (Additional file 1: Note 1). Subsequently, *T. araraticum* was found in several other locations in Armenia and Azerbaijan [11, 15-17], as well as in Iran, Iraq and Turkey. Single herbarium specimens resembling *T. araraticum* have been sporadically recorded among *T. dicoccoides* accessions collected from the Fertile Crescent [18, 19]. However, only botanical expeditions from the University of California at Riverside (USA) to Turkey in 1965, to the Fertile Crescent in 1972–73 [20, 21] and the Botanical Expedition of Kyoto University to the Northern Highlands of Mesopotamia in 1970 [22, 23] significantly expanded our understanding of the natural distribution of *T. araraticum*. More recently *T. araraticum* was found in north-western Syria [24]. Especially in south-eastern Anatolia, Turkey, the distribution area of *T. araraticum* overlaps with the distribution range of *T. dicoccoides*. From the western to eastern Fertile Crescent, it is assumed that *T. araraticum* gradually substitutes *T. dicoccoides* [20, 25], and *T. dicoccoides* is absent from Transcaucasia [26]. In most habitats, *T. araraticum* grows in patches and in mixed stands with other wild cereals [27, 28].

It is difficult to distinguish *T. araraticum* from *T. dicoccoides* by morphology under field conditions [29, 30]. However, both species can easily be differentiated based on biochemical, immunological, cytological and molecular markers [31-35]. From an archaeobotanical perspective, both species can be reliably identified based on several characteristics of charred spikelets [36]. In blind tests, it was possible to distinguish modern representatives of the two species with a c. 90% accuracy, on the basis of the primary keel of the glume, which arises just

below the rachis disarticulation scar, and the prominent vein on the secondary keel (observable at the base of the glume, which is the part of spikelet most commonly preserved by charring in archaeological material) [36].

According to cytogenetic and molecular analyses, *T. araraticum* originated as a result of hybridization between *Aegilops speltoides* Tausch (2n = 2x= 14, SS) and *Triticum urartu* Thumanjan ex Gandilyan (2n = 2x= 14, AA) independently from *T. dicoccoides* [37-39]. Similarity of the cytoplasmic genomes of *Ae. speltoides* and *T. araraticum* indicated that *Ae. speltoides* was the maternal parent of *T. araraticum* [40]. Hybrids between *T. araraticum* × *T. dicoccoides* were reported as sterile because of meiotic disturbances and gene interactions [22, 41-43], although a few authors [19, 23, 44] reported relatively good chromosome pairing in the F$_1$ hybrids in some *T. araraticum* × *T. dicoccoides* combinations.

Triticum dicoccoides is considered to be the older species than *T. araraticum* [45, 46], which is supported by higher similarity of the S-G genomes compared to the S-B genomes [38, 39, 47]. Cytogenetic and molecular data showed that the speciation (tetraploidization) of *T. araraticum* was accompanied by complex species-specific translocations involving chromosomes 1G-6A^{ت}t-4G and 3A^t-4A^t [33, 48-50] as well as with mutations of the primary DNA structure causing the divergence of homoeologous chromosomes (e.g., chromosomes 3A-3A^t) [51]. These changes of karyotype structure are specific for the whole section Timopheevii as compared to the emmer wheat lineage (BBAA, BBAADD) [31, 52, 53].

Intraspecific diversity of *T. araraticum* was detected by karyotype analysis [31, 34, 54, 55]. Most recent phylogenetic studies based on whole chloroplast genome sequences, genome-wide sequence information and enlarged taxon sampling provided increased resolution of the evolutionary history within the Triticeae tribe, thereby shedding also new light on the GG^{t}A^t wheat genepool [45, 56].

Research and pre-breeding activities have focused on *T. dicoccoides* because it gave rise to the economically most important wheats, *T. durum* Desf. and *T. aestivum* L. [57, 58]. However, *T. araraticum* and *T. timopheevii* have also contributed to bread wheat improvement. Important genes were transferred such as genes controlling resistance against stem rust, leaf rust, powdery mildew or wheat leaf blotch [59-63]. Cytoplasmic male sterility (CMS) induced by *T. timopheevii* cytoplasm showed great potential for heterotic hybrid technology [64-66]. However, despite of its potential huge value for bread and durum wheat improvement, only a comparatively small number of genes was transferred from *T. timopheevii* (even less from *T. araraticum*). Most of the gene contributions originate from only one line (D-357-1) bred by R. Allard at the University of Wisconsin in 1948 [67]   .

Understanding the population structure and the intraspecific variation of the entire tetraploid GGA^{t}A^t genepool would further its use for wheat improvement. In this study, we report the first comprehensive survey of cytogenetic and genomic diversity sampling the full breadth and depth of the tetraploid GGA^{t}A^t genepool. We provide new insights into the genetic relationships among GGA^{t}A^t wheats and its domestication, taking into account the state-of-the-art archaeobotanical finds.

Results

Genetic diversity and population structure of the GGA^{t}A^t genepool

Simultaneous amplification of multiple retrotransposon insertion sites using eight Long Terminal Repeat (LTR) primer combinations generated 656 polymorphic Sequence-Specific Amplification Polymorphism (SSAP) markers: 255 markers were obtained for *Jeli* insertions and 401 markers for *BARE-1* insertions. According to our previous study [68], *Jeli* targets mostly the A genome, while *BARE-1* is distributed between A- and B/G-genome chromosomes. Altogether, 786 wheat genotypes representing 753 genebank accessions were considered for data analysis, after excluding apparently misidentified taxa and several cases of low-yield DNA extraction (Additional file 2: Table S1). Several major observations were made: (i) the extant GGA^{t}A^t genepool consists of three distinct lineages (two *T. araraticum* lineages and one of *T. timopheevii*, TIM). Surprisingly, wild *T. araraticum* consists of two major genetic lineages, preliminarily designated as 'ARA-0' and 'ARA-1' (Fig. 1; Additional file 3: Figs. S1, S2); (ii) while ARA-0 was found to be geographically widespread, ARA-1 was only found in south-eastern Turkey (Adiyaman, Kahramanmaraş, Gaziantep, Kilis) and in north-western Syria, where the distribution ranges of *T. araraticum* and of *T. dicoccoides* overlap (Fig. 2); (iii) as expected, *T. dicoccoides* is genetically more diverse than the GGA^{t}A^t genepool, supporting a more recent origin of *T. araraticum*; (iv) among the GGA^{t}A^t lineages, ARA-0 harbors most genetic diversity; (v) differences between A^t- and A^t+G genome diversity patterns were discovered; and (vi) Nei's genetic distance between lineages based on all 656 SSAP markers or considering only the *BARE-1* markers revealed that ARA-0 is phylogenetically more closely related to TIM than ARA-1. However, considering only the *Jeli* markers, ARA-1 was more closely related to TIM. *Triticum dicoccoides* (DIC) was genetically related most closely to the ARA-1 lineage (Additional file 4: Table S2).

A carefully selected subset of 103 genotypes was used for Amplified Fragment Length Polymorphism (AFLP) analysis (Additional file 2: Table S1). A screen using the six most promising AFLP primer combinations uncovered a total of 146 polymorphic markers across all

genotypes. Major findings can be summarized as: (i) clustering of genotypes based on AFLP markers corroborates the existence of the three major and genetically distinct lineages of GGA^{t}A^t wheats (Additional file 3: Fig. S3a). One lineage comprised all *T. timopheevii* (and its derived mutant *T. militinae* Zhuk. et Migusch.) genotypes. Importantly, the two lineages of *T. araraticum* and also their distribution ranges were verified; (ii) the summary statistics highlight that, in contrast to the SSAP analysis, ARA-1 was more diverse than TIM for all parameters (Additional file 4: Table S3); (iii) ARA-1 was found to be genetically most closely related to TIM; (iv) ARA-1 was related most closely to DIC (Additional file 4: Table S3); (v) potential hybridization signals between ARA-0 and ARA-1 lineages were identified (Additional file 3: Fig. S3b, purple split); (vi) ARA-1 lines collected around Kilis, Kahramanmaraş and Gaziantep in Turkey were genetically closest to TIM (Additional file 3: Fig. S3c, light blue split in ARA-1); and (vii) shared splits between ARA-0 genotypes collected in Armenia and Azerbaijan with TIM (light blue split in Additional file 3: Fig. S3b; and brown split in Additional file 3: Fig. S3c) indicate the potential contribution of ARA-0 to the formation of TIM or probably hybridization of ARA-0 with TIM. The genetically closest ARA-0 genotype to TIM was collected from the present Ararat province of Armenia.

In total, 248 genotypes of *T. araraticum* and 17 of *T. timopheevii* collected across the entire distribution range and representing all karyotypic variants (Additional file 2: Table S1) were selected to infer the population structure based on chromosomal passports [69]

. The results based on 96 informative C-bands supported the molecular findings using SSAP and AFLP markers and were congruent with the AFLP marker results as the closest ARA-1 genotypes to TIM were collected near Gaziantep (61 km SE from Türkoğlu, SW of Karadağ), Turkey (Fig. 2; Additional file 3: Fig. S4; Additional file 4: Table S4).

Fluorescence *in situ* hybridization (FISH) using six DNA probes was carried out for 95 genotypes, which, in turn, corroborates the findings obtained using AFLP and C-banding markers (Additional file 4: Table S5).

Intraspecific genetic diversity of T. araraticum based on karyotype structure and C-banding patterns

Cytogenetic analysis of 391 *T. araraticum* genotypes representing 370 genebank accessions in comparison with 17 *T. timopheevii* genotypes provided detailed insights into the genetic composition of GGA^{t}A^t wheats. First, cytogenetic analysis highlighted significant differences between wild emmer *T. dicoccoides* and *T. araraticum/T. timopheevii* in karyotype structure and C-banding patterns (Fig. 3). Second, we revealed high diversity of the C-banding patterns and broad translocation polymorphisms within *T. araraticum* (Fig. 3; Fig. 4; Additional file 3: Fig.

S5–S12). The most frequent karyotype variant shared by *T. timopheevii* and *T. araraticum* was defined as 'normal' (N) and it was found in 175 of 391 *T. araraticum* genotypes (44.6%) (more specifically: 155 of 342 ARA-0 = 45.32%; 20 of 49 ARA-1 = 40.81%; Additional file 2: Table S1). These 175 genotypes differed from each other only in the presence/absence or size of one to several C-bands. The ratio of karyotypically normal genotypes decreased from 86.7% in Azerbaijan (excluding Nakhichevan), to 60.0% in Turkey, 46.5% in Iraq, 30.6% in Armenia, 20.0% in Nakhichevan, Azerbaijan, 16.7% in Syria, to 9.1% in Iran (Additional file 3: Fig. S5). Local populations differed in the ratio of normal/rearranged genotypes. For example, some populations from Dahuk, Iraq, possessed only karyotypically normal genotypes, while in others all genotypes possessed chromosomal rearrangements. Similarly, in Turkey the frequency of karyotypically normal genotypes varied from 100% (Mardin) to 0% (Kilis) (Additional file 2: Table S1; Additional file 4: Table S6).

C-banding patterns of *T. araraticum* were highly polymorphic. Based on the presence or absence of particular C-bands (Figs. 3, 5), all genotypes were divided into two groups. The first, larger group (ARA-0) comprised of 342 genotypes (Additional file 3: Figs. S6–S11). The second group (ARA-1) included 49 genotypes (Additional file 3: Fig. S10, *h1–h5; h6–h9;* Fig. S12). Interestingly, genebank accession TA1900 presumably collected 32 km S of Denizli, Dulkadiroğlu district, Kahramanmaraş province in the Taurus Mountain Range of Turkey in 1959 (Fig. 3c; Additional file 2: Table S1; Additional file 3: Fig. S11, *i31*) shared karyotypic features of *T. timopheevii* and ARA-0, and, probably, it is a natural or artificially produced hybrid of *T. timopheevii* with an unknown genotype of *T. araraticum.*

Six chromosomes carried diagnostic C-bands for the ARA-1 lineage at the following positions: 1A^{t}L3, 4A^{t}S7, 5A^{t}L3, 1GL5, 2GL7, 3GL7+L11 (Fig. 5; Additional file 3: Fig. S10, *h1–h5; h6–h9;* Fig. S11). All these diagnostic C-bands were present in all ARA-1 genotypes, both with normal and rearranged karyotypes. Some C-bands were also common for both ARA-0 and ARA-1 (Fig. 3), but their size was larger in ARA-1 genotypes (2A^{t}L3, 2GS15).

Only few C-bands were characteristic for the ARA-0 lineage (Fig. 3, Fig. 5). Three C-bands appeared with a frequency of over 95%, and two of them, 6A^{t}L3 and 5GL15, occurred only in the ARA-0 group. The third C-band, 2GL13, was detected in 97% of ARA-0, but also in few ARA-1 genotypes.

'Region-specific' C-bands (in terms of highest frequency) [31] were detected for ARA-0. For example, (i) the band 4A^{t}L7 dominated in Turkey and Transcaucasia (Additional file 3: Figs. S7, S11); (ii) a medium to large 3A^{t}S7 band was frequently observed in Iran and Sulaymaniyah (Iraq); (iii) one distinct 5GL13 band was frequently detected in genotypes from Erbil, Iraq (Additional file 3: Figs. S9, S10). The specificity of C-banding patterns in genotypes originating

from the same geographic region was not only determined by single bands, but usually by a particular combination of C-bands on several chromosomes. These region-specific banding patterns were observed for both, normal and translocated forms. For example, the unique banding pattern of chromosome 7G, lacking the marker C-band 7GS11, but carrying large bands for 7GS13, 7GS17 and 7GS21 on the short arm, and 7GL13 and 7GL15 on the long arm was common in Transcaucasia (Additional file 3: Fig. S6). The chromosome 6G lacking telomeric C-bands on the long arm was frequent in genotypes from Dahuk and Sulaymaniyah (both Iraq) (Additional file 3: Figs. S7, *c16–c20*, S10, *f9–f15*), and also occurred in ARA-1 genotypes from Kahramanmaraş, Turkey (Additional file 3: Fig. S12, *j2–j9*).

Chromosomal rearrangements play an important role in intraspecific diversity of T. araraticum

In total, 216 out of 391 (55.4%) *T. araraticum* accessions carried a translocated karyotype. Seventy-six variants of chromosomal rearrangements including single and multiple translocations, paracentric and pericentric inversions were identified (Fig. 4; Additional file 2: Table S1; Additional file 4: Table S6). Novel rearrangements were represented by 44 variants, while 32 variants were described earlier [31, 34, 54, 55, 70]. One-hundred-forty-seven genotypes differed from the 'normal' karyotype by one, 45 genotypes by two (double translocations), 21 genotypes by three (triple translocations) and three genotypes – by four chromosomal rearrangements (quadruple translocations).

Altogether, we revealed 52 (33 novel) variants of single chromosomal rearrangements (Fig. 4; Additional file 4: Table S6). They included paracentric (one variant) and pericentric inversions (seven variants) and 44 single translocations involving A^t-A^t, A^t-G, or G-G-genome chromosomes. Double rearrangements were represented by 16 independent and three cyclic translocations; among them 10 were novel. Triple translocations were represented by three variants, two of which – T2A^t:7G + T6A^t:5G:6G and T3G:7G:7A^t + T6A^t:6G – were found here for the first time. Both variants of quadruple translocations have been identified earlier in Transcaucasia [31, 54].

A translocation between two chromosomes could give rise to different products depending on the breakpoint position and arm combination in rearranged chromosomes. For example, a centromeric translocation between 1G and 2G resulted in two translocation variants which were distinct in arm combinations (S:S *vs.* S:L). Three translocation variants involving chromosomes 3G and 4G differed from each other in arm combination and breakpoint position. To discriminate

different translocation variants involving same chromosomes, they were designated as T1G:2G-1 and T1G:2G-2, etc. (Additional file 4: Table S6).

Most variants of chromosomal rearrangements were unique and identified in one or few genotypes, and only four variants were relatively frequent. These were a triple translocation T2A^t:4G:7G + T4A^t:7A^t, *per*Inv7A^t-1, T6G:7G, and T2G:4G:6G (21, 16, 12, and 13 genotypes, respectively). Taken together, these four variants accounted for approximately 16% of the whole materials we studied. Genotypes carrying the same rearrangement usually had similar C-banding patterns and were collected from the same, or closely located geographic regions. For example, (i) all genotypes with T2A^t:4G:7G + T4A^t:7A^t originated from Nakhichevan, Azerbaijan; (ii) T6G:7G and T2G:4G:6G were found in Erbil, Iraq; and (iii) *per*Inv7A^t-1 was collected in Iran and in the neighboring region of Sulaymaniyah, Iraq.

Other frequent translocations had a more restricted distribution and usually occurred in a single population. Only few genotypes carrying the same translocations were identified in spatially separated populations. These genotypes differed in their C-banding patterns, for example: (i) T2G:4G was found in four genotypes from Erbil, Iraq and in two genotypes from Siirt, Turkey; (ii) T1G:3G was identified in five ARA-0 genotypes from Dahuk, Iraq, one ARA-1 from Turkey and one ARA-1 genotype of a mixed accession IG 117895 collected in Syria; and (iii) T1G:5G was identified not only in three cytogenetically distinct *T. araraticum* ARA-0 genotypes from Armenia and Azerbaijan (Additional file 3: Fig. S6, *b10*) and ARA-1 from Turkey (Additional file 3: Fig. S12, *k2*), but also in *T. timopheevii* [71].

Most populations consisted of both genotypes with 'normal' karyotype and genotypes with one to several variants of chromosomal rearrangements. However, their ratio and spectra differed between regions (Fig. 6; Additional file 1: Note 2; Additional file 3: Fig. S5).

Cytogenetic diversity of GGA^{t}A^t wheats assessed using FISH markers

To further investigate the intraspecific diversity of *T. araraticum* and to assess their phylogenetic relationships with *T. timopheevii*, we carried out FISH using six DNA probes. The probes pTa-535, pSc119.2 and GAA$_n$ ensured chromosome identification [71] whereas Spelt-1, Spelt-52 and pAesp_SAT86 were used to estimate intra- and interspecific variation (Fig. 7).

The distribution of pTa-535 was monomorphic among *T. araraticum* and *T. timopheevii* (Fig. 7*a*, *b*). Only the pSc119.2 site located on 1A^{t}L discriminated ARA-1 and TIM from ARA-0 (Fig. 7*a*).

The variability of pAesp_SAT86 hybridization patterns was analyzed in four *T. timopheevii* and 26 *T. araraticum* genotypes, of them seven were from ARA-1 and 19 from ARA-0 lineages (Additional file 2: Table S1; Additional file 3: Fig. S13). Distribution of the pAesp_SAT86

probe in all *T. timopheevii* genotypes was similar except for chromosome 7A^t in genotype K-38555, which was modified due to a paracentric inversion or insertion of an unknown chromosomal fragment. Labeling patterns, however, were highly polymorphic for *T. araraticum* (Fig. 7; Additional file 3: Fig. S13). Large pAesp_SAT86 sites were found only on some G-genome chromosomes. The A^t-genome chromosomes possessed several small, but genetically informative polymorphic sites. Some of these sites were lineage-specific. Most obvious differences were observed for 3A^t, 4G and 7G chromosomes (Fig. 5). Thus, all TIM and ARA-1 genotypes carried the pAesp_SAT86 signal in the middle of 3A^{t}S, while for ARA-0 it was located sub-terminally on the long arm (Additional file 3: Fig. S13). One large pAesp_SAT86 cluster was present on the long arm of 4G in ARA-0, but on the short arm in ARA-1 and TIM. Two large and adjacent pAesp_SAT86 clusters were detected on 7GS in ARA-1 and TIM, but they were split between opposite chromosome arms in all ARA-0 genotypes. Differences between ARA-0 and ARA-1 in pAesp_SAT86 cluster position on 4G and 7G could be caused by pericentric inversions. Some other pAesp_SAT86 sites were identified in either one of the three groups. ARA-1 exhibited the largest polymorphism of pAesp_SAT86 labeling patterns among all groups (Additional file 3: Fig. S13).

FISH with Spelt-1 and Spelt-52 probes on chromosomes of 87 *T. araraticum* genotypes from different chromosomal groups and of different geographic origin, seven *T. timopheevii,* and one *T. zhukovskyi* genotypes revealed high intraspecific diversity of *T. araraticum* and low polymorphism in *T. timopheevii* (Additional file 3: Fig. S14). The broadest spectra of labeling patterns were found in genotypes from Dahuk and Sulaymaniyah (Iraq) and in the ARA-1 group from Turkey, while material from Transcaucasia exhibited the lowest variation. The polymorphism was due to variation in the number of Spelt-1 and Spelt-52 sites, their size and chromosomal location (Additional file 3: Fig. S14). The Spelt-1 signals in various combinations appeared in sub-telomeric regions of either one or both arms of 2A^t, 6A^t, and all G-genome chromosomes. The Spelt-52 signals were observed in various combinations on 2A^{t}S, 1GS, and 6GL chromosomes (Additional file 3: Fig. S14; Additional file 4: Table S7). Among them, two Spelt-1 sites (6GL and 7GL) and one Spelt-52 locus (2GS) were novel. The patterns of Spelt-1 and Spelt-52 repeats varied across geographic regions and between chromosomal groups (Additional file 1: Note 3; Additional file 4: Table S7).

Discussion

We present the most comprehensive survey of cytogenetic and genomic diversity of GGA^{t}A^t wheats. We describe the composition, distribution and characteristics of the GGA^{t}A^t genepool.

Building on our results, the latest published complementary genomics studies and state-of-the-art archaeobotanical evidence we revisit the domestication history of the GGA^{t}A^t wheats. We arrived at the following four key findings:

1. The GGA^{t}A^t genepool consists of three distinct lineages

We sampled the full breadth and depth of GGA^{t}A^t wheat diversity and discovered a clear genetic and geographic differentiation among extant GGA^{t}A^t wheats. Surprisingly, and supported by all marker types, three clearly distinct lineages were identified. The first lineage is comprised of all *T. timopheevii* genotypes (and the derived *T. militinae* and *T. zhukovskyi*; note that all *T. militinae* and all *T. zhukovskyi* accessions maintained *ex situ* in genebanks are each derived from only one original genotype). Interestingly, *T. araraticum* consists of two lineages that we preliminarily describe as 'ARA-0' and 'ARA-1'. This finding is in contrast to Kimber and Feldman [72] who concluded that *T. araraticum* does not contain cryptic species, molecularly distinct from those currently recognized. ARA-0 was found across the whole predicted area of species distribution. ARA-1 was only detected in south-eastern Turkey and in neighboring north-western Syria. It is interesting to note that only in this part of the Fertile Crescent, the two wild tetraploid wheat species, *T. dicoccoides* and *T. araraticum*, grow in abundance in mixed stands.

Based on Figure 2, collecting gaps are evident for *T. araraticum*, and future collecting missions should focus on four specific regions: (i) between the Euphrates river in the west and the Elazığ – Silvan – Mardin transect region in the east. Interestingly, so far only *T. dicoccoides* was reported from this region; (ii) between Bitlis (Turkey) – Amadiyah (Iraq) in the west and the Armenian border in the east including north-western Iran; (iii) between Adıyaman – Silvan in the south and Tunceli in the north; and (iv) between Hama in the south and west/north-west of Aleppo in Syria.

We found only two *T. araraticum* populations which contained representatives of both ARA lineages: (i) 45 km south-east of Kahramanmaraş to Gaziantep, and (ii) 4 km north of St. Simeon on the road to Afrin in Syria. However, in both cases, the ARA-1 lineage was significantly more frequent than ARA-0. This suggests at least a certain level of taxon boundary between ARA-0 and ARA1 lineages and should be investigated in the future.

Independent support for the existence of two wild *T. araraticum* lineages and their distribution comes from Mori et al. [73] based on 13 polymorphic chloroplast microsatellite markers (cpSSR). The 'plastogroup G-2' was distributed in south-eastern Turkey and northern Syria and was closely related to *Triticum timopheevii* [73]. However, Gornicki et al. [45], based on whole chloroplast genome sequence information and sufficient taxon sampling (13 Triticeae species and 1127 accessions; 163 accessions in common with our study), provided increased resolution

of the chloroplast genome phylogeny and showed that the *T. timopheevii* lineage possibly originated in northern Iraq (*and thus according to our data, belong to the ARA-0 lineage as no ARA-1 occurs in Iraq*). This was supported by Bernhardt et al. [56], who, based on re-sequencing 194 individuals at the chloroplast locus *ndhF* (2232 bp) and on whole genome chloroplast sequences of 183 individuals representing 15 Triticeae genera, showed that some ARA-0 and TIM genotypes are most closely related. All GGA^{t}A^t wheats re-sequenced by Bernhardt et al. [56] were considered in our study (3x ARA-0, 2x ARA-1, 1x TIM, 1x *T. militinae*) (Additional file 2: Table S1).

It is important to note that our results (i.e., the characteristics, composition and geographical distribution of ARA-0 and ARA-1 lineages) are not in agreement with the latest comprehensive taxonomical classification of wheat by Dorofeev et al. [11], who divided *T. araraticum* into two subspecies: subsp. *kurdistanicum* Dorof. et Migusch. and subsp. *araraticum* (Additional file 2: Table S1). We propose to re-classify the GGA^{t}A^t genepool taxonomically in the future.

2. The karyotypic composition of GGA^{t}A^t wheats is as complex as the phylogenetic history of the GGA^{t}A^t genepool

Based on karyotype analyses, translocation spectra and distribution of DNA probes, *T. araraticum* populations from Dahuk and Sulaymaniyah (both Iraq) harbored the highest karyotypic diversity among all *T. araraticum* populations studied. We consider the region around Dahuk in Northern Iraq as the center of diversity of *T. araraticum*, and this is probably the region where *T. araraticum* originated. This is supported by Bernhardt et al. [56] and Gornicki et al. [45] who traced chloroplast haplotypes from *Aegilops speltoides* growing in Iraq via *T. araraticum* (ARA-0) to *T. timopheevii* and *T. zhukovskyi*.

The karyotype 'similar' to those in 'normal' *T. timopheevii* was found in 44.6% of all *T. araraticum* genotypes. This is the group of candidates, in which the closest wild relative(s) to *T. timopheevii* is (are) expected. The frequency of the normal karyotype varied among countries and between populations. It is interesting to note that the Samaxi-Akhsu population in Azerbaijan and some populations near Dahuk (Iraq) possessed mostly karyotypically normal genotypes. Diagnostic C-bands for the ARA-1 lineage, both with normal and rearranged karyotypes, were 1A^{t}L3, 4A^{t}S7, 5A^{t}L3, 1GL5, 2GL7, 3GL7+L11. As expected, the number of C-bands characteristic for ARA-0 was smaller (due to the wide geographical distribution) and only two C-bands were lineage-specific and found in normal as well as translocated genotypes: 6A^{t}L3 and 5GL15.

Some FISH patterns suggested that *T. timopheevii* probably originated in Turkey and probably from ARA-1 (or, ARA-1 and TIM may have originated from a common ancestor, but then

diverged). This is supported by the following observations: (i) TIM and ARA-1 carry the pSc119.2 signal in the middle of $1A^t$ long arm, while this site was absent from ARA-0; (ii) all ARA-0 and most ARA-1 possessed the Spelt-52 signal on 6GL, but it is absent in TIM and five ARA-1 genotypes from Gaziantep-Kilis, Turkey. The distribution of Spelt-1 and Spelt-52 probes on chromosomes of these five genotypes was similar to, and in accession IG 116165 (ARA-1 from Gaziantep) almost identical with TIM; (iii) the pAesp_SAT86 patterns on chromosomes $3A^t$, 4G, and 7G are similar in TIM and ARA-1 but differed from ARA-0. Differences between ARA-1 and TIM based on FISH patterns of some other chromosomes as well as the results of C-banding and molecular analyses suggest that extant ARA-1 genotypes are not the direct progenitors of TIM but that the ARA-1 lineage is most closely related to it.

Based on AFLP, C-banding, FISH and *Jeli* retrotransposon markers, TIM was genetically most closely related to ARA-1. Additional evidence for the close relationship between TIM and ARA-1 lineages comes from allelic variation at the *VRN-1* locus of genome A^t [74]. This analysis revealed a 2.7 kb deletion in intron 1 of *VRN-A1* in three *T. timopheevii* and four *T. araraticum* accessions, which, according to our data, belong to the ARA-1 lineage. However, at *Vrn-G1*, TIM from Kastamonu in Turkey (PI 119442) shared the same haplotype (*Vrn1Ga*) with ARA-1 samples, while TIM from Georgia harbored haplotype *VRN-G1* as found in ARA-0. These results suggest multiple introgression events and incomplete lineage sorting as suggested by Bernhardt et al. [56, 75].

Regular chromosome pairing observed in the F_1 hybrids of lines with 'normal' karyotypes [55], identified in our study as ARA-0 $\times$ ARA-1 (Additional file 2: Table S1), suggested that karyotypic differences between ARA-0 and ARA-1 lineages are not associated with structural chromosomal rearrangements such as large translocations or inversions.

The emergence or loss of most lineage-specific Giemsa C-bands (Fig. 3) or FISH loci (Additional file 3: Fig. S13, S14) could be due to heterochromatin re-pattering: amplification, elimination or transposition of repetitive DNA sequences. Wide hybridization can also induce changes in C-banding and FISH patterns of *T. araraticum* chromosomes. Changes in pAesp_SAT86 hybridization patterns on 4G and 7G chromosomes, however, are likely to be caused by pericentric inversions, which are also frequent in common wheat [76]. The role of inversions in inter- and intraspecific divergence is probably underestimated. In our case, it seems possible that divergence between ARA-1/TIM (two inversions) from ARA-0 (no inversion) was associated with at least two pericentric inversions.

We did not find any genotype harboring both ARA-0 and ARA-1 specific FISH sites, although ARA-0 and ARA-1 genotypes co-existed in two populations in Turkey and Syria. However,

based on FISH (Spelt-1 site on chromosome 6GL and 7GS, respectively), hybridization between certain ARA-1 and ARA-0 lines can be predicted.

Iran occupies a marginal part of the distribution range of *T. araraticum*. An abundance of the pericentric inversion of the $7A^t$ chromosome in the Iranian group indicates that it is derived from Iraq. The karyotypically 'normal' genotype was probably introduced to Transcaucasia via Western Azerbaijan (Iran). The low diversity of FISH patterns and the low C-banding polymorphism of *T. araraticum* from Transcaucasia indicate that *T. araraticum* was introduced as a single event. Interestingly, the AFLP data suggested some similarity between ARA-0 from Armenia and Azerbaijan and *T. timopheevii*.

We hypothesize that homoploid hybrid speciation (HHS) [77-79] and incomplete lineage sorting may be the possible mechanisms explaining the origin of the ARA-1 lineage. ARA-1 grows in sympatry and in mixed populations with *T. dicoccoides* and is phylogenetically most closely related to *T. dicoccoides*. ARA-1 could be derived from the hybridization of *T. timopheevii* × *T. dicoccoides*; or alternatively, ARA-1 and TIM could be derived from the hybridization ARA-0 × *T. dicoccoides*.

3. Does the *T. timopheevii* population found in western Georgia represent the last remnant of a widespread ancient cultivation area of GGA^tA^t wheats?

Wild emmer *T. dicoccoides* belongs to the first cereals to be domesticated by humans in the Fertile Crescent and the evolution and domestication history of *T. dicoccoides* are relatively well studied [26, 69, 80]. Domesticated emmer *Triticum dicoccon* Schrank was a staple crop of Neolithic agriculture, was widely cultivated for over 10,000 years and harbored impressive genetic diversity [81-83]. The domestication of *T. dicoccoides* provided the key for durum wheat [84] and bread wheat evolution [85].

Much less is known about *T. timopheevii*. It is believed that *T. timopheevii* is the domesticated form of *T. araraticum* [11, 17, 18]. In contrast to *T. dicoccon*, *T. timopheevii* was, since its discovery, considered as a '*monomorphous narrowly endemic species*' [86] cultivated in few villages of western Georgia ([14, 87] Additional file 1: Note 4). Dekaprelevich and Menabde [86] noticed that the area of cultivation had probably been larger in the past. The last plants of *T. timopheevii in situ* were found by the expedition of the N.I. Vavilov Institute of Plant Genetic Resources (VIR, Russia) in 1983 near the village of Mekvena (Tskhaltubo, Georgia) and deposited in the VIR genebank under accession number K-56422 [E.V. Zuev, personal communication]. Today, the widespread view is that the cultivation area of *T. timopheevii* was restricted to Georgia in the (recent) past [10, 88].

However, hulled tetraploid wheat morphologically similar to *T. timopheevii* was identified at three Neolithic sites and one Bronze Age site in northern Greece and described by Jones et al. [36] as a **'New'** Glume Wheat (NGW). The glume bases of these archaeological finds morphologically resemble *T. timopheevii* more than any other extant domesticated wheat [36]. After these finds of NGW in Greece, this wheat was also identified at Neolithic and Bronze Age sites in Turkey, Bulgaria, Romania, Hungary, Slovakia, Austria, Italy, Poland, Germany and France [89-100]. Earlier finds of an 'unusual' glume wheat in Serbia [101] and Turkey [102] have subsequently been recognized as NGW [36, 103]. Criteria were also established for distinguishing the grains of NGW [96].

At some sites, NGW appeared as a minor component and may have been part of the accompanying weed flora of cereal fields [99, 104]. In other cases, it was probably cultivated in a mix with einkorn [36, 96] and/or emmer. The recovery of large quantities in storage deposits of whole spikelets at Çatalhöyük in Turkey, caryopses and spikelet bases at the early Bronze Age settlement of Clermont-Ferrand in France, and rich deposits including whole spikelets at bronze age sites in Italy, demonstrated that, at least in some places, NGW was a major crop in itself [92, 104-108].

Based on intensive archaeobotanical investigations at Çatalhöyük in central Anatolia, for example, NGW was the predominant hulled wheat, overtaking emmer wheat around 6500 cal BC and remaining dominant until the site's abandonment c. 5500 cal BC. The finds suggested that this wheat was a distinct crop, processed, stored, and presumably grown, separately from other glume wheats. NGW formed part of a diverse plant food assemblage at Neolithic Çatalhöyük, including six cereals, five pulses and a range of fruits, nuts and other plants, which enabled this early farming community to persist for 1500 years (c. 7100 to 5500 cal BC) [92, 105].

Recently, polymerase chain reactions specific for the wheat B and G genomes, and extraction procedures optimized for retrieval of DNA fragments from heat-damaged charred material, have been used to identify archaeological finds of NGW [109]. DNA sequences from the G genome were detected in two of these samples, the first comprising grain from the mid 7[th] millennium BC at Çatalhöyük in Turkey, and the second made up of glume bases from the later 5[th] millennium BC site of Miechowice 4 in Poland. These results provide evidence that NGW is indeed a cultivated member of the GGA^{t}A^t genepool [109]. As NGW is a recognized wheat type across a broad geographic area in prehistory, dating back to the 9[th] millennium BC in SW Asia, this indicates that *T. timopheevii* (*sensu lato, s.l. = domesticated GGA^{t}A^t wheat in general*), was domesticated from *T. araraticum* during early agriculture, and was widely cultivated in the prehistoric past [109].

This raises the question of whether the few populations of *T. timopheevii* (*sensu stricto*, *s.str.*) found in western Georgia were the last remnants of a wider GGA^{t}A^t wheat cultivation or whether the *T. timopheevii* of Georgia was a local domestication independent of the domestication of *T. araraticum* in SW Asia. To answer this question, sequence information for NGW, the Georgian *T. timopheevii*, and the two lineages of *T. araraticum* (ARA-0, ARA-1) need to be compared.

N.I.Vavilov [110] suggested that *T. timopheevii* of western Georgia was probably originally introduced from north-eastern Turkey. Menabde and Ericzjan (1942) as cited by Dorofeev et al. [11] associated the origin of *T. timopheevii* with the region of the ancient kingdom of Urartu, whence immigrant ancestors of modern-day Georgians introduced it into western Georgia. Certainly, the possibility of introduction of Timofeev's wheat into Georgia from the south should not be rejected [11]. Is the domestication history of *T. timopheevii* *s.str.* connected with other endemic wheats of Georgia, such as *T. karamyschevii* Nevski (*T. georgicum* Dekapr. et Menabde or *T. paleocolchicum* Menabde) and *T. carthlicum* Nevski, which were cultivated by Mingrelians in Western Georgia [111, 112]?

4. Was the cultivation range of *T. timopheevii* (*s.str.*) wider in the recent past?

We screened all available passport data and found three interesting cases, which could potentially help to reconstruct the recent past cultivation range of *T. timopheevii* *s.str*: Interestingly, two *T. timopheevii* accessions maintained in two *ex situ* genebanks are reported to originate from Turkey (Additional file 2: Table S1) [https://www.genesys-pgr.org/]: (i) ATRI 3433 (TRI 3433) conserved in the Federal *ex situ* Genebank of Germany hosted at the Institute of Plant Genetics and Crop Plant Research, IPK, Gatersleben. This *T. timopheevii* line was most likely collected by E. Baur in Turkey in 1926 [113]; and (ii) PI 119442 identified among a barley sample obtained from a market in Araç, near Kastamonu, Turkey (Fig. 2) in 1936 by Westover and Wellmann, and maintained at the National Plant Germplasm System, USDA-ARS, USA. Both accessions harbor the 'normal' karyotype of *T. timopheevii*, both were characterized in our studies and confirmed as a typical *T. timopheevii*. Additionally, the accession TA1900, presumably collected 32 km south of Denizli near Kahramanmaraş in Turkey on the 14th of August 1959 and maintained in the wheat germplasm collection of the Wheat Genetics Resource Center, Kansas State University, U.S.A., is interesting because it shared karyotypic features of *T. timopheevii* and the ARA-0 lineage (Fig. 3; Additional file 3: Fig. S11). However, we are not fully convinced that this line is a true natural hybrid. Based on the passport data, this accession could potentially have escaped from an experimental field or a breeding station, or received introgression(s) during *ex situ* maintenance [114]. Assuming that the passport data is correct

(Additional file 2: Table S1), we could speculate that *T. timopheevii* may have been cultivated in Turkey during the first half of the 20th century. However, is this realistic option?

We believe not. As reported by Stoletova [87], Dekaprelevich and Menabde [86, 115], Menabde [116], Dekaprelevich [117], *T. timopheevii s.str.* was part of the spring landrace Zanduri (mixture of *T. timopheevii s.str.* and *T. monococcum*) and well adapted to the historical provinces Lechkhumi and Racha of Georgia (Additional file 1: Note 4). The Zanduri landrace was cultivated in the '*humid and moderately cool climate zone 400–800*' m above the sea level [11]. Martynov et al. [67] reported that *T. timopheevii* potentially has a '*low potential for plasticity*' and is not drought tolerant. Climate at origin based on bioclimatic variables [118, 119] clearly differs between the regions of Western Georgia where the Zanduri landrace grew till the recent past, and both Kastamonu and Kahramanmaraş regions in Turkey (Additional file 1: Note 5). We speculate that the three *T. timopheevii* accessions which were collected in Turkey were probably introduced from Transcaucasia or elsewhere and may have been left over from unsuccessful cultivation or breeding experiments of *T. timopheevii s. str.* in the recent historical past. Also, based on botanical records, *T. timopheevii (s.str.)* has only been identified in Georgia, but not in Turkey or elsewhere [120, 121]. From this we conclude that the cultivation range of *T. timopheevii* (*s str.*) was not wider in the recent past.

Conclusions

The evolutionary history of the GGA^{t}A^t wheat genepool is complex. However, some pieces of the puzzle are clearly recognizable: the region around Dahuk in northern Iraq can be considered the center of origin, but also the center of diversity of *T. araraticum*.

The origin of *T. timopheevii s.str.* remains unclear, but we speculate that it was probably introduced from Turkey, on the grounds that wild *T. araraticum* does not grow in Georgia and that *T. timopheevii s.str.* is more closely related to *T. araraticum* from Turkey or northern Iraq than to the Transcaucasian types.

Based on bioclimatic variables, we predict that *T. timopheevii s.str.* is maladapted to the climate outside Western Georgia. If this speculation is correct, it suggests a sister-group relationship between (i) the Georgian *T. timopheevii* (*s.str.*) and both *T. araraticum* lineages (ARA-0, ARA-1), but also between (ii) *T. timopheevii s.str.* and the prehistoric SW Asian *T. timopheevii s.l.*

The oldest known records of prehistoric *T. timopheevii s.l.* are of Turkish origin: (i) A□ıklı Höyük in Cappadocia, and (ii) Cafer Höyük (just inside the Fertile Crescent and potentially older than A□ıklı Höyük but less thoroughly dated) [122, 123], though more prehistoric finds may be identified in the future. Emphasis for further archaeobotanical research should be given to the

central part of the Fertile Crescent, including northern Iraq because this is probably the region where wild *T. araraticum* originated.

The unexpected discovery of the ARA-1 lineage is exciting and may have implications for our understanding of the origins of agriculture in southwest Asia. Was ARA-1 involved in the formation of *T. timopheevii s.l.* (= NGW)?

The distribution area of the ARA-1 lineage requires our attention. It is interesting to note that the ARA-1 lineage (GGA^{t}A^t) and *T. dicoccoides* (BBAA) grow in sympatry and in mixed populations only in a peculiar geographical area in the Northern Levant. This specific area is located between Kahramanmaraş in the north, Gaziantep in the east, Aleppo in the south and the eastern foothills of the Nur Dağlari mountains in the west. It is interesting that the closest extant wild relatives of domesticated einkorn, barley and emmer were also collected here: (i) the beta race of wild einkorn [38] and the closest wild relatives to einkorn *btr1* type [124]; (ii) the closest wild barley to *btr2* barley [125] and to *btr1b* barley [126]. This specific region was among the areas predicted with high probability as potential refugia for wild barley during the Last Glacial Maximum (LGM) by Jakob et al. [127]; and (iii) wild emmer subgroup II of the central-eastern race [26]. On the other hand, some genetic research has suggested that domesticated emmer wheat and barley received substantial genetic input from other regions of the Fertile Crescent, resulting in hybridized populations of different wild lineages indicating a mosaic of ancestry from multiple source populations [80, 128, 129]. It will be interesting to see whether further genetic and archaeobotanical research on *T. araraticum* lineages and *T. timopheevii s.l.* can help to resolve this issue.

Finally, and fortunately, 1294 accessions of *T. araraticum* and 590 accessions of *T. timopheevii s.str.* are stored in 24 and 37 *ex situ* genebank repositories, respectively [130]. Our study provides the basis for a more efficient use of *T. araraticum* and *T. timopheevii* materials for crop improvement.

Methods

Germplasm

A comprehensive germplasm collection of tetraploid wheats was established comprising 862 genebank accessions (1–5 genotypes per accession number): 450 *T. araraticum,* 88 *T. timopheevii,* three *T. militinae* Zhuk. et Migusch.*,* one *T. zhukovskyi,* 307 *T. dicoccoides,* eight *T. dicoccon,* and five *T. durum.* Additionally, the following materials were included: (i) four samples of *T. araraticum* collected by Dr. Nelli Hovhannisyan in Armenia; (ii) 17 samples of *T. araraticum* and 12 samples of *T. dicoccoides* collected by Dr. H. Özkan in Turkey; and (iii) 26

samples of *T. dicoccoides* collected by Drs. E. Badaeva, O.M. Raskina and A. Belyayev in Israel. Altogether, 921 accessions of wild and domesticated tetraploid wheats were examined (Additional file 2: Table S1).

A subset of 786 tetraploid wheat genotypes representing 765 genebank accessions was examined using Sequence-Specific Amplification Polymorphism (SSAP) markers. The subset included 360 genotypes of *T. araraticum*, 76 *T. timopheevii* (including two *T. militinae*), while 350 genotypes of *T. dicoccoides* were considered as an outgroup. The whole collection was single-seed descended (SSD) at least twice under field conditions (2009–2012) and taxonomically re-identified in the field at IPK, Gatersleben in 2011 (Additional file 2: Table S1).

A subset of 103 genotypes, including 38 *T. araraticum*, two *T. militinae*, 13 *T. timopheevii*, 40 *T. dicoccoides,* 10 *T. dicoccon,* and four *T. durum* was selected for a complementary analysis using Amplified Fragment Length Polymorphism (AFLP) markers to infer the population structure of GGA^{t}A^t wheats (Additional file 2: Table S1).

Karyotype diversity of 370 *T. araraticum* accessions was assessed by C-banding and Fluorescence *in situ* hybridization (FISH) in comparison with 17 *T. timopheevii* and one *T. zhukovskyi* genotypes. According to C-banding analysis, most *T. araraticum* accessions (353 of 370) were karyotypically uniform and were treated as single genotype each. Seventeen accessions were heterogeneous and consisted of two (13 accessions) or even three (four accessions) cytogenetically distinct genotypes, which were treated as different entities (genotypes). Altogether 391 distinct genotypes of *T. araraticum* were distinguished. In total, 243 genotypes were shared with SSAP and 33 with AFLP analyses (Additional file 2: Table S1). In order to infer the population structure of GGA^{t}A^t wheats, 265 typical genotypes were selected representing all karyotypic variants.

Seven *T. timopheevii,* one *T. zhukovskyi,* and 87 *T. araraticum* genotypes representing all chromosomal groups were selected for Fluorescence i*n situ* hybridization (FISH) analysis (Additional file 2: Table S1).

Molecular analysis

SSAP analysis

Evolutionary relationships among wild and domesticated tetraploid wheat taxa were first inferred based on polymorphic retrotransposon insertions. For this, the highly multiplex genome fingerprinting method SSAP was implemented based on polymorphic insertions of retrotransposon families *BARE-1* and *Jeli* spread across wheat chromosomes. DNA was isolated from freeze-dried leaves of 786 SSD plants, using the Qiagen DNeasy Kit (Hilden, Germany). SSAP protocol was based on Konovalov et al. [68] with further optimizations for capillary-based

fragment detection (Additional file 1: Note 6). In total, 656 polymorphic markers were generated for BBAA- and GGA^{t}A^t-genome wheats by amplification of multiple retrotransposon insertion sites (Additional file 5). Data analysis was performed in SplitsTree 4.15.1 [131]. NeighborNet planar graphs of Dice distances [132] were constructed based on presence/absence of SSAP bands in the samples.

AFLP analysis

The AFLP protocol, as described by Zabeau and Vos [133], was performed with minor modifications according to Altıntas□ et al. [134] and Alsaleh et al. [135]. In total, six AFLP primer combinations were used to screen the collection of 103 lines (Additional file 1: Note 7). A total of 146 polymorphic AFLP markers were detected across all genotypes (Additional file 6). Neighbor-Joining (NJ) trees were computed based on Jaccard distances [136, 137]. NeighborNet planar graphs were generated based on Hamming distances [131]. Genetic diversity parameters and genetic distances were calculated using Genalex 6.5 [138].

Cytogenetic analysis

C-banding

Chromosomal preparation and C-banding procedure followed the protocol published by [31]. The A^t- and G-genome chromosomes were classified according to nomenclature proposed by [139] except for chromosomes 3A^t and 4A^t. Based on meiotic analysis of the F$_1$ *T. timopheevii* × *T. turgidum* hybrids [49] and considering karyotype structure of the synthetic wheat *T. × soveticum* (Zhebrak) [140], the chromosomes 3A^t and 4A^t were exchanged. Population structure of *T. araraticum* and the phylogenetic relationship with *T. timopheevii* were inferred based on chromosomal passports compiled for 265 genotypes with high-quality C-banding patterns (247 *T. araraticum*, 17 *T. timopheevii*, one *T. zhukovskyi*). Chromosomal passports (Additional file 7) were constructed by comparing the karyotype of the particular accession with the generalized idiogram of A^t- and G-genome chromosomes (Fig. 5), as described for *T. dicoccon* Schrank [69] (Additional file 1: Note 8). The 96 positions of informative C-bands (see Fig. 5) considered for chromosomal passport construction are indicated in Additional file 7. At each of these positions, C-bands of different sizes were treated as different entities ("polymorphic C-bands"). The full dataset based on 265 genotypes and 379 polymorphic C-bands is provided in Additional file 7.

Fluorescence *in situ* hybridization (FISH)

Two polymorphic G-genome specific DNA probes, Spelt-1 [141] and Spelt-52 [142], were used for all 95 (87 *T. araraticum,* seven *T. timopheevii* and one *T. zhukovskyi*) genotypes considered for FISH analysis. A total of 57 polymorphic FISH (46 Spelt-1, 11 Spelt-52) markers were detected across 95 genotypes (Additional file 8). Additionally, 26 of the 95 genotypes (Additional file 2: Table S1) were analyzed using the probe pAesp_SAT86 [143], which also showed differences between the accessions. The probes pSc119.2 [144], GAA_n [145], and pTa-535 [146] were subsequently hybridized to the same chromosomal spread to allow chromosome identification. Classification of pSc119.2- and pTa-535-labeled chromosomes followed the nomenclature of [47, 71]. Chromosomes hybridized with the GAA_{10} microsatellite sequence were classified according to nomenclature suggested for C-banded chromosomes (Additional file 1: Note 8).

Legends to Figures

Fig. 1 Genetic relationships between GGA^{t}A^t and BBAA wheats. NeighborNet planar graph of Dice distances representing the diversity of 786 GGA^{t}A^t and BBAA tetraploid wheat genotypes based on 656 SSAP markers.

Fig. 2 Natural geographical distribution of wild tetraploid *T. araraticum* and *T. dicoccoides*. Green dots correspond to collection sites of ARA-0 accessions, pink dots to ARA-1, and dark blue dots to *T. dicoccoides* (DIC). The collection sites of *T. timopheevii* and *T. zhukovskyi* are shown with turquoise and yellow dots, respectively. Key excavation sites in Turkey where NGW was identified are indicated with red triangles.

Fig. 3 Comparison of the C-banding patterns of *T. dicoccoides* (DIC), *T. timopheevii* (TIM, *a–c,* normal karyotypes), *T. araraticum* ARA-1 (*d–f*) and ARA-0 (*g–t*). DIC (IG 117174, Gaziantep), *a* – KU-1818 (Georgia); *b* – PI 119442; *c* – TA1900; *d* – IG 116165; *e* – PI 654340; *f* – KU-1950; *g* – CItr 17677; *h* – KU-8917; *i* – KU-8909; *j* – KU-1933 (all from Turkey); *k* – CItr 17680 (Iran); *l* – PI 427381 (Erbil, Iraq); *m* – PI 538518*; n* – PI 427425 (Dahuk, Iraq); *o* – KU-8705; *p* – KU-8695 (Shaqlawa, Erbil, Iraq); *q* – KU-8451; *r* – KU-8774 (Sulaymaniyah, Iraq); *s* – TRI 11945 (Nakhichevan); *t* – KU-1901 (Armenia). 1–7 – homoeologous groups. C-bands typical for ARA-1 are indicated with blue arrows, for ARA-0 – with green arrows, and C-bands characteristic for *T. timopheevii* – with red arrows. Black arrows point to rearranged chromosomes in genotypes.

Fig. 4 Chromosomal rearrangements identified in *T. araraticum*. The number of translocation variant corresponds to the number of the respective variant in Additional file 4: Table S6. Novel variants are designated with black numerals, and already known variants by red numerals.

Fig. 5 Generalized idiogram and nomenclature of the A^t- and G-genome chromosomes. The C-banding pattern is shown on the left, the pSc119.2 (red) and pAesp_SAT86 (green) pattern on the right side of each chromosome. 1–7 – homoeologous groups; S – short arm, L – long arm. The numerals on the left-hand side designate putative positions of C-bands/FISH sites that can be detected on the chromosome arm; C-bands specific for the ARA-1 group are shown with pink numerals, C-bands specific for the ARA-0 group are indicated by green numerals. Red asterisks on the right-hand side indicate C-bands that were considered for the "chromosomal passport".

Fig. 6 Intraspecific divergence of *T. araraticum* and *T. timopheevii*. Combinations of chromosome arms in rearranged chromosomes are designated. Line colors mark the different groups: ARA-0 (green), ARA-1 (pink) and *T. timopheevii* (black). Solid arrows designate novel rearrangements; arrows with asterisk designate previously described rearrangements [31, 54]. The numerals above/next to the arrows indicate the number of accessions carrying the respective translocation.

Fig. 7 Distribution of different families of tandem repeats on chromosomes of *T. timopheevii* and *T. araraticum*. *Triticum timopheevii,* KU-107 (*a*), and *T. araraticum,* CItr 17680, ARA-0 (*b*), KU-8944, ARA-0 (*c*), KU-1984B, ARA-1 (*d*), PI 427364, ARA-0 (*e*), and 2630, ARA-1 (*f*). The following probe combinations were used: *a, b* – pSc119.2 (green) + pTa-535 (red); *d, e* – pAesp_SAT86 (red) + GAA$_n$ (green); *c, f* – Spelt-1 (red) + Spelt-52 (green). The position of pSc119.2 site on 1A^t chromosome typical for *T. timopheevii* and ARA-1 is shown with an arrow (*a*). Translocated chromosomes (*c, d*) are arrowed. Chromosomes are designated according to genetic nomenclature; the A^t-genome chromosomes are designated with yellow numerals and the G-genome chromosomes with white numerals. Scale bar, 10 μm.

Supplementary information

Supplementary information accompanies this paper at http xxx

Additional file 1:

Note 1. Regarding the first finds of *T. timopheevii*. **Note 2.** Region-specific chromosomal rearrangements. **Note 3.** Additional findings of the FISH analysis. **Note 4.** Information extracted and translated from Dorofeev et al. [11], regarding the cultivation area and geographical distribution of *Triticum timopheevii*, and the "Zanduri" wheat complex. Translation by H. Knüpffer and colleagues, February 2020. **Note 5.** Was the cultivation range of *T. timopheevii* (*s.str.*) wider in the recent past? **Note 6.** Details about the SSAP protocol and analysis. **Note 7.** Details about the AFLP protocol and analysis. **Note 8.** Details about the C-banding and FISH protocols and analyses.

Additional file 2:

Table S1. List of materials investigated. Column 1: Species; Column 2: Additional taxonomical classification information based on passport data (when available); Column 3: Accession number; Column 4: Seed/ DNA sources; Column 5: Other identifiers; Column 6: SSAP code; Column 7: SSAP-NNet group; Column 8: AFLP-code; Column 9: AFLP-NNet group; Column 10: Group assignment of GGA^{t}A^t wheat, defined visually based on C-banding-analysis; Column 11: Translocation type - codes of chromosomal rearrangements identified in *T. araraticum*; Column 12: PASSP - "chromosome passport" codes; Column 13: FISH - combination of probes used in FISH analysis; Column 14: Translocation type in accessions identified by Kawahara et al. [55]; Column 15: Accessions considered by Mori et al. [73] and their haplotypes; Column 16: Accessions considered by Gornicki et al. [45] and their haplotype numbers; Column 17: Accessions considered by Bernhardt et al. [56]; Column 18: Country of origin; Column 19: Collection site.

Additional file 3:

Fig. S1. NeighborNet phylogenetic networks constructed for BBAA and GGA^{t}A^t-genome wheat genotypes based on Dice genetic distances and SSAP markers. **Fig. S2.** NeighborNet phylogenetic networks constructed for GGA^{t}A^t-genome wheat genotypes only based on Dice genetic distances and SSAP markers. **Fig. S3. (a)** NJ tree for 103 genotypes based on Jaccard distances; **(b)** NeighborNet (NNet) planar graph based on Hamming distances for 59 genotypes. **(c)** NNet planar graph based on Hamming distances for 52 genotypes. **Fig. S4.** Neighbor-Joining (NJ) tree for 265 genotypes of *T. araraticum* and *T. timopheevii* using 379 C-banding markers based on Jaccard distances. **Fig. S5.** The ratios of chromosomal rearrangements among geographic populations of *T. araraticum.* **Fig. S6.** Polymorphism of the C-banding patterns of *T. araraticum* from Transcaucasia. **Fig. S7.** Polymorphism of the C-banding patterns of *T. araraticum* from Iraq (I). **Fig. S8.** Polymorphism of the C-banding patterns of *T. araraticum* from Erbil, Iraq (II). **Fig. S9.** Polymorphism of the C-banding patterns of *T. araraticum* from Sulaymaniyah, Iraq (III). **Fig. S10.** Polymorphism of the C-banding patterns of *T. araraticum* from Iran (*g1–g18*) and Syria (*h1–h9*). **Fig. S11.** Polymorphism of the C-banding patterns of *T. araraticum* from Turkey (ARA-0). **Fig. S12.** Polymorphism of the C-banding patterns of *T. araraticum* from Turkey (ARA-1). **Fig. S13.** Comparison of the distribution of pAesp_SAT86 probe on chromosomes of *T. dicoccoides* (DIC), *T. timopheevii* TIM (T1–T4), *T. araraticum* ARA-1 (A1–A7) and ARA-0 (A8–A24). **Fig. S14.** Distribution of Spelt-1 (red) and Spelt-52 (green) probes on chromosomes of *T. timopheevii* (T) and *T. araraticum* representing ARA-1 (A1, A2, A4–A10) and ARA-0 (A3, A11–A29) groups.

Additional file 4:

Table S2. Summary statistics and genetic distances based on SSAP markers. **Table S3.** Summary statistics and genetic distances based on AFLP markers. **Table S4.** Summary statistics and genetic distances based on C-bands. **Table S5.** Summary statistics and genetic distances based on FISH markers. **Table S6.** Variants of chromosomal rearrangements in *T. araraticum.* **Table S7.** Distribution of Spelt-1 and Spelt-52 loci in geographic/chromosomal groups in comparison with *T. timopheevii.*

Additional file 5:

SSAP dataset. This dataset consists of 656 SSAP markers (401 *BARE-1* and 255 *Jeli* markers) for 786 genotypes.

Additional file 6:

AFLP dataset. This dataset consists of 146 polymorphic AFLP markers across 103 genotypes.

Additional file 7:

C-banding dataset. The dataset consists of 379 C-banding markers across 265 genotypes. The 96 putative positions of informative C-bands considered for passport data (Fig. 5) are indicated.

Additional file 8:

FISH dataset. The dataset consists of 57 polymorphic Spelt-1 (C-AV) and Spelt-52 (AW-BG) markers across 95 genotypes.

List of abbreviations

AFLP, Amplified Fragment Length Polymorphism
ARA-0, *T. araraticum* lineage 0 (wild, widespread)
ARA-1, *T. araraticum* lineage 1 (wild, native only to south-eastern Turkey (Adiyaman, Kahramanmaraş, Gaziantep, Kilis) and in north-western Syria)
DIC, *T. dicoccoides*
FISH, Fluorescence *in situ* hybridization
LTR, Long Terminal Repeat
NGW, new glume wheat
SSAP, Sequence-Specific Amplification Polymorphism
SSD, single seed descent
TIM, *T. timopheevii* (domesticated)

Acknowledgements

We thank the following for providing seeds, DNA and passport data: N. I. Vavilov Institute of Plant Genetic Resources (VIR, Russia), the Federal *ex situ* Genebank of Germany, Gatersleben (IPK, Germany), the International Center for Agricultural Research in the Dry Areas (ICARDA), the Max Planck Institute for Plant Breeding Research (MPIPZ, Germany), Kyoto University (National Bioresource Project, NBRP, Japan), the John Innes Centre (JIC, UK), Wheat Genetics Resource Centre, Kansas State University (WGRC, USA), Centre for Genetic Resources (WUR, CGN, Netherlands), the Australian Winter Cereal Collection Tamworth (AWCC Australia), the National Plant Germplasm System (USDA-ARS, USA), Eastern Cereal and Oilseed Research Centre, Agriculture and Agri-Food Canada (Canada), Botanical Institute Tbilisi (Georgia), Tbilisi Agricultural Institute (Georgia), Institute of Cytology and Genetics (Siberian Branch of the Russian Academy of Sciences, Russia), and the Russian State Agrarian University Moscow, Timiryazev Agricultural Academy (Russia). We also thank Drs. P.A. Gandilyan, M. Nazarova and I.G. Gukasyan (all Botanical Institute, Erevan, Armenia), Drs. I.D.O. Mustafaev and N.Kh. Aminov (both Institute of Genetic Resources of Azerbaijan, Baku, Azerbaijan) for providing materials from their collection missions. We thank the following for excellent technical assistance: Sigi Effgen (MPIPZ), Christiane Kehler, Birgit Dubsky, Heike Harms, Ute Krajewski, Marita Nix, Kerstin Wolf, Jürgen Marlow, Michael Grau, Peter Schreiber (all IPK), the 'Experimental Fields and Nurseries' and 'Genome Diversity' groups at IPK. We thank Dr. E.V. Zuev for providing us the data on collection sites of *T. araraticum* and *T. timopheevii* accessions from VIR collection. We are greatly indebted to Maarten Koornneef and George Willcox for discussions and support. We thank two anonymous reviewers, who helped us to improve an earlier version of this manuscript.

Authors' contributions

Ekaterina D. Badaeva conceived, designed and conducted the cytogenetic experiments and analyzed data, wrote the first manuscript version.

Fedor Konovalov conducted the SSAP experiments and analyzed the data.

Helmut Knüpffer verified genebank information concerning the origin of the material, translated texts from Russian into English, and contributed to the discussions and editing of the manuscript.

Agostino Fricano, supported the molecular data analysis.

Alevtina S. Ruban contributed to the FISH analysis.

Zakaria Kehel conducted the simulation of regional agro-ecological variation based on bioclimatic variables.

Svyatoslav A. Zoshchuk contributed to the work on mapping of Spelt-1 and Spelt-52 probes on *T. timopheevii* and *T. araraticum* genotypes.

Sergey A. Surzhikov designed and synthesized oligo-probes for FISH analyses.

Kerstin Neumann contributed to data analysis, supported the field trials and edited the manuscript.

Andreas Graner contributed to the discussions and editing of the manuscript.

Anna Filatenko phenotyped and taxonomically re-identified the collection, contributed to the discussions and edited the manuscript.

Karl Hammer phenotyped and taxonomically re-identified the collection, contributed to the discussions and edited the manuscript.

Amy Bogaard contributed to the discussions and editing of the manuscript.

Glynis Jones contributed to the discussions and editing of the manuscript.

Hakan Özkan conducted the AFLP experiments, analyzed the data and edited the manuscript.

Benjamin Kilian conceived the project, established the germplasm collection, analyzed data, interpreted results, wrote and edited the manuscript.

Funding

This work was supported by the European Community's Seventh Framework Programme (FP7, 2007–2013) under the grant agreement n°FP7-613556, Whealbi project [http://www.whealbi.eu/project/], the German Research Foundation (DFG) (FKZ KI 1465/1-1 and FKZ KI 1465/5-1), the Federal Office for Agriculture and Food (BLE) of Germany (FKZ 01/12-13-KAD), the Leibniz Institute of Plant Genetics and Crop Research (IPK, Gatersleben, Germany), the Max Planck Institute for Plant Breeding Research (MPIPZ, Cologne, Germany) and the Crop Wild Relatives Project (*Adapting Agriculture to Climate Change: Collecting, Protecting and Preparing Crop Wild Relatives*) which is supported by the Government of Norway and managed by the Global Crop Diversity Trust [https://www.cwrdiversity.org/]. EDB acknowledges support from the Russian State Foundation for Basic Research (projects 20-04-00284, 17-04-00087, 14-04-00247), State Budget Project and the Program "Dynamics and Preservation of Gene Pools" from the Presidium of the Russian Academy of Sciences.

Availability of data and materials

All data sets supporting the conclusions of this article are available in the Additional files. The acquisition of plant material used in this study complies with institutional, national, and international guidelines.

Ethics approval

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

References

1. Tanksley SD, McCouch SR. Seed banks and molecular maps: Unlocking genetic potential from the wild. Science (New York) 1997; doi: 10.1126/science.277.5329.1063.

2. van Heerwaarden J, Doebley J, Briggs WH, Glaubitz JC, Goodman MM, de Jesus Sanchez Gonzalez J, Ross-Ibarra J. Genetic signals of origin, spread, and introgression in a large sample of maize landraces. Proc Natl Acad Sci U.S.A. 2010; doi: 10.1073/pnas.1013011108.

3. Dempewolf H, Baute G, Anderson J, Kilian B, Smith C, Guarino L. Past and future use of wild relatives in crop breeding. Crop Sci 2017; 10.2135/cropsci2016.10.0885.

4. Kilian B, Dempewolf H, Guarino L, Werner P, Coyne C, Warburton ML. Crop Sci special issue: Adapting agriculture to climate change: A walk on the wild side. Crop Sci 2021; https://doi.org/10.1002/csc2.20418.

5. Iizumi T, Furuya J, Shen Z, Kim W, Okada M, Fujimori S, Hasegawa T, Nishimori M. Responses of crop yield growth to global temperature and socioeconomic changes. Sci Rep 2017; doi: 10.1038/s41598-017-08214-4 .

6.	Ray DK, Mueller ND, West PC, Foley JA. Yield trends are insufficient to double global crop production by 2050. PLOS ONE 2013; doi: 10.1371/journal.pone.0066428

7.	Ray DK, Ramankutty N, Mueller ND, West PC, Foley JA. Recent patterns of crop yield growth and stagnation. Nat Com 2012, https://doi.org/10.1038/ncomms2296.

8.	Placido DF, Campbell MT, Folsom JJ, Cui X, Kruger GR, Baenziger PS, Walia H. Introgression of novel traits from a wild wheat relative improves drought adaptation in wheat. Plant Phys 2013; doi: 10.1104/pp.113.214262.

9.	Zhang H, Mittal N, Leamy LJ, Barazani O, Song B-H: Back into the wild — Apply untapped genetic diversity of wild relatives for crop improvement. Evol Appl 2017; https://doi.org/10.1111/eva.12434.

10.	Zohary D, Hopf M, Weiss E. Domestication of plants in the Old World, 4th edition. Oxford: Oxford Univ. Press; 2012.

11.	Dorofeev VF, Filatenko AA, Migushova EF, Udachin RA, Jakubziner MM. Cultivated flora of the USSR. v. 1. Wheat, vol. 1. Leningrad: Kolos; 1979.

12.	Goncharov NP. Comparative genetics of wheats and their related species 2 edition. Novosibirsk: Academic Publishing Group @Geo@; 2012.

13.	Nazarova EA: To the taxonomy and nomenclature of Triticum araraticum Jakubz. (Poaceae). Flora of Plant Resources of Armenia 2007;16:32-4.

14.	Zhukovsky PM. The new wheat species Bulletin of Appled Botany (Trudy po Prikladnoi Botanike) 1928;19(2):59-66 (in Russian).

15.	Jakubziner MM. The results of the VIR expedition in Transcaucasia. Plant Industry in USSR, Series A. 1933;5-6:216 (in Russian)

16.	Jakubziner MM. New wheat species. In: Proceedings of the 1st International Wheat Genetics Symposium, August 11-15 1959: 1959; Winnipeg, Canada. Winnipeg Public Press: 207-220.

17.	Zhukovsky PM: Wild emmer in Georgia. Notes of Applied Scientific Departments of Tiflis Botanical Garden 1923;16(3):1-3 (in Russian)

18.	Jakubziner MM. Contribution to the knowledge of wild wheat in Transcaucasia. Bul Appl Bot, Gen Plant Breed 1932;5(1):147-198 (in Russian).

19.	Sachs L. Chromosome behavior in species hybrids with *Triticum timopheevi*. Heredity 1953;7(1):49-58.

20.	Johnson BL, Hall O. Report of a joint expedition through Turkey to collect species of the Triticinae under the auspices of the University of California, U.S.A., Kungl. Skogs-och Lantbruksakademien of Sweden and the Food and Agriculture Organization of the United Nations. Wheat Inf Serv 1967;25:19-29.

21. Johnson BL, Waines JG. Use of wild wheat resources. California Agriculture 1977, September:8-9.

22. Tanaka M, Ishii H. Cytogenetic evidence on the speciation of wild tetraploid wheats collected in Iraq, Turkey, and Iran. In: 4[th] International Wheat Genetics Symposium: 1973; Columbia, Mo. Mo. Agr. Exp. Stn:115-121.

23. Tanaka M, Kawahara T. Wild tetraploid wheats from Northern Iraq cytogenetically closely related to each other. Wheat Inf Serv 1976;43:3.

24. Valkoun J, Giles WJ, Konopka JP: Part 6. Conservation of wild progenitors. Current geographical distribution and habitat of wild wheats and barley. In: Damania AB, Valkoun J, Willcox G, Qualset CO, editors. The Origins of Agriculture and Crop Domestication. Aleppo, Syria: ICARDA; 1998. p. 293-299.

25. Johnson BL. Identification of the apparent B-genome donor of wheat. Can J Genet Cytol 1975; doi: 10.1139/g75-004.

26. Özkan H, Willcox G, Graner A, Salamini F, Kilian B. Geographic distribution and domestication of wild emmer wheat (*Triticum dicoccoides*). Genet Res Crop Evol 2011; doi: 10.1007/s10722-010-9581-5.

27. Troitzky NA. Botanical characteristic of the habitats of wild wheats in Armenia (Erevan). Bul Appl Bot, Genet, Plant Breed 1932;5(1):53-68 (in Russian).

28. Tumanyan MG. Wild eincorns and emmers in Armenia. Bul Appl Bot (Trudy po Prikladnoi Botanike) 1930;24(2):1-14 (in Russian).

29. Dagan J, Zohary D. Wild tetraploid wheats from West Iran cytogenetically identical with Israeli *T. dicoccoides*. Wheat Inform Serv 1970;31:15-7.

30. Tanaka M, Sakamoto S. Morphological and physiological variations in wild tetraploid wheats collected from the Zagros mountains. In: Tanaka M, editor. A preliminary report of studies on the differentiation of tetraploid wheats from different biological levels. vol. 4. Kyoto, Japan:Report of the Germ-plasm Institute; 1979: p. 12-17.

31. Badaeva ED, Badaev NS, Gill BS, Filatenko AA. Intraspecific karyotype divergence in *Triticum araraticum* (Poaceae). Plant Syst Evol 1994; doi: 10.1007/BF00985912.

32. Gill BS, Chen PD. Role of cytoplasm-specific introgression in the evolution of the polyploid wheats. Proc Natl Acad Sci U.S.A. 1987; http://www.pnas.org/cgi/content/abstract/84/19/6800.

33. Jiang J, Gill BS. Different species-specific chromosome translocations in *Triticum timopheevii* and *T. turgidum* support the diphyletic origin of polyploid wheats. Chrom Res 1994; doi: 10.1007/BF01539455.

34. Kawahara T, Tanaka M. Six chromosome types in *Triticum araraticum* Jakubz. different with reciprocal translocations. Jpn J Genet 1977;52(4):261-7.

35. Konarev VG, Gavriliuk EP, Penyova TI, Kovnarev AV, Khakimova AG, Migushova EF. On the nature and origin of wheat genomes by the data of biochemistry and immunochemistry of seed proteins. Agric Biol 1976;11(5):656-65 (in Russian).

36. Jones GH, Valamoti S, Charles M. Early crop diversity: A "new" glume wheat from northern Greece. Veget Hist Archaebot 2000; doi: 10.1007/BF01299798.

37. Dvořák J, Mcguire PE, Cassidy B. Apparent sources of the A genomes of wheats inferred from polymorphism in abundance and restriction fragment length of repeated nucleotide sequences. Genome 1988; doi: 10.1139/g88-115.

38. Kilian B, Ozkan H, Deusch O, Effgen S. Independent wheat B and G genome origins in outcrossing *Aegilops* progenitor haplotypes. Mol Biol Evol 2007; doi: 10.1093/molbev/msl151.

39. Rodríguez S, Maestra B, Perera E, Díez M, Naranjo T. Pairing affinities of the B- and G-genome chromosomes of polyploid wheats with those of *Aegilops speltoides*. Genome 2000; doi:10.1139/gen-43-5-814.

40. Tsunewaki K. Plasmon analysis as the counterpart of genome analysis. In: Jauhar PP, editor. Methods of Genome Analysis in Plant: Their Merrits and Piffals. Boca Ration:CRC Press; 1996; p. 271-299.

41. Makushina EN. New wheat species *Titicum armeniacum* (Jakubz.) sp n. Proc USSR Acad Sci (Doklady Academii Nauk SSSR) 1938;21(7):350-3 (in Russian).

42. Svetozarova BB: On the second genome of *Triticum timopheevii* Zhuk. Proc USSR Acad Sci (Doklady Academii Nauk SSSR) 1939;23(5):472-6 (in Russian).

43. Wagenaar EB. Studies on the genome constitution of *Triticum timopheevi* Zhuk. I. Evidence for genetic control of meiotic irregularities in tetraploid hybrids. Can J Genet Cytol 1961; doi: 10.1139/g61-010.

44. Noda K, Ge K. Structural differentiation of chromosomes between *Triticum dicoccoides* Korn. and *T. araraticum* Jakubz., showing high meiotic pairing homology. Jpn J Genet 1989;64:191-8.

45. Gornicki P, Zhu H, Wang J, Challa GS, Zhang Z, Gill BS, Li W. The chloroplast view of the evolution of polyploid wheat. New Phytol 2014; doi: 10.1111/nph.12931.

46. Huang S, Sirikhachornkit A, Su X, Faris J, Gill B, Haselkorn R, Gornicki P. Genes encoding plastid acetyl-CoA carboxylase and 3-phosphoglycerate kinase of the *Triticum/Aegilops* complex and the evolutionary history of polyploid wheat. Proc Natl Acad Sci U.S.A. 2002; http://www.pnas.org/cgi/content/abstract/99/12/8133.

47. Jiang J, Gill BS. Sequential chromosome banding and in situ hybridization analysis. Genome 1993; doi: 10.1139/g93-104.

48. Chen PD, Gill BS. The origin of chromosome 4A, and genomes B and G of tetraploid wheats. In: Proceedings of the 6th International Wheat Genetics Symposium, November 28 - December 3 1983: 1984; Kyoto, Japan. Plant Germ-Plasm Inst., Kyoto University; p. 39-48.

49. Rodríguez S, Perera E, Maestra B, Díez M, Naranjo T. Chromosome structure of *Triticum timopheevii* relative to *T. turgidum.* Genome 2000; doi:10.1139/gen-43-6-923.

50. Salina EA, Leonova IN, Efremova TT, Roder MS. Wheat genome structure: translocations during the course of polyploidization. Funct Integr Genom 2006;6(1):71-80.

51. Dobrovolskaya O, Sourdille P, Bernard M, Salina E. Chromosome synteny of the a genome of two evolutionary wheat lines. Rus J Genet 2009; doi: 10.1134/S1022795409110118.

52. Hutchinson J, Miller TE, Jahier J, Shepherd KW. Comparison of the chromosomes of *Triticum timopheevi* with related wheats using the techniques of C-banding and in situ hybridization. Theor Appl Genet 1982; doi: 10.1007/BF00303647 .

53. Zhang H, Bian Y, Gou X, Dong Y, Rustgi S, Zhang B, Xu C, Li N, Qi B, Han F et al. Intrinsic karyotype stability and gene copy number variations may have laid the foundation for tetraploid wheat formation. Proc Natl Acad Sci U.S.A. 2013; doi: 10.1073/pnas.1319598110.

54. Badaeva ED, Boguslavsky RL, Badaev NS, Zelenin AV. Intraspecific chromosomal polymorphism of *Triticum araraticum* (Poaceae) detected by C-banding technique. Plant Syst Evol 1990; doi: 10.1007/BF00935980.

55. Kawahara T, Badaeva ED, Badaev NS, Gill BS. Spontaneous translocations in *Triticum araraticum* Jakubz. Wheat Inf Serv 1996;83:7-14.

56. Bernhardt N, Brassac J, Kilian B, Blattner FR. Dated tribe-wide whole chloroplast genome phylogeny indicates recurrent hybridizations within Triticeae. BMC Evol Biol 2017; doi: 10.1186/s12862-017-0989-9.

57. Avni R, Nave M, Barad O, Baruch K, Twardziok SO, Gundlach H, Hale I, Mascher M, Spannagl M, Wiebe K et al. Wild emmer genome architecture and diversity elucidate wheat evolution and domestication. Science (New York) 2017; doi: 10.1126/science.aan0032.

58. El Haddad N, Kabbaj H, Zaïm M, El Hassouni K, Tidiane Sall A, Azouz M, Ortiz R, Baum M, Amri A, Gamba F et al. Crop wild relatives in durum wheat breeding: Drift or thrift? Crop Sci 2021; https://doi.org/10.1002/csc2.20223.

59. Allard RW, Shands RG. Inheritance of resistance to stem rust and powdery mildew in cytologically stable spring wheats derived from *Triticum timopheevi*. Phytopathol 1954; 44(2):266-74.

60. Brown-Guedira G, Badaeva ED, Gill BS, Cox TS. Chromosome substitutions of *Triticum timopheevii* in common wheat and some observations on the evolution of polyploid wheat species. Theor Appl Genet 1996; doi: 10.1007/BF00223462.

61. Brown-Guedira GL, Singh S, Fritz AK. Performance and mapping of leaf rust resistance transferred to wheat from *Triticum timopheevii* subsp. *armeniacum*. Phytopathol 2003; doi: 10.1094/PHYTO.2003.93.7.784

62. Dyck PL. Transfer of a gene for stem rust resistance from *Triticum araraticum* to hexaploid wheat. Genome 1992; doi: 10.1139/g92-120.

63. McIntosh RA, Gyarfas J. *Triticum timopheevii* as a source of resistance to wheat stem rust. Z Pflanzenzucht 1971;66(3):240-8.

64. Maan SS, Lucken KA. Interacting male sterility-male fertility restoration systems for hybrid wheat research. Crop Sci 1972; doi: 10.2135/cropsci1972.0011183X001200030031x.

65. Mikó P, Megyeri M, Farkas A, Molnár I, Molnár-Láng M. Molecular cytogenetic identification and phenotypic description of a new synthetic amphiploid, *Triticum timococcum* ($A^tA^tGGA^mA^m$). Genet Res Crop Evol 2015; doi: 10.1007/s10722-014-0135-0.

66. Würschum T, Leiser WL, Weissmann S, Maurer HP. Genetic architecture of male fertility restoration of *Triticum timopheevii*. Theor Appl Genet 2017; doi: 10.1007/s00122-017-2885-5 .

67. Martynov SP, Dobrotvorskaya TV, Krupnov VA. Analysis of the distribution of *Triticum timopheevii* Zhuk. genetic material in common wheat varieties (*Triticum aestivum* L.). Rus J Genet 2018; doi: 10.1134/S1022795418020126 .

68. Konovalov F, Goncharov N, Goryunova S, Shaturova A, Proshlyakova T, Kudryavtsev A. Molecular markers based on LTR retrotransposons *BARE-1* and *Jeli* uncover different strata of evolutionary relationships in diploid wheats. Mol Genet Genom 2010; doi: 10.1007/s00438-010-0539-2.

69. Badaeva ED, Keilwagen J, Knüpffer H, Waßermann L, Dedkova OS, Mitrofanova OP, Kovaleva ON, Liapunova OA, Pukhalskiy VA, Özkan H et al. Chromosomal passports

provide new insights into diffusion of emmer wheat. PLoS ONE 2015; doi: 10.1371/journal.pone.0128556 10(5):e0128556.

70. Kawahara T, Tanaka M. Intraspecific differentiation in chromosome structure in the wild tetraploid wheats. Wheat Inf Serv 1981; 52:33.

71. Badaeva ED, Ruban AS, Zoshchuk SA, Surzhikov SA, Knüpffer H, Kilian B. Molecular cytogenetic characterization of *Triticum timopheevii* chromosomes provides new insight on genome evolution of *T. zhukovskyi*. Plant Syst Evol 2016; doi: 10.1007/s00606-016-1309-3.

72. Kimber G, Feldman M. Wild wheat, an Introduction, Special Report 353, . Columbia, Mo, U.S.A.:College of Agriculture University of Missouri; 1987.

73. Mori N, Kondo Y, Ishii T, Kawahara T, Valkoun J, Nakamura C. Genetic diversity and origin of timopheevi wheat inferred by chloroplast DNA fingerprinting. Breed Sci 2009; doi: 10.1270/jsbbs.59.571 .

74. Shcherban AB, Schichkina AA, Salina EA. The occurrence of spring forms in tetraploid Timopheevi wheat is associated with variation in the first intron of the *VRN-A1* gene. BMC Plant Biol 2016; doi: 10.1186/s12870-016-0925-y.

75. Bernhardt N, Brassac J, Dong X, Willing E-M, Poskar CH, Kilian B, Blattner FR. Genome-wide sequence information reveals recurrent hybridization among diploid wheat wild relatives. Plant J 2019; doi: 10.1111/tpj.14641.

76. Qi L, Friebe B, Gill BS. Complex genome rearrangements reveal evolutionary dynamics of pericentromeric regions in the Triticeae. Genome 2006; doi: 10.1139/G06-123 .

77. Abbott RJ, Hegarty MJ, Hiscock SJ, Brennan AC. Homoploid hybrid speciation in action. TAXON 2010; doi: 10.1002/tax.595005.

78. Nieto Feliner G, Álvarez I, Fuertes-Aguilar J, Heuertz M, Marques I, Moharrek F, Piñeiro R, Riina R, Rosselló JA, Soltis PS et al. Is homoploid hybrid speciation that rare? An empiricist's view. Heredity 2017; doi: 10.1038/hdy.2017.7.

79. Soltis PS, Soltis DE. The role of hybridization in plant speciation. Ann Rev Plant Biol 2009; doi:10.1146/annurev.arplant.043008.092039.

80. Civáň P, Ivaničová Z, Brown TA. Reticulated origin of domesticated emmer wheat supports a dynamic model for the emergence of agriculture in the Fertile Crescent. PLoS ONE 2013; doi: 10.1371/journal.pone.0081955.

81. Nesbitt M, Samuel D. From staple crop to extinction? The archaeology and history of the hulled wheats In: Padulosi S, Hammer K, Heller J, editors. Hulled Wheat. Rome, Italy: IPGRI; 1996; p. 41-100.

82. Szabo AT, Hammer K. Notes on the taxonomy of farro: *Triticum monococcum, T. dicoccon* and *T. spelta*. In: Padulosi S, Hammer K, Heller J, editors. Hulled Wheat. Rome, Italy: IPGRI; 1996: 2-40.

83. Zaharieva M, Ayana N, Hakimi A, Misra S, Monneveux P. Cultivated emmer wheat (*Triticum dicoccon* Schrank), an old crop with promising future: a review. Genet Res Crop Evol 2010; doi: 10.1007/s10722-010-9572-6.

84. Maccaferri M, Harris NS, Twardziok SO, Pasam RK, Gundlach H, Spannagl M, Ormanbekova D, Lux T, Prade VM, Milner SG et al. Durum wheat genome highlights past domestication signatures and future improvement targets. Nat Genet 2019; doi: 10.1038/s41588-019-0381-3.

85. Pont C, Leroy T, Seidel M, Tondelli A, Duchemin W, Armisen D, Lang D, Bustos-Korts D, Goué N, Balfourier F et al. Tracing the ancestry of modern bread wheats. Nat Genet 2019; doi: 10.1038/s41588-019-0393-z.

86. Dekaprelevich LL, Menabde VL. Hulled wheat in Western Georgia. Bull Appl Bot, Genet Plant Breed 1932;5(1):1-46 (in Russian).

87. Stoletova EA: Emmer (An assay of study of one of the vanishing crops) Bull Appl Bot, Plant Breed 1924-25;14(1):27-111 (in Russian).

88. Feldman M. Origin of cultivated wheat. In: Bonjean AP, Angus WJ, editors. The world wheat book - A history of wheat breeding. Londres-Paris-New York:Tec & Doc/Intersept Ltd.; 2001; p. 3-56.

89. Bieniek A. Archaeobotanical analysis of some early Neolithic settlements in the Kujawy region, central Poland, with potential plant gathering activities emphasised. Veg Hist Archaeobot. 2002; doi: 10.1007/s003340200004.

90. Bieniek A. Neolithic plant husbandry in the Kujawy region of central Poland. In: Colledge S, Conolly J editors. The Origin and Spread of Domestic Plants in Southwest Asia and Europe. Walnut Creek: Left Coast Press. 2007; p. 327–342.

91. Bogaard A, Bending J, Jones G. Archaeobotanical evidence for plant husbandry and use. In: Whittle AWR, editor. The Early Neolithic on the Great Hungarian Plain Investigations of the Körös culture site of Ecsegfalva 23, County Békés. vol. 21. Budapest:Institute of Archaeology, Hungarian Academy of Sciences. 2007; p. 421–445.

92. Bogaard A, Charles M, Livarda A, Ergun M, Filipovic D, Jones G. The archaeobotany of mid-later Neolithic occupation levels at Çatalhöyük. In: Hodder I, editor. Humans and Landscapes of Çatalhöyük: Reports from the 2000-2008 seasons. Los Angeles: Monographs of the Cotsen Institute of Archaeology, University of California at Los Angeles; 2013; p. 93-128.

93.	Fairbairn A, Asouti E, Near J, Martinoli D. Macro-botanical evidence for plant use at Neolithic Çatalhöyük south-central Anatolia, Turkey. Veget Hist Archaeobot 2002; doi: 10.1007/s00334.

94.	Fischer E, Rösch A. Archäobotanische Untersuchungen. In: Schier W, Draşovean F, editors. Praehistorische Zeitschrift Vorbericht über die rumänisch-deutschen Prospektionen und Ausgrabungen in der befestigten Tellsiedlung von Uivar, jud Timiş, Rumänien (1998–2002). vol. 79; 2004; p. 209–220.

95.	Hajnalová M. Early farming in Slovakia: an archaeobotanical perspective. In: Colledge S, Connolly J, editors. The origins and spread of domestic plants in Southwest Asia and Europe. Walnut Creek:Left Coastal Press. 2007; p. 295–314.

96.	Kohler-Schneider M. Contents of a storage pit from late Bronze Age Stillfried, Austria: another record of the "new" glume wheat. Veget Hist Archaeobot 2003; 10.1007/s00334-003-0010-y.

97.	Kreuz A, Boenke N. The presence of two-grained einkorn at the time of the Bandkeramik culture. Veget Hist Archaeobot 2002; 10.1007/s003340200026.

98.	Rottoli M, Pessina A. Neolithic agriculture in Italy. In: Colledge S, Conolly J, editors. The origin and spread of domestic plants in South-west Asia and Europe. Walnut Creek: Left Coast Press. 2007; p. 141-154.

99.	Ulaş B, Fiorentino G. Recent attestations of "new" glume wheat in Turkey: a reassessment of its role in the reconstruction of Neolithic agriculture. Veget Hist Archaeobot 2020; doi: 10.1007/s00334-020-00807-w.

100.	Valamoti SM, Kotsakis K. Transitions to agriculture in the Aegean: the archaeobotanical evidence. In: Colledge S, Conolly J, editors. The origins and spread of domestic plants in Southwest Asia and Europe. Walnut Creek:Left Coast Press. 2007; p. 75-92.

101.	Borojevic K. Emmer aus Feudvar. In: Hänsel B, Medović P, editors. Vorbericht über die jugoslawisch-deutschen Ausgrabungen in der Siedlung von Feudvar bei Mosorin (Gem Titel, Vojvodina) von 1986–1990: Bronzezeit–Vorrömische Eisenzeit. Darmstadt:Verlag Philipp von Zabern. 1991; p. 171–177.

102.	de Moulins D. Agricultural changes at Euphrates and steppe sites in the mid-8[th] to the 6[th] millennium B.C. Oxford.: Archaeopress; 1997.

103.	Kroll H. Die Pflanzenfunde von Feudvar. In: Kroll H, Reed R editors. Feudvar III: Die Archäobotanik. Würzberg:Würzberg University Press. 2016; p. 37-194.

104.	Kenéz Á, Pető Á, Gyulai F. Evidence of 'new glume wheat' from the Late Neolithic (Copper Age) of south-eastern Hungary (4th millennium cal. b.c.). Veget Hist Archaeobot 2014; doi: 10.1007/s00334-013-0405-3 .

105. Bogaard A, Filipović D, Fairbairn A, Green L, Stroud E, Fuller D, Charles M. Agricultural innovation and resilience in a long-lived early farming community: the 1,500-year sequence at Neolithic to early Chalcolithic Çatalhöyük, central Anatolia. Anatolian Studies 2017; doi: 10.1017/S0066154617000072.

106. Ergun M. "Where the wild things are". Contextual insights into wild plant exploitation at Aceramic Neolithic Aşıklı Höyük, Turkey. Paléorient 2018; doi: 10.2307/26595372.

107. Perego R. Contribution to the development of the Bronze Age plant economy in the surrounding of the Alps: an archaeobotanical case study of two Early and Middle Bronze Age sites in northern Italy (Lake Garda region). Switzerland: University of Basel; 2017.

108. Toulemonde F, Durand F, Berrio L, Bonnaire E, Daoulas G, Wiethold J. Records of "new" glume wheat in France: a review. Veget Hist Archaeobot 2015; doi: 10.1007/s00334-014-0479-6.

109. Czajkowska BI, Bogaard A, Charles M, Jones G, Kohler-Schneider M, Mueller-Bieniek A, Brown TA. Ancient DNA typing indicates that the "new" glume wheat of early Eurasian agriculture is a cultivated member of the *Triticum timopheevii* group. J Archaeol Sci 2020; doi: 10.1016/j.jas.2020.105258.

110. Vavilov NI. Botaniko-geograficheskie osnovy selektsii [Botanical-geographical foundations of breeding]. In: Teoreticheskie osnovy selektsii rastenii [Theoretical foundations of plant breeding]. vol. 1. Moscow,Leningrad; 1935; p. 17-75.

111. Jorjadze M, Berishvili T, Shatberashvili E. The ancient wheats of Georgia and their traditional use in the southern part of the country. Emir J Food Agric 2014;26(2):192-202.

112. Mosulishvili M, Bedoshvili D, Maisaia I. A consolidated list of Triticum species and varieties of Georgia to promote repatriation of local diversity from foreign genebanks. Ann Agrar Sci 2017; doi: 10.1016/j.aasci.2017.02.006 .

113. Schiemann E. Erwin Baur. Berichte der Deutschen Botanischen Gesellschaft 1934; 52(2):51-114.

114. Zencirci N, Yılmaz H, Garaybayova N, Karagöz A, Kilian B, Özkan H, Hammer K, Knüpffer H. Mirza (Hacızade) Gökgöl (1897–1981): the great explorer of wheat genetic resources in Turkey. Genet Res Crop Evol 2018; doi: 10.1007/s10722-018-0606-9.

115. Dekaprelevich LL, Menabde VL. Regarding the investigation of crop plants in western Georgia. Zap Nauchno-prikl otdelov Tiflisskogo Bot Sada 1929;6:219–54 (in Russian).

116. Menabde VL. The Georgian wheats. Tbilisi: Izdatelstvo Akademii Nauk Gruzinskoy SSR; 1948 (in Georgian).

117. Dekaprelevich LL. Species, varieties and cultivars of Georgian wheat. Tr Inst Polevodstva AN GSSR 1954;8:3-61 (in Georgian).

118. Fick SE, Hijmans RJ: WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. Int J Climatol 2017; doi: 10.1002/joc.5086.

119. Team RC: R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria; 2016.

120. Davis PH. Flora of Turkey and the East Aegean Islands. Edinburgh:Edinburgh University Press; 1965-1988.

121. Hanelt P. Mansfeld's encyclopedia of agricultural and horticultural crops. In: Hanelt P editor. Mansfeld's encyclopedia of agricultural and horticultural crops. 2001.

122. Cauvin J, Aurenche O, Cauvin M-C, Balkan-Atlı N. The Pre-Pottery site of Cafer Höyük. In: Özdogan M, Başgelen N, Kuniholm P, editors. The Neolithic in Turkey. vol. 1. Istanbul: Archaeology and Art Publications. 2011; p. 1-40.

123. Quade J, Stiner MC, Copeland A, Clarke AE, Özbaşaran M. Summary of carbon-14 dating of the cultural levels of Aşıklı Hoyük. In: Özbaşaran M, Duru G, Stiner MC, editors. The early settlement at Aşıklı Hoyük: Essays in honor of Ufuk Esin. Istanbul: Ege Yayinlari. 2018; p. 43–56.

124. Pourkheirandish M, Dai F, Sakuma S, Kanamori H, Distelfeld A, Willcox G, Kawahara T, Matsumoto T, Kilian B, Komatsuda T. On the origin of the non-brittle rachis trait of domesticated einkorn wheat. Front Plant Sci 2018; doi: 10.3389/fpls.2017.02031.

125. Pourkheirandish M, Hensel G, Kilian B, Senthil N, Chen G, Sameri M, Azhaguvel P, Sakuma S, Dhanagond S, Sharma R et al. Evolution of the grain dispersal system in barley. Cell 2015; doi: 10.1016/j.cell.2015.07.002.

126. Civáň P, Brown TA. A novel mutation conferring the nonbrittle phenotype of cultivated barley. New Phytol 2017; doi: 10.1111/nph.14377 .

127. Jakob SS, Rödder D, Engler JO, Shaaf S, Özkan H, Blattner FR, Kilian B. Evolutionary history of wild barley (*Hordeum vulgare* subsp. *spontaneum*) analyzed using multilocus sequence data and paleodistribution modeling. Gen Biol Evol 2014; doi: 10.1093/gbe/evu047.

128. Oliveira HR, Jacocks L, Czajkowska BI, Kennedy SL, Brown TA. Multiregional origins of the domesticated tetraploid wheats. PLos ONE 2020; doi: 10.1371/journal.pone.0227148.

129. Poets A, Fang Z, Clegg M, Morrell P. Barley landraces are characterized by geographically heterogeneous genomic origins. Gen Biol 2015; doi: 10.1186/s13059-015-0712-3.

130. Knüpffer H. Triticeae genetic resources in ex situ genebank collections. In: Muehlbauer GJ, Feuillet C, editors. Genetics and Genomics of the Triticeae. vol. 7, part 1 (Genetics of the Triticeae), First edition. New York: Springer. 2009; p. 31-79.

131. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. Mol Biol Evol 2006; doi: 10.1093/molbev/msj030.

132. Dice LR. Measures of the amount of ecologic association between species. Ecology 1945; doi: 10.2307/1932409.

133. Zabeau M, Vos P. Selective restriction fragment amplification: a general method for DNA fingerprinting. European patent application number 92402629.7; publication number 0534858 Al; 1993.

134. Altıntaş S, Toklu F, Kafkas S, Kilian B, Brandolini A, Özkan H. Estimating genetic diversity in durum and bread wheat cultivars from Turkey using AFLP and SAMPL markers. Plant Breed 2008; doi:10.1111/j.1439-0523.2007.01424.x.

135. Alsaleh A, Baloch FS, Derya M, Azrak M, Kilian B, Özkan H, Nachit M. Genetic linkage map of Anatolian durum wheat derived from a cross of Kunduru-1149 × Cham1. Plant Mol Biol Rep 2015; doi: 10.1007/s11105-014-0749-6.

136. Jaccard P. Nouvelles recherches sur la distribution florale. Bulletin de la Societe Vaudoise des Sciences Naturelles 1908;44(163):223-270.

137. Perrier X, Flori A, Bonnot F. Data analysis methods. In: Hamon P, Seguin M, Perrier X, Glaszmann JC, editors. Genetic diversity of cultivated tropical plants. Montpellier: Enfield, Science Publishers. 2003; p. 43-76.

138. Peakall R, Smouse PE. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research – an update. Bioinformatics 2012; doi: doi:10.1093/bioinformatics/bts460.

139. Badaeva ED, Budashkina EB, Badaev NS, Kalinina NP, Shkutina FM. General features of chromosome substitutions in *Triticum aestivum x T. timopheevii* hybrids. Theoretical and Applied Genetics 1991; doi: 10.1007/BF00226218.

140. Mitrofanova O, Badaeva E, Salina EA. *T. timopheevii* and *T. zhukovskii,* the bread wheat cousins and their contribution to *T. aestivum* improvement. In: Bonjean AP, Angus WJ, VanGinkel M, editors. The World Wheat Book: A History of Wheat Breeding (v. 3). France:Doc Lavoisier; 2016.

141. Salina EA, Pestsova EG, Adonina IG, Vershinin AV. Identification of a new family of tandem repeats in Triticeae genomes. Euphytica 1998; doi: 10.1023/A:1018360324242.

142.	Salina EA, Numerova OM, Ozkan H, Feldman M. Alterations in subtelomeric tandem repeats during early stages of allopolyploidy in wheat. Genome 2004; doi: 10.1139/g04-044 .

143.	Badaeva ED, Amosova AV, Goncharov NP, Macas J, Ruban AS, Grechishnikova IV, Zoshchuk SA, Houben A. A set of cytogenetic markers allows the precise identification of all A-genome chromosomes in diploid and polyploid wheat. Cytogenet Genome Res 2015; doi: 10.1159/000433458.

144.	Bedbrook RJ, Jones J, O'Dell M, Thompson RJ, Flavell RB. A molecular description of telomeric heterochromatin in *Secale* species. Cell 1980; doi: 10.1016/0092-8674(80)90529-2.

145.	Pedersen C, Rasmussen SK, Linde-Laursen I. Genome and chromosome identification in cultivated barley and related species of the Triticeae (Poaceae) by in situ hybridization with the GAA-satellite sequence. Genome 1996; doi: 10.1139/g96-013.

146.	Komuro S, Endo R, Shikata K, Kato A. Genomic and chromosomal distribution patterns of various repeated DNA sequences in wheat revealed by a fluorescence in situ hybridization procedure. Genome 2013; doi: 10.1139/gen-2013-0003.

🔵	*T. dicoccoides*
🟢	*T. araraticum*, ARA-0
🔴	*T. araraticum*, ARA-1
🔵	*T. timopheevii*

DIC	TIM	ARA-1	←— Turkey →← Iran	←— Iraq	Caucasus →|

1

2 — T2A'L:6A'L — T2A'L:7GS

3

4 — T4A'S:7A'S

5

6 — T6A'S:2A'S

7 — inv7A'-1 inv7A'-2 — T7A'L:4A'L

a b c d e f g h i j k l m n o p q r s t

A^t genome

DIC	TIM	ARA-1	←— Turkey →← Iran	←— Iraq	Caucasus →|

1 — T1GL:5GS

2 — T2GL:6GL

3

4 — T5GL:1GS — inv4G — T4GS:2A'S

5

6 — T6GS:2GS

7 — T7GL:4GL

a b c d e f g h i j k l m n o p q r s t

G-genome

permanent, invariable C-bands

hybridization sites of pAesp_SAT86

permanent C-bands, variable in size

hybridization sites of pSc119.2

inconsistent C-bands

overlaping sites

C-bands/ pAesp_SAT86 sites typical for ARA-1 group

C-bands/ pAesp_SAT86 sites typical for ARA-0 group