(Subclinical mastitis alters dairy cow behavior and salivary SAA)

Changes in Behavior and Salivary Serum Amyloid A in cows with Subclinical Mastitis

G. Caplen[1*], S.D.E. Held[1]

gina.caplen@bristol.ac.uk; suzanne.held@bristol.ac.uk

[1]Animal Welfare and Behavior Group, Bristol Veterinary School, University of Bristol,

Langford, Bristol, BS40 5DU, UK

*Corresponding author

ABSTRACT

The aim of this study was to identify detailed changes in behavior, and in salivary serum

amyloid A (**SAA**), associated with subclinical mastitis. This included standard sickness

behaviors (which form part of an adaptive response to conserve energy, minimize heat loss

and facilitate recovery following infection and inflammation) and lesser-studied behaviors,

that are biologically predicted to change at low-level infection, and therefore particularly

relevant for detecting subclinical mastitis (here labelled 'luxury' behaviors). SAA is a

physiological marker of systemic inflammation, with levels in milk and serum already known

to increase during subclinical mastitis. We investigated whether the same was true for SAA

in cow saliva. Data were collected for n = 34 commercial barn-housed dairy cows, tested in

17 matched-pairs. Each pair comprised a cow with subclinical mastitis (**SCM)** and a healthy

control (**CTRL**), identified using somatic cell count (**SCC**) (SCM: SCC >200 x1000 cells/ml;

CTRL: SCC <100 x1000 cells/ml). SCM cows were selected for study ad-hoc, at which point

they were paired with a CTRL cow, based upon parity and calving date; consequently, the

full data set was accrued over several months. Data were collected for each pair over 3 days:

SCC (14:00-15:00h) Day 1; behavior (24h from 00:00) Day 2; salivary serum amyloid-A

(SAA) Day 3. We report, for the first time, that an increase in salivary SAA occurs during

subclinical mastitis; SAA was higher in SCM cows and demonstrated a positive (weak)

correlation with SCC. The behavioral comparisons revealed that SCM cows had reductions in

activity (behavioral transitions and distance moved), social exploration, social reactivity

(here: likelihood to move away/be displaced following receipt of agonism), performance of

social grooming and head butts, and the receipt of agonistic non-contact challenges. In

addition, SCM cows received more head swipes, and spent a greater proportion of time lying

with their head on their flank than CTRL cows. SCM cows also displayed an altered feeding

pattern; they spent a greater proportion of feeding time in direct contact with two

conspecifics, and a lower proportion of feeding time at self-locking feed barriers, than CTRL

cows. Behavioral measures were found to correlate, albeit loosely, with serum SAA in a

direction consistent with predictions for sickness behavior. These included positive

correlations with both lying duration and the receipt of all agonistic behavior, and negative

correlations with feeding, drinking, the performance of all social and all agonistic behavior,

and social reactivity. We conclude that changes in salivary SAA, social behavior, and activity

offer potential in the detection of subclinical mastitis and recommend further investigation to

substantiate and refine our findings.

Keywords: Cows; SAA; Sickness; Behavior; Subclinical mastitis

1.   INTRODUCTION

Clinical stages of infectious disease are typically easily identified by obvious physical

symptoms and behavioral changes (so-called 'sickness behaviors'; Hart, 1988). Subclinical

infection, that is infection below the level of clinical detection, by definition, is more difficult

to spot. However, based on the interactions between the immune and central nervous system

that cause sickness behavior predictions can also be made about behavioral changes during

subclinical infection (Dantzer, 2004). These behavioral changes can be used as early warning

signs of disease (Weary et al. 2009; von Keyserlingk et al. 2010), and/or to identify chronic

subclinical infection levels. Mastitis remains a major concern in dairy cows with serious

negative effects on welfare and productivity (Petersson-Wolfe et al. 2018). At subclinical

levels, inflammation is present in response to the infection and milk production drops, but no

abnormalities in the gland or milk are visible (Sordillo et al., 1997). It is therefore important

for infection at any level to be identified and treated as soon as possible. Our main aim was to

identify such changes in cows based on a detailed behavioral comparison between individuals

with spontaneously occurring subclinical mastitis and healthy controls.

Behaviors in healthy animals can be divided, on the basis of immediate survival benefits, into

'core maintenance' and 'luxury' (after e.g. Dawkins, 1990). Core behaviors have immediate,

short-term survival benefits; examples include resting, feeding, and drinking. They, therefore,

start to decline only at later disease stages (e.g. Littin et al., 2008) and have low sensitivity

during early and low-level stages (e.g. Sepúlveda-Varas et al., 2014). In our comparison we

included standard 'core' sickness behaviors, such as changes to feeding, previously shown to

be associated with experimentally-induced and spontaneously-occurring clinical mastitis

(Siivonen et al., 2011; Fogsgaard et al. 2012; Sepúlveda-Varas et al., 2016). Because of our

focus on subclinical mastitis, we also included so-called 'luxury' behaviors, such as physical

and social exploration, grooming and various social interactions. These 'luxury' behaviors

have delayed, longer-term, benefits and are not essential for immediate survival, meaning that

they are biologically predicted to change at lower levels of disease than other sickness

behaviors due to a greater sensitivity to disease challenge (Littin et al. 2008; Weary et al.

2009), especially when energy resources are diverted to fighting infection (see also 'low-

resilience behaviors' in Littin et al., 2008).

In cows, effective health monitoring is hampered by logistical difficulties associated with

direct animal observations within large open barns, reductions in human interaction linked

with the substantial uptake in robotic milking units, and the tendency of cattle (for several

biological reasons) to display only subtle indicators of pain/weakness (Gleerup et al., 2015).

Advances in image analysis now allow automated recognition of individuals within a herd

(Andrew et al., 2020), and accurate identification of health-related abnormal behaviors, e.g.

foot disease (Gu et al., 2017). The identification of behaviors associated with subclinical

disease (prior to the development of, or without, clinical symptoms) may therefore find

application in future diagnostic software algorithms targeted at early disease monitoring in

dairy cows (see e.g. Wagner et al. 2020).

A second aim was to quantify levels of salivary serum amyloid A (**SAA**), a major acute phase

protein (**APP**) in cows (Murata et al., 2004), with and without spontaneously occurring

subclinical mastitis. APPs are non-specific inflammatory markers that fluctuate in response to

infection. Increased SAA levels have been detected during subclinical mastitis within both

serum and milk (Kovac et al., 2011; Kovacevi-Filipovic et al., 2012). Other studies have

confirmed SAA presence in bovine saliva (Lecchi et al., 2012; Rahman et al., 2013),

suggesting salivary SAA has the potential for use in non-invasive detection of infectious

diseases, such as mastitis.

Raised systemic inflammation levels, whether during acute infection or as chronic

inflammatory states, are predicted to be accompanied by a feeling of sickness or 'malaise' in

animals, as they are in humans (Dantzer et al., 2008; Weary et al. 2009). De Boyer Des

Roches et al. (2017) reported correlations between (serum) SAA and several pain indicators

before and after experimentally induced intra-mammary challenge with *E. coli*, including

behavioral measures of attentiveness to surroundings. This suggested a direct link between

serum SAA, infection levels and some sickness behavior in mastitic cows. A non-invasive

means of monitoring systemic inflammation would also benefit future investigations into the

impact of spontaneously occurring infections, such as mastitis, on cow welfare. In our study

we tested for associations between salivary SAA levels and SCC (a standard measure of

mastitis severity), and for associations between these physiological measures and behavior.

In summary, the purpose of this study was to identify differences in behavior and salivary

SAA associated with subclinical mastitis. To this end we: (a) compared the behavior and

salivary SAA of cows with subclinical mastitis with that of matched healthy individuals and,

(b) correlated behavioral variables with both SCC and salivary SAA. We predicted that

salivary SAA would be higher in cows with subclinical mastitis than in matched healthy

controls. We also predicted that luxury behaviors, here including a range of social behaviors,

would decrease with subclinical mastitis, and that differences detected would correlate

negatively with the physiological measures (SCC and salivary SAA).

2.   MATERIALS AND METHODS

2.1.   Ethics Statement

The study was conducted between October 2017 and February 2018 at Bristol Veterinary

School dairy farm. The experimental procedures were approved by the Animal Welfare and

Ethical Review Board at the University of Bristol and conducted under University

Investigation Number UB/17/061 'Behavioural markers of subclinical disease in dairy cows'.

2.2.   Animals

Focal cows (n = 34) were part of an indoor commercial Holstein-Friesian dairy herd (n =

200) and resided within the low milk-yield group (approx. n = 80 cows) at the time of the

study; having been part of the group for at least one month prior to data collection they were

well-established within the social dominance hierarchy. Low yield animals were studied as

they were deemed to be of low risk for having, or developing, subclinical metritis or ketosis

during the trial. Cows were housed within a free-stall barn containing 93 lying cubicles (1.2 x

2.4m) with sand bedding, three stainless steel tip-over drinking troughs, a swinging brush

(DeLaval), and automatic floor scrapers. Figure 1 shows the lay-out and relative positioning

of resources within the pen. Cows were milked three times daily (at 06:00, 14:00 and 22:00h)

and fed a total mixed ration once daily (06:00h).

Only clinically healthy non-lame cows (mobility score ≤1; AHDB, 2015) without physical

symptoms of mastitis were selected for inclusion in the study. Clinical health was assessed

using individual health records and visual inspection of each cow by the researcher and

herdsman. Data were collected concurrently from cows in matched pairs, each comprising a

cow with subclinical mastitis (**SCM**) and a clinically healthy control (**CTRL**). Matching was

performed on the basis of the following potential confounds: parity, pregnancy status (yes/no)

and stage of pregnancy (days since insemination). Cows with a SCC of >200 (x1000 cells/ml)

were classified as SCM (Madouasse et al., 2010), while cows with a SCC of <100 (x1000

cells/ml) were classified as CTRL. The 34 focal cows comprised 24 pregnant and 10 barren

ones, 20 primiparous (n = 20) and 14 multiparous ones (of which two cows were at their 2nd

parity; two at 3rd; six at 4th; four at 5th). The time to expected calving date ranged from 55 -

236 days. While data from the two cows in a matched pair were collected at the same time,

the data for 17 pairs took several months to collect due to low mastitis incidence within the

herd at the time of the study.

INSERT FIGURE 1 NEAR HERE

2.3. Somatic Cell Count

Composite quarter milk samples were collected between 14:00-15:00h on Day 1. Although

pairs of cows had data collected at different dates, for the purpose of our experimental design,

all cows had SCC data collected on Day 1 and behavioral data on Day 2. Somatic cells were

manually counted using a standard direct microscopic methodology (ISO 13366-1, 1997)

following staining with Newman-Lampert stain solution: Levowitz-Weber modification

(Newman's Stain Solution: modified, 01375, Sigma-Aldrich).

2.4. Behavioral Measures

Each focal cow was fitted with a colored collar to facilitate individual recognition on Day 1.

Two CCTV systems (N441L1T, Annke®, CA 91748, US), including six cameras recorded

video footage from the entire low-yield pen. Continuous behavioral data were then coded

retrospectively from video for each focal cow for 24h starting from 00:01h of Day 2 by a

single experienced coder who was unaware of the health status of the cows at the time of

scoring; these data made up the 24 h ('**24h**') data set. A second behavior data set was

compiled for each cow using video recordings of the first 60 min following morning milking

('**1hPostM1'**). This hour coincided with the peak feeding time of the day, since fresh feed

was delivered to the feed corridor while the cows were in the parlour at morning milking. The

start time of the 60 min observation period was specific to each focal cow, starting

immediately following that cow's re-entrance into the home pen from the parlour exit.

All behavioural measures are described in Table 1. Three broad categories of behavior were

of interest: 'core' maintenance (including lying, feeding/drinking and activity), non-social

'luxury' behaviors (such as self-grooming and exploring the physical environment), and

social 'luxury' behaviors. The last category was the most detailed, comprising non-agonistic

(e.g. allogrooming, social exploration) and agonistic (e.g. head swipes, head butts) behaviors,

in addition to social reactivity (here defined as the likelihood of moving away/being

displaced following the receipt of an agonistic behavior).

Total behavioral transitions ('**Trans**'), a measure of activity, was calculated using all

behaviors, including those coded, but not analysed individually (i.e. not listed in Table 1).

These additional non-focal behaviors included eat sand, lick salt, paw sand, run, shake head,

stand, and walk. Proximity was investigated using nearest neighbour ('NN') scores. When the

focal cow was located at the feed barrier or resting in a cubicle, the number of other cows in

immediate 'proximity' were scored as 0NN, 1NN or 2NN (Table 1). To enable an estimation

of distance moved ('**Dist**'), the pen floorspace was hypothetically subdivided into 29 units

(each 4.8 m wide; see Figure 1) and the location of each cow was noted every five minutes

throughout the 24h period. The number of floor units crossed during these intervals was then

used to estimate distance moved.

Data were not available for 06:00, 14:00 or 22:00h as the cows were in the milking parlour or

collecting yard during these periods. To account for differences in total time visible (i.e. due

to variations in time spent within the parlour) data in the '24h' data sets were standardised to

either: number per hour visible (behavioral events) or seconds per hour visible (behavioral

states); e.g. to standardise data from a cow that was visible for 20h, 42min and 12s within a

24h period data this data would be divided by 20.703.

Table 1: Cow behavioral measures used in the study

Measure (unit)	Abbreviation – Definition
(A.) **Maintenance** (Core)	
Lie (s)	'**Lie**'[a]: horizontal recumbent resting position, with abdomen in contact with the floor.
Head on flank (s)	'**HOF**'[a]: lying with the head held in contact with the flank, pointing backwards towards the rump. Associated with active sleep.
Drink (s)	'**Drink**'[a]: standing with muzzle placed within a water trough.
Feed (s)	'**Feed**'[a]: ingesting or chewing of food while standing with the head crossing the feed barrier.
Activity: Transitions (n)	'**Trans**'[a]: the total number of changes in behavior a focal cow undergoes during the observation period.
Activity: Distance moved (units)	'**Dist**'[a]: the number of units of floor space crossed by the focal cow during the observation period.
(B.) **Non-social** (Luxury)	
Brush use (s)	'**Brush**'[a]: contact made between any body part and the mechanical brush (while standing).
Self-groom (s)	'**Self-groom**'[a] = 'Lick self' (lick own body part) + 'Rub self' (rub own body part against pen furniture) + 'Scratch self' (using own foot to scratch own body part).
Explore environment (s)	'**ExpEnv**'[a] = 'Explore food' (holding nose close to food while sniffing or moving/flicking nose around within food, without any obvious active ingestion) + 'Explore pen' (licking or holding nose very close to any part of the barn structure or pen furniture) + 'Explore sand' (holding nose close to sand bedding or flicking sand with nose).
(C.1) **Social** (Luxury)**:** **agonistic interactions and displacement**	Each behavior was further classified as: given (G), received (R), or received with displacement (+D) (i.e. recipient moves away/ is displaced) see e.g. 'Body push'.
Body push: given (n)	'**BPG**'[a]: use of sudden impact contact made with the flank delivering a sideways shunt to the head or flank of a recipient; frequently results in displacement (e.g. when accessing a crowded feed barrier).
Body push: received (n)	'**BPR**'[a]: receipt of a body push from another cow
Body push: displacement (%)	'**BPR+D**': receipt of a body push, immediately (≤2 s) prompting the focal cow to be displaced). '**%BPR+D**'[b] = ('BPR+D'/'BPR') x 100
Non-contact challenge	'**CG**'[a]: performance of a threatening non-contact gesture, aimed at displacing a conspecific (e.g. short-charging, determinedly approaching, or facing/staring at a conspecific with head lowered). [**CR**'[a], '**CR+D**', '**%CR+D**'[b]]

Head butt	'**HBG**'[a]: striking head/body of another cow using the front of the lowered head with force. Can often be observed at the feed barrier as a single event where the rear or flank of a standing conspecific is often targeted, or at the cubicles where lying conspecifics can be targeted multiple times in quick succession until a response is evoked. ['**HBR**'[a], '**HBR+D**', '**%HBR+D**'[b]]
Head push	'**HPG**'[a]: the use of prolonged/sustained contact of the forehead (occasionally the lateral aspect of the head) applied to a conspecifics head or body. ['**HPR**'[a], '**HPR+D**', '**%HPR+D**'[b]]
Head swipe	'**HSG**'[a]: quick sideways swipe of the head, whereby the lateral aspect of the head is used to make violent contact with the head of a conspecific. Usually observed at the feed barrier May be performed singularly or multiple times in quick succession. ['**HSR**'[a], '**HSR+D**', '**%HSR+D**'[b]]
Total agonistic: given (n)	'**FocSocG**'[a] = 'BPG' + 'CG' + 'HBG' + 'HPG' + 'HSG'
Total agonistic: received (n)	'**FocSocR**'[a] = 'BPR' + 'CR' + 'HBR' + 'HPR' + 'HSR'
'All social reactivity' = percentage displacement following all agonistic interactions received	'**%FocSocR+D**'[b] = [('BPR+D' + 'CR+D' + 'HBR+D' + 'HPR+D' + 'HSR+D')/'FocSocR'] x 100
Mutual head butt (n/s)	'**MutHdButt**'[a]: mutual head-to-head butting between the focal cow and conspecifc. Usually observed in prolonged bouts whereby the two cows stand head-down facing each other with/without sustained forehead contact between each butt.
(C.2) **Social** (Luxury)**:** **non-agonistic**	
Allo-groom give (n/s)	'**AlloG**'[a]: licking any body part of a conspecific (most frequently the head or main torso). Usually observed in prolonged bouts.
Allo-groom receive (n/s)	'**AlloR**'[a]: receipt of licks from a conspecific. During licking the recipient usually stops all other activity and stands still.
Explore social (s)	'**ExpSoc**'[a] = 'Explore cow' (nose positioned close to another cow in the act of sniffing – no reciprocation) + 'Mutual sniff' (reciprocal sniffing between a focal cow and conspecific, usually stood facing one-another with noses almost touching).
Mutual head rub (n/s)	'**MutHdRub**'[a]: mutual head-to-head rubbing performed by a focal cow and a conspecific. Usually observed in bouts.
(C.3) **Total social interactions**	
Total social interactions: given (n)	'**SocG**'[a] = 'AlloG' + 'BPG' + 'CG' + 'Chin rest give' (use of chin to exert pressure on the lateral posterior of a conspecific) + 'HBG' + 'HPG' + 'Head rub give' (rubbing head on conspecific without reciprocation) + 'HSG' + 'Mount give' (bulling behavior where focal cow stands on back legs and rests her chest on the back/rump of a conspecifc) + 'MutHdButt' + 'MutHdRub'

Total social interactions: received (n)	'**SocR**'[a] = 'AlloR' + 'BPR' + 'CR' + 'Chin rest receive' + 'HBR' + 'HPR' + 'Head rub receive' + 'HSR' + 'Mount receive' + 'MutHdButt' + 'MutHdRub'
(D.) **Social Proximity and feed barrier preference**	
Feed barrier: all (s)	'**FB_All**': total time spent with the head positioned over the feed barrier regardless of type (i.e. open-rail or self-locking).
Feed barrier: no near neighbours (%)	'**FB_0NN**': time spent with the head positioned over the feed barrier without any direct flank-to-flank contact with conspecifics. '**%FB_0NN**'[b] = ('FB_0NN'/'FB_All') x 100
Feed barrier: two near neighbours (%)	'**FB_2NN**': time spent with the head positioned over the feed barrier with direct flank-to-flank contact with two other cows (i.e. cows positioned on either side). '**%FB_2NN**'[b] = ('FB_2NN'/'FB_All') x 100
Lie: no near neighbours (%)	'C_0NN': time spent lying within a cubicle flanked by two unoccupied cubicles. '**%C_0NN**'[b] = ('C_0NN'/'Lie') x 100
Lie: two near neighbours (%)	'C_2NN': time spent lying within a cubicle flanked by two occupied cubicles. '**%C_2NN**'[b] = ('C_2NN'/'Lie') x 100
Feed barrier: open (%)	'**FB_Open**': time spent with the head positioned over the open-rail section of the feed barrier. '**%FB_Open**'[b] = ('FB_Open'/'FB_All') x 100

204   [a] inclusion in 24h and 1hPostM1 data sets;[b] inclusion in 24h data set only

2.5. Saliva Collection and SAA

205
Saliva was collected (Day 3) using a cotton swab (SalivaBio Children's Swab, Item No.
5001.06, Salimetrics) and then immediately stored at -80°C prior to analysis. SAA was
measured in saliva from 31 cows (saliva volumes from three cows were too small to be
analysed), diluted 1:2, using a commercially available kit (Bovine Serum amyloid A protein
ELISA Kit, EB0015, Finetest®, Wuhan Fine Biotech Co. Ltd.). To assess the suitability of the
kit for use with saliva an assay validation was performed. To determine parallelism (linearity)
a displacement curve, produced by double-diluting a pooled saliva sample with assay buffer,
was compared to a standard curve. Percentage binding (as a percentage of that recorded for
the zero standard) was calculated, in addition to the Log of the standard concentration (SAA
standard) and the Log of the inverse of the dilution factor (saliva sample), e.g. 1:4 was
transformed to Log(1/4). Parallelism was confirmed using a statistical test for the analysis of
covariance (ANCOVA, SPSS). To measure assay accuracy the percentage recovery of
exogenous SAA was calculated following the addition of 300 ng/ml SAA standard to a
pooled saliva sample. Precision was assessed via intra- and inter-assay coefficients of
variation (**CV**); the former was determined following the repeated measurement of aliquots of
pooled saliva containing either high (quality control: $\mathbf{QC_{high}}$) or low ($\mathbf{QC_{low}}$) endogenous
SAA within the same plate, while the later was determined following the assay of QC_{high} and
QC_{low} samples in different plates.

2.6. Statistics

225
Following tests for normality (Shapiro-Wilk analysis), all behavioral measures, SAA and
SCC were compared between CTRL and SCM using Paired samples t-test or Wilcoxon
Signed-Rank tests (SPSS Statistics 24.0). Behavioral data from the continuous 24h data set
and the data sub-set (60 min after the morning milking, '1hPostM1') were separately
analysed. Since the experimental design required the performance of multiple comparisons
between measures there was an increased associated risk of Type I errors. Use of Bonferroni
correction procedures has been highlighted as problematic (especially for animal behavioral
studies, where sample sizes are often small) due to their tendency to increase Type II errors
(Nakagawa, 2004). As an alternative to standard correction procedures we, therefore,
calculated measures of observed (standardised) effect size in addition to p-values. Effect size
measures the strength/magnitude of a relationship and, thereby, helps us to determine the
strength of a statistical claim and whether a difference is real (i.e. it enables us to judge
biological importance). Hedges' g-value (Equations 1 and 2), also termed 'Cohen's d-value

for paired samples' (Hedges, 1981; Cohen, 1988; Nakagawa and Cuthill, 2007) and 95%

confidence intervals (**CI**) for effect size (Equations 3 and 4), were calculated for all measures

that met the assumptions of normality.

$$g = \frac{\bar{x}_2 - \bar{x}_1}{\sigma_{\text{paired}}} \qquad (1)$$

where

$$\sigma_{\text{paired}} = \sqrt{\frac{(n_2 - 1)s_2^2 + (n_1 - 1)s_1^2}{n_1 + n_2 - 2}} \qquad (2)$$

$\bar{x}_1$ and $\bar{x}_2$ are the means of the two groups, σ_{paired} is the pooled standard deviation, n is the

number of data points, and s^2 is the sample variance.

$$95\%\text{CI} = g - 1.96 se_{\text{g}} \text{ to } g + 1.96 se_{\text{g}} \qquad (3)$$

where

$$se_{\text{g}} = \sqrt{\frac{2(1 - r_{1,2})}{n} + \frac{g^2}{2(n-1)}} \qquad (4)$$

se_{g} is the asymptotic standard error for the effect size, $n = n_1 = n_2$, and $r_{1,2}$ is the correlation

coefficient between the two groups. For all behavioral measures that did not meet the

assumptions of normality, bootstrap effect size values (Hedges' g-value with 95%CI, R =

2000) were computed using the software package 'bootES' (Gerlanc and Kirby, 2012; Kirby

and Gerlanc, 2013) and R (Version 3.2.2., www.r-project.org/). Effect size statistics were

interpreted as follows: (a) the size of the effect (based upon the estimated g-values: ≤0.39 =

small, 0.40 - 0.79 = medium, ≥0.80 = large); (b) statistical significance (attributed to all

measures where the associated 95%CI did not contain '0') (Lee, 2016).

Interpretation of statistically non-significant p-values is possible using effect size confidence

intervals in combination with the effect size (see Nakagawa & Foster, 2004). To identify

those measures that were not statistically significantly different between SCM and CTL cows

in the continuous 24h data set but could yet be biologically important, we used information

from published studies to set broad, accepted, relative difference levels (**RDL%**, Table 2).

Although these studies are not specific to subclinical mastitis, examples include the impact of

clinical levels of (non-mastitic) infection and social status on behavior, it is assumed that they

provide generous and relevant difference levels with which to compare our subclinical

findings.

Table 2: Summary of the relative difference levels (RDL%) and their sources, used in this study to

ascertain the existence of a biologically meaningful difference in behavioral measures between cows,

with and without subclinical mastitis, based on effect size

Behavioral Measure[1]	RDL%	Reference
Agonistic social given: all categories	10	Sepulveda-Vares et al., 2016
Agonistic social received: all categories	10	Neave et al., 2018
%FB_0NN, %FB_2NN	20	Manson and Appleby, 1990
%FB_Open	10	Huzzey et al., 2006
AlloG	30	Galindo and Broom, 2002
AlloR	80	Galindo and Broom, 2002; Hoonhout et al., 2017
Brush	30	Mandel et al., 2017
Self-groom	20	Fogsgaard et al., 2012
Activity: Trans, Dist	10	King et al., 2018; Steensels et al., 2017
Feed	10	Dollinger and Kaufmann, 2013
Drink	20	Huzzey et al., 2007
Lie	10	Toaff Rosenstein et al., 2016
Other	20	

275   [1]%FB_0NN = percentage of time at feed barrier with no near neighbors; %FB_2NN = percentage of time at feed
276   barrier with two near neighbors; %FB_Open = percentage of time at open-rail feed barrier; AlloG = allogroom
277   give; AlloR = allogroom receive; Brush = use of revolving brush; Self-groom = licking, rubbing, scratching self;
278   Trans = behavioral transitions; Dist = distance moved; Lie = lying in cubicle.
279

For those measures where no relevant literature was available ('Other'; Table 2), we used the

average of the known RDLs (i.e. 20%; Table 2). For each measure, relative difference values

(**RDV**) were then calculated using the RDL% and the respective mean value from the CTRL

group. $95\%CI_{RDV}$ were calculated using the confidence intervals from the effect size statistics

and the (between-group) difference in means. In those cases where the $95\%CI_{RDV}$ did not

include the RDV, we conclude (with 95% confidence) that the current study showed no

important biological effect for that measure; we refer to these as 'biologically unimportant'.

In cases where the $95\%CI_{RDV}$ did include the RDV, we conclude that a difference was

inconclusive but plausible; we refer to these as 'biologically inconclusive'. For example, if

CTRL cows performed more body pushes than the SCM cows, yet this difference failed to

reach statistical significance ($P \geq 0.10$), using the p-value alone we would dismiss this

behavior as being unaffected by subclinical inflammation. However, if the RDV for this

behavior was within the $95\%CI_{RDV}$ range (e.g. RDV = 0.08, $95\%CI_{RDV}$ = -0.08 to 0.27) we

would conclude that, although this effect is biologically inconclusive based upon our

evidence, the difference may become significant given a larger sample size. Alternatively, if

the RDV, in the above example, was 0.3, then we would conclude that the effect was

biologically unimportant.

To test for correlations between physiological (SCC and SAA) and behavioral measures, we

performed curve estimation regression statistics using the continuous 24h data set (SPSS:

ANOVA, coefficient of determination) following tests for normality. Due to the small sample

size, standard deviations for the behavioral measures were already large, so outliers (± 2SD)

deemed to be atypically/excessively low or high were considered and removed prior to data

analysis. Such outliers comprised a maximum of a single data point (i.e. one matched pair)

for any measure.

3. RESULTS

3.1. Behavioral Differences over 24 hours (24h data set), and during 60 min

immediately following first milking (1hPostM1 data subset)

3.1.1 Luxury behavior: Social

(a) Total social interactions

In the 24h data, no significant differences were found in the total performance or receipt of

social interactions(Table 3). The difference in the total performance of social interactions was

classified as biologically unimportant in the context of this dataset ('**SocG**': RDV = 0.50,

95%CI$_{RDV}$ = -0.26 to 0.43). Total receipt of social interactions, however, was biologically

inconclusive on the basis of RDV confidence intervals ('**SocR**': RDV = 0.44, 95%CI$_{RDV}$ = -

0.21 to 0.64). This means that given a larger sample size, significant differences might

become apparent in the latter measure; i.e. it may be possible to confirm that SCM cows do

receive fewer social interactions. Indeed, during the 60 minutes following morning milking,

SCM cows did receive significantly fewer social interactions (1hPostM1; Table 4).

(b) Agonistic interactions

Broken down into separate interaction types, SCM cows performed significantly fewer head

butts ('**HBG**': 24h; 1hPostM1) and head pushes ('**HPG**': 1hPostM1) than healthy controls

(Tables 3 and 4). Non-significant measures of agonistic interactions given over 24h were

classified as biologically inconclusive on the basis of RDV confidence intervals, again

suggesting a larger sample might confirm an effect. These included body pushes ('**BPG**':

RDV = 0.08, 95%CI$_{RDV}$ = -0.08 to 0.27); non-contact challenges ('**CG**': RDV = 0.02,

95%CI$_{RDV}$ = 0.00 to 0.15), head pushes ('**HPG**': RDV = 0.03, 95%CI$_{RDV}$ = -0.03 to 0.18),

and head swipes ('**HSG**': RDV = 0.17, 95%CI$_{RDV}$ = -0.12 to 0.63) as well as total agonistic

interactions ('**FocSocG**': RDV = 0.40, 95%CI$_{RDV}$ = -0.31 to 0.73).

Over the 24h period, SCM cows received significantly fewer non-contact challenges than

healthy cows, but more head swipes (Table 3). With the exception of body push received,

which was classified as biologically unimportant ('**BPR**': RDV = 0.08, 95%CI$_{RDV}$ = -0.01 -

0.02), all non-significant measures of agonistic interactions received over 24h were classified

as biologically inconclusive on the basis of RDV confidence intervals, indicating that a larger

sample might confirm an effect. This included head butts ('**HBR**': RDV = 0.07, 95%CI$_{RDV}$ =

-0.05 to 0.13), head pushes ('**HPR**': RDV = 0.02, 95%CI$_{RDV}$ = -0.00 to 0.14) and mutual

butting ('**MutHdButt**': RDV = 1.13 s, 95%CI$_{RDV}$ = -0.19 to 8.03 s), as well as total agonistic

interactions ('**FocSocR**': RDV = 0.34, 95%CI$_{RDV}$ = -0.13 to 0.40).

During the 60 minutes following morning milking cows with subclinical mastitis received

significantly fewer total agonistic interactions than healthy controls, and specifically fewer

body pushes and head butts (1hPostM1; Table 4).

Table 3: Measures of social and other 'luxury' behavior recorded from pair-matched cows with (SCM)

or without (CTRL) sub-clinical mastitis over 24h: parametric data (P) analysed using paired t-test, non-

parametric data (NP) analysed using Wilcoxon SR test. All units (except %) transformed to 'per hour

visible'.

Measure[1] (unit)	Test	CTRL		SCM		Effect Size Statistics		
		Mean	*SD*	Mean	*SD*	Hedges' g	95%CI	Size of effect[2]
SocG[‡], (n)	NP	5.0[a]	*2.5*	4.5[a]	*2.2*	0.20	-0.53 – 0.87	Small, ns
SocR[‡], (n)	P	4.4[a]	*1.1*	5.0[a]	*2.5*	-0.34	-0.34 – 1.02	Small, ns
FocSocG[†], (n)	P	4.0[a]	*2.6*	3.3[a]	*1.5*	0.32	-0.48 - 1.12	Small, ns
FocSocR[‡], (n)	P	3.4[a]	*1.2*	3.9[a]	*1.8*	-0.30	-0.30 – 0.90	Small, ns
BPG[‡], (n)	NP	0.8[a]	*0.8*	0.5[a]	*0.3*	0.47	-0.25 – 0.90	Med, ns
BPR[†], (n)	P	0.8[a]	*0.3*	0.8[a]	*0.4*	0.11	-0.31 – 0.53	Small, ns
CG[‡], (n)	P	0.2[a]	*0.2*	0.1[b]	*0.1*	0.62	-0.01 - 1.25	Med, ns
CR[‡], (n)	P	0.3[a]	*0.3*	0.2[c]	*0.1*	0.84	0.02 - 1.66	Large, sig

HBG[‡], (n)	P	1.1[a]	*0.9*	0.6[b]	*0.4*	0.65	0.10 - 1.20	Med, sig
HBR[‡], (n)	P	0.7[a]	*0.4*	0.6[a]	*0.3*	0.37	-0.45 – 1.19	Small, ns
HPG[‡], (n)	NP	0.3[a]	*0.4*	0.1[a]	*0.1*	0.52	-0.15 - 1.04	Med, ns
HPR[‡], (n)	P	0.2[a]	*0.2*	0.1[b]	*0.2*	0.64	-0.01 – 1.29	Med, ns
HSG[†], (n)	P	1.7[a]	*0.8*	2.3[a]	*1.4*	-0.47	-0.23 – 1.18	Med, ns
HSR[‡], (n)	P	1.4[a]	*0.9*	2.0[b]	*1.5*	-0.51	0.01 – 1.02	Med, sig
MutHdButt[‡], (s)	NP	11.3[a]	*11.9*	5.3[a]	*4.0*	0.66	-0.03 - 1.34	Med, ns
%FocSocR+D[†], (%)	NP	49.1[a]	*11.1*	39.2[d]	*11.6*	0.87	0.11 – 1.63	Large, sig
%BPR+D[‡], (%)	NP	51.5[a]	*14.0*	53.7[a]	*17.9*	-0.11	-0.53 – 0.85	Small, ns
%CR+D[‡], (%)	NP	84.4[a]	*13.4*	76.9[a]	*23.3*	0.39	-0.32 – 1.10	Small, ns
%HBR+D'[†], (%)	P	69.9[a]	*15.5*	52.7[d]	*16.8*	1.10	0.46 - 1.74	Large, sig
%HPR+D[§], (%)	NP	52.8[a]	*25.5*	22.3[c]	*35.0*	0.96	-0.02 – 2.23	Large, ns
%HSR+D[‡], (%)	P	32.9[a]	*13.3*	24.7[c]	*13.6*	0.63	0.03 - 1.22	Med, sig
ExpSoc[‡], (s)	P	18.7[a]	*12.3*	9.1[d]	*5.4*	1.04	0.30 – 1.78	Large, sig
%FB_0NN[†], (%)	P	41.4[a]	*12.9*	36.0[b]	*13.0*	0.43	-0.02 – 0.88	Med, ns
%FB_2NN[‡], (%)	P	24.6[a]	*9.7*	30.0[c]	*9.8*	-0.56	0.11- 1.01	Med, sig
%FB_Open[†], (%)	P	82.2[a]	*12.0*	90.1[c]	*7.7*	-0.81	0.10 - 1.53	Large, sig
%C_0NN[†], (%)	P	38.8[a]	*20.3*	35.8[a]	*20.7*	0.43	-0.02 – 0.88	Med, ns
%C_2NN[†], (%)	NP	21.9[a]	*19.6*	18.6[a]	*20.6*	0.08	-0.81 - 0.58	Small, ns
AlloG[‡], (s)	P	18.2[a]	*14.0*	10.1[c]	*9.7*	0.70	0.05 – 1.35	Med, sig
AlloR[‡], (s)	NP	13.8[a]	*11.4*	15.9[a]	*14.7*	-0.16	-0.54 – 0.79	Small, ns
Brush[‡], (s)	NP	29.0[a]	*43.9*	11.9[a]	*9.0*	0.53	-0.16 - 0.96	Med, ns
Self-groom[‡], (s)	P	21.2[a]	*7.9*	26.0[a]	*10.8*	-0.27	-0.58 - 1.12	Small, ns
ExpEnv[‡], (s)	P	55.4[a]	*20.9*	50.0[a]	*14.5*	0.31	-0.24 - 0.86	Small, ns

[a–d]Mean values in the same row with different superscripts differ as follows: [b]$p < 0.10$ (trend); [c]$p \leq 0.05$; [d]$p \leq 0.01$
[1]SocG = all social behavior given; SocR = total social interactions received; FocSocG = agonistic behavior
given; FocSocR = agonistic behavior received; BPG = body push give; BPR = body push receive; CG =
aggressive challenge give; CR = aggressive challenge receive; HBG = head butt give; HBR = head butt receive;
HPG = head push give; HPR = head push receive; HSG = head swipe give; HSR = head swipe receive;
MutHdButt = mutual head butt; %FocSocR+D = 'social reactivity', i.e. percentage of occasions when receipt of
agonism lead to displacement; %BPR+D = percentage of occasions when body push receipt lead to
displacement; %CR+D = percentage of occasions when receipt of a non-contact challenge lead to displacement;
%HBR+D = percentage of occasions when head butt receipt lead to displacement; %HPR+D = percentage of
occasions when head push receipt lead to displacement; %HSR+D = percentage of occasions when head swipe
receipt lead to displacement; %FB_0NN = percentage of time at feed barrier with no near neighbors; %FB_2NN
= percentage of time at feed barrier with two near neighbors; %FB_Open = percentage of time at open-rail feed
barrier; %C_0NN = percentage of time lying in cubicle with no near neighbors; %C_2NN = percentage of time
lying in cubicle with two near neighbors; ExpSoc = explore social; AlloG = allogroom give; AlloR = allogroom
receive; Brush = use of revolving brush; Self-groom = perform self-grooming behavior; ExpEnv = explore
environment.
[2]The size of the effect (based on estimated g-values): ≤ 0.39 = small, 0.40 - 0.79 = medium, ≥ 0.80 = large;
significance attributed only when 95%CI >0
Number of data points used in statistical analysis: [†]n = 34; [‡]n = 32; [§]n = 22

Table 4: Differences in social and other 'luxury' behavior identified between pair-matched cows with

(SCM) or without (CTRL) sub-clinical mastitis during 60 mins following morning milking

(1hPostM1): parametric data (P) analysed using paired t-test, non-parametric data (NP) analysed using
Wilcoxon SR test.
376

Measure[1]	Test	CTRL		SCM		Effect Size Statistics		
		Mean	*SD*	Mean	*SD*	Hedges' g	95%CI	Size of effect[2]
SocR, (n)	NP	12.1[a]	*7.43*	6.6[c]	*3.92*	-0.91	0.17 – 1.51	Large, sig
FocSocG, (n)	P	10.7[a]	*9.0*	5.8[b]	*4.7*	-0.70	-0.05 – 1.45	Med, ns
FocSocR, (n)	P	4.7[a]	*2.6*	2.5[c]	*1.9*	-0.99	0.04 – 1.93	Large, sig
BPR, (n)	NP	3.3[a]	*2.5*	1.2[c]	*1.5*	-0.98	0.29 – 1.67	Large, sig
HBG, (n)	NP	1.7[a]	*2.4*	0.5[b]	*0.6*	-0.65	0.10 – 1.09	Med, sig
HBR, (n)	NP	1.1[a]	*1.4*	0.3[b]	*0.6*	-0.73	0.10 – 1.40	Med, sig
HPG, (n)	NP	0.9[a]	*1.3*	0.1[c]	*0.3*	-0.84	0.24 – 1.38	Large, sig
ExpSoc, (s)	NP	21.2[a]	*28.8*	6.5[c]	*7.6*	-0.68	0.06 – 1.07	Med, sig
AlloR, (s)	NP	37.6[a]	*66.1*	1.9[c]	*3.2*	-0.74	0.44 – 1.11	Med, sig
Self-groom (s)	P	29.2[a]	*25.6*	47.4[b]	*44.2*	0.52	0.01 - 1.03	Medium, sig
ExpEnv, (s)	P	71.8[a]	*44.2*	39.3[c]	*27.8*	-0.91	0.11 - 1.71	Large, sig

[a-c]Mean values in the same row with different superscripts differ as follows: [b]p <0.10 (trend); [c]p ≤0.05.
[1]SocR = all social behavior received; FocSocG = agonistic behavior given; FocSocR = agonistic behavior
received; BPR = body push receive; HBG = head butt give; HBR = head butt receive; HPG = head push give;
ExpSoc = explore social; AlloR = allogroom receive; Self-groom = perform self-grooming behavior; ExpEnv =
explore environment.
[2]The size of the effect (based upon estimated g-values): ≤0.39 = small, 0.40 - 0.79 = medium, ≥0.80 = large;
significance attributed only when 95%CI >0 (i.e. do not contain '0')
384

(c) Social reactivity, social exploration, social proximity, and feed barrier preference
Over the 24h period, cows with subclinical mastitis were also significantly less 'socially
reactive' than healthy cows (Table 3), i.e. were less likely to move away/be displaced
following the receipt of agonistic interactions (see Table 1). They performed significantly
less social exploration than healthy controls ('**ExpSoc**') during both the 24h period, and the
60 minutes following morning milking (Tables 3 and 4), and spent a significantly greater
proportion of their time at the open section of the feed barrier over the 24h period (Table 3).
Cows with subclinical mastitis spent a significantly greater proportion of their time at the
feed barrier flanked by two neighbours over the 24h period than healthy controls (Table 3).
On the basis of RDV confidence intervals, non-significant measures of social proximity were
classified as biologically unimportant within the context of this study. This included
proportion of time at the feed barrier without neighbors ('**%FB_0NN**': RDV = 8.28%,
$95\%CI_{RDV}$ = -0.11 to 4.75%) and the proportion of time lying within cubicles with zero or

two neighbors ('**%C_0NN'**: RDV = 7.75%, 95%CI$_{RDV}$ = -0.06 to 2.58%; '**%C_2NN**': RDV:

4.38%, 95%CI$_{RDV}$ = -2.69 to 1.93%).

(d)   Allogrooming

Over the 24h period, SCM cows performed allogrooming significantly less than healthy

controls (Table 3). Although receipt of allogrooming was not significantly different during

24h, and this measure was classified as biologically unimportant on the basis of RDV

confidence intervals ('**AlloR**': RDV = 11.02 s, 95%CI$_{RDV}$ = -1.14 to 1.67 s), during

1hPostM1 the SCM cows were allogroomed less than the healthy controls (Table 4).

3.1.2   Luxury behavior: Non-social

(a)   Self-grooming and brush use

In the 60 minutes following morning milking there was a tendency for SCM cows to perform

more self-grooming than healthy controls (Table 4). For the 24h period, differences in self-

grooming and brush use were not significant but classified as biologically inconclusive on the

basis of confidence intervals for effect size differences ('Brush': RDV = 8.70 s, 95%CI$_{RDV}$ =

-2.74 to 16.45 s; 'Self-Groom': RDV = 4.23 s, 95%CI$_{RDV}$ = -2.81 to 5.42 s).

(b)   Environmental exploration

In the 60 minutes following morning milking SCM cows explored the environment

significantly less than healthy controls (Table 4). This difference was not evident over the

24h period (Table 3), instead being classified as biologically unimportant in the context of

this study, as based upon RDV confidence intervals ('**ExpEnv**': RDV = 11.08 s, 95%CI$_{RDV}$ =

-1.30 to 4.66 s).

3.1.3   Core maintenance behavior

(a)   Feeding, drinking, and lying

For both the 24h observations and the 60 minutes following morning milking, no significant

differences emerged in time spent feeding ('**Feed**'), drinking ('**Drink**'), or lying ('**Lie**')

(Tables 5 and 6), nor was subclinical mastitis considered to have biologically important (or

even inconclusive) effects on these behaviors over 24h ('Feed': RDV = 82.39 s, 95%CI$_{RDV}$ =

-0.03 to 0.03 s; 'Drink': RDV = 6.86 s, 95%CI$_{RDV}$ = -1.33 to 3.99 s; 'Lie': RDV = 200.19 s,

95%CI$_{RDV}$ = -23.54 to 79.67 s). However, over the 24h period SCM cows spent significantly

more time lying with their head on their flank than healthy controls ('**HOF'**, Table 5).

(b)   <u>Physical activity</u>

During both the 24h period, and 60 minutes following morning milking, the SCM cows were

significantly less active than healthy controls; SCM cows performed fewer behavioral

transitions ('**Trans**') and moved over a smaller distance ('**Dist**') (Tables 5 and 6).

Table 5: Measures of 'core' maintenance behavior recorded from pair-matched cows with (SCM) or

without (CTRL) sub-clinical mastitis over 24h: analysed using paired t-tests. All units transformed to

'per hour visible'.

Measure[1] (unit)	CTRL		SCM		Effect size statistics		
	Mean	*SD*	**Mean**	*SD*	**Hedges' g**	**95%CI**	**Size of effect[2]**
Trans[‡], (n)	55.6[a]	*8.2*	48.6[c]	*7.8*	0.91	0.08 - 1.74	Large, sig
Dist[‡], (unit)	8.2[a]	*1.9*	6.8[c]	*1.5*	0.88	0.11 - 1.65	Large, sig
Feed[†], (s)	823.9[a]	*167.5*	823.8[a]	*281.2*	0.00	-0.46 - 0.46	Small, ns
Drink[†], (s)	34.3[a]	*12.2*	30.2[a]	*14.6*	0.32	-0.32 - 0.96	Small, ns
Lie[†], (s)	2001.9[a]	*211.6*	2092.9[a]	*367.6*	-0.31	-0.26 - 0.88	Small, ns
HOF[‡], (s)	143.5[a]	*49.7*	178.7[c]	*74.8*	-0.57	0.06 - 1.09	Med, sig

442   [a–c]Mean values in the same row with different superscripts differ as follows: [b]$p <0.10$ (trend); [c]$p \leq 0.05$.
443   [1]Trans = behavioral transitions; Dist = distance moved; Lie = lying in cubicle; HOF = lie with head on flank.
444   [2]The size of the effect (based upon estimated g-values): ≤ 0.39 = small, 0.40 - 0.79 = medium, ≥ 0.80 = large;
445   significance attributed only when 95%CI >0
446   Number of data points used in statistical analysis: [†]n = 34; [‡]n = 32
447

448   **Table 6:** Differences in 'core' maintenance behavior identified between cows with (SCM) or without

449   (CTRL) sub-clinical mastitis during 60 mins following morning milking (1hPostM1): parametric data

450   (P) analysed using paired t-test, non-parametric data (NP) analysed using Wilcoxon SR test.

Measure[1] (unit)	Unit	Control		SCM		Effect size statistics		
		Mean	*SD*	**Mean**	*SD*	**Hedge's g (bootstrap)**	**95%CI**	**Size of effect[2]**
Trans, (n)	NP	99.1[a]	*34.8*	71.7[c]	*24.6*	-0.89	0.21 - 1.62	Large, sig
Dist, (unit)	P	15.9[a]	*6.6*	12.5[b]	*4.0*	-0.63	-0.08 - 1.34	Med, ns

452   [a–c]Mean values in the same row with different superscripts differ as follows: [b]$p <0.10$ (trend); [c]$p \leq 0.05$.
453   [1]Trans = behavioral transitions; Dist = distance moved.
454   [2]The size of the effect (based upon estimated g-values): ≤ 0.39 = small, 0.40 - 0.79 = medium, ≥ 0.80 = large;
455   significance attributed only when 95%CI >0
456

457   **3.2   Correlations Between Physiology and Behavior**

3.2.1 Assay validation

Parallelism ($F_{1,9} = 3.46$, p >0.05) was confirmed between serial dilutions of saliva (range: 1:4 to 1:64) and SAA standards (range: 0, 9.38, 18.75, 37.5, 75, 150, 300 ng/ml), indicating that the ELISA kit was suitable for use with bovine saliva. Recovery of 300 ng/ml SAA from a spiked saliva sample was $93.76 \pm 4.63\%$ (n = 10). The intra-assay CV was 3.09% (250.87 ± 7.75 ng/ml, n = 10) for QC_{low} and 4.68% (1360.33 ± 63.70 ng/ml, n = 10) for QC_{high}. The inter-assay CV was 2.77% (246.06 ± 6.81 ng/ml, n = 2) for QC_{low} and 3.89% (1323.96 ± 51.43 ng/ml, n = 2) for QC_{high}.

3.2.2 SAA and SCC

The average SCC per group was: CTRL = 48.29 ± 28.33 (x1000 cells/ml); SCM = 351.12 ± 176.73 (x1000 cells/ml). A trend was found towards a significantly higher concentration of salivary SAA in the SCM cows (CTRL = 343.42 ± 269.60 ng/ml, SCM = 519.59 ± 315.43 ng/ml; $t_{1,12} = 1.93$, p = 0.076). A weak positive relationship was evident between SCC and salivary SAA levels ($F_{1,29} = 8.81$, p = 0.006, Figure 2).

INSERT FIGURE 2 NEAR HERE

3.3.3. SAA, SCC and behavior

All behavioral measures significantly correlated with SAA and/or SCC are presented in Table 8. The majority of these relationships are weak as indicated by R^2 values.

(a)   Total social interactions and agonistic interactions

Significant positive relationships were found between salivary SAA and total social interactions received (**'SocR'**) and total agonistic interactions received (**'FocSocR'**), indicating that cows with higher levels of salivary SAA cows received more social interactions in total, and also more agonistic interactions. Significant negative correlations were found between salivary SAA and total social interactions given (**'SocG')** and head swipes given (**'HSG'**), with a trend towards significance for total agonistic interactions given (**'FocSocG'**). This indicates that cows with higher levels of salivary SAA cows initiated fewer social interactions in total, including fewer agonistic interactions.

Table 8: Significant correlations (p<0.05; plus trends towards a correlation at p<0.10) between behavioral measures (24h data) and two markers of inflammation and mastitis infection, one from saliva (serum amyloid-A, SAA), and one from milk (somatic cell counts, SCC); given are curve

492 estimation regression statistics (ANOVA, coefficient of determination) and the equation for the

493 relationship (based upon the line of best fit).

Behavioral measure[1] (unit)	Correlation with salivary SAA and SCC
SocG (n)	SAA: $F_{(1,29)} = 6.42$, p = 0.017; $R^2 = 0.181$; $y = 6.0091e^{-7E-04x}$
SocR (n)	SAA: $F_{(2,28)} = 7.40$, p = 0.003; $R^2 = 0.346$; $y = 9E-06x^2 - 0.0061x + 5.057$
	SCC: $F_{(2,31)} = 4.90$, p = 0.014; $R^2 = 0.240$; $y = 2E-05x^2 - 0.0081x + 5.259$
FocSocG (n)	SAA: $F_{(1,29)} = 3.07$, p = 0.090; $R^2 = 0.096$; $y = 4.2715e^{-6E-04x}$
FocSocR (n)	SAA: $F_{(2,28)} = 4.23$, p = 0.025; $R^2 = 0.232$; $y = 6E-06x^2 - 0.004x + 3.817$
CR (n)	SAA: $F_{(1,29)} = 3.93$, p = 0.057; $R^2 = 0.119$; $y = 0.0003x + 0.165$
HPR (n)	SCC: $F_{(1,31)} = 5.87$, p = 0.021; $R^2 = 0.159$; $y = -1.990x + 5.520$
HSG (n)	SAA: $F_{(1,29)} = 6.89$, p = 0.014; $R^2 = 0.192$; $y = 2.3287e^{-9E-04x}$
HSR (n)	SCC: $F_{(2,30)} = 3.14$, p = 0.058; $R^2 = 0.173$; $y = 6E-06x^2 - 0.0018x + 1.621$
%FocSocR+D (%)	SAA: $F_{(1,29)} = 4.63$, p = 0.040; $R^2 = 0.138$; $y = -0.0155x + 52.078$
	SCC: $F_{(1,32)} = 10.46$, p = 0.003; $R^2 = 0.246$; $y = 49.136e^{-7E-04x}$
%HBR+D (%)	SAA: $F_{(1,29)} = 5.64$, p = 0.024; $R^2 = 0.163$; $y = 72.616e^{-5E-04x}$
	SCC: $F_{(1,31)} = 8.81$, p = 0.006; $R^2 = 0.221$; $y = -7.36\ln(x) + 95.594$
%HPR+D (%)	SAA: $F_{(2,22)} = 4.14$, p = 0.030; $R^2 = 0.227$; $y = -0.0002x^2 + 0.1952x - 0.092$
%HSR+D (%)	SAA: $F_{(1,29)} = 3.291$, p = 0.080; $R^2 = 0.102$; $y = -0.015x + 34.655$
ExpSoc (s)	SAA: $F_{(1,29)} = 4.34$, p = 0.046; $R^2 = 0.130$; $y = 0.0167x + 8.517$
	SCC: $F_{(1,31)} = 5.58$, p = 0.025; $R^2 = 0.153$; $y = 15.532e^{-0.002x}$
%FB_Open (%)	SAA: $F_{(1,28)} = 4.08$, p = 0.053; $R^2 = 0.127$; $y = 3.244\ln(x) + 67.752$
	SCC: $F_{(1,32)} = 5.98$, p = 0.020; $R^2 = 0.158$; $y = 81.19e^{0.0003x}$
ExpEnv (s)	SAA: $F_{(1,29)} = 5.96$, p = 0.021; $R^2 = 0.170$; $y = 8.690\ln(x) + 7.484$
Trans (n)	SCC: $F_{(2,30)} = 5.50$, p = 0.009; $R^2 = 0.269$; $y = 0.002x^2 - 0.0934x + 59.272$
Dist (unit)	SCC: $F_{(2,30)} = 4.11$, p = 0.026; $R^2 = 0.215$; $y = 4E-05x^2 - 0.021x + 9.146$
Feed (s)	SAA: $F_{(1,29)} = 14.63$, p = 0.001; $R^2 = 0.335$; $y = 1023.5e^{-6E-04x}$
Drink (s)	SAA: $F_{(1,29)} = 22.18$, p <0.001; $R^2 = 0.433$; $y = -8.379\ln(x) + 79.816$
Lie (s)	SAA: $F_{(1,29)} = 9.10$, p = 0.005; $R^2 = 0.239$; $y = 0.5032x + 1848.3$
	SCC: $F_{(2,31)} = 2.637$, p = 0.088; $R^2 = 0.145$; $y = -0.0019x^2 + 1.1955x + 1956.8$

[1]SocG = all social behavior given; SocR = all social behavior received; FocSocG = agonistic behavior given;
FocSocR = agonistic behavior received; CR = non-contact challenge receive; HPR = head push receive; HSG =
head swipe give; HSR = head swipe receivd; %FocSocR+D = percentage of occasions when receipt of agonism
lead to displacement; %HBR+D = percentage of occasions when head butt receipt lead to displacement;
%HPR+D = percentage of occasions when head push receipt lead to displacement; %HSR+D = percentage of
occasions when head swipe receipt lead to displacement; ExpSoc = explore social; %FB_Open = percentage of
time at feed barrier: open rail; ExpEnv = explore environment; Trans = behavioral transitions; Dist = distance
moved.

The significant relationship between SCC and total social interactions received (**'SocR'**) was

best modelled using a quadratic function. This described a negative association at low SCC

levels and a positive association at higher levels, such that as cell counts rose in cows with

low SCC (i.e. healthy CTRL cows) the tendency to receive social interactions would

decrease, but in those cows with higher SCC (i.e. SCM cows) as cell counts rose further the

tendency to receive social interactions increased. The significant positive correlation between

SCC and the receipt of head swipes ('**HSR'**) indicates that cows with higher SCC received

more agonistic head swipes.

(b)   Social reactivity

Significant negative relationships were found between measures of social reactivity, (i.e. the

likelihood that a cow would be displaced/move away after receiving an agonistic interaction),

and both inflammatory markers. This is indicative that cows with higher levels of salivary

SAA and/or SCC demonstrate a reduced tendency to be displaced following the receipt of

social stimuli, including total agonistic interactions ('**%FocSocR+D** '), and specifically to

head butts ('**%HBR+D')** (SAA and SCC), and head swipes received ('%HSR+D') (SAA).

(c)   Social and environmental exploration, and feed barrier preference

Salivary SAA was significantly positively associated with both social and environmental

exploration, such that cows with higher SAA levels performed more exploratory behavior.

SCC levels, however, were significantly negatively associated with social exploration, such

that cows with higher cell counts explored other cows less. Salivary SAA and SCC were

significantly positively correlated with proportion of time spent at the open-rail feed barrier

(**'%FB_Open'**), indicating that with rising levels of either inflammation measure, cows

became more likely to feed at the open-rail feed barrier rather than the self-locking barrier.

(d)   Physical activity, feeding, drinking, and lying

The significant relationships between SCC and both the number of behavioral transitions and

distance covered were best modelled using quadratic functions. These describe an initial

decline in both measures of physical activity as SCC increased to approximately 300 (x1000

cells/ml), followed by an increase in the measures as SCC levels continued to rise (Figure 3).

Salivary SAA was significantly negatively correlated with feeding and drinking, and

significantly positively correlated with time spent lying; this indicates that, within our study

cohort, that as SAA levels increased the cows spent less time feeding and drinking, and more

time lying.

4.   DISCUSSION

The purpose of this study was to identify potential changes in salivary serum amyloid-A

(SAA) and behavior in cows with spontaneously occurring subclinical mastitis as compared

to pair-matched, healthy controls. Salivary SAA was found to be higher in individuals with

subclinical mastitis, suggesting that it may have potential as a marker of low-level systemic

inflammation in dairy cows. Higher salivary SAA levels were also correlated with several

predicted behavioral changes. Furthermore, cows with subclinical mastitis displayed a

reduction in the performance and receipt of various social behaviours (here categorised as

'luxury' behaviors), lower 'social reactivity' (i.e. they were less likely to be displaced

following the receipt of agonism), a reduction in physical activity, but no difference in

feeding, drinking or lying duration ('core maintenance' behaviors).

4.1.   Changes in salivary SAA

We report here, for the first time, a positive (although weak) correlation between SCC and

salivary SAA. Positive associations between SCC and non-salivary SAA have previously

been reported from cows with clinical and sub-clinical mastitis (serum: de Boyer Des Roches

et al., 2017; milk: O'Mahony et al., 2006; Akerstedt et al., 2007; Pyörälä et al., 2011). SAA

in saliva thus offers potential as a non-invasive means of detecting subclinical mastitis, but

further studies will be required to substantiate our preliminary findings. Different bacterial

strains, for example, can cause mastitis of different duration and degree (Verbeke et al.,

2014), and it is possible that the concentration of SAA in saliva will vary accordingly. A case

in point is Pyörälä et al.'s (2011) finding of significant differences in SAA (milk) collected

from cows with spontaneous mastitis caused by different pathogens: low SAA was associated

with *A. pyogenes*, while high concentrations were associated with *E. coli*. Consequently,

further investigations will need to establish how, and when, salivary SAA levels change over

the time course of an infection, with different pathogens, to substantiate its diagnostic

potential as an inflammatory marker for mastitis and other diseases.

4.2.   Changes in Luxury Behaviors and correlations with salivary SAA

'Luxury' behaviors, as used here, are those deemed non-essential for survival within the

short-term and, as such, we predicted that during a subclinical infection they would be down-

regulated. Within this category we included both social and non-social behaviors.

4.2.1.   Social interactions

Over the 24h period, total social behavior, given or received, remained unaltered between the

two groups. Subclinically mastitic cows did, however, receive fewer non-contact challenges

and more head swipes. Head swipes are most commonly observed at the feed barrier as a

means of displacing competitors, and our result is in agreement with previous findings. Sick

cows are typically reported to be displaced more frequently than healthy cows when feeding,

as would result from head swipes or other agonistic behavior (Schirmann et al., 2016; Lomb

et al., 2018; Neave et al., 2018;).

In the 60 minutes following morning milking, that is during the period of peak feeding

activity in the study herd, other differences emerged; cows with subclinical mastitis received

fewer social interactions in total, and specifically fewer head butts and body pushes, than the

healthy controls. No corresponding reduction in the receipt of agonistic interactions involving

physical contact were observed over the 24h period. It is therefore possible that the

subclinically mastitic cows were avoiding peak feeding when aggression and competition for

feed are highest, as do low-ranking cows (Val-Laillet et al. 2008).

Interestingly, we also found a positive correlation between salivary SAA and total social

interactions, and agonistic interactions, received, indicating that those individuals with higher

levels of systemic inflammation received more agonistic interactions. However, we cannot be

sure whether SAA upregulation occurred in the cows with subclinical mastitis due to

infection or following exposure to social stress. Upregulation of C-Reactive Protein (another

acute phase protein known to increase during illness and stress) has, for example, been

reported in zoo-housed gorillas following an aggressive encounter (Fuller & Allard, 2018). In

the current study, saliva samples were collected the day after behavioral video recordings.

Therefore, an elevation in SAA could reflect agonistic encounters experienced during the

previous day, independent of health status. Further study should address this potential

confound.

In line with the prediction that social behaviors should decrease with subclinical mastitis,

affected cows performed fewer head butts than healthy cows, and SAA was negatively (albeit

loosely) correlated with the delivery of total social and agonistic interactions, including head

swipes in particular. Our results thus provide additional behavioral detail in support of

existing literature which reports that cows with clinical (and subclinical) conditions often

perform fewer agonistic interactions and competitive displacements from the feed-bunk

(Galindo & Broom, 2002; Huzzey et al., 2007; Patbandba et al., 2012; Sepúlveda-Varas et al.,

2014; 2016), and from cubicles (Jensen & Proudfoot, 2017), than healthy individuals.

The relative social ranks of study cows and their interaction partners were not determined in

the current study. It is possible, and an interesting focus for further study, that the relative

social rank of an individual could be influenced by the effects of disease due to a loss of

competitive vigour and motivation. High-ranking animals characteristically displace lower

rankers cows from the feed barrier (DeVries et al., 2004; Huzzey et al., 2006), and lower

rankers adjust their feeding patterns accordingly (DeVries et al., 2004).

4.2.2. Social reactivity

We found that over the 24h period, cows with subclinical mastitis were less likely to move

away/be displaced than matched healthy cows following the receipt of agonism, labelled here

as less 'socially reactive'. Correspondingly, reactivity declined with increasing inflammation

levels as demonstrated by a negative correlation between SAA levels and percentage

displacement following the receipt of agonistic interactions overall, and specifically

following the receipt of head butts and head swipes. Similarly, SCC was negatively

correlated with the percentage displacement following receipt of agonistic interactions

overall, and specifically of head butts, such that cows with higher SCC were less likely to

more away after being head butted, and/or following receipt of all agonistic interactions, than

cows with lower counts. These results appear, at first hand, to contradict previous findings of

sick cows being less motivated to compete for access to feed based on displacement (e.g.

Huzzey et al, 2006; 2007; Goldhawk et al., 2009); however, where a cow positions herself at

the feeder may, in turn, be influenced by the social rank of others already feeding, as reported

by Manson & Appleby (1990). They found that cows of the most dissimilar rank (low/high)

maintained a greater distance from each other than cows of similar rank. It is thus

conceivable that subclinically sick cows proactively avoided feeding positions next to

individuals of higher competitive vigour, and preferentially selected the company of other

sick or lower ranking cows. Subsequently, the receipt of moderate competitive aggression

from an individual, with similarly moderate motivation to compete, may have been tolerable

and thus not provoked displacement. This is beyond the reach of the current data as we did

not determine the relative social ranks of our focal cows and their interaction partners, but

warrants further investigation. As others before us, we can therefore not rule out that this and

our other findings on social behavioral differences reflect underlying differences in social

status that have led to differences in health status. We here provide a first report of a

reduction in social reactivity in cows with subclinical mastitis. By matching cows by parity

we sought some control over potentially confounding social rank biases. Further study is

needed to substantiate differences in social interactions as markers of subclinical mastitis and

other diseases.

4.2.3.   Social avoidance and proximity

In the current study, cows with subclinical mastitis performed less social exploration than

healthy controls during both the 24h period and the focussed 60 min following morning

milking, which indicates social avoidance or withdrawal. Furthermore, we observed a

negative correlation between this measure and SCC, such that cows with higher counts spent

less time exploring (sniffing) others. Sickness-driven social avoidance is well documented in

lab-animals and humans and is predicted via the action of pro-inflammatory cytokines on the

central nervous system (Kent et al., 1992; Bluthe et al., 1996; Dantzer & Kelley, 2007;

Arakawa et al., 2010). Due to the limited opportunity for social avoidance to occur within

intensive systems it has been relatively understudied in farm animals. We also found an

unexpected, if weak, positive correlation between SAA and social exploration, which may, at

least partially, be explained by the presence of both pre-clinical and post-clinical individuals

amongst our subclinical group. De Boyer Des Roches et al. (2017; 2018) report behavioral

changes including reduced environmental attentiveness during the pre-clinical phase of an

experimentally-induced mastitis model (i.e. prior to SCC and serum SAA upregulation), and

during the acute phase (coinciding with raised levels of SCC and SAA), but not during the

post-clinical remission phase (when high levels of SCC and SAA are evident). This suggests

that serum SAA may peak during post-clinical remission, rather than during the acute phase

of inflammation. Further study is therefore required to investigate when, and how, social

exploration changes throughout the course of spontaneously-occurring mastitis, and other

diseases, to confirm its usefulness in the detection of subclinical disease states.

We found that over the 24h period, cows with subclinical mastitis spent a greater proportion

of their time feeding at the open-rail section of the feed barrier than healthy controls. A weak

positive correlation was also found between this measure and SAA, which suggests that this

preference increases with systemic inflammation. Barrier design is known to influence the

incidence of agonistic behavior relating to feed access. Self-locking yokes have vertical bars

which separate the necks of adjacent cows, and these are better at reducing competitive

interactions (usually measured by displacements) compared to open post-and-rail barriers

(Endres et al., 2005; Huzzey et al., 2006). Our results therefore appear to contradict previous

findings. However, other factors are likely to also contribute to the choice of feeding location.

In the open section cows have better visibility and are more quickly able to withdraw from

potential agonistic interactions. Our finding that subclinically mastitic cows spent a greater

proportion of their time than the healthy cows at the feed barrier in close contact with two

(flanking) herdmates may be at odds with the expectation of sickness-driven social

avoidance. However, sick cows may also feel more vulnerable and therefore seek to feed in

company. Future longitudinal studies could determine whether individuals change their

feeding behavior in relation to health status, including whether they opt to feed amongst

others of lower rank when their own health is compromised.

4.2.4. Allogrooming

In agreement with our predictions relating to the basis of behavioral priorities during

infection, cows with subclinical mastitis performed less allogrooming than the healthy cows

over the 24h period, and during the 60 min following morning milking they themselves were

also allogroomed less. Previous studies have reported similar findings in relation to social

rank, with low-ranking cows performing and receiving less allogrooming than high rankers

(Napolitano et al., 2009), and allogrooming decreasing more in low-ranking cows under

conditions of increased competition (Val-Laillet et al., 2008). For health-related changes,

Galindo & Broom (2002) found lame cows to be allogroomed more than non-lame cows,

which is contrary to our result in subclinically mastitic cows. Their finding was interpreted as

a self-instigated coping strategy triggered by pain/discomfort associated with clinical

lameness, which arguably applies less to subclinical mastitis.

Since social grooming serves a variety of functions in cattle, including roles in hygiene,

pleasure, maintenance of social bonds and lowering social tension (Sato et al., 1991; 1993), it

is possible that prolonged suppression of allo-grooming, for example during a low-level

chronic inflammatory condition, could negatively affect cow welfare and health itself.

4.2.5. Non-social behavior

(a) Self-grooming

Against expectation, we did not find a difference in brush use between cows with and without

subclinical mastitis, nor a correlation between this behavior and SAA. Although grooming

itself is a comfort activity that healthy cows are highly motivated to perform (McConnachie

et al., 2018), brush use has been shown to decrease during non-mastitic disease (Toaff

Rosenstein et al., 2016; Mandel et al., 2017; Weigele et al., 2017). The brush in our study was

located central to many resources including the feed barrier, a water trough and cubicles and

was, therefore, readily accessible to all cows with minimal effort. In addition to our relatively

small sample size, a trade-off between brush location and the sensitivity of brush use for

detecting stress and morbidity (Mandel et al., 2013; 2017) may help to explain why no

decline was detected in the current study.

We observed cows with subclinical mastitis to perform more self-grooming (including self-

licking) than controls immediately following morning milking, presumably as a response to

mild udder discomfort or as a substitute for allo-grooming (see section 4.2.4). However, this

observation is at odds with existing literature that reports self-licking to remain unchanged

(Siivonen et al., 2011), or even decrease (Fogsgaard et al., 2012), during clinical mastitis.

Based upon our results the potential for the use of self-grooming as a marker of subclinical

mastitis remains inconclusive and requires further investigation.

(b) Environmental exploration

The weak positive correlation between SAA and environmental exploration described in the

current study was unexpected since exploratory behavior was predicted to decline with

increasing inflammation. As our study did not use an experimental model of induced mastitis

we do not know whether our focal cows comprised a mixture of early-stage, pre-clinical

mastitic cows (potentially associated with low SAA and sickness-driven reductions in

environmental exploration), and individuals in post-clinical remission (potentially associated

with high SAA and normal baseline values) (as discussed in Section 4.2.3). Ruminants

generally display low baseline levels of exploratory behavior when maintained in intensive

housing, since it is a largely unstimulating environment (De Rosa et al., 2009). Against

expectation, we also did not find a reduction in environmental exploration in our subclinically

mastitic cows over the 24 hours. They did, however, explore less than controls during the 60

min following morning milking. Environmental exploration, as defined in our study, included

sniffing the feed as well as pen fittings and cubicle sand. Lower levels of exploration in cows

with subclinical mastitis after morning milking may therefore also reflect avoidance of

feeding at peak time.

4.3.   Core Maintenance Behavior

4.3.1. Activity

Our finding that cows with subclinical mastitis made fewer behavioral transitions and moved

over a shorter distance than healthy cows is in agreement with our predictions, and with other

studies that describe reduced activity prior to the clinical diagnosis of mastitis (Kester et al.,

2015; Stangaferro et al., 2016; Veissier et al., 2017; King et al., 2018). The quadratic

relationships between SCC and both 'behavioral transitions' and 'distance moved' described

in the current study are of interest because mastitic cows have also been reported to display

increased activity (Siivonen et al., 2011; Medrano-Galarza et al., 2012), presumably due to

udder discomfort and an associated reduction in lying time. Jadhav et al. (2018) argue that the

threshold SCC value to delineate subclinical mastitis from normal should be 310, rather than

200 (x1000 cells/ml), as conventionally judged (e.g. Madouasse et al., 2010). This higher

value closely corresponds with the parabola vertex in both quadratic plots (Figure 3), of

approx. 300 (x1000 cells/ml); i.e. the point at which activity once again begins to rise.

4.3.2. Feeding/Drinking

Changes in feeding behavior have long been used to diagnose the onset of illness (Weary et

al., 2009). We found a negative correlation between SAA and feeding duration, as would be

predicted with sickness. Although we observed a negative correlation between SAA and

feeding duration, as would be hypothesised to occur with sickness, the average inflammatory

response within our SCM group overall was not sufficiently pronounced to trigger obvious

anorexia, as compared to CTRL cows. Sepúlveda-Varas et al. (2016) observed a decrease in

feed intake (but not duration) prior to the diagnosis of clinical mastitis which may be

attributed to underlying malaise. González et al. (2008) reported individual variability in

feeding behavior relating to spontaneously occurring udder disorders. Some cows

demonstrated a decrease in feeding duration with the onset of mastitis, while others showed

no change. It is possible that aspects of feeding behavior, other than duration, may have been

affected. Barn-housed cattle demonstrate highly synchronised feeding activity, with large

peaks in both feeding and social competition coinciding with fresh food delivery, and smaller

peaks following milking (DeVries & von Keyserlingk, 2005; Dollinger & Kaufmann, 2013).

Mastitic cows, presumably to avoid adverse social interactions, have been shown to feed at

less popular times such as early afternoon (Schirmann et al., 2016). While we found no direct

effect on feeding duration, our findings on social interactions after morning milking, reported

above, may be indicative of cows with subclinical mastitis avoiding peak feeding times.

Water and feed intake are positively related in cattle (Kume et al., 2010); however, drinking

tends to be less affected by health than feeding (Hart, 1988). Water is more immediately vital

for maintaining bodily functions (Kyriazakis and Tolkamp, 2011), and since drinking takes

less time than food consumption it is less prone to being disrupted by social competition at

the trough (Huzzey et al., 2007). Although a reduction in water consumption has been

reported in cows with mastitis (Lukas et al., 2008; Siivonen et al., 2011), and we observed a

moderate negative correlation between SAA and drinking duration, the level of systemic

inflammation within our subclinically sick cows may have been too low, and/or our sample

size too small, for a difference to emerge. Overall, the persistence of feeding and drinking

levels in our subclnically mastitic cows fits the prediction that core maintenance behavior is

conserved during early mastitic disease.

4.3.3. Lying

No difference in lying duration was found between our two groups. Lying is a highly

prioritised behavior in cattle due to its importance in rumination (Jensen et al., 2004; 2005;

Munksgaard et al., 2005) and dairy cows spend approximately 11h/day recumbent (Ito et al.,

2009; 2010). Increased lying duration, as a means of conserving energy and facilitating

recovery, is a key adaptation for sickness, and a positive correlation between SAA and lying

was found in our cows. Although extended lying duration has been frequently reported during

non-mastitic clinic conditions (Toaff Rosenstein et al., 2016; Weigele et al., 2017; Barragan

et al., 2018), lying may actually decrease during mastitis (Yeiser et al., 2012; Fogsgaard et

al., 2012; 2015; Medrano-Galarza et al. 2012;), most likely due to udder pain (Cyples et al.,

2012).

UOB Open

We also observed that cows with subclinical mastitis lay with their heads held against their

flank more than the healthy controls. This posture is primarily associated with rapid eye

movement (REM) sleep, but cows are also known to display non-rapid eye movement

(NREM) sleep and drowsing in this position (Ternman et al., 2013), and NREM (deep) sleep

often increases during infection (Bryant et al., 2004; Opp, 2005). Crucially, the action of pro-

inflammatory cytokines during the sickness response also predicts postural changes, such as

'curling up', that reduce surface area and associated loss of body heat. Lying with neck and

head bent back against and resting on the body achieves a reduction in surface area in cows,

and as such fits with predictions for adaptive behavioral changes during sickness. On the

basis of our study, lying duration itself appears less promising for the detection of sub-

clinical mastitis than our novel finding of a difference in lying posture.

4.4   Limitations

This study formed a preliminary investigation, to identify behaviors with potential for use as

a marker of sub-clinical mastitis, with the intention of informing a wider schedule of focused

research; longitudinal studies are now required to track changes in key behaviors with

deviations in the health status of individual cows. Due to an absence of pre-existing literature

(luxury behavior, in this context, has been understudied), a large number of measures were

recorded over 24h. Logistically this was time consuming and limited the amount of data

available per cow; priority was given to sample size. The analysis of a single 24h period for

each cow provided a snapshot of one day, It is therefore not possible to conclude that

behaviors observed were typical of that cow at that level of SCC. Although we health-

checked study cows for inclusion in the study, as described in the Methods, to rule out

additional clinical pathologies we could not determine subclinical conditions other than

mastitis, nor could we establish whether cows classified as having subclinical mastitis were in

the pre-clinical phases of disease or in remission.

5.   CONCLUSIONS

By studying in detail the behavior of healthy cows and cows with spontaneously occurring

subclinical mastitis, pair matched by reproductive parameters and parity, over two time

periods (24h and 60 min following morning milking) we found that subclinical mastitis was

associated with reductions in activity, social exploration, the receipt of non-agonistic social

behavior, social reactivity (probability of displacement following receipt of agonism), and an

increase in the receipt of head swipes, compared to clinically healthy control cows. Many

social interactions can be considered 'luxury' behaviors during sickness, and here we provide

preliminary findings that suggest that several social measures change at subclinical levels of

mastitis, as predicted, whereas 'core' maintenance activities (including feed, drink and lie)

did not, and luxury behaviors therefore offer greater potential for use in early disease

detection. The wider social context and, specifically the rank relationships between study

cows and their interaction partners, however, were beyond the scope of this investigation, and

further study is needed on the interactions between social rank, health status and behavior in

cows.

Cows with subclinical mastitis displayed a different feeding pattern. They spent a greater

proportion of their feeding time in direct contact with two neighbours, and a lower proportion

of time feeding at the self-locking feed barriers, than the healthy cows. Although a positive

relationship between SCC and salivary SAA was observed, and several correlations between

SAA and behavioral measures were identified in a direction consistent with sickness behavior

(including positive correlations with lying duration and the receipt of total agonism, and

negative correlations with feeding, drinking, the performance of total social and agonistic

behavior, and social reactivity), the majority of associations were relatively weak.

With this study, we have taken initial steps to identify physiological and detailed behavioral

changes associated with subclinical mastitis; a number of our findings are consistent with

predictions for low-level sickness responses. Social behavior is fundamentally dependent

upon the wider social environment and necessitates interactions between focal animals and

other individuals within the herd. We consider the preliminary identification of behavioral

differences in a small group of cows within a complex and dynamic social environment very

encouraging. We now recommend that observations be replicated in longitudinal studies,

tracking individuals, and larger data sets to substantiate and refine our findings.

ACKNOWLEDGEMENTS

This study was funded by The John Oldacre Foundation (grant code R12518-102). We would

also like to thank the staff at Wyndhurst Farm for facilitating our research.

COMPETING INTERESTS STATEMENT

The authors have no competing interests to declare.

REFERENCES

AHDB, 2015. The Agriculture and horticulture development board dairy mobility scoresheet https://projectblue.blob.core.windows.net/media/Default/Dairy/Publications/Dairy%20Mobili ty%20Scoresheet_200427_WEB.pdf - accessed 14/07/20

Åkerstedt, M., K. P. Waller, and A. Sternesjö, 2007. Haptoglobin and serum amyloid A in relation to the somatic cell count in quarter, cow composite and bulk tank milk samples. J. Dairy Res. 74:198-203. https://doi.org/10.1017/S0022029906002305

Andrew, W., J. Gao, S. M. Mullan, N. W. Campbell, A. Dowsey, and T. Burghardt, 2020. Visual identification of individual holstein friesian cattle via deep metric learning. Comput. Electron. Agric. https://arxiv.org/abs/2006.09205

Arakawa, H., K. Arakawa, and T. Deak, 2010. Sickness-related odor communication signals as determinants of social behavior in rat: a role for inflammatory processes. Horm. Behav. 57:30-341. https://doi.org/10.1016/j.yhbeh.2010.01.002

Barragan A. A., J. M. Piñeiro, G. M. Schuenemann, P. J. Rajala-Schultz, D. E. Sanders, J. Lakritz, and S. Bas, 2018. Assessment of daily activity patterns and biomarkers of pain, inflammation, and stress in lactating dairy cows diagnosed with clinical metritis. J. Dairy Sci. 101(9):8248-8258. https://doi.org/10.3168/jds.2018-14510

Bluthé, R. M., B. Michaud, K. W. Kelley, and R. Dantzer, 1996. Vagotomy attenuates behavioural effects of interleukin-1 injected peripherally but not centrally. Neuroreport 7(9):1485-1488. https://doi.org/10.1097/00001756-199606170-00008

Bryant, P. A., J. Trinder, and N. Curtis, 2004. Sick and tired: Does sleep have a vital role in the immune system? Nature Rev. Immunol. 4:457-467. https://doi.org/10.1038/nri1369

Cohen, J., 1988. Chapter 6: Differences between proportions. Pages 184-185 in Statistical Power Analysis for the Behavioral Sciences, 2nd edition. Hillsdale, Lawrence Erlbaum Publishers.

Cyples, J. A., C. E. Fitzpatrick, K. E. Leslie, T. J. DeVries, D. B. Haley, and N. Chapinal, 2012. Short communication: The effects of experimentally induced *Escherichia coli* clinical mastitis on lying behavior of dairy cows. J. Dairy Sci. 95(5):2571-2575. https://doi.org/10.3168/jds.2011-5135.

Dantzer, R. 2004. Cytokine-induced sickness behaviour: A neuroimmune response to activation of innate immunity. Eur. J. Pharmacol. 500:399–411. https://doi.org/10.1016/j.ejphar.2004.07.040

Dantzer, R. and K. W. Kelley, 2007. Twenty years of research on cytokine-induced sickness behaviour. Brain Behav. Immun. 21(2):153-160. https://doi.org/10.1016/j.bbi.2006.09.006

Dantzer, R., J. C. O'Connor, G. G. Freund, R. W. Johnson, and K. W. Kelley, 2008. From inflammation to sickness and depression: When the immune system subjugates the brain. Nature Rev. Neurosci. 9(1):46-56. https://doi.org/10.1016/j.bbi.2006.09.006

920 Dawkins, M. S. 1990. From an animal's point of view: Motivation, fitness, and animal
921 welfare. Behav. Brain Sci. 13:1–9. https://doi.org/10.1017/S0140525X00077104
922
923 De Boyer des Roches, A. D. B., M. Faure, A. Lussert, V. Herry, P. Rainard, D. Durand, and
924 G. Foucras, 2017. Behavioral and patho-physiological response as possible signs of pain in
925 dairy cows during *Escherichia coli* mastitis: A pilot study. J. Dairy Sci. 100:8385–8397.
926 https://doi.org/10.3168/jds.2017-12796
927
928 De Boyer des Roches, A. D. B., A. Lussert, M. Faure, V. Herry, P. Rainard, D. Durand, F.
929 Wemelsfelder, and G. Foucras, 2018. Dairy cows under experimentally-induced *Escherichia*
930 *coli* mastitis show negative emotional states assessed through Qualitative Behaviour
931 Assessment. Appl. Anim. Behav. Sci. 206:1-11.
932 https://doi.org/10.1016/j.applanim.2018.06.004
933
934 De Rosa, G., F. Grasso, A. Braghieri, A. Bilancione, A. Di Francia, and F. Napolitano, 2009.
935 Behavior and milk production of buffalo cows as affected by housing system. J. Dairy Sci.
936 92(3):907-912. https://doi.org/10.3168/jds.2008-1157
937
938 DeVries, T. J. and M. A. G. von Keyserlingk, 2005. Time of feed delivery affects the feeding
939 and lying patterns of dairy cows. J. Dairy Sci. 88: 625–631.
940 https://doi.org/10.3168/jds.S0022-0302(05)72726-0
941
942 DeVries, T. J., M. A. G. von Keyserlingk, and D. M. Weary, 2004. Effect of feeding space on
943 the inter-cow distance, aggression, and feeding behavior of free-stall housed lactating dairy
944 cows. J. Dairy Sci. 87:1432–1438. https://doi.org/10.3168/jds.S0022-0302(04)73293-2
945
946 Dollinger, J. and O. Kaufmann, 2013. Feeding behaviour in dairy cows with and without the
947 influence of clinical diseases or subclinical disorders. Archiv. Tierzucht 56(14):149-159.
948 https://doi.org/10.7482/0003-9438-56-014
949
950 Endres, M. I., T. J. DeVries, M. A. G. von Keyserlingk, and D. M. Weary, 2005. Effect of
951 feed barrier design on the behavior of loose housed lactating dairy cows. J. Dairy Sci.
952 88:2377–2380. https://doi.org/10.3168/jds.S0022-0302(05)72915-5
953
954 Fogsgaard, K. K., C. M. Røntved, P. Sørensen, and M. S. Herskin, 2012. Sickness behavior
955 in dairy cows during *Escherichia coli* mastitis. J. Dairy Sci. 95(2):630–638.
956 https://doi.org/10.3168/jds.2011-4350
957
958 Fogsgaard, K. K., T. W. Bennedsgaard, and M. S. Herskin, 2015. Behavioral changes in
959 freestall-housed dairy cows with naturally occurring clinical mastitis. J. Dairy Sci. 98:1730–
960 1738. https://doi.org/10.3168/jds.2014-8347
961
962 Fuller, G. and S. Allard, 2018. Preliminary data showing potential for salivary C-reactive
963 protein as an indicator of welfare in western lowland gorillas (*Gorilla gorilla gorilla*). J. Vet.
964 Behav. 28:58-62. https://doi.org/10.1016/j.jveb.2018.08.005
965
966 Galindo, F. and D. M. Broom, 2002. The effects of lameness on social and individual
967 behavior of dairy cows. J. Appl. Anim. Welfare Sci. 5(3):193–201.
968 https://doi.org/10.1207/S15327604JAWS0503_03
969

970 Gerlanc, D. and K. N. Kirby, 2012. bootES (Version 1.0). Retrieved from http://cran.r-
971 project.org/web/packages/bootES/index.html
972

973 Gleerup, K. B., P. H. Andersen, L. Munksgaard, and B. Forkman, 2015. Pain evaluation in
974 dairy cattle. Appl. Anim. Behav. Sci. 171:25-32.
975 https://doi.org/10.1016/j.applanim.2015.08.023
976

977 Goldhawk, C., N. Chapinal, D. M. Veira, D. M. Weary, and M. A. G. von Keyserlingk, 2009.
978 Prepartum feeding behavior is an early indicator of subclinical ketosis. J. Dairy Sci. 92:4971–
979 4977. https://doi.org/10.3168/jds.2009-2242
980

981 González, L. A., B. J. Tolkamp, M. P. Coffey, A. Ferret, and I. Kyriazakis, 2008. Changes in
982 feeding behavior as possible indicators for the automatic monitoring of health disorders in
983 dairy cows. J. Dairy Sci. 91:1017-1028. https://doi.org/10.3168/jds.2007-0530
984

985 Gu, J. Q., Z. H. Wang, R. H. Gao, H. R. Wu, 2017. Cow behavior recognition based on image
986 analysis and activities. Int. J. Agric. Biol. Eng. 10(3):165–174. https://doi.org/10.165-
987 174.10.3965/j.ijabe.20171003.3080
988

989 Hart, B. L., 1988. Biological basis of the behavior of sick animals. Neurosci. Biobehav. Rev.
990 12:123-137. https://doi.org/10.1016/S0149-7634(88)80004-6
991

992 Hedges, L.V., 1981. Distributional theory for Glass's estimator of effect size and related
993 estimators. J. Educ. Stats 6:107–128. https://doi.org/10.3102%2F10769986006002107
994

995 Hoonhout, L., I. Reimert, and C. I. Daigle, 2017. Relationship between allogrooming and
996 disease in feedlot steers: Social interactions may provide information about individual animal
997 health. J. Anim. Sci. 95(s4):2. https://doi.org/10.2527/asasann.2017.003
998

999 Huzzey, J. M., T. J. DeVries, P. Valois, and M. A. G. von Keyserlingk, 2006. Stocking
1000 density and feed barrier design affect the feeding and social behavior of dairy cattle. J. Dairy
1001 Sci. 89:126–133. https://doi.org/10.3168/jds.S0022-0302(06)72075-6
1002

1003 Huzzey, J. M., D. M. Veira, D. M. Weary, and M. A. G. von Keyserlingk, 2007. Prepartum
1004 behavior and dry matter intake identify dairy cows at risk for metritis. J. Dairy Sci. 90:3220–
1005 3233. https://doi.org/10.3168/jds.2006-807
1006

1007 ISO 13366-1, 1997. Part 1: Microscopic method (Reference method). In: Milk – Enumeration
1008 of Somatic Cells. International Organization for Standardisation, Geneva, Switzerland.
1009 https://webstore.ansi.org/Standards/ISO/ISO133661997
1010

1011 Ito, K., D. M. Weary, and M. A. G. von Keyserlingk, 2009. Lying behavior: Assessing
1012 within- and between-herd variation in free-stall-housed dairy cows. J. Dairy Sci. 92:4412–
1013 4420. https://doi.org/10.3168/jds.2009-2235
1014

1015 Ito, K., M. A. G. Von Keyserlingk, S. J. LeBlanc, and D. M. Weary, 2010. Lying behavior as
1016 an indicator of lameness in dairy cows. J. Dairy Sci. 93(8):3553-3560.
1017 https://doi.org/10.3168/jds.2009-2951
1018

1019 Jadhav, P. V., D. N. Das, K. P. Suresh, and B. R. Shome, 2018. Threshold somatic cell count
1020 for delineation of subclinical mastitis cases. Vet World 11(6):789-793.
1021 https://doi.org/10.14202/vetworld.2018.789-793.
1022
1023 Jensen, M. B. and K. L. Proudfoot, 2017. Effect of group size and health status on behavior
1024 and feed intake of multiparous dairy cows in early lactation. J. Dairy Sci. 100:9759–9768.
1025 https://doi.org/10.3168/jds.2017-13035
1026
1027 Jensen, M. B., L. Munksgaard, L.J. Pedersen, J. Ladewig, and L. Matthews, 2004. Prior
1028 deprivation and reward duration affect the demand function for rest in dairy heifers. Appl.
1029 Anim. Behav. Sci. 88:1–11. https://doi.org/10.1016/j.applanim.2004.02.019
1030
1031 Jensen, M. B., L. J. Pederson, and L. Munksgaard, 2005. The effect of reward duration on
1032 demand functions for rest in dairy heifers and lying requirements as measured by demand
1033 functions. Appl. Anim. Behav. Sci. 90:207–217.
1034 https://doi.org/10.1016/j.applanim.2004.08.006
1035
1036 Kent, S., R. M. Bluthé, K. W. Kelley, and R. Dantzer, 1992. Sickness behavior as a new
1037 target for drug development. Trends in Pharmacol. Sci. 13:24-28.
1038 https://doi.org/10.1016/0165-6147(92)90012-U
1039
1040 Kester, H. J., D. E. Sorter, and J. S. Hogan, 2015. Activity and milk compositional changes
1041 following experimentally induced *Streptococcus uberis* bovine mastitis. J. Dairy Sci.
1042 98(2):999-1004. https://doi.org/10.3168/jds.2014-8576
1043
1044 King, M. T. M., S. J. LeBlanc, E. A. Pajor, T. C. Wright, and T. J. DeVries, 2018. Behavior
1045 and productivity of cows milked in automated systems before diagnosis of health disorders in
1046 early lactation. J. Dairy Sci. 101:4343–4356. https://doi.org/10.3168/jds.2017-13686
1047
1048 Kirby, K. N. and D. Gerlanc, 2013. BootES: An R package for bootstrap confidence intervals
1049 on effect sizes. Behav. Res. 45:905–927. https://doi.org/10.3758/s13428-013-0330-5
1050
1051 Kováč, G., C. Tóthová, O. Nagy, O., and H. Seidel, 2011. Milk amyloid A and selected
1052 serum proteins in cows suffering from mastitis. Acta Vet. Brno 80:3–9.
1053 https://doi.org/10.2754/avb201180010003
1054
1055 Kovačević-Filipović, M., V. Ilić, Z. Vujčić, B. Dojnov, M. Stevanov-Pavlović, Z. Mijačević,
1056 and T. Božić, 2012. Serum amyloid A isoforms in serum and milk from cows with
1057 *Staphylococcus aureus* subclinical mastitis. Vet. Immunol. Immunopathol. 145(1-2):120-128.
1058 https://doi.org/10.1016/j.vetimm.2011.10.015
1059
1060 Kume, S., K. Nonaka, T. Oshita, and T. Kozaka, 2010. Evaluation of drinking water intake,
1061 feed water intake and total water intake in dry and lactating cows fed silages. Livest. Sci.
1062 128:46–51. https://doi.org/10.1016/j.livsci.2009.10.012
1063
1064 Kyriazakis, I. and B. J. Tolkamp, 2011. Hunger and Thirst. Pages 44–57 in Animal Welfare,
1065 Appleby, M.C., J. A. Mench, I. A. S. Olsson, and B. O. Hughes, Eds. 2nd ed. CAB
1066 International, Wallingford, UK.
1067

1068　Lecchi, C., F. Dilda, P. Sartorelli, and F. Ceciliani, 2012. Widespread expression of SAA and
1069　Hp RNA in bovine tissues after evaluation of suitable reference genes. Vet. Immunol.
1070　Immunopathol. 145:556-562. https://doi.org/10.1016/j.vetimm.2011.12.017
1071

1072　Lee, D. K., 2016. Alternatives to P value: confidence interval and effect size. Korean J.
1073　Anesthesiol. 69(6):555-562. https://doi.org/10.4097/kjae.2016.69.6.555
1074

1075　Littin, K., A. Acevedo, W. Browne, J. Edgar, M. Mendl, D. Owen, C. Sherwin, H. Würbel,
1076　and C. J. Nicol, 2008. Towards humane end points: Behavioural changes precede clinical
1077　signs of disease in a Huntington's disease model. Proc. Biol. Sci. 275:1865–1874.
1078　https://doi.org/10.1098/rspb.2008.0388
1079

1080　Lomb, J., H. W. Neave, D. M. Weary, S. J. LeBlanc, J. M. Huzzey, and M. A. G. von
1081　Keyserlingk, 2018. Changes in feeding, social, and lying behaviors in dairy cows with
1082　metritis following treatment with a nonsteroidal anti-inflammatory drug as adjunctive
1083　treatment to an antimicrobial. J. Dairy Sci. 101(5):4400-4411.
1084　https://doi.org/10.3168/jds.2017-13812
1085

1086　Lukas, J. M., J. K. Reneau, and J. G. Linn, 2008. Water intake and dry matter intake changes
1087　as a feeding management tool and indicator of health and estrus status in dairy cows. J. Dairy
1088　Sci. 91(9):3385-3394. https://doi.org/10.3168/jds.2007-0926
1089

1090　Mandel, R., H. R. Whay, C. J. Nicol, and E. Klement, 2013. The effect of food location, heat
1091　load, and intrusive medical procedures on brushing activity in dairy cows. J. Dairy Sci.
1092　96:6506–6513. https://doi.org/10.3168/jds.2013-6941
1093

1094　Mandel, R., C. J. Nicol, H. R. Whay, and E. Klement, 2017. Short communication: Detection
1095　and monitoring of metritis in dairy cows using an automated grooming device. J. Dairy Sci.
1096　100:5724-5728. https://doi.org/10.3168/jds.2016-12201
1097

1098　Manson, F. J. and M. C. Appleby, 1990. Spacing of dairy cows at a food trough. Appl. Anim.
1099　Behav. Sci. 26:69-81. https://doi.org/10.1016/0168-1591(90)90088-U
1100

1101　Madouasse, A., J. N. Huxley, W. J. Browne, A. J. Bradley, and M. J. Green, 2010. Somatic
1102　cell count dynamics in a large sample of dairy herds in England and Wales. Prev. Vet. Med.
1103　96:56–64. https://doi.org/10.1016/j.prevetmed.2010.05.005
1104

1105　McConnachie, E., A. M. C. Smid, A. J. Thompson, D. M. Weary, M. A. Gaworski, and M. A.
1106　G. Von Keyserlingk, 2018. Cows are highly motivated to access a grooming substrate. Biol.
1107　Lett. 14:1-4. https://doi.org/10.1098/rsbl.2018.0303
1108

1109　Medrano-Galarza, C., J. Gibbons, S. Wagner, A. M. de Passillé, and J. Rushen, 2012.
1110　Behavioral changes in dairy cows with mastitis, J. Dairy Sci. 95(12):6994-7002.
1111　https://doi.org/10.3168/jds.2011-5247
1112

1113　Munksgaard, L., M. B. Jensen, L. J. Pedersen, S. W. Hansen, and L. Matthews, 2005.
1114　Quantifying behavioural priorities - Effects of time constraints on behaviour of dairy cows,
1115　*Bos taurus*. Appl. Anim. Behav. Sci. 92:3–14.
1116　https://doi.org/10.1016/j.applanim.2004.11.005
1117

1118 Murata, H., N. Shimada, and M. Yoshioka, 2004. Current research on acute phase proteins in
1119 veterinary diagnosis: An overview. Vet. J. 168:28-40. https://doi.org/10.1016/s1090-
1120 0233(03)00119-9
1121
1122 Nakagawa, S., 2004. A farewell to Bonferroni: the problems of low statistical power and
1123 publication bias. Behav. Ecol. 15(6):1044–1045. https://doi.org/10.1093/beheco/arh107
1124
1125 Nakagawa, S. and I. C. Cuthill, 2007. Effect size, confidence interval and statistical
1126 significance: a practical guide for biologists. Biol. Rev. 82:591–605.
1127 https://doi.org/10.1111/j.1469-185x.2007.00027.x
1128
1129 Nakagawa, S. and T. M. Foster, 2004. The case against retrospective statistical power
1130 analyses with an introduction to power analysis, Acta Ethol. 7:103–108.
1131 https://doi.org/10.1007/s10211-004-0095-z
1132
1133 Napolitano, F., U. Knierim, F. Grasso, and G. De Rosa, 2009. Positive indicators of cattle
1134 welfare and their applicability to on-farm protocols. Ital. J. Anim. Sci. 8 (s1):355-365.
1135 https://doi.org/10.4081/ijas.2009.s1.355
1136
1137 Neave H. W., J. Lomb, D. M. Weary, S. J. Leblanc, J. M. Huzzey, and M. A. G. Von
1138 Keyserlingk, 2018. Behavioral changes before metritis diagnosis in dairy cows. J. Dairy Sci.
1139 101(5):4388-4399. https://doi.org/10.3168/jds.2017-13078
1140
1141 O'Mahony, M. C., A. M. Healy, D. Harte, K. G. Walshe, P. R. Torgerson, and M. L.
1142 Doherty, 2006. Milk amyloid A: Correlation with cellular indices of mammary inflammation
1143 in cows with normal and raised serum amyloid A. Res. Vet. Sci. 80:155–161.
1144 https://doi.org/10.1016/j.rvsc.2005.05.005
1145
1146 Opp, M. R., 2005. Cytokines and sleep. Sleep Med. Rev. 9(5):355-364.
1147 https://doi.org/10.1016/j.smrv.2005.01.002
1148
1149 Patbandha, T. K., T. K. Mohanty, S. S. Layek, A. Kumaresan, and K. Behera, 2012.
1150 Application of pre-partum feeding and social behaviour in predicting risk of developing
1151 metritis in cross bred cows. Appl. Anim. Behav. Sci. 139:10-17.
1152 https://doi.org/10.1016/j.applanim.2012.03.014
1153
1154 Petersson-Wolfe, C.S., Leslie, K.E. and Swartz, T.H., 2018. An update on the effect of
1155 clinical mastitis on the welfare of dairy cows and potential therapies. *Veterinary Clinics:
1156 Food Animal Practice*, *34*(3), pp.525-535.
1157
1158 Pyörälä, S., M. Hovinen, H. Simojoki, J. Fitzpatrick, P. D. Eckersall, and T. Orro, 2011.
1159 Acute phase proteins in milk in naturally acquired bovine mastitis caused by different
1160 pathogens. Vet. Rec. 168(20):535. https://doi.org/10.1136/vr.d1120
1161
1162 Rahman, M., U. Müller, H. Sauerwein, P. Heimberg, and F. Ceciliani, 2013. Investigation
1163 of salivary acute phase proteins in calves. Pages 213-216 in Farm Animal Proteomics
1164 2013, de Almeida A. et al. (eds). Wageningen Academic Publishers, Wageningen.
1165

1166 Sato, S., S. Sako, and A. Maeda, 1991. Social licking patterns in cattle (*Bos taurus*): influence
1167 of environmental and social factors. Appl. Anim. Behav. Sci. 32:3-12.
1168 https://doi.org/10.1016/S0168-1591(05)80158-3
1169
1170 Sato, S., K. Tarumizu, and K. Hatae, 1993. The influence of social factors on allogrooming in
1171 cows. Appl. Anim. Behav. Sci. 38:235-244. https://doi.org/10.1016/0168-1591(93)90022-H
1172
1173 Schirmann, K., D. M. Weary, W. Heuwieser, N. Chapinal, R. L. A. Cerri, and M. A. G. von
1174 Keyserlingk, 2016. Short communication: Rumination and feeding behaviors differ between
1175 healthy and sick dairy cows during the transition period. J. Dairy Sci. 99:9917–9924.
1176 https://doi.org/10.3168/jds.2015-10548
1177
1178 Sepúlveda-Varas, P., D. M. Weary, and M. A. G. von Keyserlingk, 2014. Lying behavior and
1179 postpartum health status in grazing dairy cows. J. Dairy Sci. 97:6334–6343.
1180 https://doi.org/10.3168/jds.2014-8357
1181
1182 Sepúlveda-Varasa, P., K. L. Proudfoot, D. M. Weary, and M. A. G. Keyserlingk, 2016.
1183 Changes in behaviour of dairy cows with clinical mastitis. Appl. Anim. Behav. Sci. 175:8-13.
1184 https://doi.org/10.1016/j.applanim.2014.09.022
1185
1186 Siivonen, J., S. Taponen, M. Hovinen, M. Pastell, B. Joop Lensink, S. Pyörälä, and L.
1187 Hanninen, 2011. Impact of acute clinical mastitis on cow behaviour. Appl. Anim. Behav. Sci.
1188 132:101–106. https://doi.org/10.1016/j.applanim.2011.04.005
1189
1190 Sordillo, L. M., K. Shafer-Weaver, and D. DeRosa, 1997. Immunobiology of the mammary
1191 gland. J. Dairy Sci. 80:1851–1865. https://doi.org/10.3168/jds.S0022-0302(97)76121-6
1192
1193 Stangaferro, M. L., R. Wijma, L. S. Caixeta, M. A. Al-Abri, and J. O. Giordano, 2016. Use of
1194 rumination and activity monitoring for the identification of dairy cows with health disorders.
1195 Part II. Mastitis. J. Dairy Sci. 99:7411-7421. https://doi.org/10.3168/jds.2016-10907
1196
1197 Steensels, M., E. Maltz, C. Bahr, D. Berckmans, A. Antler, and I. Halachmi, 2017. Towards
1198 practical application of sensors for monitoring animal health: the effect of post-calving health
1199 problems on rumination duration, activity and milk yield. J. Dairy Res. 84:132–138.
1200 https://doi.org/10.1017/S0022029917000176
1201
1202 Ternman, E., L. Hänninen, M. Pastell, S. Agenäs, and P. Nielsen, 2013. Sleep in dairy cows
1203 recorded with a non-invasive EEG technique. Appl. Anim. Behav. Sci. 140:25-32.
1204 https://doi.org/10.1016/j.applanim.2012.05.005
1205
1206 Toaff-Rosenstein, R. L., L. J. Gershwin, A. J. Zanella, and C. B. Tucker, 2016. The sickness
1207 response in steers with induced bovine respiratory disease before and after treatment with a
1208 non-steroidal anti-inflammatory drug. Appl. Anim. Behav. Sci. 181:49-62.
1209 https://doi.org/10.1016/j.applanim.2016.05.016
1210
1211 Val-Laillet, D., A. M. de Passille, J. Rushen, and M. A. G. von Keyserlingk, 2008. The
1212 concept of social dominance and the social distribution of feeding-related displacements
1213 between cows. Appl. Anim. Behav. Sci. 111:158–172.
1214 https://doi.org/10.1016/j.applanim.2007.06.001
1215

1216

1217 Veissier I., M. M. Mialon, and K. H. Sloth, 2017. Short communication: Early modification
1218 of the circadian organization of cow activity in relation to disease or estrus. J. Dairy Sci.
1219 100:3969–3974. https://doi.org/10.3168/jds.2016-11853

1220

1221 Verbeke, J., S. Piepers, K. Supré, and S. De Vliegher, 2014. Pathogen-specific incidence rate
1222 of clinical mastitis in Flemish dairy herds, severity, and association with herd hygiene. J.
1223 Dairy Sci. 97:6926–6934. https://doi.org/10.3168/jds.2016-11853

1224

1225 von Keyserlingk, M.A.G. and D. M. Weary, 2010. Review: Feeding behaviour of dairy cattle:
1226 Measures and applications. Can. J. Anim. Sci. 90(3):303-309.
1227 https://doi.org/10.4141/CJAS09127

1228

1229 Wagner, N., V. Antoine, M. M. Mialon, R. Lardy, M. Silberberg, J. Koko, and I. Veissier,
1230 2020. Machine learning to detect behavioural anomalies in dairy cows under subacute
1231 ruminal acidosis. Comput. Electron. Agric. 170:105233.
1232 https://doi.org/10.1016/j.compag.2020.105233

1233

1234 Weary, D. M., J. M. Huzzey, and M. A. G. von Keyserlingk, 2009. Using behavior to predict
1235 and identify ill health in animals. J. Anim. Sci. 87:770–777. https://doi.org/10.2527/jas.2008-
1236 1297

1237

1238 Weigele, H. C., L. Gygax, A. Steiner, B. Wechsler, and J. B. Burla, 2017. Moderate lameness
1239 leads to marked behavioral changes in dairy cows. J. Dairy Sci. 101:2370–2382.
1240 https://doi.org/10.3168/jds.2017-13120

1241

1242 Yeiser, E. E., K. E. Leslie, M. L. McGilliard, and C. S. Petersson-Wolfe, 2012. The effects of
1243 experimentally induced *Escherichia coli* mastitis and flunixin meglumine administration on
1244 activity measures, feed intake, and milk parameters. J. Dairy Sci. 95:4939–4949.
1245 https://doi.org/10.3168/jds.2011-5064

1265 **Figure 1:** Plan of the home pen including CCTV camera position and virtual division of floor-space
1266 (F1-13, M1/7/13, B1-13) for logging cow position. SB = salt bin, T1-3 = water troughs, B = rotating
1267 brush
1268

1294   **Figure 2:** The significant positive relationship between somatic cell count (SCC) in milk and salivary

1295   serum amylase-A (SAA) in dairy cattle ($R^2 = 0.233$; $y = 31.715x^{0.4815}$)

1331   **Figure 3:** Quadratic relationship between somatic cell count (SCC) and two measures of activity in

1332   cows with (SCM = black), and without (CTRL = open circle) subclinical mastitis: (a) behavioral

1333   transitions ('Trans'); (b) distance covered ('Dist').

1334   **(a)**   **(b)**

1335

The STROBE-Vet statement checklist.

	Item	STROBE-Vet recommendation	Page #
Title and Abstract	1	(a) Indicate that the study was an observational study and, if applicable, use a common study design term	1
		(b) Indicate why the study was conducted, the design, the results, the limitations, and the relevance of the findings	1
Background / rationale	2	Explain the scientific background and rationale for the investigation being reported	2
Objectives	3	(a) State specific objectives, including any primary or secondary prespecified hypotheses or their absence	4
		(b) Ensure that the level of organization[a] is clear for each objective and hypothesis	4
Study design	4	Present key elements of study design early in the paper	4
Setting	5	(a) Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	4
		(b) If applicable, include information at each level of organization	n/a
Participants [b]	6	(a) Describe the eligibility criteria for the owners/managers and for the animals, at each relevant level of organization	4
		(b) Describe the sources and methods of selection for the owners/managers and for the animals, at each relevant level of organization	4
		(c) Describe the method of follow-up	n/a
		(d) For matched studies, describe matching criteria and the number of matched individuals per subject (e.g., number of controls per case)	5
Variables	7	(a) Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. If applicable, give diagnostic criteria	5
		(b) Describe the level of organization at which each variable was measured	5
		(c) For hypothesis-driven studies, the putative causal-structure among variables should be described (a diagram is strongly encouraged)	2-4

Data sources / measurement	8*	(a) For each variable of interest, give sources of data and details of methods of assessment (measurement). If applicable, describe comparability of assessment methods among groups and over time	5-10
		(b) If a questionnaire was used to collect data, describe its development, validation, and administration	n/a
		(c) Describe whether or not individuals involved in data collection were blinded, when applicable	5
		(d) Describe any efforts to assess the accuracy of the data (including methods used for "data cleaning" in primary research, or methods used for validating secondary data)	n/a
Bias	9	Describe any efforts to address potential sources of bias due to confounding, selection, or information bias	n/a
Study size	10	(a) Describe how the study size was arrived at for each relevant level of organization	30
		(b) Describe how non-independence of measurements was incorporated into sample-size considerations, if applicable	n/a
		(c) If a formal sample-size calculation was used, describe the parameters, assumptions, and methods that were used, including a justification for the effect size selected	n/a
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen, and why	10-13
Statistical methods	12	(a) Describe all statistical methods for each objective, at a level of detail sufficient for a knowledgeable reader to replicate the methods. Include a description of the approaches to variable selection, control of confounding, and methods used to control for non-independence of observations	10-13
		(b) Describe the rationale for examining subgroups and interactions and the methods used	5-6
		(c) Explain how missing data were addressed	10
		(d) If applicable, describe the analytical approach to loss to follow-up, matching, complex sampling, and multiplicity of analyses	n/a
		(e) Describe any methods used to assess the robustness of the analyses (e.g., sensitivity analyses or quantitative bias assessment)	10-13
Participants	13*	(a) Report the numbers of owners/managers and animals at each stage of study and at each relevant level of organization - e.g., numbers eligible, included in the study, completing follow-up, and analyzed	4

		(b) Give reasons for non-participation at each stage and at each relevant level of organization	10
		(c) Consider use of a flow diagram and/or a diagram of the organizational structure	n/a
Descriptive data on exposures and potential confounders	14*	(a) Give characteristics of study participants (e.g., demographic, clinical, social) and information on exposures and potential confounders by group and level of organization, if applicable	4
		(b) Indicate number of participants with missing data for each variable of interest and at all relevant levels of organization	14,15,16,18
		(c) Summarize follow-up time (e.g., average and total amount), if appropriate to the study design	n/a
Outcome data	15*	(a) Report outcomes as appropriate for the study design and summarize at all relevant levels of organization	13-22
		(b) For proportions and rates, report the numerator and denominator	
		(c) For continuous outcomes, report the number of observations and a measure of variability	
Main results	16	(a) Give unadjusted estimates and, if applicable, adjusted estimates and their precision (e.g., 95% confidence interval). Make clear which confounders and interactions were adjusted. Report all relevant parameters that were part of the model	13-19
		(b) Report category boundaries when continuous variables were categorized	n/a
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	n/a
Other analyses	17	Report other analyses done, such as sensitivity/robustness analysis and analysis of subgroups	
Key results	18	Summarize key results with reference to study objectives	1,22,30-31
Strengths and Limitations	19	Discuss strengths and limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	30-31
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	30-31
Generalizability	21	Discuss the generalizability (external validity) of the study results	31
~~Funding~~ Transparency	22	(a) Funding- Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based (b) Conflicts of interest-Describe any conflicts of interest, or lack thereof, for each author	32

| | | (c) Describe the authors' roles- Provision of an authors' declaration of transparency is recommended
 (d) Ethical approval- Include information on ethical approval for use of animal and human subjects
 (e) Quality standards-Describe any quality standards used in the conduct of the research | |

[a] Level of organization recognizes that observational studies in veterinary research often deal with repeated measures (within an animal or herd) or animals that are maintained in groups (such as pens and herds); thus, the observations are not statistically independent. This non-independence has profound implications for the design, analysis, and results of these studies.

[b] The word "participant" is used in the STROBE statement. However, for the veterinary version, it is understood that "participant" should be addressed for both the animal owner/manager and for the animals themselves.

*Give such information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.