High regional and intra-generic variation in susceptibility to

mass bleaching in Indo-Pacific coral species

Paul R. Muir[1,2], Terrance Done [1,3], J. David Aguirre[4]

[1] Biodiversity and Geosciences, Queensland Museum, 78-102 Flinders St. Townsville Qld 4810

Australia. [2] Global Change Institute, University of Queensland, Brisbane, Queensland 4072, Australia.

[3] Australian Institute of Marine Science, Townsville Qld 4810 Australia. [4] School of Natural and

Computational Sciences, Massey University, Auckland, New Zealand.

Corresponding author: paul.muir@qm.qld.gov.au

Abstract

Aim: Mass bleaching is a major threat to reef-building corals and the ecosystems they

underpin. Here, we identified regional variation in the nature of this threat in terms of

the bleaching-susceptibility of individual coral species on some Indian Ocean and

Pacific Ocean reefs.

Location: 22 sites in the central Great Barrier Reef, Australia (GBR) and 30 sites in

the central Maldives Archipelago (MA).

Time period: 2002 for the GBR and 2016 for the MA.

Major taxa studied: Corals (Order Scleractinia)**.**

Methods: Following marine heat-wave conditions, timed in-situ surveys were used to

record bleaching responses (tissue colour) of large samples of individual coral

colonies. Responses of 106 shared species were analysed for sites with similar levels

of temperature stress, depth of occurrence and mortality. In each region, phylogenetic

mixed models were used to partition the effects on responses of species of deep-time

phylogeny, contemporary history and local-scale, among-site variability.

Results: Relative susceptibility to bleaching varied widely between regions: only 27

of the 106 shared species were in the same quartile for relative susceptibility in both

regions. Few species were highly susceptible in both regions. Closely related species

varied widely in their individual susceptibilities. Phylogenetic effects were moderate

in both regions, but contemporary phenotypic effects indicative of recent evolution

and acclimatization were greater in the MA, consistent with a stronger history of

recent bleaching.

Main conclusions: The high regional and intra-generic variation in coral bleaching-

susceptibility described here suggests there may be important differences in the extent

to which these Indian and Pacific Ocean coral populations are exhibiting responses to

deep-time evolutionary changes on the one hand, versus recent adaptation, on the

other. There is a concerning scarcity of this type of data, by which coral species most

at risk from bleaching in particular regions may be more accurately identified.

1. INTRODUCTION

Coral reefs are one of the most threatened ecosystems on the planet, with mass coral

bleaching widely regarded as their greatest current threat (Heron et al., 2016).

Repeated, severe bleaching events have significantly damaged reefs across the tropics

and subtropics, and mass bleaching events are predicted to increase in frequency and

severity in the near future (Hoegh-Guldberg et al., 2007; Hughes et al., 2017; Hughes,

et al., 2018a). Indeed, in 2020, the Great Barrier Reef (GBR) underwent its third

severe event within three years (Hughes & Pratchett, 2020). Local and possible

regional species extinctions related to repeated bleaching events have been reported

(Glynn, 2011; Muir et al., 2017; Sheppard et al., 2020) and there is now a risk of

global extinctions of reef-building coral species (Carpenter et al., 2008; IUCN, 2017;

Richards & Day, 2018).

Rising sea temperatures, the ultimate cause of most mass bleaching events

(Brown, 1997), are a global phenomenon, but the search for pragmatic management

interventions are focused at local and regional scales (Anthony et al., 2015;

Wooldridge & Done, 2009). While conventional management measures continue in

the form of reducing impacts from agricultural runoff, overfishing and coastal

development (GBRMPA, 2018; Steneck et al., 2019), current biological research

seeks to support active interventions such as reef restoration, assisted evolution,

assisted migration, hybridisation, seed banking, artificial refugia and assisted

recruitment (van Oppen et al., 2017). However, given the scale of the problem, with

around 750 species of scleractinian reef-corals (Hoeksema & Cairns, 2019) and over

280,000 km^2 of reefs threatened around the globe (Spalding et al., 2001), these

management and research efforts face a daunting task.

Despite the threat to reef-building corals globally, there is surprisingly little

data on the bleaching susceptibility of individual species. Susceptibility has been

documented mostly for coral genera (Baird et al., 2018; Guest et al., 2012; Hughes et

al., 2018b; McClanahan et al., 2007; Marshall & Baird, 2000; Chou et al., 2016; Loya

et al., 2001) with only a small number of species examined. While wide geographical

variation in relative susceptibility has been established for entire coral communities

(McClanahan et al. 2020, Thompson & van Woesik, 2009), there are few comparative

studies of individual taxa and these are mostly restricted to comparisons at a genus-

level (Guest et al., 2012, McClanahan et al., 2004; Pratchett et al., 2013). While much

of the current research is aimed at active interventions to prevent loss of coral species

following repeated mass bleaching events, the species most at risk from bleaching

have not been well documented. For instance, the IUCN "Red List" which is widely

used to categorise the risk and conservation status of coral taxa, is compiled from

mainly genus-level bleaching data from a small number of regions (Carpenter et al.,

2008) and thus might be misrepresenting the status of many coral species and regional

populations. The lack of data limits the extent to which management and interventions

could be targeted to the most susceptible species.

Here, we investigate the susceptibility of a broad range of Indo-Pacific scleractinian

species to moderate bleaching conditions in shallow reef waters of the central

Maldives Archipelago (MA) during 2016 and the central Great Barrier Reef (GBR)

during 2002. To the best of the authors' knowledge, our datasets are unique in

capturing in-situ bleaching responses for diverse assemblages of coral species. For

each region, phylogenetic mixed model analyses were used to estimate longer-term

phylogenetically heritable effects, shorter-term effects commensurate with adaptation

and/or acclimatisation (hereafter, contemporary phenotypic effects) and local-scale

variation. Comparison between regions were restricted to corals exposed to similar

levels of temperature stress at a similar depth of occurrence, but since mass bleaching

is caused by a complex array of stressors that can occur over many weeks (Skirving et

al. 2019) we avoided absolute comparisons. Instead, we compared the responses of

the 106 species present in both regions in terms of their relative susceptibilities. We

then assessed how the available data might be used to estimate species risk across

wide geographic regions and closely related species. The findings have important

implications for how coral bleaching, the greatest ongoing threat to these critically

important, yet vulnerable species, is assessed and how research and management

efforts might be prioritized.

2. MATERIALS AND METHODS

For the GBR in 2002 and the MA in 2016, sea temperatures exceeded bleaching thresholds over wide areas for several weeks, resulting in moderate mortalities over the ensuing months. To control for variation due to differing levels of temperate stress we restricted our analyses to sites with a Degree Heating Week (DHW) index of between 4 and 7 derived from satellite data (NOAA, 2017). This index is widely used to estimate the duration and severity of the temperature anomaly, with 4-7 DHW currently considered a moderate event (NOAA, 2017; Skirving, et al., 2019). Colony mortalities at selected sites reached 68% in the MA (Cowburn et al. 2019) and 86% in the GBR (Done et al. 2003). Since there can be a marked depth gradient in bleaching response (Muir et al., 2017; Baird et al., 2018), we focussed on corals assayed between 3 to 11 m depth in both regions. We further confined our analyses to the proportion of colonies with severe bleaching or recently partial or complete mortality to account for the possibility that less affected individuals recovered before our surveys (see below).

2.1 Field Surveys

A total of 14 locations (22 sites) in the central GBR and 10 locations (30 sites) in the central MA met the DHW criteria for inclusion in the analyses (Figure 1). At each site, timed surveys using SCUBA or snorkel dives of 45-80 minutes duration were conducted, recording bleaching responses and recent mortality of individuals according to standard methods (Supporting Information, Table S1). Species identifications were mostly made *in situ,* with high-resolution macro photographs or small samples taken from problematic colonies (under permit) for later analysis by

comparison with material available in the Queensland Museum, consultation with

relevant experts and reference to standard texts (see Supporting Information for

details). Both datasets were converted to current valid species (Hoeksema & Cairns,

2019). For certain genera (e.g. *Montipora, Porites, Goniopora* and *Cycloseris*) it was

not possible to accurately identify many of the species *in situ,* so these were assessed

as grouped species and only considered in a supplementary analysis.

The MA locations were surveyed two months after the peak of the temperature

anomaly, whereas the GBR locations were surveyed 4-6 months after. To assess the

effect of survey timing, we analysed a supplementary dataset of 142 individual corals

(five species) monitored *in situ* after a moderate bleaching event in the GBR (Baird &

Marshall, 2002, data kindly provided by A. Baird as: DOI to be supplied). These data

showed that colonies with mild to moderate bleaching had reasonable recovery,

whereas severely bleached colonies showed little recovery or >50% tissue death

(Supporting Information, Table S2). Thus, the proportion of severely affected or

recently dead/partially dead individuals remained consistent from 2.5 to 5.5 months

after peak temperatures. We were therefore confident that our analyses were robust to

differences in the timing of our surveys.

2.2 Statistical Analyses

The proportion of severely affected or recently dead/partially dead individuals

were analysed using phylogenetic generalised linear mixed models implemented in

the MCMCglmm package (Hadfield, 2019) in R v3.6.1. Separate models were applied

to each region to accommodate different fixed effects. For the MA analysis, fixed

effects included depth (3-5 m and 9-11 m), shading (shaded microhabitats and

unshaded) and the depth × shading interaction. For the GBR analysis, the fixed effects

included two levels of depth (3-5 m and 6-9 m), seven levels of habitat/zone

(nearshore-fringing reef, coastal-fringing reef, offshore-northern lagoon, offshore-

northern reef slope, offshore- southern back reef, offshore-southern lagoon and

offshore-southern reef slope) and habitat/zone × depth interaction. The random effects

used in both analyses were: phylogenetically heritable effects, contemporary

phenotypic effects, and location. Only taxa with more than two records in each region

were included in the analyses. See S2, Supporting Information for further detail.

The super tree of Huang and Roy (2016) was used in the phylogenetic mixed

model analyses, it currently being the most comprehensive for the group. We also

explored the effect of phylogenetic uncertainty on our results and undertook analyses

using a restricted phylogeny based only on molecular data (Huang & Roy, 2016).

These analyses are not considered further here as they recovered similar results to the

analyses conducted using the super tree phylogeny, indicating our methods were

robust to uncertainty in the construction of the super tree (Supporting Information,

Tables S4 and S5). A supplementary analysis using the Bleaching Mortality Index

(BMI, McClanahan et al., 2004) with species data pooled into genera was conducted

to allow comparison with previous reports.

3. RESULTS

Analysis of 2888 individuals from 180 species, 55 genera and 14 families from the

GBR showed moderate phylogenetically heritable effects, weak contemporary effects

and weak local-scale effects (Table 1). For the MA, analysis of data from 4480

individuals representing 144 species, 47 genera and 14 families also showed moderate

phylogenetic effects and minor local-scale effects. However, contemporary effects

were 2.7 times greater than those for the GBR. In both regions, the species responses

varied widely within genera (Figure 2). Genus *Acropora* had the largest number of

species present in each region (35 species in both regions) and species were present in

both the lowest and highest quartiles of bleaching response in each region (Figure 3).

Species of the genera *Pocillopora, Echinopora, Leptastrea, Platygyra* and

Dipsastraea showed a greater range of responses in the MA relative to the GBR

(Figure 2).

Of the 218 species with sufficient data for analysis, only 106 species (64

genera, 14 families) were present in both regions. Categorizing responses into

quartiles based on their posterior median estimates from the phylogenetic mixed

model analyses showed only 27 species (25.4%) were in the same quartile in both

regions, 45 species differed by one quartile and 34 by two or more quartiles (Figure

3). Only six species were in the most susceptible quartile in both regions. The

phylogenetically heritable effects for each species showed a weak correlation between

regions, markedly different from a slope of +/-1.0 expected if bleaching responses

were phylogenetically constrained between regions (Figure 4). The contemporary

effects also showed a low correlation between regions (Figure 4).

As contemporary effects were 2.7 times greater in the MA than the GBR

(Table 1) and the region had a stronger history of recent bleaching, we examined

species with the strongest contemporary effects consistent with recent adaptation or

acclimatisation ("C" in Supporting Information, Table S3). A wide range of genera

(21 of 47 analysed) were represented in the upper quartile (38 species) for

contemporary effects in the MA, including the ecologically significant genera:

Acropora, Dipsastraea, Cycloseris, Favites, Goniastrea, Hydnophora, Lobophyllia,

Pavona and *Pocillopora*.

For many genera, re-analysis of our data using the bleaching mortality index

(BMI) at the level of genus (Supporting Information, Figure S2) showed results

somewhat similar to those of the study that defined it (McClanahan et al., 2004).

However, the relative responses were markedly different to those derived from the

phylogenetic mixed models of our main analyses.

4. DISCUSSION

We report the first regional comparison of bleaching susceptibility for a broad range

of common Indo-Pacific reef corals at the level of species. Comparing shallow-reef

populations exposed to similar levels of moderate thermal stress we found relative

responses varied only slightly at local scales, but showed wide differences between

the regions. The findings have important implications for how mass bleaching

research is conducted and how this severe threat is addressed.

Prior to this study, the small amount of data available indicated low to

moderate levels of regional variation in the relative susceptibility to mass bleaching

for Indo-Pacific coral taxa. While entire assemblages varied widely in their overall

susceptibility (McClanahan et al. 2020), the major coral genera often showed a

similar hierarchy of susceptibility for different regions and bleaching events (e.g.

Loya et al. 2001, Marshall & Baird 2000, Hughes et al. 2017). The few direct

comparative studies also showed low to moderate regional variation. For example,

relative bleaching responses between East Africa and the north/central GBR were

similar for 19 coral genera (McClanahan et al., 2004), while a comparison of sites

within SE Asia and Australia found only two of 18 genera varied significantly in their

relative susceptibility for two sites (Guest et al., 2012). Our study showed markedly

higher levels of regional variation when susceptibility was considered at the level of

species. Only 25% of the 106 species with sufficient data for analysis were present in

the same quartile of susceptibility in the MA and GBR (Figure 3). This comparison

was for an identical assemblage of currently accepted species for similar levels of

temperature stress and depth of occurrence and for events that produced similar levels

of mortality.

Phylogenetic analyses allow the dependence resulting from a shared evolutionary

history along a phylogeny to be considered (Housworth et al., 2004) and here provide

an indication of the origins of variation in bleaching susceptibility between the two

assemblages. A strong inter-regional correlation in the phylogenetic contribution to

bleaching response would have suggested that bleaching susceptibility was strongly

phylogenetically constrained. By contrast, we found only a weak correlation (Figure

4) suggesting divergence in deep-time that has affected susceptibility to bleaching in

each region. This result, and indeed our general findings, are consistent with recent

genetics studies that indicate wide divergence within many species of Indian and

Pacific Ocean reef corals, fishes and other reef fauna (summarized Richards et al.,

2016). Several cryptic regional endemic species or subspecies have also been resolved

within species previously considered pan Indo-Pacific (Richards et al., 2016; Arrigoni

et al., 2020), suggesting regional divergence of reef faunas may be greater than

currently recognised. Such cryptic regional endemics would account for some the

regional variation documented here. The existence of more regionally restricted taxa

would have implications for management and conservation: taxa with smaller range

sizes are likely to have a greater risk of extinction (Richards et al., 2016), further

strengthening the case for species-level and regional assessments of susceptibility.

Our analyses indicate that phylogeny imposes only limited constraints on the

bleaching responses in each region. This suggests that regional divergence of entire

lineages, as well as recent acclimation and adaptation, play a role in mediating

bleaching responses. The greater proportion of variation partitioned to contemporary

effects in the MA assemblage in 2016, relative to the 2002 GBR assemblage is

consistent with this view (Table 1). The MA study area experienced several bleaching

events in the two decades before our surveys (summarized Muir et al., 2017),

potentially providing a strong pressure for contemporary phenotypic responses. By

contrast, although several events had been reported on the GBR in the two decades

before our 2002 surveys (Berkelmans & Oliver, 1999; Berkelmans et al., 2004), our

study sites had not yet been exposed to the globally widespread and severe events of

the present century, and thus any associated selective tendency towards reduced

susceptibility to subsequent bleaching.

Reduced susceptibility following a series of recent bleaching events has been

reported previously, but mainly for communities (Thompson & van Woesik, 2009;

McClanahan et al. 2020) or genera. Relatively low bleaching responses for the genera

Pocillopora and *Acropora* in Singapore and Malaysia (Guest et al., 2012, Chou et al.,

2016) and for *Acropora* and *Montipora* in French Polynesia (Pratchett et al., 2013)

have been attributed to adaptation/acclimation following several bleaching events.

Here, we partitioned individual species responses between deep-time and

contemporary effects to better understand the potential for adaption and/or

acclimation. This method could provide a means of selecting species for different

streams of management and intervention. For example, in the MA the species with the

strongest contemporary phenotypic resistance following repeated bleaching events

("C" in Supporting Information, Table S3) are potential candidates for interventions

such as reef restoration (Chamberland et al., 2017; dela Cruz & Harrison, 2017),

assisted migration (van Oppen et al., 2017) and isolation of genes that confer

resistance (van Oppen et al., 2015). Conversely, species with weak contemporary

effects and higher bleaching responses ("P" in Supporting Information, Table S3) are

potentially at increased risk and require heightened monitoring and first consideration

for interventions such as artificial refugia (Coelhoa et al., 2017), assisted evolution

(van Oppen, et al., 2015) and 'seed-banking' of cryopreserved gametes or larvae

(Daly et al., 2018). Clearly, more data from other events and regions are required to

fully realize the potential offered by this approach: our categories are only applicable

for the MA and will likely change as the populations are challenged by further

bleaching events.

High regional and intra-generic variation in bleaching susceptibility have

important implications for the prioritization of reef management and conservation.

The IUCN Red List (Carpenter et al., 2008) is widely used to set these priorities, but

is currently based on mostly genus-level bleaching data from few locations. Obtaining

the data to address this issue may be challenging as many species require expert

taxonomic input for accurate identification, the dominant genera *Acropora, Porites and Montipora* being particularly difficult. In addition, high local-scale variation can be problematic for *in situ* bleaching assessments (Chou et al., 2016; Cantin & Spalding, 2008), but our methods produced low local-scale variation and consistent species responses within each region (Table 1). Thus, despite the challenges there are the means to better inform management priorities and mitigate species loss.

By assessing coral bleaching responses at greater taxonomic resolution over a wide range of taxa, we found much greater variation within genera and between regions than had previously been reported. By analysing these data using phylogenetic comparisons we found evidence for this variation having origins in both deep-time and recent adaptation and/or acclimation in response to repeated events. Overall, we conclude that there is an alarming lack of reliable data on species' susceptibility with which to address the coral bleaching phenomenon, the greatest current threat to these already endangered species. Somewhat encouragingly, we found that only a small proportion of species were highly susceptible in both regions and that bleaching responses were weakly phylogenetically constrained, with some potential for adaptation and/or acclimation to moderate levels of bleaching stress.

Acknowledgements. We thank Paul Marshall, Ameer Abdullah, Hussein Zahir, Emre Turak and Mary Wakeford for assisting with fieldwork and Michel Pichon and Carden Wallace for taxonomic advice to both teams. Emre Turak provided much of the identifications for the GBR surveys. Ray Berkelmans, Tom Schlesinger, Michel Pichon and Cynthia Riginos provided valuable advice and Andrew Baird kindly provided the supplementary bleaching recovery data.

REFERENCES

Anthony, K. R. N., Marshall, P. A., Abdulla, A., Beeden, R., Bergh, C., Black, R., … Wear, S. (2015) Operationalizing resilience for adaptive coral reef management under global environmental change marine conservation. *Global Change Biology,* 21, 48-61.

Arrigoni, R., Berumen, M. L., Mariappan, K. G, Beck, P. S. A., Hulver, A. M., Montano, S, … Benzoni F. (2020) Towards a rigorous species delimitation framework for scleractinian corals based on RAD sequencing: the case study of *Leptastrea* from the Indo-Pacific. *Coral Reefs,* 39, 1001–1025. doi.org/10.1007/s00338-020-01924-8

Baird. A. H., Madin, J. S., Álvarez-Noriega, M., Fontoura, L., Kerry, J. T., Kuo1, C. Y., … Hughes T. P. (2018) A decline in bleaching suggests that depth can provide a refuge from global warming in most coral taxa. *Marine Ecology Progress Series,* 603, 257–264. https://doi.org/10.3354/meps12732

Baird, A. & Marshall, P. (2002) Mortality, growth and reproduction in scleractinian corals following bleaching on the Great Barrier Reef. *Marine Ecology Progress Series,* 237, 133–141.

Berkelmans, R. & Oliver, J. K. (1999) Large-scale bleaching of corals on the Great Barrier Reef. *Coral Reefs,* 18, 55–60.

Berkelmans, R., De'ath, G., Kininmonth, S., Skirving, W. J. (2004) A comparison of the 1998 and 2002 coral bleaching events on the Great Barrier Reef: spatial correlation, patterns, and predictions. *Coral Reefs,* 23,74–83.

Brown, B. E. (1997) Coral bleaching: causes and consequences. *Coral Reefs* 16, S129–S138.

Cantin, N. E, Spalding, M. (2008) Detecting and monitoring coral bleaching events. In M. J. H. van Oppen & Lough, J. M. (Eds.), *Coral Bleaching, Ecological Studies* (pp. 85-110), Vol. 205. Berlin, Germany: Springer.

Carpenter, K. E., Abrar, M., Aeby, G., Abrar, M., Aeby, G., Aronson, R. B., … Wood, R. (2008) One-third of reef-building corals face elevated extinction risk from climate change and local impacts. *Science,* 321, 560–563.

Chamberland, V. F., Petersen, D., Guest, J. R., Brittsan, M., Vermeij, M. J. A. (2017) New seeding approach reduces costs and time to outplant sexually propagated corals for reef restoration. *Science Reports*, 7, 18076. doi.org/10.1038/s41598-017-17555-z

Chou, L. M., Toh, T. C., Toh, K. B., Ng, C. S. L., Cabaitan, P., …Song, T. (2016) differential response of coral assemblages to thermal stress underscores the complexity in predicting bleaching susceptibility. *PLoS ONE,* 11, e0159755. doi:10.1371/journal.pone.0159755

Coelhoa, V. R., Fenner, D., Carusoc, C., Baylesa, B. R., Huanga, Y., Birkeland, C. (2017) Shading as a mitigation tool for coral bleaching in three common Indo-Pacific species. *Journal of Experimental Marince Biology & Ecology,* 497, 152–163.

Cowburn, B., Moritz, C., Grimsditch, G., Solandt, J. L. (2019) Evidence of coral bleaching avoidance,resistance and recovery in the Maldives during the 2016 mass-bleaching event. *Marine Ecology Progress Series*, 626, 53–67. https://doi.org/10.3354/meps13044

Daly, J., Zuchowicz, N., Nuñez Lendo, C. I., Kosla, K., Lager, C., Henley, M. … Hagedorn, M. (2018) Successful cryopreservation of coral larvae using vitrification and laser warming. *Science Reports,* 8, 15714. DOI:10.1038/s41598-018-34035-0

dela Cruz, D. W. & Harrison, P. L. (2017) Enhanced larval supply and recruitment can replenish reef corals on degraded reefs. *Science Reports,* 7, 13985. DOI:10.1038/s41598-017-14546-y

Done T., Turak E., Wakeford M., De'ath G., Kininmonth S., Wooldridge S., … Mahoney M. (2003) *Testing bleaching resistance hypotheses for the 2002 Great Barrier Reef bleaching event.* Australia: Australian Institute of Marine Science.

GBRMPA (2018) *Reef 2050 Long-Term Sustainability Plan.* Canberra, Australia: Commonwealth of Australia.

Glynn, P. W. (2011) In tandem reef coral and cryptic metazoan declines and extinctions. *Bulletin of Marine Science*, *87*, 767–794.

Guest, J. R., Baird, A. H., Maynard, J., Muttaqin, E., Edwards, A. J., … Chou, L. M. (2012) Contrasting patterns of coral bleaching susceptibility in 2010 suggest and adaptive response to thermal stress. *PLoS ONE* 7, e33353.

Hadfield, J. D. (2019) *Package MCMCglmm.* See https://cran.r-project.org

Heron S., Eakin, C. M., Douvere, F. (2017) *Impacts of climate change on world heritage coral reefs: A first global scientific assessment.* Paris, France: UNESCO World Heritage Centre.

Hoegh-Guldberg, O., Mumby, P. J., Hooten, A. J., Steneck, R. S., Greenfield, P., Gomez, E., … Hatziolos, M. E. (2007) Coral reefs under rapid climate change and ocean acidification. *Science,* 318, 1737–1742. doi:101126/science1152509

Hoeksema, B. & Cairns, S. (2019) *World list of Scleractinia.* See www.marinespecies.org/scleractinia.

Housworth, E. A., Martins, E. P., Lynch, M. (2004) The phylogenetic mixed model. *American Naturalist,* 163, 84-96.

Huang, D. & Roy, K. (2016) The future of evolutionary diversity in reef corals. *Philosophical Transactions of the Royal Society B,* 370, 20140010.

Hughes, T. P., Kerry, J. T., Baird, A. H., Connolly, S. R., Dietzel, A., Eakin, M., … Torda, G. (2018a) Global warming transforms coral reef assemblages. *Nature* 556, 493-496.

Hughes T. P., Anderson K. D., Connolly S. R., Heron S. F., Kerry J. T., Lough J. M., … Wilson S. K. (2018b) Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science,* 359, 80–83.

Hughes, T. P., Kerry, J. T., Álvarez-Noriega, M., Álvarez-Romero, J. G., Anderson, K. D., & Baird, A. H., … Wilson, S. K. (2017) Global warming and recurrent mass bleaching of corals. *Nature,* 543, 373. https://doi.org/10.1038/nature21707

Hughes, T. & Pratchett, M. (2020) We just spent two weeks surveying the Great Barrier Reef. What we saw was an utter tragedy. See www. theconversation.com.

IUCN (2017) IUCN Red List of Threatened Species.Ver. 2017.1 Retrieved from www.iucnredlist.org.

Loya, Y., Sakai, K., Yamazato, K., Nakano, Y., Sambali, H., van Woesik, R. (2001) Coral bleaching: the winners and the losers. *Ecological Letters* 4, 122-131.

Marshall, P. A. & Baird, A. H. (2000) Bleaching of corals on the Great Barrier Reef: differential susceptibilities among taxa. *Coral Reefs,* 19, 155-163.

McClanahan, T. R., Maina, J. M., Darling, E. S., Mireille, M., Guillaume, M., … Grimsditch, G. (2020) Large geographic variability in the resistance of corals to thermal stress. *Global Ecology Biogeography*, 29, 2229–2247. https://doi.org/10.1111/geb.13191

McClanahan, T. R. (2017) Changes in coral sensitivity to thermal anomalies. *Marine Ecology Progress Series,* 570, 71–85.

McClanahan, T. R., Ateweberhan, M., Graham, N. A. J., Wilson, S. K., Sebastian, C. R., Bruggemann, J. H. (2007) Western Indian Ocean coral communities: bleaching responses and susceptibility to extinction. *Marine Ecology Progress Series*, 337, 1-13.

McClanahan, T. R., Baird, A. H., Marshall, P. A., Toscano, M. A. (2004) Comparing bleaching and mortality responses of hard corals between southern Kenya and the Great Barrier Reef Australia. *Marine Pollution Bulletin*, 48, 327-35.

Muir, P. R., Marshall, P. A., Abdulla, A., Aguirre, J. D. (2017) Species identity and depth predict bleaching severity in reef building corals: Shall the deep inherit the reef? *Proceedings of the Royal Society B,* 284, 20171551. doi.org/10.1098/rspb.2017.1551

NOAA Coral Reef Watch (2017) *Daily 5 km satellite coral bleaching heat stress monitoring.* Retrieved from www.coralreefwatch.noaa.gov/satellite/index.php.

Pratchett, M. S., McCowan, D., Maynard, J. A., Heron, S. F. (2013) Changes in bleaching susceptibility among corals subject to ocean warming and recurrent bleaching in Moorea French Polynesia. *PLoS ONE,* 8, e70443.

Richards, Z. T. & Day, J. C. (2018) Biodiversity of the Great Barrier Reef—how adequately is it protected? *PeerJ* 6, e4747. doi.org/10.7717/peerj.4747.

Richards, Z. T., Berry, O., van Oppen, M. J. H. (2016) Cryptic genetic divergence within threatened species of *Acropora* coral from the Indian and Pacific Oceans. *Conservation Genetics,* 17, 577–591. doi.org/10.1007/s10592-015-0807-0

Sheppard, C., Sheppard, A., Fenner, D. (2020) Coral mass mortalities in the Chagos Archipelago over 40 years: Regional species and assemblage extinctions and indications of positive feedbacks. *Marine Pollution Bulletin,* 154, 111075. doi.org/10.1016/j.marpolbul.2020.111075

Skirving, W. J., Heron, S. F., Marsh, B. L., Liu, G., De La Cour, J. L., Geiger, E. F., Eakin, C. M. (2019) The relentless march of mass coral bleaching: a global perspective of changing heat stress. *Coral Reefs,* 38, 547–557. https://doi.org/10.1007/s00338-019-01799-4

Spalding, M. D., Ravilious, C., Green, E. P. (2001) *World atlas of coral reefs.* UNEP-WCMC.

Steneck, R. S., Arnold, S. N., Boenish, R., de León, R., Mumby, P. J., Rasher, D. B. & Wilson, M. W. (2019) Managing recovery resilience in coral reefs against climate-induced bleaching and hurricanes: a 15 year case study from Bonaire Dutch Caribbean. *Frontiers of Marine Science*, 6, 265. doi:10.3389/fmars.2019.00265

Thompson, D. M., van Woesik, R. (2009) Corals escape bleaching in regions that recently and historically experienced frequent thermal stress. *Proceedings of the Royal Society B,* 276, 2893–2901. doi:10.1098/rspb.2009.0591

van Hooidonk, R., Maynard, J., Tamelander, J., Gove, J., Ahmadia, G., Raymundo, L., … Planes, S. (2016) Local-scale projections of coral reef futures and implications of the Paris Agreement. *Science Reports* 6, 39666.

van Oppen, M. J. H., Oliver, J. K., Putnam, H. M., Gates, R. D. (2015) Building coral reef resilience through assisted evolution. *Proceedings National Academy of Sciences*, 112, 2307-23313.

van Oppen, M. H., Gates, R. D., Blackall, L., Cantin, N., Chakravarti, L. J., … Putnam, H. M. (2017) Shifting paradigms in restoration of the world's coral reefs. *Global Change Biology,* 23, 3437–3448. doi:10.1111/gcb.13647

Wooldridge, S. & Done, T. (2009) Improved water quality can ameliorate effects of climate change on corals. *Ecological Applications,* 19, 1492–1499.

FIGURES AND TABLES

Region	Proportion of Variation Explained		
	Phylogenetic	Contemporary	Local-Scale
Maldives	0.367 (0.065,0.638)	0.115 (0.047, 0.258)	0.032 (0.011,0.105)
GBR	0.454 (0.264, 0.658)	0.042 (0.094,0.103)	0.025 (0.006,0.095)

Table 1. Summary of phylogenetic mixed model analyses for the bleaching responses of scleractinian species in the Maldives Archipelago and Great Barrier Reef (GBR). The proportion of variation in bleaching explained by phylogenetically heritable (Phylogenetic), recent adaptation/acclimatization (Contemporary) and local-scale effects. Proportions given as posterior medians with 0.025 and 0.975 quantiles in brackets.

Figure 1. Locations where scleractinian coral bleaching responses were surveyed in the Great Barrier Reef and Maldives Archipelago. The severity of the bleaching at each location is given as peak degree heating weeks (DHW) in brackets.

Figure 2. Species′ bleaching responses (●) varied widely within many of the main scleractinian coral genera in both the Great Barrier Reef (■) and Maldives Archipelago (▢). Responses shown as posterior medians derived from phylogenetic mixed model analyses. The genera shown had more than two shared species between regions with sufficient data for analysis.

Figure 3. Susceptibility to moderate coral bleaching for scleractinian species in the Maldives Archipelago (inner ring) and Great Barrier Reef (outer ring) plotted onto the currently accepted phylogeny of the group (Huang and Roy, 2016). Susceptibility was categorized using quartiles of the species' posterior medians derived from separate phylogenetic mixed models that analysed the proportion of the population that was severely affected.

Figure 4. The bleaching responses of scleractinian species varied widely between regions, with phylogenetic mixed model analyses indicating that the variation had origins in both deep-time, phylogenetically heritable effects and short-term contemporary effects. These effects were poorly correlated between regions. Responses are shown as posterior medians (see Supporting Information, Figure S3 for details).