Mitofusin 1 and 2 regulation of mitochondrial DNA content is a critical determinant of

glucose homeostasis

Vaibhav Sidarala[1*]; Jie Zhu[1*]; Elena Levi-D'Ancona[1]; Gemma L. Pearson[1]; Emma C. Reck[1];

Emily M. Walker[1]; Brett A. Kaufman[2]; and Scott A. Soleimanpour[1,3,4]

[1]Division of Metabolism, Endocrinology & Diabetes and Department of Internal Medicine,

University of Michigan, Ann Arbor, MI 48105, USA.

[2]Vascular Medicine Institute, Division of Cardiology, Department of Medicine, University of

Pittsburgh School of Medicine, Pittsburgh, PA, 15260, USA

[3]Department of Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI

48105, USA

[4]VA Ann Arbor Healthcare System, Ann Arbor, MI 48105, USA.

*These authors contributed equally to this work

Corresponding Author

Scott A. Soleimanpour, MD, 1000 Wall Street, Brehm Tower Room 6329, Ann Arbor, MI 48105,

USA. Phone: (734) 763-0528 E-mail: ssol@med.umich.edu

ABSTRACT

The dynamin-like GTPases Mitofusin 1 and 2 (Mfn1 and Mfn2) are essential for mitochondrial

function, which has been principally attributed to their regulation of fission/fusion dynamics.

Here, we report that Mfn1 and 2 are critical for glucose-stimulated insulin secretion (GSIS)

primarily through control of mitochondrial DNA (mtDNA) content. Whereas Mfn1 and Mfn2

individually were dispensable for glucose homeostasis, combined Mfn1/2 deletion in β-cells

reduced mtDNA content, impaired mitochondrial morphology and networking, and decreased

respiratory function, ultimately resulting in severe glucose intolerance. Importantly, gene dosage

studies unexpectedly revealed that Mfn1/2 control of glucose homeostasis was dependent on

maintenance of mtDNA content, rather than mitochondrial structure. Mfn1/2 maintain mtDNA

content by regulating the expression of the crucial mitochondrial transcription factor Tfam, as

Tfam overexpression ameliorated the reduction in mtDNA content and GSIS in Mfn1/2-deficient

β-cells. Thus, the primary physiologic role of Mfn1 and 2 in β-cells is coupled to the preservation

of mtDNA content rather than mitochondrial architecture, and Mfn1 and 2 may be promising

targets to overcome mitochondrial dysfunction and restore glucose control in diabetes.

INTRODUCTION

Mitochondrial dynamics, the balance of fusion and fission of mitochondrial networks, is essential to control mitochondrial structure. Mitochondrial fission and fusion are governed by several key proteins that maintain mitochondrial quality control [1, 2]. Fission is primarily regulated by the GTPase dynamin-related protein 1 (Drp1), while fusion is controlled by both outer and inner mitochondrial membrane machinery. The dynamin-related large GTPase optic atrophy protein 1 (Opa1) controls inner membrane fusion, while outer membrane fusion is controlled by two GTPases known as Mitofusin 1 and 2 (Mfn1 and Mfn2). Mitofusin 1 and 2 share ~80% sequence similarity and contain homologous functional domains, suggesting they play both overlapping and unique roles in metabolic function largely ascribed to their control of mitochondrial structure [3].

Imbalances in mitochondrial dynamics in metabolically active tissues have been implicated in many human diseases, including neurodegenerative conditions, cancer, cardiovascular disease, and diabetes [2]. Mitochondria provide the energy necessary for β-cell insulin release [4, 5], and abnormalities in mitochondrial structure and bioenergetics have been observed in the β-cells of humans with type 2 diabetes (T2D; [6, 7]). Further, mitochondrial defects have been recently found to precede the development of T2D in human β-cells [8]. Thus, strategies to understand and overcome mitochondrial dysfunction in β-cells are of great appeal for the treatment of diabetes. However, the role of the core fission/fusion machinery in β-cells is less clear. Reduced β-cell *Mfn2* expression has been observed secondary to the deposition of toxic islet amyloid polypeptide oligomers observed in T2D as well as in mouse models of T2D [9, 10], yet the functions of Mfn1 and Mfn2 in β-cells *in vivo* are unexplored.

Here, we elucidate a key physiologic role for Mfn1 and 2 in the maintenance of glucose homeostasis that unexpectedly occurs through the preservation of β-cell mtDNA content, rather than mitochondrial structure. Utilizing several genetic mouse models, high resolution imaging, and transcriptomic profiling, we show that Mfn1 and 2 act in tandem to preserve β-cell

mitochondrial health and, consequently, glucose-stimulated insulin secretion (GSIS). Mfn1 and

2 maintain mtDNA content by preventing loss of the mitochondrial transcription factor Tfam,

which is a master regulator of mtDNA copy number control and mtRNA expression [11, 12]. Indeed,

adenoviral overexpression of Tfam ameliorates reductions in mtDNA content and GSIS in

Mfn1/2-deficient β-cells, illustrating the critical importance of mtDNA copy number control by

Mfn1/2 to promote glucose homeostasis.

RESULTS

Combined deficiency of Mfn1 and Mfn2 in β-cells leads to severe glucose intolerance and reduced glucose-stimulated insulin secretion

Mitochondrial health is vital to support β-cell insulin secretory responses to glucose or other nutrient stimuli. We hypothesized that Mfn1 and Mfn2 are required for GSIS in β-cells through their regulation of mitochondrial function. To test this hypothesis, we generated mice bearing β-cell specific deletion of Mfn1 alone (*Mfn1*^{loxP/loxP};*Ins1*-Cre), Mfn2 alone (*Mfn2*^{loxP/loxP};*Ins1*-Cre), and combined deletion of both Mfn1 and 2 (*Mfn1*^{loxP/loxP}*Mfn2*^{loxP/loxP};*Ins1*-Cre), hereafter known as β-Mfn1^{KO}, β-Mfn2^{KO}, and β-Mfn1/2^{DKO} mice, respectively. Conditional alleles targeted the canonical G-1 GTPase motif of Mfn1 (exon 4) or Mfn2 (exon 6) (Figure S1a; [13]). *Ins1*-Cre and floxed alone controls were phenotypically indistinguishable from one another, exhibiting no differences in body weight or glycemic control (Figure S1b-c). Therefore, they were pooled as controls for subsequent analyses. β-Mfn1^{KO}, β-Mfn2^{KO}, and β-Mfn1/2^{DKO} mice each exhibited an efficient reduction in Mfn1 and/or Mfn2 islet protein expression when compared to littermate controls (Ctrl; Figures 1a-b). We did not observe significant compensatory changes in Mfn2 expression in β-Mfn1^{KO} islets, nor in Mfn1 expression in β-Mfn2^{KO} islets (Figures 1a-b). Further, we did not observe differences in expression of the other core fission/fusion proteins, Drp1 and Opa1, following deletion of Mfn1 and/or Mfn2 (Figure S1d).

While loss of either Mfn1 or Mfn2 alone has been shown to impair glucose homeostasis in other metabolic tissues [3, 14], we were surprised to find that both β-Mfn1^{KO} and β-Mfn2^{KO} mice exhibited normal glucose homeostasis during an intraperitoneal glucose tolerance test (IPGTT; Figure 1c). We also observed no changes in insulin secretion in single β-Mfn1^{KO} or β-Mfn2^{KO} mice after glucose administration *in vivo* (Figure 1d) or in isolated islets (Figure 1e). However, combined deletion of both Mfn1 and Mfn2 resulted in severe glucose intolerance (Figure 1c; p <0.005 by one-way ANOVA vs. Ctrl), due to markedly reduced GSIS (Figures 1d-e and S1e).

To evaluate other etiologies of glucose intolerance in β-Mfn1/2^{DKO} mice, we first

measured β-cell mass. We observed no significant differences in β-cell mass between

genotypes (Figure 1f). There were also no differences in body weight or peripheral insulin

sensitivity between the groups (Figures S1f-g). Taken together, these studies suggest that Mfn1

and 2 are individually dispensable for glycemic control and GSIS, but together play vital

complementary roles for the maintenance of glucose tolerance and β-cell insulin release.

Mfn1 and 2 regulate β-cell mitochondrial structure and respiratory function

Given the connections between mitochondrial function and dynamics, we asked whether

mitochondrial respiration and architecture would be defective in the islets of β-Mfn1/2^{DKO} mice,

as has been reported in other models of Mfn1/2-deficiency [15, 16, 17, 18, 19]. Indeed, islets isolated

from β-Mfn1/2^{DKO} mice displayed a reduced glucose-stimulated oxygen consumption rate (OCR;

Figures 2a and S2a) when compared to Ctrl islets. We did not observe differences in glycolysis

as measured by extracellular acidification rate (ECAR; Figure S2a). Differences in glucose-

stimulated OCR between Ctrl and β-Mfn1/2^{DKO} islets were no longer observed following

exposure to the Complex III inhibitor antimycin A, suggesting that Mfn1/2-deficient islets

primarily possess a defect in β-cell mitochondrial respiration (Figures 2a and S2a).

Loss of key mediators of mitochondrial fusion disrupt mitochondrial network balance and

favor increased mitochondrial fission, often leading to the appearance of increased punctate or

spherical mitochondria [20]. We thus examined mitochondrial structure in Mfn1- and/or Mfn2-

deficient β-cells by several independent and complementary methods. First, we assessed β-cell

mitochondrial ultrastructure by transmission electron microscopy (TEM). While β-cells from all

groups had the expected appearance of electron dense insulin granules, β-cells from β-

Mfn1/2^{DKO} mice had reductions in mitochondrial aspect ratio, form factor, and perimeter, and

increased mitochondrial circularity (Figures 2b-c), indicating smaller, punctate mitochondria. We

observed no overt differences in ER morphology and proximity to mitochondria in β-cells from

control and β-Mfn1/2^{DKO} mice (Fig. S2b), however.

Secondly, we examined mitochondrial networking utilizing high-resolution deconvolution

imaging of the mitochondrial marker succinate dehydrogenase A (SDHA). We initially used

Imaris® imaging software to generate three-dimensional (3D) reconstructions of β-cell

mitochondria to grossly evaluate mitochondrial networking [21, 22]. Consistent with our findings

using TEM, the number of 3D mitochondrial networks appeared similar between Ctrl and single

β-Mfn1^{KO} or β-Mfn2^{KO} β-cells (Figure S2c). However, Mfn1/2-double deficient mice showed a

dramatic increase in the frequency of smaller β-cell mitochondrial networks (Figure 2d).

Next, we analyzed 3D deconvolution imaging of β-cell SDHA using the Mitochondria

Analyzer, a validated and unbiased quantification tool used to evaluate mitochondrial

morphology and network/branch complexity [22, 23]. This confirmed that measures of mitochondrial

morphology (reduced mitochondrial volume and increased sphericity) and network/branch

complexity (reduced branch number and branch junctions) were significantly impaired in β-

Mfn1/2^{DKO} mice, commensurate with an increase in punctate/spherical mitochondria due to

decreased mitochondrial fusion (Figure 2e). Together, these data demonstrate that together

Mfn1 and 2 maintain β-cell mitochondrial structure and function.

Mfn1 and 2 maintain β-cell mtDNA content

Mitochondrial DNA copy number is vital for β-cell function and glucose homeostasis [24, 25]. Thus,

we next examined whether mtDNA content was altered by loss of Mfn1 and/or Mfn2 in β-cells.

We observed a small, yet significant, reduction in mtDNA content in islets of β-Mfn2^{KO} mice that

was not observed in islets of β-Mfn1^{KO} mice (Figure 3a). Interestingly, mtDNA depletion

substantially worsened in the islets of β-Mfn1/2^{DKO} mice (Figure 3a), suggesting that loss of

Mfn1 exacerbates the depletion of mtDNA caused by Mfn2 deficiency. We next assessed

expression of the 13 mitochondrially encoded genes and observed a consistent and significant

reduction in mtRNA expression in β-Mfn2[KO] islets that was exacerbated in β-Mfn1/2[DKO] mice

(Figure 3b). Concordantly, examination of protein expression of mitochondrial OXPHOS

subunits by Western blot revealed reductions of the mitochondrially encoded subunits of

Complex IV, mt-Co1, and Complex III, mt-Cytb, in β-Mfn2[KO] islets and β-Mfn1/2[DKO] islets

(Figures 3c-d). We also observed significant reductions of nuclear encoded subunits of Complex

I (Ndufb8) in both β-Mfn2[KO] and β-Mfn1/2[DKO] islets and Complex III (Uqcrc2) in β-Mfn1/2[DKO]

islets, but no significant depletion of nuclear encoded Complex II (Sdhb) and Complex V (Atp5a)

(Figures 3c-d). Depletion of mtDNA content in β-Mfn1/2[DKO] islets did not appear to be secondary

to changes in mitochondrial mass, as we did not observe reductions in Atp5a (Figures 3c-d) or

in the mitochondrial outer membrane protein Tom20 (Figure 3e), a common marker of

mitochondrial mass [26]. These studies reveal that both Mfn1 and 2 (and Mfn2 alone to a lesser

degree) are necessary to maintain mtDNA copy number in β-cells.

Defects in β-cell mitochondrial fusion induce glucose intolerance upon loss of mtDNA

content

Our observations showing distinct and overlapping contributions of Mfn1 and 2 in the regulation

of β-cell mitochondrial function led us to hypothesize that loss of mitochondrial fusion would lead

to glucose intolerance due to mtDNA depletion. However, approaches to clarify the specific

contribution of mtDNA depletion to metabolic dysfunction following imbalances in mitochondrial

dynamics have been elusive. To this end, we generated animals harboring only a single

functional allele of either *Mfn1* (β-Mfn1[+/-]Mfn2[KO]) or *Mfn2* (β-Mfn1[KO]Mfn2[+/-]) to ascertain the

relative importance of mtDNA content and structure in the maintenance of glycemic control. For

robustness of interpretation, these experiments were performed alongside littermate controls

and β-Mfn1/2[DKO] mice as previously presented (Figures 1-3). As expected, neither β-Mfn1[+/-]

Mfn2[KO] nor β-Mfn1[KO]Mfn2[+/-] mice developed changes in β-cell mass, body weight, or insulin

sensitivity (Figures S3a-c). Interestingly, 3D quantification of mitochondrial morphology and

network integrity revealed that, similar to β-Mfn1/2^{DKO} mice, a single allele of either *Mfn1* or *Mfn2*

was not sufficient to maintain β-cell mitochondrial structure, leading to increases in punctate

mitochondria consistent with loss of mitochondrial fusion (Figures 4a-b).

Despite mitochondrial morphologic imbalances in both β-Mfn1$^{+/-}$Mfn2^{KO} and β-

Mfn1^{KO}Mfn2$^{+/-}$ mice, *only* β-Mfn1$^{+/-}$Mfn2^{KO} mice developed glucose intolerance and impaired

GSIS similar to that of β-Mfn1/2^{DKO} mice (Figures 4c-d). We also found significant reductions in

mtDNA content associated with the loss of GSIS observed in β-Mfn1$^{+/-}$Mfn2^{KO} and β-Mfn1/2^{DKO}

mice (Figure 4e). Thus, maintenance of a single functional allele of *Mfn2*, but not *Mfn1*, is able

to preserve glucose homeostasis, mtDNA content, and insulin secretion, despite being

insufficient to maintain β-cell mitochondrial network integrity. Importantly, these results disjoin β-

cell function from changes in mitochondrial morphology, and implicate mtDNA copy number as

the vital purveyor of glycemic control following loss of mitochondrial fusion.

Mfn1 and 2 deficiency induces a transcriptional signature consistent with defects in

mitochondrial metabolism and alterations in mtDNA replication

To begin to understand the mechanisms underlying control of mtDNA content by Mfn1 and 2,

we analyzed bulk RNA sequencing (RNAseq) data generated from islets of β-Mfn1^{KO}, β-Mfn2^{KO},

and β-Mfn1/2^{DKO} mice as well as littermate controls. Initial hierarchical clustering analyses based

on the top 500 most highly expressed genes revealed that only β-Mfn1/2^{DKO} islets clustered by

genotype, while we did not observe discrete clustering across Ctrl or single β-Mfn1^{KO} and β-

Mfn2^{KO} islets due to very few differences in expression between these groups (Figures 5a and

S4a-b). Gene ontology and pathway analyses of differentially expressed genes from β-

Mfn1/2^{DKO} islets revealed significant changes related to endocrine hormone/insulin secretion, β-

cell signature genes, as well as several metabolic pathways (Figures 5b and S4c-d), consistent

with the defects in β-cell function and mitochondrial metabolism observed in β-Mfn1/2^{DKO} mice.

The nearly 1,000 significantly dysregulated genes in β-Mfn1/2^{DKO} islets included

downregulation of mitochondrial encoded transcripts, as well as key β-cell signature genes such

as *Slc2a2* (Glut2) and *MafA* (Figures 5c and 5e). We also observed increases in expression of

regulators of triglyceride metabolism, including *Angptl4* and *Angptl8* [27], the

glycolytic/gluconeogenic enzyme *AldoB*, which is highly upregulated in models of β-cell

mitochondrial dysfunction and in T2D islets [28, 29], as well as *Nrf1*, a key regulator of

mitochondrial biogenesis (Figures 5c, 5e, and S4d; [30]). To resolve additional mitochondrial

specific effectors that were differentially regulated following the loss of Mfn1 and 2, we next

overlaid our RNAseq data on MitoCarta 2.0 [31], which contains a compendium of targets with

strong evidence for localization to the mitochondria (Figure 5d). This revealed increased

expression of several genes related to mtDNA replication machinery or associated with the

mitochondrial nucleoid (Figures 5d-e), including the critical mtDNA topoisomerase *Top1mt* [32],

the mitochondrial co-chaperone *Dnaja3*/*Tid1*, which maintains mtDNA integrity [33], and the

mitochondrial pyrimidine transporter *Slc25a33*, which is essential for mtDNA replication [34]. We

also observed upregulation in the mitochondrial AAA+ protease *LonP1*, which interacts with and

regulates the stability of numerous proteins at the mitochondrial nucleoid (Figures 5d-e; [35, 36, 37]).

Importantly, we did not observe changes in expression of other fission/fusion genes or

regulators of mtRNA transcription, mtDNA repair, sirtuins, or mitophagy (Figure S4d). Taken

together, these data indicate that Mfn1/2-deficiency induced a transcriptional signature

consistent with an activation of mtDNA replication to potentially compensate for β-cell mtDNA

depletion.

Mfn1 and 2 act through Tfam to regulate β-cell mtDNA copy number

The mechanisms underlying regulation of mtDNA by mitochondrial fusion have been

controversial, and previous studies implicate several pathways that could reduce mtDNA

content, including genome instability/increased mtDNA mutations, enhanced mitophagy, defects

in mitochondria-associated ER membrane (MAM) function, or impaired mtDNA replication [17, 19,

38, 39, 40, 41]. As our studies were completed in young mice, which bear infrequent mtDNA

mutations [42], and we did not observe differences in expression of mtDNA repair enzymes

(Figure S4d), we focused on assessments of mitophagy and mtDNA replication to better resolve

the significant reduction of mtDNA observed in β-Mfn1/2^{DKO} islets. We first assessed rates of

mitophagy following incubation of control and β-Mfn1/2^{DKO} islets with the cell permeable

Mtphagy dye, which did not reveal differences in mitophagy between genotypes (Figure S4e).

We also did not observe a difference in frequency of autophagosomes bearing mitochondria by

TEM, or changes in essential genes in the mitophagy pathway between control and β-Mfn1/2^{DKO}

islets (Figure 2b, S2b and S4d). As an initial screen for MAM function, we measured MAM-

dependent lipids, including cholesteryl esters and phospholipids [43, 44]. However, we did not

detect differences in total cholesterol, cholesteryl esters, or phosphatidylserine concentrations in

isolated islets of control and β-Mfn1/2^{DKO} mice (Figures S4f-g).

To assess regulators of β-cell mtDNA replication, we examined expression of the

replisome proteins Ssbp1, Twinkle, and Polg. We also measured expression of Tfam, a master

regulator of mtDNA copy number that promotes mitochondrial genome abundance via effects on

mtDNA packaging, stability, and replication, and whose levels are proportional to mtDNA copy

number [11, 45, 46, 47, 48]. While replisome proteins were largely unchanged, we observed a

significant reduction in Tfam protein in β-Mfn1/2^{DKO} islets (Figures 6a-b). *Tfam* mRNA

expression was unchanged in β-Mfn1/2^{DKO} mice (Figure S4d), suggesting the reduction in Tfam

protein occurred post-transcriptionally. We further observed a significant increase in the level of

LonP1 protein, which has been shown to lead to Tfam protein turnover [49, 50]. We additionally

confirmed reductions in Tfam and increases in LonP1 protein in β-Mfn1$^{+/-}$Mfn2^{KO} islets (Figure

S5), correlating with the mtDNA depletion, glucose intolerance, and impaired GSIS observed in

these mice (Figures 4c-e). In contrast, β-Mfn1^{KO}Mfn2$^{+/-}$ mice, which develop mitochondrial

structural alterations without glucose intolerance or mtDNA depletion, did not exhibit changes in

Tfam and LonP1 (Figure S5 and Figures 4c-e). These results suggest that changes in Tfam and

LonP1 expression following Mfn1/2 deficiency were not due to increases in punctate

mitochondria.

To test for defects in mtDNA replication, we next assessed initiation of first strand

replication. First strand synthesis in mitochondrial replication begins in the D-loop. However, a

single stranded (ss) molecule that forms the classic D-loop structure called 7S, which is a non-

replicative structure, shares this sequence. It is the elongation of the 7S sequence into the

nearby region (Cytb) that demonstrates committed initiation of replication. Here we assessed

the levels of ss and dsDNA at specific sequences by quantitative PCR to overcome expected

sample limitations from studies with mouse pancreatic islets. Notably, import of nascent Tfam

has been shown to increase the generation of 7S DNA due to its transcription initiation activity

[51]. We found a trend towards reduced total ss 7S sequences normalized to ds mtDNA in β-

Mfn1/2^{DKO} islets (Figure 6c). However, the rate of 7S sequence extension into committed first

strand replication was increased in β-Mfn1/2^{DKO} islets (Figure 6c), which could represent a

compensatory response to mtDNA depletion.

Increased first strand replication intermediates heighten the levels of displaced second

strand template, which is bound by the single strand binding protein Ssbp1 [52]. Consistent with

ss qPCR studies, in situ levels of Ssbp1 at the mtDNA nucleoid (detected by perinuclear anti-

DNA antibody staining), were significantly elevated in Mfn1/2-deficient β-cells (Figures 6d-e).

The increase in Ssbp1 localization to mitochondrial nucleoids could represent binding to a

higher quantity of single-stranded mtDNA replication intermediates and could also be consistent

with observations of impaired completion of mtDNA replication previously reported following

Mfn1/2-deficiency [19, 53]. We also observed Ssbp1 nuclear localization (Figures 6d and f), which

has been previously demonstrated following mitochondrial stress [54]. Further, we detected an

increase in the volume of mtDNA nucleoid structures in Mfn1/2-deficient β-cells by high-

resolution 3D imaging (Figures 6g-h), which are similar to previous reports of Tfam-deficiency

but could also represent impaired nucleoid distribution [19, 55, 56]. Taken together, our observations

of reduced mtDNA content, mtRNA levels, as well as changes in single-stranded mtDNA

products, Ssbp1 localization, and nucleoid size following Mfn1/2-deficiency suggest a role for

Tfam in mediating the effects of Mfn1/2 in β-cell function.

To test whether increasing mtDNA copy number was sufficient to improve β-cell function

in fusion-deficient islets, we overexpressed Tfam in β-Mfn1/2^{DKO} and littermate control islets. We

transduced intact islets using adenoviral vectors encoding the human form of TFAM, thus

allowing us to track overexpression of TFAM using human-specific TFAM anti-sera (Figures 7a-

b and S6a). Transduction efficiency in β-cells was estimated at ~40-50% (Figure S6a). Despite

limitations in transduction efficiency and expression, TFAM overexpression significantly

ameliorated the reductions in both GSIS and mtDNA copy number in β-Mfn1/2^{DKO} islets (Figures

7c-d and S6b). Therefore, Mfn1 and 2 direct β-cell function, at least in part, through the

regulation of Tfam-mediated mtDNA copy number control.

Mitofusin agonists restore mtDNA content independent of changes in mitochondrial

structure

We next asked if pharmacologically targeting mitofusins could restore mtDNA levels in models

of mitofusin deficiency independent of changes in mitochondrial structure. To this end, we

employed recently described tandem pharmacological agonists, (2-{2-[(5-cyclopropyl-4-phenyl-

4H-1,2,4-triazol-3-yl)sulfanyl]propanamido}-4H,5H,6H-cyclopenta[b]thiophene-3-carboxamide

and 1-[2-(benzylsulf-anyl)ethyl]-3-(2-methylcyclohexyl)urea (hereafter called Mfn agonists),

which together have been previously shown to improve mitochondrial morphology in either

Mfn1- or Mfn2-deficient fibroblasts [57]. These agonists require some residual mitofusin levels to

be effective [57]. Thus, we initially treated control and β-Mfn2^{KO} islets, which exhibit reduced

mtDNA content without changes in mitochondrial structure (Figures 3a, 2b-c, and 2e). To avoid

systemic effects of modifying mitochondrial fission/fusion balance in other metabolic tissues [1],

we utilized Mfn agonists on isolated islets. Interestingly, Mfn agonists restored mtDNA content

to baseline in β-Mfn2^{KO} islets without affecting mitochondrial morphology or inducing

transcriptional regulators of mitochondrial biogenesis (Figures 8a and S7a-b). As anticipated,

Mfn agonists were ineffective in the setting of combined absence of Mfn1 and 2 and were

unable to rescue reductions in mtDNA content or Tfam protein in β-Mfn1/2^{DKO} islets (Figures

S7c-d).

We next evaluated the effects of Mfn agonists in other models of mitofusin deficiency, as

Mfn2 expression is reduced in several models of β-cell dysfunction [9, 10]. We assessed the

expression of Mfn1 and 2 in islets isolated from the leptin receptor deficient *db*/*db* BKS mouse

model, which develops obesity, progressive β-cell mitochondrial and insulin secretory

dysfunction, and eventual β-cell failure [58, 59, 60, 61, 62]. We observed that Mfn2, but not Mfn1,

protein levels were reduced in islets of 10–12-week-old *db*/*db* mice when compared to *db*/+

controls (Figure 8b), measured at an age prior to β-cell failure [58, 59]. Similar to observations in β-

Mfn2^{KO} islets, *db*/*db* islets exhibited reduced mtDNA content without alterations in mitochondrial

structure (Figures 8c and 8e). Concordantly, treatment with Mfn agonists rescued mtDNA levels

in *db*/*db* islets, while leaving mitochondrial structure and islet architecture unchanged (Figures

8c-e). We also observed that Mfn agonists improved GSIS in the islets of *db*/*db* mice (Figure 8f).

Taken together, these results indicate that Mfn agonists restore mtDNA content and improve

GSIS independent of changes in mitochondrial structure. Moreover, these results extend the

association between mitofusin deficiency, mtDNA depletion, and impaired GSIS in β-cells.

323
324

DISCUSSION

Here, we identify that Mfn1 and 2 promote mitochondrial fitness to fuel glucose-stimulated insulin release in pancreatic β-cells and maintain glucose homeostasis. Mfn1 or Mfn2 are individually dispensable for β-cell function, but together coordinately control mitochondrial respiration, structure, and mtDNA content. Mfn1 and 2 were thought to regulate cellular metabolism principally through the control of mitochondrial architecture. Our single allele mouse models directly demonstrate that Mfn1/2-deficiency is coupled to glucose intolerance through mtDNA loss rather than mitochondrial structure. Further, we observe that increased mtDNA copy number following Tfam overexpression is capable of ameliorating impaired GSIS in fusion-deficient β-cells. Thus, our studies position the maintenance of mtDNA content as the principal task necessary for Mfn1 and 2 to regulate β-cell function.

Our description of a central role for Mfn1 and 2 in β-cells through mtDNA content rather than mitochondrial structure elucidates a previously unappreciated role for mitochondrial fusion on cellular metabolic function. Mfn1 and 2 act in a complementary manner to support mitochondrial metabolism from development to adulthood. Both germline and cardiomyocyte-specific deletion of Mfn1 and 2 are associated with developmental or early postnatal demise [16, 18], presumed to be primarily caused by defects in mitochondrial network dynamics. Later descriptions of reduced mtDNA levels in constitutive or inducible loss of Mfn1/2 in skeletal muscle did not discern the physiologic importance of mtDNA depletion versus structure following fusion-deficiency [15, 17]. Our results clarify a primary physiologic impact of mtDNA depletion rather than mitochondrial structure following Mfn1/2 deficiency in β-cells that may also underlie the physiologic importance of Mfn1 and 2 in other metabolic tissues. To our knowledge, this is the first time that these functions of Mfn1/2 have been separated in a physiological context.

We observe that β-Mfn1/2^{DKO} islets attempt unsuccessfully to compensate for mtDNA depletion. Our analyses of first strand replication suggest that Mfn1/2-deficient β-cells increase

the activation of early steps in mtDNA replication, which is supported by increased Ssbp1

localization to nucleoids to potentially bind newly formed ss-mtDNA. Despite these changes,

mtDNA content remains low in Mfn1/2-deficient β-cells, suggesting that the increased

commitment to replication is insufficient. Indeed, increased nucleoid Ssbp1 may be engaged in

binding abnormal single-stranded replication products due to fusion and replication defects [19, 53].

The increased expression of *Top1mt* found in Mfn1/2-deficient β-cells (Figure 5e) could also

represent a response to increased mitochondrial replication stress [63].

Loss of Mfn1/2 in β-cells leads to severe hyperglycemia, reduced GSIS, and mtDNA

depletion, which bears similarity to a previous report of selective Tfam deficiency in β-cells [25].

Notably, previously described β-cell specific Tfam knockout mice developed progressive loss of

β-cell mass with age [25]. We found that β-cell mass was unchanged in young β-Mfn1/2^{DKO} mice,

but it is possible that they would similarly develop glucotoxicity and an age-related loss of β-cell

mass over time. Alternatively, complete depletion of mtDNA, which occurs in the setting of

Tfam-deficiency but does not occur in β-Mfn1/2^{DKO} mice, might be required for loss of β-cell

mass. Furthermore, the transgenic *RIP2*-Cre used in the β-cell Tfam knockout model (in

contrast to the highly specific *Ins1*-Cre knockin strain we used in β-Mfn1/2^{DKO} mice) is known to

result in off-target recombination[25, 64], which may in part underlie phenotypic differences

between the models. In future experiments, it will be intriguing to examine Tfam and Mfn1/2-

deficient mice generated using the same Cre-recombinase strains to determine the contribution

of Tfam-dependent and independent functions on Mfn1/2 in β-cell health and function.

A related concern in the Mfn1/2-deficiency models is the use of a constitutive Cre strain,

which may yield phenotypic defects due to recombination during β-cell development. However,

an independent group found that mice with inducible Mfn1/2-deficiency only in adult β-cells

developed a highly similar phenotype to constitutive β-Mfn1/2^{DKO} mice [65], suggesting that

developmental defects are unlikely to be a major contributor to the phenotype we observe.

While our data suggest that Tfam mediates the effects of Mfn1/2 on mtDNA content and

GSIS in β-cells, the mechanisms for post-transcriptional loss of Tfam in β-Mfn1/2[DKO] islets are

unclear. Loss of Tfam may be due to increased LonP1 expression, which is known to degrade

Tfam [49, 50]. Further support for a role for LonP1 as a contributor to cellular dysfunction include

studies demonstrating reduction of nuclear-encoded subunits of Complex I and III following

increases in LonP1 expression [66, 67, 68], an effect we also observe following Mfn1/2-deficiency

(Figure 3d). Mfn1/2-deficiency has also been linked to reductions in nuclear-encoded Complex I

and III subunits via instability in respiratory supercomplexes, which have been observed

following LonP1 overexpression [15, 66, 67, 68]. However, the function and transcriptional regulation

of *LonP1* in β-cells are unknown and require future study.

Mfn2 is frequently observed at MAMs [69, 70]. MAMs are essential for communication and

Ca^{2+} transfer between the ER and mitochondria, are important for β-cell function and GSIS, and

have been suggested to regulate mtDNA replication [39, 40, 41, 71, 72, 73]. Thus, alterations in MAMs in

β-Mfn1/2[DKO] islets could lead to impaired mtDNA replication. Our initial functional assessments

of MAMs by measurement of MAM-dependent lipids did not reveal differences between control

and β-Mfn1/2[DKO] islets (Figures S4f-g). We also did not observe ER dilation or overt changes in

mitochondrial-ER proximity upon analysis of transmission EM in β-Mfn1/2[DKO] islets (Figures 2b

and S2b). Additionally, a recent report found that inducible loss of Mfn1/2 in adult β-cells did not

alter MAM-dependent ER calcium release [65]. Despite these findings, our results do not exclude

the potential that impaired MAM function at β-cell mtDNA nucleoids contributes to the

impairments in mtDNA replication and content we observe. Indeed, our results demonstrating

the importance for a single allele of *Mfn2* over *Mfn1* in the maintenance of glycemic control and

mtDNA content could suggest a role for Mfn2-dependent MAM function in β-cell health and

function, as Mfn1 is not found at MAMs. Alternatively, the regulation of MAMs by Mfn2, as well

as the MAM-dependent role of Mfn2 in mtDNA replication, remains controversial [74, 75]. Recent

work also implicates mitochondrial fusion as necessary to maintain the stoichiometry of the

protein components within the mtDNA replisome [19]. Future studies will be valuable to dissect the

molecular details underlying the connections between mitochondrial fusion, MAMs, and control

of β-cell mtDNA replication.

Mitochondrial structural defects and age-related mtDNA depletion combine to impair β-

cell function to increase T2D risk [6, 7, 76, 77, 78], thus, improving mitochondrial function would have

significant translational benefits to restore β-cell health in diabetes. Our studies demonstrate the

feasibility of targeting mtDNA copy number to improve GSIS in the setting of mitofusin

deficiency. While our results in *db/db* islets are promising, loss of the leptin receptor could lead

to mitochondrial defects that are particularly amenable to improvement by Mfn agonists. Thus, it

will be intriguing to determine if Mfn agonists can also reverse β-cell dysfunction in diet-induced

models of obesity *in vivo*. Further, recently identified compounds targeting pyrimidine

metabolism to induce *Mfn1/2* transcription [79] will be of interest to determine if they can improve

β-cell function and mtDNA content. Our observation of the upregulation of mitochondrial

pyrimidine transporter *Slc25a33* in β-Mfn1/2^{DKO} islets is consistent with links between pyrimidine

metabolism and mitochondrial fusion, which suggest these compounds could hold promise to

enhance β-cell mitochondrial function. As expression of MFN1 and MFN2 is reduced in

cardiomyocytes and skeletal muscle of patients with diabetes, respectively [80, 81, 82], activators of

Mfn1/2 may have additional benefits beyond improving β-cell function. Future studies will be

required to fully characterize the efficacy of overcoming mitochondrial defects *in vivo* to treat

diabetes.

METHODS

Animals

$Mfn1^{loxP/loxP}$ and $Mfn2^{loxP/loxP}$ mice possessing loxP sites flanking exon 4 of the *Mfn1* gene and exon 6 of the *Mfn2* gene, respectively, were purchased from Jackson Laboratories [13]. $Mfn1^{loxP}$ and $Mfn2^{loxP}$ mice were mated to *Ins1*-Cre [83] to generate experimental groups. *Ins1*-Cre–alone, $Mfn1^{loxP/loxP}$ and $Mfn2^{loxP/loxP}$ mice were phenotypically indistinguishable from each other and combined as controls (Ctrl; see Figures S1b-c). *Ins1*-Cre–alone were also phenotypically indistinguishable from wild-type C57BL/6N controls, consistent with previous reports from our group and others [21, 83]. All animals were maintained on the C57BL/6N background. *db*/+ and *db*/*db* mice, maintained on the BKS background, were purchased from Jackson Laboratories. Animals were housed on a standard 12-hour light/12-hour dark cycle at room temperature and 30-70% humidity, with ad libitum access to food and water. All mice were fed PicoLab® Laboratory Rodent Diet 5L0D (LabDiet) containing 28% protein, 13% fat, and 57% carbohydrates. All studies and endpoints in all figures were completed using both male and female mice, and results from both sexes were combined in all experimental groups. Animal studies were approved by the University of Michigan Institutional Animal Care and Use Committee.

Mitochondrial respirometry

Islet respirometry was measured using an XF96 extracellular flux analyzer (Seahorse Bioscience) according to manufacturer's instructions. Briefly, 20 similarly sized islets were plated per well of a Seahorse spheroid plate pre-treated with CellTak (Corning), similar to previously published approaches [84]. Prior to respirometry profiling, islets were incubated for 1 hour in an atmospheric CO_2 incubator at 37°C and supplemented with mouse islet culture media [85, 86] comprised of pH 7.4 unbuffered RPMI1640 media (Seahorse Bioscience) and 2mM glucose prior to flux analyzer assays. Analysis was performed at baseline, upon exposure to

20mM glucose, and again after treatment with 2 μM antimycin A. OCR measurements were

normalized to islet number confirmed by light microscopy (Zeiss Stemi 2000) after the

completion of respirometry assays.

Transmission electron microscopy

Mouse islets were pelleted, fixed in 3% glutaraldehyde and 3% paraformaldehyde in 0.1 M

Cacodylate buffer (CB; pH 7.2) overnight at 4°C. Islets were then embedded in 2% agarose as

described previously [87]. Samples were then subjected to osmification in 1.5% $K_4Fe(CN)_6$ + 2%

OsO_4 in 0.1 CB for 1 h, dehydrated by serial washes in EtOH (30%, 50%, 70%, 80%, 90%, 95%

and 100%) and embedded in Spurr's resin by polymerization at 60° C for 24h. Polymerized

resins were then sectioned at 90nm thickness using a Leica EM UC7 ultramicrotome and

imaged at 70kV using a JEOL 1400 TEM equipped with an AMT CMOS imaging system.

Mitochondrial structures (Aspect Ratio, Perimeter, Circularity) were analyzed and quantified by

ImageJ. Form factor assessments were analyzed by ImageJ and quantified as described [22].

Imaging of islet and organelle morphology and subcellular localization analysis

Mitochondrial morphology and subcellular localization analyses were performed on immuno-

stained paraffin-embedded mouse pancreas tissue sections and dissociated islet cells as

previously described [21]. Briefly, pancreatic tissue sections were fixed in 4% PFA overnight,

embedded in paraffin and sectioned. Following dewaxing and rehydration, antigen retrieval was

performed with 10mM citric acid (pH 6.0). Dissociated islet cells were fixed in 4% PFA for 15

minutes and centrifuged onto glass slides. Pancreatic sections and dissociated islets were then

blocked using 5% donkey serum for 1hr at room temperature followed by immunostaining. Islet

architecture was analyzed using isolated mouse islets fixed in 4% PFA overnight followed by a

20 min incubation with methylene-blue solution (Sigma) and OCT embedding. Cryo-sectioned

islets were rinsed in 75% ethanol for 5 minutes to remove methylene blue, permeabilized with

0.2% Triton-X 100 for 15 minutes, blocked using 5% donkey serum for 1h and immunostained.

Images were captured with an IX81 microscope (Olympus) using an ORCA Flash4

CMOS digital camera (Hamamatsu). Immunostained pancreatic sections, cryo-sectioned and

dispersed mouse islets were captured with Z-stack images and subjected to deconvolution

(CellSens; Olympus). Mitochondrial morphology and nucleoids were visualized using 3D-

renderings generated with Imaris® imaging software (Bitplane). Quantitative 3D assessments of

mitochondrial morphology and network were performed on ImageJ using Mitochondria Analyzer

plugin [22]. Co-localization analyses were performed on Z-stack images of immunostained

dissociated islet cells using the Coloc2 plugin on ImageJ. Changes in relative nucleoid size

were quantified from 3D deconvolution images of immuno-stained dissociated islet cells,

detected by perinuclear anti-DNA antibody staining, using ImageJ.

mtDNA content and replication assays

Relative mtDNA content by qPCR with SYBR-based detection (Universal SYBR Green

Supermix; Biorad) was conducted as previously described [88] following DNA isolation with the

Blood/Tissue DNeasy kit (Qiagen) according to manufacturer's protocols. First strand mtDNA

replication levels were measured by qPCR following digestion by Mnl I, which site specifically

digest double stranded DNA [89]. Briefly, primers specific to 7S mtDNA (5'-

ACTCTTCTCTTCCATATGACTATCCC-3' and 5'-GGCCCTGAAGTAAGAACCAGATGT-3'),

whose amplicon contains a Mnl I site, were used to quantify relative single-stranded mtDNA

forming the D-loop. Single-stranded DNA generated from the extension of 7S region was

measured using primers specific to CYTB (5'-TGCATACGCCATTCTACGCTCA-3' and 5'-

GTGATTGGGCGGAATATTAGGCTTC-3'), which also contains an Mnl I site. Primers specific to

COXI   (5'-GCAGGAGCATCAGTAGACCTAACA-3'   and   5'-

GCGGCTAGCACTGGTAGTGATAAT-3'), whose amplicon does not contain an Mnl I site, were

used to quantify non-replicating double stranded mtDNA. The ratio of 7S or CYTB amplicons to

the COXI amplicon were then calculated after qPCR of Mnl I digested DNA to determine the

relative amount of single stranded mtDNA at the D-loop strand or initiation of first strand

replication, respectively.

RNA sequencing, differential expression, and functional analysis

Sequencing was performed by the Advanced Genomics Core at University of Michigan Medical

School. Total RNA was isolated, and DNase-treated using commercially available kits (Omega

Biotek and Ambion, respectively). Libraries were constructed and subsequently subjected to

151bp paired-end cycles on the NovaSeq-6000 platform (Illumina). FastQC (v0.11.8) was used

to ensure the quality of data. Reads were mapped to the reference genome GRCm38

(ENSEMBL), using STAR (v2.6.1b) and assigned count estimates to genes with RSEM (v1.3.1).

Alignment options followed ENCODE standards for RNA-seq. FastQC was used in an additional

post-alignment step to ensure that only high-quality data were used for expression quantitation

and differential expression. Data were pre-filtered to remove genes with 0 counts in all samples.

Differential gene expression analysis was performed using DESeq2, using a negative binomial

generalized linear model (thresholds: linear fold change >1.5 or <-1.5, Benjamini-Hochberg FDR

(P_{adj}) <0.05). Plots were generated using variations of DESeq2 plotting functions and other

packages with R version 3.3.3. Genes were annotated with NCBI Entrez GeneIDs and text

descriptions. Functional analysis, including candidate pathways activated or inhibited in

comparison(s) and GO-term enrichments, was performed using iPathway Guide (Advaita). The

RNA sequencing data generated in this study have been deposited in the Gene Expression

Omnibus (GEO) database under accession code GSE194204.

Adenoviral transfection

Isolated mouse islets were distended with Accutase for 1 min at 37° C as described previously

[90]. Islets were then transduced with 0.15 MOI of Ad.EV (expressing an empty vector) or

Ad.hTFAM (expressing human-specific TFAM) for 48 hrs (viral particles purchased from Vector

Biolabs). β-cell transduction was evaluated on dissociated islet cells by immunostaining with

Pdx1 and human-specific TFAM anti-sera.

Flow cytometry

Following isolation and culture, live mouse islets were exposed to 100 nM Mtphagy dye (Dojindo Molecular Technologies) for 3 hours to assess time-dependent accumulation of mitochondria to acidic organelles by the relative fluorescence intensity of the dye per cell as described [91, 92]. Islets were then dissociated into single cells, stained with DAPI (Thermo Fisher Scientific) and Fluozin-3 (Thermo Fisher Scientific), and resuspended in phenol red–free islet culture medium as previously described [21]. Samples were analyzed on an LSR Fortessa flow cytometer (BD Biosciences). Single cells were gated using forward scatter and side scatter (FSC and SSC, respectively) plots, DAPI staining was used to exclude dead cells, and Fluozin-3 was used to identify β-cells as previously described [21, 93]. Mtphagy measurements in β-cells were made using 488 nm excitation laser with a 710 nm emission filter and analyzed using FlowJo (Tree Star Inc.). A total of 5,000 β-cells was quantified from each independent islet preparation. Gating strategies with representative plots are provided in Supplementary Figure S8.

Glucose and insulin measurements

IPGTT, ITT, and serum insulin measurements were performed as described previously [85, 86]. Briefly, mice were fasted for 6 hrs prior to glucose and insulin measurements. For IPGTT or ITT, mice were injected intraperitoneally with 2 g/kg glucose or 0.75 U/kg Insulin (Novolin R), respectively. Tail blood was collected for measurements of blood glucose. For *in vivo* insulin release measurements, tail blood was collected following glucose injection, centrifuged at 2000 rpm for 10 min at 4 °C and stored at −20 °C. Static insulin secretion assays on isolated islets were performed following 1hr incubations with 2mM and 16.7mM glucose in HEPES-supplemented KRB buffer containing 135 mM NaCl, 4.7 mM KCl, 1.2 mM KH_2PO_4, 5 mM $NaHCO_3$, 1.2 mM $MgSO_4.7H_2O$, 1 mM $CaCl_2$, 10 mM HEPES and 0.1% BSA (pH 7.4). Samples were analyzed for insulin release using Mouse Ultrasensitive Insulin ELISA (ALPCO) as per the manufacturer's protocol.

Islet isolation and cell culture

Mouse islets were isolated and cultured as previously described [85, 86]. Cell treatments included

DMSO (Fisher) and antimycin A (Sigma). Mitofusin agonist treatment included an equimolar

mixture of two previously described compounds (0.5 µM each; [57]), which were administered for

24 hours (2-{2-[(5-cyclopropyl-4-phenyl-4H-1,2,4-triazol-3-yl)sulfanyl]propanamido}-4H,5H,6H-

cyclopenta[b]thiophene-3-carboxamide, CAS# 920868-45-7 and 1-[2-(benzylsulfanyl)ethyl]-3-(2-

methylcyclohexyl)urea, CAS# 1007818-44-1; Enamine Ltd.).

Western blotting, quantitative PCR, and immunostaining

All assays were performed as previously described [21, 85, 86]. Mouse pancreatic islets were lysed

with radioimmunoprecipitation assay buffer containing protease and phosphatase inhibitors

(Calbiochem), and insoluble material was removed by centrifugation. Equal amounts of proteins

were resolved on 4%–20% gradient Tris-glycine gels (Bio-Rad) and transferred to nitrocellulose

membranes (Bio-Rad). Membranes were then blocked in 5% milk for 1hr and immunoblotting

was performed using Cyclophilin B (1:5000; ThermoFisher, Catalog# PA1-027A), Drp1 (1:1000;

CST #8570), Lonp1 (1:1000, ProteinTech, Catalog# 15440-1-AP), Mfn1 (1:1000; Abcam,

Catalog# ab126575), Mfn2 (1:1000; Abcam, Catalog# ab56889), mt-Cytb (1:500; ProteinTech,

Catalog# 55090-1-AP), Opa1 (1:1000, BD Transduction laboratories, Catalog# 612606), Total

Oxphos antibody cocktail (1:1000; Abcam, ab110413), Polg (1:1000, Abcam, Catalog#

ab128899), Ssbp1 (1:250; Atlas antibodies; Catalog# HPA002866), Anti-human TFAM (1:1000,

PhosphoSolutions, Catalog# 1999-hTFAM), Anti-mouse TFAM (1:1000; PhosphoSolutions

2001-TFAM), Tom20 (1:1000; Cell Signaling Technology, Catalog# 42406), Twnk (1:1000,

ProteinTech, Catalog# 13435-1-AP), Vinculin (1:5000; Millipore, Catalog# CP74) and species-

specific HRP-conjugated secondary antibodies (Vector Laboratories). Immunostaining on

pancreatic sections, cryo-sectioned and dispersed mouse islets was performed using Anti-DNA

(1:50; American Research Products, Catalog# PA1-027A), Insulin (1:250; Dako, Catalog#

A0564), Insulin (1:100, Abcam, Catalog# ab7842), Glucagon (1:2000; SantaCruz, Catalog# sc-

13091), Pdx1 (1:250; Abcam, Catalog# ab47383), SDHA (1:100, Abcam, Catalog# ab14715),

Ssbp1 (1:250; Atlas antibodies, Catalog# HPA002866), Somatostatin (1:500, Abcam, Catalog#

ab30788) and species-specific Cy2, Cy3 and Cy5-conjugated secondary antibodies (Jackson

Immunoresearch). β-cell mass was quantified as total insulin-positive area/total pancreatic area

multiplied by pancreatic wet weight using stitched images of complete pancreatic sections [86]. All

antibodies used for Western blotting and immunostaining are listed in Supplementary Table 1.

RNA samples were reverse transcribed using High-Capacity cDNA Reverse

Transcription Kit (Thermo Fisher Scientific). Quantitative reverse transcription PCR (qRT-PCR)

was performed with SYBR-based detection (Universal SYBR Green Supermix; Biorad) using

primers for *Ppargc1a* (5'-ACTATGAATCAAGCCACTACAGAC-3' and 5'-

TTCATCCCTCTTGAGCCTTTCG-3'), *Nrf1* (5'-ACAGATAGTCCTGTCTGGGGAAA-3' and 5'-

TGGTACATGCTCACAGGGATCT-3') and *Tfam* (5'-AGCTTCCAGGAGGCAAAGGATGAT-3'

and 5'-ACTTCAGCCATCTGCTCTTCCCAA-3').

Cholesterol esterification and phosphatidylserine assay

Total amounts of free cholesterol and cholesterol esters were determined in mouse islet lysates

as per the manufacturer's instructions (Promega). Phosphatidylserine levels were measured in

mouse islet lysates as per the manufacturer's instructions (LSBio).

Statistics

In all figures, data are presented as means ± SEM, and error bars denote SEM, unless

otherwise noted in the legends. Statistical comparisons were performed using unpaired two-

tailed Student's t-tests, one-way or two-way ANOVA, followed by Tukey's or Sidak's post-hoc

test for multiple comparisons, as appropriate (GraphPad Prism). A *P* value < 0.05 was

considered significant.

Data Availability

The authors declare that data supporting the findings of this study are available within the article

and its supplementary information files or from the corresponding author (S.A.S.) upon request.

RNA sequencing data used for differential gene expression-, pathway- and GO- enrichment

analyses have been deposited in the GEO database under the accession number GSE194204.

The reference genome used for RNA sequencing was obtained from ENSEMBL

(Mus_musculus - Ensembl genome browser 105). All uncropped western blots and data values

for all figures are provided in the source data file.

Code Availability

High-throughput RNA sequencing data was analyzed using DESeq2 (v1.26.0) to determine

differential gene expression. iPathway Guide (Advaita version 1910) was used to perform

pathway and GO-term analysis.

614   **REFERENCES**
615
616 1.   Sebastian D, Palacin M, Zorzano A. Mitochondrial Dynamics: Coupling Mitochondrial
617   Fitness with Healthy Aging. *Trends Mol Med* **23**, 201-215 (2017).
618
619 2.   Westermann B. Mitochondrial fusion and fission in cell life and death. *Nat Rev Mol Cell
620   Biol* **11**, 872-884 (2010).
621
622 3.   Schrepfer E, Scorrano L. Mitofusins, from Mitochondria to Metabolism. *Mol Cell* **61**, 683-
623   694 (2016).
624
625 4.   Kaufman BA, Li C, Soleimanpour SA. Mitochondrial regulation of beta-cell function:
626   maintaining the momentum for insulin release. *Molecular aspects of medicine* **42**, 91-104
627   (2015).
628
629 5.   Maechler P. Mitochondrial function and insulin secretion. *Mol Cell Endocrinol* **379**, 12-18
630   (2013).
631
632 6.   Anello M*, et al.* Functional and morphological alterations of mitochondria in pancreatic
633   beta cells from type 2 diabetic patients. *Diabetologia* **48**, 282-289 (2005).
634
635 7.   Masini M*, et al.* Ultrastructural alterations of pancreatic beta cells in human diabetes
636   mellitus. *Diabetes Metab Res Rev* **33**, (2017).
637
638 8.   Wigger L*, et al.* Multi-omics profiling of living human pancreatic islet donors reveals
639   heterogeneous beta cell trajectories towards type 2 diabetes. *Nat Metab* **3**, 1017-1031
640   (2021).
641
642 9.   Safia Costes TG, Alexandra E Butler, Sam Sereda, Orian Shirihai and Peter Cawood
643   Butler. Disruption of the Endoplasmic Reticulum-Mitochondria Interface By Toxic IAPP
644   Oligomers May Compromise Beta-Cell Mitochondrial Network Integrity and Function in
645   Type 2 Diabetes. In: *The Endocrine Society Annual Meeting* (ed Society E) (2015).
646

10.   Stoehr JP, *et al.* Genetic obesity unmasks nonlinear interactions between murine type 2
diabetes susceptibility loci. *Diabetes* **49**, 1946-1954 (2000).

11.   Campbell CT, Kolesar JE, Kaufman BA. Mitochondrial transcription factor A regulates
mitochondrial transcription initiation, DNA packaging, and genome copy number.
Biochim Biophys Acta **1819**, 921-929 (2012).

12.   Filograna R, Mennuni M, Alsina D, Larsson NG. Mitochondrial DNA copy number in
human disease: the more the better? *FEBS Lett*, (2020).

13.   Chen H, McCaffery JM, Chan DC. Mitochondrial fusion protects against
neurodegeneration in the cerebellum. *Cell* **130**, 548-562 (2007).

14.   Liesa M, Shirihai OS. Mitochondrial dynamics in the regulation of nutrient utilization and
energy expenditure. *Cell Metab* **17**, 491-506 (2013).

15.   Bell MB, Bush Z, McGinnis GR, Rowe GC. Adult skeletal muscle deletion of Mitofusin 1
and 2 impedes exercise performance and training capacity. *J Appl Physiol (1985)* **126**,
341-353 (2019).

16.   Chen H, Detmer SA, Ewald AJ, Griffin EE, Fraser SE, Chan DC. Mitofusins Mfn1 and
Mfn2 coordinately regulate mitochondrial fusion and are essential for embryonic
development. *J Cell Biol* **160**, 189-200 (2003).

17.   Chen H, *et al.* Mitochondrial fusion is required for mtDNA stability in skeletal muscle and
tolerance of mtDNA mutations. *Cell* **141**, 280-289 (2010).

18.   Papanicolaou KN, *et al.* Mitofusins 1 and 2 are essential for postnatal metabolic
remodeling in heart. *Circ Res* **111**, 1012-1026 (2012).

19.   Silva Ramos E, *et al.* Mitochondrial fusion is required for regulation of mitochondrial DNA
replication. *PLoS Genet* **15**, e1008085 (2019).

20.   Twig G, *et al.* Fission and selective fusion govern mitochondrial segregation and
elimination by autophagy. *The EMBO journal* **27**, 433-446 (2008).

21.   Sidarala V, *et al.* Mitophagy protects beta cells from inflammatory damage in diabetes.
JCI Insight **5**, (2020).

22.   Chaudhry A, Shi R, Luciani DS. A pipeline for multidimensional confocal analysis of
mitochondrial morphology, function, and dynamics in pancreatic beta-cells. *Am J Physiol*
Endocrinol Metab **318**, E87-E101 (2020).

23.   Hemel I, Engelen BPH, Luber N, Gerards M. A hitchhiker's guide to mitochondrial
quantification. *Mitochondrion* **59**, 216-224 (2021).

24.   Mulder H. Transcribing beta-cell mitochondria in health and disease. *Molecular*
metabolism **6**, 1040-1051 (2017).

25.   Silva JP, *et al.* Impaired insulin secretion and beta-cell loss in tissue-specific knockout
mice with mitochondrial diabetes. *Nature genetics* **26**, 336-340 (2000).

26. Narendra D, Tanaka A, Suen DF, Youle RJ. Parkin is recruited selectively to impaired
mitochondria and promotes their autophagy. *J Cell Biol* **183**, 795-803 (2008).
701

702 27. Kersten S. Physiological regulation of lipoprotein lipase. *Biochim Biophys Acta* **1841**,
703   919-933 (2014).
704

705 28. Haythorne E*, et al.* Diabetes causes marked inhibition of mitochondrial metabolism in
706   pancreatic beta-cells. *Nat Commun* **10**, 2474 (2019).
707

708 29. Gerst F*, et al.* The Expression of Aldolase B in Islets Is Negatively Associated With
709   Insulin Secretion in Humans. *J Clin Endocrinol Metab* **103**, 4373-4383 (2018).
710

711 30. Scarpulla RC. Transcriptional paradigms in mammalian mitochondrial biogenesis and
712   function. *Physiol Rev* **88**, 611-638 (2008).
713

714 31. Calvo SE, Clauser KR, Mootha VK. MitoCarta2.0: an updated inventory of mammalian
715   mitochondrial proteins. *Nucleic Acids Res* **44**, D1251-1257 (2016).
716

717 32. Zhang H*, et al.* Human mitochondrial topoisomerase I. *Proc Natl Acad Sci U S A* **98**,
718   10608-10613 (2001).
719

720 33. Ng AC, Baird SD, Screaton RA. Essential role of TID1 in maintaining mitochondrial
721   membrane potential homogeneity and mitochondrial DNA integrity. *Mol Cell Biol* **34**,
722   1427-1437 (2014).
723

724 34. Di Noia MA*, et al.* The human SLC25A33 and SLC25A36 genes of solute carrier family
725   25 encode two mitochondrial pyrimidine nucleotide transporters. *J Biol Chem* **289**,
726   33137-33148 (2014).
727

728 35. Zurita Rendon O, Shoubridge EA. LONP1 Is Required for Maturation of a Subset of
729   Mitochondrial Proteins, and Its Loss Elicits an Integrated Stress Response. *Mol Cell Biol*
730   **38**, (2018).
731

732 36. Venkatesh S, Lee J, Singh K, Lee I, Suzuki CK. Multitasking in the mitochondrion by the
733   ATP-dependent Lon protease. *Biochim Biophys Acta* **1823**, 56-66 (2012).
734

735 37. Lu B*, et al.* Roles for the human ATP-dependent Lon protease in mitochondrial DNA
736   maintenance. *J Biol Chem* **282**, 17363-17374 (2007).
737

738 38. Sebastian D*, et al.* Mfn2 deficiency links age-related sarcopenia and impaired autophagy
739   to activation of an adaptive mitophagy pathway. *The EMBO journal* **35**, 1677-1693
740   (2016).
741

742 39. Friedman JR, Lackner LL, West M, DiBenedetto JR, Nunnari J, Voeltz GK. ER tubules
743   mark sites of mitochondrial division. *Science* **334**, 358-362 (2011).
744

745 40. Lewis SC, Uchiyama LF, Nunnari J. ER-mitochondria contacts couple mtDNA synthesis
746   with mitochondrial division in human cells. *Science* **353**, aaf5549 (2016).
747

748 41.   Murley A, *et al.* ER-associated mitochondrial division links the distribution of
749   mitochondria and mitochondrial DNA in yeast. *Elife* **2**, e00422 (2013).
750
751 42.   Ma H, *et al.* Germline and somatic mtDNA mutations in mouse aging. *PLoS One* **13**,
752   e0201304 (2018).
753
754 43.   Area-Gomez E. Assessing the function of mitochondria-associated ER membranes.
755   *Methods Enzymol* **547**, 181-197 (2014).
756
757 44.   Vance JE. Molecular and cell biology of phosphatidylserine and
758   phosphatidylethanolamine metabolism. *Prog Nucleic Acid Res Mol Biol* **75**, 69-111
759   (2003).
760
761 45.   Ekstrand MI, *et al.* Mitochondrial transcription factor A regulates mtDNA copy number in
762   mammals. *Hum Mol Genet* **13**, 935-944 (2004).
763
764 46.   Kang D, Kim SH, Hamasaki N. Mitochondrial transcription factor A (TFAM): roles in
765   maintenance of mtDNA and cellular functions. *Mitochondrion* **7**, 39-44 (2007).
766
767 47.   Kang I, Chu CT, Kaufman BA. The mitochondrial transcription factor TFAM in
768   neurodegeneration: emerging evidence and mechanisms. *FEBS Lett* **592**, 793-811
769   (2018).
770
771 48.   Larsson NG, *et al.* Mitochondrial transcription factor A is necessary for mtDNA
772   maintenance and embryogenesis in mice. *Nature genetics* **18**, 231-236 (1998).
773
774 49.   Matsushima Y, Goto Y, Kaguni LS. Mitochondrial Lon protease regulates mitochondrial
775   DNA copy number and transcription by selective degradation of mitochondrial
776   transcription factor A (TFAM). *Proc Natl Acad Sci U S A* **107**, 18410-18415 (2010).
777
778 50.   Lu B, *et al.* Phosphorylation of human TFAM in mitochondria impairs DNA binding and
779   promotes degradation by the AAA+ Lon protease. *Mol Cell* **49**, 121-132 (2013).
780
781 51.   Gensler S, *et al.* Mechanism of mammalian mitochondrial DNA replication: import of
782   mitochondrial transcription factor A into isolated mitochondria stimulates 7S DNA
783   synthesis. *Nucleic Acids Res* **29**, 3657-3663 (2001).
784
785 52.   Pohjoismaki JL, Goffart S. Of circles, forks and humanity: Topological organisation and
786   replication of mammalian mitochondrial DNA. *Bioessays* **33**, 290-299 (2011).
787
788 53.   Gustafson MA, Sullivan ED, Copeland WC. Consequences of compromised
789   mitochondrial genome integrity. *DNA Repair (Amst)* **93**, 102916 (2020).
790
791 54.   Tan K, Fujimoto M, Takii R, Takaki E, Hayashida N, Nakai A. Mitochondrial SSBP1
792   protects cells from proteotoxic stresses by potentiating stress-induced HSF1
793   transcriptional activity. *Nat Commun* **6**, 6580 (2015).
794
795 55.   Kasashima K, Sumitani M, Endo H. Human mitochondrial transcription factor A is
796   required for the segregation of mitochondrial DNA in cultured cells. *Exp Cell Res* **317**,
797   210-220 (2011).
798

799   56.   West AP*, et al.* Mitochondrial DNA stress primes the antiviral innate immune response.
800   *Nature* **520**, 553-557 (2015).

802   57.   Rocha AG*, et al.* MFN2 agonists reverse mitochondrial defects in preclinical models of
803   Charcot-Marie-Tooth disease type 2A. *Science* **360**, 336-341 (2018).

805   58.   Guo S*, et al.* Inactivation of specific beta cell transcription factors in type 2 diabetes. *J*
806   *Clin Invest* **123**, 3305-3316 (2013).

808   59.   Ishida E, Kim-Muller JY, Accili D. Pair Feeding, but Not Insulin, Phloridzin, or
809   Rosiglitazone Treatment, Curtails Markers of beta-Cell Dedifferentiation in db/db Mice.
810   *Diabetes* **66**, 2092-2101 (2017).

812   60.   Li J*, et al.* Amorphous solid dispersion of Berberine mitigates apoptosis via
813   iPLA2beta/Cardiolipin/Opa1 pathway in db/db mice and in Palmitate-treated MIN6 beta-
814   cells. *Int J Biol Sci* **15**, 1533-1545 (2019).

816   61.   Liu J*, et al.* Rhein protects pancreatic beta-cells from dynamin-related protein-1-
817   mediated mitochondrial fission and cell apoptosis under hyperglycemia. *Diabetes* **62**,
818   3927-3935 (2013).

820   62.   Zhao Z*, et al.* Protection of pancreatic beta-cells by group VIA phospholipase A(2)-
821   mediated repair of mitochondrial membrane peroxidation. *Endocrinology* **151**, 3038-3048
822   (2010).

824   63.   Pommier Y, Sun Y, Huang SN, Nitiss JL. Roles of eukaryotic topoisomerases in
825   transcription, replication and genomic stability. *Nat Rev Mol Cell Biol* **17**, 703-721 (2016).

827   64.   Wicksteed B*, et al.* Conditional gene targeting in mouse pancreatic ss-Cells: analysis of
828   ectopic Cre transgene expression in the brain. *Diabetes* **59**, 3090-3098 (2010).

830   65.   Georgiadou E*, et al.* Mitofusins Mfn1 and Mfn2 are required to
831   preserve glucose-but not incretin- stimulated beta cell connectivity and insulin secretion.
832   *bioRxiv*, 2020.2004.2022.055384 (2021).

834   66.   Besse A, Brezavar D, Hanson J, Larson A, Bonnen PE. LONP1 de novo dominant
835   mutation causes mitochondrial encephalopathy with loss of LONP1 chaperone activity
836   and excessive LONP1 proteolytic activity. *Mitochondrion* **51**, 68-78 (2020).

838   67.   Peter B*, et al.* Defective mitochondrial protease LonP1 can cause classical mitochondrial
839   disease. *Hum Mol Genet* **27**, 1743-1753 (2018).

841   68.   Quiros PM*, et al.* ATP-dependent Lon protease controls tumor bioenergetics by
842   reprogramming mitochondrial activity. *Cell Rep* **8**, 542-556 (2014).

844   69.   Sebastian D*, et al.* Mitofusin 2 (Mfn2) links mitochondrial and endoplasmic reticulum
845   function with insulin signaling and is essential for normal glucose homeostasis. *Proc Natl*
846   *Acad Sci U S A* **109**, 5523-5528 (2012).

848   70.   de Brito OM, Scorrano L. Mitofusin 2 tethers endoplasmic reticulum to mitochondria.
849   *Nature* **456**, 605-610 (2008).

71. Dingreville F*, et al.* Differential Effect of Glucose on ER-Mitochondria Ca(2+) Exchange Participates in Insulin Secretion and Glucotoxicity-Mediated Dysfunction of beta-Cells. *Diabetes* **68**, 1778-1794 (2019).

72. Madec AM, Perrier J, Panthu B, Dingreville F. Role of mitochondria-associated endoplasmic reticulum membrane (MAMs) interactions and calcium exchange in the development of type 2 diabetes. *Int Rev Cell Mol Biol* **363**, 169-202 (2021).

73. Rieusset J. Role of Endoplasmic Reticulum-Mitochondria Communication in Type 2 Diabetes. *Adv Exp Med Biol* **997**, 171-186 (2017).

74. Cosson P, Marchetti A, Ravazzola M, Orci L. Mitofusin-2 independent juxtaposition of endoplasmic reticulum and mitochondria: an ultrastructural study. *PLoS One* **7**, e46293 (2012).

75. Filadi R, Greotti E, Turacchio G, Luini A, Pozzan T, Pizzo P. Mitofusin 2 ablation increases endoplasmic reticulum-mitochondria coupling. *Proc Natl Acad Sci U S A* **112**, E2174-2181 (2015).

76. Nile DL*, et al.* Age-related mitochondrial DNA depletion and the impact on pancreatic Beta cell function. *PLoS One* **9**, e115433 (2014).

77. Cree LM*, et al.* Age-related decline in mitochondrial DNA copy number in isolated human pancreatic islets. *Diabetologia* **51**, 1440-1443 (2008).

78. Chinnery PF*, et al.* Mitochondrial DNA haplogroups and type 2 diabetes: a study of 897 cases and 1010 controls. *J Med Genet* **44**, e80 (2007).

79. Miret-Casals L*, et al.* Identification of New Activators of Mitochondrial Fusion Reveals a Link between Mitochondrial Morphology and Pyrimidine Metabolism. *Cell Chem Biol* **25**, 268-278 e264 (2018).

80. Bach D*, et al.* Expression of Mfn2, the Charcot-Marie-Tooth neuropathy type 2A gene, in human skeletal muscle: effects of type 2 diabetes, obesity, weight loss, and the regulatory role of tumor necrosis factor alpha and interleukin-6. *Diabetes* **54**, 2685-2693 (2005).

81. Montaigne D*, et al.* Myocardial contractile dysfunction is associated with impaired mitochondrial function and dynamics in type 2 diabetic but not in obese patients. *Circulation* **130**, 554-564 (2014).

82. Zorzano A, Liesa M, Palacin M. Mitochondrial dynamics as a bridge between mitochondrial dysfunction and insulin resistance. *Arch Physiol Biochem* **115**, 1-12 (2009).

83. Thorens B, Tarussio D, Maestro MA, Rovira M, Heikkila E, Ferrer J. Ins1(Cre) knock-in mice for beta cell-specific gene recombination. *Diabetologia* **58**, 558-565 (2015).

84. Taddeo EP*, et al.* Individual islet respirometry reveals functional diversity within the islet population of mice and human donors. *Molecular metabolism* **16**, 150-159 (2018).

901
902 85. Pearson G*, et al.* Clec16a, Nrdp1, and USP8 Form a Ubiquitin-Dependent Tripartite
903   Complex That Regulates beta-Cell Mitophagy. *Diabetes* **67**, 265-277 (2018).
904
905 86. Soleimanpour SA*, et al.* The diabetes susceptibility gene Clec16a regulates mitophagy.
906   *Cell* **157**, 1577-1590 (2014).
907
908 87. Soleimanpour SA*, et al.* Calcineurin signaling regulates human islet {beta}-cell survival. *J
909   Biol Chem* **285**, 40050-40059 (2010).
910
911 88. Kolesar JE, Wang CY, Taguchi YV, Chou SH, Kaufman BA. Two-dimensional intact
912   mitochondrial DNA agarose electrophoresis reveals the structural complexity of the
913   mammalian mitochondrial genome. *Nucleic Acids Res* **41**, e58 (2013).
914
915 89. Karamanlidis G, Nascimben L, Couper GS, Shekar PS, del Monte F, Tian R. Defective
916   DNA replication impairs mitochondrial biogenesis in human failing hearts. *Circ Res* **106**,
917   1541-1548 (2010).
918
919 90. Reissaus CA*, et al.* A Versatile, Portable Intravital Microscopy Platform for Studying
920   Beta-cell Biology In Vivo. *Sci Rep* **9**, 8449 (2019).
921
922 91. Iwashita H*, et al.* Live Cell Imaging of Mitochondrial Autophagy with a Novel Fluorescent
923   Small Molecule. *ACS Chem Biol* **12**, 2546-2551 (2017).
924
925 92. Konig J*, et al.* Mitochondrial contribution to lipofuscin formation. *Redox Biol* **11**, 673-681
926   (2017).
927
928 93. Jayaraman S. A novel method for the detection of viable human pancreatic beta cells by
929   flow cytometry using fluorophores that selectively detect labile zinc, mitochondrial
930   membrane potential and protein thiols. *Cytometry A* **73**, 615-625 (2008).
931

Acknowledgements

S.A.S was supported by the JDRF (CDA-2016-189, SRA-2018-539, COE-2019-861), the NIH

(R01 DK108921, U01 DK127747), the Department of Veterans Affairs (I01 BX004444), the

Brehm family, and the Anthony family. E.L.-D. was supported by the NIH (T32 AI007413 and

T32-AG000114). G.L.P. was supported by the American Diabetes Association (19-PDF-063).

B.A.K. was supported by the Department of Veterans Affairs (I01 BX004444). The JDRF Career

Development Award to S.A.S. is partly supported by the Danish Diabetes Academy and the

Novo Nordisk Foundation. We acknowledge the Microscopy, Imaging and Cellular Physiology

Core of the University of Michigan DRC (P30 DK020572) for assistance with imaging studies.

We thank the University of Michigan Flow Cytometry Core for assistance with flow cytometry

studies. Next generation sequencing was carried out in the Advanced Genomics Core at the

University of Michigan. We acknowledge support from the Bioinformatics Core of the University

of Michigan's Biomedical Research Core Facilities. We thank Drs. K. Claiborn, C. Rutledge, N.

Desai, G. Rutter, A. Garner, A. Menon, M. Torres, and members of the Soleimanpour laboratory

for helpful advice.

Author Contributions

V.S. conceived, designed and performed experiments, interpreted results, drafted and reviewed

the manuscript. J.Z. and E.L-D. designed and performed experiments, interpreted results, and

reviewed the manuscript. G.L.P. and E.C.R. designed and performed experiments and

interpreted results. E.M.W. designed and performed experiments, interpreted results, edited,

and reviewed the manuscript. B.A.K. designed studies, interpreted results and reviewed the

manuscript. S.A.S. conceived and designed the studies, interpreted results, drafted, edited, and

reviewed the manuscript.

Conflict of interest statement

The authors have declared that no conflict of interest exists.
957

 FIGURE LEGENDS

Figure 1. Loss of both Mfn1 and Mfn2 in β-cells impairs glucose tolerance and glucose-

stimulated insulin release.

a Expression of Mfn1 and Mfn2 by Western blot (WB) in islets isolated from 8-10-week-old Ctrl,

β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice. Cyclophilin B serves as a loading control.

Representative of 6 independent mice/group for Mfn1 and 10 independent mice/group for Mfn2.

Band intensity of each lane of displayed representative blot (shown as fold change compared to

Ctrl, expression normalized to cyclophilin B) above each blot in red. **b** Densitometry (normalized

to cyclophilin B) from studies in Figure 1A. n=6/group for Mfn1 and 10/group for Mfn2. Data are

presented as Mean ± SEM. *p<0.0005 by one-way ANOVA followed by Tukey's post-hoc test

for multiple comparisons. **c** Blood glucose concentrations measured during IPGTT of 8-week-

old Ctrl (n=28), β-Mfn1^{KO} (n=7), β-Mfn2^{KO} (n=6) and β-Mfn1/2^{DKO} (n=15) littermates. Data are

presented as Mean ± SEM. (p<0.0005 by two-way ANOVA for β-Mfn1/2^{DKO} mice vs Ctrl

followed by Sidak's post-hoc test for multiple comparisons). **d** Serum insulin concentrations

(n=7-24/group) measured during *in vivo* glucose-stimulated insulin release testing in 8-week-old

Ctrl (n=24), β-Mfn1^{KO} (n=7), β-Mfn2^{KO} (n=10) and β-Mfn1/2^{DKO} (n=14) littermates. Data are

presented as mean ± SEM. *p<0.0005 for β-Mfn1/2^{DKO} mice vs Ctrl and β-Mfn1^{KO}, p=0.004 for

β-Mfn1/2^{KO} vs β-Mfn2^{KO} by one-way ANOVA followed by Sidak's post-hoc test for multiple

comparisons. **e** Glucose-stimulated insulin secretion following static incubations in 2 mM and

16.7 mM glucose, performed in isolated islets of 8-week-old Ctrl (n=16), β-Mfn1^{KO} (n=8), β-

Mfn2^{KO} (n=11) and β-Mfn1/2^{DKO} (n=10) littermates. Data are presented as mean from

independent mice ± SEM. *p<0.0005 by one-way ANOVA followed by Sidak's post-hoc test for

multiple comparisons. **f** Pancreatic β-cell mass measured in 10-week-old Ctrl (n=10), β-Mfn1^{KO}

(n=4), β-Mfn2^{KO} (n=4) and β-Mfn1/2^{DKO} (n=6) littermates. Data are presented as mean ± SEM.

Uncropped western blots and source data are provided as a Source Data file.

Figure 2. Combined Mfn1 and Mfn2 deficiency impairs mitochondrial respiration and structure in β-cells.

a OCR measured following exposure to 2mM glucose, 20mM glucose, and 2µM antimycin A in islets isolated from 8-10-week-old Ctrl (n=9) and β-Mfn1/2^{DKO} (n=8) mice by a Seahorse flux analyzer. Data are presented as mean from independent mice ± SEM. *p=0.0176 by one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons. **b** Representative transmission EM images of β-cells from 12-week-old Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} littermates. n=3/group. Mitochondria are highlighted with red arrows. **c** Quantification of mitochondrial morphology in Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} β-cells from transmission EM images (~125 independent mitochondria scored/animal). n=3/group. Data are presented as mean ± SEM. *p=0.0087 for Aspect Ratio, p=0.004 for Form factor and Circularity, p=0.037 for Perimeter by one-way ANOVA followed by Tukey's post-hoc test for multiple comparisons. **d** Imaris® generated three-dimensional reconstruction of deconvolution immunofluorescence Z-stack images at 100X magnification stained for SDHA (see inset image – red) from pancreatic sections of Ctrl and β-Mfn1/2^{DKO} mice. β-cells were identified by insulin co-staining (inset: insulin – green). Each unique color represents a separate β-cell mitochondrial network cluster. Representative image of 6 independent mice/group. **e** β-cell mitochondrial morphology and network analysis of deconvolution immunofluorescence Z-stack images at 100X magnification stained for SDHA (and insulin) from pancreatic sections of Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice by MitoAnalyzer. n=6/group (~150 β-cells/animal were quantified). Data are presented as mean ± SEM. *p=0.0006 for Average mitochondrial volume, p=0.0089 for Sphericity, p=0.0032 for Branches per mitochondria, p=0.0039 for Branch junctions by one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons. Source data are provided as a Source Data file.

Figure 3. Mfn1/2-deficient β-cells exhibit reduced mtDNA content.

a Relative mtDNA content measured by qPCR (normalized to nuclear DNA expression) in isolated islets of 8-week-old Ctrl (n=7), β-Mfn1^{KO} (n=5), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO} (n=5) littermates. Data are presented as mean ± SEM. *p=0.0037 for β-Mfn2^{KO} vs Ctrl, p<0.0005 for β-Mfn1/2^{DKO} vs Ctrl by one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons. **b** Heatmap representing relative expression (compared to littermate controls; Ctrl) of mitochondrial encoded transcripts in islets isolated from β-Mfn1^{KO} (n=4), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO} (n=5) mice. §* Benjamini-Hochberg FDR (Padj) <0.05 comparing β-Mfn2^{KO} (§) and β-Mfn1/2^{DKO} (*) to Ctrl. **c** Expression of OXPHOS subunit proteins by WB in islets isolated from 8-10-week-old Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice. Representative of 4 independent mice/group. **d** OXPHOS subunit densitometry (normalized to Vinculin) from studies in Figure 3D. n=4/group; Data are presented as mean ± SEM. *p=0.02 for Uqcrc2, p<0.05 for mt-co1, p<0.001 for Ndufb8, p<0.005 for mt-Cytb by one-way ANOVA followed by Tukey's post-hoc test for multiple comparisons. **e** Expression of Tom20 by WB in islets isolated from 8-10-week-old Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice. Representative image (top) of 6 independent mice/group. Tom20 densitometry, normalized to vinculin (bottom). n=6/group. Data are presented as mean ± SEM. Uncropped western blots and source data are provided as a Source Data file.

Figure 4. A single allele of *Mfn2*, but not *Mfn1*, maintains glycemic control by preserving β-cell mtDNA content despite impaired mitochondrial structure.

a Imaris® generated three-dimensional reconstruction of deconvolution immunofluorescence Z-stack images at 100X magnification stained for SDHA (see inset image – red) from pancreatic sections of Ctrl, β-Mfn1/2^{DKO}, β-Mfn1^{KO}Mfn2$^{+/-}$ and β-Mfn1$^{+/-}$Mfn2^{KO} mice. β-cells were identified by insulin co-staining (inset: insulin – green). Each unique color represents a separate β-cell mitochondrial network cluster. Representative image of 6 independent mice/group. **b** β-cell mitochondrial morphology and network analysis of deconvolution immunofluorescence Z-stack

images from pancreatic sections stained for insulin and SDHA of Ctrl, β-Mfn1/2^{DKO}, β-

Mfn1^{KO}Mfn2$^{+/-}$ and β-Mfn1$^{+/-}$Mfn2^{KO} mice by MitoAnalyzer. n=6/group (50-200 β-cells/animal

were quantified). *p=0.005 for Average mitochondrial volume, p=0.05 for Sphericity, p=0.05 for

Branches per mitochondria, p=0.05 for Branch junctions by one-way ANOVA followed by

Sidak's post-hoc test for multiple comparisons. **c** Blood glucose concentrations measured

during IPGTT of 8-week-old Ctrl (n=28), β-Mfn1/2^{DKO} (n=15), β-Mfn1^{KO}Mfn2$^{+/-}$ (n=13) and β-

Mfn1$^{+/-}$Mfn2^{KO} (n=17) littermates. (p<0.0005 by two-way ANOVA for β-Mfn1$^{+/-}$Mfn2^{KO} and β-

Mfn1/2^{DKO} mice vs Ctrl followed by Sidak's post-hoc test for multiple

comparisons). **d** Serum insulin concentrations (n=11-24/group) measured during *in*

vivo glucose-stimulated insulin release testing in 8-week-old Ctrl (n=24), β-Mfn1/2^{DKO} (n=14), β-

Mfn1^{KO}Mfn2$^{+/-}$ (n=11) and β-Mfn1$^{+/-}$Mfn2^{KO} (n=16) littermates. *p<0.0005 by one-way ANOVA

followed by Sidak's post-hoc test for multiple comparisons. **e** Relative mtDNA content measured

by qPCR (normalized to nuclear DNA expression) in isolated islets of 8-week-old Ctrl (n=7), β-

Mfn1/2^{DKO} (n=5), β-Mfn1^{KO}Mfn2$^{+/-}$ (n=5) and β-Mfn1$^{+/-}$Mfn2^{KO} (n=5) littermates. Data are

presented as mean ± SEM. *p<0.0005 by one-way ANOVA followed by Sidak's post-hoc test for

multiple comparisons. Of note, studies in Ctrl and β-Mfn1/2^{DKO} mice were performed together

alongside all β-Mfn1^{KO}, β-Mfn2^{KO}, β-Mfn1^{KO}Mfn2$^{+/-}$ (n=11) and β-Mfn1$^{+/-}$Mfn2^{KO} (n=16) littermates

and thus may appear twice for purposes of relevant comparisons (in Figures 1C, 1D, 2D, 2E,

3A and again in Figures 4A-E). Source data are provided as a Source Data file.

Figure 5. Mfn1/2-deficiency induces expression of genes associated with mtDNA

replication.

a Hierarchical clustering of RNAseq samples displaying the 500 genes with the highest mean

expression in islets of 12-week-old Ctrl (n=6), β-Mfn1^{KO} (n=4), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO}

(n=5) mice. **b** Pathway analysis of top differentially regulated pathways in β-Mfn1/2^{DKO} islets

(n=5) compared to littermate Ctrl islets (n=6), demarcated by both false discovery rate (FDR)

and the number of significantly differentially expressed genes (DEGs) per pathway. **c** Volcano

plot depicting differential RNA expression in islets of β-Mfn1/2^{DKO} (n=5) mice compared to

littermate Ctrl mice (n=6). Significantly differentially expressed genes demarcated by -log10

FDR > or < 2 and log2 fold change (FC) > or < 0.6. **d** Volcano plot depicting differential RNA

expression of MitoCarta 2.0 targets in islets of β-Mfn1/2^{DKO} (n=5) mice compared to littermate

Ctrl mice (n=6). Significantly differentially expressed genes demarcated by -log10 FDR > or <2

and log2 fold change (FC) > or <0.6. **e** Differential RNA expression heatmap of selected genes

from islets of β-Mfn1^{KO} (n=4), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO} (n=5) mice compared to littermate

Ctrl mice (n=6). up. ‡§* Benjamini-Hochberg FDR (Padj) <0.05 comparing β-Mfn1^{KO}(‡), β-

Mfn2^{KO}(§) and β-Mfn1/2^{DKO}(*) to Ctrl.

Figure 6. Mfn1 and 2 maintain expression of Tfam in β-cells.

a Expression of Tfam, key replisome proteins, and LonP1 by Western blot (WB) in islets isolated

from 11-week-old Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice. Representative

of 4 independent mice per group. **b** Densitometry (normalized to Vinculin) from studies in Figure

6A. n=4/group; Data are presented as mean ± SEM. *p=0.03 for Tfam, p=0.01 for Lonp1 by

one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons. **c** Quantification of

single strand mtDNA products measured by qPCR in isolated islets from 8-week-old Ctrl (n=13),

β-Mfn1^{KO} (n=5), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO} (n=9) mice. Data are presented as mean ±

SEM. *p=0.0007 by one-way ANOVA followed by Sidak's post-hoc test for multiple

comparisons. **d** Deconvolution immunofluorescence image at 100X magnification of islets from

Ctrl and β-Mfn1/2^{DKO} mice stained for Ssbp1 (red), mtDNA (green), and Pdx1 (blue).

Representative of 8 independent experiments. White arrows demarcate co-localized Ssbp1+

mtDNA+ structures. **e** Quantification of mitochondrial nucleoid Ssbp1 localization

(Ssbp1+ mtDNA+ co-localization) in Ctrl and β-Mfn1/2^{DKO} β-cells from studies depicted in Figure

6D by Pearson's correlation coefficient (PCC). n=8/group. Data are presented as mean ± SEM.

*p=0.0142 by two-tailed t-test. (~100 β-cells from each animal per group were

analyzed). **f** Quantification of nuclear Ssbp1 localization in Ctrl and β-Mfn1/2^{DKO} β-cells

from studies depicted in Figure 6D by Pearson's correlation coefficient (PCC). n=8/group. Data

are presented as mean ± SEM. *p=0.0008 by two-tailed t-test. (~100 β-cells from each animal

per group were analyzed). **g** Imaris® generated 3D reconstruction of deconvolution

immunofluorescence Z-stack images at 100X magnification stained for DNA of β-cells (see inset

image – Pdx1, blue, DNA, green) from Ctrl and β-Mfn1/2^{DKO} mice. Colors represent relative

mitochondrial volume. Representative image of 8 independent mice/group. **h** Quantification of

relative nucleoid volume of Ctrl and β-Mfn1/2^{DKO} β-cells from studies depicted in Figure 6g.

n=8/group. Data are presented as mean ± SEM. *p=0.0127 by two-tailed t-test. (~100 β-cells

from each animal per group were analyzed). Uncropped western blots and source data are

provided as a Source Data file.

Figure 7. Mfn1 and 2 maintain β-cell function through Tfam-mediated mtDNA copy

number control.

a Expression of mouse-specific Tfam and human-specific TFAM by WB in islets isolated from

10-15-week-old Ctrl and β-Mfn1/2^{DKO} mice, transduced with empty vector control (Ad.EV) or

human TFAM-overexpressing (Ad.hTFAM) adenoviral particles. Representative

of 4 independent mice/group. **b** Densitometry (normalized to cyclophilin B) from studies in

Figure 7A. n=4/group; Data are presented as mean ± SEM. p=NS (non-significant) by two-tailed

t-test. **c** Glucose-stimulated insulin secretion following static incubations in 2mM and 16.7

mM glucose, performed in isolated Ctrl and β-Mfn1/2^{DKO} islets following transduction

with Ad.EV or Ad.hTFAM adenoviral particles. n=6/group. Data are presented as mean ± SEM.

*p<0.0005 for Ctrl + Ad.EV vs β-Mfn1/2^{DKO} + Ad.EV, p=0.0148 for β-Mfn1/2^{DKO} + Ad.EV vs β-

Mfn1/2^{DKO} + Ad.hTFAM by one-way ANOVA followed by Tukey's post-hoc test for multiple

comparisons. **d** Relative mtDNA content measured by qPCR (normalized to nuclear DNA

expression) from isolated Ctrl and β-Mfn1/2^{DKO} islets following transduction

with Ad.EV or Ad.hTFAM adenoviral particles. n=12/group. Data are presented as mean ± SEM.

*p<0.0005 for Ctrl + Ad.EV vs β-Mfn1/2^{DKO} + Ad.EV, p=0.007 for β-Mfn1/2^{DKO} + Ad.EV vs β-

Mfn1/2^{DKO} + Ad.hTFAM by two-tailed t-test. Uncropped western blots and source data are

provided as a Source Data file.

Figure 8. Mitofusin agonists restore mtDNA depletion following Mfn2 deficiency

independent of changes in mitochondrial structure.

a Relative mtDNA content measured by qPCR (normalized to nuclear DNA expression) from

isolated islets of Ctrl and β-Mfn2^{KO} mice following treatment with vehicle (Veh) or 0.5 μM Mfn

agonists for 24 hrs. n=4/group. Data are presented as mean ± SEM. *p=0.0138 for Ctrl + Veh vs

β-Mfn2KO + Veh, p=0.0302 for β-Mfn2^{KO} + Veh vs β-Mfn2^{KO} + Mfn agonists by one-way ANOVA

followed by Sidak's post-hoc test for multiple comparisons. **b** Expression of Mfn1 and Mfn2 by

WB in islets isolated from 10-12-week-old *db*/+ and *db*/*db* mice. Vinculin and cyclophilin B serve

as loading controls. Representative image (left) of 3 independent mice/group for Mfn1 and 4

independent mice/group for Mfn2. Mfn1 (n=3) and Mfn2 (n=4) densitometry, normalized to

vinculin or cyclophilin B (right). Data are presented as mean ± SEM. *p=0.0095 by two-tailed t-

test. **c** Relative mtDNA content measured by qPCR (normalized to nuclear DNA expression)

from isolated islets of 10-12-week-old *db*/+ (n=6) and *db*/*db* (n=5) mice following treatment with

vehicle (Veh) or 0.5 μM mitofusin agonists for 24 hrs. Data are presented as mean ± SEM.

*p=0.0038 for *db*/+ + Veh vs *db*/*db* + Veh, p=0.0428 for *db*/*db* + Veh vs *db*/*db* + Mfn agonists

by two-tailed t-test. **d** Immunofluorescence images at 20X magnification of islets from 10-12-

week-old *db*/+ and *db*/*db* mice stained for Insulin (green), Glucagon (red), Somatostatin (SST;

white), and DAPI (blue) following treatment with vehicle (Veh) or 0.5 μM mitofusin agonists for

24 hrs. n=3/group. **e** β-cell mitochondrial morphology and network analysis by MitoAnalyzer of

deconvolution immunofluorescence Z-stack images, stained for SDHA (and insulin), from

pancreatic islets of 10-12-week-old *db*/+ and *db*/*db* mice following treatment with vehicle (Veh)

or 0.5 μM mitofusin agonists for 24 hrs. n=6/group (~150-200 β-cells/animal were quantified).

Data are presented as mean ± SEM. **f** Glucose-stimulated insulin secretion following static

incubations in 2 mM and 16.7 mM glucose, performed in isolated islets of 10-12-week-old *db*/+

and *db*/*db* mice following treatment with vehicle (Veh) or 0.5 µM mitofusin agonists for 24 hrs.

n=11/group. Data are presented as mean ± SEM. *p=0.0092 by two-tailed t-test. Uncropped

western blots and source data are provided as a Source Data file.

1143

Mitofusin 1 and 2 regulation of mitochondrial DNA content is a critical determinant of

glucose homeostasis

Vaibhav Sidarala[1*]; Jie Zhu[1*]; Elena Levi-D'Ancona[1]; Gemma L. Pearson[1]; Emma C. Reck[1];

Emily M. Walker[1]; Brett A. Kaufman[2]; and Scott A. Soleimanpour[1,3,4]

[1]Division of Metabolism, Endocrinology & Diabetes and Department of Internal Medicine,

University of Michigan, Ann Arbor, MI 48105, USA.

[2]Vascular Medicine Institute, Division of Cardiology, Department of Medicine, University of

Pittsburgh School of Medicine, Pittsburgh, PA, 15260, USA

[3]Department of Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI

48105, USA

[4]VA Ann Arbor Healthcare System, Ann Arbor, MI 48105, USA.

*These authors contributed equally to this work

Corresponding Author

Scott A. Soleimanpour, MD, 1000 Wall Street, Brehm Tower Room 6329, Ann Arbor, MI 48105,

USA. Phone: (734) 763-0528 E-mail: ssol@med.umich.edu

ABSTRACT

The dynamin-like GTPases Mitofusin 1 and 2 (Mfn1 and Mfn2) are essential for mitochondrial

function, which has been principally attributed to their regulation of fission/fusion dynamics. Here,

we report that Mfn1 and 2 are critical for glucose-stimulated insulin secretion (GSIS) primarily

through control of mitochondrial DNA (mtDNA) content. Whereas Mfn1 and Mfn2 individually were

dispensable for glucose homeostasis, combined Mfn1/2 deletion in β-cells reduced mtDNA

content, impaired mitochondrial morphology and networking, and decreased respiratory function,

ultimately resulting in severe glucose intolerance. Importantly, gene dosage studies unexpectedly

revealed that Mfn1/2 control of glucose homeostasis was dependent on maintenance of mtDNA

content, rather than mitochondrial structure. Mfn1/2 maintain mtDNA content by regulating the

expression of the crucial mitochondrial transcription factor Tfam, as Tfam overexpression

ameliorated the reduction in mtDNA content and GSIS in Mfn1/2-deficient β-cells. Thus, the

primary physiologic role of Mfn1 and 2 in β-cells is coupled to the preservation of mtDNA content

rather than mitochondrial architecture, and Mfn1 and 2 may be promising targets to overcome

mitochondrial dysfunction and restore glucose control in diabetes.

INTRODUCTION

Mitochondrial dynamics, the balance of fusion and fission of mitochondrial networks, is essential to control mitochondrial structure. Mitochondrial fission and fusion are governed by several key proteins that maintain mitochondrial quality control [1, 2]. Fission is primarily regulated by the GTPase dynamin-related protein 1 (Drp1), while fusion is controlled by both outer and inner mitochondrial membrane machinery. The dynamin-related large GTPase optic atrophy protein 1 (Opa1) controls inner membrane fusion, while outer membrane fusion is controlled by two GTPases known as Mitofusin 1 and 2 (Mfn1 and Mfn2). Mitofusin 1 and 2 share ~80% sequence similarity and contain homologous functional domains, suggesting they play both overlapping and unique roles in metabolic function largely ascribed to their control of mitochondrial structure [3].

Imbalances in mitochondrial dynamics in metabolically active tissues have been implicated in many human diseases, including neurodegenerative conditions, cancer, cardiovascular disease, and diabetes [2]. Mitochondria provide the energy necessary for β-cell insulin release [4, 5], and abnormalities in mitochondrial structure and bioenergetics have been observed in the β-cells of humans with type 2 diabetes (T2D; [6, 7]). Further, mitochondrial defects have been recently found to precede the development of T2D in human β-cells [8]. Thus, strategies to understand and overcome mitochondrial dysfunction in β-cells are of great appeal for the treatment of diabetes. However, the role of the core fission/fusion machinery in β-cells is less clear. Reduced β-cell *Mfn2* expression has been observed secondary to the deposition of toxic islet amyloid polypeptide oligomers observed in T2D as well as in mouse models of T2D [9, 10], yet the functions of Mfn1 and Mfn2 in β-cells *in vivo* are unexplored.

Here, we elucidate a key physiologic role for Mfn1 and 2 in the maintenance of glucose homeostasis that unexpectedly occurs through the preservation of β-cell mtDNA content, rather than mitochondrial structure. Utilizing several genetic mouse models, high resolution imaging, and transcriptomic profiling, we show that Mfn1 and 2 act in tandem to preserve β-cell mitochondrial health and, consequently, glucose-stimulated insulin secretion (GSIS). Mfn1 and 2

65   maintain mtDNA content by preventing loss of the mitochondrial transcription factor Tfam, which

66   is a master regulator of mtDNA copy number control and mtRNA expression [11, 12]. Indeed,

67   adenoviral overexpression of Tfam ameliorates reductions in mtDNA content and GSIS in Mfn1/2-

68   deficient β-cells, illustrating the critical importance of mtDNA copy number control by Mfn1/2 to

69   promote glucose homeostasis.

RESULTS

Combined deficiency of Mfn1 and Mfn2 in β-cells leads to severe glucose intolerance and reduced glucose-stimulated insulin secretion

Mitochondrial health is vital to support β-cell insulin secretory responses to glucose or other nutrient stimuli. We hypothesized that Mfn1 and Mfn2 are required for GSIS in β-cells through their regulation of mitochondrial function. To test this hypothesis, we generated mice bearing β-cell specific deletion of Mfn1 alone (*Mfn1*^{loxP/loxP};*Ins1*-Cre), Mfn2 alone (*Mfn2*^{loxP/loxP};*Ins1*-Cre), and combined deletion of both Mfn1 and 2 (*Mfn1*^{loxP/loxP}*Mfn2*^{loxP/loxP};*Ins1*-Cre), hereafter known as β-Mfn1^{KO}, β-Mfn2^{KO}, and β-Mfn1/2^{DKO} mice, respectively. Conditional alleles targeted the canonical G-1 GTPase motif of Mfn1 (exon 4) or Mfn2 (exon 6) (Figure S1a; [13]). *Ins1*-Cre and floxed alone controls were phenotypically indistinguishable from one another, exhibiting no differences in body weight or glycemic control (Figure S1b-c). Therefore, they were pooled as controls for subsequent analyses. β-Mfn1^{KO}, β-Mfn2^{KO}, and β-Mfn1/2^{DKO} mice each exhibited an efficient reduction in Mfn1 and/or Mfn2 islet protein expression when compared to littermate controls (Ctrl; Figures 1a-b). We did not observe significant compensatory changes in Mfn2 expression in β-Mfn1^{KO} islets, nor in Mfn1 expression in β-Mfn2^{KO} islets (Figures 1a-b). Further, we did not observe differences in expression of the other core fission/fusion proteins, Drp1 and Opa1, following deletion of Mfn1 and/or Mfn2 (Figure S1d).

While loss of either Mfn1 or Mfn2 alone has been shown to impair glucose homeostasis in other metabolic tissues [3, 14], we were surprised to find that both β-Mfn1^{KO} and β-Mfn2^{KO} mice exhibited normal glucose homeostasis during an intraperitoneal glucose tolerance test (IPGTT; Figure 1c). We also observed no changes in insulin secretion in single β-Mfn1^{KO} or β-Mfn2^{KO} mice after glucose administration *in vivo* (Figure 1d) or in isolated islets (Figure 1e). However, combined deletion of both Mfn1 and Mfn2 resulted in severe glucose intolerance (Figure 1c; p <0.005 by one-way ANOVA vs. Ctrl), due to markedly reduced GSIS (Figures 1d-e and S1e).

To evaluate other etiologies of glucose intolerance in β-Mfn1/2[DKO] mice, we first measured

β-cell mass. We observed no significant differences in β-cell mass between genotypes (Figure

1f). There were also no differences in body weight or peripheral insulin sensitivity between the

groups (Figures S1f-g). Taken together, these studies suggest that Mfn1 and 2 are individually

dispensable for glycemic control and GSIS, but together play vital complementary roles for the

maintenance of glucose tolerance and β-cell insulin release.

Mfn1 and 2 regulate β-cell mitochondrial structure and respiratory function

Given the connections between mitochondrial function and dynamics, we asked whether

mitochondrial respiration and architecture would be defective in the islets of β-Mfn1/2[DKO] mice, as

has been reported in other models of Mfn1/2-deficiency [15, 16, 17, 18, 19]. Indeed, islets isolated from

β-Mfn1/2[DKO] mice displayed a reduced glucose-stimulated oxygen consumption rate (OCR;

Figures 2a and S2a) when compared to Ctrl islets. We did not observe differences in glycolysis

as measured by extracellular acidification rate (ECAR; Figure S2a). Differences in glucose-

stimulated OCR between Ctrl and β-Mfn1/2[DKO] islets were no longer observed following exposure

to the Complex III inhibitor antimycin A, suggesting that Mfn1/2-deficient islets primarily possess

a defect in β-cell mitochondrial respiration (Figures 2a and S2a).

Loss of key mediators of mitochondrial fusion disrupt mitochondrial network balance and

favor increased mitochondrial fission, often leading to the appearance of increased punctate or

spherical mitochondria [20]. We thus examined mitochondrial structure in Mfn1- and/or Mfn2-

deficient β-cells by several independent and complementary methods. First, we assessed β-cell

mitochondrial ultrastructure by transmission electron microscopy (TEM). While β-cells from all

groups had the expected appearance of electron dense insulin granules, β-cells from β-Mfn1/2[DKO]

mice had reductions in mitochondrial aspect ratio, form factor, and perimeter, and increased

mitochondrial circularity (Figures 2b-c), indicating smaller, punctate mitochondria. We observed

no overt differences in ER morphology and proximity to mitochondria in β-cells from control and

β-Mfn1/2^{DKO} mice (Fig. S2b), however.

Secondly, we examined mitochondrial networking utilizing high-resolution deconvolution

imaging of the mitochondrial marker succinate dehydrogenase A (SDHA). We initially used

Imaris® imaging software to generate three-dimensional (3D) reconstructions of β-cell

mitochondria to grossly evaluate mitochondrial networking [21, 22]. Consistent with our findings using

TEM, the number of 3D mitochondrial networks appeared similar between Ctrl and single β-

Mfn1^{KO} or β-Mfn2^{KO} β-cells (Figure S2c). However, Mfn1/2-double deficient mice showed a

dramatic increase in the frequency of smaller β-cell mitochondrial networks (Figure 2d).

Next, we analyzed 3D deconvolution imaging of β-cell SDHA using the Mitochondria

Analyzer, a validated and unbiased quantification tool used to evaluate mitochondrial morphology

and network/branch complexity [22, 23]. This confirmed that measures of mitochondrial morphology

(reduced mitochondrial volume and increased sphericity) and network/branch complexity

(reduced branch number and branch junctions) were significantly impaired in β-Mfn1/2^{DKO} mice,

commensurate with an increase in punctate/spherical mitochondria due to decreased

mitochondrial fusion (Figure 2e). Together, these data demonstrate that together Mfn1 and 2

maintain β-cell mitochondrial structure and function.

Mfn1 and 2 maintain β-cell mtDNA content

Mitochondrial DNA copy number is vital for β-cell function and glucose homeostasis [24, 25]. Thus,

we next examined whether mtDNA content was altered by loss of Mfn1 and/or Mfn2 in β-cells.

We observed a small, yet significant, reduction in mtDNA content in islets of β-Mfn2^{KO} mice that

was not observed in islets of β-Mfn1^{KO} mice (Figure 3a). Interestingly, mtDNA depletion

substantially worsened in the islets of β-Mfn1/2^{DKO} mice (Figure 3a), suggesting that loss of Mfn1

exacerbates the depletion of mtDNA caused by Mfn2 deficiency. We next assessed expression

of the 13 mitochondrially encoded genes and observed a consistent and significant reduction in

mtRNA expression in β-Mfn2^{KO} islets that was exacerbated in β-Mfn1/2^{DKO} mice (Figure 3b).

Concordantly, examination of protein expression of mitochondrial OXPHOS subunits by Western

blot revealed reductions of the mitochondrially encoded subunits of Complex IV, mt-Co1, and

Complex III, mt-Cytb, in β-Mfn2^{KO} islets and β-Mfn1/2^{DKO} islets (Figures 3c-d). We also observed

significant reductions of nuclear encoded subunits of Complex I (Ndufb8) in both β-Mfn2^{KO} and β-

Mfn1/2^{DKO} islets and Complex III (Uqcrc2) in β-Mfn1/2^{DKO} islets, but no significant depletion of

nuclear encoded Complex II (Sdhb) and Complex V (Atp5a) (Figures 3c-d). Depletion of mtDNA

content in β-Mfn1/2^{DKO} islets did not appear to be secondary to changes in mitochondrial mass,

as we did not observe reductions in Atp5a (Figures 3c-d) or in the mitochondrial outer membrane

protein Tom20 (Figure 3e), a common marker of mitochondrial mass [26]. These studies reveal that

both Mfn1 and 2 (and Mfn2 alone to a lesser degree) are necessary to maintain mtDNA copy

number in β-cells.

Defects in β-cell mitochondrial fusion induce glucose intolerance upon loss of mtDNA

content

Our observations showing distinct and overlapping contributions of Mfn1 and 2 in the regulation

of β-cell mitochondrial function led us to hypothesize that loss of mitochondrial fusion would lead

to glucose intolerance due to mtDNA depletion. However, approaches to clarify the specific

contribution of mtDNA depletion to metabolic dysfunction following imbalances in mitochondrial

dynamics have been elusive. To this end, we generated animals harboring only a single functional

allele of either *Mfn1* (β-Mfn1$^{+/-}$Mfn2^{KO}) or *Mfn2* (β-Mfn1^{KO}Mfn2$^{+/-}$) to ascertain the relative

importance of mtDNA content and structure in the maintenance of glycemic control. For

robustness of interpretation, these experiments were performed alongside littermate controls and

β-Mfn1/2^{DKO} mice as previously presented (Figures 1-3). As expected, neither β-Mfn1$^{+/-}$Mfn2^{KO} nor

β-Mfn1^{KO}Mfn2$^{+/-}$ mice developed changes in β-cell mass, body weight, or insulin sensitivity

(Figures S3a-c). Interestingly, 3D quantification of mitochondrial morphology and network integrity

revealed that, similar to β-Mfn1/2^{DKO} mice, a single allele of either *Mfn1* or *Mfn2* was not sufficient

to maintain β-cell mitochondrial structure, leading to increases in punctate mitochondria

consistent with loss of mitochondrial fusion (Figures 4a-b).

Despite mitochondrial morphologic imbalances in both β-Mfn1$^{+/-}$Mfn2^{KO} and β-

Mfn1^{KO}Mfn2$^{+/-}$ mice, *only* β-Mfn1$^{+/-}$Mfn2^{KO} mice developed glucose intolerance and impaired GSIS

similar to that of β-Mfn1/2^{DKO} mice (Figures 4c-d). We also found significant reductions in mtDNA

content associated with the loss of GSIS observed in β-Mfn1$^{+/-}$Mfn2^{KO} and β-Mfn1/2^{DKO} mice

(Figure 4e). Thus, maintenance of a single functional allele of *Mfn2*, but not *Mfn1*, is able to

preserve glucose homeostasis, mtDNA content, and insulin secretion, despite being insufficient

to maintain β-cell mitochondrial network integrity. Importantly, these results disjoin β-cell function

from changes in mitochondrial morphology, and implicate mtDNA copy number as the vital

purveyor of glycemic control following loss of mitochondrial fusion.

Mfn1 and 2 deficiency induces a transcriptional signature consistent with defects in

mitochondrial metabolism and alterations in mtDNA replication

To begin to understand the mechanisms underlying control of mtDNA content by Mfn1 and 2, we

analyzed bulk RNA sequencing (RNAseq) data generated from islets of β-Mfn1^{KO}, β-Mfn2^{KO}, and

β-Mfn1/2^{DKO} mice as well as littermate controls. Initial hierarchical clustering analyses based on

the top 500 most highly expressed genes revealed that only β-Mfn1/2^{DKO} islets clustered by

genotype, while we did not observe discrete clustering across Ctrl or single β-Mfn1^{KO} and β-

Mfn2^{KO} islets due to very few differences in expression between these groups (Figures 5a and

S4a-b). Gene ontology and pathway analyses of differentially expressed genes from β-Mfn1/2^{DKO}

islets revealed significant changes related to endocrine hormone/insulin secretion, β-cell

signature genes, as well as several metabolic pathways (Figures 5b and S4c-d), consistent with

the defects in β-cell function and mitochondrial metabolism observed in β-Mfn1/2^{DKO} mice.

The nearly 1,000 significantly dysregulated genes in β-Mfn1/2[DKO] islets included

downregulation of mitochondrial encoded transcripts, as well as key β-cell signature genes such

as *Slc2a2* (Glut2) and *MafA* (Figures 5c and 5e). We also observed increases in expression of

regulators of triglyceride metabolism, including *Angptl4* and *Angptl8* [27], the

glycolytic/gluconeogenic enzyme *AldoB*, which is highly upregulated in models of β-cell

mitochondrial dysfunction and in T2D islets [28, 29], as well as *Nrf1*, a key regulator of mitochondrial

biogenesis (Figures 5c, 5e, and S4d; [30]). To resolve additional mitochondrial specific effectors

that were differentially regulated following the loss of Mfn1 and 2, we next overlaid our RNAseq

data on MitoCarta 2.0 [31], which contains a compendium of targets with strong evidence for

localization to the mitochondria (Figure 5d). This revealed increased expression of several genes

related to mtDNA replication machinery or associated with the mitochondrial nucleoid (Figures

5d-e), including the critical mtDNA topoisomerase *Top1mt* [32], the mitochondrial co-chaperone

Dnaja3/*Tid1*, which maintains mtDNA integrity [33], and the mitochondrial pyrimidine transporter

Slc25a33, which is essential for mtDNA replication [34]. We also observed upregulation in the

mitochondrial AAA+ protease *LonP1*, which interacts with and regulates the stability of numerous

proteins at the mitochondrial nucleoid (Figures 5d-e; [35, 36, 37]). Importantly, we did not observe

changes in expression of other fission/fusion genes or regulators of mtRNA transcription, mtDNA

repair, sirtuins, or mitophagy (Figure S4d). Taken together, these data indicate that Mfn1/2-

deficiency induced a transcriptional signature consistent with an activation of mtDNA replication

to potentially compensate for β-cell mtDNA depletion.

Mfn1 and 2 act through Tfam to regulate β-cell mtDNA copy number

The mechanisms underlying regulation of mtDNA by mitochondrial fusion have been

controversial, and previous studies implicate several pathways that could reduce mtDNA content,

including genome instability/increased mtDNA mutations, enhanced mitophagy, defects in

mitochondria-associated ER membrane (MAM) function, or impaired mtDNA replication [17, 19, 38, 39,]

[40, 41]. As our studies were completed in young mice, which bear infrequent mtDNA mutations [42],

and we did not observe differences in expression of mtDNA repair enzymes (Figure S4d), we

focused on assessments of mitophagy and mtDNA replication to better resolve the significant

reduction of mtDNA observed in β-Mfn1/2^{DKO} islets. We first assessed rates of mitophagy following

incubation of control and β-Mfn1/2^{DKO} islets with the cell permeable Mtphagy dye, which did not

reveal differences in mitophagy between genotypes (Figure S4e). We also did not observe a

difference in frequency of autophagosomes bearing mitochondria by TEM, or changes in essential

genes in the mitophagy pathway between control and β-Mfn1/2^{DKO} islets (Figure 2b, S2b and

S4d). As an initial screen for MAM function, we measured MAM-dependent lipids, including

cholesteryl esters and phospholipids [43, 44]. However, we did not detect differences in total

cholesterol, cholesteryl esters, or phosphatidylserine concentrations in isolated islets of control

and β-Mfn1/2^{DKO} mice (Figures S4f-g).

To assess regulators of β-cell mtDNA replication, we examined expression of the

replisome proteins Ssbp1, Twinkle, and Polg. We also measured expression of Tfam, a master

regulator of mtDNA copy number that promotes mitochondrial genome abundance via effects on

mtDNA packaging, stability, and replication, and whose levels are proportional to mtDNA copy

number [11, 45, 46, 47, 48]. While replisome proteins were largely unchanged, we observed a significant

reduction in Tfam protein in β-Mfn1/2^{DKO} islets (Figures 6a-b). *Tfam* mRNA expression was

unchanged in β-Mfn1/2^{DKO} mice (Figure S4d), suggesting the reduction in Tfam protein occurred

post-transcriptionally. We further observed a significant increase in the level of LonP1 protein,

which has been shown to lead to Tfam protein turnover [49, 50]. We additionally confirmed reductions

in Tfam and increases in LonP1 protein in β-Mfn1$^{+/-}$Mfn2^{KO} islets (Figure S5), correlating with the

mtDNA depletion, glucose intolerance, and impaired GSIS observed in these mice (Figures 4c-

e). In contrast, β-Mfn1^{KO}Mfn2$^{+/-}$ mice, which develop mitochondrial structural alterations without

glucose intolerance or mtDNA depletion, did not exhibit changes in Tfam and LonP1 (Figure S5

and Figures 4c-e). These results suggest that changes in Tfam and LonP1 expression following

Mfn1/2 deficiency were not due to increases in punctate mitochondria.

To test for defects in mtDNA replication, we next assessed initiation of first strand

replication. First strand synthesis in mitochondrial replication begins in the D-loop. However, a

single stranded (ss) molecule that forms the classic D-loop structure called 7S, which is a non-

replicative structure, shares this sequence. It is the elongation of the 7S sequence into the nearby

region (Cytb) that demonstrates committed initiation of replication. Here we assessed the levels

of ss and dsDNA at specific sequences by quantitative PCR to overcome expected sample

limitations from studies with mouse pancreatic islets. Notably, import of nascent Tfam has been

shown to increase the generation of 7S DNA due to its transcription initiation activity [51]. We found

a trend towards reduced total ss 7S sequences normalized to ds mtDNA in β-Mfn1/2^{DKO} islets

(Figure 6c). However, the rate of 7S sequence extension into committed first strand replication

was increased in β-Mfn1/2^{DKO} islets (Figure 6c), which could represent a compensatory response

to mtDNA depletion.

Increased first strand replication intermediates heighten the levels of displaced second

strand template, which is bound by the single strand binding protein Ssbp1 [52]. Consistent with ss

qPCR studies, in situ levels of Ssbp1 at the mtDNA nucleoid (detected by perinuclear anti-DNA

antibody staining), were significantly elevated in Mfn1/2-deficient β-cells (Figures 6d-e). The

increase in Ssbp1 localization to mitochondrial nucleoids could represent binding to a higher

quantity of single-stranded mtDNA replication intermediates and could also be consistent with

observations of impaired completion of mtDNA replication previously reported following Mfn1/2-

deficiency [19, 53]. We also observed Ssbp1 nuclear localization (Figures 6d and f), which has been

previously demonstrated following mitochondrial stress [54]. Further, we detected an increase in the

volume of mtDNA nucleoid structures in Mfn1/2-deficient β-cells by high-resolution 3D imaging

(Figures 6g-h), which are similar to previous reports of Tfam-deficiency but could also represent

impaired nucleoid distribution [19, 55, 56]. Taken together, our observations of reduced mtDNA

content, mtRNA levels, as well as changes in single-stranded mtDNA products, Ssbp1

localization, and nucleoid size following Mfn1/2-deficiency suggest a role for Tfam in mediating

the effects of Mfn1/2 in β-cell function.

To test whether increasing mtDNA copy number was sufficient to improve β-cell function

in fusion-deficient islets, we overexpressed Tfam in β-Mfn1/2^{DKO} and littermate control islets. We

transduced intact islets using adenoviral vectors encoding the human form of TFAM, thus allowing

us to track overexpression of TFAM using human-specific TFAM anti-sera (Figures 7a-b and

S6a). Transduction efficiency in β-cells was estimated at ~40-50% (Figure S6a). Despite

limitations in transduction efficiency and expression, TFAM overexpression significantly

ameliorated the reductions in both GSIS and mtDNA copy number in β-Mfn1/2^{DKO} islets (Figures

7c-d and S6b). Therefore, Mfn1 and 2 direct β-cell function, at least in part, through the regulation

of Tfam-mediated mtDNA copy number control.

Mitofusin agonists restore mtDNA content independent of changes in mitochondrial

structure

We next asked if pharmacologically targeting mitofusins could restore mtDNA levels in models of

mitofusin deficiency independent of changes in mitochondrial structure. To this end, we employed

recently described tandem pharmacological agonists, (2-{2-[(5-cyclopropyl-4-phenyl-4H-1,2,4-

triazol-3-yl)sulfanyl]propanamido}-4H,5H,6H-cyclopenta[b]thiophene-3-carboxamide and 1-[2-

(benzylsulf-anyl)ethyl]-3-(2-methylcyclohexyl)urea (hereafter called Mfn agonists), which together

have been previously shown to improve mitochondrial morphology in either Mfn1- or Mfn2-

deficient fibroblasts [57]. These agonists require some residual mitofusin levels to be effective [57].

Thus, we initially treated control and β-Mfn2^{KO} islets, which exhibit reduced mtDNA content without

changes in mitochondrial structure (Figures 3a, 2b-c, and 2e). To avoid systemic effects of

modifying mitochondrial fission/fusion balance in other metabolic tissues [1], we utilized Mfn

agonists on isolated islets. Interestingly, Mfn agonists restored mtDNA content to baseline in β-

Mfn2[KO] islets without affecting mitochondrial morphology or inducing transcriptional regulators of mitochondrial biogenesis (Figures 8a and S7a-b). As anticipated, Mfn agonists were ineffective in the setting of combined absence of Mfn1 and 2 and were unable to rescue reductions in mtDNA content or Tfam protein in β-Mfn1/2[DKO] islets (Figures S7c-d).

We next evaluated the effects of Mfn agonists in other models of mitofusin deficiency, as Mfn2 expression is reduced in several models of β-cell dysfunction [9, 10]. We assessed the expression of Mfn1 and 2 in islets isolated from the leptin receptor deficient *db/db* BKS mouse model, which develops obesity, progressive β-cell mitochondrial and insulin secretory dysfunction, and eventual β-cell failure [58, 59, 60, 61, 62]. We observed that Mfn2, but not Mfn1, protein levels were reduced in islets of 10–12-week-old *db/db* mice when compared to *db/+* controls (Figure 8b), measured at an age prior to β-cell failure [58, 59]. Similar to observations in β-Mfn2[KO] islets, *db/db* islets exhibited reduced mtDNA content without alterations in mitochondrial structure (Figures 8c and 8e). Concordantly, treatment with Mfn agonists rescued mtDNA levels in *db/db* islets, while leaving mitochondrial structure and islet architecture unchanged (Figures 8c-e). We also observed that Mfn agonists improved GSIS in the islets of *db/db* mice (Figure 8f). Taken together, these results indicate that Mfn agonists restore mtDNA content and improve GSIS independent of changes in mitochondrial structure. Moreover, these results extend the association between mitofusin deficiency, mtDNA depletion, and impaired GSIS in β-cells.

DISCUSSION

Here, we identify that Mfn1 and 2 promote mitochondrial fitness to fuel glucose-stimulated insulin release in pancreatic β-cells and maintain glucose homeostasis. Mfn1 or Mfn2 are individually dispensable for β-cell function, but together coordinately control mitochondrial respiration, structure, and mtDNA content. Mfn1 and 2 were thought to regulate cellular metabolism principally through the control of mitochondrial architecture. Our single allele mouse models directly demonstrate that Mfn1/2-deficiency is coupled to glucose intolerance through mtDNA loss rather than mitochondrial structure. Further, we observe that increased mtDNA copy number following Tfam overexpression is capable of ameliorating impaired GSIS in fusion-deficient β-cells. Thus, our studies position the maintenance of mtDNA content as the principal task necessary for Mfn1 and 2 to regulate β-cell function.

Our description of a central role for Mfn1 and 2 in β-cells through mtDNA content rather than mitochondrial structure elucidates a previously unappreciated role for mitochondrial fusion on cellular metabolic function. Mfn1 and 2 act in a complementary manner to support mitochondrial metabolism from development to adulthood. Both germline and cardiomyocyte-specific deletion of Mfn1 and 2 are associated with developmental or early postnatal demise [16, 18], presumed to be primarily caused by defects in mitochondrial network dynamics. Later descriptions of reduced mtDNA levels in constitutive or inducible loss of Mfn1/2 in skeletal muscle did not discern the physiologic importance of mtDNA depletion versus structure following fusion-deficiency [15, 17]. Our results clarify a primary physiologic impact of mtDNA depletion rather than mitochondrial structure following Mfn1/2 deficiency in β-cells that may also underlie the physiologic importance of Mfn1 and 2 in other metabolic tissues. To our knowledge, this is the first time that these functions of Mfn1/2 have been separated in a physiological context.

We observe that β-Mfn1/2DKO islets attempt unsuccessfully to compensate for mtDNA depletion. Our analyses of first strand replication suggest that Mfn1/2-deficient β-cells increase the activation of early steps in mtDNA replication, which is supported by increased Ssbp1

localization to nucleoids to potentially bind newly formed ss-mtDNA. Despite these changes,

mtDNA content remains low in Mfn1/2-deficient β-cells, suggesting that the increased

commitment to replication is insufficient. Indeed, increased nucleoid Ssbp1 may be engaged in

binding abnormal single-stranded replication products due to fusion and replication defects [19, 53].

The increased expression of *Top1mt* found in Mfn1/2-deficient β-cells (Figure 5e) could also

represent a response to increased mitochondrial replication stress [63].

Loss of Mfn1/2 in β-cells leads to severe hyperglycemia, reduced GSIS, and mtDNA

depletion, which bears similarity to a previous report of selective Tfam deficiency in β-cells [25].

Notably, previously described β-cell specific Tfam knockout mice developed progressive loss of

β-cell mass with age [25]. We found that β-cell mass was unchanged in young β-Mfn1/2^{DKO} mice,

but it is possible that they would similarly develop glucotoxicity and an age-related loss of β-cell

mass over time. Alternatively, complete depletion of mtDNA, which occurs in the setting of Tfam-

deficiency but does not occur in β-Mfn1/2^{DKO} mice, might be required for loss of β-cell mass.

Furthermore, the transgenic *RIP2*-Cre used in the β-cell Tfam knockout model (in contrast to the

highly specific *Ins1*-Cre knockin strain we used in β-Mfn1/2^{DKO} mice) is known to result in off-target

recombination[25, 64], which may in part underlie phenotypic differences between the models. In

future experiments, it will be intriguing to examine Tfam and Mfn1/2-deficient mice generated

using the same Cre-recombinase strains to determine the contribution of Tfam-dependent and

independent functions on Mfn1/2 in β-cell health and function.

A related concern in the Mfn1/2-deficiency models is the use of a constitutive Cre strain,

which may yield phenotypic defects due to recombination during β-cell development. However,

an independent group found that mice with inducible Mfn1/2-deficiency only in adult β-cells

developed a highly similar phenotype to constitutive β-Mfn1/2^{DKO} mice [65], suggesting that

developmental defects are unlikely to be a major contributor to the phenotype we observe.

While our data suggest that Tfam mediates the effects of Mfn1/2 on mtDNA content and

GSIS in β-cells, the mechanisms for post-transcriptional loss of Tfam in β-Mfn1/2^{DKO} islets are

unclear. Loss of Tfam may be due to increased LonP1 expression, which is known to degrade

Tfam [49, 50]. Further support for a role for LonP1 as a contributor to cellular dysfunction include

studies demonstrating reduction of nuclear-encoded subunits of Complex I and III following

increases in LonP1 expression [66, 67, 68], an effect we also observe following Mfn1/2-deficiency

(Figure 3d). Mfn1/2-deficiency has also been linked to reductions in nuclear-encoded Complex I

and III subunits via instability in respiratory supercomplexes, which have been observed following

LonP1 overexpression [15, 66, 67, 68]. However, the function and transcriptional regulation of *LonP1* in

β-cells are unknown and require future study.

Mfn2 is frequently observed at MAMs [69, 70]. MAMs are essential for communication and

Ca^{2+} transfer between the ER and mitochondria, are important for β-cell function and GSIS, and

have been suggested to regulate mtDNA replication [39, 40, 41, 71, 72, 73]. Thus, alterations in MAMs in

β-Mfn1/2^{DKO} islets could lead to impaired mtDNA replication. Our initial functional assessments of

MAMs by measurement of MAM-dependent lipids did not reveal differences between control and

β-Mfn1/2^{DKO} islets (Figures S4f-g). We also did not observe ER dilation or overt changes in

mitochondrial-ER proximity upon analysis of transmission EM in β-Mfn1/2^{DKO} islets (Figures 2b

and S2b). Additionally, a recent report found that inducible loss of Mfn1/2 in adult β-cells did not

alter MAM-dependent ER calcium release [65]. Despite these findings, our results do not exclude

the potential that impaired MAM function at β-cell mtDNA nucleoids contributes to the impairments

in mtDNA replication and content we observe. Indeed, our results demonstrating the importance

for a single allele of *Mfn2* over *Mfn1* in the maintenance of glycemic control and mtDNA content

could suggest a role for Mfn2-dependent MAM function in β-cell health and function, as Mfn1 is

not found at MAMs. Alternatively, the regulation of MAMs by Mfn2, as well as the MAM-dependent

role of Mfn2 in mtDNA replication, remains controversial [74, 75]. Recent work also implicates

mitochondrial fusion as necessary to maintain the stoichiometry of the protein components within

the mtDNA replisome [19]. Future studies will be valuable to dissect the molecular details underlying

the connections between mitochondrial fusion, MAMs, and control of β-cell mtDNA replication.

Mitochondrial structural defects and age-related mtDNA depletion combine to impair β-

cell function to increase T2D risk [6, 7, 76, 77, 78], thus, improving mitochondrial function would have

significant translational benefits to restore β-cell health in diabetes. Our studies demonstrate the

feasibility of targeting mtDNA copy number to improve GSIS in the setting of mitofusin deficiency.

While our results in *db/db* islets are promising, loss of the leptin receptor could lead to

mitochondrial defects that are particularly amenable to improvement by Mfn agonists. Thus, it will

be intriguing to determine if Mfn agonists can also reverse β-cell dysfunction in diet-induced

models of obesity *in vivo.* Further, recently identified compounds targeting pyrimidine metabolism

to induce *Mfn1/2* transcription [79] will be of interest to determine if they can improve β-cell function

and mtDNA content. Our observation of the upregulation of mitochondrial pyrimidine transporter

Slc25a33 in β-Mfn1/2[DKO] islets is consistent with links between pyrimidine metabolism and

mitochondrial fusion, which suggest these compounds could hold promise to enhance β-cell

mitochondrial function. As expression of MFN1 and MFN2 is reduced in cardiomyocytes and

skeletal muscle of patients with diabetes, respectively [80, 81, 82], activators of Mfn1/2 may have

additional benefits beyond improving β-cell function. Future studies will be required to fully

characterize the efficacy of overcoming mitochondrial defects *in vivo* to treat diabetes.

METHODS

Animals

Mfn1[loxP/loxP] and *Mfn2*[loxP/loxP] mice possessing loxP sites flanking exon 4 of the *Mfn1* gene and exon

6 of the *Mfn2* gene, respectively, were purchased from Jackson Laboratories [13]. *Mfn1*[loxP] and

Mfn2[loxP] mice were mated to *Ins1*-Cre [83] to generate experimental groups. *Ins1*-Cre–alone,

Mfn1[loxP/loxP] and *Mfn2*[loxP/loxP] mice were phenotypically indistinguishable from each other and

combined as controls (Ctrl; see Figures S1b-c). *Ins1*-Cre–alone were also phenotypically

indistinguishable from wild-type C57BL/6N controls, consistent with previous reports from our

group and others [21, 83]. All animals were maintained on the C57BL/6N background. *db*/+ and *db*/*db*

mice, maintained on the BKS background, were purchased from Jackson Laboratories. Animals

were housed on a standard 12-hour light/12-hour dark cycle at room temperature and 30-70%

humidity, with ad libitum access to food and water. All mice were fed PicoLab® Laboratory Rodent

Diet 5L0D (LabDiet) containing 28% protein, 13% fat, and 57% carbohydrates. All studies and

endpoints in all figures were completed using both male and female mice, and results from both

sexes were combined in all experimental groups. Animal studies were approved by the University

of Michigan Institutional Animal Care and Use Committee.

Mitochondrial respirometry

Islet respirometry was measured using an XF96 extracellular flux analyzer (Seahorse Bioscience)

according to manufacturer's instructions. Briefly, 20 similarly sized islets were plated per well of

a Seahorse spheroid plate pre-treated with CellTak (Corning), similar to previously published

approaches [84]. Prior to respirometry profiling, islets were incubated for 1 hour in an atmospheric

CO_2 incubator at 37°C and supplemented with mouse islet culture media [85, 86] comprised of pH

7.4 unbuffered RPMI1640 media (Seahorse Bioscience) and 2mM glucose prior to flux analyzer

assays. Analysis was performed at baseline, upon exposure to 20mM glucose, and again after

treatment with 2 µM antimycin A. OCR measurements were normalized to islet number confirmed

by light microscopy (Zeiss Stemi 2000) after the completion of respirometry assays.

Transmission electron microscopy

Mouse islets were pelleted, fixed in 3% glutaraldehyde and 3% paraformaldehyde in 0.1 M

Cacodylate buffer (CB; pH 7.2) overnight at 4°C. Islets were then embedded in 2% agarose as

described previously [87]. Samples were then subjected to osmification in 1.5% $K_4Fe(CN)_6$ + 2%

OsO_4 in 0.1 CB for 1 h, dehydrated by serial washes in EtOH (30%, 50%, 70%, 80%, 90%, 95%

and 100%) and embedded in Spurr's resin by polymerization at 60° C for 24h. Polymerized resins

were then sectioned at 90nm thickness using a Leica EM UC7 ultramicrotome and imaged at

70kV using a JEOL 1400 TEM equipped with an AMT CMOS imaging system. Mitochondrial

structures (Aspect Ratio, Perimeter, Circularity) were analyzed and quantified by ImageJ. Form

factor assessments were analyzed by ImageJ and quantified as described [22].

Imaging of islet and organelle morphology and subcellular localization analysis

Mitochondrial morphology and subcellular localization analyses were performed on immuno-

stained paraffin-embedded mouse pancreas tissue sections and dissociated islet cells as

previously described [21]. Briefly, pancreatic tissue sections were fixed in 4% PFA overnight,

embedded in paraffin and sectioned. Following dewaxing and rehydration, antigen retrieval was

performed with 10mM citric acid (pH 6.0). Dissociated islet cells were fixed in 4% PFA for 15

minutes and centrifuged onto glass slides. Pancreatic sections and dissociated islets were then

blocked using 5% donkey serum for 1hr at room temperature followed by immunostaining. Islet

architecture was analyzed using isolated mouse islets fixed in 4% PFA overnight followed by a

20 min incubation with methylene-blue solution (Sigma) and OCT embedding. Cryo-sectioned

islets were rinsed in 75% ethanol for 5 minutes to remove methylene blue, permeabilized with

0.2% Triton-X 100 for 15 minutes, blocked using 5% donkey serum for 1h and immunostained.

467   Images were captured with an IX81 microscope (Olympus) using an ORCA Flash4 CMOS

digital camera (Hamamatsu). Immunostained pancreatic sections, cryo-sectioned and dispersed

mouse islets were captured with Z-stack images and subjected to deconvolution (CellSens;

Olympus). Mitochondrial morphology and nucleoids were visualized using 3D-renderings

generated with Imaris® imaging software (Bitplane). Quantitative 3D assessments of

mitochondrial morphology and network were performed on ImageJ using Mitochondria Analyzer

plugin [22]. Co-localization analyses were performed on Z-stack images of immunostained

dissociated islet cells using the Coloc2 plugin on ImageJ. Changes in relative nucleoid size were

quantified from 3D deconvolution images of immuno-stained dissociated islet cells, detected by

perinuclear anti-DNA antibody staining, using ImageJ.

mtDNA content and replication assays

Relative mtDNA content by qPCR with SYBR-based detection (Universal SYBR Green Supermix;

Biorad) was conducted as previously described [88] following DNA isolation with the Blood/Tissue

DNeasy kit (Qiagen) according to manufacturer's protocols. First strand mtDNA replication levels

were measured by qPCR following digestion by Mnl I, which site specifically digest double

stranded DNA [89]. Briefly, primers specific to 7S mtDNA (5'-

ACTCTTCTCTTCCATATGACTATCCC-3' and 5'-GGCCCTGAAGTAAGAACCAGATGT-3'),

whose amplicon contains a Mnl I site, were used to quantify relative single-stranded mtDNA

forming the D-loop. Single-stranded DNA generated from the extension of 7S region was

measured using primers specific to CYTB (5'-TGCATACGCCATTCTACGCTCA-3' and 5'-

GTGATTGGGCGGAATATTAGGCTTC-3'), which also contains an Mnl I site. Primers specific to

COXI (5'-GCAGGAGCATCAGTAGACCTAACA-3' and 5'-GCGGCTAGCACTGGTAGTGATAAT-

3'), whose amplicon does not contain an Mnl I site, were used to quantify non-replicating double

stranded mtDNA. The ratio of 7S or CYTB amplicons to the COXI amplicon were then calculated

after qPCR of Mnl I digested DNA to determine the relative amount of single stranded mtDNA at

the D-loop strand or initiation of first strand replication, respectively.

RNA sequencing, differential expression, and functional analysis

Sequencing was performed by the Advanced Genomics Core at University of Michigan Medical

School. Total RNA was isolated, and DNase-treated using commercially available kits (Omega

Biotek and Ambion, respectively). Libraries were constructed and subsequently subjected to

151bp paired-end cycles on the NovaSeq-6000 platform (Illumina). FastQC (v0.11.8) was used

to ensure the quality of data. Reads were mapped to the reference genome GRCm38

(ENSEMBL), using STAR (v2.6.1b) and assigned count estimates to genes with RSEM (v1.3.1).

Alignment options followed ENCODE standards for RNA-seq. FastQC was used in an additional

post-alignment step to ensure that only high-quality data were used for expression quantitation

and differential expression. Data were pre-filtered to remove genes with 0 counts in all samples.

Differential gene expression analysis was performed using DESeq2, using a negative binomial

generalized linear model (thresholds: linear fold change >1.5 or <-1.5, Benjamini-Hochberg FDR

(P_{adj}) <0.05). Plots were generated using variations of DESeq2 plotting functions and other

packages with R version 3.3.3. Genes were annotated with NCBI Entrez GeneIDs and text

descriptions. Functional analysis, including candidate pathways activated or inhibited in

comparison(s) and GO-term enrichments, was performed using iPathway Guide (Advaita). The

RNA sequencing data generated in this study have been deposited in the Gene Expression

Omnibus (GEO) database under accession code GSE194204.

Adenoviral transfection

Isolated mouse islets were distended with Accutase for 1 min at 37° C as described previously [90].

Islets were then transduced with 0.15 MOI of Ad.EV (expressing an empty vector) or Ad.hTFAM

(expressing human-specific TFAM) for 48 hrs (viral particles purchased from Vector Biolabs). β-

cell transduction was evaluated on dissociated islet cells by immunostaining with Pdx1 and

human-specific TFAM anti-sera.

Flow cytometry

Following isolation and culture, live mouse islets were exposed to 100 nM Mtphagy dye (Dojindo

Molecular Technologies) for 3 hours to assess time-dependent accumulation of mitochondria to

acidic organelles by the relative fluorescence intensity of the dye per cell as described [91, 92]. Islets

were then dissociated into single cells, stained with DAPI (Thermo Fisher Scientific) and Fluozin-

3 (Thermo Fisher Scientific), and resuspended in phenol red–free islet culture medium as

previously described [21]. Samples were analyzed on an LSR Fortessa flow cytometer (BD

Biosciences). Single cells were gated using forward scatter and side scatter (FSC and SSC,

respectively) plots, DAPI staining was used to exclude dead cells, and Fluozin-3 was used to

identify β-cells as previously described [21, 93]. Mtphagy measurements in β-cells were made using

488 nm excitation laser with a 710 nm emission filter and analyzed using FlowJo (Tree Star Inc.).

A total of 5,000 β-cells was quantified from each independent islet preparation. Gating strategies

with representative plots are provided in Supplementary Figure S8.

Glucose and insulin measurements

IPGTT, ITT, and serum insulin measurements were performed as described previously [85, 86].

Briefly, mice were fasted for 6 hrs prior to glucose and insulin measurements. For IPGTT or ITT,

mice were injected intraperitoneally with 2 g/kg glucose or 0.75 U/kg Insulin (Novolin R),

respectively. Tail blood was collected for measurements of blood glucose. For *in vivo* insulin

release measurements, tail blood was collected following glucose injection, centrifuged at 2000

rpm for 10 min at 4 °C and stored at −20 °C. Static insulin secretion assays on isolated islets were

performed following 1hr incubations with 2mM and 16.7mM glucose in HEPES-supplemented

KRB buffer containing 135 mM NaCl, 4.7 mM KCl, 1.2 mM KH_2PO_4, 5 mM $NaHCO_3$, 1.2 mM

$MgSO_4.7H_2O$, 1 mM $CaCl_2$, 10 mM HEPES and 0.1% BSA (pH 7.4). Samples were analyzed for

insulin release using Mouse Ultrasensitive Insulin ELISA (ALPCO) as per the manufacturer's

protocol.

Islet isolation and cell culture

Mouse islets were isolated and cultured as previously described [85, 86]. Cell treatments included

DMSO (Fisher) and antimycin A (Sigma). Mitofusin agonist treatment included an equimolar

mixture of two previously described compounds (0.5 µM each; [57]), which were administered for

24 hours (2-{2-[(5-cyclopropyl-4-phenyl-4H-1,2,4-triazol-3-yl)sulfanyl]propanamido}-4H,5H,6H-

cyclopenta[b]thiophene-3-carboxamide, CAS# 920868-45-7 and 1-[2-(benzylsulfanyl)ethyl]-3-(2-

methylcyclohexyl)urea, CAS# 1007818-44-1; Enamine Ltd.).

Western blotting, quantitative PCR, and immunostaining

All assays were performed as previously described [21, 85, 86]. Mouse pancreatic islets were lysed

with radioimmunoprecipitation assay buffer containing protease and phosphatase inhibitors

(Calbiochem), and insoluble material was removed by centrifugation. Equal amounts of proteins

were resolved on 4%–20% gradient Tris-glycine gels (Bio-Rad) and transferred to nitrocellulose

membranes (Bio-Rad). Membranes were then blocked in 5% milk for 1hr and immunoblotting

was performed using Cyclophilin B (1:5000; ThermoFisher, Catalog# PA1-027A), Drp1 (1:1000;

CST #8570), Lonp1 (1:1000, ProteinTech, Catalog# 15440-1-AP), Mfn1 (1:1000; Abcam,

Catalog# ab126575), Mfn2 (1:1000; Abcam, Catalog# ab56889), mt-Cytb (1:500; ProteinTech,

Catalog# 55090-1-AP), Opa1 (1:1000, BD Transduction laboratories, Catalog# 612606), Total

Oxphos antibody cocktail (1:1000; Abcam, ab110413), Polg (1:1000, Abcam, Catalog#

ab128899), Ssbp1 (1:250; Atlas antibodies; Catalog# HPA002866), Anti-human TFAM (1:1000,

PhosphoSolutions, Catalog# 1999-hTFAM), Anti-mouse TFAM (1:1000; PhosphoSolutions 2001-

TFAM), Tom20 (1:1000; Cell Signaling Technology, Catalog# 42406), Twnk (1:1000,

ProteinTech, Catalog# 13435-1-AP), Vinculin (1:5000; Millipore, Catalog# CP74) and species-

specific HRP-conjugated secondary antibodies (Vector Laboratories). Immunostaining on

pancreatic sections, cryo-sectioned and dispersed mouse islets was performed using Anti-DNA

(1:50; American Research Products, Catalog# PA1-027A), Insulin (1:250; Dako, Catalog#

A0564), Insulin (1:100, Abcam, Catalog# ab7842), Glucagon (1:2000; SantaCruz, Catalog# sc-

13091), Pdx1 (1:250; Abcam, Catalog# ab47383), SDHA (1:100, Abcam, Catalog# ab14715),

Ssbp1 (1:250; Atlas antibodies, Catalog# HPA002866), Somatostatin (1:500, Abcam, Catalog#

ab30788) and species-specific Cy2, Cy3 and Cy5-conjugated secondary antibodies (Jackson

Immunoresearch). β-cell mass was quantified as total insulin-positive area/total pancreatic area

multiplied by pancreatic wet weight using stitched images of complete pancreatic sections [86]. All

antibodies used for Western blotting and immunostaining are listed in Supplementary Table 1.

RNA samples were reverse transcribed using High-Capacity cDNA Reverse Transcription

Kit (Thermo Fisher Scientific). Quantitative reverse transcription PCR (qRT-PCR) was performed

with SYBR-based detection (Universal SYBR Green Supermix; Biorad) using primers for

Ppargc1a (5'-ACTATGAATCAAGCCACTACAGAC-3' and 5'- TTCATCCCTCTTGAGCCTTTCG-

3'), *Nrf1* (5'-ACAGATAGTCCTGTCTGGGGAAA-3' and 5'- TGGTACATGCTCACAGGGATCT-3')

and   *Tfam*   (5'-AGCTTCCAGGAGGCAAAGGATGAT-3'   and   5'-

ACTTCAGCCATCTGCTCTTCCCAA-3').

Cholesterol esterification and phosphatidylserine assay

Total amounts of free cholesterol and cholesterol esters were determined in mouse islet lysates

as per the manufacturer's instructions (Promega). Phosphatidylserine levels were measured in

mouse islet lysates as per the manufacturer's instructions (LSBio).

Statistics

In all figures, data are presented as means ± SEM, and error bars denote SEM, unless otherwise

noted in the legends. Statistical comparisons were performed using unpaired two-tailed Student's

t-tests, one-way or two-way ANOVA, followed by Tukey's or Sidak's post-hoc test for multiple

comparisons, as appropriate (GraphPad Prism). A P value < 0.05 was considered significant.

Data Availability

The authors declare that data supporting the findings of this study are available within the article

and its supplementary information files or from the corresponding author (S.A.S.) upon request.

RNA sequencing data used for differential gene expression-, pathway- and GO- enrichment

analyses have been deposited in the GEO database under the accession number GSE194204.

The reference genome used for RNA sequencing was obtained from ENSEMBL (Mus_musculus

- Ensembl genome browser 105). All uncropped western blots and data values for all figures are

provided in the source data file.

Code Availability

High-throughput RNA sequencing data was analyzed using DESeq2 (v1.26.0) to determine

differential gene expression. iPathway Guide (Advaita version 1910) was used to perform

pathway and GO-term analysis.

REFERENCES
604
605 1. Sebastian D, Palacin M, Zorzano A. Mitochondrial Dynamics: Coupling Mitochondrial
606 Fitness with Healthy Aging. *Trends Mol Med* **23**, 201-215 (2017).
607
608 2. Westermann B. Mitochondrial fusion and fission in cell life and death. *Nat Rev Mol Cell*
609 *Biol* **11**, 872-884 (2010).
610
611 3. Schrepfer E, Scorrano L. Mitofusins, from Mitochondria to Metabolism. *Mol Cell* **61**, 683-
612 694 (2016).
613
614 4. Kaufman BA, Li C, Soleimanpour SA. Mitochondrial regulation of beta-cell function:
615 maintaining the momentum for insulin release. *Molecular aspects of medicine* **42**, 91-104
616 (2015).
617
618 5. Maechler P. Mitochondrial function and insulin secretion. *Mol Cell Endocrinol* **379**, 12-18
619 (2013).
620
621 6. Anello M*, et al.* Functional and morphological alterations of mitochondria in pancreatic
622 beta cells from type 2 diabetic patients. *Diabetologia* **48**, 282-289 (2005).
623
624 7. Masini M*, et al.* Ultrastructural alterations of pancreatic beta cells in human diabetes
625 mellitus. *Diabetes Metab Res Rev* **33**, (2017).
626
627 8. Wigger L*, et al.* Multi-omics profiling of living human pancreatic islet donors reveals
628 heterogeneous beta cell trajectories towards type 2 diabetes. *Nat Metab* **3**, 1017-1031
629 (2021).
630
631 9. Safia Costes TG, Alexandra E Butler, Sam Sereda, Orian Shirihai and Peter Cawood
632 Butler. Disruption of the Endoplasmic Reticulum-Mitochondria Interface By Toxic IAPP
633 Oligomers May Compromise Beta-Cell Mitochondrial Network Integrity and Function in
634 Type 2 Diabetes. In: *The Endocrine Society Annual Meeting* (ed Society E) (2015).
635
636 10. Stoehr JP*, et al.* Genetic obesity unmasks nonlinear interactions between murine type 2
637 diabetes susceptibility loci. *Diabetes* **49**, 1946-1954 (2000).
638
639 11. Campbell CT, Kolesar JE, Kaufman BA. Mitochondrial transcription factor A regulates
640 mitochondrial transcription initiation, DNA packaging, and genome copy number. *Biochim*
641 *Biophys Acta* **1819**, 921-929 (2012).
642

643   12.   Filograna R, Mennuni M, Alsina D, Larsson NG. Mitochondrial DNA copy number in
644   human disease: the more the better? *FEBS Lett*, (2020).

646   13.   Chen H, McCaffery JM, Chan DC. Mitochondrial fusion protects against
647   neurodegeneration in the cerebellum. *Cell* **130**, 548-562 (2007).

649   14.   Liesa M, Shirihai OS. Mitochondrial dynamics in the regulation of nutrient utilization and
650   energy expenditure. *Cell Metab* **17**, 491-506 (2013).

652   15.   Bell MB, Bush Z, McGinnis GR, Rowe GC. Adult skeletal muscle deletion of Mitofusin 1
653   and 2 impedes exercise performance and training capacity. *J Appl Physiol (1985)* **126**,
654   341-353 (2019).

656   16.   Chen H, Detmer SA, Ewald AJ, Griffin EE, Fraser SE, Chan DC. Mitofusins Mfn1 and Mfn2
657   coordinately regulate mitochondrial fusion and are essential for embryonic development.
658   *J Cell Biol* **160**, 189-200 (2003).

660   17.   Chen H*, et al.* Mitochondrial fusion is required for mtDNA stability in skeletal muscle and
661   tolerance of mtDNA mutations. *Cell* **141**, 280-289 (2010).

663   18.   Papanicolaou KN*, et al.* Mitofusins 1 and 2 are essential for postnatal metabolic
664   remodeling in heart. *Circ Res* **111**, 1012-1026 (2012).

666   19.   Silva Ramos E*, et al.* Mitochondrial fusion is required for regulation of mitochondrial DNA
667   replication. *PLoS Genet* **15**, e1008085 (2019).

669   20.   Twig G*, et al.* Fission and selective fusion govern mitochondrial segregation and
670   elimination by autophagy. *The EMBO journal* **27**, 433-446 (2008).

672   21.   Sidarala V*, et al.* Mitophagy protects beta cells from inflammatory damage in diabetes.
673   *JCI Insight* **5**, (2020).

675   22.   Chaudhry A, Shi R, Luciani DS. A pipeline for multidimensional confocal analysis of
676   mitochondrial morphology, function, and dynamics in pancreatic beta-cells. *Am J Physiol*
677   *Endocrinol Metab* **318**, E87-E101 (2020).

679   23.   Hemel I, Engelen BPH, Luber N, Gerards M. A hitchhiker's guide to mitochondrial
680   quantification. *Mitochondrion* **59**, 216-224 (2021).

682   24.   Mulder H. Transcribing beta-cell mitochondria in health and disease. *Molecular*
683   *metabolism* **6**, 1040-1051 (2017).

685   25.   Silva JP*, et al.* Impaired insulin secretion and beta-cell loss in tissue-specific knockout
686   mice with mitochondrial diabetes. *Nature genetics* **26**, 336-340 (2000).

688   26.   Narendra D, Tanaka A, Suen DF, Youle RJ. Parkin is recruited selectively to impaired
689   mitochondria and promotes their autophagy. *J Cell Biol* **183**, 795-803 (2008).

691   27.   Kersten S. Physiological regulation of lipoprotein lipase. *Biochim Biophys Acta* **1841**, 919-
692   933 (2014).

694 28. Haythorne E*, et al.* Diabetes causes marked inhibition of mitochondrial metabolism in
695   pancreatic beta-cells. *Nat Commun* **10**, 2474 (2019).
696
697 29. Gerst F*, et al.* The Expression of Aldolase B in Islets Is Negatively Associated With Insulin
698   Secretion in Humans. *J Clin Endocrinol Metab* **103**, 4373-4383 (2018).
699
700 30. Scarpulla RC. Transcriptional paradigms in mammalian mitochondrial biogenesis and
701   function. *Physiol Rev* **88**, 611-638 (2008).
702
703 31. Calvo SE, Clauser KR, Mootha VK. MitoCarta2.0: an updated inventory of mammalian
704   mitochondrial proteins. *Nucleic Acids Res* **44**, D1251-1257 (2016).
705
706 32. Zhang H*, et al.* Human mitochondrial topoisomerase I. *Proc Natl Acad Sci U S A* **98**,
707   10608-10613 (2001).
708
709 33. Ng AC, Baird SD, Screaton RA. Essential role of TID1 in maintaining mitochondrial
710   membrane potential homogeneity and mitochondrial DNA integrity. *Mol Cell Biol* **34**, 1427-
711   1437 (2014).
712
713 34. Di Noia MA*, et al.* The human SLC25A33 and SLC25A36 genes of solute carrier family 25
714   encode two mitochondrial pyrimidine nucleotide transporters. *J Biol Chem* **289**, 33137-
715   33148 (2014).
716
717 35. Zurita Rendon O, Shoubridge EA. LONP1 Is Required for Maturation of a Subset of
718   Mitochondrial Proteins, and Its Loss Elicits an Integrated Stress Response. *Mol Cell Biol*
719   **38**, (2018).
720
721 36. Venkatesh S, Lee J, Singh K, Lee I, Suzuki CK. Multitasking in the mitochondrion by the
722   ATP-dependent Lon protease. *Biochim Biophys Acta* **1823**, 56-66 (2012).
723
724 37. Lu B*, et al.* Roles for the human ATP-dependent Lon protease in mitochondrial DNA
725   maintenance. *J Biol Chem* **282**, 17363-17374 (2007).
726
727 38. Sebastian D*, et al.* Mfn2 deficiency links age-related sarcopenia and impaired autophagy
728   to activation of an adaptive mitophagy pathway. *The EMBO journal* **35**, 1677-1693 (2016).
729
730 39. Friedman JR, Lackner LL, West M, DiBenedetto JR, Nunnari J, Voeltz GK. ER tubules
731   mark sites of mitochondrial division. *Science* **334**, 358-362 (2011).
732
733 40. Lewis SC, Uchiyama LF, Nunnari J. ER-mitochondria contacts couple mtDNA synthesis
734   with mitochondrial division in human cells. *Science* **353**, aaf5549 (2016).
735
736 41. Murley A*, et al.* ER-associated mitochondrial division links the distribution of mitochondria
737   and mitochondrial DNA in yeast. *Elife* **2**, e00422 (2013).
738
739 42. Ma H*, et al.* Germline and somatic mtDNA mutations in mouse aging. *PLoS One* **13**,
740   e0201304 (2018).
741
742 43. Area-Gomez E. Assessing the function of mitochondria-associated ER membranes.
743   *Methods Enzymol* **547**, 181-197 (2014).
744

44. Vance JE. Molecular and cell biology of phosphatidylserine and phosphatidylethanolamine metabolism. *Prog Nucleic Acid Res Mol Biol* **75**, 69-111 (2003).

45. Ekstrand MI*, et al.* Mitochondrial transcription factor A regulates mtDNA copy number in mammals. *Hum Mol Genet* **13**, 935-944 (2004).

46. Kang D, Kim SH, Hamasaki N. Mitochondrial transcription factor A (TFAM): roles in maintenance of mtDNA and cellular functions. *Mitochondrion* **7**, 39-44 (2007).

47. Kang I, Chu CT, Kaufman BA. The mitochondrial transcription factor TFAM in neurodegeneration: emerging evidence and mechanisms. *FEBS Lett* **592**, 793-811 (2018).

48. Larsson NG*, et al.* Mitochondrial transcription factor A is necessary for mtDNA maintenance and embryogenesis in mice. *Nature genetics* **18**, 231-236 (1998).

49. Matsushima Y, Goto Y, Kaguni LS. Mitochondrial Lon protease regulates mitochondrial DNA copy number and transcription by selective degradation of mitochondrial transcription factor A (TFAM). *Proc Natl Acad Sci U S A* **107**, 18410-18415 (2010).

50. Lu B*, et al.* Phosphorylation of human TFAM in mitochondria impairs DNA binding and promotes degradation by the AAA+ Lon protease. *Mol Cell* **49**, 121-132 (2013).

51. Gensler S*, et al.* Mechanism of mammalian mitochondrial DNA replication: import of mitochondrial transcription factor A into isolated mitochondria stimulates 7S DNA synthesis. *Nucleic Acids Res* **29**, 3657-3663 (2001).

52. Pohjoismaki JL, Goffart S. Of circles, forks and humanity: Topological organisation and replication of mammalian mitochondrial DNA. *Bioessays* **33**, 290-299 (2011).

53. Gustafson MA, Sullivan ED, Copeland WC. Consequences of compromised mitochondrial genome integrity. *DNA Repair (Amst)* **93**, 102916 (2020).

54. Tan K, Fujimoto M, Takii R, Takaki E, Hayashida N, Nakai A. Mitochondrial SSBP1 protects cells from proteotoxic stresses by potentiating stress-induced HSF1 transcriptional activity. *Nat Commun* **6**, 6580 (2015).

55. Kasashima K, Sumitani M, Endo H. Human mitochondrial transcription factor A is required for the segregation of mitochondrial DNA in cultured cells. *Exp Cell Res* **317**, 210-220 (2011).

56. West AP*, et al.* Mitochondrial DNA stress primes the antiviral innate immune response. *Nature* **520**, 553-557 (2015).

57. Rocha AG*, et al.* MFN2 agonists reverse mitochondrial defects in preclinical models of Charcot-Marie-Tooth disease type 2A. *Science* **360**, 336-341 (2018).

58. Guo S*, et al.* Inactivation of specific beta cell transcription factors in type 2 diabetes. *J Clin Invest* **123**, 3305-3316 (2013).

795 59. Ishida E, Kim-Muller JY, Accili D. Pair Feeding, but Not Insulin, Phloridzin, or Rosiglitazone
796   Treatment, Curtails Markers of beta-Cell Dedifferentiation in db/db Mice. *Diabetes* **66**,
797   2092-2101 (2017).
798
799 60. Li J*, et al.* Amorphous solid dispersion of Berberine mitigates apoptosis via
800   iPLA2beta/Cardiolipin/Opa1 pathway in db/db mice and in Palmitate-treated MIN6 beta-
801   cells. *Int J Biol Sci* **15**, 1533-1545 (2019).
802
803 61. Liu J*, et al.* Rhein protects pancreatic beta-cells from dynamin-related protein-1-mediated
804   mitochondrial fission and cell apoptosis under hyperglycemia. *Diabetes* **62**, 3927-3935
805   (2013).
806
807 62. Zhao Z*, et al.* Protection of pancreatic beta-cells by group VIA phospholipase A(2)-
808   mediated repair of mitochondrial membrane peroxidation. *Endocrinology* **151**, 3038-3048
809   (2010).
810
811 63. Pommier Y, Sun Y, Huang SN, Nitiss JL. Roles of eukaryotic topoisomerases in
812   transcription, replication and genomic stability. *Nat Rev Mol Cell Biol* **17**, 703-721 (2016).
813
814 64. Wicksteed B*, et al.* Conditional gene targeting in mouse pancreatic ss-Cells: analysis of
815   ectopic Cre transgene expression in the brain. *Diabetes* **59**, 3090-3098 (2010).
816
817 65. Georgiadou E*, et al.* Mitofusins Mfn1 and Mfn2 are required to
818   preserve glucose-but not incretin- stimulated beta cell connectivity and insulin secretion.
819   *bioRxiv*, 2020.2004.2022.055384 (2021).
820
821 66. Besse A, Brezavar D, Hanson J, Larson A, Bonnen PE. LONP1 de novo dominant
822   mutation causes mitochondrial encephalopathy with loss of LONP1 chaperone activity and
823   excessive LONP1 proteolytic activity. *Mitochondrion* **51**, 68-78 (2020).
824
825 67. Peter B*, et al.* Defective mitochondrial protease LonP1 can cause classical mitochondrial
826   disease. *Hum Mol Genet* **27**, 1743-1753 (2018).
827
828 68. Quiros PM*, et al.* ATP-dependent Lon protease controls tumor bioenergetics by
829   reprogramming mitochondrial activity. *Cell Rep* **8**, 542-556 (2014).
830
831 69. Sebastian D*, et al.* Mitofusin 2 (Mfn2) links mitochondrial and endoplasmic reticulum
832   function with insulin signaling and is essential for normal glucose homeostasis. *Proc Natl
833   Acad Sci U S A* **109**, 5523-5528 (2012).
834
835 70. de Brito OM, Scorrano L. Mitofusin 2 tethers endoplasmic reticulum to mitochondria.
836   *Nature* **456**, 605-610 (2008).
837
838 71. Dingreville F*, et al.* Differential Effect of Glucose on ER-Mitochondria Ca(2+) Exchange
839   Participates in Insulin Secretion and Glucotoxicity-Mediated Dysfunction of beta-Cells.
840   *Diabetes* **68**, 1778-1794 (2019).
841
842 72. Madec AM, Perrier J, Panthu B, Dingreville F. Role of mitochondria-associated
843   endoplasmic reticulum membrane (MAMs) interactions and calcium exchange in the
844   development of type 2 diabetes. *Int Rev Cell Mol Biol* **363**, 169-202 (2021).
845

73.   Rieusset J. Role of Endoplasmic Reticulum-Mitochondria Communication in Type 2 Diabetes. *Adv Exp Med Biol* **997**, 171-186 (2017).

74.   Cosson P, Marchetti A, Ravazzola M, Orci L. Mitofusin-2 independent juxtaposition of endoplasmic reticulum and mitochondria: an ultrastructural study. *PLoS One* **7**, e46293 (2012).

75.   Filadi R, Greotti E, Turacchio G, Luini A, Pozzan T, Pizzo P. Mitofusin 2 ablation increases endoplasmic reticulum-mitochondria coupling. *Proc Natl Acad Sci U S A* **112**, E2174-2181 (2015).

76.   Nile DL*, et al.* Age-related mitochondrial DNA depletion and the impact on pancreatic Beta cell function. *PLoS One* **9**, e115433 (2014).

77.   Cree LM*, et al.* Age-related decline in mitochondrial DNA copy number in isolated human pancreatic islets. *Diabetologia* **51**, 1440-1443 (2008).

78.   Chinnery PF*, et al.* Mitochondrial DNA haplogroups and type 2 diabetes: a study of 897 cases and 1010 controls. *J Med Genet* **44**, e80 (2007).

79.   Miret-Casals L*, et al.* Identification of New Activators of Mitochondrial Fusion Reveals a Link between Mitochondrial Morphology and Pyrimidine Metabolism. *Cell Chem Biol* **25**, 268-278 e264 (2018).

80.   Bach D*, et al.* Expression of Mfn2, the Charcot-Marie-Tooth neuropathy type 2A gene, in human skeletal muscle: effects of type 2 diabetes, obesity, weight loss, and the regulatory role of tumor necrosis factor alpha and interleukin-6. *Diabetes* **54**, 2685-2693 (2005).

81.   Montaigne D*, et al.* Myocardial contractile dysfunction is associated with impaired mitochondrial function and dynamics in type 2 diabetic but not in obese patients. *Circulation* **130**, 554-564 (2014).

82.   Zorzano A, Liesa M, Palacin M. Mitochondrial dynamics as a bridge between mitochondrial dysfunction and insulin resistance. *Arch Physiol Biochem* **115**, 1-12 (2009).

83.   Thorens B, Tarussio D, Maestro MA, Rovira M, Heikkila E, Ferrer J. Ins1(Cre) knock-in mice for beta cell-specific gene recombination. *Diabetologia* **58**, 558-565 (2015).

84.   Taddeo EP*, et al.* Individual islet respirometry reveals functional diversity within the islet population of mice and human donors. *Molecular metabolism* **16**, 150-159 (2018).

85.   Pearson G*, et al.* Clec16a, Nrdp1, and USP8 Form a Ubiquitin-Dependent Tripartite Complex That Regulates beta-Cell Mitophagy. *Diabetes* **67**, 265-277 (2018).

86.   Soleimanpour SA*, et al.* The diabetes susceptibility gene Clec16a regulates mitophagy. *Cell* **157**, 1577-1590 (2014).

87.   Soleimanpour SA*, et al.* Calcineurin signaling regulates human islet {beta}-cell survival. *J Biol Chem* **285**, 40050-40059 (2010).

896 88.   Kolesar JE, Wang CY, Taguchi YV, Chou SH, Kaufman BA. Two-dimensional intact
897   mitochondrial DNA agarose electrophoresis reveals the structural complexity of the
898   mammalian mitochondrial genome. *Nucleic Acids Res* **41**, e58 (2013).
899
900 89.   Karamanlidis G, Nascimben L, Couper GS, Shekar PS, del Monte F, Tian R. Defective
901   DNA replication impairs mitochondrial biogenesis in human failing hearts. *Circ Res* **106**,
902   1541-1548 (2010).
903
904 90.   Reissaus CA*, et al.* A Versatile, Portable Intravital Microscopy Platform for Studying Beta-
905   cell Biology In Vivo. *Sci Rep* **9**, 8449 (2019).
906
907 91.   Iwashita H*, et al.* Live Cell Imaging of Mitochondrial Autophagy with a Novel Fluorescent
908   Small Molecule. *ACS Chem Biol* **12**, 2546-2551 (2017).
909
910 92.   Konig J*, et al.* Mitochondrial contribution to lipofuscin formation. *Redox Biol* **11**, 673-681
911   (2017).
912
913 93.   Jayaraman S. A novel method for the detection of viable human pancreatic beta cells by
914   flow cytometry using fluorophores that selectively detect labile zinc, mitochondrial
915   membrane potential and protein thiols. *Cytometry A* **73**, 615-625 (2008).
916

Acknowledgements

S.A.S was supported by the JDRF (CDA-2016-189, SRA-2018-539, COE-2019-861), the NIH

(R01 DK108921, U01 DK127747), the Department of Veterans Affairs (I01 BX004444), the Brehm

family, and the Anthony family. E.L.-D. was supported by the NIH (T32 AI007413 and T32-

AG000114). G.L.P. was supported by the American Diabetes Association (19-PDF-063). B.A.K.

was supported by the Department of Veterans Affairs (I01 BX004444). The JDRF Career

Development Award to S.A.S. is partly supported by the Danish Diabetes Academy and the Novo

Nordisk Foundation. We acknowledge the Microscopy, Imaging and Cellular Physiology Core of

the University of Michigan DRC (P30 DK020572) for assistance with imaging studies. We thank

the University of Michigan Flow Cytometry Core for assistance with flow cytometry studies. Next

generation sequencing was carried out in the Advanced Genomics Core at the University of

Michigan. We acknowledge support from the Bioinformatics Core of the University of Michigan's

Biomedical Research Core Facilities. We thank Drs. K. Claiborn, C. Rutledge, N. Desai, G. Rutter,

A. Garner, A. Menon, M. Torres, and members of the Soleimanpour laboratory for helpful advice.

Author Contributions

V.S. conceived, designed and performed experiments, interpreted results, drafted and reviewed

the manuscript. J.Z. and E.L-D. designed and performed experiments, interpreted results, and

reviewed the manuscript. G.L.P. and E.C.R. designed and performed experiments and interpreted

results. E.M.W. designed and performed experiments, interpreted results, edited, and reviewed

the manuscript. B.A.K. designed studies, interpreted results and reviewed the manuscript. S.A.S.

conceived and designed the studies, interpreted results, drafted, edited, and reviewed the

manuscript.

Conflict of interest statement

The authors have declared that no conflict of interest exists.

 FIGURE LEGENDS

Figure 1. Loss of both Mfn1 and Mfn2 in β-cells impairs glucose tolerance and glucose-

stimulated insulin release.

a Expression of Mfn1 and Mfn2 by Western blot (WB) in islets isolated from 8-10-week-old Ctrl,

β-Mfn1[KO], β-Mfn2[KO] and β-Mfn1/2[DKO] mice. Cyclophilin B serves as a loading control.

Representative of 6 independent mice/group for Mfn1 and 10 independent mice/group for Mfn2.

Band intensity of each lane of displayed representative blot (shown as fold change compared to

Ctrl, expression normalized to cyclophilin B) above each blot in red. **b** Densitometry (normalized

to cyclophilin B) from studies in Figure 1A. n=6/group for Mfn1 and 10/group for Mfn2. Data are

presented as Mean ± SEM. *$p<0.0005$ by one-way ANOVA followed by Tukey's post-hoc test for

multiple comparisons. **c** Blood glucose concentrations measured during IPGTT of 8-week-old Ctrl

(n=28), β-Mfn1[KO] (n=7), β-Mfn2[KO] (n=6) and β-Mfn1/2[DKO] (n=15) littermates. Data are presented

as Mean ± SEM. ($p<0.0005$ by two-way ANOVA for β-Mfn1/2[DKO] mice vs Ctrl followed by Sidak's

post-hoc test for multiple comparisons). **d** Serum insulin concentrations (n=7-24/group) measured

during *in vivo* glucose-stimulated insulin release testing in 8-week-old Ctrl (n=24), β-Mfn1[KO] (n=7),

β-Mfn2[KO] (n=10) and β-Mfn1/2[DKO] (n=14) littermates. Data are presented as mean ± SEM.

*$p<0.0005$ for β-Mfn1/2[DKO] mice vs Ctrl and β-Mfn1[KO], $p=0.004$ for β-Mfn1/2[KO] vs β-Mfn2[KO] by

one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons. **e** Glucose-

stimulated insulin secretion following static incubations in 2 mM and 16.7 mM glucose, performed

in isolated islets of 8-week-old Ctrl (n=16), β-Mfn1[KO] (n=8), β-Mfn2[KO] (n=11) and β-Mfn1/2[DKO]

(n=10) littermates. Data are presented as mean from independent mice ± SEM. *$p<0.0005$ by

one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons. **f** Pancreatic β-cell

mass measured in 10-week-old Ctrl (n=10), β-Mfn1[KO] (n=4), β-Mfn2[KO] (n=4) and β-Mfn1/2[DKO]

(n=6) littermates. Data are presented as mean ± SEM. Uncropped western blots and source data

are provided as a Source Data file.

Figure 2. Combined Mfn1 and Mfn2 deficiency impairs mitochondrial respiration and

structure in β-cells.

a OCR measured following exposure to 2mM glucose, 20mM glucose, and 2µM antimycin A in

islets isolated from 8-10-week-old Ctrl (n=9) and β-Mfn1/2^{DKO} (n=8) mice by a Seahorse flux

analyzer. Data are presented as mean from independent mice ± SEM. *p=0.0176 by one-way

ANOVA followed by Sidak's post-hoc test for multiple

comparisons. **b** Representative transmission EM images of β-cells from 12-week-old Ctrl, β-

Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} littermates. n=3/group. Mitochondria are highlighted with red

arrows. **c** Quantification of mitochondrial morphology in Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-

Mfn1/2^{DKO} β-cells from transmission EM images (~125 independent mitochondria scored/animal).

n=3/group. Data are presented as mean ± SEM. *p=0.0087 for Aspect Ratio, p=0.004 for Form

factor and Circularity, p=0.037 for Perimeter by one-way ANOVA followed by Tukey's post-hoc

test for multiple comparisons. **d** Imaris® generated three-dimensional reconstruction of

deconvolution immunofluorescence Z-stack images at 100X magnification stained for SDHA (see

inset image – red) from pancreatic sections of Ctrl and β-Mfn1/2^{DKO} mice. β-cells were identified

by insulin co-staining (inset: insulin – green). Each unique color represents a separate β-cell

mitochondrial network cluster. Representative image of 6 independent mice/group. **e** β-cell

mitochondrial morphology and network analysis of deconvolution immunofluorescence Z-

stack images at 100X magnification stained for SDHA (and insulin) from pancreatic sections of

Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice by MitoAnalyzer. n=6/group (~150 β-cells/animal

were quantified). Data are presented as mean ± SEM. *p=0.0006 for Average mitochondrial

volume, p=0.0089 for Sphericity, p=0.0032 for Branches per mitochondria, p=0.0039 for Branch

junctions by one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons. Source

data are provided as a Source Data file.

Figure 3. Mfn1/2-deficient β-cells exhibit reduced mtDNA content.

a Relative mtDNA content measured by qPCR (normalized to nuclear DNA expression) in isolated islets of 8-week-old Ctrl (n=7), β-Mfn1^{KO} (n=5), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO} (n=5) littermates. Data are presented as mean ± SEM. *p=0.0037 for β-Mfn2^{KO} vs Ctrl, p<0.0005 for β-Mfn1/2^{DKO} vs Ctrl by one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons. **b** Heatmap representing relative expression (compared to littermate controls; Ctrl) of mitochondrial encoded transcripts in islets isolated from β-Mfn1^{KO} (n=4), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO} (n=5) mice. §* Benjamini-Hochberg FDR (Padj) <0.05 comparing β-Mfn2^{KO} (§) and β-Mfn1/2^{DKO} (*) to Ctrl. **c** Expression of OXPHOS subunit proteins by WB in islets isolated from 8-10-week-old Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice. Representative of 4 independent mice/group. **d** OXPHOS subunit densitometry (normalized to Vinculin) from studies in Figure 3D. n=4/group; Data are presented as mean ± SEM. *p=0.02 for Uqcrc2, p<0.05 for mt-co1, p<0.001 for Ndufb8, p<0.005 for mt-Cytb by one-way ANOVA followed by Tukey's post-hoc test for multiple comparisons. **e** Expression of Tom20 by WB in islets isolated from 8-10-week-old Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice. Representative image (top) of 6 independent mice/group. Tom20 densitometry, normalized to vinculin (bottom). n=6/group. Data are presented as mean ± SEM. Uncropped western blots and source data are provided as a Source Data file.

Figure 4. A single allele of *Mfn2*, but not *Mfn1*, maintains glycemic control by preserving β-cell mtDNA content despite impaired mitochondrial structure.

a Imaris® generated three-dimensional reconstruction of deconvolution immunofluorescence Z-stack images at 100X magnification stained for SDHA (see inset image – red) from pancreatic sections of Ctrl, β-Mfn1/2^{DKO}, β-Mfn1^{KO}Mfn2$^{+/-}$ and β-Mfn1$^{+/-}$Mfn2^{KO} mice. β-cells were identified by insulin co-staining (inset: insulin – green). Each unique color represents a separate β-cell mitochondrial network cluster. Representative image of 6 independent mice/group. **b** β-cell mitochondrial morphology and network analysis of deconvolution immunofluorescence Z-stack images from pancreatic sections stained for insulin and SDHA of Ctrl, β-Mfn1/2^{DKO}, β-

Mfn1^{KO}Mfn2$^{+/-}$ and β-Mfn1$^{+/-}$Mfn2^{KO} mice by MitoAnalyzer. n=6/group (50-200 β-cells/animal were

quantified). *p=0.005 for Average mitochondrial volume, p=0.05 for Sphericity, p=0.05 for

Branches per mitochondria, p=0.05 for Branch junctions by one-way ANOVA followed by Sidak's

post-hoc test for multiple comparisons. **c** Blood glucose concentrations measured during IPGTT

of 8-week-old Ctrl (n=28), β-Mfn1/2^{DKO} (n=15), β-Mfn1^{KO}Mfn2$^{+/-}$ (n=13) and β-Mfn1$^{+/-}$Mfn2^{KO}

(n=17) littermates. (p<0.0005 by two-way ANOVA for β-Mfn1$^{+/-}$Mfn2^{KO} and β-Mfn1/2^{DKO} mice vs

Ctrl followed by Sidak's post-hoc test for multiple comparisons). **d** Serum insulin concentrations

(n=11-24/group) measured during *in vivo* glucose-stimulated insulin release testing in 8-week-

old Ctrl (n=24), β-Mfn1/2^{DKO} (n=14), β-Mfn1^{KO}Mfn2$^{+/-}$ (n=11) and β-Mfn1$^{+/-}$Mfn2^{KO} (n=16)

littermates. *p<0.0005 by one-way ANOVA followed by Sidak's post-hoc test for multiple

comparisons. **e** Relative mtDNA content measured by qPCR (normalized to nuclear DNA

expression) in isolated islets of 8-week-old Ctrl (n=7), β-Mfn1/2^{DKO} (n=5), β-Mfn1^{KO}Mfn2$^{+/-}$ (n=5)

and β-Mfn1$^{+/-}$Mfn2^{KO} (n=5) littermates. Data are presented as mean ± SEM. *p<0.0005 by one-

way ANOVA followed by Sidak's post-hoc test for multiple comparisons. Of note, studies in Ctrl

and β-Mfn1/2^{DKO} mice were performed together alongside all β-Mfn1^{KO}, β-Mfn2^{KO}, β-

Mfn1^{KO}Mfn2$^{+/-}$ (n=11) and β-Mfn1$^{+/-}$Mfn2^{KO} (n=16) littermates and thus may appear twice for

purposes of relevant comparisons (in Figures 1C, 1D, 2D, 2E, 3A and again in Figures 4A-E).

Source data are provided as a Source Data file.

Figure 5. Mfn1/2-deficiency induces expression of genes associated with mtDNA

replication.

a Hierarchical clustering of RNAseq samples displaying the 500 genes with the highest mean

expression in islets of 12-week-old Ctrl (n=6), β-Mfn1^{KO} (n=4), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO}

(n=5) mice. **b** Pathway analysis of top differentially regulated pathways in β-Mfn1/2^{DKO} islets (n=5)

compared to littermate Ctrl islets (n=6), demarcated by both false discovery rate (FDR) and the

number of significantly differentially expressed genes (DEGs) per pathway. **c** Volcano plot

depicting differential RNA expression in islets of β-Mfn1/2^{DKO} (n=5) mice compared to littermate

Ctrl mice (n=6). Significantly differentially expressed genes demarcated by -log10 FDR > or < 2

and log2 fold change (FC) > or < 0.6. **d** Volcano plot depicting differential RNA expression of

MitoCarta 2.0 targets in islets of β-Mfn1/2^{DKO} (n=5) mice compared to littermate Ctrl mice (n=6).

Significantly differentially expressed genes demarcated by -log10 FDR > or <2 and log2 fold

change (FC) > or <0.6. **e** Differential RNA expression heatmap of selected genes from islets of β-

Mfn1^{KO} (n=4), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO} (n=5) mice compared to littermate Ctrl mice (n=6).

up. ‡§* Benjamini-Hochberg FDR (Padj) <0.05 comparing β-Mfn1^{KO}(‡), β-Mfn2^{KO}(§) and β-

Mfn1/2^{DKO}(*) to Ctrl.

Figure 6. Mfn1 and 2 maintain expression of Tfam in β-cells.

a Expression of Tfam, key replisome proteins, and LonP1 by Western blot (WB) in islets isolated

from 11-week-old Ctrl, β-Mfn1^{KO}, β-Mfn2^{KO} and β-Mfn1/2^{DKO} mice. Representative

of 4 independent mice per group. **b** Densitometry (normalized to Vinculin) from studies in Figure

6A. n=4/group; Data are presented as mean ± SEM. *p=0.03 for Tfam, p=0.01 for Lonp1 by one-

way ANOVA followed by Sidak's post-hoc test for multiple comparisons. **c** Quantification of single

strand mtDNA products measured by qPCR in isolated islets from 8-week-old Ctrl (n=13), β-

Mfn1^{KO} (n=5), β-Mfn2^{KO} (n=5) and β-Mfn1/2^{DKO} (n=9) mice. Data are presented as mean ± SEM.

*p=0.0007 by one-way ANOVA followed by Sidak's post-hoc test for multiple comparisons.

d Deconvolution immunofluorescence image at 100X magnification of islets from Ctrl and β-

Mfn1/2^{DKO} mice stained for Ssbp1 (red), mtDNA (green), and Pdx1 (blue). Representative of 8

independent experiments. White arrows demarcate co-localized Ssbp1+ mtDNA+

structures. **e** Quantification of mitochondrial nucleoid Ssbp1 localization (Ssbp1+ mtDNA+ co-

localization) in Ctrl and β-Mfn1/2^{DKO} β-cells from studies depicted in Figure 6D by Pearson's

correlation coefficient (PCC). n=8/group. Data are presented as mean ± SEM. *p=0.0142 by two-

tailed t-test. (~100 β-cells from each animal per group were analyzed). **f** Quantification of nuclear

Ssbp1 localization in Ctrl and β-Mfn1/2^{DKO} β-cells from studies depicted in Figure 6D by

Pearson's correlation coefficient (PCC). n=8/group. Data are presented as mean ± SEM.

*p=0.0008 by two-tailed t-test. (~100 β-cells from each animal per group were

analyzed). **g** Imaris® generated 3D reconstruction of deconvolution immunofluorescence Z-stack

images at 100X magnification stained for DNA of β-cells (see inset image – Pdx1, blue, DNA,

green) from Ctrl and β-Mfn1/2^{DKO} mice. Colors represent relative mitochondrial volume.

Representative image of 8 independent mice/group. **h** Quantification of relative nucleoid volume

of Ctrl and β-Mfn1/2^{DKO} β-cells from studies depicted in Figure 6g. n=8/group. Data are presented

as mean ± SEM. *p=0.0127 by two-tailed t-test. (~100 β-cells from each animal per group were

analyzed). Uncropped western blots and source data are provided as a Source Data file.

Figure 7. Mfn1 and 2 maintain β-cell function through Tfam-mediated mtDNA copy number

control.

a Expression of mouse-specific Tfam and human-specific TFAM by WB in islets isolated from 10-

15-week-old Ctrl and β-Mfn1/2^{DKO} mice, transduced with empty vector control (Ad.EV) or human

TFAM-overexpressing (Ad.hTFAM) adenoviral particles. Representative of 4 independent

mice/group. **b** Densitometry (normalized to cyclophilin B) from studies in Figure 7A. n=4/group;

Data are presented as mean ± SEM. p=NS (non-significant) by two-tailed t-test. **c** Glucose-

stimulated insulin secretion following static incubations in 2mM and 16.7 mM glucose, performed

in isolated Ctrl and β-Mfn1/2^{DKO} islets following transduction with Ad.EV or Ad.hTFAM adenoviral

particles. n=6/group. Data are presented as mean ± SEM. *p<0.0005 for Ctrl + Ad.EV vs β-

Mfn1/2^{DKO} + Ad.EV, p=0.0148 for β-Mfn1/2^{DKO} + Ad.EV vs β-Mfn1/2^{DKO} + Ad.hTFAM by one-way

ANOVA followed by Tukey's post-hoc test for multiple comparisons. **d** Relative mtDNA content

measured by qPCR (normalized to nuclear DNA expression) from isolated Ctrl and β-

Mfn1/2^{DKO} islets following transduction with Ad.EV or Ad.hTFAM adenoviral particles.

n=12/group. Data are presented as mean ± SEM. *p<0.0005 for Ctrl + Ad.EV vs β-Mfn1/2^{DKO} +

Ad.EV, p=0.007 for β-Mfn1/2^{DKO} + Ad.EV vs β-Mfn1/2^{DKO} + Ad.hTFAM by two-tailed t-test.

Uncropped western blots and source data are provided as a Source Data file.

Figure 8. Mitofusin agonists restore mtDNA depletion following Mfn2 deficiency

independent of changes in mitochondrial structure.

a Relative mtDNA content measured by qPCR (normalized to nuclear DNA expression) from

isolated islets of Ctrl and β-Mfn2^{KO} mice following treatment with vehicle (Veh) or 0.5 μM Mfn

agonists for 24 hrs. n=4/group. Data are presented as mean ± SEM. *p=0.0138 for Ctrl + Veh vs

β-Mfn2KO + Veh, p=0.0302 for β-Mfn2^{KO} + Veh vs β-Mfn2^{KO} + Mfn agonists by one-way ANOVA

followed by Sidak's post-hoc test for multiple comparisons. **b** Expression of Mfn1 and Mfn2 by

WB in islets isolated from 10-12-week-old *db*/+ and *db*/*db* mice. Vinculin and cyclophilin B serve

as loading controls. Representative image (left) of 3 independent mice/group for Mfn1 and 4

independent mice/group for Mfn2. Mfn1 (n=3) and Mfn2 (n=4) densitometry, normalized to vinculin

or cyclophilin B (right). Data are presented as mean ± SEM. *p=0.0095 by two-tailed t-test.

c Relative mtDNA content measured by qPCR (normalized to nuclear DNA expression) from

isolated islets of 10-12-week-old *db*/+ (n=6) and *db*/*db* (n=5) mice following treatment with vehicle

(Veh) or 0.5 μM mitofusin agonists for 24 hrs. Data are presented as mean ± SEM. *p=0.0038 for

db/+ + Veh vs *db*/*db* + Veh, p=0.0428 for *db*/*db* + Veh vs *db*/*db* + Mfn agonists by two-tailed t-

test. **d** Immunofluorescence images at 20X magnification of islets from 10-12-week-old *db*/+ and

db/*db* mice stained for Insulin (green), Glucagon (red), Somatostatin (SST; white), and DAPI

(blue) following treatment with vehicle (Veh) or 0.5 μM mitofusin agonists for 24 hrs. n=3/group.

e β-cell mitochondrial morphology and network analysis by MitoAnalyzer of deconvolution

immunofluorescence Z-stack images, stained for SDHA (and insulin), from pancreatic islets of 10-

12-week-old *db*/+ and *db*/*db* mice following treatment with vehicle (Veh) or 0.5 μM mitofusin

agonists for 24 hrs. n=6/group (~150-200 β-cells/animal were quantified). Data are presented as

mean ± SEM. **f** Glucose-stimulated insulin secretion following static incubations in 2 mM and 16.7

mM glucose, performed in isolated islets of 10-12-week-old *db*/+ and *db*/*db* mice following

treatment with vehicle (Veh) or 0.5 μM mitofusin agonists for 24 hrs. n=11/group. Data are

presented as mean ± SEM. *p=0.0092 by two-tailed t-test. Uncropped western blots and source

data are provided as a Source Data file.

1124

Gene	β-Mfn1^KO	β-Mfn2^KO	β-Mfn1/2^DKO
mt-Nd1§*			
mt-Nd2§*			
mt-Nd4l§*			
mt-Nd3§*			
mt-Nd4§*			
mt-Nd5§*			
mt-Nd6§*			
mt-Co1§*			
mt-Co2§*			
mt-Co3§*			
mt-Cytb§*			
mt-Atp8§*			
mt-Atp6§*			