A modified fluctuation assay reveals a natural mutator phenotype that drives mutation spectrum variation within *Saccharomyces cerevisiae*

Pengyao Jiang[1], Anja R. Ollodart[1,2], Vidha Sudhesh[1], Alan J. Herr[3], Maitreya J. Dunham[1], Kelley Harris[1,4]

1 Department of Genome Sciences, University of Washington, Seattle, WA
2 Molecular and Cellular Biology Program, University of Washington, Seattle, WA
3 Department of Laboratory Medicine and Pathology, University of Washington, Seattle, WA
4 Department of Computational Biology, Fred Hutchinson Cancer Research Center, Seattle, WA

Abstract

Although studies of *Saccharomyces cerevisiae* have provided many insights into mutagenesis and DNA repair, most of this work has focused on a few laboratory strains. Much less is known about the phenotypic effects of natural variation within *S. cerevisiae*'s DNA repair pathways. Here, we use natural polymorphisms to detect historical mutation spectrum differences among several wild and domesticated *S. cerevisiae* strains. To determine whether these differences are likely caused by genetic mutation rate modifiers, we use a modified fluctuation assay with a *CAN1* reporter to measure *de novo* mutation rates and spectra in 16 of the analyzed strains. We measure a 10-fold range of mutation rates and identify two strains with distinctive mutation spectra. These strains, known as AEQ and AAR, come from the panel's "Mosaic beer" clade and share an enrichment for C>A mutations that is also observed in rare variation segregating throughout the genomes of several Mosaic beer and Mixed origin strains. Both AEQ and AAR are haploid derivatives of the diploid natural isolate CBS 1782, whose rare polymorphisms are enriched for C>A as well, suggesting that the underlying mutator allele is likely active in nature. We use a plasmid complementation test to show that AAR and AEQ share a mutator allele in the DNA repair gene *OGG1*, which excises 8-oxoguanine lesions that can cause C>A mutations if left unrepaired.

Introduction

Mutations are a double-edged sword. At the molecular level, they usually arise as a spontaneous consequence of DNA replication errors or damage and are the ultimate cause of genetic diseases (Iossifov et al. 2014; Sebat et al. 2007; Antonarakis and Beckmann 2006; Crow 1997; Nei 1983). All organisms have evolved complex mechanisms for keeping mutation rates low and safeguarding their genetic information as it is passed from generation to generation (Beckman and Loeb 1993; Eisen and Hanawalt 1999); in multicellular organisms, these mechanisms also safeguard somatic tissues from mutations that can cause cancer and age-related decline (Alexandrov et al. 2013; Lawrence A. Loeb 2016; Risques and Kennedy 2018). A low mutation rate is essential for long-term population survival, and the larger and more complex a genome is, the lower the mutation rate must be to prevent deleterious

mutations from arising faster than natural selection can eliminate them (Eigen 1971; Drake 1991; Sung, Ackerman, et al. 2012; Acosta et al. 2015). Over long time scales, however, mutations also serve as the raw material for evolution. Although beneficial mutations are rare occurrences, they are essential for the acquisition of novel phenotypes and adaptations (Gompel et al. 2005; McGregor et al. 2007).

A large body of theory has been written to describe how natural selection might act on the mutation rate to balance these beneficial and deleterious effects (Sturtevant 1937; Kimura 1967; Leigh 1970; Johnson 1999; André and Godelle 2006; Sung, Ackerman, et al. 2012). One prediction is that organisms living in more changeable environments might evolve higher mutation rates than organisms living in more stable environments, assuming that the environment determines whether a higher rate of beneficial mutations is likely to counterbalance a higher rate of deleterious mutations. This prediction has been borne out in laboratory evolution experiments, where mutator phenotypes sometimes emerge in populations that are forced to tolerate challenging conditions (Good et al. 2017; Tenaillon et al. 2016; Couce et al. 2017) and mutator strains are often observed to take over chemostat populations by producing beneficial mutations at a higher rate than competing non-mutator strains (Chao and Cox 1983). Many mutator phenotypes in *E. coli* have been linked to defects in DNA repair enzymes (Horst, Wu, and Marinus 1999). Mutator phenotypes also commonly occur in cancer (L. A. Loeb 2001; Prindle, Fox, and Loeb 2010), likely either because of relaxed selection against cellular dysfunction or because it is beneficial for cancer cells to adapt rapidly to their aggressive growth niche. However, it is less clear how much mutation rate variation exists within and between natural populations, and if such variation exists, whether it is maintained by natural selection. The drift barrier hypothesis predicts that mutator alleles will usually be deleterious because they produce more damaging mutations than beneficial ones, but that mutator alleles with relatively small effects may persist in populations because they are not deleterious enough to be efficiently eliminated (Lynch et al. 2016).

Although next-generation sequencing has rapidly increased our ability to measure the genetic variation that currently exists within populations, the extent of mutation rate variation is still more difficult and expensive to measure. One of the original methods for measuring mutation rates is the Luria-Delbrück fluctuation assay (Lang and Murray 2008; Gou, Bloom, and Kruglyak 2019; Luria and Delbrück 1943), in which a population of microorganisms is allowed to grow clonally for a controlled length of time, then challenged with a form of artificial selection that kills most cells except for those that have happened to acquire specific resistance mutations. The mutation rate can then be calculated from the number of colonies that manage to grow after this artificial selection is imposed.

Though fluctuation assays are an elegant and efficient way for measuring the mutation rates of specific reporter genes, the results are potentially sensitive to the reporter gene being used and where it is located within the genome (Lang and Murray 2011, 2008); in addition, they are not applicable to multicellular organisms. These drawbacks have motivated the development of newer methods that take advantage of high-throughput sequencing, such as mutation

accumulation (MA) assays in which a laboratory population is serially bottlenecked for many generations, eliminating most effects of natural selection and allowing mutations to be directly counted by sequencing at the end of the experiment. MA studies have been used to estimate mutation rates in a wide variety of organisms (Lynch et al. 2008; Sharp et al. 2018; Zhu et al. 2014; Farlow et al. 2015; Wang et al. 2019; Keightley et al. 2009), including a recent study of 9 heterozygous *S.cerevisiae* strains (Dutta, Dutreux, and Schacherer 2021). However, higher throughput MA experiments require DNA repair genes to be knocked out to induce higher mutation rates that can be accurately measured with fewer generations of labor-intensive propagation, as in (Liu and Zhang 2021).

An alternative source of information about mutational processes in strains with low mutation rates is genetic variation among related individuals who share common ancestors. Polymorphic sites are easier and cheaper to discover than new mutations, since they are present at a higher density within the genome and often shared among several individuals. Mining polymorphisms for information about mutation rates can be difficult since their abundance is affected by genetic drift and natural selection (Scally and Durbin 2012; Ségurel, Wyman, and Przeworski 2014; Zhu, Sherlock, and Petrov 2017), but despite these limitations, they have provided surprisingly strong evidence for the existence of historical changes to the mutation spectrum, meaning the tendency of mutations to occur most often in certain nucleotide contexts (Hwang and Green 2004). In humans, for example, Europeans and South Asians have a significantly higher proportion of TCC>TTC mutations than other human groups (K. Harris 2015; K. Harris and Pritchard 2017), a pattern that is difficult to explain without a recent population-specific increase in the rate of this type of mutation. This pattern might have been caused by either a genetic mutator or an environmental mutagen, but is not explicable by the action of selection or drift or any other process that simply modulates the retention or loss of genetic variation.

Polymorphism data has revealed that each human population and great ape species appears to have a distinctive triplet mutation spectrum, which implies that genetic and/or environmental mutators likely emerge relatively often and act within localized populations to increase mutation rates in specific sequence contexts (K. Harris and Pritchard 2017; Goldberg and Harris 2021). However, identifying these hypothetical mutators is a challenging proposition, not least because some population-specific signatures such as the human TCC>TTC enrichment appear to be relics of mutators that are no longer active. A recent study of *de novo* mutations in diverse human families found some evidence of mutation rate variation between human populations (Kessler et al. 2020), but argued that most of this variation was driven by the environment rather than genetics. Given that humans from different populations tend to be born and raised in different environments, it is extremely challenging to determine the degree to which genetics and/or the environment are responsible for variation of the rates and spectrum of *de novo* mutations accumulating within human populations today.

More is known about the genetic architecture of mutagenesis in model organisms, including the single-celled organism *Saccharomyces cerevisiae,* where it is tractable to disentangle genetic mutator effects from environmental ones by accumulating mutations on different genetic

backgrounds in controlled laboratory environments (Serero et al. 2014; Lang, Parsons, and Gammie 2013; Stirling et al. 2014; Huang et al. 2003; Herr et al. 2011). Many *S. cerevisiae* mutator alleles have been discovered using genetic screens, which involve creating libraries of artificial mutants in the lab and determining which ones have high mutation rates (Stirling et al. 2014). Mutation rates can be elevated by up to a thousand-fold in lines where DNA proofreading and repair capabilities are artificially knocked out (Herr et al. 2011; Serero et al. 2014; Lang, Parsons, and Gammie 2013), and quantitative trait loci with more modest effects have been found to underlie a five-fold range of mutation rate variation among a few natural *S. cerevisiae* strains (Gou, Bloom, and Kruglyak 2019). A more complex mutator phenotype has been observed as a result of epistasis between two incompatible alleles found as natural variation in the mismatch repair genes *MLH1* and *PMS1*, although the natural isolates in which these alleles are found appear to have acquired compensatory variants that suppress this mutator phenotype (Bui et al. 2017; Argueso et al. 2003; Raghavan et al. 2018; Heck et al. 2006). An extreme case of long-term maintenance of a hypermutation lineage comes from phylogenetic analysis of bipolar budding yeasts *Hanseniaspora* which shows large amounts of gene loss, accelerated evolutionary rates and altered mutation spectra (Steenwyk et al. 2019).

Mild environmental stressors, such as high salt, ethanol, and heat, can also alter the mutation rate of *S. cerevisiae* (Liu and Zhang 2019; Voordeckers et al. 2020) as well as *Arabidopsis thaliana* (Jiang et al. 2014; Belfield et al. 2021). The same environmental perturbations can cause detectable changes to both species' mutation spectra. The mutation spectrum has also been observed to depend on whether *S. cerevisiae* is replicating in a haploid or diploid state (Sharp et al. 2018). In addition, environmental mutagens, more complex ploidy, and genetic mutation rate modifiers could all conceivably affect the mutation spectrum of natural variation as it accumulates. However, no study to our knowledge has looked at whether any mutational signatures measured in the laboratory are capable of explaining natural mutation spectrum variation observed in polymorphism data from a model species.

Recently, comprehensive sampling efforts have produced a collection of 1011 natural isolates of *S. cerevisiae* (Peter et al. 2018). This is a uniquely powerful system containing abundant natural variation that accumulated within diverse environments during the recent and ancient evolution of *S. cerevisiae*, and the panel is also amenable for experimental accumulation of mutations over laboratory growth. Many genetic polymorphisms differentiate these strains, and these are relics of mutations that accumulated over many generations on divergent genetic backgrounds adapted to diverse environmental conditions, ranging from forests to beverage fermentation pipelines. Both environmental mutagens and genetic mutators may have created differences among the mutation spectra of these 1011 strains, but only genetically determined mutation spectrum differences should have the potential to be reproduced in the spectra of mutations accumulated in a controlled lab environment.

We hypothesized that yeast strains with outlying spectra of natural polymorphisms are more likely to have distinct *de novo* mutation spectra than strains whose polymorphisms have indistinguishable mutation spectra. The same hypothesis underlies previous inferences of *de*

novo mutation spectrum variation from polymorphism data (K. Harris 2015; K. Harris and Pritchard 2017; Dumont 2019; Goldberg and Harris 2021), and rare mutations found in mouse colonies (Dumont 2019), but this hypothesis has been difficult to test in vertebrate species. To enable such direct testing in *S. cerevisiae,* we describe a new Luria-Delbrück-based assay that efficiently measures the spectra of *de novo* mutations in haploid strains using pooled amplicon sequencing. We then use this assay to identify strains with reproducibly measurable mutator phenotypes that explain the spectrum biases of these strains' polymorphisms. Some proportion of natural mutation spectrum variation might not be reproducible in the lab if it is driven by environmental mutagens, bioinformatic artifacts, or extinct genetic mutators, but our assay has the potential to identify which gradients of mutation spectrum variance are driven by extant genotypic differences.

Results

The mutation spectrum of natural variation in *S. cerevisiae*

To measure the mutation spectrum of genetic variation present in the 1011 *S. cerevisiae* natural isolates (Peter et al. 2018), we polarized single nucleotide polymorphisms using the outgroup *S. paradoxus* (Yue et al. 2017), then classified them into two transition types and four transversion types based on their ancestral and derived alleles. Closely related strains were excluded to avoid overrepresentation of certain groups (Materials and Methods). We calculated the proportion of each mutation type among the derived alleles present in each individual strain, utilizing all derived variants present below 50% frequency (a total of 1,213,508 SNPs). In order to minimize bias from ancestral allele misidentification, we excluded strains with extensive, pre-documented introgression from *S. paradoxus* (Peter et al. 2018). A final set of 729,357 SNPs with ancestral alleles that passed the derived allele frequency filter were used for the following analysis (a subset of the 838,568 SNPs with callable ancestral alleles). Principal component analysis (PCA) on these individual mutation spectra reveals that strains from the same population tend to have more similar mutation spectra than more distantly related strains (Figure 1A, Supplementary Table S1). Some of this structure disappears when SNPs are subsampled to eliminate double-counting of variants that are shared among multiple strains (Supplementary Figure S1), but several clades appear as consistent outliers in both analyses, including the African beer and European wine strains. The compact architecture of the yeast genome makes it infeasible to exclude coding regions and conserved regions, but we obtain similar PC structures when we use only synonymous protein-coding variants and when we use all polymorphisms passing quality filters (Supplementary Figure S2).

Figure 1. Mutation spectra of natural isolates of *S. cerevisiae*

Principal component analysis of segregating mutation spectrum variation from a subset of the 1011 yeast strains. **A**. Mutation spectrum PCA of all natural variants under 50% derived allele frequency. Each strain's mutation spectrum histogram is projected as a single point, colored to indicate its population of origin (Peter et al. 2018). The inset summarizes the loadings of the first and second principal component vectors. **B**. Mutation spectrum PCA of rare variants (derived allele count 2-4). Singleton variants are excluded to minimize the impact of sequencing error. Strains appearing more than 1.8 standard deviations from the origin along both PC1 and PC2 are labeled with their strain names.

Figure 1A shows that the Taiwanese and African beer populations are outliers along PC1. As seen from the principal component loadings, these two groups mainly differ from the rest in the relative proportions of the two transition types (A>G and C>T): Taiwanese strains are enriched for C>T mutations while African beer strains are enriched for A>G mutations. In contrast, PC2 separates the majority of other populations, such as human-associated strains isolated from wine, dairy, and bioethanol production, along a gradient of varying transition/transversion ratio.

Although strains from the same population tend to cluster together, this trend is less pronounced in the 1011 *S. cerevisiae* genomes than in previously reported mutation spectrum PCAs of humans, great apes, and mice (K. Harris and Pritchard 2017; Dumont 2019; Goldberg and Harris 2021). That being said, one methodological difference from these previous studies is that we only partition the yeast mutation spectra into six basic types (A>C, A>G, etc.) rather than the 96 trinucleotide-based types used in analyses of vertebrate mutation spectra, a concession to the small size of the yeast genome. We found that the trinucleotide mutation spectra of yeast exhibit similar PCA structure (Supplementary Figure S3), but that the sparsity of yeast triplet spectra appears to limit their utility.

Figure 1B shows a PCA of rare variant mutation spectra from the same collection of strains used in 1A. We define rare variants as those with derived allele counts of 2, 3, and 4 (423,312 SNPs out of 729,357 fall within this allele count range) and exclude singletons to minimize the impact of sequencing error. These spectra are noisier than the spectra computed from variants

up to 50% frequency, but are potentially more likely to reflect recently active mutational
processes. While rare variant spectra exhibit less clustering by population than those in Figure
1A, a subset of strains from several groups appear as outliers. For example, a few Mixed origin
and Mosaic beer strains are outliers along a C>A mutation gradient, and African beer and
French dairy strains separate out along an A>G mutation gradient. For completeness, we also
examined mutation spectra of singleton variants alone (Materials and Methods), which are the
youngest mutations among polymorphisms (Supplementary Figure S4). It resembles the PCA of
non-singleton rare variants, except that C>A mutation variation explains a larger variation and
becomes the PC1 axis.

Several previous studies have found a puzzling discrepancy between the spectra of *de novo*
mutations and polymorphisms in *S. cerevisiae*: polymorphisms have a transition-to-transversion
(ts/tv) ratio around 3:1, compared to only 1:1 for *de novo* mutations (Agier and Fischer 2012;
Zhu, Sherlock, and Petrov 2017). Our analysis of the 1011 strain collection replicates this finding
(Supplementary Figure S5). We also replicate the prior finding that singletons and higher
frequency variants have nearly identical ts/tv ratios, but that singletons inferred to be young
based on their presence on long shared haplotypes have a lower ts/tv ratio somewhat closer to
that of new mutations (Supplementary Figure S5). We note that a discrepancy between *de novo*
and segregating mutation spectra has also been observed in *A. thaliana*, though in the opposite
direction, as *A. thaliana de novo* mutations are surprisingly enriched for C>T transitions (Weng
et al. 2019).

A scalable experimental pipeline for measuring *de novo* mutation rates and spectra

In order to test whether any of the mutation spectrum differences evident from natural variation
in different *S. cerevisiae* strains are driven by extant genetic mechanisms that increase the rates
of specific mutation types, we set out to measure several strains' *de novo* mutation spectra and
rates experimentally. To this end, we developed an experimental pipeline using the reporter
gene *CAN1*. Traditional reporter gene fluctuation assays only estimate the overall rate of
mutations, but we introduced an extra step that utilizes Illumina sequencing of pooled amplicons
derived from *CAN1* mutants to estimate each strain's mutation spectrum as well.

The gene *CAN1* encodes a transport protein that imports arginine and arginine analogs into
yeast cells from the surrounding growth media. This means that strains with a functional *CAN1*
transporter are sensitive to poisoning by the arginine analog canavanine, while a single
loss-of-function mutation can render such cells able to survive on canavanine media (Whelan,
Gocke, and Manney 1979). Poisoning a culture with canavanine is thus a very efficient method
to select for cells with point mutations in *CAN1*. A limitation of this method is that it only works
on genomes that contain exactly one functional copy of *CAN1*, since canavanine resistance is
recessive. This means that it cannot be used to measure mutation rates in diploid or polyploid
strains directly, which unfortunately include the Taiwanese strains and African beer strains that
are PCA outliers in Figure 1A. However, 133 of the 1011 strains are haploid, leaving many

strains of interest that are amenable to the assay, including several outliers in the rare variant
PCA (Figure 1B).

A schematic overview of our experimental setup is shown in Figure 2. First, we estimated
mutation rates using established fluctuation assay methodology (Lang and Murray 2008; Gou,
Bloom, and Kruglyak 2019), which involves plating multiple independent cultures from each
strain being investigated. We then picked a single colony from each plated culture and grew it to
saturation in canavanine-containing media. We then selected mutants observed to grow in
culture to similar saturation density and pooled them in equal proportions to give each mutant a
roughly equal frequency in the pool. Individual pools of mutants from each strain were then
subjected to PCR amplification of *CAN1* followed by Illumina sequencing. Individual mutants
were called from the sequencing pools using a customized pipeline (Materials and Methods).
Mutations collected from different pools of the same strain were combined to calculate the
strain's mutation spectrum. We aimed to collect roughly 300 mutants per strain, enough to
detect mutation spectrum differences of the magnitude estimated from polymorphism data in
several of the 1011 genomes populations.

270   Figure 2. Schematic overview of the experimental pipeline
271 Overview of the experimental pipeline used to estimate the mutation rate and spectrum for each strain using the
272 reporter gene *CAN1*. First, mutation rates were estimated using fluctuation assays. Independent mutants were then
273 pooled and sequenced to estimate the mutation spectrum of each strain.

Pooling mutants across canavanine media cultures before sequencing allowed us to efficiently estimate mutation spectra at scale, yielding measurements of many individual mutations per library prep. However, pooling too many mutants during this step could have the potential to compromise the pipeline's accuracy by putting the frequency of each mutation too close to the expected frequency of Illumina sequencing errors. To test for this failure mode, we Sanger-sequenced 38 independent mutants generated using the lab strain LCTL1 (SEY6211-MATα). After pooling these 38 mutants, we performed two replicate library preps and Illumina-sequenced both using our standard procedure. Sanger sequencing identified 37 mutants with single nucleotide mutations plus one containing two adjacent mutations (Supplementary Table S3). We expect each of these mutations, which should be present at a frequency of about 1/38 in the pooled culture, to be easily distinguishable from Illumina sequencing errors that occur at a rate of less than 1% per base.

To identify *bona fide* mutations from each Illumina sequencing pool, we developed a pipeline designed to call mutations present at or above an expected frequency that is inversely proportional to the number of mutants being pooled. In order to minimize false positive mutation calls introduced by sequencing errors, we excluded low coverage regions located at the ends of the amplicons (Materials and Methods). We also identified multinucleotide mutations (MNMs) based on the co-occurrence of variation on the same reads (Averof et al. 2000; Schrider, Hourmozdi, and Hahn 2011), separating these complex mutations from single base substitutions and small indels. When we tested this Illumina sequencing pipeline on the same mutant pool (~17,000x coverage) that we previously Sanger sequenced, we detected 37 of the 38 mutations identified by Sanger sequencing, missing only one mutation that occurred at the end of the amplicon located outside our pipeline's callable region. A second Illumina sequencing replicate measured only 36 of these mutations, missing one additional true mutation. Neither Illumina replicate produced any false positives, verifying that the pipeline is accurate enough to permit pooling of up to 40 *CAN1* mutants before each library prep.

Mutation rate variation among haploid natural isolates

We used our pipeline to measure mutation rates and spectra in 16 haploid strains from a wide variety of environments (Supplementary Table S2). We selected these strains such that to the extent possible, they include two euploid strains per clade without any copy number variation at the scale of whole chromosome arms. We also selected two lab strains, LCTL1 and LCTL2, to use as controls, since their mutation rates were previously measured: the mutation rate of LCTL1 was measured using a genome-wide mutation accumulation assay by Sharp et al. (2018), while the mutation rate of LCTL2 (GIL 104, a derivative of W303) was measured by Lang et al. (2008) using a *CAN1* fluctuation assay. Two additional strains from the 1011 collection, AAA and ACS, were selected because their mutation rates had been previously measured in another study (Gou, Bloom, and Kruglyak 2019).

We observed a ten-fold range of mutation rate variation in *CAN1* among the strains we surveyed: from 2.1×10^{-7} to 2.1×10^{-6} canavanine resistance mutations per gene per cell division

(Figure 3). This range of variation is larger than the five-fold range of mutation rate variation
found among six *S. cerevisiae* strains in a recent study (Gou, Bloom, and Kruglyak 2019). All
estimates from different replicates of the same strain were generally consistent with each other,
though three strains (ACS, LCTR2, and AAR) showed close to a 2-fold difference between our
highest and lowest mutation rate measurements. This is within the margin of error observed in
previously published fluctuation assays performed at large scale (Gou, Bloom, and Kruglyak
2019). Among the three strains (LCTL2, AAA, and ACS) with previously published mutation rate
measurements, our results fall within a 1.5-fold range of those estimates, with no particular
trend of upward or downward bias.

We noticed that AAR and AEQ, the two strains with the highest mutation rates, formed larger
colonies during the fixed duration of the experimental growth period compared to many of the
other strains tested. This suggests that AAR and AEQ may have high growth rates. This
motivated us to test for correlation between the mutation rates we measured and strain-specific
growth rates reported in the literature (Peter et al. 2018), but overall, we found no significant
correlation between these attributes (R^2<0.001; *p*=0.95) (Supplementary Figure S6).

We observed that strains from the Mosaic beer, Sake, African palm wine, and Asian
fermentation clades exhibited higher mutation rates than have been previously reported for any
natural *S. cerevisiae* strains. The two strains with the highest mutation rates, roughly 10-fold
higher than that of the control strain LCTL1, were AAR and AEQ, both from the Mosaic beer
clade. While this is milder than some mutator phenotypes that have been artificially generated in
the lab, to our knowledge, no comparably high mutation rate has been previously reported in a
natural isolate of *S. cerevisiae*, with the exception of the spore derivatives of the incompatible
cMLH1-kPMS1 diploid natural isolate (Raghavan et al. 2018).

Figure 3. Haploid natural isolates exhibit a 10-fold range of mutation rate variation

Mutation rate variation measured among haploid natural isolates using our *CAN1* reporter gene Luria-Delbrück fluctuation assays. Strains are shown ordered by their mean mutation rates. Mutation rates for each strain were estimated using at least two replicates, each estimate represented here by a dot. A standard boxplot spans the interquartile confidence interval of possible mutation rates for each strain.

CAN1 sequencing reveals *de novo* mutation spectrum differences

We identified a total of 5571 *CAN1* mutations across all strains, including 4561 point mutations, 837 indels (Supplementary Table S4), and 173 multinucleotide mutations (MNMs) (Supplementary Table S5-S6). 90% of the observed indels are single base-pair indels (754 out of 837), and for simplicity we included only single base-pair indels along with point mutations when reporting each strain's mutation spectrum.

Two of the 4561 point mutations occurred at strain-specific non-reference sites. The remaining 4559 mutations consisted of repeated observations of only 727 unique mutations at 476 positions in *CAN1*. Given that each mutation was observed an average of 6.2 times, our dataset likely contains a large fraction of the mutations that are able to knock out *CAN1*'s functionality. While 23% of the mutations are observed only once, there are several hotspots that are mutated many times in our dataset (Supplementary Figure S7 and S8, Supplementary Table S7). For example, genomic location 32399 on chromosome V was hit more than 100 times across different genetic backgrounds. 85 of these hits created the same G>T mutation, while 19 created another G>C mutation. Across all mutations counted with multiplicity, we observed 2676 missense mutations, 1866 nonsense mutations and only 17 synonymous mutations. These synonymous mutations made up less than 0.37% of the total point mutations observed. Since synonymous mutations are considered *a priori* unlikely to cause *CAN1* to lose functionality, these variants are likely sequencing errors or hitchhikers that occurred in cells containing other inactivating mutations, or possibly rare examples of synonymous mutations disrupting translation (Arthur et al. 2015) in *CAN1*. This low synonymous mutation rate further demonstrates the accuracy of our pipeline.

We were able to identify MNMs, complex mutation events that create multiple nearby substitutions or indels at once (Stone et al. 2012; K. Harris and Nielsen 2014; Averof et al. 2000; Schrider, Hourmozdi, and Hahn 2011), by looking for the presence of multiple mutations on the same Illumina read (Materials and Methods). We estimate that MNMs generate 3.1% of all point mutations, similar to the 2.6% previously reported in a single strain background (Lang and Murray 2008). Most of our strains have similar ratios of MNMs to single base-pair mutations, but there exist a few outliers (Figure 4A). For example, strain AAB has disproportionately many MNMs (Figure 4A and Supplementary Figure S9) while AAR and AEQ have strikingly few MNMs.

To our knowledge, the largest previous *CAN1* fluctuation assay in *S. cerevisiae* observed point mutations at 102 distinct positions (Lang and Murray 2008). We observed mutations at 100 of these sites as well as 376 additional sites not previously known to abrogate *CAN1* function. Among the two mutations observed by Lang and Murray that are missing from our dataset, one

is located near the end of the *CAN1* amplicon in a region we exclude due to insufficient
sequencing coverage in most strains. The other site is the location of a mutation changing the
anticodon "CTA" to "TTA," which is synonymous and thus not likely to have affected *CAN1*
function.

One possible contributor to the observed differences in mutation rates and spectra are
differences in target size: the number of distinct *CAN1* mutations that are able to help the strain
survive on canavanine media. However, we were able to rule this out as a significant contributor
by comparing the ratio of observed missense to nonsense mutations among strains, which was
previously used by Lang and Murray (2008) to estimate mutational target size. We estimate that
the mutational target size varies at most two-fold among strains, from 145 bp to 299 bp
(Supplementary Table S8). Lang and Murray (2008) previously estimated a target size of 163
bp, falling within our range. This range is too narrow to explain the 10-fold range of mutation
rate variation we observe among strains, but it is large enough that we cannot confidently
translate our *CAN1*-based mutation rate estimates into genome-wide estimates of the mutation
rate per base pair per generation.

We performed hypergeometric tests to determine whether the mutation spectra we measured
from the two control lab strains LCTL1 and LCTL2 were distinct from those measured from
haploid natural isolates and from the spectrum measured from the same LCTL2 strain by Lang
et al. (2008) (Materials and Methods). We found the spectra of point mutations we measured
from the lab strains LCTL1 and LCTL2 to be statistically indistinguishable from the spectra
Lang, et al. (2008) obtained using Sanger sequencing of canavanine-resistant mutants (p = 0.82
for LCTL1 and p = 0.087 for LCTL2). With indels included, our LCTL1 spectrum appears
significantly different from that of Lang et al. (2008) (p = 0.0012, Bonferroni corrected p-value:
0.042), but the LCTL2 spectra remain indistinguishable with indels included (p = 0.0054,
Bonferroni corrected p-value: 0.189).

Figure 4. *de novo* mutation rate and spectra in natural isolates

A. Single nucleotide mutation rates plotted against MNM rates across strains. These rates were calculated by multiplying the mean mutation rate estimated using *CAN1* by the proportion of mutations in each strain measured to be either single-nucleotide mutations or MNMs. Here, single nucleotide mutations include both single base pair substitutions and indels. **B**. Mutation spectra in AEQ and AAR show significant enrichment of C>A mutations compared to the control lab strain LCTL1. Only single base-pair indels were used to generate these counts. **C**. A PCA of the same strains' *de novo* mutation spectra compared to the mutation spectrum reported in (Lang and Murray 2008).

The lab strain LCTL1 appears to have a mutation spectrum that is representative of most natural isolates (Figure 4C, Supplementary Figure S10-S11). Using Bonferroni corrected *p*-values to determine significance, we found the strains AAA, ACS, AGM, AHH, AHC, AEF, ADF, ACM, AGR, AAI, and AAB to have mutation spectra that are statistically indistinguishable from that of LCTL1. We found that CEV and AFH are distinguished only by their high proportions of insertions. ADN showed significant but subtle divergence from LCTL1 in the spectrum of single nucleotide variants (Supplementary Figure S10-S11).

A natural mutator phenotype with a distinctive mutation spectrum

We identified two strains with mutation spectra that diverged strikingly from the reference LCTL1 spectrum: AEQ (*p* < 1e-4) and AAR (*p* < 1e-4), the two strains that also have 10-fold higher mutation rates than LCTL1. Both strains appear highly enriched for C>A mutations compared to LCTL1 (Figure 4B). The strain AAR's mutation rate estimates appear somewhat bimodal (Figure 3), but C>A mutations are consistently enriched in replicate pools with both lower and higher estimated mutation rates. The main spectrum difference between the two mutation rate modes appears to be a small difference in the C>G mutation proportion (Supplementary Figure S12).

In both AEQ and AAR, the proportion of C>A mutations was measured to be elevated nearly 3-fold above the proportion of C>A mutations in LCTL1 and similar strains (Figure 4B). Remarkably, this C>A enrichment appears sufficient to explain the placement of AEQ and AAR as rare variant mutation spectrum outliers that we previously saw in Figure 1B, which was computed from polymorphisms sampled genome-wide, not just within *CAN1*. Both strains have rare variant spectra that are displaced from the population norm along a principal component vector pointing in the direction of increased C>A enrichment. These strains' high mutation rate, C>A-heavy *de novo* mutation spectrum, and concordant C>A-heavy rare variant spectrum all point to the conclusion that these Mosaic beer strains display a naturally-occurring genetically encoded mutator phenotype.

To assess whether the C>A enrichment phenotype observed in AAR and AEQ is likely shared with any of the 1011 strains for which we lack mutation spectrum measurements, we ranked all 1011 strains (excluding close relatives) according to the C>A enrichment of their rare variants. We found that SACE_YAG and BRM, the two strains closest to AAR and AEQ in the global neighbor-joining phylogeny, clustered with AAR and AEQ in having high C>A fractions within the top 5% in the dataset. The rest of the top-ranking 5% of the strains exhibit some phylogenetic clustering, but no others fall within the Mosaic beer clade (Figure 5A). Instead, they are somewhat dispersed across two large, diverse clades known as "Mosaic region 3" and the "Mixed origin" clade. We also used a bootstrapping method to find strains with enriched C>A fractions, using an empirical *p*-value threshold of 0.05. Many of the same strains are outliers in both tests, including the four Mosaic beer strains (Supplementary Figure S13). The phylogenetic clustering of C>A rare variant enrichment suggests that multiple clades may be genetically predisposed toward accumulating relatively higher rates of this mutation type.

Figure 5. Enrichment of C>A mutations in rare natural variants

A. Phylogeny of the 1011 collection with strains in the top 5% of C>A fraction shown in red. **B.** Phylogeny of AEQ, AAR, and closely related Mosaic beer strains **C.** Top panel: The C>A ratio as a function of minor allele count in Mosaic beer strains that are closely related to AEQ and AAR, including AEQ's and AAR's the diploid common ancestor CBS 1782. C>A ratios in polymorphisms are calculated across allele count (AC) bins with cutoffs of 2, 4, 6, and 8. When computing allele counts, closely-related strains are excluded, and each strain is represented as a diploid in genotype. Bottom panel: total number of variants in each AC bin.

Focusing on the small clade containing AAR, AEQ, BRM, and SACE_YAG with all members showing C>A enrichment (Figure 5B), we found that all but BRM are haploid derivatives of the diploid *Saccharomyces cerevisiae var diastaticus* strain CBS 1782, which was isolated in 1952 from super-attenuated beer (Andrews and Gilliland 1952). AEQ and AAR differ at roughly 14,000 variant sites (the median pairwise genetic distance in the 1011 strains is 64,000) and SACE_YAG differs from AEQ and AAR at about 11,000 sites each. These differences among

haploid derivatives reflect the high level of heterozygosity in the parental diploid strain. The
fourth strain with an elevated C>A mutation fraction, BRM, is derived from an independent
source: it was isolated in 1988 from a cassava flour factory in Brazil (Laluce et al. 1988).
Despite its distinct origin, BRM differs at only 14,000-17,000 sites from each of the
aforementioned three strains.

In AAR, AEQ, BRM, and SACE_YAG, rarer variants are notably C>A-enriched, but higher
frequency variants exhibit weaker C>A enrichment that declines with increasing allele frequency
such that variants of allele count greater than eight have C>A fractions more typical of other
strains. We verified that the same pattern holds for variants present in the whole genome of the
ancestral diploid isolate CBS 1782, which we sequenced after obtaining a sample from the
NCYC database (see Materials and Methods). This suggests that AAR and AEQ indeed
inherited their mutator phenotype from their wild ancestors and that this phenotype is at least
partially penetrant in the diploid state. In contrast, the four closest outgroups to AAR, AEQ,
BRM, and SACE_YAG in the 1011 yeast genomes phylogeny exhibit a consistently lower C>A
fraction that does not vary with allele frequency (Figure 5B and Supplementary Figure S14). The
concordant enrichment of C>A mutations in rare polymorphisms and *de novo* mutations from
the same strains suggests that this C>A enrichment is genetically determined and is not specific
to the *CAN1* locus but has affected the entire genome during the recent history of this clade.

A scan for candidate mutator alleles

To explore genetic variation that could underlie the C>A enrichment phenotype observed in
AEQ and AAR, we scanned for nonsense and missense mutations in a list of 158 candidate
genes known to play roles in DNA replication and repair, including genes that were previously
identified to harbor mutator alleles through genetic screens (Supplementary Table S9) (Boiteux
and Jinks-Robertson 2013; Stirling et al. 2014). No candidate premature stop codons were
found to be both present in AAR and AEQ and rare (MAF<0.05) in the 1011 yeast genomes
overall. However, we identified 40 sites with at least one rare non-synonymous allele
(MAF<0.05) shared by AEQ and AAR and absent from the other haploid strains that we
experimentally found to have normal mutation spectra (Supplementary Table S10).

One of these missense variants was observed within *OGG1*, a glycosylase involved in the
oxidative stress response that excises the guanine lesion 8-oxo-G. *ogg1* null mutants are known
to have high C>A mutation rates (Shockley et al. 2013) and mutation rates that are ten-fold
elevated above normal (Ni, Marsischky, and Kolodner 1999). The *OGG1* missense variant
present in AAR and AEQ is homozygous in the parental diploid CBS 1782; the same variant is
also shared by two other Mosaic beer strains, AQH and AAQ, which are not enriched for C>A
rare variants. A close examination of rare variant mutation spectra revealed that AQH may have
a C>A enrichment phenotype that is masked by a high rate of C>T mutations (Supplementary
Figure S15), but no evidence of C>A enrichment was found in AAQ.

Testing the candidate mutator allele *OGG1*$^{AEQ/AAR}$

We sought to experimentally test whether the *OGG1* allele present in AEQ and AAR, hereafter called *OGG1*$^{AEQ/AAR}$, contributes to the AEQ/AAR mutator phenotype. We accomplished this by expressing the *OGG1*$^{AEQ/AAR}$ allele in a lab strain with its endogenous *ogg1* allele deleted. If *OGG1*$^{AEQ/AAR}$ contributes to the mutator phenotype of AEQ and AAR, then this gene variant should be less effective at rescuing the *ogg1* deletion phenotype than the wild-type *OGG1* allele.

To test this hypothesis, we constructed two CEN plasmids. One plasmid was constructed to express *OGG1*$^{AEQ/AAR}$; the other was constructed to express *OGG1*^{wt}, with the same sequence as the *OGG1* allele present in the S288C reference. Through Sanger sequencing, we found that *OGG1*$^{AEQ/AAR}$ in fact contains two single nucleotide differences from the reference sequence (Figure 6A). One of these variants (an A>G substitution) lies 156 bp upstream of *OGG1*, presumably in the gene's promoter region, which we previously missed when examining the coding sequence. The other variant is a nonsynonymous change from C to T at position 800, which changes alanine to leucine. While the nonsynonymous mutation in *OGG1* is present in only 7 strains from the Mosaic beer clade (including AEA which is a close relative of BRM) from the 1011 collection, the promoter mutation is present in 57 strains. Since the promoter mutation is more widespread and not linked to the nonsynonymous mutation, we hypothesize that it is less likely to cause the mutator phenotype. To avoid truncating any important regulatory sequences, we used PCR to amplify two fragments with different lengths of flanking sequence from each endogenous *OGG1* locus: one shorter fragment called *OGG1-A,* (1.7kb) and one longer fragment called *OGG1-B* (2.2kb), creating a total of four plasmids. We then used these *OGG1* plasmids, as well as an empty plasmid, to transform haploid *S. cerevisiae* with an *ogg1* deletion derived from the heterozygous diploid deletion collection (SGA) (Tong and Boone 2007).

We used fluctuation assays to estimate the mutation rates of *ogg1* deletion strains complemented with these different *OGG1* plasmids. We found that the mutation rate of the *ogg1* null yeast with an empty control plasmid was elevated 4-fold compared to the mutation rate of the strain transformed with *OGG1*^{wt}. This difference is smaller than the ten-fold mutation rate difference previously measured between an *ogg1* null mutant and a strain with an intact endogenous *OGG1* allele (Ni, Marsischky, and Kolodner 1999), possibly reflecting a difference in the efficiency of endogenous *OGG1* expression versus expression of the same allele from a plasmid. We found that the strain transformed with *OGG1*$^{AEQ/AAR}$ had a 2.4-fold higher mutation rate than the strain transformed with *OGG1*^{wt}, suggesting that the AEQ/AAR allele is less effective at complementing the DNA repair deficiency of the *ogg1* deletion strain. This supports our hypothesis that the *OGG1*$^{AEQ/AAR}$ allele is at least in part responsible for the high mutation rate of the AEQ and AAR strains. We measured similar mutation rates in strains transformed by the shorter and longer PCR fragments containing each *OGG1* allele, suggesting that the additional regulatory DNA present in the longer sequence did not substantially impact *OGG1* expression.

537 Figure 6. Mutation rates of a ogg1 deletion strain complemented by *OGG1*[AEQ/AAR] *versus OGG1*[wt]

538 **A**. Locations of the genetic variants differentiating the *OGG1*[wt] and *OGG1*[AEQ/AAR] alleles. Solid and dashed lines

539 denote the boundaries of *OGG1_A* and *OGG1_B* (not scale)*. OGG1_A* contains a 1.7kb PCR fragment including

540 *OGG1*, while OGG1_B contains a 2.2kb PCR fragment. **B**. Mutation rate estimates of *ogg1* deletion strains

541 transformed by plasmids containing different OGG1 alleles.

Discussion

In this study, we compare the mutation spectra of natural variants and *de novo* mutations within

diverse strains of *S. cerevisiae.* We find both population genetic and experimental evidence

pointing to a recent increase in the rate of C>A mutations within one clade as a result of a

naturally occurring genetically encoded mutator phenotype.

Our highly accurate pipeline uses a classical fluctuation assay to quantify *de novo* mutation rate

in the reporter gene *CAN1,* then utilizes sequencing of pooled mutants to measure mutation

spectra. Sanger validations confirmed that this approach has an undetectably low false positive

rate and a very low false negative rate. The reliability of our pipeline is further supported by

concordance between our mutation spectrum measurements and previous measurements

made from the same lab strains (Lang and Murray 2008). Using this method, we identified two

extreme mutation spectrum outliers—the Mosaic beer strains AEQ and AAR—as well as several

strains with more subtle mutation spectrum phenotypes that are good candidates for future

investigation.

Although the mutation spectrum variation from natural polymorphisms we report here is

reminiscent of observed mutation spectrum variation among humans, great apes, and

laboratory mouse strains, we note that *S. cerevisiae* mutation spectra separate less by

population compared to vertebrate mutation spectra. While it is possible to infer a human

genome's continental group of origin using its mutation spectrum alone, the same is not

generally true in *S. cerevisiae*. Several factors, which are not mutually exclusive, might underlie
this difference. One is the presence of complex population structure and pervasive gene flow
between *S. cerevisiae* clades (Schacherer et al. 2009; Peter et al. 2018; Liti et al. 2009).
Another factor is the small size of the *S. cerevisiae* genome; each strain has two orders of
magnitude fewer derived alleles than most vertebrate genomes have. This limits our ability to
detect how flanking base pairs affect the mutation rate of each site in the genome and
constrains us to study a lower-dimensional mutation spectrum than the context-dependent ones
studied in species with larger genomes.

While gene flow and genome size might be responsible for the relatively modest magnitude of
mutation spectrum divergence between most strains of *S. cerevisiae*, it is also possible that
DNA replication and repair are intrinsically more uniform in *S. cerevisiae* than in vertebrates,
perhaps because of the greater efficiency of selection against weakly deleterious mutator alleles
in a unicellular organism that exists at large effective population sizes and often reproduces
asexually. Asexual reproduction should theoretically increase the efficiency of selection against
mutator alleles because deleterious variants created by the mutator cannot recombine onto
other genetic backgrounds; on the other hand, it can also limit the efficiency of selection against
individual deleterious mutations by permanently tethering them to particular genetic
backgrounds. Further measurements of mutation spectrum variation within other species will be
needed to determine whether the stability observed here is indeed characteristic of unicellular
eukaryotes. If mutation spectra tend to be stable within species that have low mutation rates
and strong selection against mutation rate modifiers, we might expect to see even less mutation
spectrum variation among populations of ciliates like *Paramecium* and *Tetrahymena*, whose
mutation rates are substantially lower than that of *S. cerevisiae* (Sung, Tucker, et al. 2012)*.

Many questions about mutation spectrum variation within *S. cerevisiae* and other species
remain unresolved and present important avenues for future work. For example, it is unclear
whether rare polymorphisms in the Mixed origin and Mosaic region 3 clades are enriched for
C>A mutations due to the same genetic mechanisms active in AEQ and AAR. Other genes
might underlie the mutation spectrum differences observed among other strains, though our
analyses suggest that some mutation spectrum gradients that dominate the common variation
PCA are unlikely to be explained by extant mutators. One such gradient is the A>G enrichment
in the African beer yeast clade, which is less pronounced in our rare variant PCA (Figure 1B)
compared to our PCA extracted from variation of all frequencies (Figure 1A). This is somewhat
reminiscent of the frequency distribution of the TCC>TTC mutation "pulse" that distinguishes
Europeans and South Asians from other human populations, and may suggest that the African
beer A>G enrichment was caused by an extinct mutator allele or a mutagen found in a past
environment.

Natural selection might contribute to the mutation spectrum variation within *S. cerevisiae* if
certain mutation types are more often beneficial than others and if such asymmetries vary
between populations. One example of how natural selection can affect mutation spectrum
variation is that transitions are more often synonymous than transversions are (Freeland and

Hurst 1998), leading to more frequent selection against transversions in genic regions.
However, we note that most of the gradient structure observed in our analyses can be
reproduced with synonymous mutations alone, meaning that selection is unlikely to explain
much of the natural yeast mutation spectrum variation we observe.

The C>A mutations enriched in AEQ, AAR, and their relatives might be a signature of oxidative
stress damage; such mutations are a known signature of failure to repair 8-oxoguanine lesions,
which is consistent with a causal role for these strains' missense substitution in the oxidative
stress response gene *OGG1* as verified through our *OGG1* plasmid assay. A recent study
knocked out *OGG1* in human cells and found that this elevated the rate of a C>A-dominated
mutational signature known in the Cosmic catalog as SBS18 (Zou et al. 2020). SBS18 was
initially identified in tumors from individuals with pathogenic variation in *Mutyh* (Viel et al. 2017),
a direct interacting partner of *OGG1* in the 8-oxoguanine repair pathway that has also been
implicated in a C>A-dominated germline mutational signature in mice (Sasani et al. 2021).

Although our *OGG1* plasmid assay provides compelling evidence that the 8-oxoguanine repair
pathway plays a role in the AAR/AEQ mutator phenotype, we note that C>A mutations do not
comprise all of the excess mutations measured in AEQ and AAR. Compared to LCTL1, the ratio
of C>A mutations to C>T mutations is elevated roughly 3-fold in AEQ and AAR, which is less
than the 10-fold overall elevation of the mutation rate in these strains. These mutation data
imply that all mutation types have higher rates in AEQ and AAR compared to other *S. cerevisiae*
strains, not just C>A. Further work will be required to verify whether the C>A enrichment and
broad-spectrum mutation rate increase are driven by the same biochemical mechanism.

Although our estimates of *de novo* mutation spectra are all based on the *CAN1* locus, we note
that *CAN1* mutation spectra are generally similar to the genome-wide spectra obtained from MA
experiments (Zhu et al. 2014; Sharp et al. 2018), except that *CAN1* spectra have more
mutations at C/G base pairs and relatively fewer mutations at A/T base pairs (Supplementary
Figure S16). The 2676 and 1866 missense and nonsense mutations in our dataset contain
numerous instances of all six mutation types that comprise our summary mutation spectrum,
suggesting that *CAN1* contains many opportunities to ascertain the full spectrum observed
genomewide. We noticed that the *CAN1* spectra we measured from the LCTL2 strain showed a
slightly higher proportion of C>A mutations than that measured from the same strain by (Lang
and Murray 2008) also using a *CAN1* reporter. This could be due to the fact that Lang et al.
used ten times the concentration of canavanine that we used, which might mean that some
mutants could have grown after plating on our canavanine media (Shor, Fox, and Broach 2013).
The fact that we do not see elevated C>A mutations in LCTL1 suggests that this could be a
LCTL2 strain-specific behavior.

In summary, the results presented in this paper provide some of the most direct evidence to
date that eukaryotic mutation spectra are variable within species (K. Harris 2015; K. Harris and
Pritchard 2017; Dumont 2019; Goldberg and Harris 2021). It has been proposed that the best
explanation for such mutation spectrum heterogeneity is the frequent emergence of nearly

neutral mutator alleles that turn over rapidly as a consequence of weak purifying selection on the mutation rate. Our *de novo* mutation spectrum measurements provide experimental verification of this claim, showing that at least one mutational signature whose activity varies among natural yeast strains is likely caused by an extant mutator allele.

Although our results show that the mutation spectrum bias shared by certain Mosaic beer yeast is genetically encoded, it is worth noting that this C>A gradient is not the principal axis of mutation spectrum variation in the 1011 yeast genomes that we computed from all variants (Figure 1A). It remains to be seen how many other mutator or antimutator alleles might exist within this strain collection and to what extent they can explain the mutation spectrum variation observed among strains from different environments. A broader question still is whether the forces that created *S. cerevisiae*'s mutation spectrum variation are similar to the forces that shaped the distinctive mutation spectra of different human populations and great ape species. If we can identify the genes that underlie natural yeast mutator phenotypes such as the one described in this study, it will likely be more straightforward to test these genes for mutator activity in humans and other species than to discover mutator alleles via any kind of agnostic genome scan.

Acknowledgements

We dedicate this paper to the memory of Dr. Alan J. Herr, a sorely missed colleague, mentor, friend and inspiring role model. We would like to thank all members of the Harris and Dunham labs for helpful comments and discussions. We thank Nathaniel Sharp and Greg Lang for sharing strains. We also thank Joseph Schacherer for sharing the 1011 strain collection with the Dunham lab. P.J was supported by a Burroughs Wellcome Fund Career Award at the Scientific Interface awarded to K.H. K.H. acknowledges additional support from a Searle Scholarship, a Sloan Research Fellowship, a Pew Biomedical Scholarship, and National Institute of General Medical Sciences Grant 1R35GM133428-01. A.J.H was supported by the National Institute for General Medical Sciences (NIH/NIGMS R01GM118854). A.R.O. was supported by the National Human Genome Research Institute of the NIH under award T32 HG00035. The research of M.J.D. was supported by NIH/NIGMS award P41 GM103533 and a Faculty Scholar grant from the Howard Hughes Medical Institute. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Burroughs Wellcome Fund, the Kinship Foundation, the Sloan Foundation, the Pew Charitable Trust, HHMI, the NIH, or NIGMS.

Materials and Methods

Variant filtering and mutation PCA analysis

We filtered the original variants from the 1011 *S. cerevisiae* collection (Peter et al. 2018) by including biallelic SNPs with less than 20% missing genotypes. We restricted to regions in the genomes where reads can be uniquely mapped ("mregions_100_annot_2011.bed" from (Jubin et al. 2014)) and excluded repeat-masked regions. Closely related strains (pairwise genetic

distance less than 8000) are excluded from the 1011 dataset. Singletons were excluded when
counting individual mutations in Figure 1 to minimize the impact of sequencing errors. Strains
with extensive introgression from *S. paradoxus* (clade 2, 9, 10 from (Peter et al. 2018)) were
excluded in order to minimize bias from errors in the inference of ancestral and derived alleles.
Ancestral states of mutations were inferred using five *S. paradoxus* sequences (Yue et al.
2017), aligned to the *S. cerevisiae* reference genome R64-1-1 using lastz v1.04.00 (R. S. Harris
2007). Only sites that are fixed in four out of five strains were inferred to be the ancestral alleles,
and other sites were ignored. When computing the mutation spectra of strains from variants for
Figure 1A, each individual strain was assumed to be diploid, with homozygous derived alleles
counted with twice the weight as heterozygous derived alleles. When counting rare variants,
homozygous derived alleles were given the same weights as heterozygous derived alleles.

To further minimize confounding of the mutation spectrum by ancestral allele misidentification,
only variants with derived allele frequency less than 0.5 were used. Variants that passed all
filtering criteria were used to compute a normalized mutation spectrum histogram for each
individual strain. When performing PCAs, no more than 30 strains from each population were
randomly sampled to minimize bias from uneven sampling. The same strains were used to
generate PCA plots in Supplementary Figures S1-S4 (Supplementary Table S1), except that
strains with fewer than 8 singletons or rare variants were further excluded when generating
Supplementary Figure S4 and Figure 1B. Our definition of singletons varied as a function of
ploidy (Supplementary Figure S4-S5): In haploids and homozygous diploids (as defined in
(Peter et al. 2018)), a singleton will be fixed in the strain where it occurs (represented as
homozygous), but in other types of strains, a singleton is required to be heterozygous. In all
cases, a singleton is a variant present in only a single strain.

Fluctuation assays and sequencing

We performed fluctuation assays according to an established protocol (Lang 2018) with the
following modifications: 4μl of overnight inoculant was diluted in 40ml SC-Arginine+2% Glucose
media. 50μl of the diluted cultures were distributed in 96-well round-bottom plates (Costar 3788)
for each strain. Plates were sealed with Breathe-Easy sealing membrane (Sigma Z380059).
SC-Arginine-Serine+Canavanine (60 mg/liter L-canavanine) Omni plates (Nunc OmniTray
242811) were used and dried for 2-4 days in a 30℃ incubator before using. Depending on the
strains, 50μl of culture were diluted one- to four-fold when plating on the Omni plates, either to
reduce the background or to avoid growth of too many mutant colonies. After plating, the plates
were dried and then incubated at 30℃ for 48 hours. Independent mutants from separate
cultures were inoculated into 200μl SC-Arginine-Serine+60mg/Liter Canavanine+2% Glucose
media, and then grown to saturation over ~43 hours at 30℃ with shaking. Optical densities
(ODs) were measured after incubation, and only mutants that reached similar saturation ODs
were pooled (150ul each) to achieve equal proportions. Genomic DNA from each pool was
extracted using the Hoffman Winston protocol (Hoffman and Winston 1987). *CAN1* was then
PCR amplified using published primers (Lang and Murray 2008) with 15 cycles. Two
independent 25μl PCR reactions were then pooled and cleaned up with a Zymo Clean &

Concentrator Kit (D4004). Sequencing libraries were prepared using the Nextera XT DNA
Library Preparation Kit with customized indices. Sequencing runs with 75 or 150 bp paired-end
reads were performed using an Illumina NextSeq 550 sequencer (BioProject: PRJNA691686).

Calculation of mutation rates

The rSalvador package (Zheng 2017) was used to estimate the number of mutation events (m)
in each fluctuation assay using maximum likelihood under the Lea-Coulson model (Lea and
Coulson 1949; Luria and Delbrück 1943; Ma, G. Vh. Sandri, and Sarkar 1992). The total number
of cells (Nt) was measured by counting colonies seeded with dilutions of cells on YPD plates
with dilutions ranging from 1:10,000 to 1:40,000. The rate of loss-of-function mutations per
CAN1 gene per cell division was estimated to be m/Nt.

Mutation calling

Sequencing reads were first mapped using bowtie v2.2.3 (Langmead and Salzberg 2012) to the
Scer3 S288C reference *CAN1* PCR fragment sequence using primers designed by Lang et al
(2008). Mutation coordinates were therefore called relative to the start of the *CAN1* amplicon.
Adapters were trimmed using the program trim_galore v0.6.6 (Krueger n.d.) and paired-end
reads were merged using pear v0.9.11 (Zhang et al. 2014). The command `fastq_quality_filter
-q 20 -p 94` was used to remove low quality reads before running bowtie. A MAPQ cutoff of 40
was used for SNPs and a cutoff of 20 was used for indels. Pysamstats v1.1.2 (Miles n.d.) was
used to compute the frequencies of all possible alleles at each base pair. Sites with read depth
less than 200 or with less than 40% coverage of the amplicon were excluded. After the first
round of mapping, sites that were fixed in each strain were called and compared to the SNPs in
the 1011 collection to confirm strain identity. We then performed a second round of read
mapping using the same pipeline except that each strain's reads were mapped to a
strain-specific *CAN1* reference sequence.

For each sequencing pool, we let N be the number of mutants that were pooled prior to
sequencing. Non-reference alleles with frequencies between $0.65 \times 1/N$ and 0.95 were included
as evidence of mutations, discarding alleles below this frequency range as likely to be
sequencing errors and alleles above this frequency range as likely to be strain-specific SNPs.
Adjacent indels were merged if their frequencies differed by less than 10%). MNMs were
identified in each pool by first flagging pairs of mutations occurring at similar frequencies (plus
or minus 9%) within 10bp of one another and then verifying the coexistence of the two
mutations on at least 70% of the paired-end reads where at least one of the two mutations
appears. Complex MNMs containing three or more variants were identified by merging MNMs
that share a SNP in common. To obtain single nucleotide mutation counts and indel counts,
mutations that are part of MNMs were first excluded from each pool. The coordinates of each
mutation were converted back from *CAN1*-specific coordinates to genomic positions. Point
mutations were further annotated using VEP (McLaren et al. 2016) to further categorize into
missense, nonsense or synonymous mutation types.

Allele frequencies were used to estimate the multiplicity of each mutant as follows: First, the mean and standard deviation of all mutant allele frequencies were calculated from each pool. Each allele frequency more than two standard deviations above the mean was then translated into a mutation count by dividing it by the mean allele frequency and then rounding to the nearest integer. Mutations with frequencies less than two standard deviations above the mean are assumed to be mutations with count 1.

Statistically quantifying mutation spectrum differentiation

To compare the mutation spectra between strains, mutations were first classified as one of the 6 general classes of base-substitutions (A>C, A>G, A>T, C>A, C>G, C>T) or as single base-pair insertions or deletions. We then compared the mutation spectra of the two control strains LCTL1 and LCTL2 to all other haploid isolates as well as one spectrum published by Lang et al (2008) (a total of 35 tests) using a pairwise hypergeometric test (Adams and Skopek 1987), a custom python script (Tracy et al. 2020). In the first round of this test, the paired mutation counts were arranged in a 2 x 8 contingency table. To test the null hypothesis that the two mutation spectra are the same, the hypergeometric probability of the observed table was calculated and compared to the hypergeometric probabilities of 10,000 random tables with the same row and column totals. The number of random tables with a higher hypergeometric probability than the observed provides an estimate of the p-value. We used the conservative Bonferroni correction to compute the significance cutoff (0.05/35=0.001429). A second set of Bonferroni-corrected p-values was calculated after excluding indels to form a 2 x 6 contingency table. These p-values were used to determine how many of the significant mutation spectrum differences were driven by the indel category (Supplementary Figure S8, S9).

Whole-genome Sequencing and variant calling of CBS 1782
A sample of CBS 1782 was obtained from the National Collection of Yeast Cultures (NCYC 361) January of 2021. Genomic library was prepared from the Illumina Nextera DNA Flex Library Prep kit (Illumina 20018704). Sequencing runs with 150 bp paired-end reads were performed using an Illumina NextSeq 550 sequencer to roughly 250x coverage. Sequencing reads were mapped using BWA mem v 0.7.17 (Li and Durbin 2009). Only reads that are uniquely mapped with MQ>20 were used. GATK4 HaplotypeCaller was used for variant calling (Poplin et al. 2018). Variant sites (+/- 2bp) that overlap with deletion were filtered out. The variants for CBS 1782 were then merged with the raw vcf file for the 1011 collection excluding closely-related strains, and underwent the same filtering as for the PCA analysis.

Constructing *OGG1* plasmids
Haploid yeast with both *ogg1* and *leu2* deleted was generated from spores dissected from the heterozygous diploid strain #20510 of the SGA collection (Tong and Boone 2007). CEN plasmids were generated using pRS415 with *LEU2* as the selection marker (Chee and Haase 2012). *OGG1-A* and *OGG1-B* were amplified by PCR from strain AAR for *OGG1*$^{AEQ/AAR}$ or from LCTL1 for *OGG1*^{wt}. Primers for *OGG1-A:* CGATAGTTTGGCGTGCGATA, CGCCTTGGTGACCGTTTT. Primers for *OGG1-B:*

794 GGTTCTTCCCAATCATCCGA,AGGGCTTATTGACGACGACA. pRS415 were linearized by
795 HindIII and BamHI, followed by Gibson assembly with the *OGG1* alleles.

References

Acosta, Sharlene, Miguelina Carela, Aurian Garcia-Gonzalez, Mariela Gines, Luis Vicens,
Ricardo Cruet, and Steven E. Massey. 2015. "DNA Repair Is Associated with Information
Content in Bacteria, Archaea, and DNA Viruses." *The Journal of Heredity* 106 (5): 644–59.
Adams, W. T., and T. R. Skopek. 1987. "Statistical Test for the Comparison of Samples from
Mutational Spectra." *Journal of Molecular Biology* 194 (3): 391–96.
Agier, Nicolas, and Gilles Fischer. 2012. "The Mutational Profile of the Yeast Genome Is Shaped
by Replication." *Molecular Biology and Evolution* 29 (3): 905–13.
Alexandrov, Ludmil B., Serena Nik-Zainal, David C. Wedge, Peter J. Campbell, and Michael R.
Stratton. 2013. "Deciphering Signatures of Mutational Processes Operative in Human
Cancer." *Cell Reports* 3 (1): 246–59.
André, Jean-Baptiste, and Bernard Godelle. 2006. "The Evolution of Mutation Rate in Finite
Asexual Populations." *Genetics* 172 (1): 611–26.
Andrews, By J., and R. B. Gilliland. 1952. "SUPER-ATTENUATION OF BEER: A STUDY OF
THREE ORGANISMS CAPABLE OF CAUSING ABNORMAL ATTENUATIONS." *Journal of*
the Institute of Brewing. Institute of Brewing 58 (3): 189–96.
Antonarakis, Stylianos E., and Jacques S. Beckmann. 2006. "Mendelian Disorders Deserve
More Attention." *Nature Reviews. Genetics* 7 (4): 277–82.
Argueso, Juan Lucas, Amanda Wraith Kijas, Sumeet Sarin, Julie Heck, Marc Waase, and Eric
Alani. 2003. "Systematic Mutagenesis of the Saccharomyces Cerevisiae MLH1 Gene
Reveals Distinct Roles for Mlh1p in Meiotic Crossing over and in Vegetative and Meiotic
Mismatch Repair." *Molecular and Cellular Biology* 23 (3): 873–86.
Arthur, Laura, Slavica Pavlovic-Djuranovic, Kristin Smith-Koutmou, Rachel Green, Pawel
Szczesny, and Sergej Djuranovic. 2015. "Translational Control by Lysine-Encoding A-Rich
Sequences." *Science Advances* 1 (6). https://doi.org/10.1126/sciadv.1500154.
Averof, M., A. Rokas, K. H. Wolfe, and P. M. Sharp. 2000. "Evidence for a High Frequency of
Simultaneous Double-Nucleotide Substitutions." *Science* 287 (5456): 1283–86.
Beckman, R. A., and L. A. Loeb. 1993. "Multi-Stage Proofreading in DNA Replication." *Quarterly*
Reviews of Biophysics 26 (3): 225–331.
Belfield, Eric J., Carly Brown, Zhong Jie Ding, Lottie Chapman, Mengqian Luo, Eleanor Hinde,
Sam W. van Es, et al. 2021. "Thermal Stress Accelerates Arabidopsis Thaliana Mutation
Rate." *Genome Research* 31 (1): 40–50.
Boiteux, Serge, and Sue Jinks-Robertson. 2013. "DNA Repair Mechanisms and the Bypass of
DNA Damage in Saccharomyces Cerevisiae." *Genetics* 193 (4): 1025–64.
Bui, Duyen T., Anne Friedrich, Najla Al-Sweel, Gianni Liti, Joseph Schacherer, Charles F.
Aquadro, and Eric Alani. 2017. "Mismatch Repair Incompatibilities in Diverse Yeast
Populations." *Genetics* 205 (4): 1459–71.
Chao, Lin, and Edward C. Cox. 1983. "Competition between High and Low Mutating Strains of
Escherichia Coli." *Evolution; International Journal of Organic Evolution* 37 (1): 125–34.
Chee, Mark K., and Steven B. Haase. 2012. "New and Redesigned pRS Plasmid Shuttle
Vectors for Genetic Manipulation of Saccharomycescerevisiae." *G3* 2 (5): 515–26.
Couce, Alejandro, Larissa Viraphong Caudwell, Christoph Feinauer, Thomas Hindré, Jean-Paul
Feugeas, Martin Weigt, Richard E. Lenski, Dominique Schneider, and Olivier Tenaillon.

2017. "Mutator Genomes Decay, despite Sustained Fitness Gains, in a Long-Term Experiment with Bacteria." *Proceedings of the National Academy of Sciences of the United States of America* 114 (43): E9026–35.

Crow, J. F. 1997. "The High Spontaneous Mutation Rate: Is It a Health Risk?" *Proceedings of the National Academy of Sciences of the United States of America* 94 (16): 8380–86.

Drake, J. W. 1991. "A Constant Rate of Spontaneous Mutation in DNA-Based Microbes." *Proceedings of the National Academy of Sciences of the United States of America* 88 (16): 7160–64.

Dumont, Beth L. 2019. "Significant Strain Variation in the Mutation Spectra of Inbred Laboratory Mice." *Molecular Biology and Evolution* 36 (5): 865–74.

Dutta, Abhishek, Fabien Dutreux, and Joseph Schacherer. 2021. "Loss of Heterozygosity Results in Rapid but Variable Genome Homogenization across Yeast Genetic Backgrounds." *eLife* 10 (June). https://doi.org/10.7554/eLife.70339.

Eigen, M. 1971. "Selforganization of Matter and the Evolution of Biological Macromolecules." *Die Naturwissenschaften* 58 (10): 465–523.

Eisen, J. A., and P. C. Hanawalt. 1999. "A Phylogenomic Study of DNA Repair Genes, Proteins, and Processes." *Mutation Research* 435 (3): 171–213.

Farlow, Ashley, Hongan Long, Stéphanie Arnoux, Way Sung, Thomas G. Doak, Magnus Nordborg, and Michael Lynch. 2015. "The Spontaneous Mutation Rate in the Fission Yeast Schizosaccharomyces Pombe." *Genetics* 201 (2): 737–44.

Freeland, S. J., and L. D. Hurst. 1998. "The Genetic Code Is One in a Million." *Journal of Molecular Evolution* 47 (3): 238–48.

Goldberg, Michael E., and Kelley Harris. 2021. "Mutational Signatures of Replication Timing and Epigenetic Modification Persist through the Global Divergence of Mutation Spectra across the Great Ape Phylogeny." *Genome Biology and Evolution*, May. https://doi.org/10.1093/gbe/evab104.

Gompel, Nicolas, Benjamin Prud'homme, Patricia J. Wittkopp, Victoria A. Kassner, and Sean B. Carroll. 2005. "Chance Caught on the Wing: Cis-Regulatory Evolution and the Origin of Pigment Patterns in Drosophila." *Nature* 433 (7025): 481–87.

Good, Benjamin H., Michael J. McDonald, Jeffrey E. Barrick, Richard E. Lenski, and Michael M. Desai. 2017. "The Dynamics of Molecular Evolution over 60,000 Generations." *Nature* 551 (7678): 45–50.

Gou, Liangke, Joshua S. Bloom, and Leonid Kruglyak. 2019. "The Genetic Basis of Mutation Rate Variation in Yeast." *Genetics* 211 (2): 731–40.

Harris, Kelley. 2015. "Evidence for Recent, Population-Specific Evolution of the Human Mutation Rate." *Proceedings of the National Academy of Sciences of the United States of America* 112 (11): 3439–44.

Harris, Kelley, and Rasmus Nielsen. 2014. "Error-Prone Polymerase Activity Causes Multinucleotide Mutations in Humans." *Genome Research* 24 (9): 1445–54.

Harris, Kelley, and Jonathan K. Pritchard. 2017. "Rapid Evolution of the Human Mutation Spectrum." *eLife* 6 (April). https://doi.org/10.7554/eLife.24284.

Harris, Robert S. 2007. "Improved Pairwise Alignment of Genomic Dna." Phd, USA: Pennsylvania State University. https://dl.acm.org/doi/book/10.5555/1414852.

Heck, Julie Akiko, Juan Lucas Argueso, Zekeriyya Gemici, Richard Guy Reeves, Ann Bernard, Charles F. Aquadro, and Eric Alani. 2006. "Negative Epistasis between Natural Variants of the Saccharomyces Cerevisiae MLH1 and PMS1 Genes Results in a Defect in Mismatch Repair." *Proceedings of the National Academy of Sciences of the United States of America* 103 (9): 3256–61.

Herr, Alan J., Masanori Ogawa, Nicole A. Lawrence, Lindsey N. Williams, Julie M. Eggington,
Mallika Singh, Robert A. Smith, and Bradley D. Preston. 2011. "Mutator Suppression and
Escape from Replication Error-Induced Extinction in Yeast." *PLoS Genetics* 7 (10):
e1002282.
Hoffman, Charles S., and Fred Winston. 1987. "A Ten-Minute DNA Preparation from Yeast
Efficiently Releases Autonomous Plasmids for Transformaion of Escherichia Coli." *Gene* 57
(2): 267–72.
Horst, J. P., T. H. Wu, and M. G. Marinus. 1999. "Escherichia Coli Mutator Genes." *Trends in
Microbiology* 7 (1): 29–36.
Huang, Meng-Er, Anne-Gaelle Rio, Alain Nicolas, and Richard D. Kolodner. 2003. "A
Genomewide Screen in Saccharomyces Cerevisiae for Genes That Suppress the
Accumulation of Mutations." *Proceedings of the National Academy of Sciences of the
United States of America* 100 (20): 11529–34.
Hwang, Dick G., and Phil Green. 2004. "Bayesian Markov Chain Monte Carlo Sequence
Analysis Reveals Varying Neutral Substitution Patterns in Mammalian Evolution."
Proceedings of the National Academy of Sciences of the United States of America 101
(39): 13994–1.
Iossifov, Ivan, Brian J. O'Roak, Stephan J. Sanders, Michael Ronemus, Niklas Krumm, Dan
Levy, Holly A. Stessman, et al. 2014. "The Contribution of de Novo Coding Mutations to
Autism Spectrum Disorder." *Nature* 515 (7526): 216–21.
Jiang, Caifu, Aziz Mithani, Eric J. Belfield, Richard Mott, Laurence D. Hurst, and Nicholas P.
Harberd. 2014. "Environmentally Responsive Genome-Wide Accumulation of de Novo
Arabidopsis Thaliana Mutations and Epimutations." *Genome Research* 24 (11): 1821–29.
Johnson, T. 1999. "Beneficial Mutations, Hitchhiking and the Evolution of Mutation Rates in
Sexual Populations." *Genetics* 151 (4): 1621–31.
Jubin, Claire, Alexandre Serero, Sophie Loeillet, Emmanuel Barillot, and Alain Nicolas. 2014.
"Sequence Profiling of the Saccharomyces Cerevisiae Genome Permits Deconvolution of
Unique and Multialigned Reads for Variant Detection." *G3* 4 (4): 707–15.
Keightley, Peter D., Urmi Trivedi, Marian Thomson, Fiona Oliver, Sujai Kumar, and Mark L.
Blaxter. 2009. "Analysis of the Genome Sequences of Three Drosophila Melanogaster
Spontaneous Mutation Accumulation Lines." *Genome Research* 19 (7): 1195–1201.
Kessler, Michael D., Douglas P. Loesch, James A. Perry, Nancy L. Heard-Costa, Daniel Taliun,
Brian E. Cade, Heming Wang, et al. 2020. "De Novo Mutations across 1,465 Diverse
Genomes Reveal Mutational Insights and Reductions in the Amish Founder Population."
Proceedings of the National Academy of Sciences of the United States of America 117 (5):
2560–69.
Kimura, Motoo. 1967. "On the Evolutionary Adjustment of Spontaneous Mutation Rates*."
Genetics Research 9 (1): 23–34.
Krueger, Felix. n.d. *TrimGalore*. Github. Accessed May 28, 2020.
https://github.com/FelixKrueger/TrimGalore.
Laluce, C., M. C. Bertolini, J. R. Ernandes, A. V. Martini, and A. Martini. 1988. "New Amylolytic
Yeast Strains for Starch and Dextrin Fermentation." *Applied and Environmental
Microbiology* 54 (10): 2447–51.
Lang, Gregory I. 2018. "Measuring Mutation Rates Using the Luria-Delbrück Fluctuation Assay."
In *Genome Instability: Methods and Protocols*, edited by Marco Muzi-Falconi and Grant W.
Brown, 21–31. New York, NY: Springer New York.
Lang, Gregory I., and Andrew W. Murray. 2008. "Estimating the Per-Base-Pair Mutation Rate in
the Yeast Saccharomyces Cerevisiae." *Genetics* 178 (1): 67–82.

———. 2011. "Mutation Rates across Budding Yeast Chromosome VI Are Correlated with Replication Timing." *Genome Biology and Evolution* 3 (June): 799–811.

Lang, Gregory I., Lance Parsons, and Alison E. Gammie. 2013. "Mutation Rates, Spectra, and Genome-Wide Distribution of Spontaneous Mutations in Mismatch Repair Deficient Yeast." *G3* 3 (9): 1453–65.

Langmead, Ben, and Steven L. Salzberg. 2012. "Fast Gapped-Read Alignment with Bowtie 2." *Nature Methods* 9 (4): 357–59.

Lea, D. E., and C. A. Coulson. 1949. "The Distribution of the Numbers of Mutants in Bacterial Populations." *Journal of Genetics* 49 (3): 264–85.

Leigh, Egbert Giles. 1970. "Natural Selection and Mutability." *The American Naturalist* 104 (937): 301–5.

Li, Heng, and Richard Durbin. 2009. "Fast and Accurate Short Read Alignment with Burrows-Wheeler Transform." *Bioinformatics* 25 (14): 1754–60.

Liti, Gianni, David M. Carter, Alan M. Moses, Jonas Warringer, Leopold Parts, Stephen A. James, Robert P. Davey, et al. 2009. "Population Genomics of Domestic and Wild Yeasts." *Nature* 458 (7236): 337–41.

Liu, Haoxuan, and Jianzhi Zhang. 2019. "Yeast Spontaneous Mutation Rate and Spectrum Vary with Environment." *Current Biology: CB* 29 (10): 1584–91.e3.

———. 2021. "The Rate and Molecular Spectrum of Mutation Are Selectively Maintained in Yeast." *Nature Communications* 12 (1): 4044.

Loeb, L. A. 2001. "A Mutator Phenotype in Cancer." *Cancer Research* 61 (8): 3230–39.

Loeb, Lawrence A. 2016. "Human Cancers Express a Mutator Phenotype: Hypothesis, Origin, and Consequences." *Cancer Research* 76 (8): 2057–59.

Luria, S. E., and M. Delbrück. 1943. "Mutations of Bacteria from Virus Sensitivity to Virus Resistance." *Genetics* 28 (6): 491–511.

Lynch, Michael, Matthew S. Ackerman, Jean-Francois Gout, Hongan Long, Way Sung, W. Kelley Thomas, and Patricia L. Foster. 2016. "Genetic Drift, Selection and the Evolution of the Mutation Rate." *Nature Reviews. Genetics* 17 (11): 704–14.

Lynch, Michael, Way Sung, Krystalynne Morris, Nicole Coffey, Christian R. Landry, Erik B. Dopman, W. Joseph Dickinson, et al. 2008. "A Genome-Wide View of the Spectrum of Spontaneous Mutations in Yeast." *Proceedings of the National Academy of Sciences of the United States of America* 105 (27): 9272–77.

Ma, W. T., G. Vh. Sandri, and S. Sarkar. 1992. "Analysis of the Luria-Delbrück Distribution Using Discrete Convolution Powers." *Journal of Applied Probability* 29 (2): 255–67.

McGregor, Alistair P., Virginie Orgogozo, Isabelle Delon, Jennifer Zanet, Dayalan G. Srinivasan, François Payre, and David L. Stern. 2007. "Morphological Evolution through Multiple Cis-Regulatory Mutations at a Single Gene." *Nature* 448 (7153): 587–90.

McLaren, William, Laurent Gil, Sarah E. Hunt, Harpreet Singh Riat, Graham R. S. Ritchie, Anja Thormann, Paul Flicek, and Fiona Cunningham. 2016. "The Ensembl Variant Effect Predictor." *Genome Biology* 17 (1): 122.

Miles, Alistair. n.d. *Pysamstats*. Github. Accessed May 28, 2020. https://github.com/alimanfoo/pysamstats.

Nei, Masatoshi. 1983. "Genetic Polymorphism and the Role of Mutation in Evolution." *Evolution of Genes and Proteins* 71: 165–90.

Ni, T. T., G. T. Marsischky, and R. D. Kolodner. 1999. "MSH2 and MSH6 Are Required for Removal of Adenine Misincorporated Opposite 8-Oxo-Guanine in S. Cerevisiae." *Molecular Cell* 4 (3): 439–44.

Peter, Jackson, Matteo De Chiara, Anne Friedrich, Jia-Xing Yue, David Pflieger, Anders

Bergström, Anastasie Sigwalt, et al. 2018. "Genome Evolution across 1,011
Saccharomyces Cerevisiae Isolates." *Nature* 556 (7701): 339–44.
Poplin, Ryan, Valentin Ruano-Rubio, Mark A. DePristo, Tim J. Fennell, Mauricio O. Carneiro,
Geraldine A. Van der Auwera, David E. Kling, et al. 2018. "Scaling Accurate Genetic Variant
Discovery to Tens of Thousands of Samples." *bioRxiv*. https://doi.org/10.1101/201178.
Prindle, Marc J., Edward J. Fox, and Lawrence A. Loeb. 2010. "The Mutator Phenotype in
Cancer: Molecular Mechanisms and Targeting Strategies." *Current Drug Targets* 11 (10):
1296–1303.
Raghavan, Vandana, Duyen T. Bui, Najla Al-Sweel, Anne Friedrich, Joseph Schacherer, Charles
F. Aquadro, and Eric Alani. 2018. "Incompatibilities in Mismatch Repair Genes MLH1-PMS1
Contribute to a Wide Range of Mutation Rates in Human Isolates of Baker's Yeast."
Genetics 210 (4): 1253–66.
Risques, Rosa Ana, and Scott R. Kennedy. 2018. "Aging and the Rise of Somatic
Cancer-Associated Mutations in Normal Tissues." *PLoS Genetics* 14 (1): e1007108.
Sasani, Thomas A., David G. Ashbrook, Lu Lu, Abraham A. Palmer, Robert W. Williams,
Jonathan K. Pritchard, and Kelley Harris. 2021. "A Wild-Derived Antimutator Drives
Germline Mutation Spectrum Differences in a Genetically Diverse Murine Family." *bioRxiv*.
https://doi.org/10.1101/2021.03.12.435196.
Scally, Aylwyn, and Richard Durbin. 2012. "Revising the Human Mutation Rate: Implications for
Understanding Human Evolution." *Nature Reviews. Genetics* 13 (10): 745–53.
Schacherer, Joseph, Joshua A. Shapiro, Douglas M. Ruderfer, and Leonid Kruglyak. 2009.
"Comprehensive Polymorphism Survey Elucidates Population Structure of Saccharomyces
Cerevisiae." *Nature* 458 (7236): 342–45.
Schrider, Daniel R., Jonathan N. Hourmozdi, and Matthew W. Hahn. 2011. "Pervasive
Multinucleotide Mutational Events in Eukaryotes." *Current Biology: CB* 21 (12): 1051–54.
Sebat, Jonathan, B. Lakshmi, Dheeraj Malhotra, Jennifer Troge, Christa Lese-Martin, Tom
Walsh, Boris Yamrom, et al. 2007. "Strong Association of de Novo Copy Number Mutations
with Autism." *Science* 316 (5823): 445–49.
Ségurel, Laure, Minyoung J. Wyman, and Molly Przeworski. 2014. "Determinants of Mutation
Rate Variation in the Human Germline." *Annual Review of Genomics and Human Genetics*
15 (June): 47–70.
Serero, Alexandre, Claire Jubin, Sophie Loeillet, Patricia Legoix-Né, and Alain G. Nicolas. 2014.
"Mutational Landscape of Yeast Mutator Strains." *Proceedings of the National Academy of
Sciences of the United States of America* 111 (5): 1897–1902.
Sharp, Nathaniel P., Linnea Sandell, Christopher G. James, and Sarah P. Otto. 2018. "The
Genome-Wide Rate and Spectrum of Spontaneous Mutations Differ between Haploid and
Diploid Yeast." *Proceedings of the National Academy of Sciences of the United States of
America* 115 (22): E5046–55.
Shockley, Arthur H., David W. Doo, Gina P. Rodriguez, and Gray F. Crouse. 2013. "Oxidative
Damage and Mutagenesis in Saccharomyces Cerevisiae: Genetic Studies of Pathways
Affecting Replication Fidelity of 8-Oxoguanine." *Genetics* 195 (2): 359–67.
Shor, Erika, Catherine A. Fox, and James R. Broach. 2013. "The Yeast Environmental Stress
Response Regulates Mutagenesis Induced by Proteotoxic Stress." *PLoS Genetics* 9 (8):
e1003680.
Steenwyk, Jacob L., Dana A. Opulente, Jacek Kominek, Xing-Xing Shen, Xiaofan Zhou, Abigail
L. Labella, Noah P. Bradley, et al. 2019. "Extensive Loss of Cell-Cycle and DNA Repair
Genes in an Ancient Lineage of Bipolar Budding Yeasts." *PLoS Biology* 17 (5): e3000255.
Stirling, Peter C., Yaoqing Shen, Richard Corbett, Steven J. M. Jones, and Philip Hieter. 2014.

        "Genome Destabilizing Mutator Alleles Drive Specific Mutational Trajectories in
        Saccharomyces Cerevisiae." *Genetics* 196 (2): 403–12.

Stone, Jana E., Scott A. Lujan, Thomas A. Kunkel, and Thomas A. Kunkel. 2012. "DNA
        Polymerase Zeta Generates Clustered Mutations during Bypass of Endogenous DNA
        Lesions in Saccharomyces Cerevisiae." *Environmental and Molecular Mutagenesis* 53 (9):
        777–86.

Sturtevant, A. H. 1937. "Essays on Evolution. I. On the Effects of Selection on Mutation Rate."
        The Quarterly Review of Biology 12 (4): 464–67.

Sung, Way, Matthew S. Ackerman, Samuel F. Miller, Thomas G. Doak, and Michael Lynch.
        2012. "Drift-Barrier Hypothesis and Mutation-Rate Evolution." *Proceedings of the National
        Academy of Sciences of the United States of America* 109 (45): 18488–92.

Sung, Way, Abraham E. Tucker, Thomas G. Doak, Eunjin Choi, W. Kelley Thomas, and Michael
        Lynch. 2012. "Extraordinary Genome Stability in the Ciliate Paramecium Tetraurelia."
        Proceedings of the National Academy of Sciences of the United States of America 109
        (47): 19339–44.

Tenaillon, Olivier, Jeffrey E. Barrick, Noah Ribeck, Daniel E. Deatherage, Jeffrey L. Blanchard,
        Aurko Dasgupta, Gabriel C. Wu, et al. 2016. "Tempo and Mode of Genome Evolution in a
        50,000-Generation Experiment." *Nature* 536 (7615): 165–70.

Tong, Amy Hin Yan, and Charles Boone. 2007. "16 High-Throughput Strain Construction and
        Systematic Synthetic Lethal Screening in Saccharomycescerevisiae." In *Methods in
        Microbiology*, edited by Ian Stansfield and Michael Stark JR, 36:369–707. Academic Press.

Tracy, Maxwell A., Mitchell B. Lee, Brady L. Hearn, Ian T. Dowsett, Luke C. Thurber, Jason Loo,
        Anisha M. Loeb, et al. 2020. "Spontaneous Polyploids and Antimutators Compete During
        the Evolution of Saccharomyces Cerevisiae Mutator Cells." *Genetics* 215 (4): 959–74.

Viel, Alessandra, Alessandro Bruselles, Ettore Meccia, Mara Fornasarig, Michele Quaia,
        Vincenzo Canzonieri, Eleonora Policicchio, et al. 2017. "A Specific Mutational Signature
        Associated with DNA 8-Oxoguanine Persistence in MUTYH-Defective Colorectal Cancer."
        EBioMedicine 20 (June): 39–49.

Voordeckers, Karin, Camilla Colding, Lavinia Grasso, Benjamin Pardo, Lore Hoes, Jacek
        Kominek, Kim Gielens, et al. 2020. "Ethanol Exposure Increases Mutation Rate through
        Error-Prone Polymerases." *Nature Communications* 11 (1): 3664.

Wang, Long, Yilun Ji, Yingwen Hu, Huaying Hu, Xianqin Jia, Mengmeng Jiang, Xiaohui Zhang,
        et al. 2019. "The Architecture of Intra-Organism Mutation Rate Variation in Plants." *PLoS
        Biology* 17 (4): e3000191.

Weng, Mao-Lun, Claude Becker, Julia Hildebrandt, Manuela Neumann, Matthew T. Rutter, Ruth
        G. Shaw, Detlef Weigel, and Charles B. Fenster. 2019. "Fine-Grained Analysis of
        Spontaneous Mutation Spectrum and Frequency in Arabidopsis Thaliana." *Genetics* 211
        (2): 703–14.

Whelan, W. L., E. Gocke, and T. R. Manney. 1979. "The CAN1 Locus of Saccharomyces
        Cerevisiae: Fine-Structure Analysis and Forward Mutation Rates." *Genetics* 91 (1): 35–51.

Yue, Jia-Xing, Jing Li, Louise Aigrain, Johan Hallin, Karl Persson, Karen Oliver, Anders
        Bergström, et al. 2017. "Contrasting Evolutionary Genome Dynamics between
        Domesticated and Wild Yeasts." *Nature Genetics* 49 (6): 913–24.

Zhang, Jiajie, Kassian Kobert, Tomáš Flouri, and Alexandros Stamatakis. 2014. "PEAR: A Fast
        and Accurate Illumina Paired-End reAd mergeR." *Bioinformatics* 30 (5): 614–20.

Zheng, Qi. 2017. "rSalvador: An R Package for the Fluctuation Experiment." *G3* 7 (12):
        3849–56.

Zhu, Yuan O., Gavin Sherlock, and Dmitri A. Petrov. 2017. "Extremely Rare Polymorphisms in

Saccharomyces Cerevisiae Allow Inference of the Mutational Spectrum." *PLoS Genetics* 13
(1): e1006455.
Zhu, Yuan O., Mark L. Siegal, David W. Hall, and Dmitri A. Petrov. 2014. "Precise Estimates of
Mutation Rate and Spectrum in Yeast." *Proceedings of the National Academy of Sciences*
of the United States of America 111 (22): E2310–18.
Zou, Xueqing, Gene Ching Chiek Koh, Arjun Scott Nanda, Andrea Degasperi, Katie Urgo,
Theodoros I. Roumeliotis, Chukwuma A. Agu, et al. 2020. "Dissecting Mutational
Mechanisms Underpinning Signatures Caused by Replication Errors and Endogenous DNA
Damage." *bioRxiv*. https://doi.org/10.1101/2020.08.04.234245.