1 Is earlier reproduction associated with higher or lower survival? Antagonistic
2 results between individual and population scales in the blue tit

AUTHORS

Olivier Bastianelli[1,2], Anne Charmantier[2], Clotilde Biard[3], Suzanne Bonamour[1,2], Céline
Teplitsky[2], Alexandre Robert[1*]

AFFILIATIONS

1. Centre d'Ecologie et des Sciences de la Conservation (CESCO), Muséum national d'Histoire
naturelle, Centre National de la Recherche Scientifique, Sorbonne Université, Paris, France

2. CEFE, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul Valéry Montpellier 3, Montpellier,
France

3. Sorbonne Université, UPEC, Paris 7, CNRS, INRA, IRD, Institut d'Ecologie et des Sciences de
l'Environnement de Paris, iEES Paris, F-75005 Paris, France.

*Author for correspondence

KEYWORDS

Survival, Capture-Mark-Recapture, Laying date, *Cyanistes caeruleus*, Phenological mismatch,
Demographic heterogeneity

ABSTRACT

Although it has been shown that phenology can respond to temporal environmental
variation in free ranging populations of several species, little is known about the mechanisms
of these responses and their effects on demography, and in particular on survival. Exploring
phenological responses and their associated consequences on survival can be achieved at
two distinct scales: the population scale, which focusses on a set of common responses to
environmental conditions, and the individual scale, focusing on the relative position of each
individual in the distribution of survival and phenology under particular conditions. In this
study, we apply capture-mark-recapture multistate modelling on a 38-year monitoring
dataset of blue tits (*Cyanistes caeruleus*) to investigate the effects of breeding phenology
and some demographic covariates (breeding density, average and individual breeding
success) on adult survival, at both population and individual scales. Our analysis revealed
that (i) at the population scale, early breeding years are followed by lower average adult
survival. (ii) At the individual level, earlier breeders within the population have higher
subsequent survival than later breeders, although this relationship is reversed in years with
very harsh conditions, e.g. warm spring and high breeding density. (iii) High individual
relative breeding success is also associated with higher subsequent survival and explains
more survival variation than relative phenology. Overall, our study indicates that, although
earlier breeding is associated with a survival cost at the population level, substantial
intrapopulation hererogeneity shapes a positive association between earlier breeding,
breeding success and survival at the individual level.

INTRODUCTION

The phenological response of organisms to variation in their local environment is a major issue in evolutionary ecology and conservation research, both because the diversity of this response reflects differences in evolutionary potential and strategies between and within species (Thackeray et al., 2016), and because advances in reproductive phenology is seen as one of the universal ecological responses to global warming (Durant et al., 2007; Parmesan, 2007; Radchuk et al., 2019). Although it has been shown that phenology can respond to meteorological variations (noise and cycles) as well as climatic trends in many plant (Bjorkman et al., 2015; Diekmann, 1996) and animal (Duursma et al., 2018; Samplonius et al., 2018; Sheriff et al., 2015; Haest et al., 2018) species, little is known about the mechanisms of these responses and their effects on demography and population dynamics (Forrest and Miller-Rushing, 2010, but see Shipley et al., 2020; Simmonds et al., 2020).

In particular, it is likely that phenological variations, which may involve variations in strategy within populations, are constrained by trade-offs (Diggle, 1999; Forrest and Miller-Rushing, 2010). For example, in birds, while early reproduction might be beneficial in terms of reproductive success (Harriman et al., 2017; in the focal population: Marrot et al., 2018), it is also potentially associated with subsequent survival cost for the breeders (e.g. Brinkhof et al., 2002; Nilsson, 1994) or the early hatched nestlings (Shipley et al., 2020). The presence of these trade-offs could partly explain phenological mismatches (*i.e.* a difference between a species phenology and its main resource phenology) frequently observed in bird populations (Visser et al., 2012). Although a review of the literature provides multiple estimations of reproductive selection acting on avian breeding phenology (Verhulst and Nilsson, 2008; Charmantier and Gienapp, 2014), similar links between survival and timing of breeding are too scarce to conclude on the presence of such trade-offs. One reason why survival has not been explored frequently in this context might be because it is often assumed that reproductive success will be affected by breeding phenology while survival will be influenced by migration phenology (e.g. Visser and Gienapp, 2019).

Additionally, the survival consequences of phenological variations may be influenced by other mechanisms such as intraspecific competition. Several ecological field studies (in both long-lived and short-lived species) have reported lower population level survival rates at high densities (Abadi et al., 2012; Fay et al., 2015; Brouwer et al., 2006; Bonenfant et al., 2009; Le Cœur et al., 2016) and competition has been shown to influence population dynamics in many studies (see e.g. Grøtan et al., 2009 and Gamelon et al., 2016 in the Great tit *Parus major*). Importantly, however, in the presence of both year-to-year environmental changes and intraspecific competition, population density can also be a factor maintaining demographic stability in a changing environment. Reed and colleagues (Reed et al., 2013) showed that in a Great tit population, the reduced reproductive success observed in years with greater mismatch between the tits and their prey was compensated by a higher overwinter survival due to relaxed competition.

Overall, exploring breeding phenological responses and their associated demographic benefits and costs can be achieved at two distinct scales: a/ the population scale, by studying for example the overall response of a population (e.g. mean or median phenology, reproductive success or survival) to interannual environmental variation, or b/ the individual level, by focusing on the relative position of each individual in the distribution of reproductive success, survival and phenology under particular conditions (e.g. for the same year, see e.g. Harriman et al., 2017 for an experimental approach focusing on breeding success). The first approach examines a set of common responses (behavioural, demographic) to environmental conditions, while the second approach focuses on individual heterogeneity within a population, in terms of fitness components. While some external constraints induced by the limitation of resources for survival and reproduction are shared by all individuals of the population, differences can arise in resource acquisition due to e.g. genotype or developmental conditions (Lindström, 1999; van Noordwijk and de Jong, 1986; Wilson and Nussey, 2010).

In the context of exploring demographic consequences of phenological changes, investigating variation within and between individuals is of primary importance as it occurs in virtually all traits, including demographic components such as reproduction and survival (Clutton-Brock, 1988). Between-individual variation may be linked with phenotypic attributes that change along an individual's life, such as age or breeding status, or with fixed attributes (such as sex or genotype, see Gimenez et al., 2018 and references therein). A significant part of the research on individual heterogeneity has focused on the covariation between demographic components such as reproduction and survival. In particular, a negative relationship between survival and reproduction is expected in the context of a trade-off (energetic or genetic) between these two components (Harshman and Zera, 2007; Roff, 1993; Stearns, 1992). On the contrary, a positive relationship is expected within the framework of dynamic (Tuljapurkar et al., 2009) and/or fixed (Cam et al., 2013) variations between individuals. Fixed individual heterogeneity is linked, for example, to intrinsic or resource acquisition differences early in life (Hamel et al., 2009; Lescroël et al., 2009), sometimes referred to as individual quality (Cam et al., 2002; Wilson and Nussey, 2010). Reproductive costs and individual quality are not mutually exclusive and have been suggested to interact with each other as well as with environmental (Robert et al., 2012) and individual (Robert et al., 2015) conditions.

Although phenological variation within populations has been well described in many organisms (Elzinga et al., 2007; Forrest and Miller-Rushing, 2010; Laaksonen et al., 2006), the relationship between such variation and demographic heterogeneity remains poorly understood both empirically and theoretically. Even in the well-studied context of temperate forest passerines where early seasonal breeding has been repeatedly associated with higher breeding success, the theoretical frameworks outlined above on trade-offs *versus* resource acquisition and individual heterogeneity can lead to two opposing predictions. First, one could predict that, in a given population a given year, the earliest breeders will show the lowest survival due to a reproduction-survival trade-off (Brinkhof et al., 2002; Wiggins et al., 1998; Nilsson, 1994; Nilsson and Svensson, 1996). Alternately, one could expect that the

high "quality", earliest breeders (e.g. with highest resource acquisition, van Noordwijk and
de Jong, 1986), will exhibit the highest survival. Additionally, the question remains whether
the observed phenology-survival pattern is consistent across years, in particular whether it is
similar in years with early or late average breeding dates. Answering these questions
requires studying the links between reproductive phenology and survival, first at the
population level, second at the individual heterogeneity level, and third the links between
these two levels.

In this study, using a long-term monitoring dataset, we investigated the effect of breeding
phenology on adult survival in a population of a temperate passerine bird, the Blue tit
(*Cyanistes caeruleus*). The data comprised 1562 adult individuals captured and ringed in nest
boxes during the reproduction period across 38 years (1979-2016). In this population,
breeding phenology is influenced by meteorological conditions. Median laying date is earlier
in warmer years (Bonamour et al., 2019) and has an impact on breeding success: earlier
breeding is correlated with higher number of fledglings at the individual and population
scales (Marrot et al., 2018). Long-term effects of climate on the phenology of the studied
population also occurred over the period of the study (Marrot et al., 2018), during which
spring temperatures have significantly increased (1979-2016: +0.49°C/decade; *p-value* < 10^{-3})
and median laying dates have significantly advanced (1979-2016: -2.97 days/decade; *p-value*
< 10^{-3}).

Using flexible capture-mark-recapture multistate modelling (Lebreton et al., 2003), we
addressed four different questions: First, is there a link between population-level phenology
and average annual adult survival in the focal Blue tit population? Since previous
experimental studies in birds (Brinkhof et al., 2002; Wiggins et al., 1998) including blue tits
(Nilsson, 1994; Nilsson and Svensson, 1996) have revealed a cost of early breeding on adult
survival, we predict a low average adult survival following early population breeding
phenology. Second, within a given year, what is the link between individual timing of
breeding (quantified as laying date quartiles of the population for this particular year,
hereafter *relative laying date*) and adult survival? We expect to find either a positive
association between early breeding and survival if "good quality" individuals are those that
invest in early breeding (Robb et al., 2008) or, on the contrary, a negative association if the
cost of breeding earlier outweighs differences in individual heterogeneity (Tarwater and
Arcese, 2017). Third, does the link between the relative laying date and adult survival vary
depending on the conditions faced in a given year (e.g., population density)? Here we expect
to find a weaker link between relative laying date and adult survival in years with low
constraints on breeding (e.g. low population density) and therefore probable relaxed cost of
breeding. Fourth and finally, since breeding early and having a high breeding success are
correlated (*i.e.* early broods are larger: review in Klomp, 1970, in the focal population
Delahaie et al., 2017) but may convey distinct costs or reflect distinct facets of individual
heterogeneity, our last interrogation is about the link between relative laying date, survival
and individual breeding success. Specifically, are large brood sizes relative to the population
associated with high or low survival, and what is the magnitude of this effect relative to the
effect of laying date?

Methods

1   Population Monitoring

Our data was collected over 38 years (1979-2016) in a natural population of blue tits *Cyanistes caeruleus* in an evergreen oak (*Quercus ilex*) forest on the Mediterranean island of Corsica (E-Pirio site: Lat 42.38; Long: 8.75; see Blondel et al., 2006 and Charmantier et al., 2016 for details). Blue tits are cavity-nesting non-migratory passerine birds that readily breed in nest boxes, which they often prefer to natural cavities (Newton, 1994). The high density of nest boxes compared to the rarity of natural cavities in the E-Pirio site (A.C., unpublished data) allowed us to monitor almost all breeding individuals in the E-Pirio study area. During the whole study period, nest boxes were checked every year at least weekly during the breeding season (March to June), allowing to accurately assess the date of the first laid egg for all clutches (hereafter, laying date), as well as the clutch size and the number of fledglings. Breeders were captured in the boxes when nestlings were 10-15 days old, *i.e.* late enough to avoid the risk of nest abandonment, but early enough to avoid early departure by the nestlings. Parents were identified with a unique numbered metal ring provided by the Centre de Recherches sur la Biologie des Populations d'Oiseaux (CRBPO, which also provided the permits under which capture and handling of birds were conducted). Sex and age at first capture were determined by plumage patterns for every breeder. As these patterns cannot be used to age individuals older than two years, only two adult age classes were used in survival modelling: 1 year and 2 or more years. Individuals whose age at first capture had not be assessed were removed (3.4% (n=55) of the individuals). In total, 1562 reproducing individuals were considered in this study. All data analysed concerned first broods only (second broods represent less than 2.5% of breeding events in this population).

2   Population-level annual covariates

For each year of the study, we computed the median of individual laying dates in the population (median laying date, medLD), the mean number of fledglings per attempted first brood in the population (reproductive success, repS) and the arcsine normalized breeding density of the population (population density, popD).

Population breeding density was obtained by measuring nest box occupation rates by blue tits in a defined area of the study site, where nest boxes number varied as little as possible over the 38 years of study (n=58 to 62 nest boxes). Only blue tit occupation was considered as the diameter of the nest boxes entry hole (26 mm) was too small to allow regular occurrence of Great tit (*Parus major*) nesting.

The variable used to study how meteorological conditions affect individual survival and reproduction was the average daily mean temperature during the period from March 31[st] to May 7[th] (hereafter, Temp). This 38 day window has been identified in a previous study using a sliding window analysis, as the time window most correlated with the average annual laying date in the focal population (see Table 1 in Bonamour et al., 2019). The temperature

210 data were obtained by regressing daily temperature measurements over 4 years in the E-
211 Pirio station with temperature data from the meteorological station of Calvi (Lat: 42.52;
212 Long: 8.79, 15.9 km away from the population) and inferring the temperature in E-Pirio over
213 the remaining years.

3   Capture-Mark-Recapture modelling

Individual capture-recapture histories for the 1562 breeding individuals were analysed to
provide robust estimates of survival and recapture probabilities (respectively φ and p,
Lebreton et al., 1992), using a logit-link function. Monostate Capture-Mark-Recapture
modelling was used for the assessment of the link between annual population-level
phenology and annual survival, while multistate capture-mark-recapture models (Lebreton
et al., 2003) were necessary for assessing the link between intra-population variation of
phenology (i.e., relative laying date), breeding success and adult survival. Multistate models
make it possible to model the survival of individuals that may be in different states (which
correspond to different categories of laying date or reproductive success in the case of our
study). These models include four types of parameters: the probabilities of being in the
various initial states (IS), the probabilities of transition among states (T), survival
probabilities (φ) and recapture probabilities (p). All analyses were conducted using the
program E-SURGE (Choquet et al., 2009b). Goodness-of-fit of models to the data was
ensured for each dataset using the program U-CARE (Choquet et al., 2009a), based on the
Cormack Jolly Seber model for monostate models and on the Jolly Move model for
multistate models.

In several years, experiments were conducted in the population including brood size
manipulations that can highly alter adult survival (Dijkstra et al., 1990; Nur, 1984). Capture-
recapture histories of the corresponding individuals were censored right after the first time
they experienced such a fitness-changing experiment. However, nestlings born during these
experiments and later recaptured as adults were not removed from the analysis. This
necessary censoring resulted in a significant decrease in observation numbers (-23.6% from
2955 to 2258 capture and recapture events in the monostate dataset). Breeding events in
experimental conditions were also removed from reproductive success estimations.

4   Data analysis and model selection

The Akaike's Information Criterion corrected for small sample size (AICc) was used for model
selection (Burnham and Anderson, 1998). A low AICc was considered revealing a good
compromise between the fit to the data (likelihood of the model) and the number of
parameters used by the model. The threshold for significant difference between two models
was set at two AICc points. In case of lower difference, the model with the lowest number of
parameters was selected. The four main questions addressed are summarised in Table 1 and
details are provided in the sections below.

a   Link between population-level phenology and average annual adult survival

Prior to assessing the effects of population-level covariates on survival, a preliminary
model selection assessed the effects of year, sex and age on recapture (p) and survival
(φ) probabilities with the monostate dataset of capture histories (Table 2A). The starting
model (Model 22) included potential effects of sex, age and yearly variations (hereafter,
year) on survival and potential effects of sex and year on recapture probabilities. Only
simple (first order) interactions between variables influencing survival and recapture
probabilities were considered, to enable robust biological interpretation of the results
and avoid overcomplexity. All subsequent models were nested in this one. The model
selection started with simplifying constraints on p. As a second step, we retained the
best model structure for recapture and considered simpler survival models. We then
used the best model to test the effects of our three demographic temporal covariates
(medLD, popD, repS) and their interactions (Table 2B) on survival.

b   Link between individual timing of breeding (relative laying date) and adult survival

To assess the effect of intra-population variation in laying dates, a multistate dataset of
capture histories for our population was created. The different states were defined as
the quartiles of laying date distribution of the population in a given year, for instance the
25% earliest breeders in the population for a given year were given the state "1" for this
year, and the latest quarter were given state "4". Observations without any laying date
information were removed (4 observations, 3 individuals, out of 2258 observations).

The starting model used for model selection (Table 3, Model 85) was set to have only sex
allowed as a possible constraint for recapture probabilities (p) based on the above
selection procedure on the monostate model. It included (1) all possible interactions
between state, sex and age for the probability estimates of being in a given initial state
(IS); (2) interactions between sex, age, departure and arrival state for between state
transition probabilities (T); (3) interactions between state, sex and age for survival
probabilities (φ). Selection was made sequentially (i.e. by retaining the best structure at
each step for the next step) firstly on initial state distribution constraints, then on
transition probabilities constraints. For survival constraints, the possible interactive
effects of sex and age with the states were tested. Then various potential groupings of
states were tested with the association of sex and age as in the best previous models,
including grouping of quartiles 1 vs 2 3 4, 1 2 3 vs 4, 1 vs 2 3 vs 4, 1 2 vs 3 4, 1 vs 2 vs 3 4,
1 2 vs 3 vs 4.

c   Effects of population-level annual covariates on the relative laying date - adult
survival link

Based on the model selection described above, we started from a model taking into
account differences in laying date quartiles (Tables 3 and 4: model 43), and ran a further
model selection by adding four population-level annual metrics (Temp, medLD, popD,
repS) and their interactions as continuous temporal covariates (Table 4).

d   Links between individual breeding success, relative laying date and adult survival

To assess the role of intra-population reproductive investment in the system, another
multistate dataset of capture histories was used. The 6 states took into account
combinations of the individual differences in reproduction in three categories (Failure:
clutch abandoned or predated, 0 fledglings; Low reproduction: number of fledglings
below the median number of fledglings per brood in the population this year; High
reproduction: number of fledglings above this median) and individual differences in
phenology in two categories (Early breeders: first quartile of laying date (state 1 in the
previous dataset), Late breeders: second to fourth quartiles of laying date (states 2,3 and
4 in the previous dataset)).

The starting model for the selection included the interactive effects of age, sex and state
for its various structural components (Table 5, model 276). Recapture probabilities were
assumed to depend on sex only. The sequential selection procedure was the same as
above. For survival rates, various groupings of states were tested (Table 5) including
grouping of states 1 2 vs 3 4 5 6, 1 2 vs 3 4 vs 5 6, 1 3 5 vs 2 4 6, 1 2 vs 3 5 vs 4 6.

	Population-level analysis	Individual-level analysis	Additional annual population-level covariates tested on survival	Type of models used
a) Link between population-level phenology and average annual adult survival	Phenology ↔ Survival		Demographic covariates (median laying date, population density, mean reproductive success)	Monostate models
b) Link between individual timing of breeding (relative laying date) and adult survival		Relative Phenology ↔ Survival		Multistate models (4 relative laying date states)
c) Effects of population-level annual covariates on the relative laying date - adult survival link		Relative Phenology ↔ Survival	Demographic covariates and spring temperature	Multistate models (4 relative laying date states)
d) Links between individual breeding success, relative laying date and adult survival		Relative Laying date and Relative Reproductive Success ↔ Survival		Multistate models (6 relative laying date and reproductive success states)

304 **Table 1:** Summary of the four main types of questions addressed and capture-mark-recapture models used in this study.

Results

Goodness-of-fit

No significant violation of standard models assumptions was found while performing the goodness-of-fit tests using program U-CARE (Choquet et al., 2009a): Detailed goodness-of-fit results are provided in Appendix 1.

a   Link between population-level phenology and average annual adult survival

The best model from our preliminary selection on the single state capture histories (without covariates) is presented in Table 2A (Model 1). This model assumes an effect of sex only on recapture probabilities, with higher recapture rates for females (p_{males} = 0.673 [0.611; 0.729]$_{95\%CI}$; $p_{females}$ = 0.859 [0.805; 0.899] $_{95\%CI}$), and of age only on survival probabilities, with higher survival probabilities for first year adults than older adults (ϕ_{1year} = 0.597 [0.546; 0.645] $_{95\%CI}$; $\phi_{2+years}$ = 0.526 [0.494; 0.558] $_{95\%CI}$).

When adding the three population-level annual demographic covariates (median laying date of the population (medLD), population density (popD) and mean number of fledglings per attempted brood in the population (repS)) as determinants of survival, we observed that adding each covariate separately to Model 1 improved the fit to the data (Table 2B, models 35, 36 and 37). The best model from this selection was Model 23, which took into account the effects of the medLD, popD and their interaction.

All models showed a positive relationship between the median laying date and adult survival and a negative relationship between density and survival, which means that a late reproduction of the population was associated with a high subsequent adult survival, and that a high density of the population was associated with a low subsequent survival (Figure 1A and B). Importantly, these relationships were not due to a change in the proportion of age classes between years and remained unchanged within each age class (see Appendix 3 for additional results within age classes). The interaction between population density and median laying date was negative: when population density increased, the strength of the (positive) relationship between median laying date and adult survival decreased and it almost became negative at very high density.

Figure 1: Relationship between annual survival probabilities and two demographic parameters. A: Survival is positively correlated with median laying date of the population in the previous spring (ß=0.11; SE=0.06). The points are the estimates of Model 3 (Table 2A) for age 1 with 95% confidence intervals, the solid line is for age 1 from Model 36 (Table 2B) which has a continuous effect of median laying date on survival, its 95% CI are represented by the dashed lines. B: Survival is negatively correlated with population density during the previous spring (ß=-0.14; SE=0.07). The points are the estimates of Model 3 (Table 2A) for age 1 with 95% confidence intervals, the solid line is for age 1 from Model 35 (Table 2B) which has a continuous effect of population density on survival, its 95% CI are represented by the dashed lines.

Model	Model Description	np	Deviance	ΔAICc
	Table 2A			
1	**φ (age) P(sex)**	**5**	**2540.29**	**0**
2	φ (sex + age) P(sex)	6	2540.20	1.92
3	φ (age + year) P(sex)	41	2470.48	3.57
4	φ (.) P(sex)	4	2545.74	3.44
5	φ (sex.age) P(sex)	7	2540.11	3.84
6	φ (sex + age + year) P(sex)	42	2470.37	5.54
7	φ (sex) P(sex)	5	2545.72	5.42
8	φ (sex.age + year) P(sex)	43	2470.18	7.41
	…			
18	φ(sex.age + sex.year + age.year) P(sex)	112	2407.88	92.23
	…			
22	**φ(sex.age + sex.year + age.year) P(sex.year)**	**180**	**2305.75**	**143.69**
	Table 2B			
23	**φ(age + popD + medLD + popD*medLD) P(sex)**	**8**	**2525.39**	**0**
24	φ(age + repS + popD + medLD + popD*medLD) P(sex)	9	2523.60	0.22
25	φ(age + repS + popD + repS*popD) P(sex)	8	2527.02	1.63
26	φ(age + repS + popD + medLD + repS*medLD + popD*medLD) P(sex)	10	2523.03	1.66
27	φ(age + repS + popD + medLD + repS*popD + popD*medLD) P(sex)	10	2523.60	2.23
28	φ(age + repS + popD + medLD + repS*popD) P(sex)	9	2526.36	2.98
29	φ(age + repS + popD + LEDmed + repS*popD + repS*medLD + popD*medLD) P(sex)	11	2523.01	3.66
	…			
35	φ(age + popD) P(sex)	6	2535.62	6.19
36	φ(age + medLD) P(sex)	6	2536.86	7.44
37	φ(age + repS) P(sex)	6	2537.07	7.65
	…			
1	**φ(age)P(sex)**	**5**	**2540.29**	**8.86**

Table 2: Monostate model selection assessing the link between population-level phenology and average annual
survival. Full tables are provided in Table A2 of Appendix 2. A: Preliminary model selection assessing the effects
of year, sex and age on recapture (p) and survival (φ) probabilities. Starting model: Model 22, best model:
Model 1. AICc(Model 1) = 2550.3171. B: Secondary model selection assessing the effect of three annual
population-level demographic covariates (repS= reproductive success, popD = population density, medLD =
median laying date) on survival probabilities. The covariates and their interactions were added to preliminary
selected model 1. Starting model: Model 1, best model: Model 23. AICc(Model 23) = 2541.4519. np: number of
parameters

b   Link between individual timing of breeding (relative laying date) and adult survival

The model selection on the state-separated laying date quartiles capture histories dataset
did not show any model stand out in terms of AICc. Seven different models were included
within the first 2 points of AICc of our selection (Table 3). All differ only by their constraints
on survival estimates, differences among phenological quartiles are included in 5 of these
models and the effect of sex is included in only one of them. A model averaging conducted
on these models provided an accurate graphic representation of the estimates they
produced (Figure 2). While little differences occurred for first year adults, survival for older
adults decreased consistently from the first to the last laying date quartile.

Figure 2: Survival probability estimates associated with different laying date quartiles (LD quartile) of the
population obtained by model averaging of the first seven models of our selection (Table 3). For both age
classes, each bar represents a different quartile, lines indicate the 95% confidence intervals of the estimates.

Model	Model Description	np	Deviance	ΔAICc
41	**IS(to.age) φ(age) T(f.to + age) P(sex)**	**23**	**8539.83**	**0**
42	**IS(to.age) φ(f(1 2 3,4) + age) T(f.to + age) P(sex)**	**24**	**8538.58**	**0.79**
43	**IS(to.age) φ(f(1,2 3 4) + age) T(f.to + age) P(sex)**	**24**	**8538.59**	**0.80**
44	**IS(to.age) φ(f(1 2,3 4) + age) T(f.to + age) P(sex)**	**24**	**8539.17**	**1.38**
45	**IS(to.age) φ(f(1,2 3 4).age) T(f.to + age) P(sex)**	**25**	**8537.51**	**1.75**
46	**IS(to.age) φ(sex + age) T(f.to + age) P(sex)**	**24**	**8539.77**	**1.98**
47	**IS(to.age) φ(f(1 4,2 3) + age) T(f.to + age) P(sex)**	**24**	**8539.83**	**2.04**
48	IS(to.age) φ(f(1,2 3,4) + age) T(f.to + age) P(sex)	25	8538.02	2.26
49	IS(to.age) φ(f(1,2 3 4) + sex + age) T(f.to + age) P(sex)	25	8538.54	2.79
	…			
74	IS(to.age) φ(f.age.sex) T(f.to + age) P(sex)	37	8528.26	17.08
	…			
84	IS(to.age) φ(f.age.sex) T(f.to.age.sex) P(sex)	72	8491.61	53.38
85	IS(to.sex.age) φ(f.age.sex) T(f.to.age.sex) P(sex)	78	8483.63	58.10
	…			
101	IS(to.age) φ(f.age.sex) T(.) P(sex)	25	8666.16	130.41
	…			

Table 3: Multistate model selection assessing the link between relative laying date and subsequent adult survival. Full table is provided in Table A3 of Appendix 2. The four states represent laying date quartiles (an individual in state 1 has laid in the earliest 25% of the population). Starting model: Model 85, best model: Model 41. AICc(Model 41) = 8586.2435. For initial state distribution: "to" means that at their first capture, individuals are not evenly distributed among the four phenological quartiles. For between states transition probabilities: "f.to" means that transition probabilities depend on the departure state (f), the arrival state (to) and their interaction. For survival: "f" means that survival depends on the current state of the individual. np: number of parameters.

c   Effects of population-level annual covariates on the relative laying date - adult survival link

Performing a new model selection after adding one meteorological (Temp) and three demographic (medLD, popD and repS) continuous temporal covariates to constrain survival probabilities yielded the same results as in the single state model selection (see a)): the most important temporal covariates were the median laying date, population density and their interaction, all included in the best model (Table 4, model 104).

In all models of this selection, the survival estimates associated with laying date quartile 1 (25% earliest breeders of the population for a given year) were higher than those associated with grouped laying date quartiles 2, 3 and 4 (e.g. Model 43 gives $\phi_{Age1,LD1}$ = 0.628 [0.553; 0.698]$_{95\%CI}$; $\phi_{Age1,LD234}$ = 0.592 [0.541; 0.641] $_{95\%CI}$).

Overall, while adult survival was globally positively correlated with median laying date, we observed differences in the slope of this correlation between early breeders and the other

individuals: early breeders tended to be almost unaffected by the median laying date
(β_{LD1}=0.02; SE=0.13 for early breeders (laying date quartile 1) and β_{LD234}=0.14; SE=0.07 for
others in model 154, Figure 3). The correlation between adult survival and population
density remained almost equal for both categories of laying date (β_{LD1}=-0.19; SE=0.14 and
β_{LD234}=-0.13; SE=0.07 in model 139, Table 4).

Figure 3: Annual survival probabilities as functions of the median laying date of the population for early
breeders (laying date quartile 1, in blue) and for other individuals (laying date quartiles 2, 3 and 4, in red). Filled
circles are the annual survival estimates for age 1 from Model 219 (Table 4), solid lines are the regression
slopes for age 1 of both laying date quartile groups (1 vs 2,3 and 4) from Model 154 (Table 4), their 95%
confidence intervals are represented by the dashed lines.

When looking at the same correlations of annual demographic covariates with survival, but
taking the most important covariates namely median laying date, population density and
their interaction into account at the same time, different effects arose (Figure 4): For all
relative laying date states, at high population density, the positive correlation between
survival and the median laying date still occurred. At low population density, the correlation

became negative. This pattern was qualitatively similar but stronger for early breeders than
for others.

Model	Model Description		T	P	np	Deviance	ΔQAIC
	IS	φ					
103	to.age	(medLD + popD + medLD*popD) + f(1,2 3 4) + age)	f.to + age	sex	27	8523.97	0
104 (Figure 4)	"	**(medLD + popD + medLD*popD).f(1,2 3 4) + age**	"	"	**29**	**8520.13**	**0.25**
105	"	(Temp + medLD + popD + medLD*popD).f(1,2 3 4) + age	"	"	31	8516.64	0.85
106	"	(medLD + popD + repS + medLD*popD).f(1,2 3 4) + age	"	"	31	8518.07	2.28
107	"	(popD + repS + popD*repS).f(1,2 3 4) + age	"	"	29	8522.57	2.69
108	"	(Temp + medLD + popD + repS + medLD*popD).f(1,2 3 4) + age	"	"	33	8515.57	3.89
		…					
139	"	(popD).f(1,2 3 4) + age	"	"	25	8535.01	6.96
		…					
154 (Figure 3)	"	**(medLD).f(1,2 3 4) + age**	"	"	**25**	**8536.04**	**7.99**
		…					
43	"	f(1,2 3 4) + age	"	"	24	8538.59	8.51
		…					
219	"	year.f(1,2 3 4) + age	"	"	97	8406.10	28.90

Table 4: Multistate model selection assessing the effects of population-level annual covariates on the relative laying date - adult survival link. Full table is provided in Table A4 of Appendix 2. The effects of 4 annual covariates (Temp = mean daily mean temperature from March 31[st] to May 7[th], repS = reproductive success, popD = population density, medLD = median laying date) and their first-order interactions as continuous temporal covariates on survival probabilities were tested on one of the best models from the previous selection (Model 43, Table 3). Starting model: Model 43, best model: Model 103. AICc(Model 103) = 8578.5326. For initial state distribution: "to" means that at their first capture, individuals are not evenly distributed among the four phenological quartiles.. For between states transitions probabilities: "f.to" means that transition probabilities depend on the departure state (f), the arrival state (to) and their interaction. For survival: "f" means that survival depends on the current state of the individual. np: number of parameters.

Figure 4: Annual survival probabilities as functions of the median laying date of the population at high (full symbols, solid lines) and low (open symbols, dashed lines) population density, for early breeders (LD quartile 1, blue) and later breeders (LD quartiles 2, 3 and 4, orange). High/low density years are the 15 years of monitoring with highest/lowest population densities. Estimates are from the best model (Model 104, Table 4).

d   Links between individual breeding success, relative laying date and adult survival

When considering combinations of breeding success and relative laying date states (see Figure 5A for a description of the states) on subsequent survival probability, model selection resulted in four models differing from less than two points of AICc (Table 5, Figure 5C), clearly indicating that the most important survival variation within our combined states occurred between individuals failing to reproduce and those that succeed (regardless of their relative laying date, Figure 5D). The average survival estimates were respectively 0.614 [0.562; 0.664]$_{95\%CI}$ for successful breeders and 0.481 [0.387; 0.576]$_{95\%CI}$ for failed breeders (estimates from model 220 in Table 5). While early breeders exhibited slightly higher survival probabilities than late breeders for the two categories of successful breeders (Figure 5C and D), our analysis indicated that the magnitude of this difference was small as compared with the difference in survival between failed and successful breeders.

450   Our analysis also revealed that transitions were asymmetrical with respect to the breeding
451   state: both failed and successful breeders tended to transition preferentially to successful
452   breeding states (model 265, comparison with null model 284 of equal transition
453   probabilities: ΔAICc=138.95, see Figure 5B for transition estimates).

A.

State	Reproductive Success	Relative laying date
1	Failure	Early Breeding (quartile 1)
2	no fledglings	Late Breeding (quartiles 2, 3 and 4)
3	Low Success	Early Breeding (quartile 1)
4	less fledglings than median of the population	Late Breeding (quartiles 2, 3 and 4)
5	High Success	Early Breeding (quartile 1)
6	more fledglings than median of the population	Late Breeding (quartiles 2, 3 and 4)

B. Transition probabilities:

| | to | | | | | |
from	state 1	state 2	state 3	state 4	state 5	state 6
state 1	NA	0.085	0.270	0.370	0.275	NA
state 2	NA	NA	0.116	0.184	0.294	0.272
state 3	0.079	0.067	0.314	0.198	0.224	0.118
state 4	0.023	0.123	0.144	0.384	0.140	0.186
state 5	0.042	0.035	0.244	0.162	0.409	0.108
state 6	0.022	0.096	0.143	0.288	0.232	0.219

C.

Model Number	Model Description	ΔQAICc	Signification of grouped states for Survival	Number of different Survival Estimates	Annual Survival at age 1					
					State 1	State 2	State 3	State 4	State 5	State 6
220	f(12,3456)+age	0	Reproduction Failure, Reproduction Success (Additive age effect)	2	0.481		0.614			
221	f(12,35,46)+age	1.1828	Reproduction Failure, Reproduction Success and Early Breeding, Reproduction Success and Late Breeding	3	0.484		0.639	0.608		
222	f(12,3456).age	1.8208	Reproduction Failure, Reproduction Success (Interactive age effect)	2	0.458		0.618			
223	f(12,34,56)+age	1.8468	Reproduction Failure, Reproduction with Low Success, Reproduction with High Success	3	0.48		0.62		0.605	
233	age	5.7687	No states	1	0.597					
235	from+age	6.7108	No grouping	6	0.465	0.489	0.658	0.61	0.617	0.603
236	f(135,246)+age	6.7556	Early Breeding, Late Breeding	2	0.623	0.591				

D.

Figure 5: Links between individual breeding success, relative laying date and subsequent adult survival. A: Description of the reproductive success and laying date states used in the multistate models. B: Estimates of transition probabilities among breeding success and relative laying date states (model 220, Table 5 and C). For example, the probability of transitioning from state 3 to state 4 is 0.198. NA correspond to transitions that could not be estimated due to insufficient data. C: Models of significant interest from the model selection (Table 5) with the interpretation of the provided estimates. D: Estimates of first year adult survival in the year following a given breeding success and relative laying date. The provided estimates were obtained by averaging the 4 best models of our selection (Table 5), 95%CIs are represented by the grey lines.

Model	Model Description	np	Deviance	ΔAICc
220	*IS(to.age) φ(f(12,3456) + age) T(f.to) P(sex)*	*45*	*9508.25*	*0*
221	*IS(to.age) φ(f(12,35,46) + age) T(f.to) P(sex)*	*46*	*9507.36*	*1.18*
222	*IS(to.age) φ(f(12,3456).age) T(f.to) P(sex)*	*46*	*9508.00*	*1.82*
223	*IS(to.age) φ(f(12,34,56) + age) T(f.to) P(sex)*	*46*	*9508.03*	*1.85*
224	IS(to.age) φ(f(12,3456) + sex + age) T(f.to) P(sex)	46	9508.19	2.01
225	IS(to.age) φ(f(12,3456).age + sex) T(f.to) P(sex)	47	9507.96	3.84
226	IS(to.age) φ(f(12,34,56) + sex + age) T(f.to) P(sex)	47	9507.98	3.86
227	IS(to.age) φ(f(12,3456).sex + age) T(f.to) P(sex)	47	9507.98	3.87
228	IS(to.age) φ(f(12,3456) + sex.age) T(f.to) P(sex)	47	9508.16	4.05
229	IS(to.age) φ(f(12,3456)) T(f.to) P(sex)	44	9514.23	3.91
	...			
233	*IS(to.age) φ(age) T(f.to) P(sex)*	*44*	*9516.09*	*5.77*
	...			
235	*IS(to.age) φ(from + age) T(f.to) P(sex)*	*49*	*9506.67*	*6.71*
236	*IS(to.age) φ(f(135,246) + age) T(f.to) P(sex)*	*45*	*9515.01*	*6.76*
	...			
265	IS(to.age) φ(f.age.sex) T(f.to) P(sex)	66	9498.25	33.79
	...			
274	IS(to.age) φ(f.age.sex) T(f.to.age.sex) P(sex)	153	9414.88	139.55
	...			
276	IS(to.sex.age) φ(f.age.sex) T(f.to.age.sex) P(sex)	163	9408.55	155.79
	...			
284	IS(to.age) φ(f.age.sex) T(.) P(sex)	37	9697.50	172.74
	...			

Table 5: Multistate model selection assessing the links between individual breeding success, relative laying date
and adult survival. Full table is provided in Table A5 of Appendix 2. The 6 states represent different
combinations of individual breeding success and relative laying date (see Figure 5 for details). Starting model:
Model 276 included the interactive effects of age, sex and state for its various structural components.
Recapture probabilities were assumed to depend on sex only. Sequential selection procedure was the same as
in previous model selection (Table 4). Best model: Model 220. AICc(Model 220) = 9599.8097. np: number of
parameters.

Discussion

The analysis conducted on the links between reproductive phenology and subsequent adult
survival in a Mediterranean forest Blue tit (*Cyanistes caeruleus*) population revealed that (a)
at the population scale, early median breeding years are followed by lower average adult
survival, (b) at the individual level, earlier breeders within the population have higher
subsequent survival than late breeders, (c) population density influences the relative
phenology ↔ survival relationship, (d) individual relative reproductive success is also
related to subsequent adult survival in our population, in particular, failed reproduction is
linked to lower subsequent survival.

a   At the population scale, early breeding years are followed by low average survival

We showed that at the population level, advanced phenology is associated with lower
overall subsequent survival in breeders (Table 2, Figure 1A). Although the link between avian
breeding phenology and survival has not been examined nearly as much as the link between
phenology and reproductive success, this first result is in agreement with previous results in
birds (Brinkhof et al., 2002; Nilsson, 1994). Such inter-annual, population-level variation in
phenology reflects the average, or common response of breeders in the year t to
environmental conditions, including spring temperatures (Bonamour et al., 2019). Similarly,
reduced survival between year t and t+1 reflects the average demographic response of
breeders to these conditions, regardless of their relative position in the laying date
distribution of year t. Reduced average survival may be the consequence of a high energetic
cost of early breeding itself, in addition to the cost of high reproduction associated with early
breeding (Brinkhof et al., 2002). These results are compatible with our preliminary
hypothesis of the importance of a cost of reproduction in our population, as well as with the
general prediction that, in short-lived species, costs of reproduction mainly occur on survival
(rather than on future reproduction, Hamel et al., 2010; Le Coeur et al., 2018). However, it is
important to note that the correlative nature of our analysis prevents us from demonstrating
fully this hypothesis. We could be observing, for example, the mortality consequences of a
high hydric or energetic stress in warm years, or of high adult predation (due to favourable
meteorological conditions for predators) in these years of early avian phenology. Another
explanation for the observed pattern could be the existence of phenological mismatches: in
particularly warm years, the advance of phenology of the tit population could be insufficient
compared to the advance of their prey (Visser and Gienapp, 2019). However, such predator-
prey mismatch would also result in a lower reproduction for blue tits, which is not what is
generally observed for early years: on the contrary, in the E-Pirio population, the mean
number of fledglings per attempted brood increased when the median laying date decreased
(+0.12 fledglings/earlier day, *p-value* < 10^{-3}).

b   At the individual level, relative early breeders have higher survival the following year

Interestingly, when examining the relationship between survival and relative laying date (i.e.,
the position of the focal individual within the laying date distribution in a given year), the
opposite result was obtained compared to the population scale. Regardless of the median
laying date of the population, individuals that breed early in the population have higher
subsequent survival than others (for ages of two years or more) (Figure 2). This result
suggests that some individual heterogeneity (Gimenez et al., 2018) occurs within our study
population: while breeding early can be seen as a strategy to improve breeding success
(Marrot et al., 2018) that may convey some survival cost (as suggested by our present
population-level results), early breeders may be more prone to survive in spite of this cost.
Similar results have been obtained by previous studies on the cost of reproduction: while
reproduction is expected to be costly, positive covariation between breeding success and
subsequent survival and other 'big house big car' correlations are frequently observed in
free ranging animal populations (van Noordwijk and de Jong, 1986; Reznick et al., 2000;
Robert et al., 2015). In our blue tit population, the high repeatability of the rank of laying
date among years (R = 0.406 [0.316; 0.493]$_{95\%CI}$ for females, see Appendix 4 for details)
suggests indeed the occurrence of some intrapopulation heterogeneity regarding the
phenological strategy. This result is confirmed by the analysis of the transitions among states
of relative laying date, which are not random (comparison of model 74 and model 101,
ΔAICc = 113.33, Table 3). In particular, the earliest breeders (quartile 1 of the laying date
distribution at year t) have a very high probability (52%) to remain in quartile 1 at year t+1
(see transition probabilities matrix in Appendix 5). These results are in line with previous
findings in many bird species that breeding phenology is repeatable and heritable (see e.g.
Postma, 2014; for the focal population Delahaie et al., 2017 estimated a heritability of laying
date of 0.125 [0.059; 0.199]$_{95\%CI}$).

When considering both the absolute (median of the population) and the relative laying date,
we found that the strength of the correlation between survival and the median laying date
varies between early breeders and later breeders (Figure 3). Early breeders, on top of having
higher survival rates, are less affected by the "quality" of the year (warm or cold spring
resulting in early or late median laying date) than late breeders. This result confirms that
earlier breeders might be high resource individuals able to secure the earliest and best
territories while also having high somatic maintenance.

c   Population density influences the relative phenology ↔ survival relationship

Population density was another important factor (negatively) correlated with annual
subsequent survival (Figure 1B), presumably reflecting the intensity of intraspecific
competition for food resources (Gamelon et al., 2016). We thus included density in the
multistate individual model to test whether population breeding density modulated the
relationship between phenology and survival probability. Results show that at high breeding
population density the suspected potential trade-off between early reproduction and
subsequent survival becomes apparent (Figure 4): this suggests that the unfavourable

environment of high intra-population competition increases the costs of early breeding. On
the contrary, at low population density, the constraints of the trade-off seem to be relaxed:
no cost in survival of breeding early can be observed, and in warm years (early median laying
date), on top of having better reproduction (Delahaie et al., 2017), early breeding individuals
have higher survival rates (Figure 4).

Two main conclusions can be drawn from these results. First, the cost of breeding early is
modulated by environmental conditions. Although, to our knowledge, it has never been
applied to phenology, this finding is in agreement with previous results showing that costs of
reproduction may be modulated by age (Rauset et al., 2015), resource availability (Le Coeur
et al., 2018) or meteorological conditions (Stoelting et al., 2014). Second, while the cost of
breeding earlier than the other breeders in a given year may be hidden by demographic
heterogeneity under average conditions (Figure 2), it becomes apparent under very harsh
conditions (here, a combination of warm spring and high population density, which are both
correlated to low adult survival). Under such harsh conditions, individuals that invest in early
breeding show particularly low survival probabilities (Figure 4). Again, similar interplays
between demographic heterogeneity, reproduction cost and environmental conditions have
been found in previous studies (Robert et al., 2012). For example, in the Alpine ibex (Capra
ibex), survival probabilities were found higher for successfully breeding females than for
unsuccessful females under average conditions. However, during a severe disease outbreak,
survival patterns according to reproductive state were reversed, with lower survival for
successful than unsuccessful females (Garnier et al., 2016).

590
591   d   Both individual laying date and individual breeding success are related to survival
592

Life-history theory predicts that, at the scale of organisms, investment in current
reproduction may constrain the energy available for maintenance, leading to trade-offs with
survival and/or future reproduction (Roff, 1993; Stearns, 1992) referred to as the cost of
reproduction. In the context of our blue tit population, if early breeding is a strategy for
breeders to improve their reproductive success, the survival cost associated with early
reproduction can be considered a form of reproductive cost. However, it is notoriously
difficult to disentangle the cost of early reproduction *per se* from the cost associated with
high reproductive success. Previous analyses in our study population indicate that early
breeders are generally more successful than late breeders (Porlier et al., 2012), which
implies that the effects of breeding success and laying date on survival are partially
confounded. When considering combinations of reproductive success and relative laying
date in the survival models, results uncover strong differences in survival probabilities
between individuals with reproduction success (at least one fledgling) or failure (Figure 5).
All individuals of our datasets were captured in the nest boxes during reproduction: they
have all at least attempted reproduction and reached the nestling stage, i.e. successfully
hatched at least one nestling. This means that individuals with failed reproduction (0
fledglings) encountered strong impediments during the breeding event (scarceness of

resources, harsh climatic events, predation of the brood, abandonment by one or the other
parent, or death).

A high proportion of these brood failures may be linked to the death of one or both parents:
as it occurs after the recapture and identification of the parents (otherwise they would not
be in the dataset), it will be analysed as death during the subsequent year and therefore we
expected to observe low survival for adults with failed reproduction compared to others.
However, the positive effect of being an early breeder is still observed here, on top of the
effects of reproductive success. Model selection indicates that (1) models assuming survival
differences between early and late breeders are part of the set of best models (ΔAICc<2,
Table 5) (2) among breeders that fledged at least one chick, the relative laying date explains
more deviance than the breeding success (model 221 vs. 223, Table 5).

e Evolutionary perspectives in a context of climate change

Previous selection analyses using reproductive success as fitness estimator concluded that
early breeding individuals are favoured by selection (e.g. Klomp, 1970; Marrot et al., 2018 in
the focal population). As laying date is a heritable trait (Delahaie et al., 2017; Postma, 2014),
and as it stands under negative selection, many studies using reproductive success were
surprised by the absence of evolution of laying date in the context of climate change (e.g.
Price et al., 1988). An often suggested explanation to this evolutionary stasis is the potential
presence of a positive selection on laying date via survival, opposed to and compensating the
negative selection via reproductive success (Merilä et al., 2001). Here, we showed that this
cannot be the case in our population, as early breeding individuals have not only higher
reproductive success, but also higher survival, which only increases the selection for early
breeding.

Another explanation behind common evolutionary stasis in heritable selected traits is the
spatio-temporal fluctuations of selection (e.g. Miehls et al., 2015). In our study population,
the only case in which selection on survival was inversed was in harsh condition years (early
median laying date and high population density), whereby later laying individuals had higher
survival rates: If the frequency of such years were to increase in the context of climate
change, the selection regime of the focal population might face significant changes.

Acknowledgements

We are grateful to Emmanuelle Cam, Aurélie Cohas, Olivier Gimenez, François Chiron, François Sarrazin, Roger Pradel and Pierre-Yves Henry for their constructive comments on an earlier version of this manuscript. We thank all the people who helped conduct the blue tit monitoring in the past four decades, in particular Jacques Blondel, Phillippe Perret, Marcel Lambrechts, Claire Doutrelant, Denis Réale, Samuel Caro, and Christophe de Franceschi (who also managed the database). We thank the APEEM, the ONF and Achille Sanroma for field logistics, and the Fango valley MAB reserve. The long term project has been supported by the OSU-OREME since 2009.

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content of this article.

References

References

Abadi, F., Gimenez, O., Jakober, H., Stauber, W., Arlettaz, R., Schaub, M., 2012. Estimating
the strength of density dependence in the presence of observation errors using
integrated population models. Ecol. Mod. 242, 1-9.
https://doi.org/10.1016/j.ecolmodel.2012.05.007

Bjorkman, A.D., Elmendorf, S.C., Beamish, A.L., Vellend, M., Henry, G.H.R., 2015. Contrasting
effects of warming and increased snowfall on Arctic tundra plant phenology over the
past two decades. Glob. Change Biol. 21, 4651–4661.
https://doi.org/10.1111/gcb.13051

Blondel, J., Thomas, D.W., Charmantier, A., Perret, P., Bourgault, P., Lambrechts, M.M.,
2006. A Thirty-Year Study of Phenotypic and Genetic Variation of Blue Tits in
Mediterranean Habitat Mosaics. BioScience 56, 661. https://doi.org/10.1641/0006-
3568(2006)56[661:ATSOPA]2.0.CO;2

Bonamour, S., Chevin Luis-Miguel, Charmantier Anne, Teplitsky Céline, 2019. Phenotypic
plasticity in response to climate change: the importance of cue variation. Philos. Trans.
R. Soc. B Biol. Sci. 374, 20180178. https://doi.org/10.1098/rstb.2018.0178

Bonenfant, C., Gaillard, J.M., Coulson, T., Festa-Bianchet, M., Loison, A., Garel, M., Loe, L.E.,
Blanchard, P., Pettorelli, N., Owen-Smith, N., Du Toit, J., 2009. Empirical evidence of
density-dependence in populations of large herbivores. Advances in ecological
research, 41, pp.313-357. https://doi.org/10.1016/S0065-2504(09)00405-X

Brinkhof, M.W.G., Cavé, A.J., Daan, S., Perdeck, A.C., 2002. Timing of Current Reproduction
Directly Affects Future Reproductive Output in European Coots. Evolution 56, 400–411.
https://doi.org/10.1111/j.0014-3820.2002.tb01349.x

Brouwer, L., Richardson, D. S., Eikenaar, C., & Komdeur, J., 2006. The role of group size and
environmental factors on survival in a cooperatively breeding tropical passerine. J.
Anim. Ecol. 75, 1321-1329.

Burnham, K.P., Anderson, D.R., 1998. Practical Use of the Information-Theoretic Approach,
in: Burnham, K.P., Anderson, D.R. (Eds.), Model Selection and Inference: A Practical
Information-Theoretic Approach. Springer New York, New York, NY, pp. 75–117.
https://doi.org/10.1007/978-1-4757-2917-7_3

Cam, E., Gimenez, O., Alpizar-Jara, R., Aubry, L.M., Authier, M., Cooch, E.G., Koons, D.N.,
Link, W.A., Monnat, J.-Y., Nichols, J.D., Rotella, J.J., Royle, J.A., Pradel, R., 2013. Looking
for a needle in a haystack: inference about individual fitness components in a
heterogeneous population. Oikos 122, 739–753. https://doi.org/10.1111/j.1600-
0706.2012.20532.x

Cam, E., Link, W.A., Cooch, E.G., Monnat, J., Danchin, E., 2002. Individual Covariation in Life-History Traits: Seeing the Trees Despite the Forest. Am. Nat. 159, 96–105. https://doi.org/10.1086/324126

Charmantier, A., Doutrelant, C., Dubuc-Messier, G., Fargevieille, A., Szulkin, M., 2016. Mediterranean blue tits as a case study of local adaptation. Evol. Appl. 9, 135–152. https://doi.org/10.1111/eva.12282

Charmantier, A., Gienapp, P., 2014. Climate change and timing of avian breeding and migration: evolutionary versus plastic changes. Evol. Appl. 7, 15–28. https://doi.org/10.1111/eva.12126

Choquet, R., Lebreton, J.-D., Gimenez, O., Reboulet, A.-M., Pradel, R., 2009a. U-CARE: Utilities for performing goodness of fit tests and manipulating CApture–REcapture data. Ecography 32, 1071–1074. https://doi.org/10.1111/j.1600-0587.2009.05968.x

Choquet, R., Rouan, L., Pradel, R., 2009b. Program E-Surge: A Software Application for Fitting Multievent Models, in: Thomson, D.L., Cooch, E.G., Conroy, M.J. (Eds.), Modeling Demographic Processes In Marked Populations, Environmental and Ecological Statistics. Springer US, Boston, MA, pp. 845–865. https://doi.org/10.1007/978-0-387-78151-8_39

Clutton-Brock, T.H., 1988. Reproductive Success: Studies of Individual Variation in Contrasting Breeding Systems. University of Chicago Press.

Delahaie, B., Charmantier, A., Chantepie, S., Garant, D., Porlier, M., Teplitsky, C., 2017. Conserved G-matrices of morphological and life-history traits among continental and island blue tit populations. Heredity 119, 76–87. https://doi.org/10.1038/hdy.2017.15

Diekmann, M., 1996. Relationship between flowering phenology of perennial herbs and meteorological data in deciduous forests of Sweden. Can. J. Bot. 74, 528–537. https://doi.org/10.1139/b96-067

Diggle, P.K., 1999. Heteroblasty and the Evolution of Flowering Phenologies. Int. J. Plant Sci. 160, S123–S134. https://doi.org/10.1086/314217

Dijkstra, C., Daan, S., Tinbergen, J.M., 1990. Family Planning in the Kestrel (Falco Tinnunculus): the Ultimate Control of Covariation of Laying Date and Clutch Size. Behaviour 114, 83–116. https://doi.org/10.1163/156853990X00068

Durant, J.M., Hjermann, D.Ø., Ottersen, G., Stenseth, N.C., 2007. Climate and the match or mismatch between predator requirements and resource availability. Clim. Res. 33, 271–283. https://doi.org/10.3354/cr033271

Duursma, D.E., Gallagher, R.V., Griffith, S.C., 2018. Effects of El Niño Southern Oscillation on avian breeding phenology. Divers. Distrib. 24, 1061–1071. https://doi.org/10.1111/ddi.12750

Elzinga, J.A., Atlan, A., Biere, A., Gigord, L., Weis, A.E., Bernasconi, G., 2007. Time after time:
flowering phenology and biotic interactions. Trends Ecol. Evol. 22, 432–439.
https://doi.org/10.1016/j.tree.2007.05.006

Fay, R., Weimerskirch, H., Delord, K., Barbraud, C., 2015. Population density and climate
shape early-life survival and recruitment in a long-lived pelagic seabird. J. Anim. Ecol.
84, 1423–1433. https://doi.org/10.1111/1365-2656.12390

Forrest, J., Miller-Rushing, A.J., 2010. Toward a synthetic understanding of the role of
phenology in ecology and evolution. Philos. Trans. R. Soc. B Biol. Sci. 365, 3101–3112.
https://doi.org/10.1098/rstb.2010.0145

Gamelon, M., Grøtan, V., Engen, S., Bjørkvoll, E., Visser, M.E., Sæther, B.-E., 2016. Density
dependence in an age-structured population of great tits: identifying the critical age
classes. Ecology 97, 2479–2490. https://doi.org/10.1002/ecy.1442

Garnier, A., Gaillard, J.-M., Gauthier, D., Besnard, A., 2016. What shapes fitness costs of
reproduction in long-lived iteroparous species? A case study on the Alpine ibex.
Ecology 97, 205–214. https://doi.org/10.1890/15-0014.1

Gimenez, O., Cam, E., Gaillard, J.-M., 2018. Individual heterogeneity and capture-recapture
models: what, why and how? Oikos 127, 664–686. https://doi.org/10.1111/oik.04532

Grosbois, V., Henry, P.-Y., Blondel, J., Perret, P., Lebreton, J.-D., Thomas, D.W., Lambrechts,
M.M., 2006. Climate impacts on Mediterranean blue tit survival: an investigation
across seasons and spatial scales. Glob. Change Biol. 12, 2235–2249.
https://doi.org/10.1111/j.1365-2486.2006.01286.x

Grøtan, V., Sæther, B. E., Engen, S., Van Balen, J. H., Perdeck, A. C., & Visser, M. E., 2009.
Spatial and temporal variation in the relative contribution of density dependence,
climate variation and migration to fluctuations in the size of great tit populations. J.
Anim. Ecol. 78, 447-459.

Haest, B., Hüppop, O., Bairlein, F., 2018. Challenging a 15-year-old claim: The North Atlantic
Oscillation index as a predictor of spring migration phenology of birds. Glob. Chang.
Biol. 24(4), 1523-1537. DOI: 10.1111/gcb.14023

Hamel, S., Côté, S.D., Gaillard, J.-M., Festa-Bianchet, M., 2009. Individual variation in
reproductive costs of reproduction: high-quality females always do better. J. Anim.
Ecol. 78, 143–151. https://doi.org/10.1111/j.1365-2656.2008.01459.x

Hamel, S., Gaillard, J.-M., Yoccoz, N.G., Loison, A., Bonenfant, C., Descamps, S., 2010. Fitness
costs of reproduction depend on life speed: empirical evidence from mammalian
populations. Ecol. Lett. 13, 915–935. https://doi.org/10.1111/j.1461-
0248.2010.01478.x

Harriman, V. B., Dawson, R. D., Bortolotti, L. E., & Clark, R. G., 2017. Seasonal patterns in
reproductive success of temperate-breeding birds: Experimental tests of the date and
quality hypotheses. Ecol. evol. 7, 2122-2132

Harshman, L.G., Zera, A.J., 2007. The cost of reproduction: the devil in the details. Trends
Ecol. Evol. 22, 80–86. https://doi.org/10.1016/j.tree.2006.10.008

Klomp, H., 1970. The Determination of Clutch-Size in Birds a Review. Ardea 55, 1–124.
https://doi.org/10.5253/arde.v58.p1

Laaksonen, T., Ahola, M., Eeva, T., Väisänen, R.A., Lehikoinen, E., 2006. Climate change,
migratory connectivity and changes in laying date and clutch size of the pied
flycatcher. Oikos 114, 277–290. https://doi.org/10.1111/j.2006.0030-1299.14652.x

Le Cœur, C., Chantepie, S., Pisanu, B., Chapuis, J.-L., Robert, A., 2016. Inter-annual and inter-
individual variations in survival exhibit strong seasonality in a hibernating rodent.
Oecologia 181, 795–807. https://doi.org/10.1007/s00442-016-3597-2

Le Coeur, C., Pisanu, B., Chapuis, J.-L., Robert, A., 2018. Within- and between-year variations
of reproductive strategy and cost in a population of Siberian chipmunks. Oecologia
188, 765–776. https://doi.org/10.1007/s00442-018-4259-3

Lebreton, J.-D., Burnham, K.P., Clobert, J., Anderson, D.R., 1992. Modeling Survival and
Testing Biological Hypotheses Using Marked Animals: A Unified Approach with Case
Studies. Ecol. Monogr. 62, 67–118. https://doi.org/10.2307/2937171

Lebreton, J.D., Hines, J.E., Pradel, R., Nichols, J.D., Spendelow, J.A., 2003. Estimation by
capture-recapture of recruitment and dispersal over several sites. Oikos 101, 253–264.
https://doi.org/10.1034/j.1600-0706.2003.11848.x

Lescroël, A., Dugger, K.M., Ballard, G., Ainley, D.G., 2009. Effects of individual quality,
reproductive success and environmental variability on survival of a long-lived seabird.
J. Anim. Ecol. 78, 798–806. https://doi.org/10.1111/j.1365-2656.2009.01542.x

Lindström, J., 1999. Early development and fitness in birds and mammals. Trends Ecol. Evol.
14, 343–348. https://doi.org/10.1016/S0169-5347(99)01639-0

Marrot, P., Charmantier, A., Blondel, J., Garant, D., 2018. Current Spring Warming as a
Driver of Selection on Reproductive Timing in a Wild Passerine. J. Anim. Ecol. 87, 754–
764. https://doi.org/10.1111/1365-2656.12794

Merilä, J., Sheldon, B.C., Kruuk, L.E.B., 2001. Explaining stasis: microevolutionary studies in
natural populations. Genetica 112, 199–222.
https://doi.org/10.1023/A:1013391806317

Miehls, A. L. J., Peacor, S. D., Valliant, L., McAdam, A. G., 2015. Evolutionary stasis despite
selection on a heritable trait in an invasive zooplankton. J. Evol. Biol. 28(5), 1091-1102.
DOI: 10.1111/jeb.12632

Newton, I., 1994. The role of nest sites in limiting the numbers of hole-nesting birds: A
review. Biol. Conserv. 70, 265–276. https://doi.org/10.1016/0006-3207(94)90172-4

Nilsson, J.-Å., 1994. Energetic Bottle-Necks During Breeding and the Reproductive Cost of
Being Too Early. J. Anim. Ecol. 63, 200–208. https://doi.org/10.2307/5595

Nilsson, J.-Å., Svensson, E., 1996. The cost of reproduction: a new link between current
reproductive effort and future reproductive success. Proc. R. Soc. Lond. B Biol. Sci. 263,
711–714. https://doi.org/10.1098/rspb.1996.0106

Nur, N., 1984. The Consequences of Brood Size for Breeding Blue Tits I. Adult Survival,
Weight Change and the Cost of Reproduction. J. Anim. Ecol. 53, 479–496.
https://doi.org/10.2307/4529

Parmesan, C., 2007. Influences of species, latitudes and methodologies on estimates of
phenological response to global warming. Glob. Change Biol. 13, 1860–1872.
https://doi.org/10.1111/j.1365-2486.2007.01404.x

Porlier, M., Charmantier, A., Bourgault, P., Perret, P., Blondel, J., Garant, D., 2012. Variation
in phenotypic plasticity and selection patterns in blue tit breeding time: between- and
within-population comparisons: Variation in plasticity among populations. J. Anim.
Ecol. 81, 1041–1051. https://doi.org/10.1111/j.1365-2656.2012.01996.x

Postma, E., 2014. Four decades of estimating heritabilities in wild vertebrate populations:
improved methods, more data, better estimates?, in: Charmantier A, Garant D, Kruuk
LEB (Eds). Quantitative Genetics in the Wild. Oxford University Press: Oxford, UK, p. pp
16–33.

Price, T., Kirkpatrick, M., Arnold, S.J., 1988. Directional selection and the evolution of
breeding date in birds. Science 240, 798–799.
https://doi.org/10.1126/science.3363360

Radchuk, V., Reed, T., Teplitsky, C., Van De Pol, M., Charmantier, A., Hassall, C., Adamík, P.,
Adriaensen, F., Ahola, M.P., Arcese, P., Avilés, J.M., 2019. Adaptive responses of
animals to climate change are most likely insufficient. Nat. Com., 10(1), 1-14.
https://doi.org/10.1038/s41467-019-10924-4

Rauset, G.R., Low, M., Persson, J., 2015. Reproductive patterns result from age-related
sensitivity to resources and reproductive costs in a mammalian carnivore. Ecology 96,
3153–3164. https://doi.org/10.1890/15-0262.1

Reed, T.E., Grotan, V., Jenouvrier, S., Saether, B.-E., Visser, M.E., 2013. Population Growth in
a Wild Bird Is Buffered Against Phenological Mismatch. Science 340, 488–491.
https://doi.org/10.1126/science.1232870

Reznick, D., Nunney, L., Tessier, A., 2000. Big houses, big cars, superfleas and the costs of
reproduction. Tr. Ecol. Evol. 15, 421-425. https://doi.org/10.1016/S0169-
5347(00)01941-8

Robb, G.N., McDonald Robbie A, Chamberlain Dan E, Reynolds S. James, Harrison Timothy
J.E, Bearhop Stuart, 2008. Winter feeding of birds increases productivity in the
subsequent breeding season. Biol. Lett. 4, 220–223.
https://doi.org/10.1098/rsbl.2007.0622

Robert, A., Bolton, M., Jiguet, F., Bried, J., 2015. The survival-reproduction association
becomes stronger when conditions are good. Proc. R. Soc. Lond. B Biol. Sci. 282,
20151529. https://doi.org/10.1098/rspb.2015.1529

Robert, A., Paiva, V.H., Bolton, M., Jiguet, F., Bried, J., 2012. The interaction between
reproductive cost and individual quality is mediated by oceanic conditions in a long-
lived bird. Ecology 93, 1944–1952. https://doi.org/10.1890/11-1840.1

Roff, D., 1993. Evolution Of Life Histories: Theory and Analysis. Springer Science & Business
Media.

Samplonius, J.M., Bartošová, L., Burgess, M.D., Bushuev, A.V., Eeva, T., Ivankina, E.V.,
Kerimov, A.B., Krams, I., Laaksonen, T., Mägi, M., Mänd, R., Potti, J., Török, J., Trnka,
M., Visser, M.E., Zang, H., Both, C., 2018. Phenological sensitivity to climate change is
higher in resident than in migrant bird populations among European cavity breeders.
Glob. Change Biol. 24, 3780–3790. https://doi.org/10.1111/gcb.14160

Sheriff, M.J., Buck, C.L., Barnes, B.M., 2015. Autumn conditions as a driver of spring
phenology in a free-living arctic mammal. Clim. Change Responses 2, 4.
https://doi.org/10.1186/s40665-015-0012-x

Shipley, J. R., Twining, C. W., Taff, C. C., Vitousek, M. N., Flack, A., Winkler, D. W., 2020. Birds
advancing lay dates with warming springs face greater risk of chick mortality. Proc.
Nat. Ac. Sc. 117(41), 25590-25594. DOI: 10.1073/pnas.2009864117

Simmonds, E. G., Cole, E. F., Sheldon, B. C., Coulson, T., 2020. Phenological asynchrony: a
ticking time-bomb for seemingly stable populations?. Ecol. Lett., 23(12), 1766-1775.
https://doi.org/10.1111/ele.13603

Stearns, S.C., 1992. The Evolution of Life Histories. Oxford University Press, Oxford, New
York.

Stoelting, R.E., Gutiérrez, R.J., Kendall, W.L., Peery, M.Z., 2014. Life-history tradeoffs and
reproductive cycles in Spotted OwlsCompromisos en la historia de vida de Strix
occidentalisLife-history tradeoffs in Spotted Owls. The Auk 132, 46–64.
https://doi.org/10.1642/AUK-14-98.1

Tarwater, C.E., Arcese, P., 2017. Young females pay higher costs of reproduction in a short-
lived bird. Behav. Ecol. Sociobiol. 71, 84. https://doi.org/10.1007/s00265-017-2309-1

Thackeray, S.J., Henrys, P.A., Hemming, D., Bell, J.R., Botham, M.S., Burthe, S., Helaouet, P.,
Johns, D.G., Jones, I.D., Leech, D.I., Mackay, E.B., Massimino, D., Atkinson, S., Bacon,
P.J., Brereton, T.M., Carvalho, L., Clutton-Brock, T.H., Duck, C., Edwards, M., Elliott,

J.M., Hall, S.J.G., Harrington, R., Pearce-Higgins, J.W., Høye, T.T., Kruuk, L.E.B.,
Pemberton, J.M., Sparks, T.H., Thompson, P.M., White, I., Winfield, I.J., Wanless, S.,
2016. Phenological sensitivity to climate across taxa and trophic levels. Nature 535,
241–245. https://doi.org/10.1038/nature18608

Tuljapurkar, S., Steiner, U.K., Orzack, S.H., 2009. Dynamic heterogeneity in life histories.
Ecol. Lett. 12, 93–106. https://doi.org/10.1111/j.1461-0248.2008.01262.x

van Noordwijk, A.J., de Jong, G., 1986. Acquisition and Allocation of Resources: Their
Influence on Variation in Life History Tactics. Am. Nat. 128, 137–142.
https://doi.org/10.1086/284547

Verhulst, S., Nilsson, J.-Å., 2008. The timing of birds' breeding seasons: a review of
experiments that manipulated timing of breeding. Philos. Trans. R. Soc. B Biol. Sci. 363,
399–410. https://doi.org/10.1098/rstb.2007.2146

Visser, M.E., Gienapp, P., 2019. Evolutionary and demographic consequences of
phenological mismatches. Nat. Ecol. Evol. 3, 879. https://doi.org/10.1038/s41559-019-
0880-8

Visser, M.E., te Marvelde, L., Lof, M.E., 2012. Adaptive phenological mismatches of birds and
their food in a warming world. J. Ornithol. 153, 75–84.
https://doi.org/10.1007/s10336-011-0770-6

Wiggins, D.A., Pärt, T., Gustafsson, L., 1998. Timing of Breeding and Reproductive Costs in
Collared Flycatchers. The Auk 115, 1063–1067. https://doi.org/10.2307/4089524

Wilson, A.J., Nussey, D.H., 2010. What is individual quality? An evolutionary perspective.
Trends Ecol. Evol. 25, 207–214. https://doi.org/10.1016/j.tree.2009.10.002

Appendix 1: Goodness-of-fit

No significant violation of standard models assumptions was found while performing the goodness-of-fit tests using program U-CARE (Choquet et al., 2009): for the monostate dataset $\chi^2(df=184)=85.34$, P-level=1; for the multistate dataset with relative laying date states: $\chi^2(df=98)=21.289$ for group 1, $\chi^2(df=131)=26.885$ for group 2, $\chi^2(df=57)=14.276$ for group 3, $\chi^2(df=36)=2.808$ for group 4, p-value=1 for all groups; for the multistate dataset with breeding success and relative laying date states: $\chi^2(df=83)=21.318$ for group 1, $\chi^2(df=132)=36.779$ for group 2, $\chi^2(df=51)=13.323$ for group 3, $\chi^2(df=30)=4.530$ for group 4, p-value=1 for all groups.

References

Choquet, R., Lebreton, J.-D., Gimenez, O., Reboulet, A.-M., Pradel, R., 2009. U-CARE: Utilities for performing goodness of fit tests and manipulating CApture–REcapture data. Ecography 32, 1071–1074. https://doi.org/10.1111/j.1600-0587.2009.05968.x

Appendix 2: Full model selection tables (Tables A2-A5)

TABLE A2 A

Model	Model Description	np	Deviance	ΔAICc
1	**ɸ (age) P(sex)**	**5**	**2540.29**	**0**
2	ɸ (sex + age) P(sex)	6	2540.20	1.92
3	ɸ (age + year) P(sex)	41	2470.48	3.57
4	ɸ (.) P(sex)	4	2545.74	3.44
5	ɸ (sex.age) P(sex)	7	2540.11	3.84
6	ɸ (sex + age + year) P(sex)	42	2470.37	5.54
7	ɸ (sex) P(sex)	5	2545.72	5.43
8	ɸ (sex.age + year) P(sex)	43	2470.18	7.41
9	ɸ (year) P(sex)	40	2476.49	7.52
10	ɸ (sex + year) P(sex)	41	2476.47	9.56
11	ɸ (age.year) P(sex)	74	2440.68	42.97
12	ɸ (age.year + sex) P(sex)	75	2440.63	45.04
13	ɸ (sex.age + age.year) P(sex)	76	2440.60	47.14
14	ɸ (sex.year + age) P(sex)	78	2439.65	50.45
15	ɸ (sex.age + sex.year) P(sex)	79	2439.22	52.15
16	ɸ (sex.year) P(sex)	77	2444.39	53.06
17	ɸ (sex.year + age.year) P(sex)	111	2408.41	90.56
18	**ɸ(sex.age + sex.year + age.year) P(sex)**	**112**	**2407.88**	**92.23**
19	ɸ(sex.age + sex.year + age.year) P(sex + year)	147	2337.96	100.25
20	ɸ(sex.age + sex.year + age.year) P(.)	111	2427.16	109.31
21	ɸ(sex.age + sex.year + age.year) P(year)	146	2361.01	121.03
22	**ɸ(sex.age + sex.year + age.year) P(sex.year)**	**180**	**2305.75**	**143.69**

TABLE A2 B

Model	Model Description	np	Deviance	ΔAICc
23	**φ(age + popD + medLD + popD*medLD) P(sex)**	**8**	**2525.39**	**0**
24	φ(age + repS + popD + medLD + popD*medLD) P(sex)	9	2523.60	0.22
25	φ(age + repS + popD + repS*popD) P(sex)	8	2527.02	1.63
26	φ(age + repS + popD + medLD + repS*medLD + popD*medLD) P(sex)	10	2523.03	1.67
27	φ(age + repS + popD + medLD + repS*popD + popD*medLD) P(sex)	10	2523.60	2.24
28	φ(age + repS + popD + medLD + repS*popD) P(sex)	9	2526.36	2.98
29	φ(age + repS + popD + LEDmed + repS*popD + repS*medLD + popD*medLD) P(sex)	11	2523.01	3.67
30	φ(age + popD + medLD) P(sex)	7	2531.07	3.66
31	φ(age + repS + popD) P(sex)	7	2531.11	3.70
32	φ(age + repS + popD + medLD) P(sex)	8	2529.77	4.38
33	φ(age + repS + popD + medLD + repS*popD + repS*medLD) P(sex)	10	2526.35	4.98
34	φ(age + repS + popD + medLD + repS*medLD) P(sex)	9	2529.57	6.19
35	**φ(age + popD) P(sex)**	**6**	**2535.62**	**6.20**
36	**φ(age + medLD) P(sex)**	**6**	**2536.86**	**7.44**
37	φ(age + repS) P(sex)	6	2537.07	7.65
38	φ(age + repS + medLD) P(sex)	7	2536.02	8.61
1	**φ(age)P(sex)**	**5**	**2540.29**	**8.87**
40	φ(age + repS + medLD + repS*medLD) P(sex)	8	2535.45	10.06

916 **Table A2**: Monostate model selection assessing the link between population-level phenology and average annual survival. **A**: Preliminary model
917 selection assessing the effects of year, sex and age on recapture (*p*) and survival (*φ*) probabilities. Starting model: Model 22, best model: Model 1.
918 AICc(Model 1) = 2550.3171. **B**: Secondary model selection assessing the effect of three annual population-level demographic covariates (repS=
919 reproductive success, popD = population density, medLD = median laying date) on survival probabilities. The covariates and their interactions were
920 added to preliminary selected model 1. Starting model: Model 1, best model: Model 23. AICc(Model 23) = 2541.4519. np: number of parameters.

TABLE A3

Model	Model Description	np	Deviance	ΔAICc
41	**IS(to.age) ɸ(age) T(f.to + age) P(sex)**	**23**	**8539.83**	**0**
42	**IS(to.age) ɸ(f(1 2 3,4) + age) T(f.to + age) P(sex)**	**24**	**8538.58**	**0.79**
43	**IS(to.age) ɸ(f(1,2 3 4) + age) T(f.to + age) P(sex)**	**24**	**8538.59**	**0.80**
44	**IS(to.age) ɸ(f(1 2,3 4) + age) T(f.to + age) P(sex)**	**24**	**8539.17**	**1.38**
45	**IS(to.age) ɸ(f(1,2 3 4).age) T(f.to + age) P(sex)**	**25**	**8537.51**	**1.75**
46	**IS(to.age) ɸ(sex + age) T(f.to + age) P(sex)**	**24**	**8539.77**	**1.98**
47	**IS(to.age) ɸ(f(1 4,2 3) + age) T(f.to + age) P(sex)**	**24**	**8539.83**	**2.04**
48	IS(to.age) ɸ(f(1,2 3,4) + age) T(f.to + age) P(sex)	25	8538.02	2.26
49	IS(to.age) ɸ(f(1,2 3 4) + sex + age) T(f.to + age) P(sex)	25	8538.54	2.79
50	IS(to.age) ɸ(f(1 2,3,4) + age) T(f.to + age) P(sex)	25	8538.55	2.79
51	IS(to.age) ɸ(f(1,2,3 4) + age) T(f.to + age) P(sex)	25	8538.57	2.82
52	IS(to.age) ɸ(.) T(f.to + age) P(sex)	22	8545.10	3.24
53	IS(to.age) ɸ(f(1,2 3 4).sex + age) T(f.to + age) P(sex)	26	8537.42	3.70
54	IS(to.age) ɸ(f(1,2 3 4).age + sex) T(f.to + age) P(sex)	26	8537.44	3.72
55	IS(to.age) ɸ(sex.age) T(f.to + age) P(sex)	25	8539.66	3.91
56	IS(to.age) ɸ(f(1 4,2,3) + age) T(f.to + age) P(sex)	25	8539.76	4.00
57	IS(to.age) ɸ(f + age) T(f.to + age) P(sex)	26	8537.94	4.22
58	IS(to.age) ɸ(f(1,2 3 4).[age + sex]) T(f.to + age) P(sex)	27	8536.19	4.51
59	IS(to.age) ɸ(f(1,2 3 4)+ sex.age) T(f.to + age) P(sex)	26	8538.46	4.74
60	IS(to.age) ɸ(f(1,2 3 4)) T(f.to + age) P(sex)	23	8544.80	4.97
61	IS(to.age) ɸ(sex) T(f.to + age) P(sex)	23	8545.07	5.24
62	IS(to.age) ɸ(f + sex + age) T(f.to + age) P(sex)	27	8537.89	6.22
63	IS(to.age) ɸ(f(1,2 3 4) + sex) T(f.to + age) P(sex)	24	8544.74	6.95
64	IS(to.age) ɸ(f.age) T(f.to + age) P(sex)	29	8534.63	7.04
65	IS(to.age) ɸ(f(1,2 3 4).sex.age) T(f.to + age) P(sex)	29	8535.45	7.86
66	IS(to.age) ɸ(f.sex + age) T(f.to + age) P(sex)	30	8533.72	8.17
67	IS(to.age) ɸ(f + sex.age) T(f.to + age) P(sex)	28	8537.82	8.19
68	IS(to.age) ɸ(f(1,2 3 4).sex) T(f.to + age) P(sex)	25	8543.96	8.20
69	IS(to.age) ɸ(f) T(f.to + age) P(sex)	25	8544.41	8.65
70	IS(to.age) ɸ(f.age + sex) T(f.to + age) P(sex)	30	8534.60	9.05
71	IS(to.age) ɸ(f.[age + sex]) T(f.to + age) P(sex)	33	8529.71	10.31
72	IS(to.age) ɸ(f + sex) T(f.to + age) P(sex)	26	8544.35	10.63
73	IS(to.age) ɸ(f.sex) T(f.to + age) P(sex)	29	8540.37	12.78
74	**IS(to.age) ɸ(f.age.sex) T(f.to + age) P(sex)**	**37**	**8528.26**	**17.08**
75	IS(to.age) ɸ(f.age.sex) T(f.to + sex + age) P(sex)	38	8526.56	17.44
76	IS(to.age) ɸ(f.age.sex) T(f.to) P(sex)	36	8531.58	18.34
77	IS(to.age) ɸ(f.age.sex) T(f.to + sex.age) P(sex)	39	8526.27	19.20
78	IS(to.age) ɸ(f.age.sex) T(f.to + sex) P(sex)	37	8530.39	19.20
79	IS(to.age) ɸ(f.age.sex) T(f.to.sex + age) P(sex)	49	8507.34	20.95
80	IS(to.age) ɸ(f.age.sex) T(f.to.sex) P(sex)	48	8510.60	22.13
81	IS(to.age) ɸ(f.age.sex) T(f.to.age) P(sex)	48	8519.76	31.30
82	IS(to.age) ɸ(f.age.sex) T(f.to.age + sex) P(sex)	49	8518.15	31.76

			np		
97	IS(to.age) φ(f.age.sex) T(age) P(sex)	26	8649.29	115.57
98	IS(to.age) φ(f.age.sex) T(s + age) P(sex)	27	8647.84	116.16
99	IS(to.age) φ(f.age.sex) T(s.age) P(sex)	28	8646.91	117.28
101	IS(to.age) φ(f.age.sex) T(.) P(sex)	25	8666.16	130.41
102	IS(to.age) φ(f.age.sex) T(sex) P(sex)	26	8665.69	131.98
83	IS(to.age + sex) φ(f.age.sex) T(f.to.age.sex) P(sex)	73	8488.61	52.49
84	**IS(to.age) φ(f.age.sex) T(f.to.age.sex) P(sex)**	**72**	**8491.61**	**53.38**
85	**IS(to.sex.age) φ(f.age.sex) T(f.to.age.sex) P(sex)**	**78**	**8483.63**	**58.10**
86	IS(to.sex + age) φ(f.age.sex) T(f.to.age.sex) P(sex)	73	8514.14	78.03
87	IS(to + age) φ(f.age.sex) T(f.to.age.sex) P(sex)	70	8525.51	83.06
88	IS(age.sex + to) φ(f.age.sex) T(f.to.age.sex) P(sex)	72	8522.48	84.25
89	IS(sex + age) φ(f.age.sex) T(f.to.age.sex) P(sex)	69	8529.57	85.02
90	IS(age) φ(f.age.sex) T(f.to.age.sex) P(sex)	68	8532.57	85.90
91	IS(sex.age) φ(f.age.sex) T(f.to.age.sex) P(sex)	70	8529.55	87.09
92	IS(to + sex) φ(f.age.sex) T(f.to.age.sex) P(sex)	70	8530.16	87.71
93	IS(to.sex) φ(f.age.sex) T(f.to.age.sex) P(sex)	72	8527.87	89.64
94	IS(to) φ(f.age.sex) T(f.to.age.sex) P(sex)	69	8542.91	98.35
95	IS(sex) φ(f.age.sex) T(f.to.age.sex) P(sex)	68	8545.83	99.16
96	IS(.) φ(f.age.sex) T(f.to.age.sex) P(sex)	67	8558.58	109.81
100	IS(to + s + age) φ(f.age.sex) T(f.to.age.sex) P(sex)	70	8561.74	119.29

Table A3: Multistate model selection assessing the link between relative laying date and subsequent adult
survival. The four states represent laying date quartiles (an individual in state 1 in year t has laid in the earliest
25% of the population that year). All models have sex as only constraint on recapture probabilities (P).
Selection was made sequentially (by retaining the best structure at each step for the next step), firstly on initial
state distribution constraints (IS), then on transition probabilities constraints (T), then on survival constraints
(φ). Starting model: Model 85, best model: Model 41. AICc(Model 41) = 8586.2435. For initial state distribution:
"to" means that at their first capture, individuals are not evenly distributed among the four phenological
quartiles. For between states transition probabilities: "f.to" means that transition probabilities depend on the
departure state (f), the arrival state (to) and their interaction. For survival: "f" means that survival depends on
the current state of the individual. np: number of parameters.

TABLE A4

Model	Model Description		T	P	np	Deviance	ΔQAIC
	IS	φ					
103	to.age	(medLD + popD + medLD*popD) + f(1,2 3 4) + age)	f.to + age	sex	27	8523.97	0
104 **(Figure 4)**	"	**(medLD + popD + medLD*popD).f(1,2 3 4) + age**	"	"	**29**	**8520.13**	**0.25**
105	"	(Temp + medLD + popD + medLD*popD).f(1,2 3 4) + age	"	"	31	8516.64	0.85
106	"	(medLD + popD + repS + medLD*popD).f(1,2 3 4) + age	"	"	31	8518.07	2.28
107	"	(popD + repS + popD*repS).f(1,2 4) + age	"	"	29	8522.57	2.69
108	"	(Temp + medLD + popD + repS + medLD*popD).f(1,2 3 4) + age	"	"	33	8515.57	3.89
109	"	(Temp + medLD + popD + Temp*medLD + medLD*popD).f(1,2 3 4) + age	"	"	33	8515.60	3.92
110	"	(Temp + medLD + popD + Temp*popD + medLD*popD).f(1,2 3 4) + age	"	"	33	8515.62	3.93
111	"	(Temp + popD + repS + Temp*repS + popD*repS).f(1,2 3 4) + age	"	"	33	8515.81	4.12
112	"	(Temp + medLD + popD + repS + Temp*repS + popD*repS).f(1,2 3 4) + age	"	"	35	8512.35	4.77
113	"	(Temp + medLD + popD + repS + popD*repS).f(1,2 3 4) + age	"	"	33	8516.40	4.71
114	"	(Temp + medLD + popD + Temp*medLD).f(1,2 3 4) + age	"	"	31	8520.73	4.95
115	"	(medLD + popD + repS + medLD*popD + popD*repS).f(1,2 3 4) + age	"	"	33	8516.76	5.07
116	"	(medLD + popD + repS + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	33	8516.91	5.22
117	"	(Temp + popD + repS + popD*repS).f(1,2 3 4) + age	"	"	31	8521.14	5.35
118	"	(Temp + medLD + popD + repS + medLD*popD + popD*repS).f(1,2 3 4) + age	"	"	35	8513.22	5.64
119	"	(Temp + medLD + popD + repS + Temp*repS + medLD*popD).f(1,2 3 4) + age	"	"	35	8513.57	5.99
120	"	(Temp + medLD + popD).f(1,2 3 4) + age	"	"	29	8525.60	5.72
121	"	(Temp + medLD + popD + repS + Temp*medLD + popD*repS).f(1,2 3 4) + age	"	"	35	8513.61	6.02
122	"	(Temp + medLD + popD + Temp*popD).f(1,2 3 4) + age	"	"	31	8521.67	5.89
123	"	(medLD + popD + repS + popD*repS).f(1,2 3 4) + age	"	"	31	8521.80	6.01
124	"	(Temp + medLD + popD + repS + Temp*popD + medLD*popD).f(1,2 3 4) + age	"	"	35	8514.09	6.51
125	"	(popD) + f(1,2 3 4) + age)	"	"	25	8534.10	6.05
126	"	(popD + repS).f(1,2 3 4) + age	"	"	27	8530.21	6.24
127	"	(Temp + popD + Temp*popD).f(1,2 3 4) + age	"	"	29	8526.29	6.41
128	"	(medLD + popD).f(1,2 3 4) + age	"	"	27	8530.30	6.33
129	"	(Temp + medLD + popD + repS + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	35	8514.34	6.75
130	"	(Temp + medLD + popD + repS + Temp*repS + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	37	8510.38	6.91
131	"	(Temp + medLD + popD + Temp*medLD + Temp*popD + medLD*popD).f(1,2 3 4) + age	"	"	35	8514.48	6.90

132	"	(Temp + medLD + popD + repS + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	37	8510.53	7.06
133	"	(Temp + medLD + popD + repS + Temp*repS + medLD*popD + popD*repS).f(1,2 3 4) + age	"	"	37	8510.56	7.09
134	"	(Temp + medLD + popD + repS + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	35	8514.64	7.06
135	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	39	8506.77	7.42
136	"	(Temp + medLD + popD + repS + Temp*medLD + medLD*popD).f(1,2 3 4) + age	"	"	35	8514.79	7.21
137	"	(medLD + popD + repS + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	35	8514.85	7.27
138	"	(Temp + medLD + Temp*medLD).f(1,2 3 4) + age	"	"	29	8526.90	7.02
139	"	**(popD).f(1,2 3 4) + age**	"	"	**25**	**8535.01**	**6.96**
140	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS + popD*repS).f(1,2 3 4) + age	"	"	37	8511.06	7.59
141	"	(Temp + medLD + popD + repS + Temp*repS + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	37	8511.22	7.75
142	"	(Temp + medLD + popD + Temp*medLD + Temp*popD).f(1,2 3 4) + age	"	"	33	8519.23	7.54
143	"	(Temp + popD + repS + Temp*repS).f(1,2 3 4) + age	"	"	31	8523.27	7.49
144	"	(Temp + repS + Temp*repS).f(1,2 3 4) + age	"	"	29	8527.34	7.46
145	"	(medLD) + f(1,2 3 4) + age)	"	"	25	8535.54	7.49
146	"	(Temp + popD + repS + Temp*popD).f(1,2 3 4) + age	"	"	31	8523.59	7.80
147	"	(Temp + popD + repS + Temp*popD + Temp*repS + popD*repS).f(1,2 3 4) + age	"	"	35	8515.61	8.02
148	"	(Temp + medLD + popD + repS + Temp*popD + medLD*popD + popD*repS).f(1,2 3 4) + age	"	"	37	8511.69	8.21
149	"	(Temp + medLD + popD + repS + Temp*medLD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	37	8511.74	8.27
150	"	(Temp + medLD + popD + repS + Temp*popD + Temp*repS + medLD*popD).f(1,2 3 4) + age	"	"	37	8511.93	8.46
151	"	(Temp + medLD + popD + repS + Temp*medLD + medLD*popD + popD*repS).f(1,2 3 4) + age	"	"	37	8511.99	8.52
152	"	(Temp + medLD).f(1,2 3 4) + age	"	"	27	8532.02	8.01
153	"	(Temp + popD + repS + Temp*popD + popD*repS).f(1,2 3 4) + age	"	"	33	8520.03	8.34
154 **(Figure 3)**	"	**(medLD).f(1,2 3 4) + age**	"	"	**25**	**8536.04**	**7.99**
155	"	(repS).f(1,2 3 4) + age	"	"	25	8536.06	8.01
156	"	(Temp + popD + repS + Temp*popD + Temp*repS).f(1,2 3 4) + age	"	"	33	8520.22	8.53
157	"	(Temp + medLD + popD + repS + Temp*medLD).f(1,2 3 4) + age	"	"	33	8520.25	8.56
158	"	(Temp + medLD + popD + repS + Temp*popD + Temp*repS + popD*repS).f(1,2 3 4) + age	"	"	37	8512.25	8.78
159	"	(Temp + medLD + popD + repS + Temp*popD + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	37	8512.42	8.95
160	"	(Temp + medLD + popD + repS + Temp*popD + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	39	8508.43	9.07
161	"	(Temp + medLD + popD + repS + Temp*repS).f(1,2 3 4) + age	"	"	33	8520.44	8.75
43	"	**f(1,2 3 4) + age**	"	"	**24**	**8538.59**	**8.51**

162	"	(Temp + medLD + popD + repS + Temp*popD + Temp*repS + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	41	8504.68	9.44
163	"	(medLD + popD + repS + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	33	8520.83	9.15
164	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	39	8508.88	9.52
165	"	(Temp + medLD + popD + repS + Temp*popD + Temp*repS + medLD*popD + popD*repS).f(1,2 3 4) + age	"	"	39	8508.91	9.55
166	"	(medLD + popD + repS).f(1,2 3 4) + age	"	"	29	8528.91	9.03
167	"	(Temp + medLD + popD + repS + Temp*popD).f(1,2 3 4) + age	"	"	33	8520.93	9.24
168	"	(Temp + popD + repS).f(1,2 3 4) + age	"	"	29	8528.94	9.06
169	"	(Temp + medLD + popD + repS).f(1,2 3 4) + age	"	"	31	8524.97	9.19
170	"	(Temp + popD).f(1,2 3 4) + age	"	"	27	8533.06	9.09
171	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + popD*repS).f(1,2 3 4) + age	"	"	37	8513.18	9.71
172	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + medLD*popD).f(1,2 3 4) + age	"	"	37	8513.23	9.76
173	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS + medLD*popD).f(1,2 3 4) + age	"	"	37	8513.24	9.77
174	"	(Temp + medLD + repS + Temp*repS).f(1,2 3 4) + age	"	"	31	8525.25	9.46
175	"	(Temp + medLD + popD + repS + Temp*popD + Temp*repS + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	39	8509.26	9.90
176	"	(Temp + medLD + popD + repS + Temp*medLD + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	39	8509.27	9.91
177	"	(Temp + medLD + popD + repS + Temp*medLD + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	37	8513.34	9.86
178	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS + medLD*popD + popD*repS).f(1,2 3 4) + age	"	"	39	8509.89	10.53
179	"	(Temp + medLD + popD + repS + Temp*popD + Temp*repS).f(1,2 3 4) + age	"	"	35	8517.92	10.38
180	"	(Temp + medLD + popD + repS + Temp*popD + Temp*repS + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	39	8509.94	10.58
181	"	(Temp).f(1,2 3 4) + age	"	"	25	8537.96	9.92
182	"	(Temp + medLD + popD + repS + Temp*popD + popD*repS).f(1,2 3 4) + age	"	"	36	8516.04	10.52
183	"	(Temp + medLD + popD + repS + Temp*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	37	8514.13	10.66
184	"	[(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + medLD*popD + popD*repS)].f(1,2 3 4) + age	"	"	39	8510.17	10.82
185	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS).f(1,2 3 4) + age	"	"	35	8518.22	10.64
186	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	41	8506.27	11.04
187	"	(Temp + medLD + repS + Temp*medLD).f(1,2 3 4) + age	"	"	31	8526.43	10.65
188	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD).f(1,2 3 4) + age	"	"	35	8518.70	11.11
189	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + Temp*repS + popD*repS).f(1,2 3 4) +	"	"	39	8510.74	11.39

		age					
190	"	(Temp + repS).f(1,2 3 4) + age	"	"	27	8534.79	10.82
191	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	39	8510.88	11.52
192	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	41	8506.98	11.75
193	"	(medLD + repS).f(1,2 3 4) + age	"	"	27	8535.00	11.03
194	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	39	8511.10	11.74
195	"	(Temp + medLD + popD + repS + Temp*repS + medLD*repS).f(1,2 3 4) + age	"	"	35	8519.15	11.57
196	"	(Temp + medLD + repS + Temp*medLD + Temp*repS).f(1,2 3 4) + age	"	"	33	8523.38	11.69
197	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	39	8511.42	12.06
198	"	(Temp + medLD + repS).f(1,2 3 4) + age	"	"	29	8531.44	11.56
199	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + Temp*repS + medLD*popD).f(1,2 3 4) + age	"	"	39	8511.57	12.21
200	"	(Temp + medLD + repS + Temp*repS + medLD*repS).f(1,2 3 4) + age	"	"	33	8523.84	12.15
201	"	(Temp + medLD + popD + repS + Temp*medLD + medLD*repS).f(1,2 3 4) + age	"	"	35	8519.98	12.40
202	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + Temp*repS + medLD*popD + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	43	8504.05	12.95
203	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + Temp*repS + medLD*popD + popD*repS).f(1,2 3 4) + age	"	"	41	8508.07	12.84
204	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + Temp*repS + medLD*repS + popD*repS).f(1,2 3 4) + age	"	"	41	8508.30	13.07
205	"	(Temp + medLD + popD + repS + Temp*popD + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	37	8516.31	12.83
206	"	(Temp + medLD + popD + repS + Temp*popD + Temp*repS + medLD*repS).f(1,2 3 4) + age	"	"	37	8516.31	12.83
207	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*repS + medLD*repS).f(1,2 3 4) + age	"	"	37	8516.61	13.14
208	"	(medLD + popD + repS + medLD*repS).f(1,2 3 4) + age	"	"	31	8528.66	12.87
209	"	(Temp + medLD + popD + repS + medLD*repS).f(1,2 3 4) + age	"	"	33	8524.72	13.03
210	"	(Temp + medLD + popD + repS + Temp*popD + medLD*repS).f(1,2 3 4) + age	"	"	35	8520.73	13.15
211	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + Temp*repS).f(1,2 3 4) + age	"	"	37	8516.80	13.33
212	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + Temp*repS + medLD*popD + medLD*repS).f(1,2 3 4) + age	"	"	41	8508.94	13.71
213	"	(Temp + medLD + repS + Temp*medLD + Temp*repS + medLD*repS).f(1,2 3 4) + age	"	"	35	8521.61	14.02
214	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + medLD*repS).f(1,2 3 4) + age	"	"	37	8518.23	14.75
215	"	(Temp + medLD + repS + Temp*medLD + medLD*repS).f(1,2 3 4) + age	"	"	33	8526.26	14.57

216	"	(medLD + repS + medLD*repS).f(1,2 3 4) + age	"	"	29	8534.61	14.74
217	"	(Temp + medLD + popD + repS + Temp*medLD + Temp*popD + Temp*repS + medLD*repS).f(1,2 3 4) + age	"	"	39	8515.05	15.70
218	"	(Temp + medLD + repS + medLD*repS).f(1,2 3 4) + age	"	"	31	8531.11	15.32
219	**"**	**year.f(1,2 3 4) + age**	**"**	**"**	**97**	**8406.10**	**28.90**

Table A4: Multistate model selection assessing the effects of population-level annual covariates on the relative laying date - adult survival link. The effects of 4 annual covariates (Temp = mean daily mean temperature from March 31st to May 7th, repS = reproductive success, popD = population density, medLD = median laying date) and their first-order interactions as continuous temporal covariates on survival probabilities are tested on one of the best models from the previous selection (Table 3): Model 43. Starting model: Model 43, best model: Model 103. AICc(Model 103) = 8578.5326. For initial state distribution: "to" means that at their first capture, individuals are not evenly distributed among the four phenological quartiles. For between states transition probabilities: "f.to" means that transition probabilities depend on the departure state (f), the arrival state (to) and their interaction. For survival: "f" means that survival depends on the current state of the individual. np: number of parameters.

TABLE A5

Model	Model Description	np	Deviance	ΔAICc
220	*IS(to.age) φ(f(12,3456) + age) T(f.to) P(sex)*	*45*	*9508.25*	*0*
221	*IS(to.age) φ(f(12,35,46) + age) T(f.to) P(sex)*	*46*	*9507.37*	*1.18*
222	*IS(to.age) φ(f(12,3456).age) T(f.to) P(sex)*	*46*	*9508.00*	*1.82*
223	*IS(to.age) φ(f(12,34,56) + age) T(f.to) P(sex)*	*46*	*9508.03*	*1.85*
224	IS(to.age) φ(f(12,3456) + sex + age) T(f.to) P(sex)	46	9508.20	2.01
225	IS(to.age) φ(f(12,3456).age + sex) T(f.to) P(sex)	47	9507.96	3.84
226	IS(to.age) φ(f(12,34,56) + sex + age) T(f.to) P(sex)	47	9507.98	3.87
227	IS(to.age) φ(f(12,3456).sex + age) T(f.to) P(sex)	47	9507.98	3.87
228	IS(to.age) φ(f(12,3456) + sex.age) T(f.to) P(sex)	47	9508.16	4.04
229	IS(to.age) φ(f(12,3456)) T(f.to) P(sex)	44	9514.23	3.91
230	IS(to.age) φ(f(12,34,56).age) T(f.to) P(sex)	48	9507.79	5.75
231	IS(to.age) φ(f(12,34,56)) T(f.to) P(sex)	45	9513.81	5.56
232	IS(to.age) φ(f(12,34,56) + sex.age) T(f.to) P(sex)	48	9507.93	5.89
233	*IS(to.age) φ(age) T(f.to) P(sex)*	*44*	*9516.09*	*5.77*
234	IS(to.age) φ(f(12,3456) + sex) T(f.to) P(sex)	45	9514.18	5.93
235	*IS(to.age) φ(from + age) T(f.to) P(sex)*	*49*	*9506.67*	*6.71*
236	*IS(to.age) φ(f(135,246) + age) T(f.to) P(sex)*	*45*	*9515.01*	*6.76*
237	IS(to.age) φ(f(12,34,56).sex + age) T(f.to) P(sex)	49	9507.45	7.49
238	IS(to.age) φ(f(12,34,56).age + sex) T(f.to) P(sex)	49	9507.75	7.78
239	IS(to.age) φ(f(12,34,56) + sex) T(f.to) P(sex)	46	9513.76	7.58
240	IS(to.age) φ(sex + age) T(f.to) P(sex)	45	9516.00	7.75
241	IS(to.age) φ(f(12,3456).sex) T(f.to) P(sex)	46	9514.04	7.86
242	IS(to.age) φ(from + sex + age) T(f.to) P(sex)	50	9506.63	8.74
243	IS(to.age) φ(f(12,3456).sex.age) T(f.to) P(sex)	50	9506.74	8.86
244	IS(to.age) φ(f(135,246).age) T(f.to) P(sex)	46	9514.84	8.66
245	IS(to.age) φ(f(135,246) + sex + age) T(f.to) P(sex)	46	9514.93	8.75
246	IS(to.age) φ(i) T(f.to) P(sex)	43	9521.53	9.15
247	IS(to.age) φ(sex.age) T(f.to) P(sex)	46	9515.91	9.73
248	IS(to.age) φ(f(135,246).sex + age) T(f.to) P(sex)	47	9514.42	10.31
249	IS(to.age) φ(from + sex.age) T(f.to) P(sex)	51	9506.59	10.78
250	IS(to.age) φ(f(135,246).age + sex) T(f.to) P(sex)	47	9514.75	10.64
251	IS(to.age) φ(f(135,246) + sex.age) T(f.to) P(sex)	47	9514.86	10.75
252	IS(to.age) φ(from) T(f.to) P(sex)	48	9513.23	11.19
253	IS(to.age) φ(f(135,246)) T(f.to) P(sex)	44	9521.31	10.99
254	IS(to.age) φ(f(12,34,56).sex) T(f.to) P(sex)	48	9513.38	11.34
255	IS(to.age) φ(sex) T(f.to) P(sex)	44	9521.51	11.19
256	IS(to.age) φ(from + sex) T(f.to) P(sex)	49	9513.17	13.20
257	IS(to.age) φ(f(135,246) + sex) T(f.to) P(sex)	45	9521.28	13.03
258	IS(to.age) φ(from.age) T(f.to) P(sex)	54	9503.62	14.05
259	IS(to.age) φ(f(135,246).sex) T(f.to) P(sex)	46	9520.97	14.79
260	IS(to.age) φ(from.age + sex) T(f.to) P(sex)	55	9503.59	16.11
261	IS(to.age) φ(f(123,456).sex.age) T(f.to) P(sex)	50	9513.77	15.89
262	IS(to.age) φ(f(12,34,56).sex.age) T(f.to) P(sex)	54	9505.88	16.31
263	IS(to.age) φ(from.sex + age) T(f.to) P(sex)	55	9505.15	17.67
264	IS(to.age) φ(from.sex) T(f.to) P(sex)	54	9512.05	22.49
265	**IS(to.age) φ(f.age.sex) T(f.to) P(sex)**	**66**	**9498.25**	**33.79**

266	IS(to.age) φ(f.age.sex) T(f.to + sex) P(sex)	67	9497.11	34.76
267	IS(to.age) φ(f.age.sex) T(f.to + age) P(sex)	67	9498.15	35.79
268	IS(to.age) φ(f.age.sex) T(f.to + sex + age) P(sex)	68	9496.89	36.64
269	IS(to.age) φ(f.age.sex) T(f.to + sex.age) P(sex)	69	9495.91	37.76
270	IS(to.age) φ(f.age.sex) T(f.to.age) P(sex)	96	9465.56	64.89
271	IS(to.age) φ(f.age.sex) T(f.to.age + sex) P(sex)	97	9464.00	65.49
272	IS(to.age) φ(f.age.sex) T(f.to.sex) P(sex)	96	9475.36	74.69
273	IS(to.age) φ(f.age.sex) T(f.to.sex + age) P(sex)	97	9475.13	76.62
284	IS(to.age) φ(f.age.sex) T(.) P(sex)	37	9697.50	172.74
285	IS(to.age) φ(f.age.sex) T(sex) P(sex)	38	9696.77	174.07
286	IS(to.age) φ(f.age.sex) T(age) P(sex)	38	9696.90	174.20
287	IS(to.age) φ(f.age.sex) T(sex + age) P(sex)	39	9695.97	175.33
288	IS(to.age) φ(f.age.sex) T(sex.age) P(sex)	40	9695.37	176.79
274	**IS(to.age) φ(f.age.sex) T(f.to.age.sex) P(sex)**	**153**	**9414.88**	**139.55**
275	IS(to.age + sex) φ(f.age.sex) T(f.to.age.sex) P(sex)	154	9414.50	141.42
276	**IS(to.sex.age) φ(f.age.sex) T(f.to.age.sex) P(sex)**	**163**	**9408.55**	**155.79**
277	IS(to.sex + age) φ(f.age.sex) T(f.to.age.sex) P(sex)	154	9445.49	172.41
278	IS(to + age) φ(f.age.sex) T(f.to.age.sex) P(sex)	149	9455.95	171.64
279	IS(to) φ(f.age.sex) T(f.to.age.sex) P(sex)	147	9461.22	172.43
280	IS(to + sex + age) φ(f.age.sex) T(f.to.age.sex) P(sex)	150	9455.57	173.51
281	IS(to.sex) φ(f.age.sex) T(f.to.age.sex) P(sex)	153	9451.12	175.79
282	IS(age.sex + to) φ(f.age.sex) T(f.to.age.sex) P(sex)	151	9455.56	175.74
283	IS(to + sex) φ(f.age.sex) T(f.to.age.sex) P(sex)	149	9461.21	176.90
289	IS(age) φ(f.age.sex) T(f.to.age.sex) P(sex)	145	10238.62	945.37
290	IS(sex + age) φ(f.age.sex) T(f.to.age.sex) P(sex)	146	10238.25	947.22
291	IS(.) φ(f.age.sex) T(f.to.age.sex) P(sex)	144	10243.89	948.40
292	IS(sex.age) φ(f.age.sex) T(f.to.age.sex) P(sex)	147	10238.23	949.44
293	IS(sex) φ(f.age.sex) T(f.to.age.sex) P(sex)	145	10243.88	950.62

Table A5: Multistate model selection assessing the links between individual breeding success, relative laying date and adult survival. The 6 states represent different combinations of individual breeding success and relative laying date (see Figure 5 for details). Starting model: Model 276 included the interactive effects of age, sex and state for its various structural components. Recapture probabilities were assumed to depend on sex only. Sequential selection procedure was the same as in the previous model selection (Table 4). Best model: Model 220. AICc(Model 220) = 9599.8097. np: number of parameters.

Appendix 3: Temporal variation in survival in the two age classes

A series of additional models were made to check that the relationship between the survival
of individuals and the variables presented in the results (median laying date medLD and
population density popD) was not due to interannual variations in the proportion of age
classes. (1 year old individuals 1yo versus 2 or more years 2yo). Additional survival models
consider that the survival of a single age class varies over time while the other is constant
(e.g., 2yo + 1yo.year) or that the survival of a single class is constrained by the median laying
date or the population density (e.g., 2yo + 1yo.medLD). The results are presented in Table A6
below and illustrated in Figure A1.

Model description	np	Deviance	ΔAICc
φ (age + popD) p(sex)	6	2535.62	0
φ (age +medLD) p(sex)	6	2536.86	1.24
φ (1yo + 2yo.popD) p(sex)	6	2537.76	2.14
φ (2yo +1yo.popD) p(sex)	6	2537.99	2.38
φ (1yo + 2yo.medLD) p(sex)	6	2538.14	2.52
φ (age) P(sex)	5	2540.29	2.67
φ (2yo + 1yo.medLD) p(sex)	6	2538.91	3.30
φ (1yo + 2yo.year) p(sex)	41	2476.02	11.78
φ (2yo + 1yo.year) p(sex)	38	2504.29	33.85
φ (age.year) p(sex)	74	2440.68	45.63

971 **Table A6**: Monostate model selection assessing the link between the median laying date medLD, the
972 population density and average annual survival. Best model: Model φ (age + popD) p(sex), AICc = 2547.6513. p:
973 number of parameters.

Figure A1: Relationship between annual survival probabilities and two demographic parameters. **A**: the survival probability of 1 year old individuals (1yo) is positively correlated with median laying date of the population in the previous spring (medLD). The points are the estimates of Model φ (2yo + 1yo.year) p(sex) in Table A6 for 1yo, with 95% confidence intervals, the solid line is for 1yo from Model φ (2yo + 1yo.medLD) p(sex) in Table A6, which has a continuous effect of medLD on survival, its 95% CI are represented by the dashed lines. **B**: the survival probability of 2 years old or older individuals (2yo) is positively correlated with medLD. The points are the estimates of Model φ (1yo + 2yo.year) p(sex) in Table A6 for 2yo, with 95% confidence intervals, the solid line is for 2yo from Model φ (1yo + 2yo.medLD) p(sex) in Table A6, which has a continuous effect of medLD on survival, its 95% CI are represented by the dashed lines. **C**: the survival probability of 1yo is negatively correlated with population density during the previous spring (popD). The points are the estimates of Model φ (2yo + 1yo.year) p(sex) in Table A6 for 1yo, with 95% confidence intervals, the solid line is for 1yo from Model φ (2yo + 1yo.popD) p(sex) in Table A6, which has a continuous effect of popD on survival, its 95% CI are represented by the dashed lines. **D**: the survival probability of 2yo is positively correlated with popD. The points are the estimates of Model φ (1yo + 2yo.year) p(sex) in Table A6 for 2yo, with 95% confidence intervals, the solid line is for 2yo from Model φ (1yo + 2yo.popD) p(sex) in Table A6, which has a continuous effect of popD on survival, its 95% CI are represented by the dashed lines.

Appendix 4: Repeatability of laying date percentile in the E-Pirio population

We performed a repeatability analysis based on 1000 parametric bootstraps as implemented in the rptR package (Stoffel et al., 2018) of the R-software v3.5.1 (R Core Team, 2018). The statistical significance of the repeatability of each metric was tested by a likelihood ratio test comparing the model fit of a model including grouping factor (here, the species or the street) and one excluding it.

A significant repeatability was found for both males and females:

For males: 712 observations, 382 individuals, R = 0.124 [0.027; 0.220]$_{95\%CI}$, likelihood ratio test: D=8.51, df=1, P=0.02

For females: 818 observations, 410 individuals, R = 0.406 [0.316; 0.493]$_{95\%CI}$, likelihood ratio test: D=95.1, df=1, P=0

References

R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Stoffel, M.A., Nakagawa, S., Schielzeth, H., 2018. rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. Methods Ecol. Evol. 1639–1644. https://doi.org/10.1111/2041-210X.12797@10.1111/(ISSN)2041-210X.CODEVI2018

1017 # Appendix 5: Matrix of probabilities of transition among
1018 the different relative laying date states as described in
1019 Table 3 and Figure 2.

1020

	to quartile 1	to quartile 2	to quartile 3	to quartile 4
from laying date quartile 1	**0.5186**	0.2495	0.1103	0.1215
from laying date quartile 2	0.4582	**0.2705**	0.1861	0.0852
from laying date quartile 3	0.3151	0.2722	**0.2103**	0.2025
from laying date quartile 4	0.1843	0.2610	0.3560	**0.1986**

mean transition probability: 0.25

1021

1022 Quartile 1 corresponds to the 25% earliest breeders of the population in a given year, quartile 4 to
1023 the latest. From all starting laying date quartiles, the probability of transitioning to the same or an
1024 earlier quartile are higher than transitioning to a later quartile.

1025