Paradoxes and synergies: optimizing management of a deadly virus in an endangered

carnivore

Marie L.J. Gilbertson[1]*, Dave Onorato[2], Mark W. Cunningham[3], Sue VandeWoude[4], Meggan E.

Craft[1,5]

*Corresponding author: jone1354@umn.edu

[1]Department of Veterinary Population Medicine, University of Minnesota, St Paul, MN 55108.

[2]Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission,

Naples, FL 34114.

[3]Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission,

Gainesville, FL 32601.

[4]Department of Microbiology, Immunology, and Pathology, Colorado State University, Fort

Collins, CO 80523

[5] Department of Ecology, Evolution and Behavior, University of Minnesota, St Paul, MN 55108.

Abstract

Management of pathogen transmission is often hindered by uncertainties in the efficacy

of and interactions between intervention strategies, sometimes resulting in unintended negative

consequences. Yet outbreaks of infectious disease can have serious consequences for wildlife

population health, especially species of conservation concern. The endangered Florida panther,

for example, experienced an outbreak of feline leukemia virus (FeLV) in 2002-2004, and

continues to be affected by this deadly virus. Ongoing management efforts aim to mitigate the

effects of FeLV on panthers with limited empirical information about which strategies are most

effective and efficient. We used a simulation-based approach to determine optimal FeLV

management strategies. We simulated use of proactive management interventions (i.e.,

proactive vaccination) as well as three reactive strategies, including vaccination in the face of an

outbreak, test and removal protocols, and temporary spatial segregation of the panther

population. Vaccination strategies included partial vaccine immunity, an understudied type of

vaccine imperfection. We compared the effectiveness of different strategies in mitigating FeLV

mortalities and duration of outbreaks. Results showed that inadequate proactive vaccination can

paradoxically increase the number of disease-induced mortalities in FeLV outbreaks, most likely

due to effects of partial vaccine immunity. Combinations of proactive vaccination with reactive

test-and-removal or vaccination had a synergistic effect in reducing impacts of FeLV outbreaks.

Temporary spatial restrictions were unlikely to be effective under realistic conditions. Our results

highlight the importance of integrating management and modeling approaches to identify

unexpected consequences and synergies in pathogen management interventions and aid in

conservation of at-risk species.

Significance Statement

Managing outbreaks of infectious disease is fraught with uncertainties such that seemingly

helpful interventions can have unintended negative consequences. We used a simulation

approach to determine optimal pathogen management strategies in an endangered carnivore,

the Florida panther (*Puma concolor coryi*), which continues to be affected by outbreaks of feline

leukemia virus. We tested interventions including proactive vaccination, reactive vaccination,

test-and-removal, and temporary spatial restrictions. We found that inadequate proactive

vaccination can counterintuitively increase disease-related mortalities. In contrast, we identified

synergies between proactive and reactive strategies that will be key for ongoing conservation of

the iconic Florida panther. The results of this study demonstrate the importance of linking

modeling and management to optimize pathogen control and avoid unexpected negative

consequences.

Keywords

disease model; simulation modeling; network modeling; feline leukemia virus; vaccination; test-

and-removal; spatial restrictions; *Puma concolor coryi*

Introduction

Outbreaks of infectious disease can have significant impacts on the population health of free-ranging wildlife, and are of heightened importance in species of conservation concern (1). For example, the feline retrovirus, feline leukemia virus (FeLV), has been the source of significant outbreaks in two endangered felids: Iberian lynx (*Lynx pardinus*) and Florida panthers (*Puma concolor coryi*). In the case of panthers, FeLV caused a deadly outbreak in 2002-2004 (2), spilling over from domestic cats, with subsequent direct transmission among panthers (3). In addition, there is recent evidence of ongoing FeLV spillover to and transmission among panthers (4), necessitating continued management of this deadly pathogen. FeLV inoculation with a domestic cat vaccine has been used previously, but with unknown efficacy in panthers (2). Further, the proportion of the population that must be vaccinated to most efficiently prevent future FeLV outbreaks is unknown, as is how proactive vaccination might interact with other reactive interventions to interrupt an FeLV outbreak. Such uncertainties are common among free-ranging wildlife threatened by infectious disease and hamper efforts to effectively control pathogen transmission.

Mathematical models of infectious disease transmission offer a powerful tool for filling such disease management knowledge gaps (5). Models allow the ethical testing of a wide range of different management approaches and can reveal unexpected consequences of disease control interventions (6). Models have been used to optimize disease management protocols in a variety of free-ranging wildlife species of conservation concern, including Ethiopian wolves (7), chimpanzees (8), and Amur tigers (9). Further, models can serve the important function of balancing the realities of fieldwork with ideal disease control protocols to provide practical, effective guidance for wildlife managers (e.g., 10, 11). Here, we use mathematical models of FeLV transmission in Florida panthers to determine optimal disease prevention and control strategies in this iconic carnivore.

Pathogen management strategies can be preventive (hereafter, *proactive*) or reactive.

Among proactive and reactive strategies, vaccination is a cornerstone in veterinary medicine (1,

12), and functions by reducing the availability of susceptible individuals. Vaccination can be

applied randomly, or targeted to high risk individuals (e.g., 13); to likely superspreaders (e.g., in

chimpanzees: 8); seasonally (e.g., 11, 14); or spatially, as with vaccine barriers or firewalls to

prevent spread of a pathogen to new areas or subsets of populations (e.g., 15–17). In panthers,

vaccination has been used for active management of FeLV (2, 4), but with uncertainties

regarding optimal distribution, individual efficacy, and level of population protection needed.

Among reactive pathogen management strategies, test-and-removal (or test-and-cull) is

commonly used in domestic species to functionally reduce the infectious period of infected

individuals (18). However, this approach is rarely used in wildlife due to the generally low

availability of field diagnostic tests and/or difficulty in recapturing individuals after a positive

diagnosis (18). Rather, pathogen control in wildlife often relies on non-selective culling in an

attempt to reduce density and—in theory—transmission (19). However, non-selective culling

can have a host of negative consequences for disease management including removal of

immune individuals (18, 20) or culling-induced perturbation (e.g., 21). Thanks to ready

availability of a field diagnostic test, FeLV management in domestic cats and Iberian lynx has

made use of test-and-removal or isolation of infected individuals (22, 23), and is part of future

FeLV mitigation plans in panthers. However, uncertainties remain regarding how test-and-

removal might interact with proactive vaccination efforts to more effectively reduce pathogen

impacts.

Larger-scale isolation or quarantine measures may also be used for disease

management in free-ranging wildlife, in the form of physical or behavioral barriers between

affected and unaffected subsets of a population. For example, fencing has been used to prevent

transmission of foot-and-mouth disease between wildlife and cattle in South Africa (24), and

winter feeding grounds may help prevent transmission of brucellosis between elk and cattle in

the Greater Yellowstone Ecosystem through behavioral separation (25). For panthers, while

physical barriers to prevent spillover from domestic cats are impractical, it may be possible to

use temporary spatial barriers to reduce the spatial spread of FeLV among panthers after a

spillover event. Specifically, the major I-75 freeway is fenced throughout Florida panther habitat

to reduce vehicle strikes, with regular wildlife underpasses the main means for wildlife to

traverse this barrier. While never before attempted, it may be possible to physically block these

underpasses under emergency conditions to prevent the spread of FeLV from the northern to

southern subsets of the panther population, or vice versa.

Given the ongoing risks of FeLV to panther conservation and the uncertainties regarding

application of proactive and reactive FeLV management strategies, the objectives of this study

were to test singly, and in combination, the effectiveness of: (1) proactive vaccination, (2)

reactive vaccination, (3) reactive test-and-removal, and (4) reactive temporary spatial

restrictions for reducing the population level impacts of FeLV in Florida panthers. We used a

mathematical modeling approach to address these objectives in order to efficiently and ethically

test a wide range of intervention protocols for the protection of this endangered carnivore.

Methods

Simulation pipeline

To examine the effect of different disease management regimes on FeLV control, we

used spatially-explicit, network simulations adapted from our research determining drivers of

retrovirus transmission in panthers (26). This approach involves two steps: (1) simulation of a

contact network among panthers, and (2) simulation of FeLV transmission on this network. In

brief, we simulated panther populations of 150 individuals (27) and used our previously

described exponential random graph model for retrovirus transmission in panthers (26) to

simulate contact networks constrained by network density among simulated populations (see

supplementary methods and Table S1 for further details).

We used a transmission model developed in Gilbertson et al. (26). Briefly, this model

used a susceptible-infectious-recovered compartmental framework, where individuals could be

progressively, regressively, or abortively infected (2). Progressive infections always resulted in

death, while regressive infections eventually recovered with immunity, and abortive infections

were always considered immune. Importantly, based on our previous work, we allowed both

progressives and regressives to be infectious, though regressives were less likely to transmit

(Table S1). Duration of infectiousness is long for FeLV (estimated as a mean of 18 weeks, 2;

but see 26 and Table S1). As in Gilbertson et al. (26), we included only disease-induced

mortality, and, in order to preserve key network structure, allowed territories vacated by deaths

to be reoccupied by new susceptible individuals (hereafter, *respawning*). Outbreaks were

initiated by a single, randomly selected non-isolate individual in the population (see supplement

discussion), and proceeded in weekly time steps for up to five years.

Our primary objective was to examine the effect of different FeLV management regimes

on epidemic outcomes, so in our primary simulations, we held network generation and

transmission parameters constant at previously supported values (see supplementary methods

and Table S1 for further details). We evaluated the consistency of our results to the choice of

parameter values with a sensitivity analysis (see below). Hereafter, a *parameter set* represents

the unique set of network, transmission, and management parameters for any given set of

simulations. A *full simulation* includes simulation of a single contact network and FeLV

transmission on that network (with or without management interventions). For each parameter

set, we performed 100 full simulations.

For a *baseline scenario*, we recorded key epidemic outcomes in the absence of

interventions. For *management scenarios* we recorded key outcomes in the presence of

interventions (Figure 1). These key outcomes were (1) the number of mortalities, (2) the

duration of an epidemic, and (3) the proportion of epidemics that "failed" per 100 successful

epidemics. A failed epidemic (i.e. stochastic fadeout, 28) was one in which fewer than 5

individuals acquired progressive or regressive infections. The outcomes of mortalities and

epidemic durations were summarized as median values per parameter set, as results were

often skewed; all outcomes were compared between baseline and management scenarios. All

simulations were performed in R version 3.6.3 (29).

Figure 1: *Steps of the simulation process across baseline and management scenarios. The*

basic network and FeLV transmission simulation steps are shown in yellow, the baseline (no

intervention) scenario in green, and the four different overarching management scenarios in

blue. Within each management scenario, we investigated several variations for the given

approach (shown in light blue) in a factorial design.

Proactive vaccination

In the first management scenario, we examined the effect of different levels of population

proactive vaccination (proportion of the population vaccinated prior to an outbreak), and

different ratios of single versus boosted vaccination. We simulated from 10-80% (in 10%

increments) of the population having some degree of vaccine-induced immunity to FeLV at the

onset of an outbreak. These vaccinations were distributed randomly in the population. Among

the vaccinated individuals, 0, 50, or 100% received a second boosting inoculation. Actual

vaccine efficacies are unknown for panthers, but based on efficacy studies in domestic cats (30,

31), we conservatively assumed that boosted vaccination would prevent 80% of infections and

single vaccination would prevent 40% of infections. We modeled this efficacy as a binomial

probability to represent partial vaccine immunity (versus binary immunity; 32). We conducted

proactive vaccination scenarios in a full factorial design, for a total of 24 proactive vaccination

parameter sets, and 2,400 full simulations (100 full simulations per parameter set).

Reactive vaccination

During simulated periods of reactive vaccination administration, panthers were selected

at a rate of one panther per week for vaccination. We assumed that managers would not know

the disease status of an individual selected for vaccination and that vaccination would be

ineffective in infectious or recovered individuals. In the case of previously vaccinated individuals,

a re-vaccination changed the vaccine efficacy of singly vaccinated individuals (efficacy of 40%)

to the efficacy for boosted individuals (efficacy of 80%). We further assumed that managers

would know which individuals had received two vaccinations, so boosted individuals were not

selected for additional vaccination attempts.

We varied the timing of the onset of reactive vaccination after the initiation of an FeLV

outbreak to reflect the difficulty of epidemic detection in this elusive carnivore. We therefore

began reactive vaccination at an optimistic, but difficult-to-attain time point of 26 weeks, and a

more realistic time point of 52 weeks. In addition, we varied the distribution of reactive

vaccination. While proactive vaccines were always distributed randomly, reactive vaccination

was either randomly distributed or spatially distributed in an attempted vaccine barrier based on

proximity to the I-75 freeway (see supplement for further details).

Because vaccination is resource and time intensive, we evaluated the effect of reactive

vaccination for 6 months per year versus year-round. Among reactive vaccination scenarios, we

also included scenarios with proactive vaccination, specifically where 0-60% of the population

was proactively vaccinated (in 20% increments) at the initiation of an outbreak. Because it is

highly unlikely that 100% of the proactively vaccinated population would have received boosted

vaccination, we used a more conservative—yet still challenging to attain—ratio of 50% of the

proactively vaccinated population being boosted (i.e. efficacy of 80%). We held this ratio of

efficacy constant for proactive vaccination in all reactive scenarios. We again used a factorial

design across all variations, resulting in 32 parameter sets and a total of 3,200 full simulations

under reactive vaccination scenarios.

Reactive test-and-removal

Test-and-removal scenarios were built around a protocol in which panthers infectious at

capture are removed from the population through humane euthanasia or temporary removal

until recovery. For simplicity, we assumed that all progressively infected individuals would show

clinical signs and be humanely euthanized at capture, while infectious regressive individuals

were temporarily removed from the population until their recovery and re-released into any open

territory. A maximum of five individuals were allowed to be temporarily removed in this way at

one time.

We expected that managers would be able to capture and test one panther per week at

most, and that captures occurred during a 17 week (about 4 month) capture season, in

accordance with current panther capture protocols. We assumed that managers would not know

the disease status of a target individual until the capture occurred, so captures were not

targeted by infection state. We varied the onset of test-and-removal, such that the intervention

began 26 or 52 weeks after the initiation of an epidemic. Captures were also random or spatially

targeted. If spatially targeted, captures (and consequent removals) only occurred on the same

side of the I-75 freeway as the initial FeLV infection.

As in the reactive vaccination scenarios, we again also included varying degrees of

proactive vaccination (0-60% in 20% increments). Reactive test-and-removal scenarios were

simulated in a full factorial design across variations, for a total of 16 parameter sets and 1,600

full simulations.

Reactive underpass closures

To determine the potential utility of closing I-75 wildlife underpasses in a FeLV outbreak,

we considered a "best case" scenario in which underpass closures were completely effective at

preventing transmission across the freeway. Underpasses were closed instantaneously either

26 or 52 weeks after the initiation of an FeLV outbreak, and remained closed for 4, 13, 26, or 52

weeks. We again included variations in proactive vaccination (0-60% in 20% increments), as in

other reactive management scenarios. The factorial design here resulted in 32 parameter sets

for underpass closure scenarios, for a total of 3,200 full simulations.

Sensitivity analysis

We used a latin hypercube sampling (LHS) approach to generate 50 sensitivity analysis

parameter sets across our 8 network and transmission parameters using the *lhs* package in R

(33). We repeated our baseline scenario simulations across these 50 parameter sets and

completed 50 simulations per parameter set for all sensitivity analyses.

Due to the high computational effort required to perform sensitivity simulations across all

management scenarios, we focused only on the proactive vaccination scenarios. Specifically,

we evaluated a subset of 12 LHS parameter sets across a subset of the proactive vaccination

conditions in a factorial design (see supplementary methods for further details), resulting in 108

parameter sets with 50 full simulations per parameter set (5,400 full simulations). This approach

allowed us to examine sensitivity of our proactive vaccination results across different outbreak

sizes and network and transmission parameters, while mitigating computational effort

associated with exploring such a wide range of parameters and scenario variations. Sensitivity

analysis simulation results were evaluated using scatterplots and Partial Rank Correlation

Coefficients (PRCC; 34, 35); proactive vaccination scenarios were evaluated for alignment with

our qualitative result that low proactive vaccination can increase disease-induced mortalities.

Results

Progressive infections are expected to result in death in panthers, and are therefore of

key concern for management efforts. For simplicity, hereafter, we refer to the number of

progressive infections in simulations as the number of mortalities. In the baseline, no-

intervention scenario, the median number of mortalities was 34 (range: 1-54); median duration

of epidemics was 119.5 weeks, and 34 epidemics failed (fewer than 5 progressive or regressive

infections) per 100 successful epidemics.

Proactive vaccination alone

Proactive vaccination paradoxically increased the number of mortalities across a range

of conditions, especially without vaccine boosting (Figures 2, S2). Even with 50% of vaccinates

receiving a booster, proactive vaccination only reduced mortalities from the baseline scenario at

high levels of population vaccination (i.e., 60-80%). With 100% boosting, proactive vaccination

increased mortalities at low levels of population vaccination (10-20%), had marginal effects at

30-40% population vaccination, and was strongly effective at about 50% population vaccination

levels and higher (at 80% population vaccination, median 17.5 mortalities versus 34 with no

interventions). Proactive vaccination consistently lengthened the duration of epidemics, relative

to the baseline scenario (up to a median duration of 143.5 weeks; Figures S3-4). When all

vaccinated individuals received a booster, vaccination reduced the probability of a successful

outbreak even at 40% population vaccination (52 failures versus 34 failures per 100 successful

epidemics; Figure S13). In contrast, when no vaccinated individuals received a booster,

proactive vaccination was largely only effective at reducing the probability of epidemics at very

high levels of population vaccination.

Figure 2: Histograms of FeLV mortalities in simulated epidemics with proactive vaccination

alone. Red vertical lines indicate the median number of mortalities from the baseline scenario

without interventions; blue lines, scenarios with proactive vaccination. Panel rows represent the

proportion of the population proactively vaccinated; columns represent ratios of vaccine efficacy

among vaccinates. Each histogram plot represents the results of 100 simulations. An expanded

version of this figure is available in the supplementary materials (Figure S2).

Proactive and reactive vaccination

Reactive vaccination alone did not reduce mortalities (Figure 3). However, reactive

vaccination appeared to work synergistically with proactive vaccination, particularly at moderate

to high levels of proactive vaccination (i.e., at least 40-60% of the population proactively

vaccinated). This largely held true regardless of the timing of intervention onset and the strategy

for reactive vaccination distribution (i.e., random versus spatial; Figures S5-6). The mortality-

reducing effects of reactive vaccination were, however, slightly reduced if reactive vaccination

occurred for only 6 months out of the year (Figure 3). A ratio of greater than 1.5 inoculations per

vaccinated individual appeared to promote the largest reductions in mortalities (e.g., as few as a

median of 25 mortalties with year-round reactive vaccination and 60% proactive vaccination

versus 34 mortalities with no interventions; Figure 3). Adding reactive vaccination largely did not

affect the durations of simulated epidemics (Figures S7-8), and had little impact on the

probability of epidemics failing, particularly in comparison to proactive vaccination alone (Figure

S13).

Figure 3: *Histograms of FeLV mortalities in simulated epidemics with both reactive and*

proactive vaccination. Results shown are for randomly distributed reactive vaccination starting

52 weeks after epidemic onset. Panel rows represent the proportion of the population

proactively vaccinated (with 50% boosted vaccination); columns represent duration of reactive

vaccination per year. Red vertical lines indicate the median number of mortalities from

simulations without interventions; blue lines for proactive vaccination alone; green lines for the

given combination of reactive and proactive vaccination (note green lines overlap red lines in

the 0% proactive vaccination scenarios). Numbers in the upper right of each histogram

represent (above) median total number of vaccinated individuals (both proactive and reactive),

and (below) median value for the total vaccinations used per total vaccinated individuals.

Proactive vaccination with test-and-removal

317  Test-and-removal alone did not reduce mortalities (Figure 4). Like reactive vaccination,

318 however, test-and-removal appeared to work synergistically with proactive vaccination,

319 especially at moderate to high levels of proactive vaccination (i.e., at least 40-60% of the

320 population proactively vaccinated; as few as a median of 26 mortalities). This largely held true

321 regardless of the timing of the onset of the intervention or the targeting of captures (i.e., random

322 versus spatial). Notably, simulated captures were only conducted for about 4 months per

323 simulation year, in contrast to at least 6 months of reactive vaccination per year. Captures were

324 marginally more likely to successfully identify actively infectious individuals when initiated earlier

325 in an outbreak (Figure S9). The addition of test-and-removal largely did not affect the durations

326 of epidemics (Figure S10). When coupled with proactive vaccination, test-and-removal had a

327 modest effect in reducing the probability of a successful epidemic (e.g., maximum of 50 failed

328 epidemics per 100 successful versus 34 failed epidemics per 100 successes with no

329 interventions; Figure S13).

Figure 4: *Histograms of FeLV mortalities in simulated epidemics with both reactive test-and-*

removal and proactive vaccination. Panel rows represent the proportion of the population

proactively vaccinated (with 50% boosted vaccination); columns represent the capture and

testing targeting strategy (random versus spatially targeted). Vertical red lines indicate the

median number of mortalities from simulations without interventions; blue lines for proactive

vaccination alone; purple lines for the given combination of test and removal and proactive

vaccination. Numbers in the upper right of each histogram represent the median value for the

proportion of capture events that resulted in a removal or humane euthanasia.

Reactive underpass closures

Reactive underpass closures were ineffective in the absence of proactive vaccination

(Figure S11). The most effective impact of underpass closures on reducing FeLV mortalities

occurred when onset of closure was early (26 weeks after epidemic initiation), lasted for at least

13 weeks, and occurred in conjunction with at least 40-60% of the population being proactively

vaccinated (though the clearest effects occurred when at least 60% were proactively

vaccinated). Under these conditions, underpass closures synergistically reduced mortalities

from baseline scenarios (as few as a median of 22 mortalities with underpass closures) and

increased the probability that an epidemic would fail (maximum of 75 failed epidemics per 100

successes with underpass closures; Figure S13). In some cases, simulations showed a marked

decrease in transmission during the period of underpass closures, but reopening often resulted

in a subsequent resurgence of infections (Figure 5). Underpass closures had no clear impact on

the duration of outbreaks (Figure S12).

Figure 5: *Epidemic curves showing the proportion of the population progressively infected over*

time when 60% of the population was proactively vaccinated. Lighter lines show individual

simulation results; dark lines show mean values across all simulations. Vertical black dashed

lines show when underpasses were closed (left) and reopened (right). Panel A shows

underpasses closed for 26 weeks starting 26 weeks after an epidemic started; panel B shows

underpasses closed for 26 weeks starting at 52 weeks.

Sensitivity analyses

Simulated epidemic sizes were variable across the full 50 sensitivity analysis parameter

sets in the absence of FeLV interventions (range: median 7.5 mortalities-median 47.5

mortalities; Figure S14). Based on PRCC, the parameters for network density, transmission

potential from regressives, weekly contact rates, and baseline transmission potential were

positively associated with median mortalities in the absence of interventions; the parameter for

the infection-induced mortality rate was negatively associated with median mortalities (see

supplementary results, Figure S15).

When focusing on a subset of parameter sets for proactive vaccination sensitivity

analysis, low levels of proactive vaccination (e.g., 20% population proactive vaccination) were

sometimes effective in reducing the number of mortalities, in contrast to our primary results.

PRCC results from proactive scenario sensitivity analysis suggested that the parameters for

network density, transmission potential from regressives, and weekly contact rates were

positively correlated with increased mortalities at low levels of vaccination (Figure S16).

However, network density did not have a clearly monotonic relationship with the difference

between mortalities with and without proactive vaccination (Figure S17). We therefore

performed additional *post-hoc* sensitivity analyses to further interrogate the relationship between

network density and our qualitative outcome of increased mortalities at low levels of proactive

vaccination (see supplementary results for detailed discussion of *post-hoc* analysis). These

additional analyses found some evidence that low levels of proactive vaccination were least

effective at reducing mortalities at intermediate values of parameters governing network

connectivity (network density and proportion adults), especially when coupled with increased

transmission potential (e.g. higher infectiousness of regressive individuals and/or increased

weekly contact rates; Figures S19-20).

Discussion

In this study we found unexpected consequences and impacts of several epidemic

management strategies for disease control in small populations of conservation concern.

Although our simulation results provide guidelines for FeLV management in Florida panthers,

they also demonstrate the power of partnering modeling approaches and population

management questions to test and optimize disease control strategies in free-ranging wildlife

(36). Furthermore, the principles of transmission and available methods of disease control

underlying our findings provide insights for pathogen control in other host-pathogen systems,

including humans and livestock.

Proactive vaccination alone may worsen epidemic outcomes under some conditions

Our simulation results showed a paradoxical increase in FeLV mortalities with low levels

of proactive vaccination. This counterintuitive finding is likely due, at least in part, to partial

vaccine immunity, a type of vaccine imperfection often overlooked in studies of wildlife disease

(32). Under partial vaccine immunity, vaccinates can act as a semi-protected susceptible pool,

contracting infection later in the course of an epidemic, ultimately prolonging epidemics and

increasing the total number of mortalities. Only with adequate herd immunity can these effects

be avoided. This counterintuitive result is consistent with findings by Rees et al. (37), who found

reduced vaccine effectiveness with heterogeneity in the spatial distribution of hosts. Importantly,

our results suggest that studies assuming full protection from vaccination may significantly

underestimate the level of vaccination needed for population protection—underestimates which

may even make epidemics worse if only partial vaccine immunity can be achieved.

Our sensitivity analysis suggests that populations with intermediate levels of network

connectivity and/or high transmission potential may be most vulnerable to these paradoxical

effects. As in spatially structured populations (38), high or low connectivity causes infection to

fade out quickly; with intermediate connectivity, vaccine failures provide a steady supply of new

susceptibles. Alternatively, in cases of high transmission potential, vaccination may shift a rapid

fade-out epidemic to a sustained epidemic scenario, as in Rees et al. (37). While our sensitivity

analysis was limited by computational demands and use of some discrete parameters (which

may affect PRCC inference; 34), our findings are consistent with this broader body of literature.

FeLV vaccine efficacy may operate differently in reality from our simulation structure: for

example, some vaccinates may have zero vaccine-induced immunity, while others have 100%

protection (binary immunity; 32). Alternatively, vaccination may not protect from infection but

could reduce viral shedding or increase survival of infected individuals (32). In the case of binary

immunity, vaccine efficacy would be unlikely to prolong and worsen epidemics as we saw here.

In contrast, increased survival of infected individuals without changes to shedding potential

could extend or worsen outbreaks, and even favor the evolution of virulence (32). Future

research should assess vaccinated panthers' immune response to FeLV infection *in vitro* to help

refine understanding of vaccine imperfection in panthers and how imperfect immunity may

further alter vaccination guidelines.

Based on our results, we argue that wildlife managers should continue vaccinating

available panthers and prioritize boosting vaccinated individuals to develop a core population of

high-immunity individuals, rather than a broadly distributed low-immunity population. This

recommendation should be most effective at increasing the probability of epidemic failure, and

aligns with other wildlife studies which have emphasized vaccination of core populations (39) or

risk-based sub-populations (13).

Temporary spatial restrictions are unlikely to be effective under realistic scenarios

Here, we examined the effect of wildlife underpass closures as a novel method to restrict

connectivity of the panther population under emergency disease control conditions. Such

temporary spatial restrictions could increase other types of panther mortality (e.g. vehicle strikes

or intraspecific conflict), so such an intervention would need to reduce FeLV mortalities at a rate

greater than these other types of panther mortality in order to be a viable disease control

strategy. Unfortunately, our simulations found that temporary underpass closures were

generally no more effective at reducing FeLV mortalities than less risky reactive interventions.

Further, underpass closures were less effective when occurring after the peak of simulated

epidemics, such that spatial restrictions would need to occur early in an epidemic to be

effective. However, it should be noted that spatial restrictions may be more effective if used in

combination with other reactive FeLV management strategies (e.g., reactive vaccination in

concert with temporary spatial restrictions to prevent transmission resurgence after restrictions

are removed).

Extensive research has considered how landscape barriers and fragmentation may

affect pathogen transmission and control in wildlife (e.g., 37, 40–42). By closing underpasses,

we artificially fragmented panther habitat, assuming complete efficacy of closures in preventing

transmission across the I-75 freeway. In reality, some individuals would likely successfully

traverse this barrier. Because habitat fragmentation can promote pathogen persistence (42), a

"semipermeable" freeway barrier may, in fact, ultimately worsen epidemic dynamics.

Alternatively, landscape restrictions to animal movement can facilitate more efficient disease

control, as in Haydon *et al.* (7), where vaccinating to control pathogen spread along habitat

corridors reduced the level of vaccine coverage needed to protect endangered Ethiopian

wolves. While panthers are generally well-connected spatially, expansion of the population north

of its current range may favor similar metapopulation-oriented disease control strategies.

However, given uncertainties in the effect of underpass closures on other sources of mortalities

and their limited effectiveness in our simulations, we suggest temporary underpass closures

should currently be considered a low priority for FeLV management in panthers.

Reactive and proactive strategies can work synergistically to reduce epidemic impacts

Our simulations showed that both reactive vaccination and test-and-removal strategies

reduced FeLV mortalities in panthers when used in combination with moderate levels of

proactive vaccination. Test-and-removal had more consistent effects with arguably less effort

than reactive vaccination (four months of captures compared to year-round reactive

vaccination). We therefore suggest that test-and-removal be prioritized over reactive

vaccination, especially if identification of actively infectious individuals can be improved.

In our simulations, captures that were most aligned with the initial wave of infectious

individuals (i.e., with earlier onset) were more likely to identify actively infectious individuals for

removal. This finding highlights the importance of targeting captures to individuals likely to be

actively infectious. However, determining infection status in cryptic wildlife is difficult, and

consequently supports the increased use of remote tracking technologies that may be able to

(1) identify behavior changes associated with sickness, and (2) detect the onset of an epidemic

more quickly. This conclusion is consistent with similar findings in Channel Island foxes, where

increasing the number and frequency of tracking of sentinel individuals was important for early

identification of epidemics (16).

A key component of the success of test-and-removal here is the selective removal we

simulated, which avoids removing immune individuals that contribute to overall herd immunity

(18, 20). However, we have simplified the field testing process in our simulations. The common

field-available FeLV diagnostic test identifies antigenemia, which is key for identifying actively

infectious individuals. However, the duration of antigenemia—and even degree of

infectiousness—in regressively infected individuals is unclear in panthers. We may therefore

overestimate the effect of removing regressive individuals, but given their reduced

infectiousness in our simulations, we still expect test-and-removal to be a key strategy for

mitigating FeLV impacts in panthers.

Notably, reactive vaccination, in concert with proactive vaccination, is also a viable

alternative strategy to test-and-removal. Our simulations showed reactive vaccination to have

the strongest effects for reducing FeLV mortalities when at least 50% of the population was

vaccinated, and with a ratio of about 1.5 vaccines per vaccinated individual (i.e., 50% of

vaccinated individuals received a booster). We therefore suggest that managers should

prioritize boosting at least half of vaccinated individuals in a reactive vaccination response

scenario. We did not see a strong effect of attempting a vaccine barrier, in contrast to Sanchez

et al (16), where a simulated vaccine barrier could effectively halt spread of a pathogen in

Channel Island foxes. This difference is likely due to the differences in home range size and

movement capacity between the two species, with foxes ranging less widely than panthers. In

reality, if an empirical outbreak of FeLV in panthers exhibited a stronger spatial signal than was

featured in our simulations, spatially targeted reactive vaccination may yet be a worthwhile

intervention strategy.

Importantly—particularly if levels of population protection from proactive vaccination are

unknown—both reactive vaccination and test-and-removal strategies mitigated the negative

effects of low levels of population protection seen with inadequate proactive vaccination. It is

therefore vital for managers to incorporate these reactive strategies in the event of future FeLV

outbreaks.

Limitations and future directions

In this study, we considered the effects of partial vaccine immunity, but we made the

simplifying assumption of no waning vaccine or infection-induced immunity over time. However,

our findings with regard to imperfect efficacy are at least partially representative of the likely

consequences of waning immunity, in that the loss of immunity supplies new susceptibles to the

population. Should immunity not outlast the course of an FeLV outbreak, this process could

prolong outbreaks and result in increased mortalities. Future research should therefore examine

the effects of waning vaccine immunity, particularly considering the value of revaccinating

individuals which may be experiencing loss of vaccine protection.

Our simulation results found that relatively high levels of vaccination were required to

reduce impacts of FeLV in panthers, in contrast to studies in other endangered species (7, 9).

However, here we are investigating a pathogen with a long duration of infectiousness and lack

the advantages of distinct corridors between panther sub-populations for reducing required

levels of vaccination. It is therefore unsurprising that panthers would require higher levels of

FeLV population vaccination than was found, for example, for rabies vaccination in Ethiopian

wolves (7) or canine distemper virus vaccination in Amur tigers (9). However, we also did not

assume the presence of preexisting population immunity prior to proactive vaccination, and

some degree of population immunity likely already exists in panthers, given ongoing exposures

(4). This would reduce necessary vaccination levels in panthers, as would higher vaccine

efficacy than we conservatively assumed here (39). Future research could prioritize better

understanding the realized individual immunity after single and boosted FeLV vaccination in

panthers to determine if inoculations are more protective than we conservatively assumed here.

Conclusions

Our simulation results highlight the risks of inadequate proactive vaccination, particularly

with partial vaccine immunity. We recommend prioritizing boosted vaccination in panthers, and

joint use of proactive vaccination and reactive strategies to mitigate the risks of imperfect

vaccination and most effectively reduce the impacts of FeLV in this iconic carnivore. This

research highlights the value of linking modeling and management priorities to identify

unexpected consequences of interventions and determine optimal pathogen management

strategies in free-ranging wildlife.

Authors' Contributions

All authors conceived the ideas and designed methodology; MLJG performed simulations,

analyzed data, and led the writing of the manuscript. All authors contributed critically to the

drafts and gave final approval for publication.

Acknowledgements

This research was supported by the National Science Foundation (DEB-1413925, 1654609, and

2030509). MLJG was supported by the Office of the Director, National Institutes of Health (NIH

T32OD010993), the University of Minnesota Informatics Institute MnDRIVE program, and the

Van Sloun Foundation. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health. Florida panther data

collected by the Florida Fish and Wildlife Conservation Commission is fully supported by

donations to the Florida Panther Research and Management Trust Fund via the registration of

"Protect the Panther" license plates. We acknowledge the efforts of National Park Service staff

in the collection of Florida panther data utilized in this study.

Data Availability

Full R code for simulations is available on GitHub

(https://github.com/mjones029/FeLV_Management_Simulations) and upon acceptance will be

archived at Zenodo.

Works Cited

1. A. C. Breed, *et al.*, "Disease Management in Endangered Mammals" in *Management of Disease in Wild Mammals*, R. J. Delahay, G. C. Smith, M. R. Hutchings, Eds. (Springer Japan, 2009), pp. 215–239.

2. M. W. Cunningham, *et al.*, Epizootiology and management of feline leukemia virus in the
Florida puma. *J. Wildl. Dis.* **44**, 537–552 (2008).

3. M. A. Brown, *et al.*, Genetic characterization of feline leukemia virus from Florida panthers.
Emerg. Infect. Dis. **14**, 252–259 (2008).

4. E. S. Chiu, *et al.*, Multiple Introductions of Domestic Cat Feline Leukemia Virus in
Endangered Florida Panthers. *Emerg. Infect. Dis.* **25**, 92–101 (2019).

5. H. McCallum, Models for managing wildlife disease. *Parasitology* **143**, 805–820 (2016).

6. J. O. Lloyd-Smith, *et al.*, Epidemic dynamics at the human-animal interface. *Science* **326**,
1362–1367 (2009).

7. D. T. Haydon, *et al.*, Low-coverage vaccination strategies for the conservation of
endangered species. *Nature* **443**, 692–695 (2006).

8. J. Rushmore, *et al.*, Network-based vaccination improves prospects for disease control in
wild chimpanzees. *J. R. Soc. Interface* **11**, 20140349 (2014).

9. M. Gilbert, *et al.*, Distemper, extinction, and vaccination of the Amur tiger. *Proc. Natl. Acad.*
Sci. U. S. A. (2020) https:/doi.org/10.1073/pnas.2000153117.

10. Robinson Stacie J., *et al.*, Model recommendations meet management reality:
implementation and evaluation of a network-informed vaccination effort for endangered
Hawaiian monk seals. *Proceedings of the Royal Society B: Biological Sciences* **285**,
20171899 (2018).

11. L. Baker, J. Matthiopoulos, T. Müller, C. Freuling, K. Hampson, Optimizing spatial and
seasonal deployment of vaccination campaigns to eliminate wildlife rabies. *Philos. Trans.*
R. Soc. Lond. B Biol. Sci. **374**, 20180280 (2019).

12. S. Cleaveland, Viral threats and vaccination: disease management of endangered species.
Anim. Conserv. **12**, 187–189 (2009).

13. H. L. Beyer, *et al.*, The implications of metapopulation dynamics on the design of
vaccination campaigns. *Vaccine* **30**, 1014–1022 (2012).

14. K. M. McClure, A. T. Gilbert, R. B. Chipman, E. E. Rees, K. M. Pepin, Variation in host
home range size decreases rabies vaccination effectiveness by increasing the spatial
spread of rabies virus. *J. Anim. Ecol.* **89**, 1375–1386 (2020).

15. E. J. Newton, *et al.*, Differential impacts of vaccination on wildlife disease spread during
epizootic and enzootic phases. *J. Appl. Ecol.* **56**, 526–536 (2019).

16. J. N. Sanchez, B. R. Hudgens, Vaccination and monitoring strategies for epidemic
prevention and detection in the Channel Island fox (*Urocyon littoralis*). *PLoS One* **15**,
e0232705 (2020).

17. D. Slate, *et al.*, Status of oral rabies vaccination in wild carnivores in the United States.
Virus Res. **111**, 68–76 (2005).

18. E. Miguel, *et al.*, A systemic approach to assess the potential and risks of wildlife culling for

infectious disease control. *Commun Biol* **3**, 353 (2020).

19.   J. O. Lloyd-Smith, *et al.*, Should we expect population thresholds for wildlife disease?
Trends Ecol. Evol. **20**, 511–519 (2005).

20.   A. Potapov, E. Merrill, M. A. Lewis, Wildlife disease elimination and density dependence.
Proc. Biol. Sci. **279**, 3139–3145 (2012).

21.   R. Woodroffe, C. A. Donnelly, D. R. Cox, Effects of culling on badger Meles meles spatial
organization: implications for the control of bovine tuberculosis. *Journal of Applied* (2006).

22.   S. Little, *et al.*, 2020 AAFP Feline Retrovirus Testing and Management Guidelines. *J. Feline*
Med. Surg. **22**, 5–30 (2020).

23.   G. López, *et al.*, Management measures to control a feline leukemia virus outbreak in the
endangered Iberian lynx. *Anim. Conserv.* **12**, 173–182 (2009).

24.   A. Mysterud, C. M. Rolandsen, Fencing for wildlife disease control. *J. Appl. Ecol.* **56**, 519–
525 (2019).

25.   G. G. Cotterill, *et al.*, Winter feeding of elk in the Greater Yellowstone Ecosystem and its
effects on disease dynamics. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **373** (2018).

26.   M. L. J. Gilbertson, *et al.*, Transmission of one predicts another: Apathogenic proxies for
transmission dynamics of a fatal virus. *bioRxiv* (2021)
https:/doi.org/10.1101/2021.01.09.426055.

27.   B. T. McClintock, D. P. Onorato, J. Martin, Endangered Florida panther population size
determined from public reports of motor vehicle collision mortalities. *J. Appl. Ecol.* **52**, 893–
901 (2015).

28.   M. E. Craft, H. L. Beyer, D. T. Haydon, Estimating the probability of a major outbreak from
the timing of early cases: an indeterminate problem? *PLoS One* **8**, e57878 (2013).

29.   R Core Team, R: A Language and Environment for Statistical Computing (2018).

30.   A. H. Sparkes, Feline leukaemia virus: a review of immunity and vaccination. *J. Small Anim.*
Pract. **38**, 187–194 (1997).

31.   A. N. Torres, C. K. Mathiason, E. A. Hoover, Re-examination of feline leukemia virus: host
relationships using real-time PCR. *Virology* **332**, 272–283 (2005).

32.   K. M. Barnett, D. J. Civitello, Ecological and Evolutionary Challenges for Wildlife
Vaccination. *Trends Parasitol.* **36**, 970–978 (2020).

33.   R. Carnell, lhs: Latin hypercube samples. *R package version 0. 10, URL http://CRAN. R-*
project.org/package=lhs (2012).

34.   S. Marino, I. B. Hogue, C. J. Ray, D. E. Kirschner, A methodology for performing global
uncertainty and sensitivity analysis in systems biology. *J. Theor. Biol.* **254**, 178–196 (2008).

35.   J. Wu, R. Dhingra, M. Gambhir, J. V. Remais, Sensitivity analysis of infectious disease
models: methods, advances and their application. *J. R. Soc. Interface* **10**, 20121018

(2013).

36. M. B. Joseph, *et al.*, Taming wildlife disease: bridging the gap between science and
management. *J. Appl. Ecol.* **50**, 702–712 (2013).

37. E. E. Rees, B. A. Pond, R. R. Tinline, D. Bélanger, Modelling the effect of landscape
heterogeneity on the efficacy of vaccination for wildlife infectious disease control. *J. Appl.*
Ecol. **50**, 881–891 (2013).

38. H. McCallum, Landscape structure, disturbance, and disease dynamics. *Infectious disease*
ecology: effects of ecosystems on disease and of disease on ecosystems, 100–122 (2008).

39. F. Vial, S. Cleaveland, G. Rasmussen, D. T. Haydon, Development of vaccination
strategies for the management of rabies in African wild dogs. *Biol. Conserv.* **131**, 180–192
(2006).

40. D. L. Smith, B. Lucey, L. A. Waller, J. E. Childs, L. A. Real, Predicting the spatial dynamics
of rabies epidemics on heterogeneous landscapes. *Proc. Natl. Acad. Sci. U. S. A.* **99**,
3668–3672 (2002).

41. J. A. Tracey, S. N. Bevins, S. VandeWoude, K. R. Crooks, An agent-based movement
model to assess the impact of landscape fragmentation on disease transmission.
Ecosphere **5**, 1–24 (2014).

42. L. A. White, J. D. Forester, M. E. Craft, Disease outbreak thresholds emerge from
interactions between movement behavior, landscape structure, and epidemiology. *Proc.*
Natl. Acad. Sci. U. S. A. **115**, 7374–7379 (2018).