Paradoxes and synergies: optimizing management of a deadly virus in an endangered

carnivore

Marie L.J. Gilbertson[1]*, Dave Onorato[2], Mark Cunningham[3], Sue VandeWoude[4], Meggan E.

Craft[1,5]

*Corresponding author: jone1354@umn.edu

[1]Department of Veterinary Population Medicine, University of Minnesota, St Paul, MN 55108.

[2]Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission,

Naples, FL 34114.

[3]Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission,

Gainesville, FL 32601.

[4]Department of Microbiology, Immunology, and Pathology, Colorado State University, Fort

Collins, CO 80523

[5]Department of Ecology, Evolution and Behavior, University of Minnesota, St Paul, MN 55108.

Keywords

adaptive management; disease model; simulation modeling; network modeling; feline leukemia

virus; vaccination; test-and-removal; *Puma concolor coryi*

Abstract

1. Outbreaks of infectious disease can have serious consequences for wildlife population

health, especially species of conservation concern. The endangered Florida panther, for

example, experienced an outbreak of feline leukemia virus (FeLV) in 2002-04, and

continues to be affected by this deadly virus. Ongoing management efforts aim to

mitigate the effects of FeLV on panthers, but with limited information about which

strategies may be most effective and efficient. Such uncertainties can benefit from

linking management with simulation models in order to optimize conservation efforts for

populations of concern.

2. We used a simulation-based approach to determine optimal FeLV management

strategies. We simulated use of proactive FeLV management strategies (i.e., proactive

vaccination) and several reactive strategies, including reactive vaccination and test-and-

removal. Vaccination strategies accounted for partial vaccine immunity, including when

using single or boosted inoculations. We compared the effectiveness of these different

strategies in mitigating the number of FeLV mortalities and the duration of outbreaks.

3. Results showed that inadequate proactive vaccination can paradoxically increase the

number of disease-induced mortalities in FeLV outbreaks, most likely due to effects of

partial vaccine immunity. Combinations of proactive vaccination with reactive test-and-

removal or vaccination had a synergistic effect in reducing impacts of FeLV outbreaks.

Temporary spatial restrictions, however, were unlikely to be effective under realistic

conditions.

4. *Synthesis and applications:* Management-informed disease simulations demonstrated

both unexpected negative consequences and synergies with active management

strategies for a deadly virus in Florida panthers. We recommend a combination of

proactive and reactive management approaches, and suggest prioritizing boosted

vaccination over broad distribution of unboosted inoculations. Our results highlight the

importance of integrating management and modeling approaches to aid in conservation

of at-risk species.

Introduction

Outbreaks of infectious diseases can have significant impacts on the population health of free-ranging wildlife and are of heightened importance in species of conservation concern (Breed et al., 2009). For example, the feline retrovirus, feline leukemia virus (FeLV), has been the source of significant outbreaks in two endangered felids: Iberian lynx (*Lynx pardinus*) and Florida panthers (*Puma concolor coryi*). In the case of panthers, FeLV caused a deadly outbreak in 2002-2004 (Cunningham et al., 2008), spilling over from domestic cats with subsequent direct transmission among panthers (Brown et al., 2008). In addition, recent evidence demonstrates ongoing FeLV spillover to and transmission among panthers (Chiu et al., 2019), necessitating continued management to prevent future epidemics of this deadly pathogen. FeLV inoculation with a domestic cat vaccine has been used previously in panthers but with unknown efficacy (Cunningham et al., 2008). Further, the proportion of the population that must be vaccinated to most efficiently prevent future FeLV outbreaks is unknown, as is how proactive vaccination might interact with other reactive interventions to interrupt an FeLV outbreak. Such uncertainties are common among free-ranging wildlife threatened by infectious disease and hamper efforts to effectively control pathogen transmission.

Mathematical models of infectious disease transmission have been used to optimize disease management protocols in a variety of free-ranging wildlife species of conservation concern, including Ethiopian wolves (Haydon et al., 2006), chimpanzees (Rushmore et al., 2014), and Amur tigers (Gilbert et al., 2020). Further, models can serve the important function of balancing the realities of fieldwork with ideal disease control protocols to provide practical, effective guidance for wildlife managers (e.g., Robinson et al., 2018). Here, we use mathematical models of FeLV transmission in Florida panthers to determine optimal disease prevention and control strategies in this iconic carnivore.

Pathogen management strategies can be preventive (hereafter, *proactive*) or reactive. Among proactive and reactive strategies, vaccination is a cornerstone in veterinary medicine

(Breed et al., 2009) and functions by reducing the availability of susceptible individuals.

Vaccination can be applied randomly, or targeted to: high risk individuals or populations (e.g.,

Beyer et al., 2012); likely superspreaders (e.g., in chimpanzees: Rushmore et al., 2014);

seasonal dynamics (e.g., Baker et al., 2019); or spatial risk, as with vaccine barriers to prevent

spread of a pathogen to new areas or populations (e.g., Sanchez & Hudgens, 2020; Slate et al.,

2005).

Among reactive pathogen management strategies, test-and-removal (or test-and-cull) is

commonly used in domestic species to functionally reduce the infectious period of infected

individuals (Miguel et al., 2020). However, this approach is rarely used in wildlife due to the

generally low availability of field diagnostic tests and difficulty in recapturing individuals after a

positive diagnosis (Miguel et al., 2020). Rather, pathogen control in wildlife often relies on non-

selective culling in an attempt to reduce density and—in theory—transmission (Lloyd-Smith et

al., 2005). However, non-selective culling can have myriad negative consequences for disease

management including removal of immune individuals (Miguel et al., 2020; Potapov et al., 2012)

or culling-induced perturbation (e.g., Woodroffe et al., 2006). Thanks to ready availability of a

field diagnostic test, FeLV management in domestic cats and Iberian lynx has made use of test-

and-removal or isolation of infected individuals (Little et al., 2020; López et al., 2009), and is part

of future FeLV mitigation plans in panthers. However, uncertainties remain regarding sufficiency

of test-and-removal alone to mitigate pathogen spread or how it might interact with proactive

vaccination efforts to more effectively reduce pathogen impacts.

Given the ongoing risks of FeLV to panther conservation and the uncertainties regarding

application of proactive and reactive FeLV management strategies, the objectives of this study

were to test singly, and in combination, the effectiveness of: (1) proactive vaccination, (2)

reactive vaccination, and (3) reactive test-and-removal for reducing the population level impacts

of FeLV in Florida panthers. In addition, we evaluated the effectiveness of temporary panther

spatial restrictions via closure of wildlife highway underpasses, which we describe and discuss

in the supplementary materials. We used a simulation modeling approach to address our

objectives in order to efficiently and ethically test a wide range of intervention protocols for the

protection of this endangered carnivore.

Methods

Simulation pipeline

To examine the effect of different disease management regimes on FeLV control, we

used spatially explicit, network simulations adapted from our research determining drivers of

retrovirus transmission in panthers (Gilbertson et al., unpublished preprint). This approach

involves two steps: (1) simulation of a contact network among panthers, and (2) simulation of

FeLV transmission through this network. In brief, we simulated panther populations of 150

individuals (McClintock et al., 2015) and used our previously described exponential random

graph model for retrovirus transmission in panthers (Gilbertson et al., unpublished preprint) to

simulate contact networks constrained by network density among simulated populations (see

supplementary methods for further details and Table S1 for parameters).

Briefly, we used a stochastic chain binomial transmission model with a modified

susceptible-infectious-recovered compartmental framework, where individuals could be

progressively, regressively, or abortively infected (Cunningham et al., 2008; model developed in

Gilbertson et al., unpublished preprint). Progressive infections always resulted in death, while

regressive infections eventually recovered with immunity, and abortive infections were always

considered immune. Importantly, based on our previous work, we allowed both progressives

and regressives to be infectious, though regressives were less likely to transmit (Table S1). The

infectious period is long for FeLV (estimated as a mean of 18 weeks, though based on limited

observations: Cunningham et al., 2008; see also Gilbertson et al., unpublished preprint, and

Table S1). We included only disease-induced mortality, and, in order to preserve key network

structure, allowed territories vacated by deaths to be reoccupied by new susceptible individuals

(hereafter, *respawning*). Outbreaks were initiated by a single, randomly selected non-isolate

individual in the population (see supplement discussion), and proceeded in weekly time steps

for up to five years.

Our primary objective was to examine the effect of different FeLV management regimes

on epidemic outcomes, so in our primary simulations, we held network generation and

transmission parameters constant at previously supported values (see supplementary methods

and Table S1 for further details). We evaluated the consistency of our results to the choice of

parameter values with a sensitivity analysis (see below). Hereafter, a *parameter set* represents

the unique set of network and transmission parameters for any given set of simulations. A *full*

simulation includes simulation of a single contact network and FeLV transmission through that

network (with or without management interventions). For each parameter set, we performed 100

full simulations.

For *baseline* and *management scenarios*, we recorded key epidemic outcomes in the

absence and presence of interventions, respectively (Figure 1). These key outcomes were (1)

the number of mortalities, (2) the duration of an epidemic, and (3) the proportion of epidemics

that "failed" per 100 successful epidemics. A failed epidemic (i.e. stochastic fadeout; Craft et al.,

2013) was one in which fewer than 5 individuals acquired progressive or regressive infections.

As at least 5 individuals acquired progressive infections during the 2002-04 outbreak

(Cunningham et al., 2008), this fadeout cutoff allowed for realism in simulated outbreak sizes,

while also reducing noise among "successful" epidemics resulting from especially small

outbreaks. The outcomes of mortalities and epidemic durations were summarized as median

values per parameter set, as results were often skewed; all outcomes were compared between

baseline and management scenarios, including quantification with Mann-Whitney *U* tests

(supplementary results). All simulations were performed in R version 3.6.3 (R Core Team,

2018).

157

Figure 1: *Steps of the simulation process across baseline and management scenarios. The*

basic network and FeLV transmission simulation steps are shown in yellow, the baseline (no

intervention) scenario in green, and the different overarching management scenarios in blue.

Within each management scenario, we investigated several variations for the given approach

(shown in light blue) in a factorial design. Note that methods and results for reactive underpass

closures are described in the supplementary materials.

Management scenarios

We examined the effect of different levels of population proactive vaccination (proportion

of the population vaccinated prior to an outbreak) and different ratios of single versus boosted

vaccination. We simulated from 10-80% (in 10% increments) of the population having some

degree of vaccine-induced immunity to FeLV prior to the onset of an outbreak. These proactive

vaccinations were distributed randomly in the population. Among the vaccinated individuals, 0,

50, or 100% received a second boosting inoculation. Actual vaccine efficacies are unknown for

panthers, but based on efficacy studies in domestic cats (Sparkes, 1997; Torres et al., 2005),

we conservatively assumed that boosted vaccination would prevent 80% of infections and single

vaccination would prevent 40% of infections. We modeled this efficacy as a binomial probability

to represent partial vaccine immunity, wherein increased exposures to infectious individuals

increase the likelihood of vaccine failure (versus binary immunity wherein some vaccinated

individuals are 100% protected and others receive 0% protection; Barnett & Civitello, 2020).

In contrast, for reactive vaccination management scenarios, panthers were selected for

vaccination *during* outbreaks at a rate of one panther per week. We assumed that managers

would not know the disease status of an individual selected for vaccination and that vaccination

would be ineffective in infectious or recovered individuals. In the case of previously vaccinated

individuals, re-vaccination changed the vaccine efficacy of singly vaccinated individuals (efficacy

of 40%) to the efficacy for boosted individuals (efficacy of 80%). We varied the timing of the

onset of reactive vaccination after the initiation of an FeLV outbreak to reflect the difficulty of

epidemic detection in this elusive carnivore. We therefore began reactive vaccination at an

optimistic, but difficult-to-attain time point of 26 weeks, and a more realistic time point of 52

weeks. In addition, we varied the distribution of reactive vaccination. While proactive vaccines

were always distributed randomly, reactive vaccination was either randomly distributed or

spatially distributed in an attempted vaccine barrier based on proximity to the I-75 freeway,

which runs east-west through primary panther habitat (see supplement for further details).

Because vaccination is resource and time intensive, we evaluated the effect of reactive

vaccination for 6 months per year versus year-round.

Test-and-removal management scenarios were built around a protocol in which panthers

infectious at capture were removed from the population through humane euthanasia or

temporary removal until recovery. For simplicity, based on expected fatality of progressive

cases, we assumed that all progressively infected individuals would be humanely euthanized at

capture, while infectious regressive individuals were temporarily removed from the population

until their recovery and re-released into any open territory. A maximum of five individuals were

allowed to be temporarily removed in this way at one time based on expected capacity for

housing and care.

We expected that managers would be able to capture and test one panther per week at

most, and that captures occurred during a 17 week (about 4 month) capture season, in

accordance with current panther capture protocols. We assumed that managers would not know

the disease status of a target individual until the capture occurred, so captures were not

targeted by infection state, and removed individuals were able to transmit prior to removal. We

varied the onset of test-and-removal, such that the intervention began 26 or 52 weeks after the

initiation of an epidemic. Captures were also random or spatially targeted. If spatially targeted,

captures (and consequent removals) only occurred on the same side of the I-75 freeway as the

initial FeLV infection.

In addition, we simulated temporary spatial segregation of panthers via closure of wildlife

highway underpasses along the I-75 freeway, methods and results for which are reported in the

supplementary materials. To examine the effects of proactive and reactive management

interventions conducted in concert, we included proactive vaccination in all reactive

management scenarios: specifically, with 0-60% of the population proactively vaccinated (in

20% increments) prior to initiation of an outbreak. Because it is highly unlikely that 100% of the

proactively vaccinated population would have received boosted vaccination, in reactive

management scenarios we used a more conservative—yet still challenging to attain—ratio of

50% of the proactively vaccinated population being boosted (i.e. efficacy of 80%). We

conducted all management scenarios in a full factorial design resulting in 24, 32, and 16

parameter sets for proactive vaccination, reactive vaccination, and reactive test-and-removal

scenarios, respectively (100 full simulations per set for a total of 7,200 simulations).

Sensitivity analysis

We used a latin hypercube sampling (LHS) approach to generate 50 sensitivity analysis

parameter sets across our eight network and transmission parameters (Table S1) using the *lhs*

package in R (Carnell, 2012). We repeated our baseline scenario simulations across these 50

parameter sets and completed 50 simulations per parameter set for all sensitivity analyses.

Due to the high computational effort required to perform sensitivity simulations across all

management scenarios, we focused only on the proactive vaccination scenarios. Specifically,

we evaluated a subset of 12 LHS parameter sets across a subset of the proactive vaccination

conditions in a factorial design (see supplementary methods for further details), resulting in 108

parameter sets with 50 full simulations per parameter set (5,400 full simulations). This approach

allowed us to examine sensitivity of our proactive vaccination results across different outbreak

sizes and network and transmission parameters, while mitigating computational effort

associated with exploring such a wide range of parameters and scenario variations. Sensitivity

analysis simulation results were evaluated using scatterplots and Partial Rank Correlation

Coefficients (PRCC; Marino et al., 2008; Wu et al., 2013); proactive vaccination scenarios were

evaluated for alignment with our qualitative result that low proactive vaccination can increase

disease-induced mortalities.

Results

Progressive infections are expected to result in death in panthers, and are therefore of

key concern for management efforts. For simplicity, hereafter, we refer to the number of

progressive infections in simulations as the number of mortalities. In the baseline, no-

intervention scenario, the median number of mortalities was 34 (range: 1-54); median duration

of epidemics was 119.5 weeks, and 34 epidemics failed (fewer than 5 progressive or regressive

infections) per 100 successful epidemics.

Proactive vaccination alone

Proactive vaccination paradoxically increased the number of mortalities across a range

of conditions, especially without vaccine boosting (Figures 2, S2; Table S2). Even with 50% of

vaccinated individuals receiving a booster, proactive vaccination only reduced mortalities from

the baseline scenario at high levels of population vaccination (i.e., 60-80%). With 100%

boosting, proactive vaccination increased mortalities at low levels of population vaccination (10-

20%), had marginal effects at 30-40% population vaccination, and was strongly effective at

about 50% population vaccination levels and higher (at 80% population vaccination, median

17.5 mortalities versus 34 with no interventions). Proactive vaccination consistently lengthened

the duration of epidemics, relative to the baseline scenario (up to a median duration of 143.5

weeks; Figures S3-5, Table S3). Of those individuals which became infected in proactive

vaccination scenarios, susceptibles were typically infected earliest, followed by vaccinated

individuals, and susceptible respawned individuals (Figures S6). When all vaccinated individuals

received a booster, vaccination reduced the probability of a successful outbreak even at 40%

population vaccination (52 failures versus 34 failures per 100 successful epidemics; Figure

S16). In contrast, when no vaccinated individuals received a booster, proactive vaccination was

largely only effective at reducing the probability of epidemics at very high levels of population

vaccination.

Figure 2: *Histograms of FeLV mortalities in simulated epidemics with proactive vaccination*

alone. Red vertical lines indicate the median number of mortalities from the baseline scenario

without interventions; blue lines, scenarios with proactive vaccination. Panel rows represent the

proportion of the population proactively vaccinated; columns represent ratios of vaccine efficacy

among vaccinated individuals. Each histogram plot represents the results of 100 simulations. An

expanded version of this figure is available in the supplementary materials (Figure S2).

Proactive and reactive vaccination

Reactive vaccination alone did not reduce mortalities (Figure 3, Table S4). However,

reactive vaccination appeared to work synergistically with proactive vaccination, particularly at

moderate to high levels of proactive vaccination (i.e., at least 40-60% of the population

proactively vaccinated). This largely held true regardless of the timing of intervention onset and

the strategy for reactive vaccination distribution (i.e., random versus spatial; Figures S7-8). The

mortality-reducing effects of reactive vaccination were, however, slightly reduced if reactive

vaccination occurred for only 6 months out of the year (Figure 3). A ratio of greater than 1.5

inoculations per vaccinated individual appeared to promote the largest reductions in mortalities

(e.g., as few as a median of 25 mortalties with year-round reactive vaccination and 60%

proactive vaccination versus 34 mortalities with no interventions; Figure 3). Adding reactive

vaccination largely did not affect the durations of simulated epidemics (Figures S9-10, Table

S5), and had little impact on the probability of epidemics failing, particularly in comparison to

proactive vaccination alone (Figure S16).

290

Figure 3: *Histograms of FeLV mortalities in simulated epidemics with both reactive and*

proactive vaccination. Results shown are for randomly distributed reactive vaccination starting

52 weeks after epidemic onset. Panel rows represent the proportion of the population

proactively vaccinated (with 50% boosted vaccination); note that the bottom row, with 0%

proactive vaccination, demonstrates reactive vaccination alone. Columns represent duration of

reactive vaccination per year. Red vertical lines indicate the median number of mortalities from

simulations without interventions; blue lines for proactive vaccination alone; green lines for the

given combination of reactive and proactive vaccination (note green lines overlap red lines in

the 0% proactive vaccination scenarios). Numbers in the upper right of each histogram

represent (above) median total number of vaccinated individuals (both proactive and reactive),

and (below) median value for the total vaccinations used per total vaccinated individuals.

Proactive vaccination with test-and-removal

Test-and-removal alone did not reduce mortalities (Figure 4, Table S6). Like reactive

vaccination, however, test-and-removal appeared to work synergistically with proactive

vaccination, especially at moderate to high levels of proactive vaccination (i.e., at least 40-60%

of the population proactively vaccinated; as few as a median of 26 mortalities). This largely held

true regardless of the timing of the onset of the intervention or the targeting of captures (i.e.,

random versus spatial). Notably, simulated captures were only conducted for about 4 months

per simulation year, in contrast to at least 6 months of reactive vaccination per year. Captures

were marginally more likely to successfully identify actively infectious individuals when initiated

earlier in an outbreak (Figure S11). The addition of test-and-removal largely did not affect the

durations of epidemics (Figure S12, Table S7). When coupled with proactive vaccination, test-

and-removal had a modest effect in reducing the probability of a successful epidemic (e.g.,

maximum of 50 failed epidemics per 100 successful versus 34 failed epidemics per 100

successes with no interventions; Figure S16).

Figure 4: *Histograms of FeLV mortalities in simulated epidemics with both reactive test-and-*

removal and proactive vaccination. Panel rows represent the proportion of the population

proactively vaccinated (with 50% boosted vaccination); note that the bottom row, with 0%

proactive vaccination, demonstrates reactive vaccination alone. Columns represent the capture

and testing targeting strategy (random versus spatially targeted). Vertical red lines indicate the

median number of mortalities from simulations without interventions; blue lines for proactive

vaccination alone; purple lines for the given combination of test and removal and proactive

vaccination. Numbers in the upper right of each histogram represent the median value for the

proportion of capture events that resulted in a removal or humane euthanasia.

Sensitivity analyses

Simulated epidemic sizes were variable across the full 50 sensitivity analysis parameter

sets in the absence of FeLV interventions (range: median 7.5 mortalities–median 47.5

mortalities; Figure S17). Based on PRCC, the parameters for network density, transmission

potential from regressives, weekly contact rates, and baseline transmission potential were

positively associated with median mortalities in the absence of interventions; the parameter for

the infection-induced mortality rate was negatively associated with median mortalities (see

supplementary results, Figure S18).

When focusing on a subset of parameter sets for proactive vaccination sensitivity

analysis, low levels of proactive vaccination (e.g., 20% population proactive vaccination) were

sometimes effective in reducing the number of mortalities, in contrast to our primary results.

PRCC results from proactive scenario sensitivity analysis suggested that the parameters for

network density, transmission potential from regressives, and weekly contact rates were

positively correlated with increased mortalities at low levels of vaccination (Figure S19).

However, network density did not have a clearly monotonic relationship with the difference

between mortalities with and without proactive vaccination (Figure S20). We therefore

performed additional *post-hoc* sensitivity analyses to further interrogate the relationship between

network density and our qualitative outcome of increased mortalities at low levels of proactive

vaccination (see supplementary results for detailed discussion of *post-hoc* analysis). These

additional analyses found some evidence that low levels of proactive vaccination were least

effective at reducing mortalities at intermediate values of parameters governing network

connectivity (network density and proportion adults), especially when coupled with increased

transmission potential (e.g. higher infectiousness of regressive individuals and/or increased

weekly contact rates; Figures S22-23).

Discussion

In this study we found unexpected consequences and impacts of several epidemic

management strategies for disease control in small populations of conservation concern.

Although our simulation results provide guidelines for FeLV management in Florida panthers,

they also demonstrate the power of partnering modeling approaches and population

management questions to test and optimize disease control strategies in free-ranging wildlife

(Joseph et al., 2013). Furthermore, the principles of transmission and available methods of

disease control underlying our findings provide insights for pathogen control in other host-

pathogen systems, including humans and livestock.

Proactive vaccination alone may exacerbate epidemic outcomes under some conditions

Our simulation results showed a paradoxical increase in FeLV mortalities with low levels

of proactive vaccination. This counterintuitive finding is likely due, at least in part, to partial

vaccine immunity, a type of vaccine imperfection often overlooked in studies of wildlife disease

(Barnett & Civitello, 2020). Under partial vaccine immunity, vaccinated individuals acted as a

semi-protected susceptible pool, contracting infection later in the course of an epidemic,

ultimately prolonging epidemics and increasing the total number of mortalities. Further, these

effects likely interacted with the respawning process we used, which is similar to density-

dependent birth processes which have been shown to increase pathogen prevalence and

mortality under culling conditions (Choisy & Rohani, 2006). In the case of our simulations,

proactive vaccination flattened the epidemic curve (Figure S5), leading to slow, steady

introductions of both new susceptibles and vaccine "failures" which could themselves become

infected and maintain epidemics. Only with adequate herd immunity could increased mortalities

be avoided. This counterintuitive result is consistent with findings by Rees et al. (2013), who

found reduced vaccine effectiveness in some conditions in which hosts had heterogeneous

spatial distributions. Importantly, our results suggest that studies assuming full protection from

vaccination may significantly underestimate the level of vaccination needed for population

protection—underestimates which may even make epidemics worse if only partial vaccine

immunity can be achieved.

Our sensitivity analysis suggests that populations with intermediate levels of network

connectivity and/or high transmission potential may be most vulnerable to these paradoxical

effects. As in spatially structured populations (McCallum, 2008), high or low connectivity causes

infection to fade out quickly; with intermediate connectivity, vaccine failures provide a steady

supply of new susceptibles. Alternatively, in cases of high transmission potential, vaccination

may shift a rapid fade-out epidemic to a sustained epidemic scenario, as in Rees et al. (2013).

While our sensitivity analysis was limited by computational demands and use of some discrete

parameters (which may affect PRCC inference; Marino et al., 2008), our findings are consistent

with this broader body of literature.

FeLV vaccine efficacy may operate differently in reality from our simulation structure: for

example, some vaccinated individuals may have zero vaccine-induced immunity, while others

have 100% protection (binary immunity; Barnett & Civitello, 2020). Alternatively, vaccination

may not protect from infection but could reduce viral shedding or increase survival of infected

individuals (Barnett & Civitello, 2020). In the case of binary immunity, vaccine efficacy would be

unlikely to prolong and worsen epidemics as we saw here. In contrast, increased survival of

infected individuals without changes to shedding potential could extend or worsen outbreaks,

and even favor the evolution of virulence (Barnett & Civitello, 2020). Future research should

assess vaccinated panthers' immune response to FeLV infection *in vitro* to help refine

understanding of vaccine imperfection in panthers and how imperfect immunity may further alter

vaccination guidelines.

Based on our results, we argue that wildlife managers should continue vaccinating

available panthers and prioritize boosting vaccinated individuals to develop a core population of

high-immunity individuals, rather than a broadly distributed low-immunity population. This

recommendation should be most effective at increasing the probability of epidemic failure, and

aligns with other wildlife studies which have emphasized vaccination of core populations (Vial et

al., 2006) or risk-based sub-populations (Beyer et al., 2012). Given ongoing uncertainties in

vaccine efficacy and the duration of immunity, in the event of an outbreak, we recommend

updating model parameterizations to reflect on-the-ground realities in an adaptive management

framework in order to provide the most useful predictions and guidance for managers. For

example, observed "breakthrough" infection rates (infections in vaccinated individuals) would be

a key observation for updating model predictions and adapting management responses.

Reactive and proactive strategies can work synergistically to reduce epidemic impacts

Our simulations showed that both reactive vaccination and test-and-removal strategies

reduced FeLV mortalities in panthers when used in combination with moderate levels of

proactive vaccination. Test-and-removal had more consistent effects with arguably less effort

than reactive vaccination (4 months of captures compared to year-round reactive vaccination).

We therefore suggest that test-and-removal be prioritized over reactive vaccination, especially if

identification of actively infectious individuals can be improved.

In our simulations, captures that were most aligned with the initial wave of infectious

individuals (i.e., with earlier onset) were more likely to identify actively infectious individuals for

removal. This finding highlights the importance of targeting captures to individuals likely to be

actively infectious. However, determining infection status in cryptic wildlife is difficult, and

consequently supports the increased use of remote tracking technologies that may be able to

(1) identify behavior changes associated with sickness, and (2) detect the onset of an epidemic

more quickly. This conclusion is consistent with similar findings in Channel Island foxes, where

increasing the number and frequency of tracking of sentinel individuals was important for early

identification of epidemics (Sanchez & Hudgens, 2020).

A key component of the success of test-and-removal here is the selective removal we

simulated, which avoids removing immune individuals that contribute to overall herd immunity

(Miguel et al., 2020; Potapov et al., 2012). However, we have simplified the field testing process

in our simulations. The common field FeLV diagnostic test identifies antigenemia, which is key

for identifying actively infectious individuals. However, the duration of antigenemia—and even

degree of infectiousness—in regressively infected individuals is unclear in panthers. We may

therefore overestimate the effect of removing regressive individuals, but given their reduced

infectiousness in our simulations, we still expect test-and-removal to be a key strategy for

mitigating FeLV impacts in panthers.

Notably, reactive vaccination, in concert with proactive vaccination, is also a viable

alternative strategy to test-and-removal. Our simulations showed reactive vaccination to have

the strongest effects for reducing FeLV mortalities when at least 50% of the population was

vaccinated, and with a ratio of about 1.5 vaccines per vaccinated individual (i.e., 50% of

vaccinated individuals received a booster). We therefore suggest that managers should

prioritize boosting at least half of vaccinated individuals in a reactive vaccination response

scenario. We did not see a strong effect of attempting a vaccine barrier, in contrast to Sanchez

et al (2020), where a simulated vaccine barrier could effectively halt spread of a pathogen in

Channel Island foxes. This difference is likely due to the differences in home range size and

movement capacity between the two species, with foxes ranging less widely than panthers. In

reality—and as part of an adaptive management response—if an empirical outbreak of FeLV in

panthers exhibited a stronger spatial signal than was featured in our simulations, spatially

targeted reactive vaccination may yet be a worthwhile intervention strategy.

Importantly—particularly if levels of population protection from proactive vaccination are

unknown—both reactive vaccination and test-and-removal strategies mitigated the negative

effects of low levels of population protection seen with inadequate proactive vaccination. It is

therefore vital for managers to incorporate these reactive strategies in the event of future FeLV

outbreaks.

Limitations and future directions

In this study, we considered the effects of partial vaccine immunity, but we made the

simplifying assumption of no waning vaccine or infection-induced immunity over time. However,

our findings with regard to imperfect efficacy are representative of the likely consequences of

waning immunity in that the loss of immunity supplies new susceptibles to the population.

Should immunity not outlast the course of an FeLV outbreak, this process could prolong

outbreaks and result in increased mortalities. Future research should therefore examine the

effects of waning vaccine immunity, particularly considering the value of revaccinating

individuals that may be experiencing loss of vaccine protection. Furthermore, an adaptive

management framework should monitor for infection in individuals previously considered

immune, which could represent vaccine failure or fading immunity. Model predictions could then

be updated with these observations and revaccination rates adjusted accordingly.

Our simulation results found that relatively high levels of vaccination were required to

reduce impacts of FeLV in panthers, in contrast to studies in other endangered species (Gilbert

et al., 2020; Haydon et al., 2006). However, here we are investigating a pathogen with a long

duration of infectiousness and lacking the advantages of distinct corridors between panther sub-

populations for reducing required levels of vaccination. It is therefore unsurprising that panthers

would require higher levels of FeLV population vaccination than was found, for example, for

rabies vaccination in Ethiopian wolves (Haydon et al., 2006) or canine distemper virus

vaccination in Amur tigers (Gilbert et al., 2020). However, we also did not assume the presence

of preexisting population immunity prior to proactive vaccination, and some degree of population

immunity likely already exists in panthers, given ongoing exposures (Chiu et al., 2019). This

would reduce necessary vaccination levels in panthers, as would higher vaccine efficacy than

we conservatively assumed here (Vial et al., 2006). Future research could prioritize better

understanding the realized individual immunity after single and boosted FeLV vaccination in

panthers to determine if inoculations are more protective than we conservatively assumed here.

Conclusions

Our simulation results highlight the risks of inadequate proactive vaccination, particularly

with partial vaccine immunity. We recommend prioritizing boosted vaccination in panthers, and

joint use of proactive vaccination and reactive strategies to mitigate the risks of imperfect

vaccination and most effectively reduce the impacts of FeLV in this iconic carnivore. This

research highlights the value of linking modeling and management priorities to identify

unexpected consequences of interventions and determine optimal pathogen management

strategies in free-ranging wildlife.

Authors' Contributions

496   All authors conceived the ideas and designed methodology; MLJG performed simulations,

497   analyzed data, and led the writing of the manuscript. All authors contributed critically to the

498   drafts and gave final approval for publication.

Acknowledgements

This research was supported by the National Science Foundation (DEB-1413925, 1654609, and

2030509). MLJG was supported by the Office of the Director, National Institutes of Health (NIH

T32OD010993), the University of Minnesota Informatics Institute MnDRIVE program, and the

Van Sloun Foundation. The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health. Florida panther data

collected by the Florida Fish and Wildlife Conservation Commission is fully supported by

donations to the Florida Panther Research and Management Trust Fund via the registration of

"Protect the Panther" license plates. We acknowledge the efforts of National Park Service staff

in the collection of Florida panther data utilized in this study.

Data Availability

Full R code for simulations is available on GitHub

(https://github.com/mjones029/FeLV_Management_Simulations) and upon acceptance will be

archived at Zenodo.

Works Cited

Baker, L., Matthiopoulos, J., Müller, T., Freuling, C., & Hampson, K. (2019). Optimizing spatial

and seasonal deployment of vaccination campaigns to eliminate wildlife rabies.

Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences,

374(1776), 20180280.

Barnett, K. M., & Civitello, D. J. (2020). Ecological and Evolutionary Challenges for Wildlife

Vaccination. *Trends in Parasitology*, *36*(12), 970–978.

Beyer, H. L., Hampson, K., Lembo, T., Cleaveland, S., Kaare, M., & Haydon, D. T. (2012). The

implications of metapopulation dynamics on the design of vaccination campaigns. *Vaccine*,

30(6), 1014–1022.

Breed, A. C., Plowright, R. K., Hayman, D. T. S., Knobel, D. L., Molenaar, F. M., Gardner–

Roberts, D., Cleaveland, S., Haydon, D. T., Kock, R. A., Cunningham, A. A., Sainsbury, A.

W., & Delahay, R. J. (2009). Disease Management in Endangered Mammals. In R. J.

Delahay, G. C. Smith, & M. R. Hutchings (Eds.), *Management of Disease in Wild Mammals*

(pp. 215–239). Springer Japan.

Brown, M. A., Cunningham, M. W., Roca, A. L., Troyer, J. L., Johnson, W. E., & O'Brien, S. J.

(2008). Genetic characterization of feline leukemia virus from Florida panthers. *Emerging*

Infectious Diseases, *14*(2), 252–259.

Carnell, R. (2012). lhs: Latin hypercube samples. *R Package Version 0. 10, URL http://CRAN.*

R-Project.org/package= Lhs.

Chiu, E. S., Kraberger, S., Cunningham, M., Cusack, L., Roelke, M., & VandeWoude, S. (2019).

Multiple Introductions of Domestic Cat Feline Leukemia Virus in Endangered Florida

Panthers. *Emerging Infectious Diseases*, *25*(1), 92–101.

Choisy, M., & Rohani, P. (2006). Harvesting can increase severity of wildlife disease epidemics.

Proceedings. Biological Sciences / The Royal Society, *273*(1597), 2025–2034.

Craft, M. E., Beyer, H. L., & Haydon, D. T. (2013). Estimating the probability of a major outbreak

from the timing of early cases: an indeterminate problem? *PloS One*, *8*(3), e57878.

Cunningham, M. W., Brown, M. A., Shindle, D. B., Terrell, S. P., Hayes, K. A., Ferree, B. C.,

McBride, R. T., Blankenship, E. L., Jansen, D., Citino, S. B., Roelke, M. E., Kiltie, R. A.,

Troyer, J. L., & O'Brien, S. J. (2008). Epizootiology and management of feline leukemia

virus in the Florida puma. *Journal of Wildlife Diseases*, *44*(3), 537–552.

Gilbert, M., Sulikhan, N., Uphyrkina, O., Goncharuk, M., Kerley, L., Hernandez Castro, E.,

Reeve, R., Seimon, T., McAloose, D., Seryodkin, I. V., Naidenko, S. V., Davis, C. A., Wilkie,

G. S., Vattipally, S. B., Adamson, W. E., Hinds, C., Thomson, E. C., Willett, B. J., Hosie, M.

J., … Cleaveland, S. (2020). Distemper, extinction, and vaccination of the Amur tiger.

Proceedings of the National Academy of Sciences of the United States of America.

https://doi.org/10.1073/pnas.2000153117

Gilbertson, M. L. J., Fountain-Jones, N. M., Malmberg, J. L., Gagne, R. B., Lee, J. S.,

Kraberger, S., Kechejian, S., Petch, R., Chiu, E., Onorato, D., Cunningham, M. W., Crooks,

K. R., Funk, W. C., Carver, S., VandeWoude, S., VanderWaal, K., & Craft, M. E. (2021).

Transmission of one predicts another: Apathogenic proxies for transmission dynamics of a

fatal virus. *bioRxiv*. https://doi.org/10.1101/2021.01.09.426055

Haydon, D. T., Randall, D. A., Matthews, L., Knobel, D. L., Tallents, L. A., Gravenor, M. B.,

Williams, S. D., Pollinger, J. P., Cleaveland, S., Woolhouse, M. E. J., Sillero-Zubiri, C.,

Marino, J., Macdonald, D. W., & Laurenson, M. K. (2006). Low-coverage vaccination

strategies for the conservation of endangered species. *Nature*, *443*(7112), 692–695.

Joseph, M. B., Mihaljevic, J. R., Arellano, A. L., Kueneman, J. G., Preston, D. L., Cross, P. C., &

Johnson, P. T. J. (2013). Taming wildlife disease: bridging the gap between science and

management. *The Journal of Applied Ecology*, *50*(3), 702–712.

Little, S., Levy, J., Hartmann, K., Hofmann-Lehmann, R., Hosie, M., Olah, G., & Denis, K. S.

(2020). 2020 AAFP Feline Retrovirus Testing and Management Guidelines. *Journal of*

Feline Medicine and Surgery, *22*(1), 5–30.

Lloyd-Smith, J. O., Cross, P. C., Briggs, C. J., Daugherty, M., Getz, W. M., Latto, J., Sanchez,

M. S., Smith, A. B., & Swei, A. (2005). Should we expect population thresholds for wildlife

disease? *Trends in Ecology & Evolution*, *20*(9), 511–519.

López, G., López-Parra, M., Fernández, L., Martínez-Granados, C., Martínez, F., Meli, M. L.,

Gil-Sánchez, J. M., Viqueira, N., Díaz-Portero, M. A., Cadenas, R., Lutz, H., Vargas, A., &

Simón, M. A. (2009). Management measures to control a feline leukemia virus outbreak in

the endangered Iberian lynx. *Animal Conservation*, *12*(3), 173–182.

Marino, S., Hogue, I. B., Ray, C. J., & Kirschner, D. E. (2008). A methodology for performing

global uncertainty and sensitivity analysis in systems biology. *Journal of Theoretical*

Biology, *254*(1), 178–196.

McCallum, H. (2008). Landscape structure, disturbance, and disease dynamics. *Infectious*

Disease Ecology: Effects of Ecosystems on Disease and of Disease on Ecosystems, 100–

122.

McClintock, B. T., Onorato, D. P., & Martin, J. (2015). Endangered Florida panther population

size determined from public reports of motor vehicle collision mortalities. *The Journal of*

Applied Ecology, *52*(4), 893–901.

Miguel, E., Grosbois, V., Caron, A., Pople, D., Roche, B., & Donnelly, C. A. (2020). A systemic

approach to assess the potential and risks of wildlife culling for infectious disease control.

Communications Biology, *3*(1), 353.

Potapov, A., Merrill, E., & Lewis, M. A. (2012). Wildlife disease elimination and density

dependence. *Proceedings. Biological Sciences / The Royal Society*, *279*(1741), 3139–

3145.

R Core Team. (2018). *R: A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Rees, E. E., Pond, B. A., Tinline, R. R., & Bélanger, D. (2013). Modelling the effect of landscape

heterogeneity on the efficacy of vaccination for wildlife infectious disease control. *The*

Journal of Applied Ecology, *50*(4), 881–891.

Robinson Stacie J., Barbieri Michelle M., Murphy Samantha, Baker Jason D., Harting Albert L.,

Craft Meggan E., & Littnan Charles L. (2018). Model recommendations meet management

reality: implementation and evaluation of a network-informed vaccination effort for

endangered Hawaiian monk seals. *Proceedings of the Royal Society B: Biological*

Sciences, *285*(1870), 20171899.

Rushmore, J., Caillaud, D., Hall, R. J., Stumpf, R. M., Meyers, L. A., & Altizer, S. (2014).

Network-based vaccination improves prospects for disease control in wild chimpanzees.

Journal of the Royal Society, Interface / the Royal Society, *11*(97), 20140349.

Sanchez, J. N., & Hudgens, B. R. (2020). Vaccination and monitoring strategies for epidemic

prevention and detection in the Channel Island fox (*Urocyon littoralis*). *PloS One*, *15*(5),

e0232705.

Slate, D., Rupprecht, C. E., Rooney, J. A., Donovan, D., Lein, D. H., & Chipman, R. B. (2005).

Status of oral rabies vaccination in wild carnivores in the United States. *Virus Research*,

111(1), 68–76.

Sparkes, A. H. (1997). Feline leukaemia virus: a review of immunity and vaccination. *The*

Journal of Small Animal Practice, *38*(5), 187–194.

Torres, A. N., Mathiason, C. K., & Hoover, E. A. (2005). Re-examination of feline leukemia virus:

host relationships using real-time PCR. *Virology*, *332*(1), 272–283.

Vial, F., Cleaveland, S., Rasmussen, G., & Haydon, D. T. (2006). Development of vaccination

strategies for the management of rabies in African wild dogs. *Biological Conservation*,

131(2), 180–192.

Woodroffe, R., Donnelly, C. A., & Cox, D. R. (2006). Effects of culling on badger *Meles meles*

spatial organization: implications for the control of bovine tuberculosis. *Journal of Applied*

Ecology, 43(1), 1-10.

Wu, J., Dhingra, R., Gambhir, M., & Remais, J. V. (2013). Sensitivity analysis of infectious

disease models: methods, advances and their application. *Journal of the Royal Society,*

Interface / the Royal Society, *10*(86), 20121018.