Plant-derived benzoxazinoids act as antibiotics and shape bacterial communities

Authors

Niklas Schandry[1,2,*], Katharina Jandrasits[1], Ruben Garrido-Oter[3], Claude Becker[1,2,*]

[1] Gregor Mendel Institute of Molecular Plant Biology, Austrian Academy of Sciences, Vienna

BioCenter (VBC), 1030 Vienna, Austria

[2] Genetics, Faculty of Biology, Ludwig Maximilians University Munich, 82152 Martinsried,

Germany

[3] Department of Plant Microbe Interactions, Max Planck Institute for Plant Breeding Research,

50829 Cologne, Germany

* Correspondence should be addressed to claude.becker@bio.lmu.de and

niklas.schandry@bio.lmu.de

Abstract

Plants synthesize and release specialized metabolites into their environment that can serve as chemical cues for other organisms. Metabolites that are released from the roots are important factors in determining which microorganisms will colonize the root and become part of the plant rhizosphere microbiota. Root exudates are often further converted by soil microorganisms, which can result in the formation of toxic compounds. How individual members of the plant rhizosphere respond to individual compounds and how the differential response of individual microorganisms contributes to the response of a microbial community remains unclear. Here, we investigated the impact of derivatives of one class of plant root exudates, benzoxazinoids, which are released by important crops such as wheat and maize, on a collection of 180 root-associated bacteria. We show that phenoxazine, derived from benzoxazinoids, inhibits the growth of root-associated bacteria *in vitro* in an isolate-specific manner, with sensitive and resistant isolates in most of the studied clades. Synthetic bacterial communities that were assembled from only resistant isolates were more resilient to chemical perturbations than communities composed of only sensitive members. Isolates that were shared between different communities showed stable interactions, independent of the overall community composition. On the other hand, we could attribute differential community development to differences in interactions formed by closely related representatives of the same bacterial genus. Our findings highlight the fact that profiling isolate collections can aid the rational design of synthetic communities. Moreover, our data show that simplified *in vitro* community systems are able to recapitulate observations on the influence of metabolite exudation on the structure of root-associated communities, thus providing an avenue for reductionist explorations of the rhizosphere biology in defined, host-free settings.

Introduction

In the rhizosphere, the space directly surrounding the roots, plants recruit and maintain a diverse community of microorganisms that differs from the surrounding bulk soil (Bulgarelli et al., 2013, 2012; Fitzpatrick et al., 2020; Stringlis et al., 2018). These rhizospheric microbiota, in particular bacteria and fungi, contribute to plant nutrition, growth, and defense (Fitzpatrick et al., 2020).

While it is clear that there are biochemical interactions at play in the establishment and the qualitative and quantitative modulation of the microbiota, the chemical ecology of the rhizosphere is not well understood, in part owing to the biological, chemical, and physical complexity of the soil environment. Recent studies have shown, however, that some plant specialized metabolites contribute to plant species-specific community profiles in the rhizosphere. For example, salicylic acid, camalexin, coumarins, and glucosinolates play an important role in shaping the *Arabidopsis thaliana* rhizosphere (Harbort et al., 2020; Jacoby et al., 2020; Koprivova et al., 2019; Lebeis et al., 2015; Stringlis et al., 2018; Voges et al., 2019). Members of the Poaceae family (grasses), including wheat and maize, produce a particular class of specialized metabolites, the benzoxazinoids (BX) (Frey et al., 1997; Handrick et al., 2016). Chemically, BX and some of their breakdown products chelate trivalent ions and have been suggested to act as phytosiderophores in iron uptake or as protective ion scavengers (L. Hu et al., 2018). BX and their derivatives affect different branches of the tree of life: they act as herbivore repellents when stored in plant leaves, can be hijacked by nematodes to deter predators, or interfere with the growth of nearby plants (Robert et al., 2011; Schulz et al., 2013; Venturelli et al., 2016, 2015). Beyond acting on these larger-sized eukaryotes, BX exert a major influence on root-associated fungal and bacterial microbiota. For example, BX exudation by maize plants into the surrounding soil leads to conditioning of the microbiome, which in turn results in soil-memory effects that affect plants of the following sowing (Hu et al., 2018). Another

63 study showed that BX exudation specifically recruits beneficial pseudomonads to the

64 rhizosphere whereas *Comamonadaceae* are often depleted by BX exudation (Cadot et al.,

65 2020; Cotton et al., 2019; Kudjordjie et al., 2019; Neal et al., 2012; Schandry and Becker, 2020).

66 Conversely, the presence and composition of the microbiota can alter BX chemistry. Post-

67 release breakdown and metabolism of benzoxazinoid compounds by soil microbiota leads to the

68 sequential formation of 1,4-benzoxazin-3-one (BOA) and 2-amino-3-phenoxazinone (APO),

69 among other compounds, in non-sterile soils (Kumar et al., 1993; Macías et al., 2005, 2004).

70 APO in return exhibits antibiotic activity similar to other phenoxazines and limits plant root

71 growth at concentrations corresponding to those detected in soil (Anzai et al., 1960; Cheng et

72 al., 2016; Imai et al., 1990; Venturelli et al., 2016, 2015).

73 To date, the selectivity and antimicrobial activity of BX and their non-benzoxazinoid derivatives

74 such as APO on soil-dwelling and root-associated bacteria has not been systematically

75 addressed, and it remains elusive how bacterial communities respond to these compounds.

76 Given the long-term microbiome-conditioning capacity of BX exudation described above (Hu et

77 al., 2018), it is tempting to speculate that some of the plant-growth-inhibitory effects of BX

78 compounds could be mediated by selective effects on the plant-associated microbiota (Belz and

79 Hurle, 2005). However, quantitative analyses of the degree to which microbial isolates differ in

80 their response to root exudates are extremely difficult to perform in soil, because (*i*) plant-

81 associated microbiota are highly diverse, and (*ii*) plant exudates are typically complex mixtures,

82 making it difficult to attribute effects to a particular chemical. In contrast, such studies can be

83 easily carried out *in vitro* using clonal bacterial populations (isolates) and pure compounds at

84 defined concentrations.

85 Collections of culturable bacterial isolates from diverse habitats are an important resource to

86 investigate how diverse microorganisms respond to biotic and abiotic environmental cues or to

87 chemical exposure (e.g. antibiotics, toxins, or pollutants). For example, studies on the effect of

88 therapeutic drugs on an isolate collection derived from the human gut revealed isolate-specific

effects of non-antibiotic therapeutics on bacterial growth (Maier et al., 2018). Such collections

can moreover form the basis for rationally designed synthetic microbial communities (SynComs)

(Herrera Paredes et al., 2018; Vorholt et al., 2017). SynComs have been proposed and used as

reduced complexity systems to study how various microbiomes assemble and develop over

time (De Roy et al., 2014; Finkel et al., 2020; Liu et al., 2019; Voges et al., 2019; Vrancken et

al., 2019). In these studies, the function of individual members has emerged as a key predictor

of community function, whereas taxonomic composition appears to be less crucial. SynComs

have gained traction as simplified models for naturally occurring communities and are an

emerging tool for the targeted modulation of plant health, through holobiont (the organism

formed by host and microbiome) design (Trivedi et al., 2020). Bacterial SynComs in particular

have been adopted in plant sciences to study various aspects of plant biology, including plant

phenotypes under limiting phosphate conditions (Herrera Paredes et al., 2018), assembly of the

phylosphere microbiome (Bodenhausen et al., 2014; Carlström et al., 2019), or assembly cues

of the root microbiome (Bai et al., 2015; Finkel et al., 2020; Voges et al., 2019). SynComs

assembled from plant-associated bacteria have not yet been studied for their ability to assemble

stable communities *in vitro* in the absence of a host, although such a system would significantly

reduce the experimental complexity. Such *in vitro* systems have already been adopted in other

systems, for example for gut microbiota research (Li et al., 2019).

The assembly of a community from individual isolates is a complex process involving basic

metabolic (competitive or mutualistic) and chemical interactions between the different members

that form the community. In addition, environmental perturbations may destabilize a community,

shift its composition, or affect which individuals can establish themselves. The stability of

rhizospheric communities towards environmental perturbations is an important aspect of

rhizosphere biology. With the advent of precision agriculture and highly controlled growth

environments, engineering of the rhizosphere is an emerging topic that encompasses plant

biology, microbiology, microbial ecology, as well as genetic and biochemical biological

disciplines (Dessaux et al., 2016; Ke et al., 2020).

Here, we investigated the impact of BX and their derivatives on bacterial members of the plant

root and rhizosphere. We used the *AtRSphere* collection, derived from the root and rhizosphere

of the model plant species *Arabidopsis thaliana* that includes around 180 culturable isolates

distributed across four bacterial phyla (Bai et al., 2015) for dose-dependent *in vitro* responses to

the BX APO and BOA, which can be found in soil surrounding BX-releasing plants.

A. thaliana is not a BX producer, therefore the ability of an isolate to colonize *A. thaliana* roots

should be independent of its BX tolerance. In line with this assumption, our study revealed a

wide range of growth responses, particularly to APO, among the *A. thaliana* derived *AtRSphere*

collection. This occurred both within and across genera, ranging from complete insensitivity to

strongly impacted growth. Information on the individual isolates' response guided our design of

SynComs to investigate how BX and BX derivatives shape community assembly *in vitro.* We

made use of the isolate-level diversity we observed in the collection with regards to BX

exposure, to explore their potential to form communities differing in resilience to BX-mediated

community perturbations. We found that the ability of an isolate to establish as a community

member is often uncorrelated to growth in pure culture, and that sensitivity of an isolate to BX

may change unpredictably when embedded in a community. Nevertheless, we also found that

SynComs underwent compositional changes *in vitro* that reflected those previously observed in

soil. Since the included isolates partially overlapped between generated SynComs, we

conducted a comparative analysis to understand how chemical treatment and identity of

member isolates shape the composition of *in vitro* communities.

Results

BX derivatives selectively inhibit the growth of root associated bacteria *in vitro*

To gain an overview of how benzoxazinoids and their derivatives affect individual plant-associated bacteria, we screened the *AtRSphere* collection comprising 180 isolates of *A. thaliana* rhizospheric bacteria (Bai et al., 2015) (Figure 1). We monitored the growth of each isolate over 70 h in half-strength TSB medium after supplementation with either BOA (Figure 1A, inner heatmap) or APO (Figure 1A, outer heatmap), two plant-BX-derived compounds commonly found in soil, and under control conditions (equivalent concentration of solvent only). By determining cell density at regular intervals and comparing the area under the curve for the control and BX treatments, we assessed each isolate's response to different compound concentrations (Figure 1A).

The growth of most isolates was not inhibited by BOA, even at high concentrations of 100 µM. Some isolates even seemed to benefit (darker color in Figure 1) from the presence of BOA compared with control conditions; this was most consistent for isolates within the *Rhizobium* and *Paenibacillus* genera (Figure 1B, C). However, such growth promotion was rare overall. This might be attributable to the use of a nutrient-rich medium in this study, which would make BOA superfluous as a potential energy source.

In stark contrast to the overall weak impact of BOA, exposure to APO severely inhibited growth of 43% of the isolates (77/180 isolates grew to less than 50% of control levels in 50 µM APO) (Figure 1A). Some isolates were insensitive to low APO concentrations but showed growth

inhibition at concentrations ≥10 µM, while others showed inhibition already at 5 µM APO (Figure

1). Generally, the impact of APO on growth was dose dependent. Overall, sensitivity to APO

appeared to be isolate-specific, as sensitivities varied as widely within as they did across

families and genera. APO-sensitive isolates (defined as showing <50% relative growth at 50 µM

APO) were present in all families covered by the *AtRSphere* collection (Figure 1A), with the

exception of the *Pseudomonadaceae* and *Alcaligenaceae* (Figure 1D, E). Within those clades

that contained APO-sensitive isolates, the response pattern did not follow an obvious

phylogenetic structure (16S rRNA-based phylogeny), indicating that sensitivity to APO is

species- or even isolate-specific within one genus.

Synthetic bacterial communities are structured by BX treatment

Since we were interested in the effect of BX on communities, we designed four different

SynComs based on the responses of the individual isolates: one that fully represented the

isolate collection at the genus level by randomly choosing one isolate per genus (referred to

hereafter as *random*) and three reduced-complexity SynComs (Supplemental Table 1). Of

these, the *tolerant* SynCom only contained isolates (max. one per genus) that had grown to at

least 50% of the control level in 50 µM APO. As can be seen in Figure 1A, several isolates that

were part of *tolerant* were also included in *random*. The *sensitive* SynCom consisted only of

isolates that had grown to less than 50% of control levels in 50 µM APO. Finally, the *mixed*

SynCom contained both APO-sensitive and APO-tolerant isolates (Figure 1A). To understand

the temporal and treatment-specific community dynamics, we grew each SynCom in three

different environments: control (supplemented with DMSO, the solvent used in the other two

treatments), 100 µM BOA, and 50 µM APO. We monitored SynCom composition in 24 h

intervals over 4 d via 16S rRNA sequencing (see methods).

Figure 1. Isolate sensitivity to BX compounds and derivatives. A) Heatmap arranged around a circular phylogenetic tree computed from 16S rRNA sequences. Nodes represent isolates from the collection, node shape indicates taxonomic class, node color indicates taxonomic family. The inner heatmap displays average relative growth when treated with BOA (from inner to outer: 10 μM, 50 μM and 100 μM BOA, rings 1-3). The central heatmap displays relative growth upon APO treatment (inner to outer: 1 μM, 5 μM, 10 μM and 50 μM, rings 4-7). The outer heatmap indicates whether or not the isolate was selected as a member of one of the 4 synthetic communities (see main text for details).
B-E) Sub-tree of isolates from the *Rhizobiaceae* family **(B)**, *Bacillales* order **(C)**, *Pseudomonadaceae* family **(D)**, and *Alcaligenaceae* family **(E)**; the heatmap displays relative growth in BOA (Orange, from top to bottom: 10 μM, 50 μM, 100 μM) and APO (Blue, from top to bottom: 1 μM, 5 μM, 10 μM and 50 μM). Node color represents taxonomic assignment at the genus level (legend below the heatmap).

As all bacteria analyzed here had originally been isolated from complex plant root microbiomes,

we first assessed to which degree they could reassemble into communities *in vitro* in the

absence of a host. Constrained correspondence analysis (CCA) of Jaccard distances calculated

from relative abundances revealed that SynComs assembled reproducibly (Figure 2A-D).

We next assessed if communities containing only APO-sensitive isolates would be more

strongly restructured by the addition of APO than one assembled only from APO-tolerant

isolates. Generally, we observed treatment-specific changes in all SynComs. Overall, BOA-

treated samples were similar to the control in all SynComs, with the exception of the *sensitive*

one. APO-treated samples clustered separately from both control and BOA (Figure 2 A-D). The

separation between APO and the remaining samples in the ordination was most obvious for the

mixed and *random* and less striking for the *tolerant* SynCom; it decreased over time in the

sensitive one (Figure 2 A-D).

We performed permutational multivariate analysis of variance (PERMANOVA) to see if we could

corroborate these observations. Treatment explained a significant proportion of variance in all

SynComs: 40% in *random*, 22% in *tolerant*, 75% in *mixed*, and 41% in *sensitive*. This means

that the proportion of between-sample variance explained by BX treatments was lowest in the

community assembled from APO-tolerant isolates. Time point explained a significant proportion

of variance in *random* and *tolerant* (31% and 33%, respectively) but not in *mixed* or *sensitive*

(Supplemental Table 2). We then performed another PERMANOVA analysis on each SynCom,

subsetted to a binary comparison between control and either APO or BOA treatment, to

estimate the amount of variance explained by the individual treatments (Supplemental Table 3).

This confirmed that most of the variance observed between treatments in each SynCom was

driven by APO (Figure 2A-D). The amount of variance explained by APO treatment was lowest

in the *tolerant* SynCom (22%), while APO treatment explained approx. 40% in *random* and

sensitive and 70% in *mixed* (Supplemental Table 3). In other words, the SynCom composed of

only the most APO-tolerant individual isolates showed the highest resilience to APO.

Figure 2. Community analysis. Constrained correspondence analysis (CCA, constrained by 'Treatment' and 'Time point') of the Jaccard distances computed on relative genus abundances for all samples. **A-D)** CCA per SynCom (**A**: random; **B**: tolerant; **C**: mixed; **D**: sensitive); faceted by time point (columns) and colored by treatment. **E)** Relationship between change in bacterial growth in the presence of BX compounds in pure culture (x-axis) plotted against the change in relative abundance when embedded in a community (y-axis) for APO (blue) and BOA (orange). Each dot shows the average change in relative abundance for one isolate across all time points.

Having confirmed that the different isolate mixtures reliably assembled into reproducible

communities, we asked if within-sample species diversity (alpha-diversity) was affected by time

or treatment by applying analysis of variance (ANOVA). In the *random*, *tolerant*, and *mixed*

SynComs, APO (but not BOA) treatment significantly reduced the total number of taxa at one or

more time points, while in the *sensitive* SynCom, the taxon number remained constant across

treatments and time points (Supplemental Figure 1, Supplemental Tables S4 and S5). However,

while the total number of taxa remained the same, different taxa were driven to extinction (ratio

of 0 on the relative abundance scale axis, Figure 2E) by the individual treatments, while the

presence of other taxa was specific for one of the compounds (appearing at the very top of each

panel in Figure 2E). Time alone, however, did not significantly influence the species number in

any of the SynComs. The Shannon measure of alpha diversity, which in addition to the number

of species also incorporates the evenness of species abundances, was significantly influenced

by both time and treatment (Supplemental Figure 2, Supplemental Table 5). While APO

treatment significantly increased Shannon diversity of the *mixed* and *tolerant* communities, it

decreased Shannon diversity of the *sensitive* one. In contrast, BOA significantly increased

Shannon diversity of the *sensitive* SynCom after 48 h (Supplemental Table 6).

Overall, there was no clear relationship between the growth of an isolate *in vitro* and the ability

of an isolate to establish itself as a SynCom member. In fact, some isolates that barely grew

when cultured individually reached relative abundances of >90% when grown in a community

(Figure 2E, *mixed* and *sensitive*, top left corner of the plot). The overall magnitude of magnitude

of change, excluding extinction in treatment or absence of control, in relative isolate abundance

was lowest in the *tolerant* SynCom (Figure 2E). The relative abundance ratios in the *tolerant*

community ranged from 0.03 to 2.8, while the lower bound of this range was lower (*random*:

0.0001; *mixed*: 0.005; *sensitive*: 0.008) and the upper bound was higher (*random*: 5.9; *mixed*:

23.2; *sensitive*: 13.6) in all other SynComs. This result indicates that isolates that were found in

all conditions in the *tolerant* SynCom were generally less affected in their relative abundance by

APO treatment, than isolates that were included in the other SynComs (Figure 2E).

Response of community-embedded isolates to BX treatment

Since previous studies reported a substantial impact of plant-derived BX on soil microbiota (Hu

et al., 2018; Cotton et al., 2019; Kudjordjie et al., 2019; Cadot et al., 2020), we were interested

in assessing if the reported effects could be recapitulated using reduced complexity

communities *in vitro*. To understand which taxa drove the observed differences between

treatments (Figure 2), and if similar changes occurred upon APO treatment in the different

communities, we estimated the fold-change in taxon abundance per SynCom and time point

between APO- and BOA-treated samples as well as between APO and control, and between

BOA and control (Figure 3). In this first approach, we focused on those taxa that showed a

significant fold-change between two treatments for at least one time point.

Overall, we observed APO-dependent shifts in relative abundance of different genera in all

SynComs, the comparison of relative abundances between BOA and control conditions only

yielded significant differences in the *sensitive* community.

Some genera were consistently reduced in APO-treated communities. These were *Bacillus* in

random (Figure 3A)*, tolerant* (Figure 3B) and *sensitive* (Figure 3D), *Flavobacterium* in *random*

(Figure 3A) and *mixed* (Figure 3C) and *Acidovorax* in every SynCom, irrespective of the

particular isolate used (Fig 3A-D). Other genera thrived in APO-treated samples: *Lysobacter (*in

mixed and *sensitive*) and *Hoeflea* (identical isolate Root268 in *random* and *sensitive*)*. Only the

genus *Microbacterium* displayed opposing behavior between SynComs: abundance was

increased when treated with APO in *mixed* and *sensitive*, while abundance in the *tolerant*

community was significant reduced in APO-treated samples at 48 h. While *Microbacterium* was

significantly reduced in *tolerant* after 48 h (see confidence intervals in Fig 3B), it is possible that

this is attributable to measurement error, considering that this isolate was only lowly abundant in

this community.

Figure 3: Treatment-specific and temporal isolate abundances in SynComs. Fold-changes between treatments and relative abundances of isolates that were significantly different for at least one time point, estimated by DESeq2. Plots are divided by SynCom: *random* **(A)**, *tolerant* **(B)**, *mixed* **(C)**, and *sensitive* **(D)**. In fold-change plots (top) only significant changes are shown for each time point. Line graphs (below) show relative abundance for all time points for isolates with significant fold-changes. Dots mark the mean across replicates, error bars indicate the 95% confidence interval.

Interactions between isolates are modulated by community members and chemical treatment

In the next approach, we wanted to move beyond comparing only treatment-specific effects and investigate whether interactions between community members were influenced by community context and chemical treatment. The assignment of isolates to the *random* community had been done blindly. However, several of the isolates included in the *random* community were also part of *tolerant* (Figure 1A). We therefore reasoned that differences in the relative abundance of these isolates between the two SynComs likely resulted from interactions with other, non-shared SynCom members. To test this hypothesis, we calculated Pearson's correlation between sequence counts assigned to isolates in each SynCom at 24 h and 96 h. We then constructed correlation networks per treatment and time point, including all correlations with an absolute r > 0.5. The networks for BOA and APO at the 24 h and 96 h time points are shown in Figure 4B, the remaining networks in Supplemental Figure 3.

The *random* and *tolerant* communities became more similar over time when treated with APO, overlapping in 5 genera (out of a total 8) included in the network, confirming the observation in Figure 2A. *Hoeflea* and *Rhodanobacter* were detectable only in the *random* SynCom, *Rhizobium* only in the *tolerant* SynCom. Some of the shared isolates were not detectable in APO but only in BOA and control conditions: *Massilia* Root133 was detected in the early time point, *Acidovorax* Root217 established itself at the late time point (Figure 4B).

When looking at the interactions between isolates, we observed that after treatment with APO, *Bordetella* Root565, *Noviherbaspirillum* Root189, and *Acinetobacter* Root1280 displayed strong, positive correlations after 96 h in both *tolerant* and *random* SynComs. This suggested that this group of isolates engaged in similar interactions, irrespective of the other isolates in the community. However, the relative abundance *Acinetobacter* Root1280 appeared to be more affected by APO treatment in the *tolerant* than in the *random* community (Figure 4A). Based on

310 the correlation networks, we propose that this differential behavior is related to the presence of

311 different *Pseudomonas isolate*s in those SynComs. *Pseudomonas* Root9 negatively correlated

312 with *Acinetobacter* Root1280 in the APO- but not in the BOA-treated *tolerant* SynCom at 96 h.

313 In fact, in the BOA-treated samples, this correlation was positive. In the APO-treated *random*

314 SynCom, *Pseudomonas* Root329 was positively correlated with *Acinetobacter* Root1280 at 96h.

Figure 4: Differential isolate interactions in the *random* and *tolerant* SynComs are driven by isolate identity and chemical treatment. A) Relative abundance of isolates present in both SynComs, over time and by treatment. **B)** Pearson's correlations of isolate abundances in the *tolerant* and *random* SynComs. Each node represents one isolate; node size reflects relative abundance. Color of the node label indicates if an isolate was a member of both (white) or only one (grey) of the SynComs. Each edge indicates a correlation where |r| > 0.5 between two isolates; edge color indicates the correlation value, negative correlations have a blue hue, while positive correlations are orange to yellow.

323 Discussion

324 Diversity of BX sensitivity in culturable rhizosphere bacteria

In our screen of the *AtRSphere* collection for sensitivity to BX derivatives (Figure 1), we

observed broad diversity of growth responses to BX exposure. While some clades were non-

responsive to BOA and APO, we observed diverse responses in most clades. APO exerted a

strong negative influence on the growth of some isolates. These findings are in line with

previous results showing the antibiotic activity of APO on anaerobic microorganisms (Atwal et

al., 1992). In our study, BOA influenced bacterial growth only marginally compared to APO,

similar to what has been described for their effects on fungi and plants. In general, this

corroborates the previously suggested relevance of APO, which is more stable than BOA in soil,

in BX-mediated plant-environment interactions (Kumar et al., 1993; Schandry and Becker, 2020;

Schulz et al., 2013; Schütz et al., 2019; Venturelli et al., 2015; Voloshchuk et al., 2020).

However, our data showed that bacterial response to APO was largely isolate-specific and not

dictated by phylogenetic relationship (Figure 1A). This indicates that even within genera, natural

genetic diversity can result in substantial differences in BX sensitivity. Indeed, previous studies

reported functional overlap related to responses to plant-derived compounds between isolates

from different clades, while not all isolates of the same clade shared these functionalities (Bai et

al., 2015; Harbort et al., 2020). The broad phylogenetic spectrum of the *AtRSphere* collection

prevented the identification of genetic components underlying bacterial response to BOA and

APO. However, the strongly differential responses we observed among closely related isolates

warrant a more comprehensive analysis of the genetic basis of the BX response in the

respective clades in future experiments.

One major advantage of the *AtRSphere* collection is that it was isolated from a non-BX-

producing host plant. The soil from which this collection was originally derived (Cologne soil)

was collected from an agricultural site, and in absence of a complete record of the field's history,

we cannot exclude that there might have been an input of BX in the past. Our findings on the

standing diversity of BX tolerance further imply that this diversity is maintained in plant-

associated bacteria, also in the absence of a BX-releasing host. Strain specific tolerance or

sensitivity to plant specialized metabolites may thus contribute to the colonization profiles

observed for different plant species, where the composition is similar at higher taxonomic ranks,

but some isolates show host preference (Bulgarelli et al., 2013; Schneijderberg et al., 2020)

354 Evaluation of bottom-up design strategy

Others have previously proposed that synthetic communities should be built with a focus on

functional features instead of on phylogenetic coverage (Lemanceau et al., 2017; Vorholt et al.,

2017). We employed a simple, function-guided strategy, compiling SynComs by binning

individual isolates based on a specific phenotype and contrasting these binned assemblies to a

randomly assembled SynCom. The decisive phenotype measured at the level of individual

isolates and used for binning them into SynComs was growth upon exposure to APO. The

phenotype measured at the community level, variance explained by treatment, was lowest

(22%) in the SynCom assembled solely from APO-tolerant isolates, indicating that resistance of

SynCom members leads to overall community resilience. While this approach worked well to

create an overall resilient community, the amount of variance explained by treatment was

similar in *random* and *sensitive* (~40%) and highest in the *mixed* SynCom (75%), indicating that

mixing tolerant and sensitive isolates does not necessarily result in a community with

intermediate resilience.

Stability of communities *in vitro*

Bacterial communities that assemble *in vitro* are less complex, with respect to the genera they

comprise, than those found on and around plant roots yet still contain a considerable number of

different isolates in coexistence. One might expect that in an *in vitro* system containing a

mixture of isolates, one isolate outgrows the others and reaches very high abundance. This

would mean that there are no interactions between the individual isolates and that growth rate is

a strong predictor of final abundance. We observed dominance of an isolate in the *sensitive*

community, where *Lysobacter* reached >90% relative abundance. However, upon the addition

of BOA we observed that *Flavobacterium* was able to establish itself at >10% relative

abundance (corresponding to an eight-fold increase over control). In the *mixed* community,

Bacillus reached >90% abundance in the BOA and control conditions; however, in the APO

treatment, *Bacillus* was strongly reduced in relative abundance, while *Hoeflea* and *Lysobacter*

proliferated. In all other SynComs, several isolates reproducibly formed a community, with

relative abundances staying within a narrow range across replicates but varying by orders of

magnitude between genera.

The growth of an isolate by itself was only poorly correlated with its ability to establish itself as a

SynCom member or with its relative abundance in a community (Figure 2E). In fact, several

isolates that grew poorly on their own reached high relative abundance when grown in

communities, possibly indicating facilitation by other members. However, it should be noted that

we confirmed a maximum of 15 isolates per SynCom by 16S rRNA sequencing.

BX dependent community alterations *in vitro* mimic those described in soil

Our work complements recent research efforts to understand the effect of benzoxazinoids on plant-associated microbiota in soil (Cadot et al., 2020; Cotton et al., 2019; Hu et al., 2018; Kudjordjie et al., 2019; Neal et al., 2012). Ultimately, for *in vitro* SynCom systems to be a useful model for plant microbiota research, they should behave similarly to microbiota in soil. BX constitute a prime system to assess this, as the influence of BX release on the root and rhizosphere microbiome has been investigated via mutant studies in different soils (Cadot et al., 2020; Cotton et al., 2019; Hu et al., 2018; Kudjordjie et al., 2019). Among the bacterial taxa that were found in different studies, *Flavobacteriales* were observed as depleted in multiple studies. Hu et al (Hu et al., 2018) found *Flavobacterium* depleted on the roots of BX-exuding plants and the surrounding soils when comparing wildtype and *bx1* mutant plants. Extending these observations, Cadot et al (Cadot et al., 2020) reported that *Flavobacteriaceae* were consistently depleted on roots of BX-producing plants, independent of soil type and location. Similarly, members of the *Flavobacteriales* family were more abundant on the roots of maize BX-synthesis mutants than on wildtype in another, independent study (Cotton et al., 2019). In all of the afore-mentioned studies, *Comamonadaceae* were also reported to be depleted on roots of BX-releasing plants when compared to BX-deficient mutants (Cadot et al., 2020; Cotton et al., 2019; Hu et al., 2018; Kudjordjie et al., 2019).

Our data show that the growth of *Flavobacterium* isolates was only slightly reduced across APO levels, with some isolates displaying mild sensitivity to BOA. However, *Flavobacterium* was significantly depleted from the *random* and *tolerant* SynComs once the community was exposed to APO. *Flavobacterium* was not found in the other two communities. Similarly, *Acidovorax* (*Comamonadaceae*) was found to be reduced in the APO treatment in all SynComs. Thus, even

though *in vitro* cultivation strongly differs from soils in physical and chemical parameters and in

overall complexity, our study suggests that plant-exudate-mediated effects on soil communities

can be recapitulated *in vitro*, highlighting the applicability of such reduced-complexity systems

for future studies. In addition, such simplified systems may aid in the identification of microbial

isolates that act as community keystones, or hubs, under specific chemical challenges in a short

experimental time frame (Carlström et al., 2019). These data can then be used as a basis for

the design of host-associated communities that are able to remain stable even in the presence

of toxic metabolites, which may increase plant productivity in mixed-cultivation settings.

Chemical stability of isolate interactions

Partial overlap in the communities that arose from the *random* and *tolerant* SynCom mixtures

allowed us to investigate to which degree the identity of community members influences the

interactions between isolates. The *random* and *tolerant* SynComs shared five isolates, three of

which were detectable in all treatments and time points. After 96 h, the *random* and *tolerant*

communities overlapped in 5 genera, including 3 identical isolates and differing representatives

of *Pseudomonas* and *Bacillus*. The identical isolates *Bordetella*, *Acinetobacter*, and

Noviherbaspirillum positively correlated in both SynComs, indicating that these isolates may

form a rather stable network *in vitro*. However, the influence of *Pseudomonas* on this network

was negative in the *tolerant* SynCom, while the *Pseudomonas* isolate in the *random* SynCom

was positively correlated with the network formed by *Acinetobacter*, *Bordetella*, and

Noviherbasprillium. This indicates that the exchange of a single isolate for a closely related

isolate from the same genus can have a broad influence on the overall community composition.

This supports the view that some community members can facilitate or compete with a larger

group of interacting members, i.e., acting as keystones (Banerjee et al., 2018), although this

ability seems to be variable within a clade. We speculate that understanding the connections

between environmental factors and the genetics underlying the ability of an isolate to function as

a key-stone would open up genetic engineering of community stability, for example by stacking

"key-stone genes" into one isolate. Such an isolate then could be used to stabilize communities

used as inoculants in agriculture.

440 Methods

441 Isolate profiling

Individual isolates were pre-cultured in ½ strength TSB medium in 96-Well 2 ml deep-well plates

(Semadeni) covered with a Breathe-Easy (Diversified Biotech) membrane until stationary phase

for 6 d at 28°C and 180 rpm. Cultures were then resuspended by pipetting up and down 10

times and inoculated into 96-well flat-bottom, transparent cell culture plates (Nunc), containing

either solvent control (DMSO), 1 µM APO, 5 µM APO, 10 µM APO, 50 µM APO, 10 µM BOA, 50

µM BOA or 100 µM BOA in ½-strength TSB. Different media were prepared by mixing a stock

containing the highest concentration of the respective chemical, with medium containing the

solvent at the same concentration (DMSO; 1:20.000). Pipetting was carried out using an Agilent

Bravo pipetting robot equipped with a 96-tip head. Plates were stacked without lids, with one

empty plate at the top and the bottom of the stack and placed into a BioTek plate stacker

connected to a BioTek Synergy2 plate reader. Plates were shaken for 120 s before reading, and

absorption (600 nm) was measured in each well. Plates were continuously shaken, measured,

and stacked over the course of at least 48 h per replicate. For each replicate, the relative growth

per isolate and condition was calculated by dividing the area under the growth curve (x = time in

minutes; y = increase in OD600) in the respective condition by the area under the growth curve

of the control condition. The experiment was carried out in eight replicates per condition.

Synthetic bacterial communities

Design

To investigate to which degree the overall resilience of a bacterial community to chemical changes is determined by the properties of its individual members, we followed two separate design strategies. In the first strategy, we aimed at creating a diverse community spanning all genera of the isolate collection. The *random* community was created by picking one random isolate per genus.

In the second approach, we built communities based on the growth profiles of the individual isolates. These communities span a limited number of genera, but the member growth profile is more strictly defined than in the "full" community. One of the reduced representation communities was created from isolates that grew to 75% to 100% of control conditions in 50µM APO (*tolerant*). The other reduced community was assembled from isolates that grew to 0 to 25% of control in 50µM APO (*sensitive*). If there was more than one isolate per genus that fulfilled the condition, we selected one at random. In addition, we compiled one limited representation SynCom containing a mix of sensitive and tolerant isolates (*mixed*).

Cultivation and sequencing

Individual isolates were pre-cultured in ½-strength TSB medium in 96-Well 2ml deep-well plates covered with a Breathe-Easy membrane (Diversified Biotech) with shaking at 28°C for 6 days until stationary phase, as described previously (Bai et al., 2015). Individual cultures were resuspended by pipetting up and down 10 times, and then 100 µL of each culture were mixed with 10 mL ½-strength TSB medium containing solvent at the same concentration as the treatment cultures (1:20.000 DMSO). 100 µL of the mixed culture were used to inoculate 20 mL cultures containing either a solvent control (DMSO), 100 µM BOA, or 50 µM APO in 100 mL

Erlenmayer flasks. Flasks were cultured at 28°C, 180 rpm; 1 ml culture was sampled every 24

h. Sampled cultures were centrifuged at 12.000 rpm, the supernatant was discarded and the

pellet was stored at -70 °C until further use. DNA was extracted using the DNA PowerSoil kit

(Qiagen) following the standard protocol. 16S amplicons were generated according to the

Illumina 16S protocol, indexed using Nextera XT dual indexing (Illumina), and pooled per time

point for sequencing on a MiSeq (Paired end 300bp, Illumina).

487 Data analysis

RStudio with R version 4.0.2 was used for data analysis. Reads processed and ASVs were

inferred using the dada2 pipeline (Ben J. Callahan et al., 2016). Taxonomic assignment of ASVs

was done using SynCom specific 16SrRNA databases and agglomerated at the genus level.

ASVs that could not be assigned were discarded. Scripts used for data analysis are available at

https://github.com/nschan/BxSyncomMs. The following R packages were used with R version

4.0.2:

Tidyverse (Wickham et al., 2019), Broom (Robinson et al., 2019), dada2 (Benjamin J. Callahan

et al., 2016), DECIPHER (Wright, 2016), DESeq2 (Love et al., 2014), emmeans (Lenth et al.,

2019), ggforce (Pedersen, 2020a), ggthemes (Arnold, 2019), ggtree (Yu, 2020; Yu et al., 2018,

2017), gt (Iannone et al., 2019), igraph (Csardi and Nepusz, 2006), MESS (Ekstrøm, 2016),

multcomp (Hothorn et al., 2008), patchwork (Pedersen, 2020b), phangorn (Schliep et al., 2017;

Schliep, 2011), phyloseq (McMurdie and Holmes, 2013), vegan (Oksanen et al., 2019),

wesanderson (Ram and Wickham, 2018) in combination with some custom functions.

501 Data availability

502 Sequencing reads have been deposited to the European Nucleotide Archive (ENA;

503 www.ebi.ac.uk) under the accession no. PRJEB42439.

Acknowledgements

The authors gratefully acknowledge support from the Vienna BioCenter Core Facility (VBCF)

Molecular Biology Services, especially Dr. Robert Heinen, the VBCF Media Kitchen, the VBCF

Sequencing facility and the VBCF High Performance Computing facility. The authors also

gratefully acknowledge the Leibniz Supercomputing Centre (LRZ) for funding this project by

providing computing time on its Linux-Cluster and Rstudio Server.

The authors thank Eva Knoch and Patrick Hüther for constructive feedback on the manuscript,

and Iacopo Gentile and Karina Weiser-Lobao for conducting preliminary experiments and for

technical support. The authors also thank Stijn Staepen for help with the *AtRSphere* collection

and Paul Schulze-Lefert for helpful discussion.

This work was funded by the Austrian Academy of Sciences (ÖAW) and the European

Research Council (ERC) under the European Union's Horizon 2020 research and innovation

programme (Grant agreement No. 716823, "FEAR-SAP").

Supplemental Material

Supplemental figures

Supplemental Figure 1. Observed alpha diversity

Supplemental Figure 2. Shannon diversity indices of the synthetic communities

Supplemental Figure 3. Correlation networks

Supplemental Tables

Supplemental Table 1. SynCom isolates

Supplemental Table 2. PERMANOVA

Supplemental Table 3. ANOVA: observed taxa

Supplemental Table 4. Observed diversity means and pairwise comparisons

Supplemental Table 5. ANOVA: Shannon Diversity

Supplemental Table 6. Shannon diversity means and pairwise comparisons

Literature

Anzai K, Isono K, Okuma K, Suzuki S. 1960. The new antibiotics, questiomycins A and B. *J Antibiot* **13**:125–132.

Arnold JB. 2019. ggthemes: Extra Themes, Scales and Geoms for "ggplot2."

Atwal AS, Teather RM, Liss SN, Collins FW. 1992. Antimicrobial activity of 2-aminophenoxazin-3-one under anaerobic conditions. *Can J Microbiol* **38**:1084–1088.

Bai Y, Müller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M, Dombrowski N, Münch PC, Spaepen S, Remus-Emsermann M, Hüttel B, McHardy AC, Vorholt JA, Schulze-Lefert P. 2015. Functional overlap of the Arabidopsis leaf and root microbiota. *Nature* 1–19.

Banerjee S, Schlaeppi K, van der Heijden MGA. 2018. Keystone taxa as drivers of microbiome structure and functioning. *Nat Rev Microbiol* **16**:567–576.

Belz RG, Hurle K. 2005. Differential exudation of two benzoxazinoids--one of the determining factors for seedling allelopathy of Triticeae species. *J Agric Food Chem* **53**:250–261.

Bodenhausen N, Bortfeld-Miller M, Ackermann M, Vorholt JA. 2014. A synthetic community approach reveals plant genotypes affecting the phyllosphere microbiota. *PLoS Genet* **10**:e1004283.

Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R, Eickhorst T, Schulze-Lefert P. 2012. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. *Nature* **488**:91–95.

Bulgarelli D, Schlaeppi K, Spaepen S, van Themaat EVL, Schulze-Lefert P. 2013. Structure and Functions of the Bacterial Microbiota of Plants. *Annu Rev Plant Biol* **64**:807–838.

Cadot S, Guan H, Bigalke M, Walser J-C, Jander G, Erb M, van der Heijden M, Schlaeppi K. 2020. Specific and conserved patterns of microbiota-structuring by maize benzoxazinoids in the field. *bioRxiv*. doi:10.1101/2020.05.03.075135

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**:581–583.

Callahan BJ, Sankaran K, Fukuyama JA, McMurdie PJ, Holmes SP. 2016. Bioconductor Workflow for Microbiome Data Analysis: from raw reads to community analyses. *F1000Res* **5**:1492.

Carlström CI, Field CM, Bortfeld-Miller M, Müller B, Sunagawa S, Vorholt JA. 2019. Synthetic microbiota reveal priority effects and keystone strains in the Arabidopsis phyllosphere. *Nat Ecol Evol* **3**:1445–1454.

Cheng C, Othman EM, Fekete A, Krischke M, Stopper H, Edrada-Ebel R, Mueller MJ, Hentschel U, Abdelmohsen UR. 2016. Strepoxazine A, a new cytotoxic phenoxazin from the marine sponge-derived bacterium Streptomyces sp. SBT345. *Tetrahedron Lett* **57**:4196–4199.

Cotton TEA, Pétriacq P, Cameron DD, Meselmani MA, Schwarzenbacher R, Rolfe SA, Ton J. 2019. Metabolic regulation of the maize rhizobiome by benzoxazinoids. *ISME J*. doi:10.1038/s41396-019-0375-2

Csardi G, Nepusz T. 2006. The igraph software package for complex network research. *InterJournal*.

De Roy K, Marzorati M, Van den Abbeele P, Van de Wiele T, Boon N. 2014. Synthetic microbial ecosystems: an exciting tool to understand and apply microbial communities. *Environ Microbiol* **16**:1472–1481.

Dessaux Y, Grandclément C, Faure D. 2016. Engineering the Rhizosphere. *Trends Plant Sci* **21**:266–278.

Ekstrøm C. 2016. MESS: Miscellaneous Esoteric Statistical Scripts.

Finkel OM, Salas-González I, Castrillo G, Conway JM, Law TF, Teixeira PJPL, Wilson ED,
Fitzpatrick CR, Jones CD, Dangl JL. 2020. A single bacterial genus maintains root growth
in a complex microbiome. *Nature*. doi:10.1038/s41586-020-2778-7
Fitzpatrick CR, Salas-González I, Conway JM, Finkel OM, Gilbert S, Russ D, Teixeira PJPL,
Dangl JL. 2020. The Plant Microbiome: From Ecology to Reductionism and Beyond. *Annu
Rev Microbiol*. doi:10.1146/annurev-micro-022620-014327
Frey M, Chomet P, Glawischnig E, Stettner C, Grün S, Winklmair A, Eisenreich W, Bacher A,
Meeley RB, Briggs SP, Simcox K, Gierl A. 1997. Analysis of a chemical plant defense
mechanism in grasses. *Science* **277**:696–699.
Handrick V, Robert CAM, Ahern KR, Zhou S, Machado RAR, Maag D, Glauser G, Fernandez-
Penny FE, Chandran JN, Rodgers-Melnik E, Schneider B, Buckler ES, Boland W,
Gershenzon J, Jander G, Erb M, Köllner TG. 2016. Biosynthesis of 8-O-Methylated
Benzoxazinoid Defense Compounds in Maize. *Plant Cell* **28**:1682–1700.
Harbort CJ, Hashimoto M, Inoue H, Niu Y, Guan R, Rombolà AD, Kopriva S, Mathias J E E,
Sattely ES, Garrido-Oter R, Schulze-Lefert P. 2020. Root-Secreted Coumarins and the
Microbiota Interact to Improve Iron Nutrition in Arabidopsis. *Cell Host Microbe* **0**.
doi:10.1016/j.chom.2020.09.006
Herrera Paredes S, Gao T, Law TF, Finkel OM, Mucyn T, Teixeira PJPL, Salas González I,
Feltcher ME, Powers MJ, Shank EA, Jones CD, Jojic V, Dangl JL, Castrillo G. 2018. Design
of synthetic bacterial communities for predictable plant phenotypes. *PLoS Biol*
16:e2003962.
Hothorn T, Bretz F, Westfall P. 2008. Simultaneous inference in general parametric models.
Biom J **50**:346–363.
Hu L, Mateo P, Ye M, Zhang X, Berset JD, Handrick V, Radisch D, Grabe V, Köllner TG,
Gershenzon J, Robert CAM, Erb M. 2018. Plant iron acquisition strategy exploited by an
insect herbivore. *Science* **361**:694–697.
Hu L, Robert CAM, Cadot S, Zhang X, Ye M, Li B, Manzo D, Chervet N, Steinger T, van der
Heijden MGA, Schlaeppi K, Erb M. 2018. Root exudate metabolites drive plant-soil
feedbacks on growth and defense by shaping the rhizosphere microbiota. *Nat Commun*
9:2738.
Iannone R, Cheng J, Schloerke B. 2019. gt: Easily Create Presentation-Ready Display Tables.
R package version 0.1. 0.
Imai S, Shimazu A, Furihata K, Furihata K, Hayakawa Y, Seto H. 1990. Isolation and structure
of a new phenoxazine antibiotic, exfoliazone, produced by Streptomyces exfoliatus. *J
Antibiot* **43**:1606–1607.
Jacoby RP, Koprivova A, Kopriva S. 2020. Pinpointing secondary metabolites that shape the
composition and function of the plant microbiome. *J Exp Bot*. doi:10.1093/jxb/eraa424
Ke J, Wang B, Yoshikuni Y. 2020. Microbiome Engineering: Synthetic Biology of Plant-
Associated Microbiomes in Sustainable Agriculture. *Trends Biotechnol*.
doi:10.1016/j.tibtech.2020.07.008
Koprivova A, Schuck S, Jacoby RP, Klinkhammer I, Welter B, Leson L, Martyn A, Nauen J,
Grabenhorst N, Mandelkow JF, Zuccaro A, Zeier J, Kopriva S. 2019. Root-specific
camalexin biosynthesis controls the plant growth-promoting effects of multiple bacterial
strains. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1818604116
Kudjordjie EN, Sapkota R, Steffensen SK, Fomsgaard IS, Nicolaisen M. 2019. Maize
synthesized benzoxazinoids affect the host associated microbiome. *Microbiome* **7**:59.
Kumar P, Gagliardo RW, Chilton WS. 1993. Soil transformation of wheat and corn metabolites
mboa and DIM2BOA into aminophenoxazinones. *J Chem Ecol* **19**:2453–2461.
Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, Malfatti S,
Glavina del Rio T, Jones CD, Tringe SG, Dangl JL. 2015. Salicylic acid modulates
colonization of the root microbiome by specific bacterial taxa. *Science* **349**:860–864.

628 Lemanceau P, Blouin M, Muller D, Moënne-Loccoz Y. 2017. Let the Core Microbiota Be
629   Functional. *Trends Plant Sci* **22**:583–595.
630 Lenth R, Singmann H, Love J, Buerkner P, Herve M. 2019. emmeans: Estimated Marginal
631   Means, aka Least-Squares Means (Version 1.3. 4).
632 Li L, Abou-Samra E, Ning Z, Zhang X, Mayne J, Wang J, Cheng K, Walker K, Stintzi A, Figeys
633   D. 2019. An in vitro model maintaining taxon-specific functional activities of the gut
634   microbiome. *Nat Commun* **10**:4146.
635 Liu Y-X, Qin Y, Bai Y. 2019. Reductionist synthetic community approaches in root microbiome
636   research. *Curr Opin Microbiol* **49**:97–102.
637 Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
638   RNA-seq data with DESeq2. *Genome Biol* **15**:550.
639 Macías FA, Oliveros-Bastidas A, Marín D, Castellano D, Simonet AM, Molinillo JMG. 2005.
640   Degradation studies on benzoxazinoids. Soil degradation dynamics of (2R)-2-O-beta-D-
641   glucopyranosyl-4-hydroxy-(2H)- 1,4-benzoxazin-3(4H)-one (DIBOA-Glc) and its
642   degradation products, phytotoxic allelochemicals from Gramineae. *J Agric Food Chem*
643   **53**:554–561.
644 Macías FA, Oliveros-Bastidas A, Marín D, Castellano D, Simonet AM, Molinillo JMG. 2004.
645   Degradation Studies on Benzoxazinoids. Soil Degradation Dynamics of 2,4-Dihydroxy-7-
646   methoxy-(2 H)-1,4-benzoxazin-3(4 H)-one (DIMBOA) and Its Degradation Products,
647   Phytotoxic Allelochemicals from Gramineae. *J Agric Food Chem* **52**:6402–6413.
648 Maier L, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, Brochado AR, Fernandez
649   KC, Dose H, Mori H, Patil KR, Bork P, Typas A. 2018. Extensive impact of non-antibiotic
650   drugs on human gut bacteria. *Nature* **555**:623–628.
651 McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive analysis
652   and graphics of microbiome census data. *PLoS One* **8**:e61217.
653 Neal AL, Ahmad S, Gordon-Weeks R, Ton J. 2012. Benzoxazinoids in root exudates of maize
654   attract Pseudomonas putida to the rhizosphere. *PLoS One* **7**:e35498.
655 Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB,
656   Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2019. vegan: Community
657   Ecology Package.
658 Pedersen TL. 2020a. ggforce: Accelerating "ggplot2."
659 Pedersen TL. 2020b. patchwork: The Composer of Plots.
660 Ram K, Wickham H. 2018. wesanderson: A Wes Anderson Palette Generator.
661 Robert CAM, Veyrat N, Glauser G, Marti G, Doyen GR, Villard N, Gaillard MDP, Köllner TG,
662   Giron D, Body M, Babst BA, Ferrieri RA, Turlings TCJ, Erb M. 2011. A specialist root
663   herbivore exploits defensive metabolites to locate nutritious tissues. *Ecol Lett* **14**:55–64.
664 Robinson D, Hayes A, Couch S. 2019. Broom: Convert statistical objects into tidy tibbles.
665 Schandry N, Becker C. 2020. Allelopathic Plants: Models for Studying Plant–Interkingdom
666   Interactions. *Trends Plant Sci* **25**:176–185.
667 Schliep, Klaus, Potts, J. A, Morrison, A. D, Grimm, W. G. 2017. Intertwining phylogenetic trees
668   and networks. *Methods in Ecology and Evolution*. doi:10.1111/2041-210X.12760
669 Schliep KP. 2011. phangorn: phylogenetic analysis in R. *Bioinformatics*.
670   doi:10.1093/bioinformatics/btq706
671 Schneijderberg M, Cheng X, Franken C, de Hollander M, van Velzen R, Schmitz L, Heinen R,
672   Geurts R, van der Putten WH, Bezemer TM, Bisseling T. 2020. Quantitative comparison
673   between the rhizosphere effect of Arabidopsis thaliana and co-occurring plant species with
674   a longer life history. *ISME J.* doi:10.1038/s41396-020-0695-2
675 Schulz M, Marocco A, Tabaglio V, Macias FA, Molinillo JMG. 2013. Benzoxazinoids in rye
676   allelopathy - from discovery to application in sustainable weed control and organic farming.
677   *J Chem Ecol* **39**:154–174.
678 Schütz V, Bigler L, Girel S, Laschke L, Sicker D, Schulz M. 2019. Conversions of

Benzoxazinoids and Downstream Metabolites by Soil Microorganisms. *Frontiers in Ecology*
and Evolution **7**:238.
Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, Berendsen RL,
Bakker PAHM, Feussner I, Pieterse CMJ. 2018. MYB72-dependent coumarin exudation
shapes root microbiome assembly to promote plant health. *Proc Natl Acad Sci U S A*.
doi:10.1073/pnas.1722335115
Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. 2020. Plant-microbiome interactions: from
community assembly to plant health. *Nat Rev Microbiol*. doi:10.1038/s41579-020-0412-1
Venturelli S, Belz RG, Kamper A, Berger A, von Horn K, Wegner A, Bocker A, Zabulon G,
Langenecker T, Kohlbacher O, Barneche F, Weigel D, Lauer UM, Bitzer M, Becker C.
2015. Plants Release Precursors of Histone Deacetylase Inhibitors to Suppress Growth of
Competitors. *Plant Cell* **27**:3175–3189.
Venturelli S, Petersen S, Langenecker T, Weigel D, Lauer UM, Becker C. 2016. Allelochemicals
of the phenoxazinone class act at physiologically relevant concentrations. *Plant Signal*
Behav **11**:e1176818.
Voges MJEEE, Bai Y, Schulze-Lefert P, Sattely ES. 2019. Plant-derived coumarins shape the
composition of an Arabidopsis synthetic root microbiome. *Proc Natl Acad Sci U S A*
116:12558–12565.
Voloshchuk N, Schütz V, Laschke L, Gryganskyi AP, Schulz M. 2020. The Trichoderma viride F-
00612 consortium tolerates 2-amino-3H-phenoxazin-3-one and degrades nitrated
benzo[d]oxazol-2(3H)-one. *Chemoecology*. doi:10.1007/s00049-020-00300-w
Vorholt JA, Vogel C, Carlström CI, Müller DB. 2017. Establishing Causality: Opportunities of
Synthetic Communities for Plant Microbiome Research. *Cell Host Microbe* **22**:142–155.
Vrancken G, Gregory AC, Huys GRB, Faust K, Raes J. 2019. Synthetic ecology of the human
gut microbiota. *Nat Rev Microbiol* **17**:754–763.
Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A,
Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D,
Seidel D, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome to
the Tidyverse. *JOSS* **4**:1686.
Wright ES. 2016. Using DECIPHER v2. 0 to analyze big biological sequence data in R. *R J* **8**.
Yu G. 2020. Using ggtree to Visualize Data on Tree-Like Structures. *Current Protocols in*
Bioinformatics. doi:10.1002/cpbi.96
Yu G, Lam TT-Y, Zhu H, Guan Y. 2018. Two methods for mapping and visualizing associated
data on phylogeny using ggtree. *Molecular Biology and Evolution*.
doi:10.1093/molbev/msy194
Yu G, Smith D, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods in*
Ecology and Evolution. doi:10.1111/2041-210X.12628