Plant-derived benzoxazinoids act as antibiotics and shape bacterial communities

Authors

Niklas Schandry[1,2,*], Katharina Jandrasits[1], Ruben Garrido-Oter[3], Claude Becker[1,2,*]

[1] Gregor Mendel Institute of Molecular Plant Biology, Austrian Academy of Sciences, Vienna

BioCenter (VBC), 1030 Vienna, Austria

[2] Genetics, Faculty of Biology, Ludwig Maximilians University Munich, 82152 Martinsried,

Germany

[3] Department of Plant Microbe Interactions, Max Planck Institute for Plant Breeding Research,

50829 Cologne, Germany

* Correspondence should be addressed to claude.becker@bio.lmu.de and

niklas.schandry@bio.lmu.de

Abstract

Plants synthesize and release specialized metabolites into their environment that can serve as chemical cues for other organisms. Metabolites that are released from the roots are important factors in determining which microorganisms will colonize the root and become part of the plant rhizosphere. Root exudates can be converted by soil microorganisms, which can result in the formation of toxic compounds. How individual members of the plant rhizosphere respond to individual compounds and how the differential response of individual microorganisms contributes to the response of a microbial community remains an open question. Here, we investigated the impact of derivatives of benzoxazinoids, a class of plant root exudates released by important crops such as wheat and maize, on a collection of 180 root-associated bacteria. Phenoxazine, derived in soil from benzoxazinoids, inhibited the growth of root-associated bacteria *in vitro* in an isolate-specific manner, with sensitive and resistant isolates present in most of the studied clades. Using synthetic communities, we show that community stability is a consequence of the resilience of its individual members, with communities assembled from tolerant isolates being overall more tolerant to benzoxazinoids. However, the performance of an isolate in a community context was not correlated with its individual performance but appeared to be shaped by interactions between isolates. These interactions were independent of the overall community composition and were strain-specific, with interactions between different representatives of the same bacterial genera accounting for differential community composition.

Introduction

In the rhizosphere, the space directly surrounding the roots, plants recruit and maintain a diverse community of microorganisms that differs from the surrounding bulk soil (Bulgarelli et al., 2013, 2012; Fitzpatrick et al., 2020; Stringlis et al., 2018). These rhizospheric microbiota, in particular bacteria and fungi, contribute to plant nutrition, growth, and defense (Fitzpatrick et al., 2020). Recent studies have shown that some plant specialized metabolites contribute to plant species-specific community profiles in the rhizosphere. For example, salicylic acid, camalexin, coumarins, and glucosinolates play an important role in shaping the *Arabidopsis thaliana* rhizosphere (Harbort et al., 2020; Jacoby et al., 2020; Koprivova et al., 2019; Lebeis et al., 2015; Stringlis et al., 2018; Voges et al., 2019).

Members of the Poaceae family, including wheat and maize, produce a particular class of specialized metabolites, the benzoxazinoids (BX) (Frey et al., 1997; Handrick et al., 2016). BX exert a major influence on root-associated fungal and bacterial microbiota. For example, BX exudation by maize plants into the surrounding soil conditions the microbiome, which in turn results in soil-memory effects that affect plants of the following sowing (Hu et al., 2018). Other studies showed that BX exudation specifically recruits beneficial pseudomonads to the rhizosphere, whereas *Comamonadaceae* are often depleted by BX exudation (Cadot et al., 2020; Cotton et al., 2019; Kudjordjie et al., 2019; Neal et al., 2012; Schandry and Becker, 2020). Conversely, the presence and composition of the microbiota can alter BX chemistry. Post-release breakdown and metabolism of benzoxazinoid compounds by soil microbiota leads to the sequential formation of 1,4-benzoxazin-3-one (BOA) and 2-amino-3-phenoxazinone (APO), among other compounds, in non-sterile soils (Kumar et al., 1993; Macías et al., 2005, 2004). APO in return exhibits antibiotic activity similar to other phenoxazines and limits plant root growth at concentrations corresponding to those detected in soil (Anzai et al., 1960; Cheng et al., 2016; Imai et al., 1990; Venturelli et al., 2016, 2015).

Given the long-term microbiome-conditioning capacity of BX exudation (Hu et al., 2018), it is

tempting to speculate that some of the plant-growth-inhibitory effects of BX compounds could be

mediated by selective effects on the plant-associated microbiota (Belz and Hurle, 2005).

However, quantitative analyses of the degree to which microbial isolates differ in their response

to root exudates are extremely difficult to perform in soil, because (*i*) plant-associated microbiota

are highly diverse, and (*ii*) plant exudates are typically complex mixtures, making it difficult to

attribute effects to a particular chemical. In contrast, such studies can be carried out *in vitro*

using clonal bacterial populations (isolates) and pure compounds at defined concentrations.

Collections of culturable bacterial isolates from diverse habitats are an important resource to

investigate how diverse microorganisms respond to chemical exposure (e.g., antibiotics, toxins,

or pollutants). For example, studies on the effect of therapeutic drugs on an isolate collection

derived from the human gut revealed isolate-specific effects of non-antibiotic therapeutics on

bacterial growth (Maier et al., 2018).

Such collections can moreover form the basis for rationally designed synthetic microbial

communities (SynComs) (Herrera Paredes et al., 2018; Vorholt et al., 2017). SynComs have

been proposed and used as reduced complexity systems to study how various microbiomes

assemble and develop over time (De Roy et al., 2014; Finkel et al., 2020; Liu et al., 2019; Voges

et al., 2019; Vrancken et al., 2019). Bacterial SynComs in particular have been adopted in plant

sciences to study various aspects of plant biology, including plant phenotypes under limiting

phosphate conditions (Herrera Paredes et al., 2018), assembly of the phylosphere microbiome

(Bodenhausen et al., 2014; Carlström et al., 2019), or assembly cues of the root microbiome

(Bai et al., 2015; Finkel et al., 2020; Voges et al., 2019). While host-free *in vitro* systems have

already been adopted in other systems, e.g. gut microbiota (Li et al., 2019), SynComs

assembled from plant-associated bacteria have not yet been studied for their ability to assemble

stable communities *in vitro* in the absence of a host, although such a system would significantly

reduce the experimental complexity.

Here, we investigated the impact of BX and their derivatives on bacterial members of the plant

root and rhizosphere by exposing these isolates to varying concentrations of APO and BOA, two

BX detectable in soil surrounding BX-releasing plants. We used the *AtRSphere* collection of

culturable *Arabidopsis thaliana*-derived bacterial isolates, spanning four bacterial phyla (Bai et

al., 2015); *A. thaliana* is not a BX producer and hence the isolates' capacity to colonize its roots

should be independent of their BX tolerance. In line with this assumption, growth responses,

most notably those to APO, of *AtRSphere* isolates varied substantially both within and across

genera, ranging from complete insensitivity to strongly impeded growth.

We designed different SynComs to investigate how isolate-level responses to BX influence *in*

vitro community assembly and resilience to BX-mediated community perturbations. An isolate's

individual growth behaviour did not correlate with its ability to establish as a community member

or with its sensitivity to BX when growing in a community. Moreover, there was no phylogenetic

signal for BX tolerance within most bacterial orders, indicating that BX tolerance is strain- rather

than genus- or family-specific. Finally, through comparative analyses of isolates and genera

shared between differently composed SynComs, we show that BX-mediated restructuring of

bacterial communities is attributable to the selective inhibition of some members that relieves

antagonistic interactions with other community members.

102 Results

103 BX derivatives selectively inhibit the growth of root associated

104 bacteria *in vitro*

To gain an overview of how benzoxazinoids and their derivatives affect individual plant-

associated bacteria, we screened the *AtRSphere* collection comprising 180 isolates of *A.*

thaliana rhizospheric bacteria (Bai et al., 2015) (Figure 1). We monitored the growth of each

isolate over 70 h in half-strength TSB medium after supplementation with either BOA (Figure

1A, inner heatmap) or APO (Figure 1A, outer heatmap), two plant-BX-derived compounds

commonly found in soil, and under control conditions (equivalent concentration of solvent only).

By determining cell density at regular intervals and comparing the area under the curve for the

control and BX treatments, we assessed each isolate's response to different compound

concentrations (Figure 1A).

Because the strains had been isolated from *A. thaliana*, a species that does not release BX, we

hypothesized that the collection should cover a broad diversity in terms of BX tolerance. We

used 50% relative growth compared to control as the threshold to classify isolates as either

sensitive (<50%) or tolerant (>50%).

The growth of most isolates was not inhibited by BOA, even at high concentrations of 100 µM.

Some isolates even seemed to benefit (darker color in Figure 1) from the presence of BOA

compared with control conditions; this was most consistent for isolates within the *Rhizobium* and

Paenibacillus genera (Figure 1A). However, such growth promotion was rare overall; this might

be attributable to the use of a nutrient-rich medium in this study, which would make BOA

superfluous as a potential energy source.

The BX derivative APO was previously shown to have antibiotic effects, so we anticipated a

more pronounced negative and dose-dependent effect of APO on isolate growth (Anzai et al.,

1960). Indeed, some isolates were insensitive to low APO concentrations but showed growth

inhibition at concentrations ≥10 µM, while the growth of others was inhibited already at 5 µM

APO. Generally, the impact of APO on growth was dose dependent (Figure 1A). In stark

contrast to the overall weak impact of BOA, exposure to APO severely inhibited growth of 43%

of isolates at concentrations of 50 µM: 77 out of 179 isolates grew to less than 50% of control

levels in 50 µM APO (Figure 1A). All isolates in the collection that belonged to the families

Alcaligenaceae (β-Proteobacteria) or *Pseudomonaceae* and *Moraxellaceae* (γ-Proteobacteria)

were resistant to APO, while all *Flavobacteriaceae* (Bacteroidia) were APO sensitive. The

remaining twenty families contained both sensitive and resistant isolates (Figure 1A, B).

From the overall phylogeny (Figure 1A), it appeared that there was some phylogenetic

structuring of APO tolerance; within lower taxonomic ranks, however, this signal appeared to

become less pronounced, indicating that APO tolerance might be a strain- rather than a

species- or genus-specific trait in most families and orders (Figure 1B). To test this hypothesis,

we calculated the strength of the correlation between the relative growth at 50 μM and the

phylogenetic distance using Pagel's λ and tested against the null hypothesis $\lambda = 0$ (likelihood

ratio test, Supplemental Table 1; Pagel, 1999, Münkemüller et al., 2012). A value of λ close to 0

indicates no phylogenetic signal; conversely λ values closer to 1 indicate that the trait is

correlated with the phylogenetic structure. For the overall collection, λ was 0.76 ($p < 0.001$). We

next tested the strength of the phylogenetic signal within gram-positive bacteria (classes

Actinomycetia and Bacilli), which returned a weaker but significant λ of 0.46 ($p < 0.05$).

However, within both Actinomycetia and Bacilli, or within families belonging to those classes,

this signal disappeared ($\lambda < 0.1$, $p > 0.05$). Within the phylum Proteobacteria (α-, β- and γ-

proteobacteria), λ was high (0.81, $p < 0.05$). Within phyla, response correlated with phylogenetic

structure only in β- and γ-proteobacteria ($\lambda = 1$, $p < 0.05$) but not in α-proteobacteria ($\lambda < 0.1$, p

> 0.05). With the exception of the *Xanthomonadaceae* ($\lambda = 1$, $p < 0.05$), λ values within

proteobacteria families were not significantly different from zero. Within the *Xanthomonadaceae*,

where we observed a phylogenetic signal for APO response, four out of five sensitive isolates

belonged to the genus *Lysobacter* (Supplemental Table 1, Supplemental Figure 1).

These data corroborated our initial observation that APO tolerance is largely a strain-specific

feature, and that diversity in this trait exists in the AtRSphere collection. While there was a

phylogenetic signal in the dataset at broad phylogenetic and taxonomic levels, this signal often

weakened or disappeared in lower taxonomic ranks (Figure 1A, B, Supplemental Table 1).

Figure 1. Isolate sensitivity to BX compounds and derivatives. A) Heatmap arranged around a
circular unrooted phylogenetic tree based on several marker genes. Nodes represent isolates from the
collection, node shape indicates taxonomic class, node color indicates taxonomic family (genera
Rhizobacter and Methylibium are not assigned to a taxonomic family). The inner heatmap displays
average relative growth upon BOA treatment (from inner to outer: 10 µM, 50 µM and 100 µM BOA, rings
1-3). The central heatmap displays relative growth upon APO treatment (inner to outer: 1 µM, 5 µM, 10
µM and 50 µM, rings 4-7). Relative growth was estimated from at least 6 replicate experiments per
isolate. The outer heatmap indicates whether or not the isolate was selected as a member of one of the
four synthetic communities (see main text for details).
B) Density plots of relative growth at 50 uM APO. Densities are shown per family (same colors as in A),
per class. Dots show the mean relative growth of individual isolates belonging to the respective family.

Synthetic bacterial communities are structured by BX treatment

To study the effect of BX on communities, we designed four different SynComs based on the responses of the individual isolates: one that fully represented the isolate collection at the genus level by randomly choosing one isolate per genus (referred to hereafter as *random*) and three reduced-complexity SynComs (Supplemental Table 2). Of these, the *tolerant* SynCom only contained isolates (max. one per genus) that had grown to at least 50% of the control level in 50 μM APO. As can be seen in Figure 1A, several isolates that were part of *tolerant* were also included in *random*. The *sensitive* SynCom consisted only of isolates that had grown to less than 50% of control levels in 50 μM APO. Finally, the *mixed* SynCom contained both APO-sensitive and APO-tolerant isolates (Figure 1A). To understand the temporal and treatment-specific community dynamics, we grew each SynCom in three different environments: control (supplemented with DMSO, the solvent used in the other two treatments), 100 μM BOA, and 50 μM APO. We monitored SynCom composition in 24 h intervals over 4 d via 16S rRNA sequencing (see methods).

As all bacteria analyzed here had originally been isolated from complex plant root microbiomes, we first assessed to which degree they could reassemble into communities *in vitro* in the absence of a host. Constrained correspondence analysis (CCA) of Jaccard distances calculated from relative abundances revealed that *in vitro* SynComs assembled reproducibly (Figure 2A-D).

We next assessed if communities containing only APO-sensitive isolates would be more strongly restructured by the addition of APO than one assembled only from APO-tolerant isolates. Generally, we observed treatment-specific changes in all SynComs. Overall, BOA-treated samples were similar to the control in all SynComs, with the exception of the *sensitive* one. APO-treated samples clustered separately from both control and BOA (Figure 2 A-D). The separation between APO and the remaining samples in the ordination was most obvious for the

mixed and *random* and less striking for the *tolerant* SynCom; it decreased over time in the

sensitive one (Figure 2 A-D).

We performed permutational multivariate analysis of variance (PERMANOVA) to see if we could

confirm these observations. Treatment explained a significant proportion of variance in all

SynComs: 40% in *random*, 22% in *tolerant*, 75% in *mixed*, and 41% in *sensitive*. This means

that the proportion of between-sample variance explained by BX treatments was lowest in the

community assembled from APO-tolerant isolates. Time point explained a significant proportion

of variance in *random* and *tolerant* (31% and 33%, respectively) but not in *mixed* or *sensitive*

(Supplemental Table 3). We then performed another PERMANOVA analysis on each SynCom,

subset to a binary comparison between control and either APO or BOA treatment, to estimate

the amount of variance explained by the individual treatments (Supplemental Table 4). This

confirmed that most of the variance observed between treatments in each SynCom was driven

by APO (Figure 2A-D). The amount of variance explained by APO treatment was lowest in the

tolerant SynCom (22%), while APO treatment explained approx. 40% in *random* and *sensitive*

and 70% in *mixed* (Supplemental Table 4). In other words, the SynCom composed of only the

most APO-tolerant individual isolates showed the highest resilience to APO treatment.

Having confirmed that the different isolate mixtures reliably assembled into reproducible

communities, we asked if within-sample species diversity (alpha-diversity) was affected by time

or treatment by applying analysis of variance (ANOVA). In the *random*, *tolerant*, and *mixed*

SynComs, APO (but not BOA) treatment significantly reduced the total number of taxa at one or

more time points, while in the *sensitive* SynCom, the taxon number remained constant across

treatments and time points (Supplemental Figure 2, Supplemental Tables 5 and 6).

Figure 2. Community analysis. Constrained correspondence analysis (CCA, constrained by 'Treatment' and 'Timepoint') of the Jaccard distances computed on relative genus abundances for all samples. **A-D)** CCA per SynCom (**A**: random; **B**: tolerant; **C**: mixed; **D**: sensitive); faceted by time point (columns) and colored by treatment. **E)** Relationship between change in bacterial growth in the presence of BX compounds in pure culture (x-axis) plotted against the change in relative abundance when embedded in a community (y-axis) for APO (blue) and BOA (orange). Each dot shows the average change in relative abundance for one isolate across all time points.

Time alone, however, did not significantly influence the species number in any of the SynComs.

The Shannon measure of alpha diversity, which in addition to the number of species also

incorporates the evenness of species abundances, was significantly influenced by both time and

treatment (Supplemental Figure 2, Supplemental Tables 5 and 6). While APO treatment

significantly increased Shannon diversity of the *mixed* and *tolerant* communities, it decreased

Shannon diversity of the *sensitive* one. In contrast, BOA significantly increased Shannon

diversity of the *sensitive* SynCom after 48 h (Supplemental Tables 7 and 8).

Isolates respond to APO in a community-specific fashion

Having substantiated that different SynComs display different resilience to BX treatment, we

asked if the tolerance of an isolate to BX would predict how well this isolate could establish itself

in a BX-treated community. Overall, there was no relationship between the relative growth of an

individual isolate in treatment conditions and and treatment-dependent changes in relative

abundance in a community (Figure 3A; adjusted R^2 = 0.01 for APO and BOA, p > 0.05, Table

S9). We also tested if there was a relationship between *absolute* growth in treatment and

changes in relative abundance and found that absolute growth was not predictive of

performance in a community either (adjusted R^2 = 0.02 for APO, and R^2 = -0.01 for BOA, p >

0.05, Table S9). In fact, some isolates that barely grew when cultured individually increased in

relative abundance when grown in a treated community (Figure 3A, top left corner of the plot).

After establishing that individual performance was a poor predictor of performance when

embedded in a community, we asked if the ability of an isolate to establish itself in a BX-treated

community instead depended on interactions with other community members. For this, we

exploited the partial isolate overlap between the communities (Figure 1A). We assumed that if

there were no community effects, an isolate would undergo the same treatment-related changes

in every community, independent of social context. We therefore explored the treatment-specific

changes in relative abundance for all isolates that were recovered in more than one community

(Figure 3B). In total, 10 isolates that were included in more than one community significantly

changed in relative abundance upon BX treatment (Figure 3B, FDR adjusted p < 0.05). Of

these, seven isolates changed strongly (absolute log2FC > 1) in relative abundance in a

treatment- and community-specific manner. For example, *Bacillus* Root239 did not change in

relative abundance when the *tolerant* community was APO treated, but was strongly reduced

under these conditions in the *mixed* community. Similarly, *Flavobacterium* Root420 was

unaffected by APO treatment when embedded in the *mixed* community but was strongly

reduced in relative abundance in the *random* community. *Acinetobacter* Root1280, *Bordetella*

Root565 and *Noviherbaspirillum* Root189 underwent mild (absolute log2FC < 1) fold-changes in

relative abundance when the communities they were embedded in were treated with APO.

While for these three isolates the direction remained the same, the amplitude of the change was

community-specific. This indicates that community-specific interactions between isolates can

modulate the impact of chemical exposure on the individual isolate.

264
Figure 3: Comparison between isolate growth alone and in different communities. A) Dotplot of
relative growth of an isolate in pure culture (ratio of area under the respective growth curves) on x plotted
against log2(Foldchange) in relative abundance when embedded in a community (Treatment / Control) on
y. The black line shows the regression line (y as the dependent variable, x as predictor) with 95%
confidence intervals as grey shading. **B)** Dotplots of log2(Foldchange) in relative abundance by treatment
for isolates found in multiple SynComs. Errors indicate s.e of the log2(Foldchange), colors indicate
SynCom. Changes in relative abundance were calculated across all time points; a breakdown by time
point is shown in Supplemental Figure 4.
273

274 However, it is important to consider not only the relative change per isolate but also the impact

275 of this change on overall community composition. For example, an isolate dropping from 50% to

276 25% relative abundance (l2fc of -1), may be considered a more important change in the larger

277 community context than an increase from 0.001% to 0.008% (l2fc of 3).

Interactions between isolates are modulated by community context and chemical treatment

Interactions between isolates are modulated by community context and

chemical treatment

In the next approach, we wanted to further investigate which isolates interacted in a treatment-

and community-specific manner. We focused our analysis on the *random* and *tolerant*

communities, since these shared 5 highly abundant genera, allowing for a comparative

approach (Figure 3B, Figure 4). To understand how SynCom-specific isolate interactions drove

condition-specific compositional changes at the genus level, we plotted the relative abundance

of highly abundant isolates (>1% average relative abundance) that belonged to genera shared

between *random* and *tolerant* (Figure 4A).

In APO conditions, the relative abundances of these highly abundant genera were similar in

both SynComs, irrespective of isolate identity (Figure 4A, blue shapes). However, when there

was no chemical selection (i.e., in BOA or Control treatments), the communities diverged: the

Bacillus isolate in *random* (Soil531) reached higher abundance than the one in *tolerant*

(Root239). Conversely, the genera *Acinetobacter* (Root1280) and *Pseudomonas* (Root9)

increased in abundance in *tolerant* compared with either the same strain (*Acinetobacter*

Root1280) or a closely related counterpart (*Pseudomonas* Root329) in *random*. Both

Pseudomonas isolates reached a lower relative abundance in control and BOA conditions than

in APO treatment (Figure 3B, 4A), suggesting that *Pseudomonas* profited from APO irrespective

of the community or strain identity, in line with the observation that *Pseudomonadaceae* in the

collection were generally APO insensitive (Figure 1B). The effect of APO on *Bacillus* was strain-

and community-specific: Root239 (*tolerant* SynCom) was lowly abundant in all SynCom /

treatment combinations, while Soil531 (*random*) was low in APO-treated samples but reached

20% relative abundance in the samples that were not treated with APO. Correlation analysis

using partial Pearson's correlation coefficients revealed that the relative abundance of *Bacillus*

Soil531 in the *random* SynCom was negatively correlated with the relative abundance of

Pseudomonas Root329 in the absence of APO treatment (Figure 4B). This suggested that there

was antagonism between these isolates: the high relative abundance of *Pseudomonas* Root329

in the APO-treated *random* community could be the result of a combination of its insensitivity to

APO with an APO-dependent reduction of its antagonist *Bacillus* Soil531 (which was less

abundant in APO). We also observed that both *Bacillus* isolates displayed community-specific

chemical sensitivity (Figure 3A, Supplemental Figure 3): while *Bacillus* Root239 had low relative

abundance in the *tolerant* community in all conditions, it reached >90% rel. abundance in BOA

and control conditions when part of the *mixed* community (Figure 4A, Supplemental Figure 4).

Similar to Soil531 in *random*, Root239 in *mixed* was strongly reduced in relative abundance

upon APO treatment (Figure 3A), coinciding with increased relative abundances of *Hoeflea*

Root268 and *Lysobacter* Root96 in the APO treated *mixed* community (Supplemental Figure 4),

further highlighting how treatment-specific isolate interactions affect community composition.

Figure 4: Differential isolate interactions in the *random* and *tolerant* SynComs are driven by isolate identity and chemical treatment. A) Relative abundance of Genera reaching more than 1% relative abundance and present in both SynComs. **B)** Partial Pearson's correlations (FDR adjusted p < 0.1) of isolate abundances in the *tolerant* and *random* SynComs. Each node represents one isolate: the inner edge of the circle indicates mean-sd, the outer edge shows mean + sd of the relative abundance. Color of the node label indicates if an isolate was a member of both (white) or only one (grey) of the SynComs. Each edge indicates a significant correlation between two isolates; edge color indicates the correlation value, negative correlations have a blue hue, while positive correlations are orange to yellow.

324 Discussion

325 Diversity of BX sensitivity in culturable rhizosphere bacteria

In our screen of the *AtRSphere* collection for sensitivity to BX derivatives (Figure 1), we

observed broad diversity of growth responses to BX exposure. While some clades were non-

responsive to BOA and APO, we observed diverse responses in most clades. APO in particular

exerted a strong negative influence on the growth of some isolates. These findings are in line

with previous results showing the antibiotic activity of APO on anaerobic microorganisms (Atwal

et al., 1992). In our study, BOA influenced bacterial growth only marginally compared to APO,

similar to what has been described for their effects on fungi and plants. In general, this

corroborates the previously suggested relevance of APO, which is more stable than BOA in soil,

in BX-mediated plant-environment interactions (Kumar et al., 1993; Schandry and Becker, 2020;

Schulz et al., 2013; Schütz et al., 2019; Venturelli et al., 2015; Voloshchuk et al., 2020).

However, our data showed that bacterial response to APO was largely isolate-specific and not

dictated by phylogenetic relationship (Figure 1B). This indicates that even within genera, natural

genetic diversity can result in substantial differences in BX sensitivity. Indeed, previous studies

reported functional overlap related to responses to plant-derived compounds between isolates

from different clades, while not all isolates of the same clade shared these functionalities (Bai et

al., 2015; Harbort et al., 2020). The broad phylogenetic spectrum of the *AtRSphere* collection

prevented the identification of genetic components underlying bacterial response to BOA and

APO. However, the strongly differential responses we observed among closely related isolates

warrant a more comprehensive analysis of the genetic basis of the BX response in the

respective clades in future experiments.

One major advantage of the *AtRSphere* collection is that it was isolated from a non-BX-

producing host plant. The soil from which this collection was originally derived (Cologne soil)

was collected from an agricultural site, and in absence of a complete record of the field's history,

we cannot exclude that there might have been an input of BX in the past. Our findings on the

standing diversity of BX tolerance imply that this diversity is maintained in plant-associated

bacteria, also in the absence of a BX-releasing host. Strain-specific tolerance or sensitivity to

plant specialized metabolites may thus contribute to the colonization profiles observed for

different plant species, where the composition is similar at higher taxonomic ranks but some

isolates show host preference (Bulgarelli et al., 2013; Schneijderberg et al., 2020).

355   Evaluation of bottom-up design strategy

Others have previously proposed that synthetic communities should be built with a focus on

functional features instead of on phylogenetic coverage (Lemanceau et al., 2017; Vorholt et al.,

2017). We employed a simple, function-guided strategy, compiling SynComs by binning

individual isolates based on a specific phenotype and contrasting these binned assemblies to a

randomly assembled SynCom. The decisive phenotype measured at the level of individual

isolates and used for binning them into SynComs was growth upon exposure to APO. The

phenotype measured at the community level, variance explained by treatment, was lowest

(22%) in the SynCom assembled solely from APO-tolerant isolates, indicating that resistance of

individual SynCom members leads to overall community resilience. While this approach worked

well to create an overall resilient community, the amount of variance explained by treatment was

similar in *random* and *sensitive* (~40%) and highest in the *mixed* SynCom (75%), indicating that

mixing tolerant and sensitive isolates does not necessarily result in a community with

intermediate resilience.

BX dependent community alterations *in vitro* mimic those described in soil

Our work complements recent research efforts to understand the effect of benzoxazinoids on plant-associated microbiota in soil (Cadot et al., 2020; Cotton et al., 2019; Hu et al., 2018; Kudjordjie et al., 2019; Neal et al., 2012). Ultimately, for *in vitro* SynCom systems to be a useful model for plant microbiota research, they should behave similarly to microbiota in soil. BX constitute a prime system to assess this, as the influence of BX release on the root and rhizosphere microbiome has been investigated via mutant studies in different soils (Cadot et al., 2020; Cotton et al., 2019; Hu et al., 2018; Kudjordjie et al., 2019). Among the bacterial taxa that were found in different studies, Flavobacteriales were observed as depleted in multiple studies. Hu et al (Hu et al., 2018) found *Flavobacterium* depleted on the roots of BX-exuding plants and the surrounding soils when comparing wildtype and *bx1* mutant plants. Extending these observations, Cadot et al (Cadot et al., 2020) reported that *Flavobacteriaceae* were consistently depleted on roots of BX-producing plants, independent of soil type and location. Similarly, members of the Flavobacteriales family were more abundant on the roots of maize BX-synthesis mutants than on wildtype in another, independent study (Cotton et al., 2019). In all of the afore-mentioned studies, *Comamonadaceae* were also reported to be depleted on roots of BX-releasing plants when compared to BX-deficient mutants (Cadot et al., 2020; Cotton et al., 2019; Hu et al., 2018; Kudjordjie et al., 2019).

Our data show that the growth of *Flavobacterium* isolates was only slightly inhibited in growth across APO levels, with some isolates displaying mild sensitivity to BOA (Figure 1). However, *Flavobacterium* was significantly depleted from the *random* and *sensitive* SynComs once the community was exposed to APO (Supplemental Figure 4). Similarly, *Acidovorax* (*Comamonadaceae*) was found to be reduced in the APO treatment in all SynComs. Thus, even

though *in vitro* cultivation strongly differs from soils in physical and chemical parameters and in

overall complexity, our study suggests that plant-exudate-mediated effects on soil communities

can be recapitulated *in vitro*, highlighting the applicability of such reduced-complexity systems

for future studies. In addition, such simplified systems may aid in the identification of microbial

isolates that act as community keystones, or hubs, under specific chemical challenges in a short

experimental time frame (Carlström et al., 2019).

Chemical stability of isolate networks

One might expect that in an *in vitro* system containing a mixture of isolates, one isolate

outgrows the others and reaches very high abundance. Under that assumption, there would be

no interactions between the individual isolates and growth rate would be a strong predictor of

final abundance. However, we found that the growth of an isolate by itself was only poorly

correlated with its ability to establish itself as a SynCom member *in vitro* or with its relative

abundance in a community (Figure 3A). In fact, several isolates that grew poorly when cultured

alone reached high relative abundance when grown in communities, possibly indicating

facilitation by other members or other types of positive interactions (Sanchez-Gorostiaga, et al.,

2019)

We found that some isolates that reached only low relative abundances in control communities

were able to establish themselves much better when the community was treated with APO

(Figure 4, Supplemental Figure 4). We propose that the positive effects of APO on certain

community members is not due to a direct positive effect of APO, but instead due to a negative

effect of APO on a competitor or antagonist in the same community. Interestingly, we have

observed two instances where the APO-dependent reduction of a *Bacillus* isolate coincided with

improved growth of other isolates (Figure 4, Supplemental Figure 4). *Bacillus* species are

generally competitive, and often produce a range of compounds that inhibit the growth of other

bacteria (Shafi et al., 2017) lending further support to the hypothesis that a negative impact of

APO on *Bacillus* isolates contributes to the APO-specific changes in community composition.

Bacillus members thus seem to shape the synthetic communities in a manner opposite to a

"keystone", whose presence typically facilitates the growth of other community members. This

phenomenon has been previously observed for *Bacillus* species and probably contributes to the

Bacillus-mediated biocontrol against certain plant-pathogens observed in natural soils (Lee et

al., 2021).

Methods
424

Isolate profiling
425

Individual isolates were pre-cultured in ½ strength TSB medium in 96-Well 2 ml deep-well plates

(Semadeni) covered with a Breathe-Easy (Diversified Biotech) membrane until stationary phase

for 6 d at 28°C and 180 rpm. Cultures were then resuspended by pipetting up and down 10

times and inoculated into 96-well flat-bottom, transparent cell culture plates (Nunc), containing

either solvent control (DMSO), 1 µM APO, 5 µM APO, 10 µM APO, 50 µM APO, 10 µM BOA, 50

µM BOA or 100 µM BOA in ½-strength TSB. Different media were prepared by mixing a stock

containing the highest concentration of the respective chemical, with medium containing the

solvent at the same concentration (DMSO; 1:20.000). Pipetting was carried out using an Agilent

Bravo pipetting robot equipped with a 96-tip head. Plates were stacked without lids, with one

empty plate at the top and the bottom of the stack and placed into a BioTek plate stacker

connected to a BioTek Synergy2 plate reader. Plates were shaken for 120 s before reading, and

absorption (600 nm) was measured in each well. Plates were continuously shaken, measured,

and stacked over the course of at least 48 h per replicate. For each replicate, the relative growth

per isolate and condition was calculated by dividing the area under the growth curve (x = time in

minutes; y = increase in OD600) in the respective condition by the area under the growth curve

of the control condition. The experiment was carried out in eight replicates per condition.

Synthetic bacterial communities

Design

To investigate to which degree the overall resilience of a bacterial community to chemical

changes is determined by the properties of its individual members, we followed two separate

design strategies. In the first strategy, we aimed at creating a diverse community spanning all

genera of the isolate collection. The *random* community was created by picking one random

isolate per genus.

In the second approach, we built communities based on the growth profiles of the individual

isolates. These communities span a limited number of genera, but the member growth profile is

more strictly defined than in the "full" community. One of the reduced representation

communities was created from isolates that grew to 75% to 100% of control conditions in 50µM

APO (*tolerant*). The other reduced community was assembled from isolates that grew to 0 to

25% of control in 50µM APO (*sensitive*). If there was more than one isolate per genus that

fulfilled the condition, we selected one at random. In addition, we compiled one limited

representation SynCom containing a mix of sensitive and tolerant isolates (*mixed*).

Cultivation and sequencing

Individual isolates were pre-cultured in ½-strength TSB medium in 96-Well 2ml deep-well plates

covered with a Breathe-Easy membrane (Diversified Biotech) with shaking at 28°C for 6 days

until stationary phase, as described previously (Bai et al., 2015). Individual cultures were

resuspended by pipetting up and down 10 times, and then 100 µL of each culture were mixed

with 10 mL ½-strength TSB medium containing solvent at the same concentration as the

treatment cultures (1:20.000 DMSO). 100 µL of the mixed culture were used to inoculate 20 mL

cultures containing either a solvent control (DMSO), 100 µM BOA, or 50 µM APO in 100 mL

Erlenmayer flasks. Flasks were cultured at 28°C, 180 rpm; 1 ml culture was sampled every 24

h. Sampled cultures were centrifuged at 12.000 rpm, the supernatant was discarded and the

pellet was stored at -70 °C until further use. DNA was extracted using the DNA PowerSoil kit

(Qiagen) following the standard protocol. 16S amplicons were generated according to the

Illumina 16S protocol, indexed using Nextera XT dual indexing (Illumina), and pooled per time

point for sequencing on a MiSeq (Paired end 300bp, Illumina).

Data analysis

RStudio with R version 4.0.2 was used for data analysis. The phylogenetic tree in Figure 1 was

computed from concatenated 16SrRNA, gyrB, gyrB, rpoB, recA, atpD, dnaJ and thrC DNA

sequences. These sequences were extracted from the fasta files containing nucleotide ORFs

(obtained from https://at-sphere.com/) using tBLASTn of the respective *E. coli* protein sequence.

The concatenated sequences were aligned using the DECIPHER R package (Wright, 2016) and

a maximum likelihood phylogeny with the general time reversible model was fitted using the

phangorn package (Schliep et al., 2017; Schliep 2011). Pagel's λ and the likelihood ratio test

results were obtained using the phylosig function from phytools (Revell, 2012). For the analysis

of 16S amplicon data, reads were processed and ASVs were inferred using the dada2 pipeline

(Callahan et al., 2016). Taxonomic assignment of ASVs was done using SynCom specific

16SrRNA databases and agglomerated at the genus level. ASVs that could not be assigned

were discarded. Partial Pearson's correlation coefficients were calculated and tested for isolates

with >100 reads per sample using the correlate R package. Scripts used for data analysis are

available at https://github.com/nschan/BxSynComMs. The following R packages were used with

R version 4.0.2:

Tidyverse (Wickham et al., 2019), Broom (Robinson et al., 2019), dada2 (Callahan et al., 2016),

DECIPHER (Wright, 2016), DESeq2 (Love et al., 2014), emmeans (Lenth et al., 2019), ggforce

(Pedersen, 2020a), ggthemes (Arnold, 2019), ggtree (Yu, 2020; Yu et al., 2018, 2017), gt

(Iannone et al., 2019), igraph (Csardi and Nepusz, 2006), MESS (Ekstrøm, 2016), multcomp

(Hothorn et al., 2008), patchwork (Pedersen, 2020b), phangorn (Schliep et al., 2017; Schliep,

2011), phyloseq (McMurdie and Holmes, 2013), phytools (Revell, 2012), vegan (Oksanen et al.,

2019), wesanderson (Ram and Wickham, 2018) in combination with some custom functions.

494 Data availability

495 Sequencing reads have been deposited to the European Nucleotide Archive (ENA;

496 www.ebi.ac.uk) under the accession no. PRJEB42439.

497 Acknowledgements

The authors gratefully acknowledge support from the Vienna BioCenter Core Facility (VBCF)

Molecular Biology Services, especially Dr. Robert Heinen, the VBCF Media Kitchen, the VBCF

Sequencing facility and the VBCF High Performance Computing facility. The authors also

gratefully acknowledge the Leibniz Supercomputing Centre (LRZ) for funding this project by

providing computing time on its Linux-Cluster and Rstudio Server.

The authors thank Patrick Hüther, Talia Karasov and Eva Knoch for helpful discussion and

constructive feedback on the manuscript, and Iacopo Gentile and Karina Weiser-Lobao for

conducting preliminary experiments and for technical support. The authors also thank Stijn

Staepen for help with the *AtRSphere* collection and Paul Schulze-Lefert for helpful discussion.

This work was funded by the Austrian Academy of Sciences (ÖAW) and the European

Research Council (ERC) under the European Union's Horizon 2020 research and innovation

programme (Grant agreement No. 716823, "FEAR-SAP").

510 Supplemental Material

511 The following supplemental material is contained in the file

512 Schandry_et_al_supplemental_material.pdf

513 Supplemental Tables

Supplemental Table 1. Phylogenetic signal

Supplemental Table 2. Syncom strains

Supplemental Table 3. PERMANOVA

Supplemental Table 4 PERMANOVA on treatment subsets

Supplemental Table 5. ANOVA: observed taxa

Supplemental Table 6. Observed diversity means and pairwise comparisons

Supplemental Table 7. ANOVA: Shannon Diversity

Supplemental Table 8. Shannon diversity means and pairwise comparisons

Supplemental Table 9. Correlation between change in relative abundance and change in

growth

524 Supplemental figures

525 **Supplemental Figure 1. Bx sensitivity of Xanthomonadales**

526 **Supplemental Figure 2. Observed alpha diversity**

527 **Supplemental Figure 3. Shannon diversity indices of the synthetic communities**

528   **Supplemental Figure 4. Changes in relative abundance by community**

Literature

Anzai K, Isono K, Okuma K, Suzuki S. 1960. The new antibiotics, questiomycins A and B. *J*
Antibiot **13**:125–132.
Arnold JB. 2019. ggthemes: Extra Themes, Scales and Geoms for "ggplot2."
Atwal AS, Teather RM, Liss SN, Collins FW. 1992. Antimicrobial activity of 2-aminophenoxazin-
3-one under anaerobic conditions. *Can J Microbiol* **38**:1084–1088.
Bai Y, Müller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M, Dombrowski N, Münch PC,
Spaepen S, Remus-Emsermann M, Hüttel B, McHardy AC, Vorholt JA, Schulze-Lefert P.
2015. Functional overlap of the Arabidopsis leaf and root microbiota. *Nature* 1–19.
Banerjee S, Schlaeppi K, van der Heijden MGA. 2018. Keystone taxa as drivers of microbiome
structure and functioning. *Nat Rev Microbiol* **16**:567–576.
Belz RG, Hurle K. 2005. Differential exudation of two benzoxazinoids--one of the determining
factors for seedling allelopathy of Triticeae species. *J Agric Food Chem* **53**:250–261.
Bodenhausen N, Bortfeld-Miller M, Ackermann M, Vorholt JA. 2014. A synthetic community
approach reveals plant genotypes affecting the phyllosphere microbiota. *PLoS Genet*
10:e1004283.
Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf
P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R, Eickhorst T,
Schulze-Lefert P. 2012. Revealing structure and assembly cues for Arabidopsis root-
inhabiting bacterial microbiota. *Nature* **488**:91–95.
Bulgarelli D, Schlaeppi K, Spaepen S, van Themaat EVL, Schulze-Lefert P. 2013. Structure and
Functions of the Bacterial Microbiota of Plants. *Annu Rev Plant Biol* **64**:807–838.
Cadot S, Guan H, Bigalke M, Walser J-C, Jander G, Erb M, van der Heijden M, Schlaeppi K.
2020. Specific and conserved patterns of microbiota-structuring by maize benzoxazinoids in
the field. *bioRxiv*. doi:10.1101/2020.05.03.075135
Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2:
High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**:581–583.
Callahan BJ, Sankaran K, Fukuyama JA, McMurdie PJ, Holmes SP. 2016. Bioconductor
Workflow for Microbiome Data Analysis: from raw reads to community analyses. *F1000Res*
5:1492.
Carlström CI, Field CM, Bortfeld-Miller M, Müller B, Sunagawa S, Vorholt JA. 2019. Synthetic
microbiota reveal priority effects and keystone strains in the Arabidopsis phyllosphere. *Nat*
Ecol Evol **3**:1445–1454.
Cheng C, Othman EM, Fekete A, Krischke M, Stopper H, Edrada-Ebel R, Mueller MJ, Hentschel
U, Abdelmohsen UR. 2016. Strepoxazine A, a new cytotoxic phenoxazin from the marine
sponge-derived bacterium Streptomyces sp. SBT345. *Tetrahedron Lett* **57**:4196–4199.
Cotton TEA, Pétriacq P, Cameron DD, Meselmani MA, Schwarzenbacher R, Rolfe SA, Ton J.
2019. Metabolic regulation of the maize rhizobiome by benzoxazinoids. *ISME J*.
doi:10.1038/s41396-019-0375-2
Csardi G, Nepusz T. 2006. The igraph software package for complex network research.
InterJournal.
De Roy K, Marzorati M, Van den Abbeele P, Van de Wiele T, Boon N. 2014. Synthetic microbial
ecosystems: an exciting tool to understand and apply microbial communities. *Environ*
Microbiol **16**:1472–1481.

Dessaux Y, Grandclément C, Faure D. 2016. Engineering the Rhizosphere. *Trends Plant Sci*
21:266–278.

Ekstrøm C. 2016. MESS: Miscellaneous Esoteric Statistical Scripts.

Finkel OM, Salas-González I, Castrillo G, Conway JM, Law TF, Teixeira PJPL, Wilson ED,
Fitzpatrick CR, Jones CD, Dangl JL. 2020. A single bacterial genus maintains root growth
in a complex microbiome. *Nature*. doi:10.1038/s41586-020-2778-7

Fitzpatrick CR, Salas-González I, Conway JM, Finkel OM, Gilbert S, Russ D, Teixeira PJPL,
Dangl JL. 2020. The Plant Microbiome: From Ecology to Reductionism and Beyond. *Annu*
Rev Microbiol. doi:10.1146/annurev-micro-022620-014327

Frey M, Chomet P, Glawischnig E, Stettner C, Grün S, Winklmair A, Eisenreich W, Bacher A,
Meeley RB, Briggs SP, Simcox K, Gierl A. 1997. Analysis of a chemical plant defense
mechanism in grasses. *Science* **277**:696–699.

Handrick V, Robert CAM, Ahern KR, Zhou S, Machado RAR, Maag D, Glauser G, Fernandez-
Penny FE, Chandran JN, Rodgers-Melnik E, Schneider B, Buckler ES, Boland W,
Gershenzon J, Jander G, Erb M, Köllner TG. 2016. Biosynthesis of 8-O-Methylated
Benzoxazinoid Defense Compounds in Maize. *Plant Cell* **28**:1682–1700.

Harbort CJ, Hashimoto M, Inoue H, Niu Y, Guan R, Rombolà AD, Kopriva S, Mathias J E E,
Sattely ES, Garrido-Oter R, Schulze-Lefert P. 2020. Root-Secreted Coumarins and the
Microbiota Interact to Improve Iron Nutrition in Arabidopsis. *Cell Host Microbe* **0**.
doi:10.1016/j.chom.2020.09.006

Herrera Paredes S, Gao T, Law TF, Finkel OM, Mucyn T, Teixeira PJPL, Salas González I,
Feltcher ME, Powers MJ, Shank EA, Jones CD, Jojic V, Dangl JL, Castrillo G. 2018. Design
of synthetic bacterial communities for predictable plant phenotypes. *PLoS Biol*
16:e2003962.

Hothorn T, Bretz F, Westfall P. 2008. Simultaneous inference in general parametric models.
Biom J **50**:346–363.

Hu L, Mateo P, Ye M, Zhang X, Berset JD, Handrick V, Radisch D, Grabe V, Köllner TG,
Gershenzon J, Robert CAM, Erb M. 2018. Plant iron acquisition strategy exploited by an
insect herbivore. *Science* **361**:694–697.

Hu L, Robert CAM, Cadot S, Zhang X, Ye M, Li B, Manzo D, Chervet N, Steinger T, van der
Heijden MGA, Schlaeppi K, Erb M. 2018. Root exudate metabolites drive plant-soil
feedbacks on growth and defense by shaping the rhizosphere microbiota. *Nat Commun*
9:2738.

Iannone R, Cheng J, Schloerke B. 2019. gt: Easily Create Presentation-Ready Display Tables.
R package version 0.1. 0.

Imai S, Shimazu A, Furihata K, Furihata K, Hayakawa Y, Seto H. 1990. Isolation and structure
of a new phenoxazine antibiotic, exfoliazone, produced by Streptomyces exfoliatus. *J*
Antibiot **43**:1606–1607.

Jacoby RP, Koprivova A, Kopriva S. 2020. Pinpointing secondary metabolites that shape the
composition and function of the plant microbiome. *J Exp Bot*. doi:10.1093/jxb/eraa424

Ke J, Wang B, Yoshikuni Y. 2020. Microbiome Engineering: Synthetic Biology of Plant-
Associated Microbiomes in Sustainable Agriculture. *Trends Biotechnol*.
doi:10.1016/j.tibtech.2020.07.008

Koprivova A, Schuck S, Jacoby RP, Klinkhammer I, Welter B, Leson L, Martyn A, Nauen J,
Grabenhorst N, Mandelkow JF, Zuccaro A, Zeier J, Kopriva S. 2019. Root-specific
camalexin biosynthesis controls the plant growth-promoting effects of multiple bacterial
strains. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1818604116

Kudjordjie EN, Sapkota R, Steffensen SK, Fomsgaard IS, Nicolaisen M. 2019. Maize
synthesized benzoxazinoids affect the host associated microbiome. *Microbiome* **7**:59.

Kumar P, Gagliardo RW, Chilton WS. 1993. Soil transformation of wheat and corn metabolites
mboa and DIM2BOA into aminophenoxazinones. *J Chem Ecol* **19**:2453–2461.

Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, Malfatti S,
Glavina del Rio T, Jones CD, Tringe SG, Dangl JL. 2015. Salicylic acid modulates
colonization of the root microbiome by specific bacterial taxa. *Science* **349**:860–864.
Lee, S.-M., Kong, H.G., Song, G.C., Ryu, C.-M. 2021) Disruption of Firmicutes and
Actinobacteria abundance in tomato rhizosphere causes the incidence of bacterial wilt
disease. *ISME J.* **15**: 330–347
Lemanceau P, Blouin M, Muller D, Moënne-Loccoz Y. 2017. Let the Core Microbiota Be
Functional. *Trends Plant Sci* **22**:583–595.
Lenth R, Singmann H, Love J, Buerkner P, Herve M. 2019. emmeans: Estimated Marginal
Means, aka Least-Squares Means (Version 1.3. 4).
Li L, Abou-Samra E, Ning Z, Zhang X, Mayne J, Wang J, Cheng K, Walker K, Stintzi A, Figeys
D. 2019. An in vitro model maintaining taxon-specific functional activities of the gut
microbiome. *Nat Commun* **10**:4146.
Liu Y-X, Qin Y, Bai Y. 2019. Reductionist synthetic community approaches in root microbiome
research. *Curr Opin Microbiol* **49**:97–102.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**:550.
Macías FA, Oliveros-Bastidas A, Marín D, Castellano D, Simonet AM, Molinillo JMG. 2005.
Degradation studies on benzoxazinoids. Soil degradation dynamics of (2R)-2-O-beta-D-
glucopyranosyl-4-hydroxy-(2H)- 1,4-benzoxazin-3(4H)-one (DIBOA-Glc) and its
degradation products, phytotoxic allelochemicals from Gramineae. *J Agric Food Chem*
53:554–561.
Macías FA, Oliveros-Bastidas A, Marín D, Castellano D, Simonet AM, Molinillo JMG. 2004.
Degradation Studies on Benzoxazinoids. Soil Degradation Dynamics of 2,4-Dihydroxy-7-
methoxy-(2 H)-1,4-benzoxazin-3(4 H)-one (DIMBOA) and Its Degradation Products,
Phytotoxic Allelochemicals from Gramineae. *J Agric Food Chem* **52**:6402–6413.
Maier L, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, Brochado AR, Fernandez
KC, Dose H, Mori H, Patil KR, Bork P, Typas A. 2018. Extensive impact of non-antibiotic
drugs on human gut bacteria. *Nature* **555**:623–628.
McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive analysis
and graphics of microbiome census data. *PLoS One* **8**:e61217.
Münkemüller, T., Lavergne, S., Bzeznik, B., Dray, S., Jombart, T., Schiffers, K., and Thuiller, W.
2012. How to measure and test phylogenetic signal: How to measure and test phylogenetic
signal. Methods Ecol. Evol. **3**: 743–756.
Neal AL, Ahmad S, Gordon-Weeks R, Ton J. 2012. Benzoxazinoids in root exudates of maize
attract Pseudomonas putida to the rhizosphere. *PLoS One* **7**:e35498.
Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB,
Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2019. vegan: Community
Ecology Package.
Pagel, M. 1999. Inferring the historical patterns of biological evolution. Nature **401**: 877–884.
Pedersen TL. 2020a. ggforce: Accelerating "ggplot2."
Pedersen TL. 2020b. patchwork: The Composer of Plots.
Ram K, Wickham H. 2018. wesanderson: A Wes Anderson Palette Generator.
Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and
other things). *Methods Ecol. Evol.* **3** 217-223.
Robert CAM, Veyrat N, Glauser G, Marti G, Doyen GR, Villard N, Gaillard MDP, Köllner TG,
Giron D, Body M, Babst BA, Ferrieri RA, Turlings TCJ, Erb M. 2011. A specialist root
herbivore exploits defensive metabolites to locate nutritious tissues. *Ecol Lett* **14**:55–64.
Robinson D, Hayes A, Couch S. 2019. Broom: Convert statistical objects into tidy tibbles.
Sanchez-Gorostiaga, A., Bajić, D., Osborne, M.L., Poyatos, J.F., Sanchez, A. 2019. High-order
interactions distort the functional landscape of microbial consortia. PLoS Biol. **17**:

e3000550.
Schandry N, Becker C. 2020. Allelopathic Plants: Models for Studying Plant–Interkingdom
Interactions. *Trends Plant Sci* **25**:176–185.
Schliep, Klaus, Potts, J. A, Morrison, A. D, Grimm, W. G. 2017. Intertwining phylogenetic trees
and networks. *Methods in Ecology and Evolution.* doi:10.1111/2041-210X.12760
Schliep KP. 2011. phangorn: phylogenetic analysis in R. *Bioinformatics.*
doi:10.1093/bioinformatics/btq706
Schneijderberg M, Cheng X, Franken C, de Hollander M, van Velzen R, Schmitz L, Heinen R,
Geurts R, van der Putten WH, Bezemer TM, Bisseling T. 2020. Quantitative comparison
between the rhizosphere effect of Arabidopsis thaliana and co-occurring plant species with
a longer life history. *ISME J.* doi:10.1038/s41396-020-0695-2
Schulz M, Marocco A, Tabaglio V, Macias FA, Molinillo JMG. 2013. Benzoxazinoids in rye
allelopathy - from discovery to application in sustainable weed control and organic farming.
J Chem Ecol **39**:154–174.
Schütz V, Bigler L, Girel S, Laschke L, Sicker D, Schulz M. 2019. Conversions of
Benzoxazinoids and Downstream Metabolites by Soil Microorganisms. *Frontiers in Ecology*
and Evolution **7**:238.
Shafi, J., Tian, H., and Ji, M. 2017. Bacillus species as versatile weapons for plant pathogens: a
review. Biotechnol. Biotechnol. Equip. **31**: 446–459.
Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, Berendsen RL,
Bakker PAHM, Feussner I, Pieterse CMJ. 2018. MYB72-dependent coumarin exudation
shapes root microbiome assembly to promote plant health. *Proc Natl Acad Sci U S A.*
doi:10.1073/pnas.1722335115
Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. 2020. Plant-microbiome interactions: from
community assembly to plant health. *Nat Rev Microbiol.* doi:10.1038/s41579-020-0412-1
Venturelli S, Belz RG, Kamper A, Berger A, von Horn K, Wegner A, Bocker A, Zabulon G,
Langenecker T, Kohlbacher O, Barneche F, Weigel D, Lauer UM, Bitzer M, Becker C.
2015. Plants Release Precursors of Histone Deacetylase Inhibitors to Suppress Growth of
Competitors. *Plant Cell* **27**:3175–3189.
Venturelli S, Petersen S, Langenecker T, Weigel D, Lauer UM, Becker C. 2016. Allelochemicals
of the phenoxazinone class act at physiologically relevant concentrations. *Plant Signal*
Behav **11**:e1176818.
Voges MJEEE, Bai Y, Schulze-Lefert P, Sattely ES. 2019. Plant-derived coumarins shape the
composition of an Arabidopsis synthetic root microbiome. *Proc Natl Acad Sci U S A*
116:12558–12565.
Voloshchuk N, Schütz V, Laschke L, Gryganskyi AP, Schulz M. 2020. The Trichoderma viride F-
00612 consortium tolerates 2-amino-3H-phenoxazin-3-one and degrades nitrated
benzo[d]oxazol-2(3H)-one. *Chemoecology.* doi:10.1007/s00049-020-00300-w
Vorholt JA, Vogel C, Carlström CI, Müller DB. 2017. Establishing Causality: Opportunities of
Synthetic Communities for Plant Microbiome Research. *Cell Host Microbe* **22**:142–155.
Vrancken G, Gregory AC, Huys GRB, Faust K, Raes J. 2019. Synthetic ecology of the human
gut microbiota. *Nat Rev Microbiol* **17**:754–763.
Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A,
Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D,
Seidel D, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome to
the Tidyverse. *JOSS* **4**:1686.
Wright ES. 2016. Using DECIPHER v2. 0 to analyze big biological sequence data in R. *R J* **8**.
Yu G. 2020. Using ggtree to Visualize Data on Tree-Like Structures. *Current Protocols in*
Bioinformatics. doi:10.1002/cpbi.96
Yu G, Lam TT-Y, Zhu H, Guan Y. 2018. Two methods for mapping and visualizing associated
data on phylogeny using ggtree. *Molecular Biology and Evolution.*

doi:10.1093/molbev/msy194
Yu G, Smith D, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods in*
Ecology and Evolution. doi:10.1111/2041-210X.12628