Deficiency of gluconeogenic enzyme PCK1 promotes non-alcoholic steatohepatitis progression and fibrosis through PI3K/AKT/PDGF axis activation

Qian Ye[1,#] Yi Liu[1,#], Guiji Zhang[1,#], Haijun Deng[1,#], Xiaojun Wang[2,#], Lin Tuo[3,†],Chang Chen[4], Xuanming Pan[1], Kang Wu[1], Jiangao Fan[5], Qin Pan[5], Kai Wang[1,*], Ailong Huang[1,*], Ni Tang[1,*]

[1]Key Laboratory of Molecular Biology for Infectious Diseases (Ministry of Education), Institute for Viral Hepatitis, Department of Infectious Diseases, The Second Affiliated Hospital, Chongqing Medical University, Chongqing, China

[2]Institute of Hepatobiliary Surgery, Southwest Hospital, Third Military Medical University (Army Medical University), Chongqing, China

[3]Department of Infectious Disease, Hospital of the University of Electronic Science and Technology of China and Sichuan Provincial People's Hospital, Chengdu, China

[4]Institute of Life Sciences, Chongqing Medical University, Chongqing, China

[5]Department of Gastroenterology, Xin Hua Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai, China

[#]These authors contributed equally：Qian Ye, Yi Liu, Guiji Zhang, Haijun Deng, Xiaojun Wang, Lin Tuo

Corresponding author: Ni Tang, Ailong Huang, Kai Wang, Key Laboratory of Molecular Biology for Infectious Diseases (Ministry of Education), Institute for Viral Hepatitis, Department of Infectious Diseases, The Second Affiliated Hospital, Chongqing Medical University, Chongqing, 400016, China. Tel: 86-23-68486780, Fax: 86-23-68486780, E-mail: nitang@cqmu.edu.cn (N.T.), ahuang@cqmu.edu.cn (A.H.), wangkai@cqmu.edu.cn (K.W.)

Abstract:

Nonalcoholic steatohepatitis (NASH) is a chronic liver disease characterized by hepatic lipid accumulation, inflammation, and progressive fibrosis. We demonstrated that phosphoenolpyruvate carboxykinase 1 (PCK1) plays a central role in NASH progression. Mice with liver *Pck1* deficiency fed a normal diet displayed hepatic lipid disorder and liver injury, whereas fibrosis and inflammation were aggravated in mice fed a NASH diet. Forced expression of PCK1 by adeno-associated virus in the liver ameliorated NASH in mice. PCK1 deficiency stimulated lipogenic gene expression and lipid synthesis. Moreover, loss of hepatic PCK1 activated the RhoA/PI3K/AKT pathway by increasing intracellular GTP levels, increasing secretion of platelet-derived growth factor-AA (PDGF-AA), and promoting hepatic stellate cell activation. Treatment with RhoA and AKT inhibitors or gene silencing of RhoA or AKT1 alleviated NASH progression *in vivo*. Hepatic PCK1 deficiency may be important in hepatic steatosis and fibrosis development through paracrine secretion of PDGF-AA, highlighting a therapeutic strategy for NASH.

Introduction

Non-alcoholic fatty liver disease (NAFLD) is the most common chronic liver disease worldwide, affecting nearly 25% of US and European adults [1]. NAFLD is characterized by aberrant lipid accumulation in hepatocytes in the absence of excessive alcohol consumption. Nonalcoholic fatty liver (NAFL) can progress to non-alcoholic steatohepatitis (NASH), a more serious form of liver damage hallmarked by irreversible pathological changes such as inflammation, varying degrees of fibrosis, and hepatocellular damage, which is more likely to develop into cirrhosis and hepatocellular carcinoma [2]. Although multiple parallel insults, including oxidative damage, endoplasmic reticulum stress, and hepatic stellate cell (HSC) activation, have been proposed to explain the pathogenesis of NASH, the underlying mechanisms remain unclear [3].

In gluconeogenesis, glucose is generated from non-carbohydrate substrates, such as glycerol, lactate, pyruvate, and glucogenic amino acids, mainly in the liver, to maintain glucose levels and energy homeostasis. Phosphoenolpyruvate carboxykinase 1 (PCK1) is the first rate-limiting enzyme in gluconeogenesis and converts oxaloacetate to phosphoenolpyruvate in the cytoplasm [4]. Our previous studies showed that PCK1 deficiency promotes hepatocellular carcinoma progression by enhancing the hexosamine-biosynthesis pathway [5]. However, PCK1 regulates not only glucose homeostasis but also lipogenesis by activating sterol regulatory

element-binding proteins [6]. Patients lacking PCK1 function present diffuse

hepatic macrosteatosis concomitant with hypoglycemia and hyperlactacidemia

[7]. Similarly, mice with reduced *Pck1* expression develop insulin resistance,

hypoglycemia, and hepatic steatosis, indicating the important role of PCK1 in

regulating both glucose homeostasis and lipid metabolism [8,9]. However, the

precise role of PCK1 in NASH progression is not well-understood.

The phosphoinositide 3-kinase/protein kinase B (PI3K/ATK) pathway plays a

critical role in regulating cell growth and metabolism. This pathway is activated

in response to insulin, growth factors, energy, and cytokines and, in turn,

regulates key metabolic processes such as glucose and lipid metabolism and

protein synthesis [10]. AKT promotes *de novo* lipogenesis primarily by activating

sterol regulatory element-binding protein [11]. PI3K/AKT dysregulation leads to

numerous pathological metabolic conditions, including obesity and type 2

diabetes [12]. NAFLD is characterized by disordered glucose and lipid

metabolism in the liver. Although the PI3K/AKT pathway is a key regulator for

sensing metabolic stress, its exact role in NAFLD/NASH progression is unclear

[13,14].

In this study, we explored the role of *Pck1* in a mouse NASH model. We

determined the molecular mechanisms underlying disordered lipid metabolism,

inflammation, and fibrosis induced by *Pck1* depletion. We also delineated the

functional importance of the PI3K/AKT pathway and paracrine secretion of

PDGF-AA as its effectors in steatohepatitis, providing a potential therapeutic

strategy for treating NASH.

Results

PCK1 is downregulated in patients with NASH and mouse models of NASH

To determine whether PCK1 is involved in NAFLD, we first examined hepatic gene expression in a published transcriptome dataset [Gene Expression Omnibus (GEO): GSE126848] containing samples from 14 healthy subjects, 12 patients with obesity, 15 patients with NAFLD, and 16 patients with NASH [15]. Bioinformatics analysis showed that 32 genes were markedly changed in obesity, NAFLD, and NASH; 12 genes were considerably downregulated and 20 genes were upregulated (Supplementary Fig. 1a–c). Notably, *PCK1* was gradually reduced in patients with obesity, NAFLD, and NASH (Fig. 1a, b). Downregulation of *PCK1* mRNA was also observed in a similar dataset (GSE89632) (Fig. 1b). Moreover, immunohistochemistry (IHC) assays showed that hepatic PCK1 protein levels were significantly lower in patients with NASH than in healthy subjects (Fig. 1c). Similarly, *PCK1* mRNA and protein levels decreased in the liver of mice fed a high-fat diet with drinking water containing fructose and glucose (NASH diet) for 24 weeks (Fig. 1d,e).

Next, palmitic acid (PA) was used to mimic the liver steatosis of patients with NAFLD *in vitro* [16]. Cell growth was assessed in a CCK8 assay after treatment with different concentrations of PA (Supplementary Fig. 1d). Interestingly, the PCK1 mRNA and protein levels were downregulated in a dose-dependent

manner during 24 h PA stimulation (Fig. 1f, g), suggesting that the transcription

of PCK1 was inhibited in response to lipid overload. We screened several

known regulators of *PCK1* (Supplementary Fig. 1e, f) and found that activating

transcription factor 3 (*ATF3*), a transcriptional repressor of *PCK1* [17], was

upregulated upon PA stimulation (Fig. 1h). Similarly, ATF3 expression was

remarkably upregulated in liver samples derived from patients with NASH and

NASH model mice (Fig. 1i, j, Supplementary Fig. 1g, h). Chromatin

immunoprecipitation assays revealed that the binding of ATF3 to the *PCK1*

promoter increased following PA administration (Fig. 1k). ATF3 knockdown

restored PCK1 expression in human hepatocytes treated with PA (Fig. 1l).

Furthermore, correlation analysis revealed that *ATF3* was negatively

correlated with the *PCK1* mRNA level based on the GEO database (GEO:

GSE135251) (Fig. 1m). These results indicate that increased lipid intake led to

the upregulation of the repressor ATF3, impairing *PCK1* transcription in

patients with NASH and mouse models.

L-KO mice exhibit a distinct hepatic steatosis phenotype

To explore the role of *Pck1* in fatty liver disease, wild-type (WT) and

liver-specific *Pck1*-knockout mice (L-KO) mice were fed a chow diet for 24

weeks (Fig. 2a). Hepatic-specific depletion of PCK1 was confirmed by

performing immunoblotting (Fig. 2b). Starting at 16 weeks, L-KO mice showed

an increased body weight compared with WT mice, however, the results of the

glucose tolerance test (GTT) and insulin tolerance test (ITT) did not

significantly differ (Fig. 2c). Moreover, significant hepatomegaly and an

increased liver weight were observed in L-KO mice (Fig. 2d). Alanine

transaminase (ALT) and aspartate transaminase (AST) levels were higher in

L-KO mice, indicating liver injury (Fig. 2e). In addition, total triglyceride (TG),

total cholesterol (TC), and free fatty acids (FFAs) in the liver tissues and serum

were elevated in L-KO mice compared to those in WT mice (Fig. 2f, g).

Histochemistry and enzyme-linked immunosorbent assay (ELISA) showed that

L-KO mice had prominent hepatic steatosis, increased inflammatory infiltration,

and high levels of TNF-α, whereas hepatic fibrosis was not observed (Fig.

2h–j). Additionally, PCK1 deficiency significantly increased the mRNA levels of

genes related to fatty acid transport and inflammation, whereas there were no

significant changes in fibrosis-related genes in L-KO mice fed the chow diet

(Fig. 2k). These data suggest that L-KO mice exhibited a distinct hepatic

steatosis phenotype and liver injury even when fed normal chow.

Hepatic loss of *Pck1* promotes inflammation and fibrogenesis in NASH

mice

To explore whether an unhealthy diet could exacerbate pathologic changes in

L-KO mice, WT and L-KO mice were fed NASH diet (Fig. 3a) [18,19]. Starting at 4

weeks, L-KO mice showed significant weight gain (Fig. 3b). The GTT and ITT

showed that L-KO mice developed a more severe form of glucose intolerance

and insulin resistance compared with those in WT mice (Supplementary Fig.

2a, b). L-KO mice had pale and heavier livers compared with WT mice (Fig.

3c), although there was no significant difference in the liver weight ratio

(Supplementary Fig. 2c). Insulin, AST, ALT, TC, TG, and FFAs increased in

the serum and liver homogenates of L-KO mice, suggesting more serious liver

injury and lipid metabolism disorder (Fig. 3d, Supplementary Fig. 2d, e).

Analyses of L-KO liver sections revealed increased fat droplets, more severe

fibrosis, and greater macrophage infiltration (Fig. 3e, Supplementary Fig. 2f).

Furthermore, L-KO mice exhibited higher NAFLD activity scores (NAS score)

and higher TNF-α and IL-6 levels (Fig. 3f, g). In addition, the expression of

inflammatory factors, lipogenic enzymes, and fibrogenesis-associated genes

was upregulated in L-KO mice (Supplementary Fig. 2g). In summary, mice

lacking hepatic *Pck1* showed substantial liver inflammation and fibrosis after

being fed the NASH diet.

AAV-mediated restoration of hepatic PCK1 alleviates the NASH

phenotype in *Pck1*-deficient mice

We then investigated whether adeno-associated virus (AAV)-based *Pck1*

replacement therapy could reverse ongoing liver derangement, which is

typically observed in patients with NASH. After 10 weeks of chow or NASH diet

feeding, WT and L-KO mice were injected through the tail vein with AAV

serotype 8 (AAV8) vector expressing *Pck1* under the control of a liver-specific

promoter (thyroxine-binding globulin, TBG), AAV8-TBG-*Pck1* or

AAV8-TBG-*control* (Fig. 3h). The expression of PCK1 in various tissues was

identified by Western blotting (Supplementary Fig. 2h). Interestingly, mice with

PCK1 re-expression showed a lower liver weight, body weight, serum liver

enzymes, and serum lipid contents (Fig. 3i, j, Supplementary Fig. 2i, j).

Moreover, lipid deposition, inflammation, and fibrosis significantly improved in

L-KO mice injected with AAV8-TBG-*Pck1* (Fig. 3k, l). Hepatic gene expression

analyses indicated that the expression of genes involved in inflammation and

liver fibrosis was greatly attenuated by PCK1 restoration in L-KO mice

(Supplementary Fig. 2k). Overall, these data support that forced PCK1

expression in the liver protects against NASH in mice.

Transcriptomic and metabolomics analyses confirmed that the loss of

***Pck1* promotes hepatic lipid accumulation**

To comprehensively investigate the role of *Pck1* deficiency in NASH

development, we performed RNA-seq analysis of liver samples from L-KO and

WT mice fed the normal chow or NASH diet for 24 weeks. Gene Ontology

analysis indicated that lipid metabolic processes were remarkably upregulated

in L-KO mice fed the NASH diet (Fig. 4a). The volcano plot showed that genes

involved in fatty acid uptake, such as solute carrier family 27 member 1

(*Slc27a1*) and fatty acid translocase (*Cd36*), and lipid droplet synthesis, such

as cell death-inducing DFFA like effector C (*Cidec*) and cell death-inducing

DFFA like effector A (*Cidea*), were upregulated in response to the NASH diet

(Fig. 4b). Gene Set Enrichment Analysis (GSEA) revealed that the PPAR

signaling pathway was prominently upregulated in L-KO mice fed either diet

(Fig. 4c, Supplementary Fig. 3a, b). Several genes selected from the dataset

were independently validated by quantitative polymerase chain reaction

(qPCR) and immunoblotting and found to be significantly overexpressed in

L-KO mice (Fig. 4d, e). Furthermore, genes involved in the glycerol

3-phosphate (G3P) pathway were upregulated in L-KO mice (Fig. 4f).

Metabolomics analysis showed that compared with WT mice fed the NASH

diet, L-KO mice had significantly higher G3P and PA levels (Fig. 4g, h). As G3P

is a substrate for TG synthesis and PA is a key intermediate metabolite in *de*

novo lipogenesis, *Pck1* ablation may promote substrate accumulation for lipid

synthesis.

To further examine the function of PCK1 in steatosis *in vitro*, we overexpressed

(*PCK1*-OE) using the AdEasy adenoviral vector system and knocked out

PCK1 (*PCK1*-KO) using the CRISPR-Cas9 system in human hepatocytes

(Supplementary Fig. 3c, d). We found that *PCK1*-OE attenuated the

accumulation of lipid droplets, whereas *PCK1*-KO facilitated lipid accumulation

(Supplementary Fig. 3e, f). Collectively, these results suggest that hepatic

Pck1 deficiency leads to lipid accumulation by promoting the expression of

lipogenic genes and accumulation of substrates related to lipid synthesis

(Supplementary Fig. 3g).

Hepatic *Pck1* deficiency leads to HSC activation via PI3K/AKT pathway

RNA-seq analysis indicated that the PI3K/AKT pathway was specifically

activated in L-KO mice fed the NASH diet (Fig. 5a, b). Immunoblotting

revealed p-AKT (S473) and p-AKT (T308), which are two activated forms of

AKT, and downstream c-MYC were significantly upregulated in both the livers

and primary hepatocytes of L-KO mice fed the NASH diet (Fig. 5c, d). qPCR

confirmed that genes related to the PI3K/AKT pathway were highly expressed

in L-KO mice (Supplementary Fig. 4a). Similarly, p-AKT (S473 and T308)

significantly decreased in human *PCK1*-OE cells but increased in *PCK1*-KO

cells after 0.2 mM PA treatment (Fig. 5e, f).

To clarify the role of PI3K/AKT pathway activation, transcriptome data were

further analyzed. Interestingly, *Col1a1*, *Col3a1*, and *Lama2*, which are primary

components of the extracellular matrix (ECM), were upregulated, as shown in

the heat map of the PI3K/AKT pathway (Supplementary Fig. 4b). Moreover,

GSEA analysis revealed that ECM-receptor interaction was upregulated in

L-KO mice (Supplementary Fig. 4c). Because ECM deposition is typically

considered as the key event underlying liver fibrosis, we predicted that the

activation of the PI3K/AKT pathway promotes fibrosis in L-KO mice. HSCs are

major ECM secretors; thus, we performed co-culture assays with human

hepatocyte (MIHA) and human hepatic stellate cell lines (LX-2) (Fig. 5g).

Interestingly, the mRNA levels of *ACTA2* (α-SMA, an HSC activation marker),

COL1A1, and *COL3A1* increased in LX-2 cells co-cultured with *PCK1*-KO cells

but decreased in LX-2 cells co-cultured with *PCK1*-OE cells (Fig. 5h, i).

Similarly, COL1A1, COL3A1, and α-SMA expression increased in the liver

tissues and primary HSCs of L-KO mice (Fig. 5j, k), which was confirmed in

IHC analysis of COL3A1 (Supplementary Fig. 4d). However, these increases

were partially reversed by MK2206, an AKT inhibitor (Fig. 5l, m). Collectively,

these data suggest that the loss of PCK1 in hepatocytes induces HSC

activation and ECM formation by activating the PI3K/AKT pathway.

259 Paracrine PDGF-AA from hepatocytes promotes HSC activation

Hepatocytes elicit several fibrogenic actions in a paracrine manner to promote

HSC activation [20]. Thus, PCK1-mediated hepatic fibrosis may be involved in

paracrine disorders. To test this hypothesis, several pro-fibrotic factors were

screened; *Pdgfa* was found to be significantly elevated in the liver tissues of

L-KO mice (Fig. 6a). Bioinformatics analysis confirmed that *PDGFA* was

significantly increased in patients with NAFLD and NASH (Fig. 6b). *Pdgfa*

encodes a dimer disulfide-linked polypeptide (PDGF-AA), and the chronic

elevation of PDGF-AA in the mouse liver induces fibrosis [21]. Immunoblotting

and ELISA revealed increased PDGF-AA expression in the liver tissues,

primary hepatocytes, and plasma of L-KO mice (Fig. 6c–f). Moreover, the

PDGF-AA concentration markedly increased in the culture medium of

PCK1-KO cells but decreased in that of *PCK1*-OE cells treated with 0.2 mM

PA (Fig. 6g, h). Correspondingly, platelet-derived growth factor receptor alpha

(*PDGFRA*), which encodes the PDGF-AA receptor, increased in LX-2 cells

co-cultured with *PCK1*-KO cells but decreased in LX-2 cells co-cultured with

PCK1-OE cells (Fig. 6i, j). To determine whether the pro-fibrogenic effect was

mediated by PDGF-AA secretion, we treated the cells with a neutralizing

antibody against PDGF-AA. As expected, the increases in α-SMA, COL1A1,

and COL3A1 in LX-2 cells co-cultured with *PCK1*-KO cells were reversed by

anti-PDGF-AA treatment (Fig. 6k).

A review of the transcriptome data showed that *Pdgfa* appeared in the heat

map of the PI3K/AKT pathway (Supplementary Fig. 4b). The IHC results

showed that p-AKT (S473) was positively correlated with PDGF-AA (Fig. 6l).

The AKT inhibitor MK2206 significantly blocked the increase in *PDGFA* levels

in the supernatants or cells lysates of *PCK1*-KO cells (Fig. 6m–o). Taken

together, these data confirm that PCK1 deficiency promoted PDGF-AA

expression through the PI3K/AKT pathway and activated HSCs through

hepatocyte-HSC crosstalk.

PCK1 deficiency promotes the activation of the PI3K/AKT/PDGF-AA axis

by activating RhoA signaling in hepatocytes

Rho GTPases, which cycle between active GTP-bound and inactive GDP-bound conformations, activate the PI3K/AKT pathway [22-24]. Considering that PCK1 catalyzes the conversion of oxaloacetate to phosphoenolpyruvate, consuming GTP to generate GDP, we predicted that PCK1 deficiency alters intracellular GTP homeostasis. To test this hypothesis, high-performance liquid chromatography (HPLC) analysis was conducted to detect the intracellular levels of GTP. Interestingly, intracellular GTP levels decreased in *PCK1*-OE cells (Supplementary Fig. 5a) but increased in *PCK1*-KO cells (Supplementary Fig. 5b). Considering that Rho GTPases are activated when combined with GTP[25], we examined the proteins levels of several Rho GTPases in the mouse liver tissues and found that GTP-bound RhoA significantly increased and inactivated RhoA, p-RhoA (S188), decreased in L-KO mice (Fig. 7a–c). Similar results were observed in the primary hepatocytes of mice fed the NASH diet (Supplementary Fig. 5c). Consistently, after PA treatment, the levels of GTP-bound RhoA decreased and p-RhoA (S188) expression increased in *PCK1*-OE cells, whereas the opposite results were observed in *PCK1*-KO cells (Fig. 7d–g). Next, RhoA inhibitor (Rhosin) or shRhoA was used to determine whether PI3K/AKT activation depends on RhoA. Immunoblotting, ELISA, and qPCR assays showed that both Rhosin and shRhoA blocked the increase in the activated forms of AKT and PDGF-AA

in the *PCK1*-KO cell lysate and supernatant, as well as *ACTA2*, *COL1A1*, and

COL3A1 expression in LX-2 cells co-cultured with *PCK1*-KO hepatocytes (Fig.

7h–j, Supplementary Fig. 5d–f). To further evaluate the involvement of RhoA in

HSC activation, we isolated primary hepatocytes from NASH-fed WT or L-KO

mice; treated the cells with MK2206, Rhosin, or dimethyl sulfoxide vehicle; and

then co-cultured the cells with primary HSCs from chow-fed WT mice. The

activation of co-cultured HSCs was partially eliminated by the inhibition of AKT

or RhoA in primary hepatocytes (Supplementary Fig. 5g). Moreover, PCK1

and p-RhoA (S188) were downregulated in samples from patients with NASH,

whereas p-AKT (S473) and PDGF-AA levels were upregulated (Fig. 7k, l).

These data indicate that PCK1 ablation stimulated the PI3K/AKT/PDGF-AA

axis by activating RhoA.

Genetic or pharmacological disruption of RhoA and AKT1 reduced

progressive liver fibrosis *in vivo*

To explore whether blocking RhoA/PI3K/AKT could rescue the NASH

phenotype in L-KO mice, the genetic and pharmacological disruption of RhoA

and AKT1 was performed *in vivo* (Fig. 8a). Pharmacological inhibition of AKT1

or RhoA led to improved glucose intolerance (Supplementary Fig. 6a) and

insulin resistance (Supplementary Fig. 6b). The increase in liver weight was

also prevented (Fig. 8b), whereas the body weight decreased only in the

MK2206 treatment group (Supplementary Fig. 6c). Additionally, Rhosin or

MK2206 administration attenuated AST and ALT levels, as well as TG and

FFA levels, in the serum and liver tissues (Figs. 8c, Supplementary Fig. 6d, e).

Similarly, histochemistry revealed reduced liver steatosis, inflammation, and

fibrosis in Rhosin- or MK2206-treated mice (Figs. 8d, Supplementary Fig. 6f),

which was confirmed based on the decreased liver TNF-α and IL-6 levels (Fig.

8e). Additionally, α-SMA, COL1A1, COL3A1, PDGF-AA, and p-AKT (S473,

T308) expression and GTP-bound RhoA levels decreased, whereas p-RhoA

(S188) expression increased in the treatment group (Supplementary Fig. 6g–i).

MK2206 or Rhosin treatment also reduced the expression of genes related to

inflammation and fibrosis (Supplementary Fig. 6h). These results indicate that

the pharmacological inhibition of AKT or RhoA can alleviate the clinical

phenotypes of NASH.

Next, we used pSECC, a lentiviral-based system that combines the CRISPR

system and Cre recombinase, to silence RhoA or AKT1 in L-KO mice [26]. In

agreement with the results of the experiments described above, AKT or RhoA

depletion *in vivo* partially prevented and liver weight and body weight gain (Fig.

8f,g). Moreover, serum AST, ALT, TC, TG, and FFA levels significantly

decreased in sgAkt1- or sgRhoA-treated mice compared with those in

sgCtrl-treated L-KO mice (Fig. 8h, Supplementary Fig. 7a). Histological

analysis revealed that the NAS score, hepatic fibrosis, and F4/80[+]

macrophage counts decreased in sgAkt1 and sgRhoA mice, indicating the

attenuation of the NASH phenotypes (Fig. 8i, Supplementary Fig. 7b, c).

Consistently, genes or proteins involved in hepatic inflammation and fibrosis

were significantly downregulated in the livers of sgAkt1 and sgRhoA mice,

accompanied by the reduced expression of PDGF-AA (Supplementary Fig. 7d,

e). These results indicate that the genetic inhibition of AKT or RhoA can

alleviate the clinical phenotypes of NASH.

Discussion

We found that the hepatic gluconeogenic enzyme PCK1 plays an important

role in NASH progression. The expression of PCK1 was diminished in the

livers of patients or mice with NASH. Moreover, deletion of PCK1 significantly

exacerbated hepatic steatosis, fibrosis, and inflammation in mouse models fed

the NASH diet. Mechanistically, loss of PCK1 not only promotes steatosis by

enhancing lipid deposition, but also induces fibrosis through HSC activation via

paracrine secretion of PDGF-AA, thus promoting NASH progression (Fig. 8j).

Abnormal lipid metabolism is characteristic of NAFLD and NASH. Previous

studies predicted that altered lipid homeostasis was caused by abnormal

expression of genes related to lipid metabolism [27]. However, recent studies

demonstrated that disruption of gluconeogenesis also leads to abnormal lipid

metabolism. A deficiency of fructose-1,6-bisphosphatase 1 and

glucose-6-phosphatase catalytic subunit, which are key enzymes in

gluconeogenesis, results in severe hepatic steatosis and hypoglycemia,

indicating that the suppression of gluconeogenesis also disrupts lipid

homeostasis [16,28]. As the first rate-limiting enzyme in gluconeogenesis, it is

currently unclear whether PCK1 plays a critical role in NAFLD/NASH

development. We identified a robust decrease in PCK1 expression in the livers

of NASH mice and patients with NAFLD/NASH, causing severe hepatic

steatosis and confirming that disordered hepatic gluconeogenesis affects lipid

homeostasis.

Previous reports showed that PCK1 expression is increased in several

obesity/diabetes mouse models, such as ZDF rats and *ob/ob* and *db/db* mice,

and the disease progression of NASH is positively correlated with obesity and

type 2 diabetes mellitus [29-31]. Interestingly, we found that PCK1 expression

was downregulated in a diet-induced murine NASH model. This discrepancy

may be related to differences in the animal models used in different studies.

Widely used rodent models for genetic forms of obesity and diabetes, such as

ob/ob and *db/db* mice, exhibit increased plasma glucocorticoids, which may

drive PCK1 expression [31,32]. Another explanation is that the high-fat diet

supplemented with high fructose/glucose in drinking water could have

suppress PCK1 expression [33]. Under a high-fat diet, PA can decrease the

expression of PCK1 by inhibiting the expression of SIRT3 [34]. Additionally,

acetylation, ubiquitination, and phosphorylation modulate PCK1 expression [35].

A high glucose level can reportedly destabilize PCK1 by stimulating its

acetylation, thus promoting its ubiquitination and subsequent degradation [36]. In

this study, we found that ATF3, a member of the basic leucine zipper family of

transcription factors [37], transcriptionally repressed *PCK1* upon PA overload *in*

vitro and in the NASH mouse model. This result agrees with those of previous

studies suggesting that ATF3 is upregulated in patients with NAFLD and

murine NASH models and inhibits the expression of PCK1 in alcoholic fatty

liver disease [16,38,39]. Therefore, in this study, we found that PCK1 markedly

decreased in NASH, and PA inhibited *PCK1* transcription by upregulating

ATF3.

Numerous studies using PCK1 agonists or whole-body *Pck1* knockdown mice

have verified that PCK1 can affect lipid metabolism [40,41]. In the present study,

liver-specific *Pck1* knockout induced significant hepatic steatosis even under

normal feeding conditions. This is very important because it is uncommon for

single gene ablation to cause spontaneous steatosis unless a high-fat diet is

used. Moreover, mice with liver *Pck1* deficiency present aggravated

inflammation when fed a high-fat high-fructose diet, contrasting with the results

of a previous study showing that whole-body *Pck1* knockdown prevents

hepatic inflammation [42]. This discrepancy may be related to differences

between diets and animal models, as whole-body *Pck1* knockdown may have

unexpected effects on glucolipid metabolism. However, whether the

restoration of PCK1 expression in the NASH model of L-KO mice can reverse

NASH progression requires further investigation.

Lipid accumulation is characteristic of steatosis. Emerging evidence indicates

that increased fatty acid uptake is associated with lipid accumulation [41,42].

Previous studies demonstrated that the loss of PCK1 in the liver disturbed

hepatic cataplerosis and led to the accumulation of TCA cycle intermediates.

The slowed TCA cycle impaired fatty acid oxidation, resulting in fat

accumulation in the liver [43]. Additionally, elevated plasma TGs and FFAs could

contribute to body weight gain in L-KO mice. Notably, NASH diet-induced

glucose intolerance and insulin resistance in L-KO mice play essential roles in

obesity and whole-body metabolism. In this study, genes involved in fatty acid

uptake such as *Cd36* and *Slc27a1* were highly expressed in L-KO mice. In

addition, a lipid droplet-associated protein Cidec was increased by both the

chow and NASH diets, and was shown to be upregulated in patients with

NALFD and L-KO mice, suggesting that PCK1 ablation promotes lipid droplet

formation [44,45]. Abnormal levels of metabolites also contribute to TG

accumulation in the liver, with the G3P pathway contributing to over 90% of TG

synthesis [46]. As our metabolomics data showed that G3P and PA were

significantly upregulated in L-KO mice, PCK1 deficiency may promote hepatic

lipid accumulation by enhancing the expression of Cd36, Slc27a1, and Cidec

and the levels of metabolic substrates such as G3P and PA. However, the

precise mechanism by which PCK1 regulates the G3P pathway and

expression levels of *Cd36* and *Slc27a1* must be further analyzed.

Fibrosis is another characteristic of NASH and drives the transition from simple

steatosis to NASH. Activation of HSCs through the secretion of profibrotic

cytokines, such as TGF-β and PDGF, is a key event in liver fibrosis [43]. A recent

study identified high mobility group protein B1, secreted by

fructose-1,6-bisphosphatase 1-deficient hepatocytes, as the main mediator

activating HSCs, revealing important crosstalk between hepatocytes and

HSCs via paracrine signaling. Herein, PDGF-AA was secreted by

PCK1-deficient hepatocytes and acted in a paracrine manner to activate HSCs.

Increased deposition of ECM and activation of HSCs were observed in

PDGFA-transgenic mice; however, the mechanism mediating PDGF-AA

upregulation in fibrosis remains unclear [21]. Here, we demonstrated that PCK1

deficiency promoted PDGF-AA secretion by activating the RhoA/PI3K/AKT

pathway. Mechanistically, PCK1 deletion may increase intracellular GTP levels,

thus promoting the activation of RhoA and further activating the PI3K/AKT

pathway.

Most Rho GTPases cycle between an active GTP-bound and an inactive

GDP-bound form, a process that is regulated by guanine nucleotide exchange

factors, GTPase-activating proteins, and guanine nucleotide dissociation

inhibitors. Guanine nucleotide exchange factors can activate Rho GTPases by

catalyzing the exchange of GDP for GTP when the intracellular concentration

of GTP is high [44]. Several members of the Rho-GTPase family, such as Rac1,

RhoA, and RhoC, can be activated by increased concentrations of intracellular

GTP [45-47] [48]. We found that PCK1 deficiency activated RhoA by increasing the

levels of intracellular GTP, therefore activating the downstream PI3K/AKT

pathway. Moreover, the genetic and pharmacological disruption of RhoA and

AKT1 can effectively mitigate NASH phenotypes in L-KO NASH mice. In

addition, RhoA and AKT inhibitors can reportedly inhibit the progression of

NASH through other pathways, such as the NF-κB [49], Hippo [50,51], and Notch [52]

signaling pathways. Although RhoA and AKT inhibitors are currently only in

phase 3 trials or preclinical studies for the treatment of liver fibrosis or clinical

tumors, these compounds also show therapeutic potential for NASH [53-55].

In conclusion, hepatic PCK1 deficiency promoted lipid deposition and fibrosis

in a murine NASH model. Moreover, hepatic PCK1 loss activated the

RhoA/PI3K/AKT pathway, which increased PDGF-AA secretion and promoted

HSC activation. AKT/RhoA inhibitors reduced progressive liver fibrosis,

providing a therapeutic strategy for NASH treatment.

Materials and Methods

Animal models

$Pck1^{loxp/loxp}$ mice on a 129S6/SvEv background were purchased from the Mutant Mouse Resource & Research Center (MMRRC: 011950-UNC; Bar Harbor, ME, USA) and *Alb-Cre* mice on a C57BL/6 background were purchased from Model Animal Research Center of Nanjing University (Nanjing, China). To generate liver-specific *Pck1*-knockout mice (L-KO), *Alb-Cre* mice were crossed with $Pck1^{loxp/loxp}$ mice. $Pck1^{loxp/loxp}$ mice from the same breeding step were used as controls (wild-type, WT). Male WT and L-KO mice at 7–9 weeks old were fed the NASH diet (D12492: 60% Kcal fat, with drinking water containing 23.1 g/L fructose and 18.9 g/L glucose; Research Diets, New Brunswick, NJ, USA) (n = 11 per group) or control chow diet (D12450J: 10% Kcal fat, with tap water; Research Diets) (n = 10 per group) for 24 weeks. Food and drinking water were provided *ad libitum*. All mice were housed in temperature-controlled (23 °C) pathogen-free facilities with a 12 h light-dark cycle.

For AAV8 transduction, AAV8-TBG-*control* and AAV8-TBG-*Pck1* were purchased from the Shanghai Genechem Co., Ltd. (Shanghai, China) and injected via the tail vein following 10 weeks of NASH diet feeding (2×10^{11} genome copies/mouse). The efficiency of virus infection and expression of PCK1 in mouse hepatocytes were confirmed by western blot analysis. After a

total of 24 weeks of NASH diet feeding, the mice were sacrificed for analysis.

Genetic and pharmacological inhibition of AKT1 or RhoA were performed *in*

vivo. Genetic depletion of mouse RhoA or AKT1 *in vivo* was conducted using

pSECC (#60820; Addgene, Watertown, MA, USA), a lentiviral-based system

that combined both the CRISPR system and Cre recombinase. After NASH

diet feeding for 16 weeks, male L-KO mice were injected with pSECC-sgCtrl,

pSECC-sgAkt1, or pSECC-sgRhoA through the tail vein at 1×10^9 genome

copies per mouse, with two booster injections at 7 day intervals. The

pharmacological inhibition of AKT1 or RhoA was conducted using MK2206 or

Rhosin. After NASH diet feeding for 16 weeks, the mice were divided into 3

groups and intraperitoneally injected with vehicle solution (n = 6), MK2206

(AKT inhibitor, 50 mg/kg, every 3 days) (n = 5), or Rhosin (RhoA inhibitor, 20

mg/kg, every 3 days) (n = 6) for 8 weeks. All mice were sacrificed for further

study after NASH diet feeding for 24 weeks. Animal experiments were

approved by the Animal Experimentation Ethics Committees of Chongqing

Medical University and performed in accordance with the Guide for the Care

and Use of Laboratory Animals.

Liver tissues from patients with NASH

Paraffin-embedded normal (n = 10) and NASH human liver samples (n = 36)

were kindly provided by Dr. Jiangao Fan, Dr. Xiaojun Wang, and Dr. Yalan

Wang. Exclusion criteria were the presence of other causes of liver disease,

including alcoholic fatty liver disease (>30 g/day for men, >20 g/day for

women), chronic infection with hepatitis B and/or C virus, primary biliary

cirrhosis, haemochromatosis, auto-immune hepatitis, and Wilson's disease, as

well as the use of anti-obesity, glucose-lowering, and/or lipid lowering

pharmacological treatments. Liver tissue collection was approved by the

Institutional Ethics Committees of Chongqing Medical University and Xin Hua

Hospital. Informed consent was obtained from all subjects. The general

characteristics of the NASH human liver samples are listed in **Supplementary**

Table 1.

Cell culture and treatment

All cell lines were grown in Dulbecco's modified eagle medium (DMEM)

supplemented with 10% fetal bovine serum, 100 μg/mL of streptomycin, and

100 U/mL of penicillin at 37 °C in 5% CO_2. All cells were negative for

mycoplasma. Short tandem repeat tests were performed to ensure the

authenticity of the cells. Bovine serum albumin (BSA,10%), different

concentrations of palmitic acid (PA), 10 μM MK2206 (AKT inhibitor), 40 μM

Rhosin (RhoA inhibitor), or blocking antibody against PDGF-AA (2 μg/mL) was

added to the medium. For *in vitro* co-culture assays, the human hepatic

stellate cell (HSC) line LX-2 and human hepatocyte line MIHA cells (provided

by Dr Ben C.B. Ko, The Hong Kong Polytechnic University, Hongkong, China)

were used. LX-2 was pre-cultured in the lower chamber for 12 hours, and

PCK1-OE or PCK1-KO cells (with or without indicated treatment) were seeded

in Transwell inserts (#3401, Corning, NY, USA) that were subsequently loaded

into the LX2-containing wells. The cells and supernatants were harvested for

further analysis after 48 h of co-culture.

Isolation of primary mouse hepatocytes

Primary hepatocytes were isolated and cultured from the livers of WT and

L-KO mice fed the NASH diet as described previously [56]. Briefly, following

anesthesia, the inferior vena cava was cannulated and the liver was perfused

in situ with 40 mL pre-warmed EGTA solution and 40 mL solution containing

0.35 mg/mL pronase (no. P5147, Sigma-Aldrich, St. Louis, MO, USA), followed

by 40 mL solution containing 0.55 mg/mL collagenase (no. V900893,

Sigma-Aldrich). After perfusion, the liver was crushed, and hepatocytes were

released into the DMEM. The cell suspension was filtered through a 100 μm

cell strainer and centrifuged at 50 ×g at 25°C for 3 min. After washing three

times, the cells were suspended in DMEM supplemented with 10 mM glucose,

10% fetal bovine serum, 100 nM insulin (P3376, Beyotime Biotechnology,

Shanghai, China), and 100 nM dexamethasone (D8040, Solarbio Life

Sciences, Beijing, China), and then plated on 60 mm diameter plastic plates [56].

After cell attachment, the medium was replaced with serum-free media, and

the cells were used for experiments on the following day.

Isolation of primary mouse hepatocytes and primary HSCs

HSCs were isolated from the mice as described previously [57]. Briefly, after

perfusion with solutions containing protease and collagenase, the crushed liver

was digested with a solution containing 1% DNase (10104159001, Roche

Diagnostics GmbH, Mannheim, Germany), 0.5 mg/mL protease, and 0.55

mg/mL collagenase for 25 min. The cell suspension was filtered through a 70

µm cell strainer, centrifuged at 580 ×g for 10 min at 4 °C, and washed twice

with Gey's balanced salt solution. The cells were subjected to gradient

centrifugation on a 9.7% Nycodenz (1002424, Axis-Shield, Oslo, Norway) to

isolate HSCs, which were then plated onto collagen-coated plates. The cells

were cultured in DMEM containing 10% (vol/vol) fetal bovine serum and 1%

penicillin-streptomycin [58].

Construction of adenovirus, lentivirus, and stable cell lines

AdGFP and AdPCK1 adenoviruses were generated using the AdEasy system

as described previously [4]. The *PCK1* knockout (*PCK1*-KO) MIHA cell line was

constructed using the CRISPR-Cas9 system (from Prof. Ding Xue, the School

of Life Sciences, Tsinghua University, Beijing, China), as described previously

[4]. To knock down *ATF3* expression in MIHA cells, three pairs of

oligonucleotides encoding short hairpin RNAs (shRNAs) targeting *ATF3* or

negative control shRNA were cloned into the pLL3.7 vector (from Prof. Bing

Sun, Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of

Sciences, China). The lentiviruses were obtained by transient transfection of

the psPAX2 packaging plasmid and pMD2.G envelope plasmid in HEK293T

cells by using Lipofectamine 3000 Transfection Reagent (Thermo Fisher

Scientific, Waltham, MA, USA). Transfection efficiency was validated by

western blotting. Information on the reagents is listed in **Supplementary Table**

2. The pSECC lentiviral vector cloning and packaging strategy have been

described previously[26].

RNA extraction and real-time PCR

Total RNA was extracted from the liver tissues or cell lines using Trizol reagent

(15596018, Invitrogen, Carlsbad CA, USA) according to the manufacturer's

instructions. RNA was reverse-transcribed using a PrimeScript RT reagent Kit

with gDNA Eraser (Cat: RR047A, TAKARA, Shiga, Japan), and qPCR was

performed on a Bio-Rad CFX96 machine (Hercules, CA, USA). The mRNA

levels of selected genes were calculated after normalization to β-actin by using

the $2^{(-\Delta\Delta C(T))}$ method. Primer sequences are provided in **Supplementary Table**

3.

Immunoblotting

Mouse liver tissues and cells were homogenized and lysed in Protein

Extraction Reagent (P0013, Beyotime Biotechnology) supplemented with

protease inhibitor cocktail (04693159001, Roche, Basel, Switzerland). Equal

amounts of protein lysates were separated by sodium dodecyl

sulfate-polyacrylamide gel electrophoresis and transferred onto polyvinylidene

fluoride membranes (Millipore, Billerica, MA, USA). The antibodies are listed in

Supplementary Table 4. β-Actin was used as a loading control.

BODIPY staining

The cells were fixed in 4% formaldehyde for 30 min at room temperature,

permeabilized with 0.1% Triton X-100 for 5 min at room temperature, and

stained with BODYPY (D3823, Invitrogen) for 60 min. Stained sections were

analyzed using a Leica confocal microscope (Leica TCS SP8, Wetzlar,

Germany).

CCK8 assay

The cells were seeded at 5×10^3 cells /well in 96-well plates and treated with

different concentrations of PA for 24 h. 10 µL CCK-8 solution (#C0005,

Topscience, Shanghai, China) was added to each well and incubated at 37 °C

for 1 h. The absorbance was measured at 450 nm.

ELISA and G-LISA

Serum insulin (P3376, Beyotime Biotechnology), TNFα (PT512, Beyotime

Biotechnology), IL-6 (PI326, Beyotime Biotechnology), and PDGF-AA

(SEA523Mu, Cloud-Clone Corp., Wuhan, China) concentrations in the liver

tissue and plasma were quantified using mouse ELISA kits according to the

manufacturer's instructions. The levels of secreted PDGF-AA in the cell culture

supernatants were determined using a human ELISA kit (SEA523Hu,

Cloud-Clone Corp). Active RhoA was detected in a colorimetric RhoA

activation assay (G-LISA) (BK124, Cytoskeleton, Denver, CO, USA).

Immunofluorescence

LX-2 cells were fixed in 4% formaldehyde for 25 min and then incubated in 10%

normal goat serum for 1 h. The cells were incubated with primary α-SMA

antibodies. Specific signals were visualized using secondary antibodies

(ZF-0316, Zsbio, Beijing, China). For nuclear staining, the cells were treated

with 1 µg/mL DAPI (10236276001, Roche Diagnostics). The samples were

detected with a laser-scanning confocal microscope (Leica TCS SP8).

Chromatin immunoprecipitation (ChIP) assay

Chromatin immunoprecipitation (ChIP) and ChIP-quantitative real-time PCR

(ChIP-qPCR) assays for MIHA cells were performed as described previously [59].

Briefly, sonicated chromatin was used for the immunoprecipitation assay. The

pre-cleared supernatants were incubated with a monoclonal antibody against

ATF3 overnight at 4 °C, followed by a 4 h of incubation with protein A/G

agarose beads (LOT: 3460992, Millipore). The purified DNA fragments bound

by ATF3 were analyzed using qPCR. IgG and histone H3 were used as

negative and positive controls, respectively. The primers used for real-time

PCR analysis of PCK1 are listed in **Supplementary Table 3**.

Glucose and insulin tolerance tests

The glucose tolerance test (GTT) and insulin tolerance test (ITT) were

performed at 2 or 1 weeks prior to sacrifice. The animals were fasted for 16 or

6 h, and then glucose solution (2 g/kg body weight) or insulin (0.75 U/kg body

weight) was administered via an intraperitoneal injection, respectively. Venous

tail blood samples were collected at 0, 30, 60, 90, and 120 min

post-administration to assess blood glucose levels using a glucose meter

(HGM-114, OMRON, Kyoto, Japan).

Biochemical analysis

The serum levels of aspartate transaminase (AST), alanine transaminase

(ALT), total triglyceride (TG), total cholesterol (TC), and free fatty acids (FFA)

were determined with an automated biochemical analyzer (Hitachi 7600, Tokyo,

Japan). TG, TC, and FFA levels in the mouse liver tissues were measured with

commercial kits according to the manufacturer's protocol (TG: cat. no. BC0625;

TC: cat. no. BC1985; FFA: cat. no. BC0595, Solarbio Life Sciences).

Histological analysis

Paraffin blocks were sectioned into 4 µm slices and used for hematoxylin and

eosin (HE) staining, Sirius red staining, and immunohistochemistry (IHC)

assay according to standard protocols. Frozen liver tissue sections were

stained with Oil Red O (G1260, Solarbio). For pathological grading, all liver

specimens were scored by two experienced pathologists according to the

NAFLD activity score (NAS), defined as the sum of steatosis (0–3),

inflammation (0–3), and hepatocyte ballooning (0–2). An NAS score ≥5 was

considered to indicate NASH. Samples were scanned using a slide scanner

(Pannoramic DESK, 3D Histech kft, Hungary). For quantitative analysis, the

areas of lipid droplets and Sirius red staining were quantified using ImageJ

software (version 1.6.0; NIH, Bethesda, MD, USA). Immunohistochemical

staining was semi-quantitatively analyzed using the immunoreactive scoring

system [60]. The percentage of positive cells was graded on a scale of 0–4: (0:

negative, 1: 0–25%, 2: 26–50%, 3: 51–75%, 4: 76–100%). The signal intensity

was scored on a scale of 0–3: 0 = negative; 1 = weak; 2 = moderate; and 3 =

strong. Thus, the final immunoreactive score = (score of staining intensity) ×

(score of percentage of positive cells).

Transcriptomic analyses

Using Trizol reagent, total RNA was isolated from the liver tissues of WT and

L-KO mice fed a chow or NASH diet. The RNA quality was checked with a

Bioanalyzer 2200 (Agilent Technologies, Santa Clara, CA, USA). cDNA

libraries were prepared using an NEBNext[®] Ultra™ Directional RNA Library

Prep Kit, NEBNext[®] Poly (A) mRNA Magnetic Isolation Module, NEBNext[®]

Multiplex Oligos according to the manufacturer's instructions (New England

Biolabs, Ipswich, MA, USA). Genes with fold-change >2.0 or <0.5 and false

discovery rate <0.05 were considered to be significantly differentially

expressed. The volcano, heat, and bubble maps were generated using the

'ggplot2' or 'ggpubr' packages in R (version 3.6.3; The R Project for Statistical

Computing, Vienna, Austria). Gene set enrichment analysis was performed

using 'enrichplot' packages [61]. The RNA-seq data files have been deposited to

the Gene Expression Omnibus database (www.ncbi.nlm.nih.gov/geo/) under

accession number GSE162211.

Untargeted metabolomics

Following sacrifice, the livers were snap-frozen in liquid nitrogen and stored at

–80 °C until analysis. Untargeted metabolomics was performed using an

ultra-high performance liquid chromatography apparatus (Agilent 1290 Infinity

LC, Agilent Technologies) coupled to a quadrupole time-of-flight (TripleTOF

6600, AB SCIEX, Framingham, MA, USA) at Shanghai Applied Protein

Technology Co., Ltd. (Shanghai, China). Metabolites with a variable

importance in projection value >1 were evaluated using Student's t-test. $p <$

0.05 was considered to indicate statistically significant results.

HPLC analysis of cellular nucleotides

Cellular nucleotides were extracted according to published procedures [62].

Briefly, 1×10^6 cells were washed with phosphate-buffered saline and

quenched with liquid nitrogen, and then vigorously mixed with methanol and

acetonitrile (1:1, v:v). After incubation on ice for 15 min, the samples were

centrifuged at 12,000 ×g at 4 °C for 10 min. Cellular nucleotides were

separated and quantified using a C18 column (Agilent Eclipse XDB-C18, 4.6 ×

250 mm, average particle size 5 μm) assembled on the Waters Alliance e2695

Separations Module (Milford, MA, USA). Acetonitrile (5%) and 50 mM KH_2PO_4

(pH 6.5) containing 10 mM tetrabutylammonium bromide were used as mobile

phase A, and acetonitrile was used as mobile phase B. All samples were

separated in the mobile phase at a flow rate of 1 mL/min for 30 min at 22 °C.

No degradation of individual nucleotides or changes in the ratios of nucleotide

mixtures was detected during the experiments. The GTP concentration in the

samples was calculated based on the slope of the calibration curves generated

using pooled authentic samples (to mimic the matrix), and

guanosine-5′-triphosphoric acid disodium salt was used as a standard.

GEO database mining

Raw data in the GSE126848, GSE89632, and GSE135251 datasets were

downloaded from the GEO database (https://www.ncbi.nlm.nih.gov/geo/). R

package DESeq2 was used to analyze gene expression levels between

different samples, such as NASH vs. healthy, NAFLD vs. healthy, and obesity

vs. healthy. Genes showing |$\log_2$ fold-change|>1 and false discovery rate

<0.05 were considered to present differential expression.

Statistics

Statistical analyses were performed using GraphPad Prism 6.0 software

(GraphPad, Inc., La Jolla, CA, USA). Data were represented as the mean ±

standard error of the mean unless otherwise stated. Significant differences

between the means of two groups were determined using Student's t-test.

One-way analysis of variance was used to determine statistical significance for

experiments with more than two groups followed by Tukey's post hoc test. For

correlation analysis, Spearman's correlation coefficient was used. $p < 0.05$

was considered to indicate statistically significant results.

Acknowledgements

We would like to thank Dr. T.-C He (University of Chicago, USA) and Prof. Ding Xue (Tsinghua University, China) for providing the pAdEasy and CRISPR/Cas9 system, respectively. We thank Prof. Youde Cao and Yalan Wang (Chongqing Medical University, China) for providing samples and pathological analysis support. This work was supported by the National Natural Science Foundation of China (grant no. U20A20392, 82073251, 82072286, 81872270), the 111 Project (No. D20028), the Natural Science Foundation Project of Chongqing (cstc2018jcyjAX0254, cstc2019jscx-dxwtBX0019), the Leading Talent Program of CQ CSTC (CSTCCXLJRC201719), the Science and Technology Research Program of Chongqing Municipal Education Commission (HZ2021006, KJZD-M202000401), the Future Medical Youth Innovation Team of Chongqing Medical University (W0036, W0101), the Kuanren talents program of the second affiliated hospital of Chongqing Medical University, the Scientific Research Innovation Project for Postgraduate in Chongqing (CYB19168, CYS19193), and the Youth talent fund of Sichuan Provincial People's Hospital in 2020 (No. 2020QN03).

Author contributions

NT, AH, and KW conceived and designed the study. QY, YL and GZ performed most experiments and analyzed the data. HD, CC and KW conducted bioinformatics analysis. XP assisted with mice experiments. XW, JF,

787 and QP provided human NAFLD/NASH samples. QY, KW, and NT wrote the

788 manuscript with all authors providing feedback. The order of the co-first

789 authors was assigned on the basis of their relative contributions to the study.

791 **Conflict of interest:**

792 The authors declare no competing financial interests.

794 **Ethical approval**

795 Animal experiments were approved by Animal Experimentation Ethics

796 Committees of Chongqing Medical University and were carried out in

797 accordance with the Guide for the Care and Use of Laboratory Animals. Liver

798 tissue collection was approved by the Institutional Ethics Committees of

799 Chongqing Medical University and Xin Hua Hospital. Informed consent was

800 obtained from all subjects.

802 **Data availability**

803 RNA-seq data files have been deposited into the Gene Expression Omnibus

804 database (www.ncbi.nlm.nih.gov/geo/) with accession number GSE162211.

805 All other data are available from the corresponding author upon reasonable

806 request. Source data are provided with this paper.

References

1   Younossi, Z. *et al.* Global burden of NAFLD and NASH: trends, predictions, risk factors and
prevention. *Nat Rev Gastroenterol Hepatol* **15**, 11-20 (2018).

2   Schuster, S., Cabrera, D., Arrese, M. & Feldstein, A. E. Triggering and resolution of
inflammation in NASH. *Nat Rev Gastroenterol Hepatol* **15**, 349-364 (2018).

3   Arab, J. P., Arrese, M. & Trauner, M. Recent Insights into the Pathogenesis of Nonalcoholic
Fatty Liver Disease. *Annu Rev Pathol* **13**, 321-350 (2018).

4   Tuo, L. *et al.* PCK1 negatively regulates cell cycle progression and hepatoma cell proliferation
via the AMPK/p27(Kip1) axis. *J Exp Clin Cancer Res* **38**, 50 (2019).

5   Xiang, J. *et al.* Gluconeogenic enzyme PCK1 deficiency promotes CHK2 O-GlcNAcylation and
hepatocellular carcinoma growth upon glucose deprivation. *J Clin Invest* **131** (2021).

6   Xu, D. *et al.* The gluconeogenic enzyme PCK1 phosphorylates INSIG1/2 for lipogenesis. *Nature*
580, 530-535 (2020).

7   Santra, S. *et al.* Cytosolic phosphoenolpyruvate carboxykinase deficiency presenting with
acute liver failure following gastroenteritis. *Mol Genet Metab* **118**, 21-27 (2016).

8   She, P. *et al.* Phosphoenolpyruvate carboxykinase is necessary for the integration of hepatic
energy metabolism. *Mol Cell Biol* **20**, 6508-6517 (2000).

9   Millward, C. A. *et al.* Phosphoenolpyruvate carboxykinase (Pck1) helps regulate the
triglyceride/fatty acid cycle and development of insulin resistance in mice. *J Lipid Res* **51**,
1452-1463 (2010).

10   Hoxhaj, G. & Manning, B. D. The PI3K-AKT network at the interface of oncogenic signalling
and cancer metabolism. *Nat Rev Cancer* **20**, 74-88 (2020).

11   Porstmann, T. *et al.* PKB/Akt induces transcription of enzymes involved in cholesterol and
fatty acid biosynthesis via activation of SREBP. *Oncogene* **24**, 6465-6481 (2005).

12   Huang, X., Liu, G., Guo, J. & Su, Z. The PI3K/AKT pathway in obesity and type 2 diabetes. *Int J*
Biol Sci **14**, 1483-1496 (2018).

13   Chen, J. *et al.* HIF-2α upregulation mediated by hypoxia promotes NAFLD-HCC progression by
activating lipid synthesis via the PI3K-AKT-mTOR pathway. *Aging (Albany NY)* **11**,
10839-10860 (2019).

14   Chi, Y., Gong, Z., Xin, H., Wang, Z. & Liu, Z. Long noncoding RNA lncARSR promotes
nonalcoholic fatty liver disease and hepatocellular carcinoma by promoting YAP1 and
activating the IRS2/AKT pathway. *J Transl Med* **18**, 126 (2020).

15   Suppli, M. P. *et al.* Hepatic transcriptome signatures in patients with varying degrees of
nonalcoholic fatty liver disease compared with healthy normal-weight individuals. *Am J*
Physiol Gastrointest Liver Physiol **316**, G462-g472 (2019).

16   Fang, J. *et al.* Hepatic IRF2BP2 Mitigates Nonalcoholic Fatty Liver Disease by Directly
Repressing the Transcription of ATF3. *Hepatology* **71**, 1592-1608 (2020).

17   Allen-Jennings, A. E., Hartman, M. G., Kociba, G. J. & Hai, T. The roles of ATF3 in liver
dysfunction and the regulation of phosphoenolpyruvate carboxykinase gene expression. *J Biol*
Chem **277**, 20020-20025 (2002).

18   Asgharpour, A. *et al.* A diet-induced animal model of non-alcoholic fatty liver disease and
hepatocellular cancer. *J Hepatol* **65**, 579-588 (2016).

19   Liu, X. J. *et al.* Characterization of a murine nonalcoholic steatohepatitis model induced by

851		high fat high calorie diet plus fructose and glucose in drinking water. *Lab Invest* **98**, 1184-1199
852		(2018).
853	20	Kucukoglu, O., Sowa, J. P., Mazzolini, G. D., Syn, W. K. & Canbay, A. Hepatokines and
854		adipokines in NASH-related hepatocellular carcinoma. *J Hepatol* **74**, 442-457 (2021).
855	21	Thieringer, F. *et al.* Spontaneous hepatic fibrosis in transgenic mice overexpressing PDGF-A.
856		*Gene* **423**, 23-28 (2008).
857	22	Higuchi, M., Masuyama, N., Fukui, Y., Suzuki, A. & Gotoh, Y. Akt mediates
858		Rac/Cdc42-regulated cell motility in growth factor-stimulated cells and in invasive PTEN
859		knockout cells. *Curr Biol* **11**, 1958-1962 (2001).
860	23	Dou, C. *et al.* P300 Acetyltransferase Mediates Stiffness-Induced Activation of Hepatic Stellate
861		Cells Into Tumor-Promoting Myofibroblasts. *Gastroenterology* **154**, 2209-2221.e2214 (2018).
862	24	Calvayrac, O. *et al.* The RAS-related GTPase RHOB confers resistance to EGFR-tyrosine kinase
863		inhibitors in non-small-cell lung cancer via an AKT-dependent mechanism. *EMBO Mol Med* **9**,
864		238-250 (2017).
865	25	Hodge, R. G. & Ridley, A. J. Regulating Rho GTPases and their regulators. *Nat Rev Mol Cell Biol*
866		**17**, 496-510 (2016).
867	26	Sánchez-Rivera, F. J. *et al.* Rapid modelling of cooperating genetic events in cancer through
868		somatic genome editing. *Nature* **516**, 428-431 (2014).
869	27	Snaebjornsson, M. T., Janaki-Raman, S. & Schulze, A. Greasing the Wheels of the Cancer
870		Machine: The Role of Lipid Metabolism in Cancer. *Cell Metab* **31**, 62-76 (2020).
871	28	Vily-Petit, J. *et al.* Intestinal gluconeogenesis prevents obesity-linked liver steatosis and
872		non-alcoholic fatty liver disease. *Gut* **69**, 2193-2202 (2020).
873	29	Yoon, J. C. *et al.* Control of hepatic gluconeogenesis through the transcriptional coactivator
874		PGC-1. *Nature* **413**, 131-138 (2001).
875	30	Friedman, S. L., Neuschwander-Tetri, B. A., Rinella, M. & Sanyal, A. J. Mechanisms of NAFLD
876		development and therapeutic strategies. *Nat Med* **24**, 908-922 (2018).
877	31	Samuel, V. T. *et al.* Fasting hyperglycemia is not associated with increased expression of
878		PEPCK or G6Pc in patients with Type 2 Diabetes. *Proc Natl Acad Sci U S A* **106**, 12121-12126
879		(2009).
880	32	Imai, E. *et al.* Characterization of a complex glucocorticoid response unit in the
881		phosphoenolpyruvate carboxykinase gene. *Mol Cell Biol* **10**, 4712-4719 (1990).
882	33	Lundsgaard, A. M. *et al.* Mechanisms Preserving Insulin Action during High Dietary Fat Intake.
883		*Cell Metab* **29**, 50-63.e54 (2019).
884	34	Guo, X. *et al.* The Role of Palmitoleic Acid in Regulating Hepatic Gluconeogenesis through
885		SIRT3 in Obese Mice. *Nutrients* **14** (2022).
886	35	Wang, Z. & Dong, C. Gluconeogenesis in Cancer: Function and Regulation of PEPCK, FBPase,
887		and G6Pase. *Trends Cancer* **5**, 30-45 (2019).
888	36	Jiang, W. *et al.* Acetylation regulates gluconeogenesis by promoting PEPCK1 degradation via
889		recruiting the UBR5 ubiquitin ligase. *Mol Cell* **43**, 33-44 (2011).
890	37	Hai, T., Wolfgang, C. D., Marsee, D. K., Allen, A. E. & Sivaprasad, U. ATF3 and stress responses.
891		*Gene Expr* **7**, 321-335 (1999).
892	38	Tsai, W. W. *et al.* ATF3 mediates inhibitory effects of ethanol on hepatic gluconeogenesis. *Proc*
893		*Natl Acad Sci U S A* **112**, 2699-2704 (2015).
894	39	Tu, C. *et al.* Cardiolipin Synthase 1 Ameliorates NASH Through Activating Transcription Factor

3 Transcriptional Inactivation. *Hepatology* **72**, 1949-1967 (2020).

40   Gut, P. *et al.* Whole-organism screening for gluconeogenesis identifies activators of fasting
metabolism. *Nat Chem Biol* **9**, 97-104 (2013).

41   Hakimi, P. *et al.* Phosphoenolpyruvate carboxykinase and the critical role of cataplerosis in
the control of hepatic metabolism. *Nutr Metab (Lond)* **2**, 33 (2005).

42   Satapati, S. *et al.* Mitochondrial metabolism mediates oxidative stress and inflammation in
fatty liver. *J Clin Invest* **125**, 4447-4462 (2015).

43   Tsuchida, T. & Friedman, S. L. Mechanisms of hepatic stellate cell activation. *Nat Rev*
Gastroenterol Hepatol **14**, 397-411 (2017).

44   Zhang, B., Zhang, Y., Shacter, E. & Zheng, Y. Mechanism of the guanine nucleotide exchange
reaction of Ras GTPase--evidence for a GTP/GDP displacement model. *Biochemistry* **44**,
2566-2576 (2005).

45   Wawrzyniak, J. A. *et al.* A purine nucleotide biosynthesis enzyme guanosine monophosphate
reductase is a suppressor of melanoma invasion. *Cell Rep* **5**, 493-507 (2013).

46   Hallett, M. A., Dagher, P. C. & Atkinson, S. J. Rho GTPases show differential sensitivity to
nucleotide triphosphate depletion in a model of ischemic cell injury. *Am J Physiol Cell Physiol*
285, C129-138 (2003).

47   Mondin, M. *et al.* Alterations in cytoskeletal protein expression by mycophenolic acid in
human mesangial cells requires Rac inactivation. *Biochem Pharmacol* **73**, 1491-1498 (2007).

48   Bianchi-Smiraglia, A. *et al.* Microphthalmia-associated transcription factor suppresses
invasion by reducing intracellular GTP pools. *Oncogene* **36**, 84-96 (2017).

49   Han, M. H. *et al.* Flavonoids Isolated from Flowers of Lonicera japonica Thunb. Inhibit
Inflammatory Responses in BV2 Microglial Cells by Suppressing TNF-α and IL-β Through
PI3K/Akt/NF-kb Signaling Pathways. *Phytother Res* **30**, 1824-1832 (2016).

50   Zhao, Y. *et al.* PI3K Positively Regulates YAP and TAZ in Mammary Tumorigenesis Through
Multiple Signaling Pathways. *Mol Cancer Res* **16**, 1046-1058 (2018).

51   Jeong, S. H. & Lim, D. S. Insulin receptor substrate 2: a bridge between Hippo and AKT
pathways. *BMB Rep* **51**, 209-210 (2018).

52   Gutierrez, A. & Look, A. T. NOTCH and PI3K-AKT pathways intertwined. *Cancer Cell* **12**,
411-413 (2007).

53   Chien, A. J. *et al.* MK-2206 and Standard Neoadjuvant Chemotherapy Improves Response in
Patients With Human Epidermal Growth Factor Receptor 2-Positive and/or Hormone
Receptor-Negative Breast Cancers in the I-SPY 2 Trial. *J Clin Oncol* **38**, 1059-1069 (2020).

54   Schmid, P. *et al.* Capivasertib Plus Paclitaxel Versus Placebo Plus Paclitaxel As First-Line
Therapy for Metastatic Triple-Negative Breast Cancer: The PAKT Trial. *J Clin Oncol* **38**, 423-433
(2020).

55   Yoon, C. *et al.* Chemotherapy Resistance in Diffuse-Type Gastric Adenocarcinoma Is Mediated
by RhoA Activation in Cancer Stem-Like Cells. *Clin Cancer Res* **22**, 971-983 (2016).

56   Hernández-Alvarez, M. I. *et al.* Deficient Endoplasmic Reticulum-Mitochondrial
Phosphatidylserine Transfer Causes Liver Disease. *Cell* **177**, 881-895.e817 (2019).

57   Mederacke, I., Dapito, D. H., Affò, S., Uchinami, H. & Schwabe, R. F. High-yield and high-purity
isolation of hepatic stellate cells from normal and fibrotic mouse livers. *Nat Protoc* **10**,
305-315 (2015).

58   Choi, W. M. *et al.* Glutamate Signaling in Hepatic Stellate Cells Drives Alcoholic Steatosis. *Cell*

Metab **30**, 877-889.e877 (2019).

59   Yang, F. *et al.* GSTZ1-1 Deficiency Activates NRF2/IGF1R Axis in HCC via Accumulation of
Oncometabolite Succinylacetone. *Embo j* **38**, e101964 (2019).

60   Yao, F. *et al.* A targetable LIFR-NF-κB-LCN2 axis controls liver tumorigenesis and vulnerability
to ferroptosis. *Nat Commun* **12**, 7333 (2021).

61   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for comparing biological
themes among gene clusters. *Omics* **16**, 284-287 (2012).

62   Hannan, J. P. *et al.* HPLC method to resolve, identify and quantify guanine nucleotides bound
to recombinant ras GTPase. *Anal Biochem* **631**, 114338 (2021).

Figures and Figure legends

Fig. 1 PCK1 is downregulated in patients with NASH and mouse models

of NASH. a Genes downregulated in patients with obesity (n = 12), NAFLD (n

= 15), and NASH (n = 16) from GSE126848 dataset. **b** Relative *PCK1* mRNA

levels in GSE126848 and GSE89632 datasets. **c** PCK1 expression in normal

individuals and patients with NASH and semi-quantitative analyses of

immunohistochemistry (IHC) data (healthy = 10, NASH = 36). Scale bars: 50

μm. **d, e** mRNA (**d**) and protein (**e**) levels of PCK1 in the livers of WT mice fed

with chow or NASH diet (n = 6). **F, g** PCK1 mRNA (**f**) and protein (**g**) levels in

MIHA cells treated with BSA or PA-BSA. **h** Relative levels of indicated genes in

MIHA cells treated with 0.2 mM PA. **i, j** Representative ATF3 expression in

normal individuals and patients with NASH (**i**) and semi-quantitative analyses

of IHC data (**j**) (healthy = 10, NASH = 36). Scale bars: 50 μm. **k** Chromatin

immunoprecipitation assays were performed in MIHA cells with or without PA

treatment using an antibody against ATF3, IgG, or H3. **l** Protein levels of PCK1

in MIHA cells infected with either shControl or shATF3 treated with 0.2 mM PA.

m Correlation analysis of *ATF3* mRNA level with *PCK1* in human

NAFLD/NASH liver samples (GSE135251, n = 206). Data are expressed as

the mean ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. *p* values obtained via

two-tailed unpaired Student's *t* tests, one-way analysis of variance with

Tukey's post hoc test, or non-parametric Spearman's test.

Figure 2

Fig. 2 L-KO mice fed chow diet exhibited a distinct hepatic steatosis

phenotype. a Schematic diagram of the mouse model fed the chow diet, n =

10/group. **b** PCK1 protein expression in WT and L-KO mouse heart, liver,

spleen, cecum and kidney were confirmed by immunoblotting. **c** Body weight,

glucose tolerance test (GTT), and insulin tolerance test (ITT) were measured

in WT and L-KO mice. **d** Representative liver image and liver weight of WT and

L-KO mice. **e–g** Determination of alanine aminotransferase (ALT), aspartate

aminotransferase (AST), total triglycerides (TG), total cholesterol (TC), and

free fatty acid (FFA) levels in the serum or liver tissues. **h** Paraffin-embedded

liver sections were stained with hematoxylin and eosin (H&E), Sirius red, and

F4/80. Frozen sections stained with Oil Red O. Scale bars: 50 μm. **i**

Quantification of liver sections of WT and L-KO mice fed the chow diet. **j**

Levels of TNF-α and IL-6 in the liver tissues. **k** Quantitative PCR analysis of

liver mRNA expression. Data are expressed as the mean ± SEM; *$p < 0.05$, **p

< 0.01, ***$p < 0.001$; n.s., not significant. p values obtained via two-tailed

unpaired Student's *t* tests.

Fig. 3 PCK1 ablation accelerates inflammation and fibrogenesis in NASH model. a Schematic diagram of mouse model fed the NASH diet. **b** Body weight was measured in WT and L-KO mice. **c** Representative gross liver morphology, whole body photo, and liver weight. **d** Serum levels of insulin, alanine aminotransferase (ALT), and aspartate aminotransferase (AST) were

measured. **e** Paraffin-embedded liver sections were stained with hematoxylin

and eosin (H&E), Sirius red, α-SMA immunostaining, and F4/80

immunostaining. Frozen sections stained with Oil Red O. Scale bars: 50 µm. **f**

Normalized activity scores (NAS) for each group. **g** Levels of TNF-α and IL-6 in

the liver tissues detected using enzyme-linked immunosorbent assay (ELISA).

h Schematic showing the administration protocol for AAV8-TBG-*control* or

AAV8-TBG-*Pck1* in WT and L-KO mice for experiments shown in **i**–**l**, n =

6/group. **i** Representative whole-body photo and gross liver morphology. **j**

Liver weight and body weight. **k** Paraffin-embedded liver sections were stained

with H&E, Sirius red immunostaining and Oil Red O staining. Scale bars: 50

µm. **l** Quantifications of Sirius red staining, Oil Red O staining, and NAS. Data

are expressed as the mean ± SEM; *p < 0.05, **p < 0.01, ***p < 0.001. *p*

values obtained via two-tailed unpaired Student's *t* tests.

1011
1012

Figure 4

Fig. 4 Loss of PCK1 promotes lipid accumulation according to transcriptome and metabolome analyses. RNA sequencing was performed on the livers of WT (n = 4) and L-KO (n = 5) mice fed the NASH diet. **a** Gene Ontology analysis of all significantly changed genes in top 10 biological

processes. **b** Volcano plot representation of significantly up- and

downregulated genes. **c** Gene Set Enrichment Analysis plot (left) of

enrichment in "PPAR signaling pathway" signature; heatmap (right) of

significantly upregulated PPAR target genes. **d, e** Quantitative PCR (**d**) and

immunoblot (**e**) analysis of indicated genes or protein expression in mouse

liver tissues. **f** Relative mRNA expression of key genes in G3P pathway (n = 8).

g Upregulated metabolites detected by untargeted metabolomics (n = 6). **h**

Relative level of G3P and PA in mouse liver tissues (n = 6). Data are

expressed as the mean ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. p values

obtained via two-tailed unpaired Student's t tests.

1028

1029

Figure 5

1030

Fig. 5 Hepatic PCK1 deficiency leads to HSC activation via PI3K/AKT

pathway. a Pathway enrichment analysis of significantly upregulated genes in

L-KO mice. **b** Gene Set Enrichment Analysis (GSEA) plot of enrichment in

PI3K/AKT pathway. **c–f** Immunoblot analysis of AKT and p-AKT (S473 or T308)

in mouse liver tissues (**c**), primary hepatocytes from NASH feeding mice(**d**),

PCK1-OE (**e**), and *PCK1*-KO (**f**) MIHA cells with or without 0.2 mM palmitic

acid (PA) treatment. **g** Schematic flow chart of co-culture models. **h, i**

Quantitative PCR analysis of fibrosis-related genes in LX-2 cells co-cultured

with *PCK1*-KO (**h**) or *PCK1*-OE (**i**) MIHA cells. **j, k** Western blotting of fibrosis

related protein in liver tissues (**j**) or primary HSCs from NASH feeding mice (**k**)

(n = 3) **l, m** Relative mRNA expression (**l**) and immunofluorescence images (**m**)

of *ACTA2/α*-SMA, *COL1A1*, and *COL3A1* in LX-2 cells co-cultured with

PCK1-KO MIHA cells treated with AKT inhibitor MK2206 (10 μM). Scale bars:

25 μm. Data are expressed as the mean ± SEM; $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p <$

0.001. *p* values obtained via two-tailed unpaired Student's *t* tests or one-way

ANOVA with Tukey's post hoc test.

Figure 6

1047

Fig. 6 Paracrine PDGF-AA from hepatocytes promotes hepatic stellate cell (HSC) activation. a Expression levels of genes related to fibrogenesis. **b** Relative *PDGFA* mRNA levels in GSE126848 and GSE89632 datasets. **c, d** PDGF-AA protein levels in primary hepatocytes (**c**) or liver tissues (**d**) detected by western blotting. **e, f** PDGF-AA levels in liver tissues (**e**) or serum (**f**) were detected using enzyme-linked immunosorbent assay (ELISA). **g, h** Secreted PDGF-AA levels in conditioned medium with *PCK1*-KO (**g**) or *PCK1*-OE (**h**)

MIHA cells treated with 0.2 mM palmitic acid (PA). **i**, **j** mRNA levels of *PDGFRA*

in cell lysate of LX-2 cells co-cultured with *PCK1*-KO (**i**) or *PCK1*-OE (**j**) MIHA

cells treated with PA. **k** Protein level in LX-2 cells co-cultured with *PCK1*-KO

MIHA cells containing nonspecific rabbit IgG or a PDGF-AA blocking antibody.

l Immunohistochemistry (IHC) analysis of PCK1, p-AKT (S473), and PDGF-AA

in mouse liver sections (from serial sections). Scale bars: 50 µm. **m, n** Levels

of PDGF-AA (**m**) or *PDGFA* (**n**) in conditioned medium or cell lysate of

PCK1-KO MIHA cells treated with AKT inhibitor MK2206 (10 µM). **o** Protein

levels in *PCK1*-KO MIHA cells treated with AKT inhibitor MK2206 (10 µM).

Data are expressed as the mean ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. *p*

values obtained via two-tailed unpaired Student's *t* tests or one-way ANOVA

with Tukey's post hoc test.

1067

1068

Figure 7

1069

Fig. 7 PCK1 deficiency promotes activation of PI3K/AKT/PDGF-AA axis

by activating RhoA in hepatocytes. a Immunoblotting analysis of indicated

proteins in mouse liver tissues. **b** Immunohistochemistry (IHC) analysis of

p-RhoA (S188) in mouse liver tissues. Scale bars: 50 μm. **c–e** Relative levels

of active RhoA were measured using G-LISA colorimetric RhoA activation

assay in mouse liver tissues (**c**), *PCK1*-OE (**d**) and *PCK1*-KO (**e**) MIHA cells

treated with 0.2 mM palmitic acid (PA). **f, g** Immunoblots of p-RhoA (S188) and

RhoA in *PCK1*-OE (**f**) and *PCK1*-KO (**g**) MIHA cells with or without 0.2 mM PA

treatment. **h** Expression of indicated proteins in *PCK1*-KO MIHA cells after

addition of Rhosin (30 µM). **i** Levels of PDGF-AA in the supernatant of

PCK1-KO MIHA cells treated with Rhosin (30 µM). **j** Relative mRNA

expression of *ACTA2*, *COL1A1*, and *COL3A1* in LX-2 cells co-cultured with

PCK1-KO MIHA cells treated with Rhosin (30 µM). **k** IHC analysis of PCK1,

p-RhoA (S188), p-AKT (S473), and PDGF-AA in normal individuals and

patients with NASH (from serial sections). Scale bars: 50 µm. **l**

Semi-quantitative analyses of immunohistochemistry data of healthy and

NASH human tissues for indicated proteins (healthy = 10, NASH = 10). Data

are expressed as the mean ± SEM; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. *p*

values obtained via one-way ANOVA with Tukey's post hoc test.

1089

1090

Figure 8

1091

Fig. 8 Pharmacological inhibition or genetic silencing of AKT1 or RhoA

prevents NASH development *in vivo.* L-KO mice were fed the NASH diet for

24 weeks, and therapeutic treatments were initiated at different times. **a**

Schematic diagram of *in vivo* pharmacological inhibition (**b**–**e**) and pSECC

lentivirus-mediated silencing (**f**–**i**) of AKT1 or RhoA. **b, c** Representative

whole-body, gross liver morphology, liver weight (**b**), and serum alanine

aminotransferase (ALT) and aspartate aminotransferase (AST) (**c**). **d**

Paraffin-embedded liver sections were stained with hematoxylin and eosin, or

immunostained for F4/80, COL3A1, and α-SMA. Frozen sections stained with

Oil Red O. Scale bars: 50 μm. **e** Levels of TNF-α and IL-6 in liver tissues. **f**

Representative gross images of the livers from different groups of mice. **g**

Quantification of body weight and liver weight in pSECC-sgAkt1 and

pSECC-sgRhoA L-KO mice. **h** Plasma levels of total triglycerides (TG), total

cholesterol (TC), and free fatty acids (FFA) were measured. **i** Normalized

activity scores of liver sections. **j** Model depicting the critical role of PCK1 in

controlling NASH progression. Data are expressed as the mean ± SEM; *$p <$

0.05, **$p < 0.01$, ***$p < 0.001$; n.s., not significant. p values obtained via

one-way ANOVA with Tukey's post hoc test.

1110