The SLIM1 transcription factor is required for arsenic resistance in *Arabidopsis*

thaliana

Timothy O. Jobe[1], Qi Yu[2,3], Felix Hauser[2], Qingqing Xie[2,4], Yuan Meng[2], Tim

Maassen[1], Stanislav Kopriva[5], Julian I. Schroeder[2]

[1] Biocenter, Botanical Institute, University of Cologne, Cologne, 50674 Cologne,

Germany

[2] Division of Biological Sciences, Cell and Developmental Biology Section, University

of California, San Diego, La Jolla, California 92093-0116, USA

[3] School of Life Sciences, and Hubei Key Laboratory of Genetic Regulation and

Integrative Biology, Central China Normal University, Wuhan 430079, Hubei, P. R.

China

[4] Plant Molecular and Cellular Biology Laboratory, Salk Institute for Biological

Studies, La Jolla, CA 92037, USA

[5] Institute for Plant Sciences and Cluster of Excellence on Plant Sciences (CEPLAS),

University of Cologne, 50674 Cologne, Germany

E-mail addresses: Timothy O. Jobe (tjobe@uni-koeln.de), Qi Yu (yuq084@ucsd.edu), Felix Hauser

(felix.hauser@outlook.com), Qingqing Xie (qxie@salk.edu), Yuan Meng

(mengyuan1991@outlook.com), Tim Maassen (tim.hps.maassen@gmail.com), Stanislav Kopriva

(skopriva@uni-koeln.de), Julian I. Schroeder (jischroeder@ucsd.edu)

Corresponding authors :

Julian I. Schroeder, jischroeder@ucsd.edu, Phone 858 534 7759

and

Timothy O. Jobe, tjobe@uni-koeln.de, Phone +49 (0)221 470-8294

Date of submission: 18 April 2021

Number of figures: 4

Total word count: 5,148

Supplementary Data: 2 Figures, 14 Tables

Running Title: SLIM1 regulates arsenic sensitivity in *Arabidopsis thaliana*

Abstract

The transcriptional regulators of arsenic-induced gene expression remain largely

unknown. Sulfur assimilation is tightly linked with arsenic detoxification. Here we

report that mutant alleles in the SLIM1 transcription factor are substantially more

sensitive to arsenic than cadmium. Arsenic treatment caused high levels of oxidative

stress in the *slim1* mutants, and *slim1* alleles were impaired in both thiol and sulfate

accumulation. We further found enhanced arsenic accumulation in roots of *slim1*

mutants. Transcriptome analyses indicate an important role for SLIM1 in arsenic-

induced tolerance mechanisms. The present study identifies the SLIM1 transcription

factor as an essential component in arsenic tolerance and arsenic-induced gene

expression. Our results suggest that the severe arsenic sensitivity of the *slim1* mutants is

caused by altered redox status.

Keywords

arsenic, sulfur limitation, transcription factor, *Arabidopsis thaliana*

Introduction

Many advanced technologies used by modern society rely on heavy metals and arsenic. These elements are toxic and pose a significant risk to the environment and human health if consumed. However, unlike animals, plants are often partially tolerant to heavy metals and arsenic and can accumulate large amounts in diverse tissues [1]. Arsenic is a highly toxic substance commonly found in anthropogenic wastes (electronics and fertilizers) and can also be found at high levels in certain rocks, soils, and waters globally [2–5]. While this toxic metalloid has no recognized role in plant or animal nutrition, plant-based products are the main entry point for arsenic into the food chain [6]. Thus, understanding the molecular mechanisms underlying plant uptake, transport, detoxification, and accumulation of arsenic is vital for enhancing the nutritional value and safety of our food.

We previously described the development of a plant genetic reporter line that fused the promoter of a cadmium and arsenic-inducible high-affinity sulfate transporter to firefly luciferase (*pSULTR1;2::LUC*) to identify mutants in signaling [7]. A major goal of this work was to identify the transcriptional regulators mediating rapid arsenic-induced gene expression in Arabidopsis. This approach was successful in identifying new alleles of the glutathione biosynthesis genes gamma-glutamylcysteine synthetase (γ-ECS) and glutathione synthetase (GS), as being required for cadmium and arsenic-induced gene expression [7]. Glutathione is necessary for the synthesis of phytochelatins, which detoxify many toxic compounds, including cadmium and arsenic, by chelation and sequestration in the vacuole[1,8–11]. Phytochelatins are short polymers of glutathione

synthesized in the cytosol in response to toxic metal(loid)s. Thus, arsenic exposure can

rapidly deplete glutathione levels, creating a high demand for glutathione in plant cells.

Because the tripeptide glutathione (Glu-Cys-Gly) contains the sulfur-containing amino

acid cysteine, the sulfate assimilation pathway is inextricably linked to glutathione

biosynthesis. Sulfate assimilation takes oxidized sulfur in the form of sulfate and,

through a series of energy-dependent reducing steps, produces sulfide. Due to the

toxicity of sulfide, this intermediate quickly reacts with O-acetylserine to produce the

amino acid cysteine[12]. Thus, unlike animals, plants do not require exogenous sulfur-

containing amino acids and proteins for survival[13]. More importantly, this creates a

direct link between the sulfate assimilation pathway and the ability of plants to detoxify

arsenic.

While our luciferase genetic reporter approach has not identified transcriptional

regulators of arsenic-induced gene induction to date, a similar reporter gene approach

successfully identified a transcriptional regulator of the sulfur deficiency response in

Arabidopsis. This genetic screen used the same high-affinity sulfate transporter

promoter element fused to the green fluorescent protein (*pSULTR1;2::GFP*) and

identified four allelic mutants in an ethylene insensitive-like transcription factor called

Sulfur Limitation 1 (SLIM1) that failed to induce the reporter construct under sulfur

limiting conditions [14]. All of the allelic *slim1* mutants identified in this screen resulted

in missense mutations altering single amino acid residues [14]. In *slim1-1* and *slim1-2*,

high-affinity sulfate uptake was decreased by ~60%, and sulfur-dependent microarray

analyses on *slim1-1* and *slim1-2* showed a decrease in the induction of many sulfur

limitation-induced transcripts compared to controls suggesting that SLIM1 is a positive

regulator of sulfate uptake and assimilation [14].

While the transcription factors that control arsenic-induced gene expression remain

largely unknown, arsenic exposure is known to rapidly deplete cellular glutathione

levels, increasing the demand for reduced sulfur compounds from the sulfur assimilation

pathway[7,15,16]. A similar situation occurs under sulfur deficiency. As sulfate supply

decreases, cellular levels of cysteine and glutathione become depleted. Thus, because of

the similarities in glutathione depletion and subsequent upregulation of the high-affinity

sulfate transporter *SULTR1;2* under arsenic stress [7] and sulfur limitation [14], we

investigated the hypothesis that SLIM1 plays a role in arsenic-induced transcriptional

responses. Interestingly, we found that *slim1-1* and *slim1-2* seedlings were highly

sensitive to arsenic. Here, we show that under arsenic treatment, *slim1* mutants

accumulate arsenic, experience high levels of oxidative stress, and fail to induce sulfate

uptake and assimilation. Our results suggest that SLIM1 appears to play an important

role in arsenic sensitivity due primarily to its role in regulating sulfur metabolism and

the cellular redox state.

Results

slim1 **mutants are sensitive to arsenic in root growth assays**

In a previous screen for regulators of cadmium and arsenic-induced gene expression

using a *pSULTR1;2::LUC* reporter construct, we identified new alleles in well-

characterized glutathione biosynthesis genes that play an essential role in cadmium and

arsenic detoxification [7]. Because glutathione is a significant sink of reduced sulfur in

plants, we hypothesized that the transcriptional regulator of sulfur deficiency, SLIM1,

might also play a role in regulating cadmium and arsenic sensitivity in plants. To test

this hypothesis, we performed root growth assays to evaluate the sensitivity of the *slim1-*

1 and *slim1-2* mutant alleles [14] to cadmium and arsenic (Figure 1A-1D).

The root lengths of wild-type (WT) (3.06 ± 0.09 cm, n=22), *slim1-1* (3.19 ± 0.07 cm,

n=19), and *slim1-2* (3.10 ± 0.11 cm, n=21) were not different in the control nutrient

media (see Methods) [17] without addition of cadmium or arsenic (Figure 1; p=0.99997

slim1-1 & p=1.0 *slim1-2*, one-way ANOVA). When grown on plates containing 30 μM

cadmium, WT root growth was inhibited growing only 1.88 ± 0.15 cm (n=10). This

inhibition was similar to that observed for *slim1-1* with a final root length of 1.86 ± 0.13

cm (n=12) and *slim1-2* having a root length of 1.82 ± 0.11 cm (p=1.0 for *slim1-1* &

p=0.99998 for *slim1-2*, n=13) (Figure 1A-1D, Table S1). However, when grown on

minimal media plates containing 10 μM arsenite (As (III)), the root length of WT (1.72

± 0.12 cm, n=14) was longer than both *slim1-1* (0.68 ± 0.06 cm, p=7×10^{-9}, n=14) and

slim1-2 (0.75 ± 0.06 cm, p=1.2×10^{-6}, n=10). These observations suggested that SLIM1

is involved in arsenic signaling. Thus, we further investigated possible mechanisms

underlying *slim1* sensitivity to arsenic.

Arsenic accumulation and antioxidant responses of *slim1* mutants

To determine if arsenic accumulates in the *slim1* mutants, we measured root and shoot

arsenic levels using ICP-MS. In As(III) treated seedlings, we observed no significant

increase in the accumulation of arsenic in the shoots of *slim1-1* (174.8 ± 1.97 mg/Kg

DW, n=3) or *slim1-2* (189.7 ± 3.78 mg/Kg DW, n=3) compared to WT (173.9 ± 4.32

mg/Kg DW, n=3, p=1) (Figure 2A). However, in As(V) treated seedlings, both *slim1-1*

(309.0 ± 47.5 mg/Kg DW, n=3, p=0.01) and *slim1-2* (253.9 ± 19.9 mg/Kg DW, n=3,

p=0.6) accumulated more arsenic in shoots than WT (205.6 ± 11.6 mg/Kg DW, n=3),

although the difference was only significant in *slim1-1* (Figure 2A). These results

suggest an increased root-to-shoot translocation of As(V) in the *slim1* mutants.

In the roots, we found arsenic accumulation in *slim1-1* (1420.0 ± 281.3 mg/Kg DW,

n=3, p=6.77E-3) and *slim1-2* (1473.0 ± 187.9 mg/Kg DW, n=3, p=3.69E-3) compared to

WT (420.1 ± 17.1 mg/Kg DW, n=3) in As(III) treated seedlings (Figure 2B, Table S3).

In comparison, there was no difference in root arsenic accumulation in As(V) treated

seedlings (Figure 2B, Table S3).

Because arsenic is known to cause oxidative stress and induce reactive oxygen species

(ROS) production, we also tested the activity of the key antioxidant enzymes peroxidase

(POD) and superoxide dismutase (SOD) in the *slim1-1* and *slim1-2* mutants. Basal

superoxide dismutase activity in seedlings was similar between WT (70.9 ± 9.65 units/g

FW, n=3), *slim1-1* (77.7 ± 9.44 units/g FW, p=1.0, n=3), and *slim1-2* (Figure 2C; 75.8 ±

4.20 units/g FW, p=1.0, n=3, p=1.0 *slim1-1* and p=1.0 *slim1-2*) grown under control

conditions. When exposed to arsenite (As(III)), WT superoxide dismutase increased to

306.2 ± 7.78 units/g FW (n=3) while the superoxide dismutase activity in the *slim1*

mutants increased dramatically to 709.5 ± 4.85 units/g FW (p=3.6x10-6, n=3) in *slim1-1*

and 621.1 ± 17.7 units/g FW in *slim1-2* (p=2.0x10-8, n=3). Similarly, arsenate (As(V))

treatment increased the WT superoxide dismutase activity to 234.8 ± 27.2 units/g FW

while the *slim1-1* superoxide dismutase activity increased to 543.2 ± 39.4 units/g FW

(p=2.9x10-8, n=3) and the *slim1-2* superoxide dismutase activity increased to 492.1 ±

17.7 units/g FW (p=4.7x10-7, n=3) (Figure 2C, Table S4).

In seedlings, the peroxidase activity was higher under control conditions in *slim1-1*

(94.8 ± 5.95 units/g FW, p=0.009, n=3) and *slim1-2* (120.6 ± 6.04 units/g FW,

p=5.4x10-5, n=3) compared to WT (50.7 ± 1.51 units/g FW, n=3) (Figure 2D). As (III)

exposure increased the peroxidase activity in WT to 107.5 ± 8.91 units/g FW (n=3)

(Figure 2D), while the peroxidase activity in *slim1-1* seedlings increased to 173.6 ± 1.79

units/g FW (p=0.0001, n=3) (Figure 2D). Similar values were observed for As (III)-

treated *slim1-2* seedlings (139.3 ± 4.49 units/g FW, p=0.10, n=3) (Figure 2D, Table S2).

Peroxidase activities showed similar trends under As(V) treatment (Figure 2D, Table

S5).

Decreased shoot glutathione in arsenic-treated *slim1-1* and *slim1-2*

To determine if thiol production might also be altered by arsenic treatment in the *slim1*

mutants, we measured root and shoot cysteine and glutathione levels using fluorescence

HPLC of seedlings exposed to arsenite (As(III)) or arsenate (As(V)) for 48 hours

(Figure 3A-3D). Shoot cysteine levels were lower in *slim1-1* (10.2 ± 0.5 pmol/mg FW,

p=0.04, n=3) and *slim1-2* (12.2 ± 1.0 pmol/mg FW, 0.19, n=3) than WT (23.7 ± 2.7

pmol/mg FW, n=3) in control conditions (Figure 3A, Table S6). No clear decrease in

the cysteine concentration was observed in response to As(III) or As(V) treatment

(Figure 3A, Table S6).

Root cysteine levels were statistically similar for WT, *slim1-1*, and *slim1-2* in control

conditions and were not significantly changed by As(III) or As(V) treatments (Figure

3B, Table S7; One-way ANOVA, Tukey HSD).

Under control conditions, shoot glutathione levels were lower in *slim1-1* (163.6 ± 24.2

pmol/mg FW, n=8) and *slim1-2* (190.2 ± 34.4 pmol/mg FW, n=8) than in WT (382.4 ±

36.2 pmol/mg FW, n=8) (Figure 3C; p=4.99x10-5 for *slim1-1* and p=3.9x10-4 for *slim1-

2*). Shoot glutathione levels decreased in WT from 382.4 ± 36.2 pmol/mg FW (n=8) in

control conditions to 278.1 ± 23.2 pmol/mg FW (n=3) in the As(III) treatment and 269.4

± 12.2 pmol/mg FW (n=3) in the As(V) treatment (Figure 3C). Similarly, shoot

glutathione decreased in the *slim1* mutants under As(III) and As(V) treatments with

slim1-1 having only 110.4 ± 17.5 pmol/mg FW of glutathione in As(III) and 31.4 ± 1.62

pmol/mg FW of glutathione in As(V). Furthermore, *slim1-2* had 83.8 ± 30.9 pmol/mg

FW (n=3) shoot glutathione in As(III) treatment and 43.5 ± 14.7 pmol/mg FW (n=3) in

As(V) treatment – an 80% decrease compared to control (Figure 3C, Table S8).

Root glutathione levels decreased under both As(III) and As(V) treatments for all

genotypes. However, glutathione levels in roots showed no differences between

genotypes within each treatment (Figure 3D, Table S9; One-way ANOVA, Tukey

HSD). In summary, thiol measurements showed that while cysteine and glutathione

levels were not dramatically decreased in the roots of the *slim1* mutant alleles compared

to WT (Figure 3B and D), glutathione levels were decreased in shoots of *slim1-1* and

slim1-2 compared to WT plants (Figure 3C).

Shoot sulfate and phosphate accumulation in *slim1* mutants

Arsenic is thought to be actively taken up by phosphate transporters as As(V); however,

once inside plant cells, it is reduced to As(III) and can move within plants through

aquaporins [18,19]. Mutants in SLIM1 were previously shown to be impaired in root-to-

shoot translocation of sulfate [14], but the translocation of other anions, including

phosphate, was not reported. Thus, based on the slight arsenic accumulation in shoots of

As(V) treated plants noted by ICP-MS (Figure 2 A and B), we hypothesized that

phosphate transport might also be impaired in the *slim1* mutants.

To determine if phosphate and sulfate translocation are impaired in the *slim1* mutants

under arsenic treatment, we measured sulfate and phosphate accumulation in both roots

and shoots of plants treated with As (V) for 48 hours. Interestingly, shoot phosphate

accumulation was higher in *slim1-1* and *slim1-2* than WT in all treatments (Figure 4A,

Table S10; p=5x10-6 for *slim1-1* and p=0.004 for *slim1-2*; One-way ANOVA, Tukey

HSD).

Root phosphate accumulation was similar for WT (5.03 ± 0.27 nmol/mg FW, n=5),

slim1-1 (4.95 ± 1.07 nmol/mg FW, n=3), and *slim1-2* (6.33 ± 0.80 nmol/mg FW, n=4) in

control conditions and was not different under As(V) treatment (Figure 4B). Thus, the

enhanced root-vs.-shoot phosphate accumulation observed in *slim1-1* and *slim1-2*

suggests an indirect role for SLIM1 in regulating phosphate and arsenate transport

(Figure 4A, 4B, Table S11).

Furthermore, sulfate accumulation in shoots was impaired in *slim1-1* (0.73 ± 0.10

nmol/mg FW, $p=1.7 \times 10^{-6}$, n=5) and *slim1-2* (0.61 ± 0.20 nmol/mg FW, $p=7.3 \times 10^{-7}$,

n=5) relative to WT (3.09 ± 0.18 nmol/mg FW, n=5) in control conditions (Figure 4A),

consistent with previous findings [14]. WT seedlings showed a decrease in shoot sulfate

upon As(V) treatment decreasing to 1.86 ± 0.44 nmol/mg FW (n=5) (Figure 4C, Table

S12, $p=0.008$, One-way ANOVA, Tukey HSD).

Root sulfate accumulation was similar between WT (2.11 ± 0.14 nmol/mg FW, n=5),

slim1-1 (1.95 ± 0.43 nmol/mg FW, n=3), and *slim1-2* (1.45 ± 0.21 nmol/mg FW, n=4) in

control conditions. Furthermore, WT (2.04 ± 0.16 nmol/mg FW, n=4), *slim1-1* ($1.72 \pm$

0.13 nmol/mg FW, n=5), and *slim1-2* (1.85 ± 0.15 nmol/mg FW, n=3) root sulfate were

not different in the As(V) treatment (Figure 4D, Table S13).

Microarray analyses of *slim1* mutants under As treatment

The current model for arsenic uptake and tolerance in plants suggests that arsenic is

taken up from the soil in the form of arsenate (As(V)). Once it has entered the plant, it is

rapidly reduced to arsenite (As(III)) by the arsenate reductase HAC1[20]. It has been

proposed that As(III) can be removed from the root by an unidentified efflux

transporter[21]. In rice, the aquaporin LSI1 is known to mediate As(III) efflux; however,

additional efflux transporters remain elusive[21]. A recent RNA-seq experiment using a

T-DNA mutant allele of SLIM1 (*eil3*) did not find misregulation of any aquaporin genes

in the roots of the *slim1* mutant under control or sulfur deficiency conditions[22]. Thus,

due to the observed arsenic accumulation in the roots of *slim1* mutants, we hypothesized

that the elusive As(III) efflux transporter, or alternatively an As(III) uptake transporter,

might be disrupted in an arsenic-dependent manner in the *slim1* mutant background.

To test this hypothesis and uncover genes disrupted in an arsenic-dependent manner in

the *slim1-1* mutant, we performed microarray analyses on WT and *slim1-1* seedlings

exposed to arsenite for 48 hours. Raw expression values were normalized via the R

'affy' package using the Robust Multi-Array Average (RMA) Expression Measure.

Differential gene expression was evaluated using the R package 'limma', including a

multiple test correction. We then performed a significance analysis to identify genes

disrupted under arsenic treatment and compared these to previously published putative

targets of SLIM1 obtained by DNA affinity purification sequencing (DAP-seq) [23].

From the microarray analyses, we identified 11 genes significantly differentially

upregulated by arsenic (WT +As vs. *slim1-1* + As) (Supplemental Table S14). Ten of

the 11 genes (AT3G49580, AT1G04770, AT1G12030, AT4G04610, AT4G21990,

AT5G24660, AT5G26220, AT5G48850, AT4G20820, AT1G36370) were identified as

putative targets of SLIM1 by DNA affinity purification sequencing (DAP-Seq)

(Supplemental Table S14). Many of the 11 upregulated genes in *slim1-1* are associated

with sulfur metabolism.

Genes that appear to be negatively regulated by SLIM1 include CGCT2;1

(AT5G26220), APR1 (AT4G04610) and APR3 (AT4G21990), which were upregulated

in *slim1-1* compared to WT in the presence of arsenic (WT +As vs. *slim1-1* + As).

APR1 and APR3 are involved in the reduction of sulfate into sulfide [24] and have been

shown to be induced by toxic metal stress [7]. Similarly, the LOW SULFUR 1 (LSU1,

AT3G49580) and LOW SULFUR 2 (LSU2, AT5G24660) genes were expressed at

higher levels in *slim1-1* than WT under arsenic treatment (WT +As vs. *slim1-1* + As).

Interestingly, Six of the 11 genes (GGCT2;1, APR3, LSU1, LSU2, SDI1, & SHM7)

belong to a highly co-regulated cluster of genes that respond to O-acetylserine

treatment[25].

Microarray analyses also identified 10 significantly down-regulated genes under arsenic

treatment compared to WT (WT +As vs. *slim1-1* +As) (Supplemental Figure S1)

($p<0.05$, Fold Change >2). Only one gene - SULTR1; 2 (At1G78000) - was identified as

a putative target of SLIM1 by DAP-Seq (Supplemental Table S14). Thus, our analyses

confirm the reported function of SLIM1 as a transcriptional activator of *SULTR1;2* and

show that this role is conserved under arsenic treatment and sulfur deficiency. The

remaining ten genes are involved in hormone signaling (AT1G63030, AT5G13220 &

AT5G52050), redox regulation (AT3G06590 & AT1G03020), iron homeostasis

(AT3G25190 & AT5G01600), glucosinolate biosynthesis (AT5G23020), ubiquitination

(AT1G24330), and an uncharacterized protein (AT2G17660). Based on their putative

functions, these genes encode stress response-related genes. More experiments are

needed to determine if SLIM1 is a direct transcriptional regulator of these genes under

arsenic stress.

The present transcriptome data suggest that SLIM1 can function as both a transcriptional

enhancer as well as a transcriptional repressor of specific genes in a condition-specific

manner. Furthermore, the present study provides evidence that SLIM1 plays an essential

role in the regulation of sulfur metabolism gene expression in response to arsenic.

Discussion

Plant exposure to arsenic causes rapid changes in gene expression [7,26,27]. However,

the transcription factors that function in arsenic-induced gene expression remain largely

unknown. The few transcriptional regulators that have been identified, such as WRKY6,

WRKY45, and OsARM1 (Arsenite-Responsive Myb1) [27–29], have been implicated in

the regulation of arsenic transporters while regulators of arsenic detoxification remain

unknown. To test the hypothesis that the SLIM1 transcription factor is involved in

arsenic resistance and signaling, we evaluated the sensitivity of *slim1-1* and *slim1-2* to

arsenic exposure. We found the *slim1* mutants were more sensitive to arsenic than

control plants. Arsenic treatment caused high levels of oxidative stress in the *slim1*

mutant alleles based on superoxide dismutase and peroxidase activities. Furthermore,

thiol and sulfate measurements show that *slim1* mutants are impaired in both thiol and

sulfate accumulation. Arsenic treatment did not further decrease sulfate levels in roots.

In contrast, the concentration of the thiol GSH was greatly decreased in *slim1* mutant

alleles. Furthermore, peroxidase and superoxide dismutase measurements show that

arsenic treatments cause increased levels of oxidative stress in the *slim1* mutants.

We also observed a slight increase in arsenic accumulation in the shoots of *slim1*

mutants treated with arsenic. This arsenic accumulation was accompanied by a

significant increase in shoot phosphate translocation in the *slim1* mutants. Because of

the chemical similarity between phosphate and arsenic oxyanions, future research could

investigate the hypothesis that the misregulation of phosphate transporters may

contribute to the observed increase in shoot arsenic in the *slim1* mutants. A recent study

identified mutants in Ethylene Response Factor genes (ERF34 & ERF35) that are

sensitive to both arsenite (As(III)) and arsenate (As(V)) [30]. Interestingly, similar to the

slim1 mutants, the double *erf34erf35* mutants were far less sensitive to cadmium than

arsenic suggesting the arsenic sensitivity is not exclusively due to thiol accumulation.

Furthermore, gene expression studies showed that several phosphate transporters were

down-regulated in *erf34erf35* suggesting PHTs may play a role in both As(III) and

As(V) sensitivity and/or transport [30].

Thiol measurements confirmed the role of SLIM1 in sulfate metabolism and thiol

production [14,22,31], as *slim1* mutants contained lower cysteine and glutathione levels

in shoots than WT. We hypothesized that the weaker cadmium sensitivity of *slim1*

mutant alleles might be linked to thiol accumulation, but we observed no significant

differences decrease in shoot GSH in the *slim1* mutants under Cd treatment

(Supplemental Figure S2). However, SLIM1 upregulates the root-to-shoot transport of

sulfate, which restricts sulfate assimilation mainly to the roots in *slim1* mutants. Root

sulfate levels are maintained by the high-affinity sulfate transporter SULTR1;1, which is

regulated in a SLIM1-independent manner [14]. Thus cysteine and glutathione

biosynthesis can occur in the roots. As described previously, glutathione is essential for

producing phytochelatins –arsenic chelating compounds necessary for detoxification and

storage. The heavy metal cadmium also binds to phytochelatins. Interestingly, recent

research has shown a less dramatic effect of cadmium exposure in *slim1* mutants than

wild-type controls [31], which we have also observed (Figure 1C, D). Thus, the present

study shows that the SLIM1 transcription factor plays a more central role in mediating

arsenic resistance relative to cadmium resistance. A possible hypothesis that may

contribute to this observation is that cadmium can be sequestered in vacuoles via two

independent transport pathways: via phytochelatin transport [8,32] and via thiol-

independent HMA3-mediated cadmium transport [33].

Sulfate measurements confirmed that SLIM1 is a major transcriptional regulator of

sulfate uptake and translocation [14]. Our microarray analyses also identified 11 genes

significantly differentially upregulated by arsenic (Supplemental Table S14), of which

ten of the 11 genes were identified as putative targets of SLIM1 by DNA affinity

purification sequencing (DAP-Seq). Interestingly, nine of these genes are involved in

sulfur assimilation or redox signaling. One of these genes, GGCT2;1, is involved in

glutathione recycling and has also been implicated in arsenic tolerance[34–36]

Furthermore, six of these sulfur metabolism genes belong to a highly co-regulated

cluster of genes that respond to O-acetylserine treatment [25]. While previous studies

show these genes can regulate sulfur assimilation in a SLIM1 independent manner

[25,37], results from DAP-Seq and microarray results from the current study suggest

SLIM1 may act as a negative regulator of these genes during arsenic stress

Shoot sulfate accumulation was significantly lower in the *slim1* mutants under all

conditions tested. Decreased shoot sulfate was accompanied by an increase in shoot

phosphate in the *slim1* mutants. Similar anion compensation was noted in the

Arabidopsis *phr1* mutant, which accumulates higher sulfate levels when grown under

low phosphate conditions indicating crosstalk between phosphate and sulfate transport

[38]. In fact, *PHR1* has been proposed to act both positively in the regulation of root-to-

shoot sulfate translocation via the sulfate transporter *SULTR1;3*, and negatively to

repress other sulfate transporters under phosphate deficiency [39]. We did not identify

any significantly misregulated phosphate transporters (PHTs) in our microarray

analyses. One possible explanation is that PHTs belong to a large gene family and

demonstrate a high degree of genetic redundancy. Thus, a small decrease in the

expression of several PHTs may result in measurable changes in phosphate

accumulation without any individual transcript misregulation meeting the stringent

criteria used in our microarray analyses. Xie et al. identified an artificial microRNA

mutant targeting three high-affinity phosphate transporters showing a similar sensitivity

to arsenite [30]. A recent study showing that sulfate deficiency increases phosphate

accumulation in Arabidopsis further supports this hypothesis [40].

In summary, we show here that the SLIM1 transcription factor plays an important role

in mediating arsenic resistance and in arsenic-induced gene expression. Our results

suggest that the arsenic sensitivity of *slim1* mutants can be explained by decreased thiol

production resulting in increased oxidative stress and in increased arsenic accumulation.

Interestingly, we found that the *slim1* mutant alleles do not show a strong cadmium

sensitivity, consistent with a recent study [31] indicating a difference in the rate-limiting

functions of the thiol synthesis pathway in processing arsenic and cadmium that we

discuss here. We also identify a number of genes regulated by SLIM1 in an arsenic-

dependent manner with DAP-seq data set analyses indicating direct binding of SLIM1 to

arsenic-dependent differentially-expressed genes. Taken together, our data support a

model in which SLIM1 is both a positive and negative regulator of gene expression in

response to arsenic.

Experimental Procedures

Arabidopsis accessions

The WT *Arabidopsis thaliana* ecotype used in this study is Columbia (Col-0).

The *slim1-1* and *slim1-2* mutants were generated in the Col-0 genetic background and

were kindly provided by Dr. Akiko Maruyama-Nakashita [14].

Plant Growth Media & Conditions

Seeds were surface sterilized by briefly soaking in 70% ethanol before allowing them to

dry in a sterile hood. For root growth experiments and enzymatic assay experiments,

surface-sterilized seeds were plated on minimal media containing 1/10-strength

Hoagland solution, 1% phytoagar (Duchefa, http://www.duchefa.com), pH 5.6. For the

microarray experiments, seeds were plated on1/2-strength MS standard medium

(M5519; Sigma-Aldrich, http:// www.sigmaaldrich.com) buffered with 1 mM 2-(N-

morpholine)-ethanesulphonic acid (MES), 1% phytoagar

(Duchefa, http://www.duchefa.com) and the pH was adjusted to 5.6 with 1.0 M KOH.

Seeds were then stratified with cold treatment at 4°C for 48 h, and grown under

controlled conditions (150 μmol m^{-2} s^{-1}, 70% humidity, 16-h light at 21°C/8-h dark at

18°C) for the specified time. For toxic metal(loid) treatments, the specified amounts of

either cadmium or arsenic were added to the autoclaved base media in a sterile hood

prior to pouring the plates. Concentrated stock solutions of cadmium and arsenic were

filter-sterilized prior to use.

Statistical Analyses

The root growth, thiol, peroxidase, superoxide, and anion data were all analyzed using

one-way ANOVA followed by a Tukey posthoc test to determine significance.

Significance groups are indicated in the figures, and key p-values are stated in the text.

Root Length Measurements

For root growth experiments, surface-sterilized seeds of WT, *slim1-1,* and *slim1-2* were

plated on minimal media (2.5 mM H_3PO_4, 5 mM KNO_3, 2 mM $MgSO_4$, 1 mM

$(CaNO_3)_2$, 1 mM MES, 1% phytoagar pH 5.7) supplemented with 30 μM Cd or 10 μM

As (III) [17]. Plates were placed in the dark two days at 4°C for vernalization and then

transferred to a growth chamber. After 7 days of growth, seedlings were photographed,

and root length was measured using ImageJ.

Antioxidant Enzyme Assays

Seedling samples were weighed and pulverized in liquid nitrogen after treatment. The

powder was dissolved in pre-cooled 50 mM phosphate buffer (pH 7.8) to extract the

superoxide dismutase (SOD). The extract was then centrifuged at 12 000g for 10 min,

resulting in a crude enzyme supernatant solution. In a separate 10 ml tube, 1.9 ml

reaction buffer (50 mM phosphate buffer, pH 7.8, 9.9 mM L-methionine, 57 μM NBT

solution, 1 M EDTA-Na_2 solution, 0.0044% (w/v) riboflavin) and 0.1 ml enzyme

solution were mixed and placed into 250 μmol m-2s-1 light for 20 min. Additionally,

another separate 10 ml tube was procured, where the enzyme solution was replaced with

water as a control. The reagent was added according to the above steps, where one tube

was placed in the light together with the sample, and the other was placed in the dark

where the reaction was allowed to complete. The control tube that was placed in the

dark was blanked, and the absorbance of each tube was measured at 560 nm. Peroxidase

(POD) was extracted in 50 mM phosphate buffer (pH 7.0). 30 μl of enzyme solution was

mixed with reaction buffer containing 1.77 mL of 50 mM sodium phosphate buffer (pH

7.0), 0.1 mL of 4% guaiacol and 0.1 mL of 1% (v/v) H_2O_2. Increased absorbance was

recorded at 470 nm for 1 min. All reported enzyme activities are means of 3-5

biologically independent samples, and error bars indicate the standard error of the mean

(SEM).

Arsenic Determination by ICP-MS

Plant material was harvested, dried at $70^{o}C$ for at least 48 hours before being aliquoted

and weighed. Approximately 10 mg of dried plant material was mixed with 1 ml of

concentrated nitric acid and digested by heating at $100^{o}C$ for approximately 30 minutes

or until the solution became transparent and particle-free. These digests were diluted

with deionized water and measured by ICP-MS for total arsenic concentrations at the

University of Cologne Biocenter Mass Spectrometry Platform. All reported ion

quantities are means of 3-5 biologically independent samples, and error bars indicate the

standard error of the mean (SEM).

Anion Extraction and Measurement by Ion Chromatography

To quantify the water-soluble anion concentrations (phosphate and sulfate) in plant

tissues, 10-30 mg of fresh tissue was harvested and flash-frozen in liquid nitrogen.

Frozen tissue was then pulverized using a bead mill (make & model), and anions were

extracted by addition of 1000 μL of sterile Milli-Q-water and incubating for 60 minutes

at $4^{o}C$ while shaking at 1500 rpm. The extraction process was stopped by incubating at

$95^{o}C$ for 15 minutes. Cell debris was removed by centrifugation at $4^{o}C$ for 15 minutes,

and 100-200μL of supernatant was used for anion exchange chromatography. An

automatic ion analyzer (DX 120, Dionex Corporation, Sunnyvale, CA, United States)

equipped with an IonPac^{TM} column (AS9-SC, 4×250 mm; Dionex, Thermo Fisher

Scientific GmbH; Waltham, MA, United States) was used to separate and quantify the

anions. Anions were eluted with an elution buffer of 2.0 mM Na_2CO_3 and 0.75 mM

$NaHCO_3$. Ion concentrations were detected using a conductivity detector module (CDM,

Dionex Corporation, CA, United States). All reported anion quantities are means of 3-5

biologically independent samples of tissue pooled from 4-6 individual seedlings (12-30

seedlings in total), and error bars indicate the standard error of the mean (SEM).

Thiol Detection By Fluorescence HPLC

The thiol-containing compounds cysteine and GSH were analyzed using fluorescence

detection HPLC as described by [41]. To analyze the levels of these thiol compounds,

plants were grown on minimal growth media plates for 12 days then transferred to fresh

media plates containing either 20 μM cadmium, 100 μM arsenate, or control minimal

media. To minimize the oxidation of thiol compounds during the extraction, plant

seedlings were flash-frozen in liquid nitrogen immediately after harvesting and then

pulverized using a bead mill and extracted as described by [42]. Thiols were extracted

from homogenized plant material with 1 mL 0.1 M HCl for 40 min at 25°C. After

centrifugation for 5 min at 14,000 g and 4°C, thiols in the supernatant were reduced by

mixing 60 μL of the supernatant with 100 μL 2-(cyclohexylamino)ethanesulfonic acid

(0.25 M, pH 9.4) and 35 μL DTT (10 mM, freshly prepared). The mixture was incubated

at 25°C for 40 min. Thiols were derivatized by adding 5 μL (25 mM) monobromobimane

(SigmaAldrich, Cat#B4380). Derivatization was stopped by adding 110 μL methane

sulfonic acid (100 mM) and clarified by centrifugation for 15 min at 14,000 g and 4°C.

Forty microliters of the derivatization mix were used for HPLC analysis using the

Dionex Ultimate 3000 HPLC System. Derivatized thiols were separated in a Eurosphere

100-3 C18, 150×4 mm column (Knauer), and were detected by fluorescence detection

with an excitation of 380 nm and emission detection at 480 nm. The peaks of thiol

compounds were identified and quantified by comparison with cysteine and glutathione

standards purchased from Sigma-Aldrich. All reported thiol quantities are statistical

means of 4-5 biologically independent experiments (16-30 seedlings per experiment).

Error bars indicate the standard error of the mean (SEM).

Microarray Analyses

To evaluate transcriptional differences in the *slim1* mutants under cadmium and arsenic

stress, we performed microarray analyses. To obtain tissue for the microarray analysis,

plants were grown on ¼ MS plates for 12 days then transferred to fresh media plates

containing either 100 μM cadmium or 20 μM arsenite. Whole seedlings were then

harvested in 2mL Eppendorf tubes, flash-frozen in liquid nitrogen, and stored at -80$^{\text{o}}$C

until further processing. The tissue was subsequently pulverized using a bead mill by

adding three 2.5mm glass beads to each tube and grinding for 15 seconds. RNA was

extracted using the Qiagen RNEasy mini kit (Cat#74104) per the manufacturer's

instructions (www.qiagen.com). RNA quality was assessed by spectrophotometer and

gel electrophoresis before submission to the University of California, San Diego Gene

Expression Core facility for processing. Results were analyzed using R and the

Bioconductor suite of microarray analytical packages as indicated in the text.

Acknowledgments

We thank Dr. Akiko Maruyama-Nakashita for kindly providing *slim1* mutant seeds.

This research was funded by the National Institute of Environmental Health Sciences of

the National Institutes of Health under Award Number P42ES010337 (JIS). The

microarray data are available through the NCBI GEO database (series record

GSE138943).

Conflicts of Interests

538   The authors have no conflicts of interest to declare. All co-authors have seen and agree

539   with the contents of the manuscript, and there is no financial interest to report. We

540   certify that the submission is original work and is not under review at any other

541   publication.

References

1 Clemens S (2006) Toxic metal accumulation, responses to exposure and mechanisms

of tolerance in plants. *Biochimie* **88**, 1707–1719.

2 Ogunseitan OA, Schoenung JM, Saphores J-DM & Shapiro AA (2009) The

electronics revolution: from e-wonderland to e-wasteland. *Science (80-)* **326**, 670–

671.

3 Satarug S, Garrett SH, Sens MA & Sens DA (2009) Cadmium, environmental

exposure, and health outcomes. *Environ Health Perspect* **118**, 182–190.

4 Han FX, Banin A, Su Y, Monts DL, Plodinec JM, Kingery WL & Triplett GE (2002)

Industrial age anthropogenic inputs of heavy metals into the pedosphere.

Naturwissenschaften **89**, 497–504.

5 Larison JR, Likens GE, Fitzpatrick JW & Crock JG (2000) Cadmium toxicity among

wildlife in the Colorado Rocky Mountains. *Nature* **406**, 181.

6 Mendoza-Cózatl DG, Xie Q, Akmakjian GZ, Jobe TO, Patel A, Stacey MG, Song L,

Demoin DW, Jurisson SS, Stacey G & Schroeder JI (2014) OPT3 is a component

of the iron-signaling network between leaves and roots and misregulation of OPT3

leads to an over-accumulation of cadmium in seeds. *Mol Plant* **7**.

7 Jobe TO, Sung D, Akmakjian G, Pham A, Komives EA, Mendoza-Cózatl DG &

Schroeder JI (2012) Feedback inhibition by thiols outranks glutathione depletion: a

luciferase-based screen reveals glutathione-deficient γ-ECS and glutathione

synthetase mutants impaired in cadmium-induced sulfate assimilation. *Plant J* **70**,

783–795.

8 Mendoza-Cózatl DG, Zhai Z, Jobe TO, Akmakjian GZ, Song W-Y, Limbo O, Russell

MR, Kozlovskyy VI, Martinoia E, Vatamaniuk OK, Russell P & Schroeder JI (2010) Tonoplast-localized Abc2 transporter mediates phytochelatin accumulation in vacuoles and confers cadmium tolerance. *J Biol Chem* **285**.

Verbruggen N, Hermans C & Schat H (2009) Molecular mechanisms of metal hyperaccumulation in plants. *New Phytol* **181**, 759–776.

Rea PA (2007) Plant ATP-binding cassette transporters. *Annu Rev Plant Biol* **58**, 347–375.

Mendoza-Cózatl DG & Moreno-Sánchez R (2005) Cd2+ transport and storage in the chloroplast of Euglena gracilis. *Biochim Biophys Acta (BBA)-Bioenergetics* **1706**, 88–97.

Jobe TO, Zenzen I, Rahimzadeh Karvansara P & Kopriva S (2019) Integration of sulfate assimilation with carbon and nitrogen metabolism in transition from C3 to C4 photosynthesis. *J Exp Bot* **70**, 4211–4221.

Jobe, Timothy O., Kopriva S (2018) Sulfur Metabolism in Plants. *Encycl Life Sci*.

Maruyama-Nakashita A, Nakamura Y, Tohge T, Saito K & Takahashi H (2006) Arabidopsis SLIM1 is a central transcriptional regulator of plant sulfur response and metabolism. *Plant Cell* **18**, 3235–3251.

Foyer CH & Noctor G (2011) Ascorbate and Glutathione: The Heart of the Redox Hub. *Plant Physiol* **155**, 2 LP – 18.

NOCTOR G, MHAMDI A, CHAOUCH S, HAN YI, NEUKERMANS J, MARQUEZ-GARCIA B, QUEVAL G & FOYER CH (2012) Glutathione in plants: an integrated overview. *Plant Cell Environ* **35**, 454–484.

Lee DA, Chen A & Schroeder JI (2003) ars1, an Arabidopsis mutant exhibiting

increased tolerance to arsenate and increased phosphate uptake. *Plant J* **35**, 637–

646.

18 Catarecha P, Segura MD, Franco-Zorrilla JM, García-Ponce B, Lanza M, Solano R,

Paz-Ares J & Leyva A (2007) A mutant of the Arabidopsis phosphate transporter

PHT1; 1 displays enhanced arsenic accumulation. *Plant Cell* **19**, 1123–1133.

19 Bienert GP, Thorsen M, Schüssler MD, Nilsson HR, Wagner A, Tamás MJ & Jahn

TP (2008) A subgroup of plant aquaporins facilitate the bi-directional diffusion of

As (OH) 3 and Sb (OH) 3 across membranes. *BMC Biol* **6**, 26.

20 Shi S, Wang T, Chen Z, Tang Z, Wu Z, Salt DE, Chao D-Y & Zhao F-J (2016)

OsHAC1;1 and OsHAC1;2 Function as Arsenate Reductases and Regulate Arsenic

Accumulation. *Plant Physiol* **172**, 1708 LP – 1719.

21 Chen Y, Han Y-H, Cao Y, Zhu Y-G, Rathinasabapathi B & Ma LQ (2017) Arsenic

Transport in Rice and Biological Solutions to Reduce Arsenic Risk from Rice   .

Front Plant Sci   **8**, 268.

22 Dietzen C, Koprivova A, Whitcomb SJ, Langen G, Jobe TO, Hoefgen R & Kopriva S

(2020) The Transcription Factor EIL1 Participates in the Regulation of Sulfur-

Deficiency Response. *Plant Physiol* **184**, 2120 LP – 2136.

23 O'Malley RC, Huang SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M,

Gallavotti A & Ecker JR (2016) Cistrome and epicistrome features shape the

regulatory DNA landscape. *Cell* **165**, 1280–1292.

24 Lee B, Koprivova A & Kopriva S (2011) The key enzyme of sulfate assimilation,

adenosine 5′-phosphosulfate reductase, is regulated by HY5 in Arabidopsis. *Plant J*

67, 1042–1054.

25 Hubberten H, Klie S, Caldana C, Degenkolbe T, Willmitzer L & Hoefgen R (2012)

Additional role of O-acetylserine as a sulfur status-independent regulator during

plant growth. *Plant J* **70**, 666–677.

26 Fu S-F, Chen P-Y, Nguyen QTT, Huang L-Y, Zeng G-R, Huang T-L, Lin C-Y &

Huang H-J (2014) Transcriptome profiling of genes and pathways associated with

arsenic toxicity and tolerance in Arabidopsis. *BMC Plant Biol* **14**, 94.

27 Castrillo G, Sánchez-Bermejo E, de Lorenzo L, Crevillén P, Fraile-Escanciano A, TC

M, Mouriz A, Catarecha P, Sobrino-Plata J, Olsson S, Leo del Puerto Y, Mateos I,

Rojo E, Hernández LE, Jarillo JA, Piñeiro M, Paz-Ares J & Leyva A (2013)

WRKY6 Transcription Factor Restricts Arsenate Uptake and Transposon

Activation in Arabidopsis *Plant Cell* **25**, 2944 LP – 2957.

28 Wang H, Xu Q, Kong Y-H, Chen Y, Duan J-Y, Wu W-H & Chen Y-F (2014)

Arabidopsis WRKY45 Transcription Factor Activates PHOSPHATE

TRANSPORTER1;1 Expression in Response to Phosphate Starvation.

Plant Physiol **164**, 2020 LP – 2029.

29 Wang F-Z, Chen M-X, Yu L-J, Xie L-J, Yuan L-B, Qi H, Xiao M, Guo W, Chen Z,

Yi K, Zhang J, Qiu R, Shu W, Xiao S & Chen Q-F (2017) OsARM1, an R2R3

MYB Transcription Factor, Is Involved in Regulation of the Response to Arsenic

Stress in Rice. *Front Plant Sci* **8**, 1868.

30 Xie Q, Yu Q, Jobe TO, Pham A, Ge C, Guo Q, Liu J, Liu H, Zhang H, Zhao Y, Xue

S, Hauser F & Schroeder JI (2021) An amiRNA screen uncovers redundant CBF

& ERF34/35 transcription factors that differentially regulate arsenite and

cadmium responses. *bioRxiv*.

31 Yamaguchi C, Khamsalath S, Takimoto Y, Suyama A, Mori Y, Ohkama-Ohtsu N &

Maruyama-Nakashita A (2020) SLIM1 Transcription Factor Promotes Sulfate

Uptake and Distribution to Shoot, Along with Phytochelatin Accumulation, Under

Cadmium Stress in Arabidopsis thaliana. *Plants* **9**, 163.

32 Song W-Y, Park J, Mendoza-Cózatl DG, Suter-Grotemeyer M, Shim D,

Hörtensteiner S, Geisler M, Weder B, Rea PA & Rentsch D (2010) Arsenic

tolerance in Arabidopsis is mediated by two ABCC-type phytochelatin transporters.

Proc Natl Acad Sci **107**, 21187–21192.

33 Takahashi R, Bashir K, Ishimaru Y, Nishizawa NK & Nakanishi H (2012) The role

of heavy-metal ATPases, HMAs, in zinc and cadmium transport in rice. *Plant*

Signal Behav **7**, 1605–1607.

34 Joshi NC, Meyer AJ, Bangash SAK, Zheng Z-L & Leustek T (2019) Arabidopsis γ-

glutamylcyclotransferase affects glutathione content and root system architecture

during sulfur starvation. *New Phytol* **221**, 1387–1397.

35 Paulose B, Chhikara S, Coomey J, Jung H, Vatamaniuk O & Dhankher OP (2013) A

γ-Glutamyl Cyclotransferase Protects Arabidopsis Plants

from Heavy Metal Toxicity by Recycling Glutamate to Maintain Glutathione

Homeostasis. *Plant Cell* **25**, 4580 LP – 4595.

36 Kumar S, Kaur A, Chattopadhyay B & Bachhawat AK (2015) Defining the cytosolic

pathway of glutathione degradation in Arabidopsis thaliana: role of the ChaC/GCG

family of γ-glutamyl cyclotransferases as glutathione-degrading enzymes and

AtLAP1 as the Cys-Gly peptidase. *Biochem J* **468**, 73–85.

37 Aarabi F, Naake T, Fernie AR & Hoefgen R (2020) Coordinating sulfur pools under

sulfate deprivation. *Trends Plant Sci*.

38 Rouached H, Secco D, Arpat B & Poirier Y (2011) The transcription factor PHR1

plays a key role in the regulation of sulfate shoot-to-root flux upon phosphate

starvation in Arabidopsis. *BMC Plant Biol* **11**, 19.

39 Rouached H (2011) Multilevel coordination of phosphate and sulfate homeostasis in

plants. *Plant Signal Behav* **6**, 952–955.

40 Allahham A, Kanno S, Zhang L & Maruyama-Nakashita A (2020) Sulfur Deficiency

Increases Phosphate Accumulation, Uptake, and Transport in Arabidopsis thaliana.

Int J Mol Sci **21**.

41 Fahey RC & Newton GL (1987) Determination of low-molecular-weight thiols using

monobromobimane fluorescent labeling and high-performance liquid

chromatography. In *Methods in enzymology* pp. 85–96. Elsevier.

42 Krueger S, Niehl A, Lopez Martin MC, Steinhauser D, Donath A, Hildebrandt T,

Romero LC, Hoefgen R, Gotor C & Hesse H (2009) Analysis of cytosolic and

plastidic serine acetyltransferase mutants and subcellular metabolite distributions

suggests interplay of the cellular compartments for cysteine biosynthesis in

Arabidopsis. *Plant Cell Environ* **32**, 349–367.

 Figures & Figure Legends

Figure 1. Root growth inhibition of *slim1* mutants grown on cadmium or arsenic-containing media. The *slim1-1* and *slim1-2* mutant alleles were compared to wild-type controls (WT) grown on control minimal media and media containing 30 μM Cd or 10 μM As(III) for 7 days (A - D). Root growth was quantified using ImageJ (one-way ANOVA, Tukey HSD).

Figure 2. *slim1* **mutants accumulate arsenic in roots and have high antioxidant**

activity when exposed to arsenic. *slim1* mutants grown on arsenic-containing media

accumulate arsenic in the shoots when grown on As(V) (A) but accumulate arsenic in

the roots when grown on As(III) (B). Growth on arsenic-containing media caused an

increase in superoxide dismutase (C) enzyme and peroxidase dismutase enzyme (D)

activities in both the *slim1-1* and *slim1-2* mutants compared to WT controls.

Figure 3. T**hiol accumulation of *slim1* mutants grown on arsenic.** Total shoot

cysteine levels in *slim1-1* and *slim1-2* compared to WT (A). Total root cysteine levels in

slim1-1 and *slim1-2* compared to WT (B). Total shoot glutathione levels for *slim1-1* and

slim1-2 compared to WT (C). Total root glutathione levels for WT, *slim1-1*, and *slim1-2*

(D).

Figure 4. Anion accumulation in *slim1* mutants grown on arsenic. Total shoot

phosphate levels in *slim1-1* and *slim1-2* compared to WT (A). Total root phosphate

levels in *slim1-1* and *slim1-2* compared to WT (B). Total shoot sulfate levels for *slim1-1*

and *slim1-2* compared to WT (C). Total root sulfate levels for WT, slim1-1, and *slim1-2*

(D).