A potent tumour-suppressor TCF11, than its prototypic Nrf1α, is defined by similar yet different regulatory profiles of both isoforms, with a striking disparity from Nrf2

Meng Wang[1], Yonggang Ren[1,2], Shaofan Hu[1], Keli Liu[1], Lu Qiu[1,3] and Yiguo Zhang[1]*

1 The Laboratory of Cell Biochemistry and Topogenetic Regulation, College of Bioengineering, Chongqing University, No. 174 Shazheng Street, Shapingba District, Chongqing 400044, China.

2 Department of Biochemistry, North Sichuan Medical College, Nanchong 637000, China.

3 School of Life Sciences, Zhengzhou University, No. 100 Kexue Avenue, Zhengzhou 450001, Henan, China

* Correspondence: yiguozhang@cqu.edu.cn or eaglezhang64@gmail.com

Abstract

Nrf1 and Nrf2, as two principal CNC-bZIP transcription factors, regulate similar but different targets involved in a variety of biological functions for maintaining cell homeostasis and organ integrity. Of note, the unique topobiological behavior of Nrf1 makes its functions more complicated than Nrf2, because it is allowed for alternatively transcribing and selectively splicing to yield multiple isoforms (e.g., TCF11, Nrf1α). Here, to gain a better understanding of their similarities and differences in distinct regulatory profiles, four distinct cell models for stably expressing *TCF11*, *TCF11ᐤN*, *Nrf1α* or *Nrf2* have been established by an Flp-In™ T-REx™-293 system and identified by transcriptomic sequencing. Further analysis revealed that Nrf1α and TCF11 have similar yet different regulatory profiles, though both contribute basically to positive regulation of their co-targets, which are disparate from those regulated by Nrf2. Such disparity in gene regulation by Nrf1 and Nrf2 was also corroborated by scrutinizing comprehensive functional annotation of their specific and/or common target genes. Conversely, the mutant TCF11ᐤN, resulting from a deletion of the N-terminal amino acids 2-156 from TCF11, resembles Nrf2 with largely consistent structure and function. Interestingly, our further experimental evidence demonstrates that TCF11 acts as a potent tumour-repressor relative to Nrf1α, albeit both isoforms possess a congruous capability to prevent tumour growth and upregulate those genes critical for improving the survival of patients with hepatocellular carcinoma.

Keywords: TCF11, Nrf1α, Nrf2, transcriptomic sequencing, regulatory profiling, hepatocellular carcinoma

Introduction

In all life forms, a variety of cell identifications with specialized topological shapes are evolutionarily selectively determined by diverse hub sets of transcription factors (TFs)-regulated gene expression profiles. Thereof, activation or inhibition of distinct transcription factors is essential for regulation of their target gene expression by binding a specific DNA sequence to maintain and perpetuate the normal and orderly operation of given organisms. In human, there exist over 1600 known and likely putative TFs; they have comprised nearly forty of distinct families[1]. Amongst them, the basic leucine zipper (bZIP) TFs consist of a larger group of the diverse basic-domain superfamily (in accordance with a new classification of TFs as described at http://www.edgar-wingender.de/huTF_classification.html). All bZIP proteins share a common bZIP domain structure, which is composed of a basic amino acid-enriched DNA-binding domain and an immediately-adjoining leucine zipper domain for distinct functional dimeration with cognate partners [2,3]. Of the bZIP factors, the Cap'n'collar (CNC) subfamily members are also characterized by a highly conserved 43-aa CNC domain, located N-terminally to the basic DNA-binding domain [4,5]. All CNC-bZIP orthologues share a highly conservatism of evolution from marine bacteria, ascidian, sea urchin, octopus, fly, hydra,

worm, bird, insect, fish, frog to mammals including human[6] for an indispensable role in executing cytoprotective transcriptional responses to changing environmental stress, and thus preserving the cellular homeostasis during development and growth of distinct life forms. As such, all other CNC-bZIP proteins, except SKN-1, alone can rarely serve as an exclusive transcriptional activator or repressor. This is due to the fact that they have to form a functional heterodimer with a sMaf (i.e., MAFG, MAFK and MAFF) [7,8] or other bZIP partners (e.g., JUN, JUND, FOS and ATF4) [9,10] in distinct combination for their specific DNA-binding demands [5,11,12]. Only such a CNC-bZIP heterodimer can be allowed for functional binding to target genes, containing one of specialized antioxidant/electrophile response elements (AREs/EpREs), Maf recognition element, activator protein 1-binding or other *cis*-regulatory homologous consensus sequences to drive distinct gene expression profiles and shape relevant biological functions.

Notably, Nrf1 (also called Nfe2l1, with its long TCF11 and short Nrf1β/LCR-F1 isoforms) and Nrf2 (also called Nfe2l2) are two principal CNC-bZIP transcription factors expressed in various cell types and tissues of mammals, including mouse and human [4,11]. Analysis of a neighbor-joining CNC-bZIP phylogenetic tree has unveiled that the membrane-bound Nrf1 orthologues should have emerged at a more ancient stage of the earlier evolution from marine bacteria to humans, and is thereby considered as a living fossil, than the water-soluble Nrf2[6]. This discovery supports a notion that Nrf1, than Nrf2, has a potent capability to fulfill more biological functions far beyond redox regulation that was originally identified. In fact, accumulating evidence reveals that Nrf1 exerts an important role in embryonic development [13,14], osteoblastogenes [15-17], life quality control of proteostasis by proteasome[18-20] and metabolism[21-24], anti-inflammatory immune response [25-27], and anti-tumour cytoprotection against hepatoma [28,29], in addition to redox stress defense [30-32]. Conversely, loss of Nrf1's function by gene-targeting in mice leads to severe oxidative stress and spontaneous development of distinct pathological phenotypes, resembling human non-alcoholic steatohepatitis (NASH) with progressive hepatoma, neurodegenerative diseases or diabetes mellitus[5]. By contrast, Nrf2 is dispensable based on the fact that animal development and growth are unaffected by its functional loss, without any pathological phenotypes. Such being the case, Nrf2 is still accepted as a master regulator of redox homeostasis [33,34], metabolism[35-37] and DNA repair [38,39] in order to meet the healthy needs of life. Rather, Nrf2 acts as a two-edged sword to shape significant biological functions in proliferation [40,41], resistance to apoptosis[42-44], angiogenesis [45-47], carcinogenesis [48-50] and invasion [51-53]. Overall, these demonstrate that Nrf1 and Nrf2 elicit similar yet different physiological functions. For an instance, our previous work revealed an inter-regulatory crosstalk between Nrf1 and Nrf2 at distinct levels [29], implying that they may regulate each other as a competitive player in similar biological process by distinct ways, but the details require to be further identified.

As a master of fact, so less attention has been paid on Nrf1 than Nrf2, though the indispensable Nrf1 is highly valued as a robust deterministic transcription factor and identified as an important endoplasmic reticulum (ER) sensor for changes in the intracellular redox, glucose, protein and lipid including cholesterol, status[54,55]. This is owing to the existing objective obstacles to be unresolved for the membrane-bound Nrf1, such that its functions are much less explored or verified than those of water-soluble Nrf2. Of note, a unique capability of single *Nrf1* gene confers it to be alternatively transcribed with alternative translational products; they are further subjected to selective splicing to yield multiple isoforms with different tempo-spatial topological properties. Consequently, distinct lengths of Nrf1 isoforms (e.g., TCF11, TCF11$^{\Delta N}$, Nrf1α, Nrf1β/LCR-F1) are differentially expressed in distinct type of cells, which makes diverse biological functions of Nrf1 more complicated [5,11,56]. In response to biological cues, the ER-resident Nrf1/TCF11 is

topologically dislocated from the lumen into extra-ER compartments, where their glycoproteins are deglycosylated and then subjected to selective juxtamembrane proteolytic processing to yield a mature factor before transactivating target genes (e.g., those encoding proteasomal subunits, antioxidant and cytoprotective proteins). Of note, a longer isoform than Nrf1α was referred to as TCF11 [57], consisting of 772 aa, but it is absent in the mouse, while the human prototypic Nrf1α of 742 aa lacks the Neh4L domain, due to alternative splicing of the *TCF11* transcript to remove its exon 4 (Figures 1A and S1A). As such, Nrf1α retains relative complete structural domains from Neh1L to Neh6L, of which equivalents exist in Nrf2 [5,58]. Thereby, it is postulated that Nrf1α and TCF11 should be two main players to exert differential transcriptional regulation of distinct Nrf1-target genes, but this remains to be proved.

To data, growing evidence indicates that Nrf1 and Nrf2, as two versatile leading players in maintaining cellular homeostasis, are essential for important pathophysiological processes in human diseases. Yet, which specific target genes regulated by both CNC-bZIP factors, and specific biological processes in which such genes are implicated, are required for further in-depth study, albeit two recent reports also revealed   different portions of between NRF-target expression profiles [59,60]. Herein, to refine distinct functions of Nrf1, it remains important to distinguish TCF11 and its N-terminally-truncated TCF11$^{\Delta N}$ (that can also occur naturally with the reminiscent Nrf2-like structural domains) from Nrf1α and Nrf2. As for this end, we have established four different cell lines stably expressing *TCF11, TCF11$^{\Delta N}$, Nrf1α* or *Nrf2*, respectively, by using an Flp-In™ T-REx™-293 system (as deciphered in Figure 1A). When required, this controllable system is turned on by tetracycline to induce each factor-specific transcriptional expression, while all others are unaffected in each of the indicated cell lines. Subsequently, transcriptomic sequencing of these cell modes unraveled that those common genes regulated by Nrf1 and Nrf2 are more responsible for regulating transcriptional expression and signal transduction in response to stimulus and diseases. The other differential expression genes regulated by Nrf1, but not Nrf2, are preferentially enriched in carbohydrate metabolism and cellular processes, while Nrf2-specific genes strikingly prefer to developmental process. Notably, Nrf1α and TCF11 share similar regulatory patterns, but they are disparate from those of Nrf2 or TCF11$^{\Delta N}$. Besides, TCF11 can also mediate rather more than target genes regulated by Nrf1α, displaying distinct biological functions in development, transport and catabolism. This notion is evidenced by our further supportive experiments, demonstrating that TCF11 serves as a potent tumour-repressor relative to Nrf1α, albeit both factors possess a congruous capability to prevent tumour growth, as accompanied by up-regulation of those target genes significantly improving the survival rate of patients with hepatocellular carcinoma (HCC).

Results

Controllable model cell lines for stably expressing *TCF11, TCF11$^{\Delta N}$, Nrf1α, or Nrf2* are established.

As shown in Figure 1A, four distinct expression constructs for *TCF11, TCF11$^{\Delta N}$, Nrf1α* or *Nrf2*, together with a recombinase Flp-expressing plasmid, were co-transfected into the Flp-In™ T-REx™-293 host cells and also integrated into this host cells by Flp-mediated homologous recombination at FRT sites. Then, putative positive cell clones stably expressing the indicated gene of interest (GOI) were selected and maintained by hygromycin B (150 μg/ml) and blasticidin (15 μg/ml) to establish these cell models for expression of *TCF11, TCF11$^{\Delta N}$, Nrf1α* or *Nrf2*. Such model cell lines are controllable because their GOI are tightly monitored by interaction of Tet (tetracycline) with its repressor TetR; thus, only after TetR will be released from the TetO$_2$ operator, transcriptional expression of GOI and down-stream target genes can be

induced. In subsequent parallel experiments, an additional cell line co-transfected with the empty expression vector served as an internal negative control. In order to further validate such stable controllable expression of *TCF11*, *TCF11^ΔN*, *Nrf1α* or *Nrf2,* respectively*,* all these indicated model cell lines were treated with 1 μg/ml Tet for 12 h and then determined by real-time quantitative PCR (Figure 1, B-D) and immunoblotting (Figure 1, E-G).

Figure 1. Establishment of four distinct model cell lines to stabilize expression of *TCF11*, *TCF11^ΔN*, *Nrf1α* and *Nrf2*. (A) A schematic diagram of the Flp-In™ T-REx™-293 system. The system allows the Flp recombinase-mediated homologous recombination of each indicated pcDNA5/FRT/TO-V5 expression constructs (for TCF11, TCF11^ΔN, Nrf1α or Nrf2) with the

Flp-In™ T-REx™-293 host cells through the FRT sites. (B-D) After incubation of *TCF11*, *Nrf1α*, *Nrf2* or *TCF11^{ΔN}*, as well as control cell lines with 1 μg/ml Tet for 12 h, total RNAs were isolated and then reversely transcribed into the first strand of cDNA. Subsequently, quantitative real-time PCR was employed to identify the mRNA expression levels of *Nrf1α* (B), *TCF11* (C), *TCF11^{ΔN}* (C) and *Nrf2* (D) in each of indicated cell lines. The data are shown as mean ± SEM (n = 3 × 3, $\$\$$, *p* < 0.01). (E-G) Total lysates of each cell line that had been treated with 1 μg/ml Tet (+) or not (−) were subjected to protein separation by SDS-PAGE gels, and then visualized by immunoblotting with distinct primary antibodies against V5, Nrf1 or Nrf2 to identify the protein levels of TCF11, Nrf1α, Nrf2 and TCF11^{ΔN}. (H-K) Total lysates of experimental cells, which had been induced with 1 μg/ml Tet for 12 h before being treated with CHX (50 μg/ml) alone or in combination with MG132 (10 μmol/L) for distinct times as indicated, were resolved by SDS-PAGE and then analyzed by Wester blotting with V5 antibody to identify the stability of TCF11 (H), Nrf1α (I), TCF11^{ΔN} (J) and Nrf2 (K) respectively.

Notably, a pair of specific TCF11-recognized primers, including part of the Neh4L-coding nucleotides, were designed by distinguishing it from *Nrf1α*, because the Neh4L-missing nucleotides of *Nrf1α* remains present in *TCF11* and also in its N-terminally-truncated *TCF11^{ΔN}*. Thus, the latter two factors-shared same primers were employed for quantitative PCR examinations of *TCF11* and *TCF11^{ΔN}*. As anticipated, real-time qPCR revealed that each specific transcriptional expression of *TCF11-*, *TCF11^{ΔN}*, *Nrf1α* or *Nrf2,* respectively, was induced by Tet in their indicated model cell lines (Figure 1, B-D). Such Tet-induced protein expression abundances of TCF11, TCF11^{ΔN}, Nrf1α and Nrf2 were evaluated by immunoblotting with antibodies against Nrf1/TCF11, Nrf2 and their C-terminally-tagged V5 epitope, respectively. The resulting data (Figure 1, E-G) demonstrated that those model cell lines had a strong capability to stably express each of interested genes, one of which had almost no effects on all the others examined, though each factor-specific expression was induced under Tet control. Of note, a major protein of TCF11 exhibited a slightly slower mobility, than that of Nrf1α, on electrophoretic gels, whereas the V5-tagged TCF11^{ΔN} mobility appeared to coincide closely to the electrophoretic band of Nrf2 (Figure 1E). Besides, a few of putative C-terminally-truncated isoforms of Nrf1, TCF11 or TCF11^{ΔN} were also immunoblotted with Nrf1/TCF11-specfic antibody (Figure 1F).

For further time-course analysis of *TCF11*, *TCF11^{ΔN}*, *Nrf1α* and *Nrf2*, their indicated cell lines that had been pre-treated for 12 h with 1 μg/ml Tet were treated with 50 μg/ml cycloheximide (CHX, that inhibits biosynthesis of nascent proteins) alone or plus a proteasomal inhibitor MG132 (at 10 μmol/L). As shown in Figure 1 (H-K), the N-terminal truncation of TCF11 (to remove both its ER-targeting sequence and adjacent juxtamembrane proteolytic degron) caused the resulting TCF11^{ΔN} isoform to become stabilized relatively. Subsequent calculation of their half-lives in CHX-treated cells suggested that a major TCF11^{ΔN} protein was conferred with a relative higher stability than those of TCF11, Nrf1 and Nrf2, each with distinct expression isoforms (as illustrated graphically in Figure 1, H-K). Upon co-treatment of cells with CHX and MG132, all those examined protein half-lives were markedly enhanced. Such being the case, intact TCF11 protein-A became gradually fainter and then disappeared by 2 h after co-treatment (Figure 1H). Such disappearance of TCF11 protein-A seemed to be accompanied by gradual emergence and increment of its protein-B and -C until stopping 8-h experiments. Similar yet different conversion of Nrf1α protein-A into short isoforms-B, -C and -D was observed (Figure 1I). However, no similar changes were determined in two cases of Nrf2 and TCF11^{ΔN}, because both proteins gradually decreased with increasing time of co-treatment until they finally disappeared from 4 h to 8 h after co-treatment with CHX and MG132 (Figure 1, J & K). Collectively, these demonstrate that not an ER-targeting sequence in TCF11^{ΔN} and Nrf2 allows them to display distinguishable behaviors from the ER-resident TCF11 and Nrf1α, both of which are manifested in similar, but different, topobiological processing within and around this organelle before being dislocated into the nucleus.

Differential expression profiles of genes regulated by TCF11, TCF11$^{\Delta N}$, Nrf1α and Nrf2 are defined.

To identify differential expression profiles of genes regulated by TCF11, TCF11$^{\Delta N}$, Nrf1α and/or Nrf2, relevant RNAs extracted from the established model cell lines were subjected to transcriptome sequencing. As a result, all those detectable genes, if upregulated or downregulated respectively with fold changes ≥ 2 or ≤ 0.5 plus false discovery rate (FDR) ≤ 0.001 (Figure 2A), were defined as differentially expressed genes (DEGs), by comparison with equivalents measured from control cells. Thereof, 2845 DEGs were detected in *TCF11*-expressing cells, of which 2786 target genes were upregulated by TCF11 (Figure 2A, and Table S1). By contrast, *Nrf1α*-expressing cells only yielded 1001 DEGs, i.e., 957 upregulated plus 44 downregulated (Table S2), whereas *Nrf2*-expressing cells led to a significantly decreased number of upregulated genes (i.e., 276) but as accompanied by down-regulation of 457 DEGs (Table S3). Notably, 1459 DEGs were identified in *TCF11$^{\Delta N}$*-expressing cells, with so many as 989 genes downregulated (Table S4). Interestingly, TCF11$^{\Delta N}$ appeared to be endowed with a regulatory trend of its target genes similar to that of Nrf2 (Figure 2). Such changed DEGs with distinct trends among different groups were further explicated by scatterplots of gene expression profiles (Figure S1B). Together, these data indicate that TCF11 makes a greater impact on overall gene expression than Nrf1α, albeit both contribute to basically positive regulation of their DEGs, whereas Nrf2 and TCF11$^{\Delta N}$ can also make a dominant negative contribution to their DEGs.

The intersections of these four groups of DEGs regulated by TCF11, TCF11$^{\Delta N}$, Nrf1α and/or Nrf2 were shown by the Venn diagram (Figure S1C). Those common DEGs between every two cell groups and each specific DEGs were taken into account (in *orthogonal table*, Figure 1B), with distinct regulatory tendencies of DEGs even in each subgroup. Of note, the common DEGs between *Nrf2*-expressing cells and the others were more likely to be downregulated rather than up-regulated by Nrf2, with a roughly similar number of its upregulated genes to down-regulated genes amongst *Nrf2*-specific DEGs. Another similar situation also occurred in *TCF11$^{\Delta N}$*-expressing cells.

Approximately 90 DEGs were identified to share in between all four cell lines (as shown in the Venn diagram, Figure S1C). Differential expression levels of these DEGs were presented by their heatmap with hierarchical clustering (Figure 1C). In the shared DEGs, only four genes *HMOX1* (heme oxygenase 1), *PCSK4* (proprotein convertase subtilisin/kexin type 4), *SALL3* (spalt like transcription factor 3) and *GAL* (galanin and GMAP prepropeptide) were upregulated or downregulated, with each of similar expression trends in all four cell lines. Most of other shared DEGs were up-regulated by TCF11 and Nrf1α, but down-regulated by Nrf2 and TCF11$^{\Delta N}$, besides two exceptions of *EMILIN2* (elastin microfibril interface 2) and *HSPB8* (heat shock protein family B member 8) regulated by opposite ways (Figure 1C). In addition, no evident effects of *Nrf2*-expressing cell model on *TCF11*, *TCF11$^{\Delta N}$* or *Nrf1α* was examined by transcriptome sequencing, but conversely, only a marginal increase in *Nrf2* expression in *Nrf1α-*, *TCF11-*, rather than *TCF11$^{\Delta N}$*-expressing cell models (Figure S1D). Amongst another CNC-bZIP members, only *BACH1* was up-regulated by TCF11, whilst all *sMAF* partners were up-regulated by Nrf2 and TCF11$^{\Delta N}$, but *Keap1* was unaffected.

In order to gain a further insight into similarities and differences in biological functions of between TCF11, TCF11$^{\Delta N}$, Nrf1α and Nrf2, their common and different regulatory DEGs were scrutinized in distinct combinations of every two groups as indicated by Groups A to L (Figure 2D). Similar or opposite trends in those common DEGs of every two cell groups were schematically shown. For an example of Group B, 121 common DEGs between *Nrf1α* and *Nrf2* were subdivided into 109 oppositely-regulated genes and the other 12 genes with the same directional tend. In Group E, all those common DEGs were manifested only

with the same directional trend to be regulated by both *Nrf2* and *TCF11^ΔN*. By contrast, a largely opposing expression trend in 619 of the common 671 DEGs co-regulated by *TCF11^ΔN* and *TCF11* and also in 946 of the other common 949 DEGs shared by *TCF11* and *Nrf1α* was shown in Groups H and K, respectively, but with two opposite regulatory directions. Such groups of these DEGs were also subjected to comprehensive analysis of their functional annotations as described below.

Figure 2. Statistical analysis of the data obtained from transcriptome sequencing. (A) Differentially expressed genes (DEGs) in distinct cell lines were analyzed by transcriptome sequencing, and relevant differences in the number of those increased or decreased DEGs are shown in the histogram. The DEGs were selected according to the following criteria: fold change ≥ 2 or ≤ 0.5 and FDR ≤ 0.001 (as compared to the Control cells). (B) The specific DEGs in each cell line and their common DEGs between every two cell lines were also counted as indicated in the chart, and the number of increased and decreased DEGs in each group is shown separately in black font, and the total is shown in white. In addition, the change trends of DEGs in each group were indicated in red or green, which represent up-regulated or down-regulated in the cells in the first row, respectively. (C) The heatmap with hierarchical clustering of 90 DEGs shared in all four cells lines. (D) Distinct groupings of the subsequent functional annotation and also the Venn diagram of DEGs between every two cell lines.

Nrf1α and Nrf2 have diverse regulatory patterns of their target genes.

In Group A, 880 DEGs were identified in *Nrf1α*-expressing rather than *Nrf2*-expressing cells, and further subjected to their functional annotation by the DAVID (*d*atabase for *a*nnotation, *v*isualization and *i*ntegrated *d*iscovery) (Figure 3A), for the data mining in order to delineate unique biological functionality

Figure 3. Functional annotation of specific or common DEGs in *Nrf1α* and *Nrf2* cells. (A) The top 10 of significant biological process terms and pathways enriched by DEGs in Groups A, B, and C were exhibited in histograms and scatterplots, respectively. (B) After *Nrf1α*, *Nrf2* and Control cell lines were incubated with 1 μg/ml Tet for 12 h, total RNAs were isolated and reversely transcribed into the first strand of cDNA. Subsequently, the mRNA levels of DEGs that were associated with more functions, along with high expression levels and well significance in Groups A to C, were determined by quantitative real-time PCR analysis of *Nrf1α*, *Nrf2* and Control cell lines. The data are shown as mean ± SEM (n = 3 × 3, * $p < 0.05$; ** $p < 0.01$; $, $p < 0.05$; $$, $p < 0.01$).

of Nrf1α in regulating genes preferentially than Nrf2. Besides their shared common 121 DEGs in Group B, the other 612 DEGs in Group C were also annotated to identify those biological functions of Nrf2 that were responsible preferentially than Nrf1α (Figure 3A, and Table S5). The details of all top significant biological process terms and pathways enriched by DEGs in three different groups were deciphered in histograms and scatterplots (Figure 3A). Furthermore, the biological process terms and pathways were classified by using QuickGO and KEGG databases. Such functional annotation analysis implied that DEGs regulated by Nrf1α (Group A) were predominantly involved in cellular metabolic process, response to stimulus, replication and repair, cell growth and death, folding, sorting and degradation, signal transduction, immune system and cancers. In Group B, DEGs co-regulated by Nrf1α and Nrf2 took part in distinct cellular metabolic process, RNA processing, regulation of biological process, response to stimulus, apoptotic process, regulation of transcription, translation and cancers. Nrf2-regulated DEGs (in Group C) were also responsible for cellular metabolic process, apoptotic process, response to stimulus, regulation of transcription, development and regeneration, signal transduction, endocrine system, infectious diseases and cancers.

Based on certain association with multiple functions of target genes with higher expression levels and their significance, 19 DEGs were selectively verified by further quantitative PCR analysis (Figure 3B). The results demonstrated that 4 DEGs, including *BARD1* (*BRCA1 associated RING domain 1*), *CASP3* (*caspase 3*), *HSP90AA1* (*heat shock protein 90 alpha family class A member 1*) and *BPTF* (*bromodomain PHD finger transcription factor*) were upregulated by Nrf1α, but not significantly affected by Nrf2. By contrast, 9 DEGs, including *BLM* (*Bloom syndrome, RecQ like helicase*), *CCNG1* (*cyclin G1*), *PALB2* (*partner and localizer of BRCA2*), *STAG2* (*stromal antigen 2*), *DICER1* (*dicer 1, ribonuclease III*), *PNN* (*pinin, desmosome associated protein*), *RB1* (*RB transcriptional corepressor 1*), *SMARCA1* (*SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 1*) and *SOS1* (*SOS Ras/Rac guanine nucleotide exchange factor 1*) were upregulated by Nrf1α, but downregulated by Nrf2. Another 6 DEGs, such as *FOXO1* (*forkhead box O1*), *KRAS* (*KRAS proto-oncogene, GTPase*), *MAP2K6* (*mitogen-activated protein kinase kinase 6*), *NF1* (*neurofibromin 1*), *NRP1* (*neuropilin 1*) and *TGFBR1* (*transforming growth factor beta receptor 1*) were reduced in *Nrf2*-expressing cells but, but no significant changes of them was detected in *Nrf1α*-expressing cells. In addition, putative functions of the examined genes as well as of their encoding proteins were also extracted (Figure 3B, *on the bottom*).

TCF11$^{\Delta N}$ exhibits a similar regulatory profile to that of Nrf2

All those DEGs regulated by *Nrf2* or *TCF11$^{\Delta N}$* alone or both were divided into three Groups D, E and F, respectively, and then functionally annotated with the above-described methods (Figure 4A and Table S6). In Group D, 194 DEGs regulated by Nrf2, but not by TCF11$^{\Delta N}$, were generally involved in cellular metabolic process, localization, regulation of biological process, developmental process, response to stimulus, signal transduction and cancers. In the intersected Group E, 539 DEGs co-regulated by *Nrf2* and *TCF11$^{\Delta N}$* were also associated with cellular metabolic process, development and regeneration, regulation of transcription, response to stimulus, apoptotic process, endocrine system, signal transduction, infectious diseases and cancers. In Group F, 920 DEGs regulated by *TCF11$^{\Delta N}$* were significantly enriched in cellular metabolic process, cell motility, cellular community, developmental process, regulation of transcription, signal transduction, signaling molecules and interaction, cardiovascular diseases and cancers. Intriguingly, 8 common biological process terms and additional 8 common pathways were predicted to exist between

the top 10 functions significantly enriched in Group C (including DEGs regulated by *Nrf2* but not by Nrf1α) and Group E (with an intersection of DEGs shared by *Nrf2* and *TCF11ΔN*) (Figure S1E). This implies that Nrf2 and TCF11ΔN share the common regulatory profiles, but they are likely differential from those of Nrf1α and Nrf2.

Figure 4. Functional annotation of specific or common DEGs in *Nrf2* and *TCF11ΔN* cells. (A) Top 10 of significant biological process terms and pathways enriched by DEGs in Groups D, E, and F were exhibited in histograms and scatterplots, respectively. (B) After induced with 1 μg/ml Tet for 12 h, total RNAs were isolated from *Nrf2*, *TCF11ΔN* or Control cell lines and then reversely transcribed into the first strand of cDNA. Subsequently, the mRNA levels of DEGs that were associated with more functions as annotated, along with high expression levels and well significance in Groups D to F, were determined by quantitative real-time PCR analysis of *Nrf2*, *TCF11ΔN* and Control cells. The data are shown as mean ± SEM (n = 3 × 3, * *p* < 0.05; ** *p* < 0.01; $, *p* < 0.05; $$, *p* < 0.01).

The results of quantitative PCR validation (Figure 4B) revealed that expression of *FGF21* (*fibroblast growth factor 21*), *FOSB* (a *subunit of AP-1 transcription factor*) and *JUNB* (*another subunit* of *AP-1*) were upregulated by Nrf2 and TCF11ΔN, with downregulation of *MAP2K6* (*mitogen-activated protein kinase*

kinase 6) and *NRP1* (*neuropilin 1*). By contrast, *GLI1* (*GLI family zinc finger 1*) was upregulated by Nrf2, but not significantly altered by TCF11$^{\Delta N}$. Conversely, *TGFBR1* (*transforming growth factor beta receptor 1*) was downregulated by Nrf2, but upregulated by TCF11$^{\Delta N}$. However, reduced expression of *CAV1* (*caveolin 1*) was accompanied by increased *COL6A2* (*collagen type VI alpha 2 chain*) in TCF11$^{\Delta N}$-expressing cells, but almost unaffected by Nrf2. In addition, putative functions of such target genes were mapped, as indicated by the histogram (Figure 4B, *on the bottom*).

TCF11 and its truncated TCF11$^{\Delta N}$ regulate similar yet different subsets of target genes.

The common and distinct target DEGs in *TCF11$^{\Delta N}$*- and/or *TCF11*-expressing cell lines were assigned into three groups, and functionally annotated with the aforementioned method as visualized in bar charts

Figure 5. Functional annotation of specific or common DEGs in *TCF11$^{\Delta N}$* and *TCF11* cells. (A) Top 10 of significant biological process terms and pathways enriched by DEGs in Groups G, H, and I were exhibited in histograms and scatterplots, respectively. (B) After induced with 1 μg/ml Tet for 12 h, total RNAs were isolated from Control, *TCF11$^{\Delta N}$* or *TCF11* cell lines before being reversely transcribed into the first strand of cDNA. Subsequently, relative mRNA levels of DEGs that were associated with more functions, along with high expression levels and well significance in Groups G to I, were determined by quantitative real-time PCR in Control, *TCF11$^{\Delta N}$* and *TCF11* cells. The data are shown as mean ± SEM (n = 3 × 3, * $p < 0.05$; $, $p < 0.05$).

and scatterplots (Figure 5A, and Table S7). In Group G, 788 DEGs were identified as targets of TCF11$^{\Delta N}$, but not of TCF11, and enriched with distinct functions in cellular metabolic process, regulation of biological process, cellular community, apoptotic process, developmental process, development and regeneration, response to stimulus, signal transduction, signaling molecules and interaction, infectious diseases and cancers. Their commonly-shared 671 DEGs in Group H were functionally responsible for cellular metabolic process, cell cycle, cell motility, signal transduction, protein folding, sorting and degradation, transport and catabolism, endocrine system and cancers. In Group I, 2174 DEGs were regulated preferentially by *TCF11*, rather than by *TCF11$^{\Delta N}$*, and functionally associated with cellular metabolic process, cell growth and death, replication and repair, response to stimulus, signal transduction, protein folding, sorting and degradation, regulation of transcription, translation, transport and catabolism, carbohydrate metabolism and infectious diseases. Notably, 6 identical pathways were found by comparison between top 10 significantly enriched pathways from Group C (i.e., DEGs regulated by *Nrf2* but not by Nrf1α) and Group G (i.e., DEGs regulated by *TCF11$^{\Delta N}$* but not by TCF11) (Figure S1E). This indicates that TCF11$^{\Delta N}$-regulated genes are much likely to execute somewhat combinational or overlapping functions with Nrf2-target genes.

Subsequently, several unique or common target genes regulated by TCF11 and/or TCF11$^{\Delta N}$ were also validated by quantitative PCR (Figure 5B). The results showed that *FOSB*, *JUNB* and *NFKB2* (*nuclear factor kappa B, subunit 2*) were upregulated, while *WNT5A* (*Wnt family member 5A*) was downregulated, by TCF11$^{\Delta N}$, but unaffected by TCF11. Conversely, *COL4A6* (*collagen type IV alpha 6*) and *HIF1A* (*hypoxia inducible factor 1 alpha*) were upregulated by TCF11, but downregulated by TCF11$^{\Delta N}$. Besides, *ATM* (*ATM serine/threonine kinase*) was also upregulated by TCF11, but roughly unaltered by CF11$^{\Delta N}$. The putative functions relative to these examined genes were exhibited (as shown in Figure 5B, *on the bottom*).

TCF11 and Nrf1α display similar but differential regulatory profiles

Those DEGs regulated by TCF11 and/or Nrf1α were grouped by J to L, and then functionally annotated by DAVID, with histograms and scatterplots exhibited (Figure 6A, and Table S8). Comprehensive analysis of the top significantly enriched biological process terms and pathways showed that 1896 DEGs of Group J, by identifying *TCF11-*, but not *Nrf1α -,* expressing cells, were associated with distinct functions in cellular metabolic process, cell cycle, subcellular localization, transport and catabolism, carbohydrate metabolism, regulation of transcription, translation, signal transduction, endocrine system, development and regeneration, infectious diseases and cancers. In Group K, TCF11 and Nrf1α co-regulated 949 DEGs that were involved in cellular metabolic process, cell growth and death, replication and repair, folding, sorting and degradation, regulation of transcription, response to stimulus, endocrine system, immune system and cancers. In Group L, only 52 DEGs were identified by transcriptional regulation by Nrf1α, but unaffected by TCF11 expression. Their putative functions were associated with cellular community and metabolic process, and subcellular localization, regulation of biological process, signal transduction and response to stimulus. Besides, many of overlapping functions were predicted to exist among distinct

combinations of top significantly enriched functions exerted by Group A (i.e., DEGs regulated by Nrf1α but not by Nrf2), Group I (i.e., DEGs regulated TCF11 but not by TCF11^ΔN) and Group K (i.e., DEGs co-regulated by TCF11 and Nrf1α) (Figure S1E). Thus, it is inferable that TCF11 and Nrf1α displays similar regulatory profiles, with a striking disparity from those of TCF11^ΔN or Nrf2.

Figure 6. Functional annotation of specific or common DEGs in *TCF11* and *Nrf1α* cells. (A) Top 10 of significant biological process terms and pathways enriched by DEGs in Groups J, K, and L were exhibited in histograms and scatterplots, respectively. (B) After induced with 1 μg/ml Tet for 12 h, total RNAs were isolated from Control, *TCF11* or *Nrf1α* cell lines and then reversely transcribed into the first strand of cDNA. Subsequently, relevant mRNA levels of DEGs that were associated with more functions as annotated, along with high expression levels and well significance in Groups J to L, were determined by quantitative real-time PCR in Control, *TCF11* and *Nrf1α* cells. The data are shown as mean ± SEM (n = 3 × 3, ** $p < 0.01$; $, $p < 0.05$; $$, $p < 0.01$).

Amongst the above DEGs, 14 were selected for further quantitation by real-time PCR (Figure 6B). The results revealed that 8 DEGs were upregulated by both TCF11 and Nrf1α, which included *ATF2* (*activating transcription factor 2*), *BRCA1* (*BRCA1 DNA repair associated*), *HSP90AA1* (*heat shock protein 90 alpha family class A member 1*), *HSP90B1* (*heat shock protein 90 beta family member 1*), *RAD50* (*RAD50 double strand break repair protein*), *SMC3* (*structural maintenance of chromosomes 3*), *SOS1* (*SOS Ras/Rac guanine nucleotide exchange factor 1*) and *USP16* (*ubiquitin specific peptidase 16*). Additional 4 DEGs, i.e., *ARFGAP3* (*ADP ribosylation factor GTPase activating protein 3*), *IMPA1* (*inositol monophosphatase 1*), *PRKCE* (*protein kinase C epsilon*) and *PTEN* (*phosphatase and tensin homolog*) were upregulated by TCF11, but almost unaffected by Nrf1α. Conversely, *Nrf1α* expression caused a decrease in *CGREF1* (*cell growth regulator with EF-hand domain 1*) as accompanied by increased *ICAM5* (*intercellular adhesion molecule 5*), but such both genes were unaltered by TCF11. These examined genes were responsible for their putative functions as indicated (Figure 6B, *on the bottom*). Altogether, TCF11 and Nrf1α exhibit a similar regulatory profile, but with some of quietly different target genes.

TCF11 is a more potent player than Nrf1α at preventing tumour xenografts in nude mice.

As analyzed above, distinct subset of DEGs regulated by Nrf1, TCF11 and/or Nrf2 were annotated for their functional relevancies to cancer development or prevention. In fact, our previous work had revealed that knockdown of Nrf1 caused a significant malignant growth of subcutaneous tumour xenografts in nude mice [61]. Thereof, Nrf1α was indicated to act as a dominant tumour-repressor insomuch as to confine oncogenicity of Nrf2 [28,29]. Herein, to corroborate the putative tumour-preventing effects of Nrf1α and TCF11, both CNC-bZIP factors were restored by transfecting the lentivirus expression constructs into HepG2 cells with a specific loss of *Nrf1α*, respectively, as described elsewhere [28]. As a consequence, the resulting *Nrf1α-* or *TCF11*-restored cell lines were confirmed by quantitative PCR and immunoblotting to be definitely true (Figure 7A).

Subsequently, both *Nrf1α-* and *TCF11*-restored cell lines, alongside with *Nrf1$\alpha^{-/-}$* cells and wild-type (*WT*) HepG2 cells, were heterotransplanted into distinct groups of those immunodeficient nude mice at their subcutaneous loci as indicated. After tumour formation, the sizes of growing tumours were measured at one-day intervals within ensuing five weeks before the tumour-bearing mice were sacrificed. The resulting data were calculated and shown graphically (Figure 7B), as a consequence demonstrating that restoration of *Nrf1α* or *TCF11* enables a significant tumour-preventing effect on the subcutaneous human carcinoma xenografts in nude mice, when compared with those obtained from *Nrf1$\alpha^{-/-}$* and WT cell lines. In addition, it should be also noted that both *Nrf1$\alpha^{-/-}$* and WT hepatoma cell lines, as reported previously [28], were herein used as only two co-references in this parallel animal experiments to strengthen their comparability (Figure 7B). Furthermore, a series of comparative experiments revealed that *Nrf1$\alpha^{-/-}$* cell proliferation, migration and invasion were all significantly suppressed by restoration of *Nrf1α* and *TCF11* (Figure 7, C-H), and the cell-cycle arrest at S-phase, along with the increase in early apoptosis, due to the restoration of them (Figure S2). Collectively, TCF11 acts as a more potent tumour-repressor than Nrf1α, albeit both isoforms are endowed with an intrinsic capability to prevent tumour development and malgrowth.

Figure 7. Malgrowth of *Nrf1α−/−*-derived hepatoma cells were significantly suppressed by restoration of *Nrf1α* and *TCF11*.
(A) Both quantitative real-time PCR (*up*) and Western blotting (*down*) were employed to identify the protein and mRNA levels of Nrf1α and TCF11 in *Nrf1α*- and *TCF11*-Restored hepatoma cells. The experimental cells had been treated with or without 10 μmol/L MG132 for 4 h before being harvested for Western blotting. The data are shown as mean ± SEM (n = 3 × 3, ** $p < 0.01$; $$, $p < 0.01$). (B) Differences in mouse subcutaneous xenograft tumours derived from wild type HepG2 (*WT*), *Nrf1α/-*, *Nrf1α*-Restored and *TCF11*-Restored cells were measured in size every two days, before being sacrificed on the 42nd day. The data are shown as mean ± SD (n = 7 per group, * $p < 0.05$; ** $p < 0.01$; $, $p < 0.05$; $$, $p < 0.01$). (C-H) The soft agar colony formation (C, D), as well as migration (E, F) and invasion (G, H), of wild type HepG2 (WT), *Nrf1α/-*, *Nrf1α*-Restored and *TCF11*-Restored cells were examined as described in 'Materials and Methods'. The data are shown as mean ± SD (n = 9, * $p < 0.05$, ** $p < 0.01$).

Nrf1α and TCF11 regulate critical genes for improving the survival rate of HCC patients.

To weigh the practical effects of TCF11, Nrf1α and Nrf2 on human hepatocellular carcinoma (HCC), their relevancies to the overall survival (OS) of HCC patients were firstly investigated (Figure 8A). For this,

Figure 8. Diverse effects of Nrf1 and Nrf2 on those HCC-relevant proteins that are significantly correlated with the overall survival (OS). (A) The correlation analysis between of hepatocellular carcinoma (HCC) associated genes and the relevant OS rates of patients with HCC. (B) The protein expression levels of HCC-associated proteins in *Nrf1α, TCF11, TCF11^ΔN* and *Nrf2* cell models were also examined by Western blotting analysis of these experimental cell lines that had been induced with or without 1 μg/ml Tet for 12 h before being harvested. (C) Both wild type HepG2 and its derived *Nrf1α^-/-* cells were transfected with either Nrf1α or TCF11 expression plasmids, and then allowed for 24-recovery from transfection in the fresh medium before being subjected to Western blotting, to identify the protein levels of HCC-associated proteins. (D) Both wild type HepG2 and its derived *Nrf2^-/-* cells were transfected with an Nrf2 expression plasmid, and then allowed for 24-recovery from transfection in the fresh medium before being subjected to Western blotting. (E) MHCC97L or MHCC97H cells were transfected with each of Nrf1α, TCF11 and Nrf2 expression constructs, and then allowed for 24-recovery from transfection in the fresh medium before being examined by Western blotting to identify abundances of HCC-associated proteins as described above.

HCC-relevant molecules were selected from the UniProt database and their genes were further parsed by the Kaplan-Meier Plotter[62] to find those markers of being significantly correlated with OS of patients with HCC. As shown in Figure 8 (*A1-A12*), a better OS was predicted to couple with increased expression of Nrf1, *CHP2* (*calcineurin like EF-hand protein 2*), *CPS1* (*carbamoyl-phosphate synthase 1*), *FOXO1* (*forkhead box O1*), *IRS4* (*insulin receptor substrate 4*) and/or *NKX2-8* (*NK2 homeobox 8*); this was also accompanied by reduced expression of *AKR1B10* (*aldo-keto reductase family 1 member B10*), *EPO* (*erythropoietin*), *MUTYH* (*mutY DNA glycosylase*) and/or *PKM* (*pyruvate kinase M1/*2). Besides, GP73 (also called GOLM1, golgi membrane protein 1) has been shown as a potential diagnostic marker for primary HCC [63,64], while GPC3 (glypican 3) acts as another key biomarker for early diagnosis of human HCC and a rational immunotherapeutic target for HCC [65,66].

The immunoblotting results showed that TCF11 and Nrf1α enabled the normal cells to increase CHP2 and CPS1 abundances, respectively (Figure 8, *B3* & *B4*). Although FOXO1 was slightly increased, NKX2-8 was markedly increased, by both TCF11 and Nrf1α (*B5* & *B7*), but IRS4 was down-regulated by TCF11 but not Nrf1α (*B6*). Furtherly, AKR1B10 and EPO were, to greater or less extents, diminished by TCF11 and Nrf1α (*B8* & *B9*). Notably, significant decreases in the two HCC biomarkers GP73 and GPC3 were caused by TCF11 and Nrf1α (*B12* & *B13*). By sharp contrast, most of the above-examined proteins were reduced by TCF11$^{\Delta N}$ and Nrf2 (Figure 8B, *right panels*), with an exception that GPC3 was significantly up-regulated by Nrf2, but not TCF11$^{\Delta N}$.

Further examinations revealed that overexpression of TCF11 and Nrf1α in HepG2 cells enabled CHP2, CPS1, FOXO1, IRS4 and NKX2-8 to be enhanced to varying extents, but they were markedly suppressed by loss of *Nrf1$\alpha^{-/-}$* (Figure 8C, *left panels*). Such loss of *Nrf1α* also gave rise to constructive enhancement of Nrf2, AKR1B10 and GPC3, as accompanied by constructive abolishment of EPO and GP73, besides CHP2 and NKX2-8. However, it is intriguing to note that all these constructive changes could not be ameliorated by modest restoration of ectopic TCF11 or Nrf1α into *Nrf1$\alpha^{-/-}$* cells (Figure 8C, *right panels*). By contrast, forced expression of Nrf2 in HepG2 cells caused only a marginal increase in FOXO1 or GP73, but the other examined genes were unaffected (Figure 8D, *left panels*). Conversely, loss of *Nrf2* caused a slight increase in CHP2 and GPC3; but the former CHP2 was reduced by ectopic Nrf2 restoration to its basal levels, while the latter GPC3 was not mitigated by ectopic Nrf2 (Figure 8D, *right panels*).

The above-described data indicate distinct contributions of TCF11, Nrf1α and Nrf2 to differential or opposite regulation of endogenous expression levels of different, even the same, target genes. This notion was herein substantiated by our further comparative experiments of TCF11, Nrf1α and Nrf2 that had been transfected for their respective overexpression in either MHCC97L or MHCC97H cell lines. As anticipated, abundances of CHP2, CPS1, FOXO1, IRS4, NKX2-8, MUTYH, and GP73 proteins were increased with their respectively-varying trends by ectopic TCF11 or Nrf1α, but less affected by Nrf2, expression in MHCC97L cells (Figure 8E, *left panels*). Such events did not occur in MHCC97H cells, even though TCF11, Nrf1α and Nrf2 were overexpressed (Figure 8E, *right panels*). Besides, both EPO and GPC3 were suppressed by TCF11, Nrf1α and Nrf2 in MHCC97L cells, but almost unaffected in MHCC97H cells with hyper-expression of Nrf2 and AKR1B10. Such discrepancies in these gene expression are likely contributable to distinct metastatic potentials and other malignant properties of between MHCC97L and MHCC97H cell lines. Taken together, it is demonstrated that Nrf1α and TCF11, but not Nrf2, are conferred for an intrinsic capability to regulate those genes critical for improving the survival rate of patients with HCC, such that Nrf1α or TCF11 perform a strikingly disparate effect from that of Nrf2 on human hepatoma (Figure 9).

Figure 9. Relationship and difference between the regulation patterns of Nrf1 and Nrf2 on their targets. (A) The functional protein association networks of targets regulated by Nrf1 or Nrf2. Of note, the protein-protein associations are determined by various ways, which are thus represented by different colored edges as indicated. (B) The heatmap of the sequencing expression of genes, which are composed of the network, with distinct expression levels in *TCF11*, *Nrf1α*, *Nrf2* and *TCF11^{ΔN}* cell lines. The color of the nodes in the heatmap represents the value of $\log_2$ (fold change) as shown in the color bars, indicating the gene expression trend as compared with the control group (upregulation or downregulation, were marked in red or green, respectively). (C) The heatmap of the sequencing expression of genes in *Nrf1α^{-/-}* and *Nrf2^{-/-}* cell lines. The color of the nodes in the heatmap represents the value of $\log_2$ (fold change) as shown in the color bars, indicating the gene expression trend as compared with the wild-type HepG2 cells. (D) A comprehensive regulatory model is proposed to reveal the different effects of Nrf1 and Nrf2 on hepatoma. In addition, the *Hox* hub abutting the distinct *Nrf* loci was indicated (*at the low left corner*).

Discussion

In this study, we have established four controllable cell models for stably expressing *TCF11*, *TCF11^{ΔN}*, *Nrf1α* or *Nrf2*, which all occur naturally with their respective intact structural domains, albeit they were tagged C-terminally by a neutral V5 epitope. Of striking note, Nrf1α and TCF11 are two longer isoforms of the ER-resident Nrf1 with changing gears in their overall transcriptional activity, because both proteins are tightly controlled by their unique topobiological rheostat modules from the ER to enter the nucleus. For an instance, the NST domain transactivation of Nrf1α/TCF11, as well as SKN-1 and CncC, is required for re-editing of their indicated asparagines to aspartates during dynamic dislocation from the lumen to extra-ER subcellular compartments[67,68]. Upon lack of such amino acid re-editing by peptide:N-glycosidase (PNG1), this causes an evident decrease in the rheostat capacity of Nrf1α/TCF11,

and even loss of *PNG1* or its mutations results in inactivation of Nrf1α/TCF11, manifesting inflammation and adrenal insufficiency[69-71]. Furthermore, rheostat capacity of Nrf1α/TCF11 is also monitored by its reversible ubiquitination and deubiquitination during its selective ER-associated proteasome-regulated processing to multiple isoforms[72,73].

During our manuscript preparation, removal of the N-terminal ER signal peptide-containing 104-aa or 121-aa regions from Nrf1α/TCF11 to yield two artificially-truncated mutants (i.e., Nrf1$^{\Delta N104}$ or Nrf1$^{\Delta N121}$) had been reported by both independent Bollong's and Ooi's laboratories[59,60]. It is, to our great surprise, seen that such artificial mutants Nrf1$^{\Delta N104}$ or Nrf1$^{\Delta N121}$ were asserted to represent two constitutive processed Nrf1 activators, although they each retain a negative PEST region. Thereby, our study of *TCF11$^{\Delta N}$* showed an N-terminal 2-156 residues-truncated mutant (i.e., TCF11$^{\Delta N2-156}$), which may also arise from a naturally-splicing transcript[58]. However, TCF11$^{\Delta N}$ is herein identified as a mimic Nrf2 factor, because both share conserved structural domains (Figure 1A). This seems to totally contradict the purpose of Nrf1$^{\Delta N104}$ or Nrf1$^{\Delta N121}$ by both Bollong's and Ooi's colleagues[59,60]. Additional two mutants of caNrf2 to block its Keap1-binding activity were also utilized in their experimental settings. Besides, it should be of crucial importance to note that a strong acidic property of 3×Flag, wherever it was tagged at the N-terminal or C-terminal ends of a given protein, exerts distinctive or even opposing effects on its topobiological folding and dynamic moving in and out of membranes and other subcellular compartments[72,74]. Overall, our study has provided four cell tools to evaluate similarities yet differences in between intact TCF11, TCF11$^{\Delta N}$, Nrf1α and Nrf2-regulatory profiles, which are stimulated at the Tet-inducibly controllable levels.

Subsequently, these four cell lines were determined by transcriptomic sequencing in an integrative combination with routine reductionist approaches. As expected, comprehensive functional annotation of TCF11, TCF11$^{\Delta N}$, Nrf1α and Nrf2 unraveled that they are likely to perform diverse biological functions by combinationally overlapping or competitively opposing modules within distinct cellular networks, apart from their unique functions. Thereby, a further understanding of their respectively regulated targets and functions should be achieved only by making one of them function alone. For this end, our established cell lines stably expressing *TCF11*, *TCF11$^{\Delta N}$*, *Nrf1α* or *Nrf2* respectively, are an invaluable tool to gain insights into distinguishing their regulatory profiles, while they are allowed to function alone. Consequently, the regulatory patterns of Nrf1α and TCF11 are similar yet different, which are strikingly disparate from that of Nrf2, even with certain opposite effects on the same targets. Such distinctions between Nrf1α/TCF11 and Nrf2 are attributable to differences in their primary structures and functional subcellular locations. This is due to the objective fact that the ER-associated domains of Nrf1α/TCF11 determine its unique membrane topobiology and post-synthetic processing mechanisms, which enables it to be distinguishable from Nrf2. Yet, once the ER-targeting signal peptide-adjoining region were deleted to yield TCF11$^{\Delta N}$, this N-terminally truncated isoform does exhibit a similar regulatory pattern to that of Nrf2. However, this notion appears to totally contradict the conclusions drawn from two so-called constitutive activators Nrf1$^{\Delta N104}$ or Nrf1$^{\Delta N121}$ by Bollong's and Ooi's groups[59,60]. As such, their analyses by combining ChIP-seq and RNA-seq data revealed that Nrf1$^{\Delta N121}$ (or Nrf1$^{\Delta N104}$) had 3.19-fold numbers of target-binding peaks than those of caNrf2-binding targets. Of note, over 75% of Nrf1$^{\Delta N121}$-binding targets are focused preferentially on the consensus ARE sequence (5′-TGAC/GnnnGC-3′) flanked by AT-enriched motifs, while only less than 45% of caNrf2-binding targets are recruited on ARE-like sequences flanked by GC-enriched motifs[59]. It is inferable that Nrf1 exerts its unique biological functions by predominantly regulating ARE-driven cognate

genes, whereas Nrf2 has a widely-varying capacity to elicit its promiscuous roles in diversely mediating non-ARE-battery genes. Altogether, with our previous work[29,55] demonstrating that both Nrf1α/TCF11 and Nrf2 can mutually influence each other by inter-regulating at distinct levels, this leads to the formation of a steady-state regulatory network system finely monitored by a negative feedback loop to maintain the cell homeostasis and organ integrity.

Since early discovery of Nrf2 required for regulating antioxidant and detoxification genes[75], the overwhelming majority of investigations in this field have been focused disproportionately on this CNC-bZIP factor and its negative regulator Keap1 in response to redox stress, but also provided myriad insights into its promiscuous roles in biology and medicine, as well as drug development[76-78]. However, aside from these studies by routine reductionist approaches, the whole genome-widely integrative analyses uncovered the underlying facts that Nrf1, but not Nrf2, is essentially required for regulating important homeostatic and metabolic genes involved in the normal growth and development throughout life process. This is fully consistent with the experimental evidence showing that loss of Nrf1 results in mouse embryonic lethality and also causes adult pathological phenotypes varying within its gene-targeting mutant organs (reviewed by[73]). Such bad consequences are attributable to severe endogenous oxidative stress and fatal defects in redox metabolism reprogramming and relevant constitutive gene expression profiles of *Nrf1*-deficient cells (this study and[21]). Collectively, these facts demonstrate that Nrf1 is a predominant determinist factor of cellular constitutive redox metabolic homeostasis with organ integrity. That is to say, Nrf1 has a potent capacity to contribute to the steady-state robustness of cell physiological homeostasis. By contrast, Nrf2 is much likely to make a major contribution to the sensitive plasticity of cell homeostasis, because it has been accepted as a master regulator in response to diverse stresses. This is also supported by the facts of no obvious phenotypes resulting from global loss of *Nrf2* in mice, but its deficiency leads to more susceptibility to chemical carcinogens and diverse stresses than wild-type controls [79,80].

Such differential contributions of Nrf1 and Nrf2 to cell homeostasis robustness and plasticity are also likely resulted from a striking evolutionary conservation of their CNC-bZIP family in distinct species during nature selection. Comparative genomics analyses revealed that, although this CNC-bZIP family expansion and diversification have occurred in vertebrates, Nrf1 is viewed as a living fossil that is more ancient than Nrf2, because it is conserved closely to the only CNC in *Drosophila melanogaster*, SKN-1 in *Caenorhabditis elegans* and each of newly-identified Nach factors in simple multicellular eukaryotes, but not in unicellular protozoans or other prokaryotes beyond *Endozoicomonas*[6]. This implies that they have evolved for multicellular cooperative selection. For instance, the *C. elegans* SKN-1, like Nrf1, is selectively processed by post-transcriptional and post-translational ways to yield distinct isoforms, one of which is N-terminally truncated (by analogy of TCF11$^{\Delta N}$) but manifested Nrf2-like functions, albeit with no existence of Keap1-like orthologue[68,81]. The evolutional conservativity of Nrf1, but not of Nrf2, strongly demonstrates that it is indispensable for basal constitutive contributions to orchestrating the ensemble of critical gene regulatory networks, such that the normal cell homeostasis and organ integrity have been perpetuated in the entire course of life.

In fact, the CNC-bZIP family members (p45, Nrf1, Nrf2 and Nrf3) are, though diversified in vertebrates, topologically organized together with the developmental *Hox* gene clusters (as shown in Figure 9D), upon forming the PcG hub to maintain their genes in a silent-but-poised state[82]. A recent ChIP data analysis revealed that most of Nrf1-binding sites are focused closely to the canonical ARE sequences that are

widely located in its cognate gene promoter and distal intergenic enhancer regions (i.e., dynamic genome-topology as a functional primer is built by spatial interactions of distal enhancers with the proximal promoters in the same genes or even between different genes in chromosomal architecture), whereas Nrf2-binding sites are promiscuously loosed to ARE-like or no-ARE sequences that are, however, constrained narrowly in its target gene promoters[59]. Collectively, the interplay between selective transcription factor (e.g., CNC-bZIP)-regulated gene programming and genome conformation is surmised as a driving force for cell-fate decision with distinct type-featured identifications.

Notably, the versatile Nrf2 acts *de facto* as a promiscuous, not essential, player in its biology, because it is dispensable for normal growth and development in mice with no phenotypes (e.g., cancer) resulting from its genetic loss[77,80]. Contrarily, accumulating evidence clearly demonstrates that hyperactive Nrf2 promoted cancer development and malignance, because it is relevant to most of cancer hallmarks[76]. This implies that, except that Nrf2 can exert a significant cytoprotective function against diverse stresses so as to confer the cells to be acquired for the adaptive responses, its long-term hyper-activation can also critically inspire potential cancerous cells to be extricated from being rigidly controlled and confined by inevitability of the host multicellular cooperative evolution. The notion is supported by transcriptome sequencing of *Nrf2*-deficient cells, because this loss of Nrf2 causes hepatoma to be significantly ameliorated or completely prevented[29]. In the other way round, aberrant accumulation of hyperactive Nrf2 in *Nrf1*-deficient cells (or livers) leads to further malignant transformation of human hepatocellular carcinoma (HCC).

In particular, the spontaneous development of NASH-based inflammation and hepatoma are resulted from severe endogenous oxidative stress and fatal defects in basal constitutive gene expression affected by its genetic instability in mouse *Nrf1*-deficient livers[30,83,84]. Of note, TCF11 is absent in mice and all human hepatoma cell lines, but it, together with Nrf1α at a 1:1 ratio, exists in human normal cells (Figure S3). Upon *Nrf1α*-specific knockout from hepatoma cells, this results in cancer malgrowth and metastasis to the lungs in xenograft model mice[28,61]. The *Nrf1$\alpha^{-/-}$*-derived cancer deterioration is definitely resulted from severe oxidative stress, fateful defects in the redox metabolism reprogramming, and marked dysfunctions of fate-decisive gene regulatory networks, along with critical aberrant signaling transduction networks[21,29]. However, it is, to our great surprise, found that *Nrf1$\alpha^{-/-}$*-led malignant transformation is markedly alleviated and prevented by silencing of Nrf2; this is also accompanied by blocking oxidative stress upon Nrf2 knockdown[21,29]. More excitingly, restoration of *Nrf1α* is allowed for significant mitigation of *Nrf1$\alpha^{-/-}$*-exacerbated tumour to similar wild-type extents, while its malgrowth is further suppressed or even abolished by *TCF11* restoration (Figure 7B). This demonstrates that TCF11 is a potent tumour-suppressor than Nrf1α, while Nrf2 is a tumour-promotor, particularly in the *Nrf1$\alpha^{-/-}$* case. Altogether, loss of Nrf1α/TCF11's function, with hyperactive Nrf2 accumulation, results in liver cancer initiation and progression. This consequence is likely originated from endogenous oxidative stress-induced damages of mutant cells in aberrant metabolic inflammatory microenvironments. This gives rise to an evolvable selection force in so much of *Darwinian* dynamics, so that a clade of the mutant cancerous-prone cells is endowed with a self-defined fitness function and also specified to acquire for an independent of the host team optimum cooperative confinements, in order to behave themselves with own properties of cell division, proliferation and migration during carcinogenesis and progression.

Moreover, this work further demonstrates that the tumour-preventing effect of Nrf1α and TCF11 is accompanied by the constitutive activation or repression of critical genes for improving the overall survival

of patients with hepatocellular carcinoma (Figure 9). However, the activation or repression could not be ameliorated by compensation of ectopic *Nrf1α* and *TCF11* transfection into *Nrf1α/TCF11*-deficient cancer cells. This is surmised to be the relevance of given genes closely to the contexts within genome topological conformations, enabling ectopic Nrf1α/TCF11 to be allowed or forbidden for direct access to target genes.

Conclusion

In summary, four useful model cell lines stably expressing *TCF11*, *TCF11^{ΔN}*, *Nrf1α* or *Nrf2*, not mutants, are yielded herein. Their transcriptional expression levels are controlled by a tetracycline-inducible switch, but not monitored by one of redox inducers (e.g., sulforaphane or *tert*-Butylhydroquinone) or ER stressors (e.g., tunicamycin or thapsigargin). These cells were subjected to the integrative systems biology analyses of their omics data with routine reductionist approaches. To explore the essential distinctions between Nrf1α/TCF11 and Nrf2 in their contributions to critical constitutive gene expression profiles for basal redox metabolism, normal growth, development, cell homeostasis and organ integrity, we have provided holistic relevant data as much as possible, in the present digital network era. Notably, some seemingly-paradoxical data are still retained, because they may serve as vital nodes of a few negative feedback (and feedforward) regulatory circuits existing in the self-organizing systems of life. This is for the sake of an objective truth in life. Significantly, it is demonstrated that TCF11 serves as a more potent tumour-suppressor than Nrf1α at preventing cancer development and progression. This is defined by similar yet different regulatory profiles of both isoforms, with a striking disparity from Nrf2. Rather, a naturally-spliced mutant TCF11^{ΔN} resembles Nrf2 with largely consistent structure and function in regulating similar sets of target genes. Interestingly, the tumour-preventing effect of Nrf1α/TCF11 seems to be accompanied by certain constitutive activation or repression of critical genes for improving the overall survival rates of patients with hepatoma. Once loss of Nrf1α/TCF11's function, with hyperactive Nrf2 accumulated, this leads to severe endogenous oxidative damages, aberrant redox metabolic inflammation, and ultimate spontaneous hepatoma. Such genetic and nongenetic drivers could be integrated as a selection force in *Darwinian* dynamics to enable for stochastic speciation of *Nrf1*-deficient cells during carcinogenesis and ensuing progression, albeit this is required for deeply studies. Taken together, this study provides a holistic perspective to give a better understanding of essential differences between Nrf1α/TCF11 and Nrf2 in biology and medicine. Thereby, this facilitates drug discovery to induce Nrf1α/TCF11 as a new potent chemoprevention target against cancer.

Materials and Methods
Chemicals and antibodies

All chemicals were of the highest quality commercially available. Hygromycin B and blasticidin were purchased from Invitrogen Ltd, which were employed as double drug-screening to select putative positive clones from those transfected Flp-In™ T-REx™-293 expression cells. Tetracycline from Sangon Biotech Co (Shanghai, China) was utilized as an inducible reagent at a concentration of 1 μg/ml. Both cycloheximide (CHX) and MG132 were purchased from Sigma-Aldrich (St Louis, MO, USA). The antibody against Nrf1 proteins was acquired from our own lab (as indicated in Zhang's [85]), and all other antibodies were also employed against a V5 epitope (Invitrogen), Nrf2 (Abcam), CHP2 (Sangon Biotech), CPS1 (Abcam), FOXO1 (Cell Signaling Technology), IRS4 (Abcam), NKX2-8 (Sangon Biotech), AKR1B10 (Abcam), EPO (Proteintech Group), MUTYH (Sangon Biotech), PKM (Sangon Biotech), GP73 (Proteintech Group) or GPC3 (Proteintech Group), whilst β-Actin and secondary antibodies were from ZSGB-BIO (Beijing, China).

Cell lines, cell culture and transfection

These cell lines expressing *TCF11*, *TCF11^{ΔN}*, *Nrf1α*, *Nrf2*, as well as an empty control, were established by using the Flp-In™ T-REx™ system (Invitrogen). Their cDNA fragments, encoding human TCF11, TCF11^{ΔN} (with a deletion to remove the N-terminal 2-156 residues prior to the Neh2L subdomain from TCF11), Nrf1α and Nrf2, respectively, were cloned into pcDNA5/FRT/TO-V5 expression vector, before being cotransfected with a Flp-expressing pOG44 plasmid into the Flp-In™ T-REx™-293 host cells. The Flp recombinase was allowed for its homologous recombination at FRT sites existing in the host cells with each of the expression vectors. Of note, an empty expression vector was cotransfected into the host cell, to generate a negative control cell line. Then, the positive expression clones, as indicated, were selected by co-treatment of 150 μg/ml hygromycin B and 15 μg/ml blasticidin. All positively-selected cell lines were allowed for stably expression of *TCF11*, *TCF11^{ΔN}*, *Nrf1α* or *Nrf2*, beyond the negative control line, and thus referred to simply as *TCF11*, *TCF11^{ΔN}*, *Nrf1α*, *Nrf2* or *Control*, respectively. Moreover, HepG2, MHCC97L and MHCC97H were originated from ATCC (Zhong Yuan Ltd., Beijing, China), while both *Nrf1α^{-/-}* and *Nrf2^{-/-}* cell lines were created from wild-type HepG2 cells [28,29].

All experimental cells, except elsewhere indicated, were allowed for growth in DMEM basic medium (GIBCO, Life technologies), with being supplemented with 10% (v/v) foetal bovine serum (FBS, Biological Industries, Israel) and 100 units/ml of either of penicillin and streptomycin, in the 37℃ incubator with 5% CO_2. Notably, those expression constructs for human Nrf1α, TCF11 and Nrf2 were made by inserting each of their cDNA-encoding sequences into a pcDNA3 vector, respectively. The primer pairs used for this study were provided as shown in Table S9. The cell transfection with one of those indicated plasmids, alone or in combination, were carried out by using Lipofectamine®3000 Transfection Kit (Invitrogen) for 8 h, and then allowed for a 24-h recovery from transfection in the fresh medium before being subjected to the indicated experiments.

Quantitative real-time PCR

Each of experimental cell lines was subject to its total RNAs isolated by employing an RNAsimple Kit (Tiangen Biotech CO. LTD, Beijing, China). Total RNAs (1 μg) were added in a reverse-transcriptase reaction to yield the first strand of cDNAs (by another RevertAid First Strand Synthesis Kit, from Thermo), which served as the template of quantitative PCR in the GoTaq® qPCR Master Mix (Promega). Then, each pairs of all forward and reverse primers (as listed in Table S9) were also added in an indicated PCR, that was carried out in the following conditions at 95℃ for 3 min, followed by 40 cycles of 15 s at 95℃, and the last 30 s at 60℃. The final melting curve was validated to examine the amplification quality, and β-Actin at its mRNA expression levels served as an internal control for normalization.

Western blotting

Each of experimental cell lines was harvested in a lysis buffer (0.5% SDS, 0.04 mol/L DTT, pH 7.5) containing the protease inhibitor EASYpacks (Roche, Germany). The lysates were denatured immediately at 100℃ for 10 min, sonicated sufficiently, and diluted in 3 × loading buffer (187.5 nmol/L Tris-HCl, pH 6.8, 6% SDS, 30% Glycerol, 150 nmol/L DTT, 0.3% Bromophenol blue) at 100℃ for 5 min. Subsequently, equal amounts of protein extracts were subjected to separation by SDS-PAGE containing 4-15% polyacrylamide, followed by immunoblotting with each of distinct antibodies as indicated. On some occasions, the blotted

membranes were also stripped for 30 min and then re-probed with an additional antibody, whilst β-Actin served as an internal control to verify equal loading of protein in each of electrophoretic wells.

Transcriptome sequencing analysis

Total RNAs extracted from each of cell lines, that had incubated with 1 μg/ml of tetracycline for 12 h, were subjected to the transcriptome sequencing by Beijing Genomics Institute (BGI, Shenzhen, China) on an Illumina HiSeq 2000 sequencing system (Illumina, San Diego, CA). All detected mRNAs were fragmented into short fragments (~200bp). The clean reads were obtained during data filtering to remove the low-quality reads, and subjected to sequence mapping to the reference of human genome (GRCh37/hg19 from UCSC database) by using SOAP2 [86]. The resulting expression levels of given genes were calculated by the RPKM method [87]. All those differentially expressed genes (DEGs) were identified, with the criteria fold changes ≥ 2 or ≤ 0.5 and FDR (false discovery rate) ≤ 0.001, by the Poisson distribution model method (PossionDis) [88,89]. Such sequencing metadata have also been submitted to NCRI SRA (PRJNA501789). In addition, the DEGs were functional annotated by using the online tool DAVID (https://david.ncifcrf.gov/) to search their involved GO (gene ontology) terms and pathways, which were further classified with QuickGO (https://www.ebi.ac.uk/QuickGO/) and KEGG (https://www.kegg.jp/) databases.

Lentivirus-mediated restoration of *Nrf1α* or *TCF11*

One of the lentiviral-mediated expression constructs for *Nrf1α* or *TCF11*, that was designed by a help with supplier (GeneCopoeia, Guangzhou, China), together with the GFP-expressing lentiviral control vector, were co-transfected into our *Nrf1α*$^{-/-}$ cells, to establish *Nrf1α*-restored and *TCF11*-restored cell lines. Briefly, the lentiviral-packaging 293T cells (1×10^6) were seeded in a 10-cm dish and cultured in 10 ml DMEM supplemented with 10% FBS. Then, the mixture of 2.5 μg of each lentiviral ORF expression plasmid and 0.25 μg of the Lenti-Pac HIV plasmid in 15 μl of EndoFectin Lenti was incubated with 200 μl of Opti-MEM® (Invitrogen). The DNA-EndoFectin Lenti complex was directly added into cultured cells before being allowed for overnight incubation at 37℃ in a CO_2 incubator, followed by replaced by fresh medium supplemented with 5% FBS. Subsequently, a 1:500 volume of the TiterBoost reagent was further added to the above cultured media and allowed for continuous culture. The pseudovirus-containing culture media for 48 h post-transfection were collected by centrifuging as 500×*g* for 10 min. Then, the resulting lentivirus titer was estimated, prior to being subjected to efficient transfection of *Nrf1α*$^{-/-}$ hepatoma cells.

Subcutaneous tumour xenografts in nude mice

Mouse xenograft models were made by subcutaneous heterotransplantation of wild-type HepG2 (*WT*), *Nrf1α*$^{-/-}$, *Nrf1α*-restored and *TCF11*-restored cells, respectively, into nude mice as described [90]. Each line of experimental cells (1×10^7) were allowed for its exponential growth and then suspended in 0.2 ml of serum-free DMEM, before being inoculated subcutaneously into the right upper back region of male nude mice (BALB/C$^{nu/nu}$, 6 weeks, 18 g, from HFK Bioscience, Beijing) at a single site. The procedure of injection into all mice was complete within 30 min, and subsequent formation of the subcutaneous tumour xenografts was observed. The tumour sizes, after emerged, were measured every two days, until the 42nd day when all those mice were sacrificed and the transplanted tumours were excised. The sizes of growing tumours were calculated by a standard formula (i.e., $V = ab^2/2$) and shown graphically (n = 7 per group). Notably, all the mice were maintained under standard animal housing conditions with a 12-h dark

cycle and allowed access *ad libitum* to sterilized water and diet. All relevant studies were carried out on 8-week-old mice (with the license No. PIL60/13167) in accordance with United Kingdom Animal (Scientific Procedures) Act (1986) and the guidelines of the Animal Care and Use Committees of Chongqing University and the Third Military Medical University, both of which were also subjected to the local ethical review (in China). All relevant experimental protocols were approved by the University Laboratory Animal Welfare and Ethics Committee (with two institutional licenses SCXK-PLA-20120011 and SYXK-PLA-20120031). As for additional ethical concerns about the xenograft model mice bearing so big tumours insomuch as to give rise to certain bleeding ulcers, such a bad health condition of mice was only emerged from only day 2 prior to being sacrificed, and also such relevant study was indeed conducted according to the valid ethical regulations that have been approved.

The colony formation assay on soft agar

The cell culture plates (each with a diameter of 100 mm) were coated by the basement gel containing 0.6% soft agar mixed in the complete medium, upon which the upper gel containing 0.35% soft agar. Then, experimental cells (2×10^4, that had been growing in the exponential phase) was allowed for two-layer gel formation. Thereafter, the plates were cultured for 2-3 weeks in the incubator at 37℃ with 5% CO_2 before being stained with 1% crystal violet reagent (Sigma) and counted. This was performed according to the standard procedures [91].

The *in vitro* scratch assays

When experimental cells (1×10^5) grown in 6-well plates reached 70% confluency, they were allowed for synchronization by 12-h starvation in serum-free medium and then treated with 1 μg/ml of mitomycin C (from Cayman, USA) for 6 h. Subsequently, a clear 'scratch' in the cell monolayer was created and then allowed for being healed in the continuous culture at 37℃ with 5% CO_2. Thereafter, the cell migration was quantified as described previously [92].

The transwell-based migration and invasion assays

The transwell-based migration and invasion assays were conducted in the modified Boyden chambers (Transwell; Corning Inc. Lowell, MA, USA) as described previously [93]. When the growing cells reached 70% confluency, they were starved for 12 h in serum-free medium. The experimental cells (5×10^3) were suspended in 0.5 ml medium containing 5% FBS and seeded in the upper chamber of a transwell, which allows the cells to grow on the microporous polycarbonate membrane that is tissue culture-treated to enhance the cellular attachment to the bottom. The cell-seeded transwells were placed in each well of 24-well plates containing 1 ml of complete medium (i.e., the lower chamber), and then cultured for 24h in the incubator at 37℃ with 5% CO_2. Of note, the bottom of upper transwell was pre-coated by matrigel basement matrix (BD, Biosciences, USA), before the cells were placed in the invasion assay. The remaining cells in the upper chamber were removed, and the cells attached to the lower surface of the transwell membranes were fixed with 4% paraformaldehyde (AR10669, BOSTER) and stained with 1% crystal violet reagent (Sigma) before being counted.

Statistical analysis

Statistical significances were determined using either Student's *t*-test (for a comparison between two groups) or two-way ANOVA (for comparison among multiple groups). The relevant data presented herein are shown as a fold changes (mean $\pm$ SEM or $\pm$ SD) with significant differences that were calculated by the value of $p < 0.05$).

Data availability

All data needed to evaluate the conclusions in the paper are present in this publication along with the supplementary information documents that can be found online. Additional data related to this paper may also be requested from the corresponding author on reasonable request.

Supplementary Materials

Figure S1: Alignment of missing sequences between *Nrf1α* and *TCF11*, and statistical analysis of transcriptome sequencing data. Figure S2: Alterations in the cell cycle and apoptosis caused by restoration of *Nrf1α* and *TCF11*. Figure S3: Differential expression of *Nrf1α* and *TCF11* in different liver cancer cells and normal cells. Table S1: The differentially expressed genes regulated by TCF11. Table S2: The differentially expressed genes regulated by Nrf1α. Table S3: The differentially expressed genes regulated by Nrf2. Table S4: The differentially expressed genes regulated by TCF11$^{\Delta N}$. Table S5: The functional annotation of DEGs from group A to C. Table S6: The functional annotation of DEGs from group D to F. Table S7: The functional annotation of DEGs from group G to I. Table S8: The functional annotation of DEGs from group J to L. Table S9: The primers for construction or qPCR used in this paper.

Author Contributions

M.W. designed and performed most of the experiments and bioinformatic analyses, made all figures and wrote this manuscript draft. Y.R. constructed the *Nrf1α*- and *TCF11*-restored cell lines and performed the subcutaneous tumour xenografts experiment. Both S.H. and K.L. helped M.W. with molecular cloning to create several expression constructs and performed western blotting. L.Q. established both *Nrf1α$^{-/-}$* and *Nrf2$^{-/-}$* cell lines. Lastly, Y.Z. designed and supervised this study, analyzed and parsed the data, helped to prepare all figures, wrote and revised this manuscript.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (projects 81872336 and 82073079) awarded to Prof. Yiguo Zhang (University of Chongqing, China).

Conflicts of Interest

The authors declare no conflicts of interest.

References

1. Lambert, S.A.; Jolma, A.; Campitelli, L.F.; Das, P.K.; Yin, Y.; Albu, M.; Chen, X.; Taipale, J.; Hughes, T.R.; Weirauch, M.T. The Human Transcription Factors. *Cell* **2018**, *175*, 598-599, doi:10.1016/j.cell.2018.09.045.

2. Alber, T. Structure of the leucine zipper. *Current opinion in genetics & development* **1992**, *2*, 205-210, doi:10.1016/s0959-437x(05)80275-8.

3. Bader; Andreas, G. Leucine Zipper Transcription Factors: bZIP Proteins. **2006**, *10.1007/3-540-29623-9*, 964-967.

4. Sykiotis, G.P.; Bohmann, D. Stress-activated cap'n'collar transcription factors in aging and human disease. *Science signaling* **2010**, *3*, re3, doi:10.1126/scisignal.3112re3.

5. Zhang, Y.; Xiang, Y. Molecular and cellular basis for the unique functioning of Nrf1, an indispensable transcription factor for maintaining cell homoeostasis and organ integrity. *The Biochemical journal* **2016**, *473*, 961-1000, doi:10.1042/BJ20151182.

6. Zhu, Y.P.; Wang, M.; Xiang, Y.; Qiu, L.; Hu, S.; Zhang, Z.; Mattjus, P.; Zhu, X.; Zhang, Y. Nach Is a Novel Subgroup at an Early Evolutionary Stage of the CNC-bZIP Subfamily Transcription Factors from the Marine Bacteria to Humans. *International journal of molecular sciences* **2018**, *19*, doi:10.3390/ijms19102927.

7. Marini, M.G.; Chan, K.; Casula, L.; Kan, Y.W.; Cao, A.; Moi, P. hMAF, a small human transcription factor that heterodimerizes specifically with Nrf1 and Nrf2. *The Journal of biological chemistry* **1997**, *272*, 16490-16497, doi:10.1074/jbc.272.26.16490.

8. Katsuoka, F.; Yamamoto, M. Small Maf proteins (MafF, MafG, MafK): History, structure and function. *Gene* **2016**, *586*, 197-205, doi:10.1016/j.gene.2016.03.058.

9. Venugopal, R.; Jaiswal, A.K. Nrf2 and Nrf1 in association with Jun proteins regulate antioxidant response element-mediated expression and coordinated induction of genes encoding detoxifying enzymes. *Oncogene* **1998**, *17*, 3145-3156, doi:10.1038/sj.onc.1202237.

10. Murphy, P.; Kolsto, A. Expression of the bZIP transcription factor TCF11 and its potential dimerization partners during development. *Mechanisms of development* **2000**, *97*, 141-148, doi:10.1016/s0925-4773(00)00413-5.

11. Kim, H.M.; Han, J.W.; Chan, J.Y. Nuclear Factor Erythroid-2 Like 1 (NFE2L1): Structure, function and regulation. *Gene* **2016**, *584*, 17-25, doi:10.1016/j.gene.2016.03.002.

12. Chevillard, G.; Blank, V. NFE2L3 (NRF3): the Cinderella of the Cap'n'Collar transcription factors. *Cellular and molecular life sciences : CMLS* **2011**, *68*, 3337-3348, doi:10.1007/s00018-011-0747-x.

13. Chan, J.Y.; Kwong, M.; Lu, R.; Chang, J.; Wang, B.; Yen, T.S.; Kan, Y.W. Targeted disruption of the ubiquitous CNC-bZIP transcription factor, Nrf-1, results in anemia and embryonic lethality in mice. *The EMBO journal* **1998**, *17*, 1779-1787, doi:10.1093/emboj/17.6.1779.

14. Leung, L.; Kwong, M.; Hou, S.; Lee, C.; Chan, J.Y. Deficiency of the Nrf1 and Nrf2 transcription factors results in early embryonic lethality and severe oxidative stress. *The Journal of biological chemistry* **2003**, *278*, 48021-48029, doi:10.1074/jbc.M308439200.

15. Narayanan, K.; Ramachandran, A.; Peterson, M.C.; Hao, J.; Kolsto, A.B.; Friedman, A.D.; George, A. The CCAAT enhancer-binding protein (C/EBP)beta and Nrf1 interact to regulate dentin sialophosphoprotein (DSPP) gene expression during odontoblast differentiation. *The Journal of biological chemistry* **2004**, *279*, 45423-45432, doi:10.1074/jbc.M405031200.

16. Kim, J.; Xing, W.; Wergedal, J.; Chan, J.Y.; Mohan, S. Targeted disruption of nuclear factor erythroid-derived 2-like 1 in osteoblasts reduces bone size and bone formation in mice. *Physiological genomics* **2010**, *40*, 100-110, doi:10.1152/physiolgenomics.00105.2009.

17. Xing, W.; Singgih, A.; Kapoor, A.; Alarcon, C.M.; Baylink, D.J.; Mohan, S. Nuclear factor-E2-related factor-1 mediates ascorbic acid induction of osterix expression via interaction with antioxidant-responsive element in bone cells. *The Journal of biological chemistry* **2007**, *282*, 22052-22061, doi:10.1074/jbc.M702614200.

18. Lee, C.S.; Lee, C.; Hu, T.; Nguyen, J.M.; Zhang, J.; Martin, M.V.; Vawter, M.P.; Huang, E.J.; Chan, J.Y. Loss of nuclear factor E2-related factor 1 in the brain leads to dysregulation of proteasome gene expression and neurodegeneration. *Proceedings of the National Academy of Sciences of the United States of America* **2011**, *108*, 8408-8413, doi:10.1073/pnas.1019209108.

19. Lee, C.S.; Ho, D.V.; Chan, J.Y. Nuclear factor-erythroid 2-related factor 1 regulates expression of proteasome genes in hepatocytes and protects against endoplasmic reticulum stress and steatosis in mice. *The FEBS journal* **2013**, *280*, 3609-3620, doi:10.1111/febs.12350.

20. Sha, Z.; Goldberg, A.L. Proteasome-mediated processing of Nrf1 is essential for coordinate induction of all proteasome subunits and p97. *Current biology : CB* **2014**, *24*, 1573-1583, doi:10.1016/j.cub.2014.06.004.

21. Zhu, Y.P.; Zheng, Z.; Xiang, Y.; Zhang, Y. Glucose Starvation-Induced Rapid Death of Nrf1alpha-Deficient, but Not Nrf2-Deficient, Hepatoma Cells Results from Its Fatal Defects in the Redox Metabolism Reprogramming. *Oxidative medicine and cellular longevity* **2020**, *2020*, 4959821, doi:10.1155/2020/4959821.

22. Hirotsu, Y.; Hataya, N.; Katsuoka, F.; Yamamoto, M. NF-E2-related factor 1 (Nrf1) serves as a novel regulator of hepatic lipid metabolism through regulation of the Lipin1 and PGC-1beta genes. *Molecular and cellular biology* **2012**, *32*, 2760-2770, doi:10.1128/MCB.06706-11.

23. Hirotsu, Y.; Higashi, C.; Fukutomi, T.; Katsuoka, F.; Tsujita, T.; Yagishita, Y.; Matsuyama, Y.; Motohashi, H.; Uruno, A.; Yamamoto, M. Transcription factor NF-E2-related factor 1 impairs glucose metabolism in mice. *Genes to cells : devoted to molecular & cellular mechanisms* **2014**, *19*, 650-665, doi:10.1111/gtc.12165.

24. Hou, Y.; Liu, Z.; Zuo, Z.; Gao, T.; Fu, J.; Wang, H.; Xu, Y.; Liu, D.; Yamamoto, M.; Zhu, B., et al. Adipocyte-specific

deficiency of Nfe2l1 disrupts plasticity of white adipose tissues and metabolic homeostasis in mice. *Biochemical and biophysical research communications* **2018**, *503*, 264-270, doi:10.1016/j.bbrc.2018.06.013.

25. Novotny, V.; Prieschl, E.E.; Csonga, R.; Fabjani, G.; Baumruker, T. Nrf1 in a complex with fosB, c-jun, junD and ATF2 forms the AP1 component at the TNF alpha promoter in stimulated mast cells. *Nucleic acids research* **1998**, *26*, 5480-5485, doi:10.1093/nar/26.23.5480.

26. Prieschl, E.E.; Novotny, V.; Csonga, R.; Jaksche, D.; Elbe-Burger, A.; Thumb, W.; Auer, M.; Stingl, G.; Baumruker, T. A novel splice variant of the transcription factor Nrf1 interacts with the TNFalpha promoter and stimulates transcription. *Nucleic acids research* **1998**, *26*, 2291-2297, doi:10.1093/nar/26.10.2291.

27. Berg, D.T.; Gupta, A.; Richardson, M.A.; O'Brien, L.A.; Calnek, D.; Grinnell, B.W. Negative regulation of inducible nitric-oxide synthase expression mediated through transforming growth factor-beta-dependent modulation of transcription factor TCF11. *The Journal of biological chemistry* **2007**, *282*, 36837-36844, doi:10.1074/jbc.M706909200.

28. Ren, Y.G.; Qiu, L.; Lu, F.L.; Ru, X.F.; Li, S.J.; Xiang, Y.C.; Yu, S.W.; Zhang, Y.G. TALENs-directed knockout of the full-length transcription factor Nrf1 alpha that represses malignant behaviour of human hepatocellular carcinoma (HepG2) cells. *Sci Rep-Uk* **2016**, *6*, doi:10.1038/Srep23775.

29. Qiu, L.; Wang, M.; Hu, S.F.; Ru, X.F.; Ren, Y.G.; Zhang, Z.W.; Yu, S.W.; Zhang, Y.G. Oncogenic Activation of Nrf2, Though as a Master Antioxidant Transcription Factor, Liberated by Specific Knockout of the Full-Length Nrf1 alpha that Acts as a Dominant Tumor Repressor. *Cancers* **2018**, *10*, doi:10.3390/cancers10120520.

30. Ohtsuji, M.; Katsuoka, F.; Kobayashi, A.; Aburatani, H.; Hayes, J.D.; Yamamoto, M. Nrf1 and Nrf2 play distinct roles in activation of antioxidant response element-dependent genes. *The Journal of biological chemistry* **2008**, *283*, 33554-33562, doi:10.1074/jbc.M804597200.

31. Kwong, M.; Kan, Y.W.; Chan, J.Y. The CNC basic leucine zipper factor, Nrf1, is essential for cell survival in response to oxidative stress-inducing agents. Role for Nrf1 in gamma-gcs(l) and gss expression in mouse fibroblasts. *The Journal of biological chemistry* **1999**, *274*, 37491-37498, doi:10.1074/jbc.274.52.37491.

32. Chen, L.; Kwong, M.; Lu, R.; Ginzinger, D.; Lee, C.; Leung, L.; Chan, J.Y. Nrf1 is critical for redox balance and survival of liver cells during development. *Molecular and cellular biology* **2003**, *23*, 4673-4686, doi:10.1128/mcb.23.13.4673-4686.2003.

33. Osburn, W.O.; Wakabayashi, N.; Misra, V.; Nilles, T.; Biswal, S.; Trush, M.A.; Kensler, T.W. Nrf2 regulates an adaptive response protecting against oxidative damage following diquat-mediated formation of superoxide anion. *Archives of biochemistry and biophysics* **2006**, *454*, 7-15, doi:10.1016/j.abb.2006.08.005.

34. Ma, Q. Role of nrf2 in oxidative stress and toxicity. *Annual review of pharmacology and toxicology* **2013**, *53*, 401-426, doi:10.1146/annurev-pharmtox-011112-140320.

35. Kitteringham, N.R.; Abdullah, A.; Walsh, J.; Randle, L.; Jenkins, R.E.; Sison, R.; Goldring, C.E.; Powell, H.; Sanderson, C.; Williams, S., et al. Proteomic analysis of Nrf2 deficient transgenic mice reveals cellular defence and lipid metabolism as primary Nrf2-dependent pathways in the liver. *Journal of proteomics* **2010**, *73*, 1612-1631, doi:10.1016/j.jprot.2010.03.018.

36. Mitsuishi, Y.; Taguchi, K.; Kawatani, Y.; Shibata, T.; Nukiwa, T.; Aburatani, H.; Yamamoto, M.; Motohashi, H. Nrf2 redirects glucose and glutamine into anabolic pathways in metabolic reprogramming. *Cancer cell* **2012**, *22*, 66-79, doi:10.1016/j.ccr.2012.05.016.

37. Lee, S.B.; Sellers, B.N.; DeNicola, G.M. The Regulation of NRF2 by Nutrient-Responsive Signaling and Its Role in Anabolic Cancer Metabolism. *Antioxidants & redox signaling* **2018**, *29*, 1774-1791, doi:10.1089/ars.2017.7356.

38. Singh, B.; Chatterjee, A.; Ronghe, A.M.; Bhat, N.K.; Bhat, H.K. Antioxidant-mediated up-regulation of OGG1 via NRF2 induction is associated with inhibition of oxidative DNA damage in estrogen-induced breast cancer. *BMC cancer* **2013**, *13*, 253, doi:10.1186/1471-2407-13-253.

39. Jayakumar, S.; Pal, D.; Sandur, S.K. Nrf2 facilitates repair of radiation induced DNA damage through homologous recombination repair pathway in a ROS independent manner in cancer cells. *Mutation research* **2015**, *779*, 33-45, doi:10.1016/j.mrfmmm.2015.06.007.

40. Fan, Z.; Wirth, A.K.; Chen, D.; Wruck, C.J.; Rauh, M.; Buchfelder, M.; Savaskan, N. Nrf2-Keap1 pathway promotes cell proliferation and diminishes ferroptosis. *Oncogenesis* **2017**, *6*, e371, doi:10.1038/oncsis.2017.65.

41. Murakami, S.; Motohashi, H. Roles of Nrf2 in cell proliferation and differentiation. *Free radical biology & medicine* **2015**, *88*, 168-178, doi:10.1016/j.freeradbiomed.2015.06.030.

42. Sun, X.; Ou, Z.; Chen, R.; Niu, X.; Chen, D.; Kang, R.; Tang, D. Activation of the p62-Keap1-NRF2 pathway protects against ferroptosis in hepatocellular carcinoma cells. *Hepatology* **2016**, *63*, 173-184, doi:10.1002/hep.28251.

43. Torrente, L.; Sanchez, C.; Moreno, R.; Chowdhry, S.; Cabello, P.; Isono, K.; Koseki, H.; Honda, T.; Hayes, J.D.; Dinkova-Kostova, A.T., et al. Crosstalk between NRF2 and HIPK2 shapes cytoprotective responses. *Oncogene* **2017**, *36*, 6204-6212, doi:10.1038/onc.2017.221.

44. Niture, S.K.; Jaiswal, A.K. Nrf2-induced antiapoptotic Bcl-xL protein enhances cell survival and drug resistance. *Free radical biology & medicine* **2013**, *57*, 119-131, doi:10.1016/j.freeradbiomed.2012.12.014.

45. Li, L.; Pan, H.; Wang, H.; Li, X.; Bu, X.; Wang, Q.; Gao, Y.; Wen, G.; Zhou, Y.; Cong, Z., et al. Interplay between VEGF and Nrf2 regulates angiogenesis due to intracranial venous hypertension. *Sci Rep* **2016**, *6*, 37338, doi:10.1038/srep37338.

46. Ji, X.; Wang, H.; Zhu, J.; Zhu, L.; Pan, H.; Li, W.; Zhou, Y.; Cong, Z.; Yan, F.; Chen, S. Knockdown of Nrf2 suppresses glioblastoma angiogenesis by inhibiting hypoxia-induced activation of HIF-1alpha. *International journal of cancer*

2014, *135*, 574-584, doi:10.1002/ijc.28699.

47.   Kim, T.H.; Hur, E.G.; Kang, S.J.; Kim, J.A.; Thapa, D.; Lee, Y.M.; Ku, S.K.; Jung, Y.; Kwak, M.K. NRF2 blockade suppresses colon tumor angiogenesis by inhibiting hypoxia-induced activation of HIF-1alpha. *Cancer research* **2011**, *71*, 2260-2275, doi:10.1158/0008-5472.CAN-10-3007.

48.   Jaramillo, M.C.; Zhang, D.D. The emerging role of the Nrf2-Keap1 signaling pathway in cancer. *Genes & development* **2013**, *27*, 2179-2191, doi:10.1101/gad.225680.113.

49.   Bauer, A.K.; Cho, H.Y.; Miller-Degraff, L.; Walker, C.; Helms, K.; Fostel, J.; Yamamoto, M.; Kleeberger, S.R. Targeted deletion of Nrf2 reduces urethane-induced lung tumor development in mice. *PloS one* **2011**, *6*, e26590, doi:10.1371/journal.pone.0026590.

50.   Menegon, S.; Columbano, A.; Giordano, S. The Dual Roles of NRF2 in Cancer. *Trends in molecular medicine* **2016**, *22*, 578-593, doi:10.1016/j.molmed.2016.05.002.

51.   Zhang, L.; Wang, N.; Zhou, S.; Ye, W.; Jing, G.; Zhang, M. Propofol induces proliferation and invasion of gallbladder cancer cells through activation of Nrf2. *Journal of experimental & clinical cancer research : CR* **2012**, *31*, 66, doi:10.1186/1756-9966-31-66.

52.   Pan, H.; Wang, H.; Zhu, L.; Mao, L.; Qiao, L.; Su, X. The role of Nrf2 in migration and invasion of human glioma cell U251. *World neurosurgery* **2013**, *80*, 363-370, doi:10.1016/j.wneu.2011.06.063.

53.   Shen, H.; Yang, Y.; Xia, S.; Rao, B.; Zhang, J.; Wang, J. Blockage of Nrf2 suppresses the migration and invasion of esophageal squamous cell carcinoma cells in hypoxic microenvironment. *Diseases of the esophagus : official journal of the International Society for Diseases of the Esophagus* **2014**, *27*, 685-692, doi:10.1111/dote.12124.

54.   Widenmaier, S.B.; Snyder, N.A.; Nguyen, T.B.; Arduini, A.; Lee, G.Y.; Arruda, A.P.; Saksi, J.; Bartelt, A.; Hotamisligil, G.S. NRF1 Is an ER Membrane Sensor that Is Central to Cholesterol Homeostasis. *Cell* **2017**, *171*, 1094-1109 e1015, doi:10.1016/j.cell.2017.10.003.

55.   Yu-ping Zhu, Z.Z., Shaofan Hu, Xufang Ru, Zhuo Fan, Lu Qiu and Yiguo Zhang Unification of Opposites between Two Antioxidant Transcription Factors Nrf1 and Nrf2 in Mediating Distinct Cellular Responses to the Endoplasmic Reticulum Stressor Tunicamycin. *antioxidants* **2019**, *9*, doi:10.3390/antiox9010004.

56.   Wang, M.; Qiu, L.; Ru, X.; Song, Y.; Zhang, Y. Distinct isoforms of Nrf1 diversely regulate different subsets of its cognate target genes. *Sci Rep* **2019**, *9*, 2960, doi:10.1038/s41598-019-39536-0.

57.   Luna, L.; Skammelsrud, N.; Johnsen, O.; Abel, K.J.; Weber, B.L.; Prydz, H.; Kolsto, A.B. Structural organization and mapping of the human TCF11 gene. *Genomics* **1995**, *27*, 237-244, doi:10.1006/geno.1995.1037.

58.   Zhang, Y.; Hayes, J.D. The membrane-topogenic vectorial behaviour of Nrf1 controls its post-translational modification and transactivation activity. *Sci Rep* **2013**, *3*, 2006, doi:10.1038/srep02006.

59.   Liu, P.; Kerins, M.J.; Tian, W.; Neupane, D.; Zhang, D.D.; Ooi, A. Differential and overlapping targets of the transcriptional regulators NRF1, NRF2, and NRF3 in human cells. *The Journal of biological chemistry* **2019**, *294*, 18131-18149, doi:10.1074/jbc.RA119.009591.

60.   Ibrahim, L.; Mesgarzadeh, J.; Xu, I.; Powers, E.T.; Wiseman, R.L.; Bollong, M.J. Defining the Functional Targets of Cap'n'collar Transcription Factors NRF1, NRF2, and NRF3. *Antioxidants (Basel)* **2020**, *9*, doi:10.3390/antiox9101025.

61.   Chen, J.; Wang, M.; Xiang, Y.; Ru, X.; Ren, Y.; Liu, X.; Qiu, L.; Zhang, Y. Nrf1 Is Endowed with a Dominant Tumor-Repressing Effect onto the Wnt/beta-Catenin-Dependent and Wnt/beta-Catenin-Independent Signaling Networks in the Human Liver Cancer. *Oxidative medicine and cellular longevity* **2020**, *2020*, 5138539, doi:10.1155/2020/5138539.

62.   Menyhart, O.; Nagy, A.; Gyorffy, B. Determining consistent prognostic biomarkers of overall survival and vascular invasion in hepatocellular carcinoma. *Royal Society open science* **2018**, *5*, 181006, doi:10.1098/rsos.181006.

63.   Liang, R.; Liu, Z.; Piao, X.; Zuo, M.; Zhang, J.; Liu, Z.; Li, Y.; Lin, Y. Research progress on GP73 in malignant tumors. *OncoTargets and therapy* **2018**, *11*, 7417-7421, doi:10.2147/OTT.S181239.

64.   Ye, J.Z.; Yan, S.M.; Yuan, C.L.; Wu, H.N.; Zhang, J.Y.; Liu, Z.H.; Li, Y.Q.; Luo, X.L.; Lin, Y.; Liang, R. GP73 level determines chemotherapeutic resistance in human hepatocellular carcinoma cells. *Journal of Cancer* **2018**, *9*, 415-423, doi:10.7150/jca.19185.

65.   Yu, M.; Luo, H.; Fan, M.; Wu, X.; Shi, B.; Di, S.; Liu, Y.; Pan, Z.; Jiang, H.; Li, Z. Development of GPC3-Specific Chimeric Antigen Receptor-Engineered Natural Killer Cells for the Treatment of Hepatocellular Carcinoma. *Molecular therapy : the journal of the American Society of Gene Therapy* **2018**, *26*, 366-378, doi:10.1016/j.ymthe.2017.12.012.

66.   Zhang, Q.; Han, Z.; Tao, J.; Zhao, M.; Zhang, W.; Li, P.; Tang, L.; Gu, Y. An innovative peptide with high affinity to GPC3 for hepatocellular carcinoma diagnosis. *Biomaterials science* **2018**, *7*, 159-167, doi:10.1039/c8bm01016a.

67.   Zhang, Y.; Ren, Y.; Li, S.; Hayes, J.D. Transcription factor Nrf1 is topologically repartitioned across membranes to enable target gene transactivation through its acidic glucose-responsive domains. *PloS one* **2014**, *9*, e93458, doi:10.1371/journal.pone.0093458.

68.   Lehrbach, N.J.; Breen, P.C.; Ruvkun, G. Protein Sequence Editing of SKN-1A/Nrf1 by Peptide:N-Glycanase Controls Proteasome Gene Expression. *Cell* **2019**, *177*, 737-750 e715, doi:10.1016/j.cell.2019.03.035.

69.   Tomlin, F.M.; Gerling-Driessen, U.I.M.; Liu, Y.C.; Flynn, R.A.; Vangala, J.R.; Lentz, C.S.; Clauder-Muenster, S.; Jakob, P.; Mueller, W.F.; Ordonez-Rueda, D., et al. Inhibition of NGLY1 Inactivates the Transcription Factor Nrf1 and Potentiates Proteasome Inhibitor Cytotoxicity. *ACS central science* **2017**, *3*, 1143-1155, doi:10.1021/acscentsci.7b00224.

70. Owings, K.G.; Lowry, J.B.; Bi, Y.; Might, M.; Chow, C.Y. Transcriptome and functional analysis in a Drosophila model of NGLY1 deficiency provides insight into therapeutic approaches. *Human molecular genetics* **2018**, *27*, 1055-1066, doi:10.1093/hmg/ddy026.

71. van Keulen, B.J.; Rotteveel, J.; Finken, M.J.J. Unexplained death in patients with NGLY1 mutations may be explained by adrenal insufficiency. *Physiological reports* **2019**, *7*, e13979, doi:10.14814/phy2.13979.

72. Xiang, Y.; Halin, J.; Fan, Z.; Hu, S.; Wang, M.; Qiu, L.; Zhang, Z.; Mattjus, P.; Zhang, Y. Topovectorial mechanisms control the juxtamembrane proteolytic processing of Nrf1 to remove its N-terminal polypeptides during maturation of the CNC-bZIP factor. *Toxicology and applied pharmacology* **2018**, *360*, 160-184, doi:10.1016/j.taap.2018.09.039.

73. Xiang, Y.; Wang, M.; Hu, S.; Qiu, L.; Yang, F.; Zhang, Z.; Yu, S.; Pi, J.; Zhang, Y. Mechanisms controlling the multistage post-translational processing of endogenous Nrf1alpha/TCF11 proteins to yield distinct isoforms within the coupled positive and negative feedback circuits. *Toxicology and applied pharmacology* **2018**, *360*, 212-235, doi:10.1016/j.taap.2018.09.036.

74. Zhang, Y.; Li, S.; Xiang, Y.; Qiu, L.; Zhao, H.; Hayes, J.D. The selective post-translational processing of transcription factor Nrf1 yields distinct isoforms that dictate its ability to differentially regulate gene expression. *Sci Rep* **2015**, *5*, 12983, doi:10.1038/srep12983.

75. Itoh, K.; Chiba, T.; Takahashi, S.; Ishii, T.; Igarashi, K.; Katoh, Y.; Oyake, T.; Hayashi, N.; Satoh, K.; Hatayama, I., et al. An Nrf2/small Maf heterodimer mediates the induction of phase II detoxifying enzyme genes through antioxidant response elements. *Biochemical and biophysical research communications* **1997**, *236*, 313-322, doi:10.1006/bbrc.1997.6943.

76. Rojo de la Vega, M.; Chapman, E.; Zhang, D.D. NRF2 and the Hallmarks of Cancer. *Cancer cell* **2018**, *34*, 21-43, doi:10.1016/j.ccell.2018.03.022.

77. Yamamoto, M.; Kensler, T.W.; Motohashi, H. The KEAP1-NRF2 System: a Thiol-Based Sensor-Effector Apparatus for Maintaining Redox Homeostasis. *Physiological reviews* **2018**, *98*, 1169-1203, doi:10.1152/physrev.00023.2017.

78. Cuadrado, A.; Rojo, A.I.; Wells, G.; Hayes, J.D.; Cousin, S.P.; Rumsey, W.L.; Attucks, O.C.; Franklin, S.; Levonen, A.L.; Kensler, T.W., et al. Therapeutic targeting of the NRF2 and KEAP1 partnership in chronic diseases. *Nature reviews. Drug discovery* **2019**, *18*, 295-317, doi:10.1038/s41573-018-0008-x.

79. Xu, C.; Huang, M.T.; Shen, G.; Yuan, X.; Lin, W.; Khor, T.O.; Conney, A.H.; Kong, A.N. Inhibition of 7,12-dimethylbenz(a)anthracene-induced skin tumorigenesis in C57BL/6 mice by sulforaphane is mediated by nuclear factor E2-related factor 2. *Cancer research* **2006**, *66*, 8293-8296, doi:10.1158/0008-5472.CAN-06-0300.

80. Chan, K.; Lu, R.; Chang, J.C.; Kan, Y.W. NRF2, a member of the NFE2 family of transcription factors, is not essential for murine erythropoiesis, growth, and development. *Proceedings of the National Academy of Sciences of the United States of America* **1996**, *93*, 13943-13948, doi:10.1073/pnas.93.24.13943.

81. Lehrbach, N.J.; Ruvkun, G. Endoplasmic reticulum-associated SKN-1A/Nrf1 mediates a cytoplasmic unfolded protein response and promotes longevity. *eLife* **2019**, *8*, doi:10.7554/eLife.44425.

82. Schoenfelder, S.; Sugar, R.; Dimond, A.; Javierre, B.M.; Armstrong, H.; Mifsud, B.; Dimitrova, E.; Matheson, L.; Tavares-Cadete, F.; Furlan-Magaril, M., et al. Polycomb repressive complex PRC1 spatially constrains the mouse embryonic stem cell genome. *Nature genetics* **2015**, *47*, 1179-1186, doi:10.1038/ng.3393.

83. Xu, Z.; Chen, L.; Leung, L.; Yen, T.S.; Lee, C.; Chan, J.Y. Liver-specific inactivation of the Nrf1 gene in adult mouse leads to nonalcoholic steatohepatitis and hepatic neoplasia. *Proceedings of the National Academy of Sciences of the United States of America* **2005**, *102*, 4120-4125, doi:10.1073/pnas.0500660102.

84. Oh, D.H.; Rigas, D.; Cho, A.; Chan, J.Y. Deficiency in the nuclear-related factor erythroid 2 transcription factor (Nrf1) leads to genetic instability. *The FEBS journal* **2012**, *279*, 4121-4130, doi:10.1111/febs.12005.

85. Zhang, Y.; Hayes, J.D. Identification of topological determinants in the N-terminal domain of transcription factor Nrf1 that control its orientation in the endoplasmic reticulum membrane. *The Biochemical journal* **2010**, *430*, 497-510, doi:10.1042/BJ20100471.

86. Li, R.; Yu, C.; Li, Y.; Lam, T.W.; Yiu, S.M.; Kristiansen, K.; Wang, J. SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics* **2009**, *25*, 1966-1967, doi:10.1093/bioinformatics/btp336.

87. Mortazavi, A.; Williams, B.A.; McCue, K.; Schaeffer, L.; Wold, B. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nature methods* **2008**, *5*, 621-628, doi:10.1038/nmeth.1226.

88. Benjamini, Y.; Yekutieli, D. The control of the false discovery rate in multiple testing under dependency. *Ann Stat* **2001**, *29*, 1165-1188.

89. Audic, S.; Claverie, J.M. The significance of digital gene expression profiles. *Genome Res* **1997**, *7*, 986-995.

90. Morton, C.L.; Houghton, P.J. Establishment of human tumor xenografts in immunodeficient mice. *Nature protocols* **2007**, *2*, 247-250, doi:10.1038/nprot.2007.25.

91. Dong, Z.; Cmarik, J.L. Harvesting cells under anchorage-independent cell transformation conditions for biochemical analyses. *Science's STKE : signal transduction knowledge environment* **2002**, *2002*, pl7, doi:10.1126/stke.2002.130.pl7.

92. Liang, C.C.; Park, A.Y.; Guan, J.L. In vitro scratch assay: a convenient and inexpensive method for analysis of cell migration in vitro. *Nature protocols* **2007**, *2*, 329-333, doi:10.1038/nprot.2007.30.

93. Berginc, K.; Kristl, A. Transwell-grown HepG2 cell monolayers as in vitro permeability model to study drug-drug or drug-food interactions. *Journal of medicinal food* **2011**, *14*, 135-139, doi:10.1089/jmf.2010.0041.